Seasonal Effects of Environmental Change on Yellow-Bellied Marmot

Population Dynamics

<u>Running title</u>: Seasonal mechanisms of population dynamics

<u>Authors</u>: Maria Paniw[1], Dylan Childs[2], Kenneth B Armitage[3], Daniel T Blumstein[4,5], Julien

Martin[6], Madan Oli[7], Arpat Ozgul[1]

1 - Department of Evolutionary Biology and Environmental Studies, University of Zurich,
Winterthurerstrasse 190, CH-8057 Zurich, Switzerland
2 - Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield S10 2TN,
UK
3 - Ecology & Evolutionary Biology Department, The University of Kansas, Lawrence, KS 66045-7534,
USA
4 - Department of Ecology and Evolutionary Biology, University of California Los Angeles, Los Angeles,
CA 90095, USA
5 - The Rocky Mountain Biological Laboratory, Crested Butte, CO 81224, USA
6 - School of Biological Sciences, University of Aberdeen, Aberdeen, AB24 2TZ, UK
7 - Department of Wildlife Ecology, University of Florida, Gainesville, FL 32611, USA
18
19
20 [*]<u>Corresponding author</u>: Tel.: +34-671-246-338; email: maria.paniw@ieu.uzh.ch; ORCID ID: 0000-0002-
21 1949-4448
22

ABSTRACT

Environmental factors drive the persistence of natural populations by causing complex,

covarying responses in demographic processes (*i.e.*, survival, growth, and reproduction). As

most natural populations inhabit seasonal environments, overlooking seasonal differences in this

covariation may obscure the mechanisms that buffer or amplify population responses to

environmental change. We investigated how environment-driven covariation of seasonal

demographic processes affects population dynamics using 40 years of individual-based data from

a population of yellow-bellied marmots (*Marmota flaviventer*). We first used a factor-analytic

approach to jointly model demographic processes as a function of a latent variable describing

yearly environmental quality. We then parameterized, perturbed, and projected into the future a

seasonal population model that explicitly incorporated the covarying demographic responses to

the environment. The projections were based on various scenarios of environmental change,

including changes in the trend and temporal patterning of environmental quality. We show that,

although demographic processes in both the summer and winter season contributed to population

fitness, yearly environmental quality exerted the strongest effect on winter demography only.

Simultaneous, negative responses to declining average environmental quality of winter survival

and mass and reproductive-status change resulted in an increased risk of population quasi-

extinction, regardless of summer demography. We emphasize that assessing seasonal

demographic effects is key to understanding population persistence in the face of environmental

change and should receive much more attention in ecological theory and conservation

management.

INTRODUCTION

Effects of environmental change on survival, growth, and reproduction are typically investigated

based on annual transitions among life-history stages in structured population models (Paniw,

Ozgul, & Salguero-Gómez, 2018; Salguero-Gómez et al., 2016). However, all natural

ecosystems show some level of seasonal fluctuations in environmental conditions, and numerous

species have evolved life cycles that are cued to such seasonality (Ruf, Bieber, Arnold, &

Millesi, 2012; Varpe, 2017). For example, most temperate- and many arid-environment species

show strong differences in survival and growth among seasons, with reproduction being confined

mostly to one season (Childs, Coulson, Pemberton, Clutton-Brock, & Rees, 2011; Rushing et al.,

2017; Woodroffe, Groom, & McNutt, 2017). Species with highly adapted, seasonal life cycles

are likely to be particularly vulnerable to environmental change, even if they are relatively long-

lived (Campos et al., 2017; Jenouvrier et al., 2012; Paniw, Maag, Cozzi, Clutton-Brock, &

Ozgul, 2019). This is because adverse environmental conditions in one season may carry-over to

the next season in which key life-history events (e.g., reproduction) occur, and can negate

positive season-specific environmental effects (Marra, Cohen, Loss, Rutter, & Tonra, 2015). For

instance, in species where individual traits such as body mass determine demographic rates,

environment-driven changes in the trait distribution in one season can affect trait-dependent

demographic rates in the next season (Bassar, Letcher, Nislow, & Whiteley, 2016; Paniw et al.,

2019). Investigating annual dynamics, averaged over multiple seasons, may, therefore, obscure

the mechanisms that allow populations to persist under environmental change.

Despite the potential to gain a more mechanistic view of population dynamics, modeling

the effects of seasonal environmental change is an analytically complex and data-hungry

endeavor (Bassar et al., 2016; Benton, Plaistow, & Coulson, 2006). This is in part because

multiple environmental factors that change throughout the year can interact with each other and

individual-level (e.g., body mass) or population-level factors (e.g., density dependence) to

influence demographic rates (Benton et al., 2006; Lawson, Vindenes, Bailey, & van de Pol,

2015; Ozgul, Oli, Olson, Blumstein, & Armitage, 2007; Paniw et al., 2019; Töpper et al., 2018).

One major analytical challenge for population ecologists is that typically only a small subset of

potential environmental drivers of population dynamics are known and measured (Ehrlén,

Morris, von Euler, & Dahlgren, 2016; Teller, Adler, Edwards, Hooker, & Ellner, 2016). Another

challenge is that key environmental factors simultaneously affect multiple demographic rates

causing them to covary within and among seasons (Maldonado-Chaparro, Blumstein, Armitage,

& Childs, 2018; Paniw et al., 2019). Positive environment-driven covariation in demographic

rates can then amplify the population-level effects of environmental change (Jongejans, de

Kroon, Tuljapurkar, & Shea, 2010; Robert, Bolton, Jiguet, & Bried, 2015). Antagonistic

demographic responses on the other hand can buffer populations from environmental change

(Knops, Koenig, & Carmen, 2007; Van de Pol et al., 2010); for instance, when population-level

effects of decreased reproduction are offset by increases in survival or growth (Connell &

Ghedini, 2015; Reed, Grøtan, Jenouvrier, Sæther, & Visser, 2013; Villellas, Doak, García, &

Morris, 2015). Thus, explicit consideration of environment-driven patterns in demographic

covariation can allow to gain a fuller picture of population persistence in a changing world. Such

a consideration remains scarce (Ehrlén & Morris, 2015; Ehrlén et al., 2016) but see (Bassar et al.,

2016; Compagnoni et al., 2016).

A flexible framework to model population responses to environment-driven demographic

covariation are integral projection models (IPMs; Ellner, Childs, & Rees, 2016). Seasonal IPMs

are an extension of periodic matrix population models (Caswell, 2001, ch. 13 and references

therein) (Ripley & Caswell, 2006), and simultaneously project trait distributions and population

dynamics across seasons. The construction of IPMs is typically done by fitting simple models

describing demographic rates as functions of a continuous state variable (e.g., body mass) and

(environmental) covariates. Thus, this framework can integrate demographic covariation in

response to environmental drivers (Hindle et al., 2018; Merow et al., 2014; Rees & Ellner, 2009).

IPMs are increasingly used to project population dynamics under environmental change (Paniw

et al., 2019; Paniw, Quintana-Ascencio, Ojeda, & Salguero-Gómez, 2017b; Simmonds &

Coulson, 2015); however, periodic IPMs have been developed in only a few cases (Bassar et al.,

2016; Paniw et al., 2019), despite their potential to provide a mechanistic understanding of

seasonal carry-over effects (Harrison, Blount, Inger, Norris, & Bearhop, 2010; Hostetler, Sillett,

& Marra, 2015).

101   Here, we investigated the population-level effects of environment-driven seasonal

covariation among trait-mediated demographic rates (*i.e.*, collectively referred to as demographic

processes), capitalizing on 40 years (1976-2016) of individual-based data from a population of

yellow-bellied marmots (*Marmota flaviventer*). Yellow-bellied marmots have adapted to a highly

seasonal environment; individuals spend approximately eight months in hibernation during the

cold winter (September/October-April/May), and use the short summer season (April/May-

September/October) to reproduce and replenish fat reserves (Fig. 1). One challenge that the

marmot study shares with numerous other natural systems is the identification of key proximal

environmental factors driving population dynamics. In marmots such factors are numerous and

affect population dynamics through complex, interactive pathways (Maldonado-Chaparro, Read,

& Blumstein, 2017; Oli & Armitage, 2004), which include interactions with phenotypic-trait

structure (Ozgul, Childs, Oli, Armitage, & Blumstein, 2010). As a result, measures of

environmental covariates (e.g., temperature) have previously shown little effect on the

covariation of marmot demographic processes (Maldonado-Chaparro et al., 2018). To address

this challenge, we used a novel method, a hierarchical factor analysis (Hindle et al., 2018), to

model the covariation of demographic processes as a function of a shared measure of latent

environmental quality (*i.e.*, environmental condition at a given time point), quantified in a

Bayesian modeling framework. We then built seasonal stage-, mass-, and environment-specific

IPMs for the marmot population and used prospective stochastic perturbation analyses and

population projections to assess how environmental change may affect population dynamics via

the observed demographic covariation.

METHODS

Study species

Yellow-bellied marmots are large, diurnal, burrow-dwelling rodents, occupying montane regions

of the western North America (Armitage, 2003; Frase & Hoffmann, 1980). They experience

strong seasonal fluctuations in environmental conditions (Armitage, 2014). The biology of

yellow-bellied marmots in Colorado is described in detail by Armitage (2014). Our study was

conducted in the Upper East River Valley near the Rocky Mountain Biological Laboratory,

Gothic, Colorado (38° 57' N, 106° 59' W). Climatic conditions in both winter and summer have

been shown to influence reproduction and survival in the subsequent season (Lenihan & Van

Vuren, 1996; Van Vuren & Armitage, 1991). These climatic effects on the demography of

yellow-bellied marmots are mediated through body mass, with heavier individuals more likely to

survive hibernation and reproduce in summer (Armitage, Downhower, & Svendsen, 1976; Ozgul

et al., 2010). Population dynamics of marmots are therefore likely to be susceptible to changes in

seasonal patterns of environmental drivers. However, numerous interacting climatic factors, such

as temperature extremes, snowfall, and length of snow cover, determine both winter and summer

environmental conditions, and their effect on marmot demography have been shown to be

difficult to disentangle (Inouye, Barr, Armitage, & Inouye, 2000; Schwartz & Armitage, 2002;

Schwartz & Armitage, 2005).

Seasonal demographic rates and trait transitions

For this study, we focused on the population dynamics of eight major colonies continuously

monitored since 1976 (Armitage, 2014; Supporting Material S1). Each year, marmots were live-

trapped throughout the growing season in summer (and ear-tagged upon first capture), and their

sex, age, mass, and reproductive status were recorded (Armitage & Downhower, 1974;

Schwartz, Armitage, & Van Vuren, 1998). We focused on seasonal demographic processes of

females only. All young males disperse from their natal colonies, and female immigration into

existing colonies is extremely rare; as such, local demography can be accurately represented by

the female segment of the population (Armitage, 2010). We classified female marmots by age

and reproductive status: juveniles (< 1 year old), yearlings (1 year old), and non-reproductive (≥

2 years old; not observed pregnant or with offspring) and reproductive adults (≥ 2 years old;

observed pregnant or with offspring) (Armitage & Downhower, 1974).

Figure 1: Seasonal life-cycle transitions modelled for yellow-bellied marmots. The two seasons correspond to the
main periods of mass loss (winter) and gain (summer). Solid and dashed arrows represent discrete-time stage
transitions and recruitment, respectively. Transitions among winter (W) and summer (S) stages (marked by arrows
in different colors) depend on demographic rates (survival [θ], reproduction [φ_0], and recruitment [φ_1]) and trait
transitions (mass change [γ], and offspring mass [φ_2]). Stages are: juveniles, J, yearlings, Y, non-reproductive
adults, N, and reproductive adults, R. All stage-specific demographic rates and trait transitions are modeled using
generalized mixed effects models in a Bayesian framework and include body mass and a common latent variable
representing environmental quality as covariates.

We determined all demographic rates of females (survival, reproduction, and recruitment) for

two discrete growing seasons: winter (August - June) and summer (June - August) (Fig. 1).

Although marmots begin emerging from hibernation in April and some do not hibernate until

October (Armitage, 2014), the two seasons delineate the main periods of mass loss and gain,

respectively (Maldonado-Chaparro et al., 2017). We assumed that females that permanently

disappeared from a colony had died. This measure of apparent survival may overestimate the

death of yearlings in the summer, which disperse from their natal colonies (Van Vuren &

Armitage, 1994). At the same time, the intensive trapping protocol ensured a high capture

probability of yearlings (Oli & Armitage, 2004), decreasing the discrepancies between their

apparent and true survival.

Female marmots give birth to one litter from mid-May to mid-June. In our population

model, females ≥ 2 year of age that survived the winter were considered reproductive adults at

the beginning of summer if they were observed to be pregnant of with pups, or non-reproductive

adults otherwise (Fig. 1). Upon successful reproduction, weaned offspring emerge from burrows

ca. 35 days after birth (Armitage et al., 1976); we therefore defined recruitment as the number of

female juveniles weaned by reproductive females that survive the summer (Fig. 1). The sex ratio

of female:male recruits was assumed to be 1:1 (Armitage & Downhower, 1974). Observations

and pedigree analyses allowed to determine the mother of each new juvenile recruited into the

population (Ozgul et al., 2010).

To assess changes in body mass from one season to the next, we estimated body mass of

every female at the beginning of each season: June 1 (beginning of the summer season when

marmots begin foraging) and August 15 (beginning of the winter season in our models when

foraging activity decreases). Mid-August is the latest that body mass for the vast majority of

individuals can be measured and has been shown to be a good estimate of pre-hibernation mass

(Maldonado-Chaparro et al., 2017). Body-mass estimates on the two specific dates were obtained

using linear mixed effect models as described in Ozgul *et al.* (2010; see also Maldonado-

Chaparro et al., 2017). Body mass of juvenile females was estimated for August 15.

Modelling demographic and mass change (trait-transition) rates

We jointly modeled all seasonal demographic and mass change rates (*i.e.*, demographic

processes) as a function of stage and body mass - or mother's mass in the case of juvenile mass -

at the beginning of a season, using a Bayesian modeling framework (Table 1; Supporting

Material S1). All mass estimates were cube-root transformed to stabilize the variance and

improve the normality of the residuals in the Gaussian submodels (Maldonado-Chaparro et al.,

2017). We fitted all demographic-process submodels as generalized linear mixed effects models

(GLMMs). We assumed a binomial error distribution (logit link function) for the probability of

winter (θ_W) and summer (θ_S) survival and of probability of reproducing (*i.e.*, being in the

reproductive adult stage at the beginning of summer; φ_0); a Poisson error distribution (log link

function) for the number of recruits (φ_1); and a Gaussian error distribution (identity link) for the

masses (z^*) at the end of each season (Table 1). Mass-change (*i.e.*, mass gain or loss) rates (γ)

were then defined as functions of current (z) and next (z^*) mass using a normal probability

density function. For the juvenile mass distribution (φ_2), the density function depended on the

mother's mass (z_M) (see below; Supporting Material S2).

To describe environment-driven temporal covariation in seasonal demography in the

absence of explicit knowledge on environmental drivers, we quantified a model-wide latent

variable (Q_y) affecting all demographic processes in a given year (y) (Table 1). One latent

variable was enough to capture covariation among the demographic processes (Supporting

Material S1). Details of the parameterization of Q_y can be found in Supporting Material S1 (see

also Hindle *et al*., 2018). Briefly, this covariate effectively estimates the overall annual

environmental quality that affects the various demographic processes simultaneously, thereby

capturing correlated responses (within and among seasons) to environmental fluctuations (Hindle

et al., 2018). In the absence of causal variables that can be directly measured, Q_y was

incorporated as a covariate in all seven demographic-process submodels; and the year-specific

values of Q_y were drawn from a normal distribution with mean = 0 and SD =1. Higher Q_y

indicate a better quality of the environment in a given year, and the associated β_q slope

parameters determine the magnitude and sign of the effect of Q_y on a given, season-specific

demographic process (Table 1). Thus, similar to large-scale environmental indices such as NAO

or ENSO, Q_y captures to what extent a bad ($Q_y < 0$) or good ($Q_y > 0$) environment is likely to

affect both summer and winter demography. To make the Bayesian model identifiable, we

constrained the standard deviation of Q_y to equal 1 and arbitrarily set the β_q for summer survival

(θ_S) to be positive. The β_q of the remaining submodels can, therefore, be interpreted as

correlations of demographic processes with θ_S due to environmental variation.

Aside from the environmental covariate Q_y simultaneously affecting all demographic

processes, we included a random year effect ($\varepsilon_{Ysubmodel}$) as a covariate in each submodel. This

year effect accounted for additional temporal variation not captured by Q_y. We also tested for the

effect of population density (measured as total abundance, abundance of adults, or abundance of

yearling and adults) in all submodels. However, like previous studies, we could not detect any

clear density effects (Armitage, 1984; Maldonado-Chaparro et al., 2018).

The prior distributions of the Bayesian model and posterior parameter samples obtained

are detailed in Supporting Material S1. For each demographic-process submodel, we chose the

most parsimonious model structure by fitting a full model that included all covariates (mass,

stage, and Q_y) and two-way interactions between mass and stage and stage and Q_y, and retaining

only those parameters for which the posterior distribution (± 95 % C.I.) did not overlap 0 (Table

1; Table S2.1).

Table 1: Parameterization of the most parsimonious models describing winter (W) and summer (S)

demographic processes in marmots. The distributions B, N, and P correspond to the Bernoulli, normal,

and Poisson distributions, respectively. The logit link, for winter survival as an example, corresponds to 1-

$^{(1/1+\exp(-\theta W))}$. *Stage* – life cycle stage. Q – latent environmental variable. z – season-specific mass. z_M –mass

of the mother.

Demographic process	Function	Likelihood distribution
Winter (W):		
Survival (θ_W)	$\text{logit}(\theta_W) = \alpha_{0\theta W} + \alpha_{a\theta W}[\text{stage}] + \beta_{z\theta W} \times z$ $+ \beta_{q\theta W} \times Q_y[\text{year}] + \varepsilon_{y\theta W}[\text{year}]$	$B(\theta_W)$
Mass next (z_W^*)	$z_W^* = \alpha_{0z*W} + \alpha_{az*W}[\text{stage}] + (\beta_{zz*W} +$ $\beta_{zaz*W}[\text{stage}]) \times z$ $+ \beta_{qz*W} \times Q_y[\text{year}] + \varepsilon_{yz*W}[\text{year}]$	$N(z_W^*, \tau_{z*W})$
Reproduction (φ_0)	$\text{logit}(\varphi_0) = \alpha_{0\varphi0} + \alpha_{a\varphi0}[\text{stage}] + \beta_{z\varphi0} \times z$ $+ \beta_{q\varphi0} \times Q_y[\text{year}] + \varepsilon_{y\varphi0}[\text{year}]$	$B(\varphi_0)$
Summer (S):		
Survival (θ_S)	$\text{logit}(\theta_S) = \alpha_{0\theta S} + \alpha_{a\theta S}[\text{stage}] + \beta_{z\theta S} \times z$ $+ \beta_{q\theta S} \times Q_y[\text{year}] + \varepsilon_{y\theta S}[\text{year}]$	$B(\theta_S)$
Mass next (z_S^*)	$z_S^* = \alpha_{0z*S} + \alpha_{az*S}[\text{stage}] + (\beta_{zz*S} +$ $\beta_{zaz*S}[\text{stage}]) \times z + \beta_{qz*S} \times Q_y[\text{year}]$ $+ \varepsilon_{yz*S}[\text{year}]$	$N(z_S^*, \tau_{z*S})$
Number of recruits (φ_1)	$\log(\varphi_1) = \alpha_{0\varphi1} + \beta_{z\varphi1} \times z$ $+ \beta_{q\varphi1} \times Q_y[\text{year}] + \varepsilon_{y\varphi1}[\text{year}]$	$P(\varphi_1)$
Juvenile mass (z_J^*)	$z_J^* = \alpha_{0z*J} + \beta_{zz*J} \times z_M + \beta_{qz*J} \times Q_y[\text{year}]$ $+ \varepsilon_{yz*J}[\text{year}]$	$N(z_J^*, \tau_{z*J})$

Seasonal Integral Projection Models

We used the most parsimonious models of demographic processes (Table 1) to parameterize

density-independent, stage-mass-structured, seasonal and environment-specific Integral

Projection Models (IPMs) (Easterling, Ellner, & Dixon, 2000; Ellner et al., 2016). For each stage

a, the IPMs track the number of individuals (n_a) in the mass range [z, z+dz] at time *t*. The fate of

these individuals at time *t*+1 is described by a set of coupled integral equations, which differ for

each season and are a function of the latent environmental variable Q_y. In the winter season,

individuals can survive (θ_w) and change mass (γ_w) according to their stage, mass, and

environment. Conditional on survival, juveniles (J) transition to yearlings (Y), while all other

stages are distributed to either the reproductive (R) or non-reproductive (N) adult stage at the

beginning of summer, depending on the stage-specific probability of reproducing (φ_0). During

the summer season, individuals in stages Y, N, and R survive (θ_S) and change mass (γ_S)

according to their stage and mass at the beginning of summer and according to the environment;

but, in summer, transitions to another stage do not occur. Reproductive individuals (R) of a given

mass also produce φ_1/2 female juveniles (J), *i.e.*, half of the total number of recruits. Female

recruits are distributed across *z* mass classes by the end of summer, given by φ_2. The

mathematical descriptions of the IPMs for the winter and summer seasons are provided in

Supporting Material S2.

We numerically integrated the summer and winter IPMs using the 'midpoint rule'

(Easterling et al., 2000) with upper and lower integration limits of 7.8 (472 g) and 17.1 (5000 g),

respectively. To avoid unintended eviction of individuals from the model (*i.e.*, for a given mass

class *z*, the sum of the probabilities to transition to $z^* < 1$), we applied a constant correction (*i.e.*,

equally redistributing evicted individuals among all z^*) when constructing the IPMs as suggested

in Merow *et al.*, (2014) (see also Williams, Miller, & Ellner, 2012). For each stage-specific IPM,

we chose a bin size of 100 (*i.e.*, dividing masses into 100 classes), as further increasing the bin

size did not significantly improve the precision of estimates of the long-term population growth

rate. The IPMs we constructed accurately reproduced observed population dynamics from 1976-

2016 (Supporting Material S2).

Sensitivity of population dynamics to seasonal demographic processes: prospective

perturbations

Changes in population dynamics in response to changes in environmental fluctuations are

determined by the response of demographic processes to the environment and, in turn, of

population dynamics to demographic processes (Maldonado-Chaparro et al., 2018). To explore

these two sources of variation in the long-term fitness of the marmot population, we first

quantified the proportional change in the demographic processes (Table 1) to changes in Q_y, *i.e.*,

$\partial(\log \rho)/\partial Q_y$, where ρ is a demographic process. We calculated these elasticities for different

values of Q_y (from -1 to 1), increasing each value by 0.01 and keeping mass at its stage-specific

average and ε_Y fixed to the value estimated for 2015. To assess the effect of parameter

uncertainty on our estimates, we repeated these calculations for a sample of 1000 parameter

values drawn from the posterior distribution (Paniw, Quintana-Ascencio, Ojeda, & Salguero-

Gómez, 2017a).

We next assessed which demographic processes most affected the stochastic population

fitness under observed (1976-2016) environmental fluctuations. We used a simulation of 100,000

years to assess asymptotic, long-term stochastic population dynamics (see section below for

short-term viability simulations). Each simulation started with the population of individuals with

the standing stage and mass distribution at the beginning of the winter season in 2016. At each

run of the simulation, a seasonal IPM was constructed with resampled, year-specific Q_y and ε_Y

values. We then calculated the periodic IPM product of the winter and summer IPMs, describing

the demographic and trait transitions from the beginning of one winter season to the next

(Caswell, 2001; chapter 13). Lastly, we obtained the stochastic population growth rate, logλs, (a

measure of fitness) as the mean of the logs of annual realized population growth rates, $\lambda = n_{t+1}/n_t$,

where n_t and n_{t+1} are the total number of individuals in the population at time (year) t and $t+1$,

respectively.

During the simulations, we adapted the approach described in Ellner *et al.* (2016; chapter

7) to calculate the elasticity of log λs to changes in the mean (e_S^{μ}) and standard deviation (e_S^{σ}) of

stage-specific demographic processes ρ, e.g., juvenile survival. To do so, we perturbed a

demographic-process function for a given stage over an interval of masses [z, $z+dz$] by its mean

and standard deviation obtained across 40 years of observed dynamics (Supporting Material S3).

Observed dynamics were obtained by resampling year-specific values of Q_y and ε_Y for 1976-

2016 from the Bayesian model. We then built a winter or summer IPM using the perturbed vital

rate and obtained a new, perturbed periodic IPM product (*i.e.*, the perturbation kernel). Lastly,

we integrated this product into eq. 7.5.2 in Ellner *et al.* (2016) in order to estimate e_S^{μ} and

e_S^{σ} from the simulations of population dynamics for T $= 90,000$ years (*i.e.*, discarding the first 10,

000 years of simulations to ensure no effect of transient population fluctuations):

$$e = E\left[\frac{\langle v_{t+1}, C_t w_t \rangle}{\langle v_{t+1}, K_t w_t \rangle}\right]$$

,

where C_t is the perturbation kernel, K_t is the unperturbed annual IPM kernel at each simulation

iteration t, v and w are the left and right eigenvectors associated with K at $t+1$ and t, respectively.

For instance, if average winter survival in a given mass interval (*i.e.*, IPM bin) for reproductive

individuals from 1976-2016 was 0.95, each winter IPM during T simulations was perturbed by

subtracting a focal value of winter survival from its 40-year mean; and e_S^{μ} was calculated by

integrating this perturbed IPM as C_t into the above equation. The two elasticities quantify the

strength of selection pressures on lower-level vital rates in stochastic environments (Haridas &

Tuljapurkar, 2005; Rees & Ellner, 2009). We repeated the elasticity calculations for a sample of

100 parameter values from the posterior distribution.

Population viability under changes in environmental quality

To assess how the combined effects of (i) seasonal demographic responses to environmental

fluctuations and (ii) population sensitivity to seasonal demography impact population viability,

we simulated population dynamics under environmental change. We ran 200 independent

simulations each projecting population dynamics for 50 years. The projections were based on

several scenarios of changes in the distribution of environmental quality, Q_y. We first created

base simulations (*i.e.*, no environmental change) where Q_y was picked from a normal distribution

with $\mu_Q = 0$ and $\sigma_Q = 1$ across all demographic processes. Next, we approximated random future

fluctuations in Q_y under different average environmental conditions. To do so, we sampled Q_y

from a normal distribution fixing the average environmental quality ($\mu_Q = -1, -0.5, 0.5, 1$) and its

variation ($\sigma_Q = 0.6, 1.2$) over the 50 years of projections. We then explored how a trend in μ_Q

would affect viability and mass distribution. To do so, we decreased the four μ_Q by 0.01 in each

year of the projections, keeping σ_Q unaltered. Lastly, as temporal autocorrelation in

environmental states can affect population dynamics (Paniw et al., 2018), we projected

autocorrelated changes in Q_y. That is, each year, marmots could experience either a good or bad

environment where Q_y was sampled from a normal distribution with either $\mu_Q = 1$ or $\mu_Q = -1$,

respectively (σ_Q was fixed at 0.6). The sequence of good and bad environments was determined

by a Markov chain with various autocorrelation coefficients (see Supporting Material S4 for

details). All simulations were repeated for a random sample of 1,000 parameters from the

posterior distribution to account for parameter uncertainty.

For all environmental-change scenarios, we recorded the probability of quasi-extinction

across the 200 simulations. Quasi-extinction was defined conservatively as the number of non-

juvenile individuals (*i.e.*, yearlings and non-reproductive and reproductive individuals) in the

population to be < 4, which corresponded to 10 % of their lowest observed number.

RESULTS

Seasonal environment-driven covariation in demographic processes

Our GLMMs showed a strong effect of environmental quality (measured by the latent variable

Q_y) on winter but not summer demographic processes. This effect was positive for all winter

demographic processes, as evidenced by the positive β_q (Fig. S1.3 in Supporting Material S1).

The β_q for demographic processes in the summer, however, were comparatively small and were

not significantly different from 0 (95 % posterior C.I.s overlapped 0). This environment-driven

positive covariation in demographic processes therefore indicated that Q_y provided a measure of

environmental quality (Hindle et al., 2018) that did not correlate with any single environmental

variable measured at the study site and correlated well with observed annual changes in

population size (Supporting Material S1).

In accordance with the posterior distribution of β_q parameters, only winter demographic

processes were significantly affected by small changes in Q_y (Fig. 2). Among the winter

demographic processes, changes in Q_y affected reproduction across stages the most, followed by

survival of juveniles (Fig. 2).

Figure 2: The sensitivity of seasonal demographic processes to environmental quality in marmots. Sensitivity is assessed as proportional changes in demographic processes, ρ, as environmental quality, Q_y, increases slightly. This sensitivity is measured with respect to different average values of Q_y and across four different life-cycle stages: juveniles (J), yearlings (Y), non-reproductive adults (N), and reproductive adults (R). The demographic processes include winter (W; blue color tones) and summer (S; red color tones) survival (θ) and mass change (γ); and probability of reproducing (φ_0), recruitment (φ_1), and juvenile mass (φ_2). Points and error bars show averages ± 95 % C.I. across 1,000 posterior parameter samples obtained from the Bayesian population model.

Aside from the effects of environmental quality, our models confirmed previous findings on the importance of body mass and stage on yellow-bellied marmot demography (Maldonado-Chaparro et al., 2017; Ozgul et al., 2010; Ozgul, Oli, & Armitage, 2009). The most parsimonious GLMMs (Table 1; Table S1.1 in Supporting Material S1) showed a positive effect of mass on all demographic processes, with the weakest effect of mass on summer survival (θ_S) of reproductive

adults. Survival, in particular θ_S, was highest for reproductive adults; reproduction was also

highest for adults that reproduced before (Fig. S1.3 in Supporting Material S1).

Sensitivity of population dynamics to seasonal demographic processes

While environmental quality affected winter demographic processes only, our prospective

perturbation analyses showed that winter and summer demography equally determine long-term

population fitness. Stochastic elasticity analyses (e_S^{μ} and e_S^{σ}) showed that relative increases in the

mean (μ) of winter (θ_W) and summer (θ_S) survival for reproductive adults (R), would lead to

substantial relative increases of the stochastic population growth rate, $\log\lambda s$ (Fig. 3a). Highest e_S^{μ}

were found at intermediate and large mass classes; e_S^{μ} was negative for small masses when mass

changes (γ) and offspring mass (φ2) were perturbed (Fig. S3.1a in Supporting Material S3),

explaining the overall small e_S^{μ} for γ and φ2 summed over all mass classes (Fig. 3a). Overall,

relative changes in $\log\lambda s$ due to increases in the standard deviation of demographic processes

(e_S^{σ}) were much smaller compared to e_S^{μ} (Fig. 3b) and didn't differ significantly between vital

rates, as 95 % posterior C.I. crossed 0 (Fig. S3.1b).

Figure 3 Sensitivity of the average long-term population fitness to changes in the average and variability of

demographic processes modeled for the yellow-bellied marmots. The sensitivity measure is obtained analytically as

elasticities (e) of the stochastic population growth rate, log λ_s, to changes in (a) the mean (μ) and (b) standard

deviation (σ) of stage-specific demographic processes summed over all mass classes. Stages are juveniles (J),

yearlings (Y), non-reproductive adults (N), and reproductive adults (R). Demographic processes include winter (W)

and summer (S) survival (θ) and mass change (γ); reproduction (φ_0); recruitment (φ_1), and offspring mass

distribution (φ_2). Elasticities were calculated at the mean posterior values of parameters obtained from the Bayesian

demographic model.

Population viability under changes in environmental quality

While population fitness was equally sensitive to demographic processes over winter and

summer, environmental fluctuations strongly affected viability through winter demography.

Using base simulations (*i.e.*, obtaining Q_y from a normal distribution with $\mu_Q = 0$ and $\sigma_Q = 1$),

the probability of quasi-extinction, at an average of 0.1 [0.0, 0.3 C.I.] across posterior

parameters, were relatively low. Simulations of population dynamics based on scenarios of

environmental change resulted in substantial decreases (0 at $\mu_Q = 1$) and increases (0.9 [0.6, 1.0

C.I.] at $\mu_Q = -1$) in quasi-extinction, compared to base simulations, when the population

experienced a high and low average environmental quality (Q_y), respectively (Fig. 4). Average

quasi-extinction further increased and its uncertainty across posterior parameters decreased when

a declining trend in average environmental quality (Q_y) was simulated (Fig. S4.1). Changes in

the standard deviation of Q_y had comparatively little effect on quasi-extinction (Fig. 4). The

same was true for simulations of autocorrelated environmental fluctuations, where changes in the

frequency of good environmental conditions (effectively changing the average environmental

quality) had a strong effect on viability, while changes in the patterning of environmental states

had little effect (Supporting Material S4).

Figure 4: Probability of quasi-extinction (*i.e.*, < 4 non-juveniles in the population) of yellow-bellied marmots under different scenarios of environmental change. The scenarios consisted of projecting population dynamics for 50 years fixing a different mean (μ) and standard deviation (σ) of environmental quality (Q) in all demographic processes. Points and error bars show averages ± 95 % C.I. across 1,000 posterior parameter samples obtained from the Bayesian population model. Base simulations ($\mu_Q = 0$; $\sigma_Q = 1$) are depicted in red.

DISCUSSION

One important pathway through which environmental change can act on population dynamics is through seasonal direct and carry-over effects on survival, growth, and reproduction (Harrison et al., 2010; Paniw et al., 2019). Using a novel, factor-analytic approach to quantify environment-demography relationships, we show that positive responses in several demographic processes to winter environmental conditions can drive annual population dynamics in a winter-adapted mammal. This sensitivity to winter conditions occurs despite the fact that offspring are recruited in summer and both summer and winter demographic processes determine population fitness. As

whole-year, population-level effects of environmental change can be filtered by seasonal

processes in the absence of density-dependent feedbacks, we highlight the need to assess such

seasonal processes to gain a mechanistic understanding of population persistence (Paniw et al.,

2019; Picó, de Kroon, & Retana, 2002).

Empirical studies have typically used annual models to demonstrate how positive

demographic covariation can amplify environmental effects on population dynamics

(Compagnoni et al., 2016); or how opposing demographic responses can buffer such effects

(Gamelon et al., 2017; Reed et al., 2013). However, in marmots, as in numerous other

populations (Bassar et al., 2016; Jenouvrier et al., 2018), seasonal demographic processes play an

important role in life-cycle dynamics (Armitage, 2017). Our prospective perturbations show that

changes in both mean winter and summer survival of reproductive adults have the strongest

effect on population fitness, confirming the critical role of this life-cycle stage (Maldonado-

Chaparro et al., 2018; Ozgul et al., 2009). At the same time, environmental fluctuations do not

affect adult survival or other demographic processes throughout the year. That is, although the

environment has been shown to drive particularly recruitment in numerous temperate species

(e.g., Bonardi, Corlatti, Bragalanti, & Pedrotti, 2017; Nouvellet, Newman, Buesching, &

Macdonald, 2013), such effects are not evident in marmots; here, environmental quality shows

little impact on summer demography, including recruitment, and little covariation with winter

demography. Instead, only the joint responses of demographic processes in winter are affected

by environmental quality. The joint effects of the sensitivity of winter demography to

environmental quality and the sensitivity of population fitness to winter demography then

determine population persistence under environmental change.

The seasonal effects of environmental quality on population persistence must be

understood in terms of the role of reproductive females in the marmot population (Ozgul et al.,

2009). In our simulations, a good winter quality, which has previously been shown to have the

strongest effect on individual survival (Armitage, Blumstein, & Woods, 2003), would, in

particular, result in more reproductive females in the summer. In turn, summer survival and

reproduction of these females are important to long- and short-term demography (Maldonado-

Chaparro et al., 2018; Ozgul et al., 2009), but are not driven by environmental conditions. As is

the case in other socially complex mammals (Morris et al., 2011), reproduction in yellow-bellied

marmots is governed primarily by social interactions, in particular the behavior of dominant

females (Armitage, 2010; Blumstein & Armitage, 1998). Even under optimal summer

conditions, the reproductive output of the population may not be increased as dominant females

suppress reproduction in younger subordinates and therefore regulate the size of colonies

(Armitage, 1991). Dominant females, in addition, may skip reproduction themselves if they enter

hibernation with a relatively low mass (Armitage, 2017). Thus, the necessity of meeting the

physiological requirements of hibernation profoundly affects life-history traits of yellow-bellied

marmots that are expressed during the active season. Here, an explicit seasonal population

analysis allowed us to recognize a potentially important mechanism through which the

environment filters demography: survival in or transition to the reproductive stage in a good

winter can guarantee successful reproduction in summer, in addition to direct summer processes.

Unlike the direct effects of seasonal survival and reproduction, trait transitions between

seasons had a smaller effect on annual population dynamics, even if winter mass changes were

mediated by environmental quality. In this sense, the prospective perturbation analyses

performed in our study are in agreement with retrospective analyses described by Maldonado-

Chaparro and colleagues (2018), who found a relatively small contribution of growth to past

variation in annual population growth. These relatively small effects are likely due to the fact

that marmots compensate for winter mass loss with increased growth in the summer, creating a

zero net effect on annual trait change (Maldonado-Chaparro et al., 2017). Although the strength

of compensatory effects may differ within seasons or among life-history stages (Monclús, Pang,

& Blumstein, 2014), such effects are common in rodents and other species that have a short

window for mass gain (Morgan & Metcalfe, 2001; Orizaola, Dahl, & Laurila, 2014), and

highlight the advantages of assessing seasonal dynamics when studying population-level global-

change effects (Bassar et al., 2016).

Under environmental change, the persistence of marmots was mostly affected by changes

in mean environmental quality, whereas changes in the variance and temporal patterning of the

mean showed little effects. This supports previous conclusions that yellow-bellied marmots are

partly buffered against increases in environmental variation (Maldonado-Chaparro et al., 2018;

Morris et al., 2008) or autocorrelation (Engen et al., 2013). Further support for demographic

buffering comes from the fact that changes in the mean environmental quality most strongly

affected those demographic processes to which the stochastic population growth rate was least

sensitive. That is, yearlings gaining reproductive status in particular, albeit contributing relatively

little to population dynamics, contributed strongly via environment-driven effects. It is well

known that in species where vital rates of adults are relatively buffered, juveniles are much more

sensitive to environmental variation (Gaillard & Yoccoz, 2003; Jenouvrier et al., 2018).

Although demographic buffering has been investigated in other animals (Bjørkvoll et al., 2016;

Morris et al., 2008; Pfister, 1998; Rotella, Link, Chambert, Stauffer, & Garrott, 2012), we are, to

the best of our knowledge, the first to show that it is likely to persist across the seasonal

environments and different masses for a high-altitude specialist; see Jenouvrier *et al*., (2018) for

similar analyses in a long-lived bird.

Our results emphasize that declines in environmental quality in one season alone can

strongly affect annual population dynamics of a mammal highly adapted to seasonal

environments and buffered from environmental variation. Therefore, positive demographic

covariation under environmental change may threaten populations even if it affects demographic

process to which the stochastic growth rate is least sensitive, *i.e.*, processes that are under low

selection pressure (Coulson, Gaillard, & Festa-Bianchet, 2005; Iles, Rockwell, & Koons, 2019).

Studies of global change ecology that focus on the effects of environmental factors on

demographic processes considered to strongly affect both short- and long-term population

dynamics may therefore miss the important role of seasonal demographic covariation. In

addition, such seasonal mechanisms are potentially critical for conservation management (Paniw

et al., 2019) but would be missed by annual-step analyses where demographic processes are

averaged across the entire year (Hostetler et al., 2015).

Given the strong overlap of our results with previous findings and overall robust model

performance, the factor analytic approach we used to quantify environment-driven covariation in

demographic processes appears to be a promising alternative to approaches linking specific vital

rates directly to environmental drivers (Evans & Holsinger, 2012; Hindle, Pilkington,

Pemberton, & Childs, 2019; Hindle et al., 2018). In the case of yellow-bellied marmots, where

demographic rates were relatively easy to parameterize and the assumption of positive

covariation holds in part due to a lack of density feedbacks, the factor analytic approach can be

used effectively to assess population-level effects of perturbing a global measure of

environmental quality. In cases where environment-demography relationships are more complex

and include negative demographic covariation (e.g., where trade-offs dominate), this approach

may find limited applications. However, positive covariation of demographic patterns is common

(Jongejans et al., 2010; Paniw et al., 2019); and, given the short time series of most demographic

datasets (Salguero-Gómez et al., 2015; Salguero-Gómez et al., 2016) or little knowledge on the

actual environmental drivers of population dynamics (Pol et al., 2016; Teller et al., 2016), the

factor analytic approach can be particularly useful in comparative studies.

Most species inhabit seasonal environments. Under global environmental change, it may

therefore be critical to understand how seasonality patterns mediate persistence of natural

populations. Our analyses point to an important role of seasonal differences in demographic

covariance in driving population responses to environmental change. Novel methods such as the

factor analytic approach allow researchers to overcome some challenges associated with such

mechanistic approaches, and we encourage more seasonal demographic analyses across different

taxa.

ACKNOWLEDGMENTS

We thank the many volunteers and researchers of Rocky Mountain Biological Laboratory for

collecting and providing us with the data on individual life histories of yellow-bellied marmots.

M.P. was supported by an ERC Starting Grant (33785) and a Swiss National Science Foundation

Grant (31003A_182286) to A.O.; and by a Spanish Ministry of Economy and Competitiveness

Juan de la Cierva-Formación grant FJCI-2017-32893; D.T.B. was supported by the National

Geographic Society (grant # 8140-06), the UCLA Faculty Senate and Division of Life Sciences,

a RMBL research fellowship, the National Science Foundation (IDBR-0754247, DEB-1119660

and 1557130 to D.T.B.; DBI 0242960, 0731346, and 1226713 to the RMBL).

AUTHOR CONTRIBUTIONS

K.B.A., D.T.B., and M.O. led the long-term study and data collection; M.P., D.C., and A.O.

conceived the ideas for the paper and designed the analyses; M.P. conducted the analyses with

substantial help from J.M.; M.P. wrote the manuscript; all authors discussed the results and

commented on the manuscript.

DATA AVAILABILITY

The datasets generated and analyzed during the current study as well as the R scripts to run all

analyses are available in the *Dryad* repository, [PERSISTENT WEB LINK TO DATASETS].

REFERENCES

Armitage, K. B. (1984). Recruitment in yellow-bellied marmot populations: kinship, philopatry,

and individual variability. In J. O. Murie & G. R. Michener (Eds.), *The Biology of Ground-*

Dwelling Squirrels (pp. 377–403), Lincoln: Univ. Nebraska Press.

Armitage, K. B. (1991). Social and population dynamics of yellow-bellied marmots: results from

long-term research. *Annual Review of Ecology and Systematics*, 22, 379–407.

Armitage, K. B. (2003). Observations on plant choice by foraging yellow-bellied marmots.

Oecologia Montana, 12, 25–28.

Armitage, K. B. (2010). Individual fitness, social behavior, and population dynamics of yellow-

bellied marmots. I. Billick & M.V. Price (Eds.) *The Ecology of Place: Contributions of*

Place-Based Research to Ecological Understanding (pp. 132–154), Chicago: University of

Chicago Press.

Armitage, K. B. (2014). *Marmot Biology: Sociality, Individual Fitness, and Population*

Dynamics. Cmbridge: Cambridge University Press.

Armitage, K. B. (2017). Hibernation as a major determinant of life-history traits in marmots.

Journal of Mammalogy, 98, 321–331.

Armitage, K. B., Blumstein, D. T., & Woods, B. C. (2003). Energetics of hibernating yellow-

bellied marmots (*Marmota flaviventris*). *Comparative Biochemistry and Physiology. Part A,*

Molecular & Integrative Physiology, 134, 101–114.

Armitage, K. B., & Downhower, J. F. (1974). Demography of yellow-bellied marmot

populations. *Ecology*, 55, 1233–1245.

Armitage, K. B., Downhower, J. F., & Svendsen, G. E. (1976). Seasonal changes in weights of

marmots. *The American Midland Naturalist*, 96, 36–51.

Bassar, R. D., Letcher, B. H., Nislow, K. H., & Whiteley, A. R. (2016). Changes in seasonal

climate outpace compensatory density-dependence in eastern brook trout. *Global Change*

Biology, 22, 577–593.

Benton, T. G., Plaistow, S. J., & Coulson, T. N. (2006). Complex population dynamics and

complex causation: devils, details and demography. *Proceedings. Biological Sciences*, 273,

1173–1181.

Bjørkvoll, E., Lee, A. M., Grøtan, V., Saether, B.-E., Stien, A., Engen, S., Albon, S., Loe, L.E.,

& Hansen, B. B. (2016). Demographic buffering of life histories? Implications of the choice

of measurement scale. *Ecology*, 97, 40–47.

Blumstein, D. T., & Armitage, K. B. (1998). Life history consequences of social complexity a

comparative study of ground-dwelling sciurids. *Behavioral Ecology*, 9, 8-19.

Bonardi, A., Corlatti, L., Bragalanti, N., & Pedrotti, L. (2017). The role of weather and density

dependence on population dynamics of Alpine-dwelling red deer. *Integrative Zoology*, 12,

61-76.

Campos, F. A., Morris, W. F., Alberts, S. C., Altmann, J., Brockman, D. K., Cords, M., Pusey,

A., Stoinski, T. S., Strier, K. B., & Fedigan, L. M. (2017). Does climate variability influence

the demography of wild primates? Evidence from long-term life-history data in seven

species. *Global Change Biology*, 23, 4907–4921.

Caswell, H. (2001). *Matrix Population Models*. Sunderland: Sinauer.

Childs, D. Z., Coulson, T. N., Pemberton, J. M., Clutton-Brock, T. H., & Rees, M. (2011).

Predicting trait values and measuring selection in complex life histories: reproductive

allocation decisions in Soay sheep. *Ecology Letters*, 14, 985–992.

Compagnoni, A., Bibian, A. J., Ochocki, B. M., Rogers, H. S., Schultz, E. L., Sneck, M. E.,

Elderd, B. D., Iler, A. M., Inouye, D. W., Jacquemin, H., & Miller, T. E. X. (2016). The

effect of demographic correlations on the stochastic population dynamics of perennial

plants. *Ecological Monographs*, 86, 480–494.

Connell, S. D., & Ghedini, G. (2015). Resisting regime-shifts: the stabilising effect of

compensatory processes. *Trends in Ecology & Evolution*, 30, 513–515.

Coulson, T., Gaillard, J.-M., & Festa-Bianchet, M. (2005). Decomposing the variation in

population growth into contributions from multiple demographic rates. *The Journal of*

Animal Ecology, 74, 789–801.

Easterling, M. R., Ellner, S. P., & Dixon, P. M. (2000). Size-specific sensitivity: applying a new

structured population model. *Ecology*, 81, 694–708.

Ehrlén, J., & Morris, W. F. (2015). Predicting changes in the distribution and abundance of

species under environmental change. *Ecology Letters*, 18, 303–314.

Ehrlén, J., Morris, W. F., von Euler, T., & Dahlgren, J. P. (2016). Advancing environmentally

explicit structured population models of plants. *The Journal of Ecology*, 104, 292–305.

Ellner, S. P., Childs, D. Z., & Rees, M. (2016). *Data-driven Modelling of Structured*

Populations: A Practical Guide to the Integral Projection Model. Springer.

Engen, S., Sæther, B.-E., Armitage, K. B., Blumstein, D. T., Clutton-Brock, T. H., Dobson, F. S.,

Festa-Bianchet, M., Oli, M. K., & Ozgul, A. (2013). Estimating the effect of temporally

autocorrelated environments on the demography of density-independent age-structured

populations. *Methods in Ecology and Evolution*, 4, 573–584.

Evans, M. E. K., & Holsinger, K. E. (2012). Estimating covariation between vital rates: a

simulation study of connected vs. separate generalized linear mixed models (GLMMs).

Theoretical Population Biology, 82, 299–306.

Frase, B. A., & Hoffmann, R. S. (1980). Marmota flaviventris. *Mammalian Species*,135, 1–8.

Gaillard, J.-M., & Yoccoz, N. G. (2003). Temporal variation in survival of mammals: A case of

environmental canalization? *Ecology*, 84, 3294–3306.

Gamelon, M., Grøtan, V., Nilsson, A. L. K., Engen, S., Hurrell, J. W., Jerstad, K., Phillips, A. S.,

Røstad, O. W., Slagsvold, T., Walseng, B., Steneth, N. C., & Sæther, B.-E. (2017).

Interactions between demography and environmental effects are important determinants of

population dynamics. *Science Advances*, 3, e1602298.

Haridas, C. V., & Tuljapurkar, S. (2005). Elasticities in variable environments: properties and

implications. *The American Naturalist*, 166, 481–495.

Harrison, X. A., Blount, J. D., Inger, R., Norris, R. D., & Bearhop, S. (2010). Carry-over effects

as drivers of fitness differences in animals. *The Journal of Animal Ecology*, 80, 4–18.

Hindle, B. J., Pilkington, J. G., Pemberton, J. M., & Childs, D. Z. (2019). Cumulative weather

effects can impact across the whole life cycle. *Global Change Biology*, in press.

Hindle, B. J., Rees, M., Sheppard, A. W., Quintana-Ascencio, P. F., Menges, E. S., & Childs, D.

Z. (2018). Exploring population responses to environmental change when there is never

enough data: a factor analytic approach. *Methods in Ecology and Evolution*, 147, 115.

Hostetler, J. A., Sillett, T. S., & Marra, P. P. (2015). Full-annual-cycle population models for

migratory birds. *The Auk*, 132, 433–449.

Iles, D. T., Rockwell, R. F., & Koons, D. N. (2019). Shifting vital rate correlations alter

predicted population responses to increasingly variable environments. *The American*

Naturalist, E57–E64.

Inouye, D. W., Barr, B., Armitage, K. B., & Inouye, B. D. (2000). Climate change is affecting

altitudinal migrants and hibernating species. *Proceedings of the National Academy of*

Sciences of the United States of America, 97, 1630–1633.

Jenouvrier, S., Desprez, M., Fay, R., Barbraud, C., Weimerskirch, H., Delord, K., & Caswell, H.

(2018). Climate change and functional traits affect population dynamics of a long-lived

seabird. *The Journal of Animal Ecology*, 87, 906–920.

Jenouvrier, S., Holland, M., Stroeve, J., Barbraud, C., Weimerskirch, H., Serreze, M., & Caswell,

H. (2012). Effects of climate change on an emperor penguin population: analysis of coupled

demographic and climate models. *Global Change Biology*, 18, 2756–2770.

Jongejans, E., de Kroon, H., Tuljapurkar, S., & Shea, K. (2010). Plant populations track rather

than buffer climate fluctuations. *Ecology Letters*, 13, 736–743.

Knops, J. M. H., Koenig, W. D., & Carmen, W. J. (2007). Negative correlation does not imply a

tradeoff between growth and reproduction in California oaks. *Proceedings of the National*

Academy of Sciences of the United States of America, 104, 16982–16985.

Lawson, C. R., Vindenes, Y., Bailey, L., & van de Pol, M. (2015). Environmental variation and

population responses to global change. *Ecology Letters*, 18, 724–736.

Lenihan, C., & Vuren, D. V. (1996). Growth and survival of juvenile yellow-bellied marmots

(*Marmota flaviventris*). *Canadian Journal of Zoology*, 74, 297–302.

Maldonado-Chaparro, A. A., Blumstein, D. T., Armitage, K. B., & Childs, D. Z. (2018).

Transient LTRE analysis reveals the demographic and trait-mediated processes that buffer

population growth. *Ecology Letters*, 23, 1353.

Maldonado-Chaparro, A. A., Read, D. W., & Blumstein, D. T. (2017). Can individual variation

in phenotypic plasticity enhance population viability? *Ecological Modelling*, 352, 19–30.

Marra, P. P., Cohen, E. B., Loss, S. R., Rutter, J. E., & Tonra, C. M. (2015). A call for full

annual cycle research in animal ecology. *Biology Letters*, 11, 20150552.

Merow, C., Dahlgren, J. P., Metcalf, C. J. E., Childs, D. Z., Evans, M. E. K., Jongejans, E.,

Record, S., Rees, M., Salguero-Gómez, R., & McMahon, S. M. (2014). Advancing

population ecology with integral projection models: a practical guide. *Methods in Ecology*

and Evolution, 5, 99–110.

Monclús, R., Pang, B., & Blumstein, D. T. (2014). Yellow-bellied marmots do not compensate

for a late start: the role of maternal allocation in shaping life-history trajectories.

Evolutionary Ecology, 28, 721–733.

Morgan, I. J., & Metcalfe, N. B. (2001). Deferred costs of compensatory growth after autumnal

food shortage in juvenile salmon. *Proceedings. Biological Sciences*, 268, 295–301.

Morris, W. F., Altmann, J., Brockman, D. K., Cords, M., Fedigan, L. M., Pusey, A. E., Stoinski,

T. S., Bronikowski, A. M., Alberts, S. C., & Strier, K. B. (2011). Low demographic

variability in wild primate populations: fitness impacts of variation, covariation, and serial

correlation in vital rates. *The American Naturalist*, 177, E14–E28.

Morris, W. F., Pfister, C. A., Tuljapurkar, S., Haridas, C. V., Boggs, C. L., Boyce, M. S., Bruna,

E. M., Church, D. R., Coulson, T., Doak, D. F., Forsyth, S., Gaillard, J.-M., Horvitz, C. C.,

Kalisz, S., Kendall, B. E., Knight, T. M., Lee, C. T., & Menges, E. S. (2008). Longevity can

buffer plant and animal populations against changing climatic variability. *Ecology*, 89, 19–

25.

Nouvellet, P., Newman, C., Buesching, C. D., & Macdonald, D. W. (2013). A multi-metric

approach to investigate the effects of weather conditions on the demographic of a terrestrial

mammal, the european badger (*Meles meles*). *PloS One*, 8, e68116.

Oli, M. K., & Armitage, K. B. (2004). Yellow-bellied marmot population dynamics:

demographic mechanisms of growth and decline. *Ecology*, 85, 2446–2455.

Orizaola, G., Dahl, E., & Laurila, A. (2014). Compensatory growth strategies are affected by the

strength of environmental time constraints in anuran larvae. *Oecologia*, 174, 131–137.

Ozgul, A., Childs, D. Z., Oli, M. K., Armitage, K. B., & Blumstein, D. T. (2010). Coupled

dynamics of body mass and population growth in response to environmental change. *Nature*,

466, 482-485.

Ozgul, A., Oli, M. K., & Armitage, K. B. (2009). Influence of local demography on asymptotic

and transient dynamics of a yellow-bellied marmot metapopulation. *The American*

Naturalist, 173, 517-530.

Ozgul, A., Oli, M. K., Olson, L. E., Blumstein, D. T., & Armitage, K. B. (2007). Spatiotemporal

variation in reproductive parameters of yellow-bellied marmots. *Oecologia*, 154, 95–106.

Paniw, M., Maag, N., Cozzi, G., Clutton-Brock, T., & Ozgul, A. (2019). Life history responses

of meerkats to seasonal changes in extreme environments. *Science*, 363, 631–635.

Paniw, M., Ozgul, A., & Salguero-Gómez, R. (2018). Interactive life-history traits predict

sensitivity of plants and animals to temporal autocorrelation. *Ecology Letters*, 21, 275–286.

Paniw, M., Quintana-Ascencio, P. F., Ojeda, F., & Salguero-Gómez, R. (2017a). Accounting for

uncertainty in dormant life stages in stochastic demographic models. *Oikos* , 126, 900–909.

Paniw, M., Quintana-Ascencio, P. F., Ojeda, F., & Salguero-Gómez, R. (2017b). Interacting

livestock and fire may both threaten and increase viability of a fire-adapted Mediterranean

carnivorous plant. *The Journal of Applied Ecology*, 54, 1884–1894.

Pfister, C. A. (1998). Patterns of variance in stage-structured populations: evolutionary

predictions and ecological implications. *Proceedings of the National Academy of Sciences of*

the United States of America, 95, 213–218.

Picó, F. X., de Kroon, H., & Retana, J. (2002). An extended flowering and fruiting season has

few demographic effects in a Mediterranean perennial herb. *Ecology*, 83, 1991–2004.

Pol, M., Bailey, L. D., McLean, N., Rijsdijk, L., Lawson, C. R., & Brouwer, L. (2016).

Identifying the best climatic predictors in ecology and evolution. *Methods in Ecology and*

Evolution, 7, 1246–1257.

Reed, T. E., Grøtan, V., Jenouvrier, S., Sæther, B.-E., & Visser, M. E. (2013). Population growth

in a wild bird is buffered against phenological mismatch. *Science*, 340, 488–491.

Rees, M., & Ellner, S. P. (2009). Integral projection models for populations in temporally

varying environments. *Ecological Monographs*, 79, 575–594.

Ripley, B. J., & Caswell, H. (2006). Recruitment variability and stochastic population growth of

the soft-shell clam, *Mya arenaria*. *Ecological Modelling*, 193, 517–530.

Robert, A., Bolton, M., Jiguet, F., & Bried, J. (2015). The survival–reproduction association

becomes stronger when conditions are good. *Proceedings. Biological Sciences*, 282,

20151529.

Rotella, J. J., Link, W. A., Chambert, T., Stauffer, G. E., & Garrott, R. A. (2012). Evaluating the

demographic buffering hypothesis with vital rates estimated for Weddell seals from 30 years

of mark-recapture data. *The Journal of Animal Ecology*, 81, 162–173.

Ruf, T., Bieber, C., Arnold, W., & Millesi, E. (2012). *Living in a Seasonal World:*

Thermoregulatory and Metabolic Adaptations. Springer Science & Business Media.

Rushing, C. S., Hostetler, J. A., Sillett, T. S., Marra, P. P., Rotenberg, J. A., & Ryder, T. B.

(2017). Spatial and temporal drivers of avian population dynamics across the annual cycle.

Ecology, 98, 2837–2850.

Salguero-Gómez, R., Jones, O. R., & Archer, C. R., et al. (2015). The COMPADRE Plant Matrix

Database: an open online repository for plant demography. *Journal of Ecology*, 103, 202-

218.

Salguero-Gómez, R., Jones, O. R., Archer, C. R., et al. (2016). COMADRE: a global data base

of animal demography. *The Journal of Animal Ecology*, 85, 371–384.

Schwartz, O. A., & Armitage, K. B. (2002). Correlations between weather factors and life-

history traits of yellow-bellied marmots. *Proceedings of the 3rd International Marmot*

Conference, Cheboksary, Russia.

Schwartz, O. A., & Armitage, K. B. (2005). Weather influences on demography of the yellow-

bellied marmot (*Marmota flaviventris*). *Journal of Zoology*, 265, 73–79.

Schwartz, O. A., Armitage, K. B., & Van Vuren, D. (1998). A 32-year demography of yellow-

bellied marmots (*Marmota flaviventris*). *Journal of Zoology*, 246, 337–346.

Simmonds, E. G., & Coulson, T. (2015). Analysis of phenotypic change in relation to climatic

drivers in a population of Soay sheep *Ovis aries*. *Oikos*, 124, 543–552.

Teller, B. J., Adler, P. B., Edwards, C. B., Hooker, G., & Ellner, S. P. (2016). Linking

demography with drivers: climate and competition. *Methods in Ecology and Evolution*, 7,

171–183.

Töpper, J. P., Meineri, E., Olsen, S. L., Rydgren, K., Skarpaas, O., & Vandvik, V. (2018). The

devil is in the detail: non-additive and context-dependent plant population responses to

increasing temperature and precipitation. *Global Change Biology*, 24, 4657-4666.

Van de Pol, M., Vindenes, Y., Sæther, B.-E., Engen, S., Ens, B. J., Oosterbeek, K., & Tinbergen,

J. M. (2010). Effects of climate change and variability on population dynamics in a long-

lived shorebird. *Ecology*, 91, 1192–1204.

Van Vuren, D., & Armitage, K. B. (1994). Survival of dispersing and philopatric yellow-bellied

marmots: what is the cost of dispersal? *Oikos*, 69, 179–181.

Varpe, Ø. (2017). Life history adaptations to seasonality. *Integrative and Comparative Biology*,

57, 943–960.

Villellas, J., Doak, D. F., García, M. B., & Morris, W. F. (2015). Demographic compensation

among populations: what is it, how does it arise and what are its implications? *Ecology*

Letters, 18, 1139–1152.

Van Vuren, D., & Armitage, K. B. (1991). Duration of snow cover and its influence on life-

history variation in yellow-bellied marmots. *Canadian Journal of Zoology*, 69, 1755–1758.

Williams, J. L., Miller, T. E. X., & Ellner, S. P. (2012). Avoiding unintentional eviction from

integral projection models. *Ecology*, 93, 2008–2014.

Woodroffe, R., Groom, R., & McNutt, J. W. (2017). Hot dogs: High ambient temperatures

impact reproductive success in a tropical carnivore. *The Journal of Animal Ecology*, 86,

1329–1338.

Williams J. L., Miller T. E. X. & Ellner S. P. (2012). Avoiding unintentional eviction from

integral projection models. *Ecology*, 93, 2008–2014.