Molecular Basis for G-protein-Coupled Receptor (GPCR) Activation and Biased Signalling at the Platelet Thrombin Receptor Proteinase Activated Receptor-4 (PAR4).

Pierre E. Thibeault[1], Jordan C. LeSarge[2], D'Arcy Arends[1], Michaela Fernandes[2], Peter Chidiac[1], Peter B. Stathopulos[1], Leonard G. Luyt[2,3,4], and Rithwik Ramachandran[1*]

[1] Department of Physiology and Pharmacology, University of Western Ontario, London, ON, N6A5C1, Canada.

[2] Department of Chemistry, The University of Western Ontario, London, ON, N6A5C1, Canada.

[3] Departments of Oncology, The University of Western Ontario, London, ON, N6A5C1, Canada.

[4] London Regional Cancer Program, Lawson Health Research Institute, London, ON, N6C2R5, Canada.

* For correspondence:

Rithwik Ramachandran,
Department of Physiology and Pharmacology,
Schulich School of Medicine and Dentistry,
University of Western Ontario, London,
Ontario, Canada.
Phone: 519-661-2142
email: rramach@uwo.ca

Abstract:

Proteinase Activated Receptor-4 (PAR4) is a member of the proteolytically-activated PAR family of G-Protein-coupled Receptors (GPCRs). PARs are activated following proteolytic cleavage of the receptor N-terminus by enzymes such as thrombin, trypsin, and cathepsin-G to reveal the receptor-activating motif termed the tethered ligand. The tethered ligand binds intramolecularly to the receptor and triggers receptor signalling and cellular responses. In spite of this unusual mechanism of activation, PARs are fundamentally peptide receptors and can also be activated by exogenous application of short synthetic peptides derived from the tethered ligand sequence. In order to gain a better understanding of the molecular basis for PAR4-dependent signalling, we examined signalling responses to a library of peptides derived from the canonical PAR4 activating peptide (PAR4-AP), AYPGKF-NH$_2$. We examined residues involved in activation of the G$\alpha_{q/11}$-coupled calcium signalling pathway, β-arrestin recruitment, and mitogen-activated protein kinase pathway activation. The peptide *N*-methyl-alanine-YPGKF-NH$_2$ was identified as a compound that is a poor activator of PAR4-dependent calcium signalling but was fully competent in recruiting β-arrestin-1 and -2. In order to gain a better understanding of the ligand-binding pocket, we used *in silico* docking to identify key residues involved in PAR4 interaction with AYPGKF-NH$_2$. The predicted interactions were verified by site-directed mutagenesis and analysis of calcium signalling and β-arrestin-1/-2 recruitment following proteolytic activation (with thrombin) or activation with the synthetic agonist peptide (AYPGKF-NH$_2$). We determined that a key extracellular loop-2 aspartic acid residue (Asp^{230}) is critical for signalling following both proteolytic and peptide activation of PAR4. Finally, we investigated platelet aggregation in response to A**y**PGKF-NH$_2$ (a peptide with D-tyrosine in position two) which is unable to activate calcium signalling, and AYPG**R**F-NH$_2$ a peptide that is equipotent to the parental peptide AYPGKF-NH$_2$ for calcium signalling but is more potent at recruiting β-arrestins. We found that A**y**PGKF-NH$_2$ fails to activate platelets while AYPG**R**F-NH$_2$ causes a platelet aggregation response that is greater than that seen with the parental peptide and is comparable to that seen with thrombin stimulation. Overall, these studies uncover molecular determinants for agonist binding and signalling through a non-canonically activated GPCR and provide a template for development of small molecule modulators of PAR4.

Introduction:

G-Protein-coupled receptors (GPCRs) are the largest family of cell surface receptors and regulate a host of important physiological responses (Fredriksson and Schioth, 2005; Pierce et al., 2002). GPCRs respond to a variety of extracellular signals and regulate cellular behaviour through engaging intracellular effector molecules to activate various cell-signalling pathways (Erlandson et al., 2018; Hilger et al., 2018). In drug discovery, GPCRs are a valuable and highly tractable class of drug targets with over 30% of all currently approved drugs acting on GPCRs (Sriram and Insel, 2018). GPCR activation occurs typically through binding of a soluble ligand and conformational changes that enable engagement of G-protein-dependent or β-arrestin-mediated signal transduction. The Proteinase Activated Receptors (PARs) are a four-member family of GPCRs with a distinct mechanism of activation that involves limited proteolysis and unmasking of a receptor activating motif called the tethered ligand. PARs were first discovered in an effort to identify the cellular receptors responsible for actions of thrombin that were independent from its role in the coagulation cascade (Vu et al., 1991). Various proteinases have now been described as activators of the different members of the PAR family, but canonically PAR1 and PAR4 are described as thrombin-activated receptors, while PAR2 is activated by trypsin and other trypsin-like serine proteinases. PAR4 can also be activated by trypsin. Whether PAR3 signals independently is unclear, though thrombin can cleave the PAR3 N-terminus to reveal a tethered ligand (Adams et al., 2011; Ishihara et al., 1997). Following the discovery of PARs it was quickly realized that two receptors, PAR1 and PAR4, serve as the thrombin receptors on human platelets (Coughlin, 2005, 1999). PAR1 and PAR4 both trigger platelet activation and aggregation. Given that anti-platelet agents are an important class of drugs for the treatment of various cardiovascular diseases, there has been a concerted effort aimed at understanding the role of these receptors in platelet activation and in identifying molecules that can act as PAR1 and PAR4 antagonists. A PAR1 antagonist was recently identified and approved for clinical use, however significant bleeding side effects were noted in patients administered this drug (Morrow et al., 2012, 2009; Scirica et al., 2012). While bleeding is a frequently encountered side effect of therapeutically targeting pathways involved in blood coagulation, it was hoped that targeting PAR1 would not present such liability. Much attention has now shifted to understanding and targeting the other thrombin receptor PAR4. Though both PAR1 and PAR4 are activated by thrombin, there are some key differences reported in the activation of these two receptors on platelets that argue for PAR4 being a better anti-platelet drug target than PAR1. PAR1 possesses a hirudin-like binding site that enables it to bind thrombin with relatively high affinity (Jacques et al., 2000; Liu et al., 1991). In lacking this thrombin-binding site, PAR4 can only be activated at much higher concentrations of thrombin, such as may be encountered on a growing platelet thrombus. There are also reported differences in the kinetics of PAR1 and PAR4-dependent calcium signalling, with PAR4 causing a delayed but more sustained signal compared to PAR1 (Shapiro et al., 2000). This is thought to be due to dual proline residues in the PAR4 exodomain that provide low-affinity interactions with the thrombin active site and an anionic cluster that slows dissociation from thrombin (Jacques and Kuliopulos, 2003). Overall, PAR1 is thought to initiate platelet activation in response to low concentrations of thrombin while PAR4 is engaged at higher concentrations to further consolidate and stabilize the clot.

Given the liabilities seen with targeting PAR1 as a platelet thrombin receptor there has been
renewed interest in developing small molecule PAR4 antagonists (Rwibasira Rudinga et al.,
2018; Wu et al., 2000; Young et al., 2010). Recently small molecule antagonists of PAR4 have
been described with efficacy in inhibiting injury induced thrombosis in both non-human
primates and humans (Miller et al., 2019; Wilson et al., 2017; Wong et al., 2017). Despite these
promising advances there remains the question of whether a complete inhibition of PAR4 might
also lead to bleeding liability. Our efforts have therefore focused on examining biased
signalling through PAR4 in order to identify pathways that are responsible for platelet
aggregation and that could be selectively manipulated for therapeutic efficacy. PAR4 can
activate multiple signalling pathways through coupling to $G\alpha_{q/11}$, $G\alpha_{12/13}$ and β-arrestin
(Holinstat et al., 2006; Li et al., 2011; Ramachandran et al., 2017). In previous work, we and
others showed that pepducins targeting different intracellular motifs in PAR4 could be
employed to differentially inhibit platelet activation by blocking G-protein-coupled or β-
arrestin-dependent signalling (Covic et al., 2002; Ramachandran et al., 2017). The ability to
selectively activate or inhibit GPCR signalling in a biased manner is of great interest and has
been proposed as a strategy for obtaining therapeutically superior drugs for a number of
conditions (Bar-Shavit et al., 2016; Hollenberg et al., 2014; Seyedabadi et al., 2019). In the
case of the finely balanced signalling systems involved in coagulation and platelet activation,
such subtle perturbations may be key to obtaining drugs which do not result in bleeding
liability.
122
Here we seek to gain an understanding of the molecular interactions responsible for agonist
docking to PAR4 and to identify interactions that can be manipulated to obtain biased ligands
for PAR4. By screening a peptide library derived from the most widely used PAR4 agonist
peptide, AYPGKF-NH$_2$, we have identified key residues that confer selective activation of
$G\alpha_{q/11}$ coupled calcium signalling and/or β-arrestin recruitment. Through *in silico* docking and
confirmatory mutagenesis experiments we further identify an extracellular loop 2 residue that
is critical for receptor activation and signalling. We also examine platelet activation by peptides
that show differential activation of PAR4 signalling. Overall, these studies advance our
understanding of PAR4 activation and signalling and will help guide future drug discovery
efforts targeting this receptor.

Results:

Experimental design and data analysis. In order to understand the rules governing agonist
peptide activation of PAR4 we synthesized 37 hexapeptides through modification to the
synthetic PAR4 agonist peptide AYPGKF-NH$_2$. A number of different modifications were
examined at each of the six positions, including alanine substitutions, D-isomer substitutions,
N-methyl substitutions, as well as other modifications aimed at further probing each of the six
positions. Henceforth, the parental PAR4 agonist peptide for this study, AYPGKF-NH$_2$ (Ala[1]-
Tyr[2]-Pro[3]-Gly[4]-Lys[5]-Phe[6]-NH$_2$; superscript denotes position from N-terminus), will be referred
to as PAR4-AP. G$\alpha_{q/11}$-coupled calcium signalling, β-arrestin-1/-2 recruitment, and p44/42
mitogen-activated protein kinase (MAPK) signalling by each of these peptides was monitored
and compared to responses triggered by PAR4-AP (Table 1). Whereas PAR4-AP triggered
G$\alpha_{q/11}$ signalling reached a clear plateau over the range of concentrations tested (0.3 − 300
µM), β-arrestin recruitment did not and therefore the effective concentration required to
achieve half-maximal response (EC$_{50}$) values could not be accurately assessed. In contrast, with
the substituted peptides, G$\alpha_{q/11}$ signalling concentration-effect curves in some cases did not
clearly plateau, whereas β-arrestin recruitment did. Where possible, EC$_{50}$ values are compared
between PAR4-AP and substituted peptides. However, since clear maxima were not always
obtained, comparisons of signal magnitude throughout the present study are presented as the
response elicited by a given compound at 100 µM shown as a percentage of responses elicited
by PAR4-AP at the same concentration. Since the concentration-effect curve for PAR4-AP
exhibits a saturating calcium signal, peptides that failed to saturate calcium signalling responses
are depicted as having EC$_{50}$ values that were 'not determined' (*n.d.*). Maximal calcium signal
responses are shown as a percentage of the response obtained with the calcium ionophore
A23187 (Table 1, Max; % of A23187 calcium ionophore) and are referred to throughout the
results as a change in maximal calcium signalling as a measure of the efficacy of the peptides.
Throughout the results, β-arrestin recruitment values are presented as a percentage (%) of the
PAR4-AP-induced recruitment at 100 µM concentration. For ease of comparison,
concentration-effect curves for PAR4-AP-mediated G$\alpha_{q/11}$ calcium signalling and β-arrestin
recruitment are displayed as reference values in each of the Figures 1 through 7.

Effect of alanine substitution on PAR4-AP-mediated signalling responses. Our first set of
changes involved alanine substitutions at positions 2-6 of the PAR4-AP (calcium EC$_{50}$ = 12.1
± 3.1 µM; β-arrestin-1 recruitment at 100 µM for comparison is considered 100 ± 3.9%, and
100 ± 3.5% for β-arrestin-2) (Fig. 1, Table 1). Potency was decreased with respect to calcium
signalling when tyrosine[2] was substituted with alanine, and failed to reach a plateau over the
concentration range tested (A**A**PGKF-NH$_2$; EC$_{50}$ = *n.d.*). Correspondingly, β-arrestin
recruitment was reduced compared to PAR4-AP at the same concentration (β-arr-1 47.3 ±
3.6%; β-arr-2 46.4 ± 3.2%). Substitution of proline[3] with alanine resulted in a significant
decrease in potency for calcium signalling (AY**A**GKF-NH$_2$ EC$_{50}$ = 63.7 ± 20.8 µM) and yielded
reduced β-arrestin recruitment (β-arr-1 48.5 ± 7.9%; β-arr-2 43.4 ± 7.1%). Either glycine[4] or
phenylalanine[6] substitution to alanine resulted in reduced maximal calcium signalling
compared to PAR4-AP at the same concentration, with no significant effect on potency (EC$_{50}$

$= 21.2 \pm 6.6\ \mu M$; $EC_{50} = 17.3 \pm 5.0\ \mu M$, respectively). Additionally, alanine substitutions in
these positions resulted in significantly reduced β-arrestin recruitment, to levels approximately
half of those observed with PAR4-AP (AYP**A**KF-NH$_2$ β-arr-1 $56.4 \pm 5.6\%$, β-arr-2 $54.6 \pm$
5.1%; AYPGK**A**-NH$_2$ β-arr-1 $47.9 \pm 3.8\%$, β-arr-2 $52.6 \pm 3.5\%$). Interestingly, we observed
that lysine[5] to alanine substitution had no appreciable impact on potency with respect to
calcium signalling (AYPG**A**F-NH$_2$ $EC_{50} = 9.8 \pm 2.9\ \mu M$); however, this substitution did reduce
recruitment of β-arrestin-1 ($59.3 \pm 7.2\%$) and -2 ($70.8 \pm 6.5\%$). Collectively, alanine
substitutions of positions 2-6 resulted in decreased β-arrestin recruitment compared to the
parental peptide (Fig. 1 B. and C.; Table 1). Interestingly, we observe differential effects on
calcium signalling with significant detriment to potency attributed to tyrosine substitution (Fig.
1 A.). Additionally, substitution of glycine and phenylalanine moderately decreased maximal
calcium signalling responses, perhaps indicating that these residues are important for full
agonism of this pathway comparative to that achieved with the parental peptide. Notably, all
single alanine substitutions of positions 2-6 of the PAR4-AP retained some agonist activity
indicating that signalling is not wholly reliant on any single residue alone.
190
Effect of D-isomer amino acid substitutions on PAR4-AP-mediated signalling responses.
To assess the impact of stereochemical inversion at each residue we generated five peptides
with single amino acid substitutions to their respective D-isomers (Fig. 2, Table 1). Glycine
possesses only one isomeric form and thus is not substituted. Both L-alanine to D-alanine
(**a**YPGKF-NH$_2$) and L-lysine to D-lysine (AYPG**k**F-NH$_2$) substitutions resulted in significant
losses of potency in calcium signalling assays compared to PAR4-AP. Consistent with calcium
data, we observed a significant decrease in β-arrestin recruitment compared to the parental
PAR4-AP (**a**YPGKF-NH$_2$ β-arr-1 $48.7 \pm 2.6\%$, β-arr-2 $47.5 \pm 2.4\%$; AYPG**k**F-NH$_2$ β-arr-1
$47.4 \pm 1.0\%$, β-arr-2 $36.5 \pm 0.9\%$). Interestingly, both D-tyrosine (A**y**PGKF-NH$_2$) and D-
proline (AY**p**GKF-NH$_2$) substitutions completely abolished calcium signalling. Comparably,
β-arrestin recruitment was significantly reduced with stimulation by either peptide (A**y**PGKF-
NH$_2$ β-arr-1 $50.7 \pm 4.6\%$, β-arr-2 $46.4 \pm 4.2\%$; AY**p**GKF-NH$_2$ β-arr-1 $45.2 \pm 10.5\%$, β-arr-2
$38.1 \pm 9.5\%$). In keeping with what we observed with phenylalanine to alanine substitution
(Fig. 1), D-phenylalanine substitution in position 6 (AYPGK**f**-NH$_2$) resulted in less efficacious
agonism of both calcium ($EC_{50} = 22.2 \pm 4.7\ \mu M$) and β-arrestin recruitment (β-arr-1 $80.3 \pm$
6.6%, β-arr-2 $74.1 \pm 6.0\%$). Overall, D-isomers substitutions of amino acids alanine[1], tyrosine[2],
proline[3], and lysine[5] all result in significant reduction of (D-alanine, D-lysine) or abolition of
(D-tyrosine, D-proline) calcium signalling while causing significant reductions in β-arrestin
recruitment. Substitution of L-phenylalanine[6] for D-phenylalanine was the only D-isomer
substitution that is somewhat tolerated, resulting in modest reductions in potency with respect
to both calcium signalling and β-arrestin recruitment. These data reveal that the sidechain
stereochemistry of residues within the AYPGKF-NH$_2$ peptide perform a role in agonism.
213
Effect of *N*-methylation on PAR4-AP-mediated signalling responses. We generated a series
of peptides to investigate the pharmacological impact of *N*-methylation along the peptide
backbone (Fig. 3, Table 1). In contrast to the parental PAR4-AP, all but one of the N-
methylated peptides evaluated yielded clear plateaus and discernible EC_{50} values in β-arrestin
recruitment assays, and where applicable these are indicated below along with their relative

responses compared to PAR4-AP at 100 µM. Substitution of alanine[1] with *N*-methyl-alanine
[(**N-Me-A**)-YPGKF-NH2] almost completely abolished calcium signalling (EC_{50} = *n.d.*).
Interestingly, this peptide was able to initiate β-arrestin-1/-2 recruitment comparable to levels
obtained with PAR4-AP (β-arr-1 119.2 ± 11.9%, EC_{50} = 6.8 ± 2.6 µM; β-arr-2 80.6 ± 10.8%;
EC_{50} = 2.0 ± 0.7 µM). These results suggest that increasing steric bulk and hydrophobicity of
the backbone at amide positions through the addition of an N-terminal methyl group drives
PAR4 signalling towards β-arrestin pathways and causes only modest activation of calcium
signalling. Substitution of tyrosine[2] for *N*-methyl-tyrosine [A-(**N-Me-Y**)-PGKF-NH2]
abolished calcium signalling up to 300 µM (EC_{50} = *n.d.*). Consistent with calcium signalling,
we observed decreased signal magnitude in β-arrestin recruitment assays compared to PAR4-
AP (β-arr-1 71.4 ± 10.8%, EC_{50} = 14.2 ± 8.6 µM; β-arr-2 61.5 ± 9.8%, EC_{50} = 11.8 ± 5.7 µM).
Substitution of the PAR4-AP glycine[4] with sarcosine (Sar., a.k.a *N*-methyl-glycine; AYP-**Sar**-
KF-NH2) resulted in significantly reduced calcium potency (EC_{50} = *n.d.*) and significantly
reduced β-arrestin-1/-2 recruitment (35.9 ± 4.3%, 36.1 ± 3.9% respectively). *N*-methylation of
the alpha (α) amine of lysine[5] [AYPG-(**N**$_{α}$**-Me-K**)-F-NH2] resulted in significantly reduced
calcium signalling (EC_{50} = *n.d.*) with no significant effect on maximal β-arrestin recruitment
(β-arr-1 126.7 ± 18.6%, EC_{50} = 12.2 ± 4.3 µM; β-arr-2 77.1 ± 16.9%, EC_{50} = 2.5 ± 1.0 µM).
Substitution of phenylalanine[6] with *N*-methyl-phenylalanine [AYPGK-(**N-Me-F**)-NH2] was
equipotent for calcium signalling compared to PAR4-AP (EC_{50} = 7.6 ± 1.9 µM). Interestingly,
this peptide showed increased recruitment of both β-arrestins (β-arr-1 136.9 ± 18.0%, EC_{50} =
10.4 ± 3.3 µM; β-arr-2 96.6 ± 16.4%, EC_{50} = 15.3 ± 4.6 µM). Together, these data reveal that
N-methylation of positions 1, 2, 4, and 5 are deleterious to calcium signalling.

We were excited to observe evidence of β-arrestin-biased peptides within this group, all of
which showed decreased calcium signalling and/or increased β-arrestin recruitment. Moreover,
the leftward shifts in their associated β-arrestin concentration-effect profiles imply that
substitution of *N*-methylated alanine[1], tyrosine[2], lysine[5], or phenylalanine[6] residues increases
peptide potency with respect to G-protein-independent signalling. Only *N*-methylation of
glycine[4] (sarcosine) resulted in a reduced overall ability to activate PAR4, with relative calcium
and β-arrestin signals being reduced by approximately 90% and 60%, respectively.

Effect of N-terminal and position 1 modifications on PAR4-AP-mediated signalling
responses. To further investigate the impact of *N*-terminal alterations to the PAR4-AP we
examined the impact of N-terminal acetylation and substitutions of position 1 (Fig. 4, Table 1).
Previously it was demonstrated that protonation of the N-terminus of the PAR1 tethered ligand
(SFLLRN) is critical for agonist function (Scarborough et al., 1992). Additionally, substitution
of glycine for mercaptoproprionic acid (Mpr), which lacks an amino-terminal protonated amine
in the native tethered-ligand-mimicking peptide GYPGQV, resulted in complete loss of agonist
activity (Faruqi et al., 2000). Given the evidence for N-terminal protonation in PAR agonism,
we sought to determine the impact of N-terminal acetylation on PAR4-AP activity. N-terminal
acetylation of the PAR4-AP (**Ac**-AYPGKF-NH2) negatively impacted both calcium signalling
(EC_{50} = *n.d.*) as well as the magnitude of β-arrestin recruitment to PAR4 as expected (β-arr-1
47.3 ± 3.1%, EC_{50} = 15.0 ± 4.3 µM; β-arr-2 45.5 ± 2.8%, EC_{50} = 26.7 ± 5.6 µM). In contrast

to the previously reported loss of activity with Mpr-YPGQV peptide, we observe partial
agonism with the loss of protonation in the PAR4-AP.
264
In both the native human PAR4 tethered-ligand sequence and the PAR4-AP, position 1 is
occupied by small (tethered ligand – glycine, PAR4-AP – alanine) and non-polar (PAR4-AP –
alanine) residues. To investigate the role of position 1 in the PAR4-AP, we generated seven
peptides with position 1 substitutions. Introduction of steric bulk and increased hydrophobicity
through substitution of alanine[1] for valine resulted in partial agonism of both calcium signalling
and β-arrestin recruitment (**VYPGKF-NH$_2$** EC$_{50}$ = 22.9 ± 8.3 µM; β-arr-1 73.6 ± 13.0%, EC$_{50}$
= 33.8 ± 13.7 µM; β-arr-2 55.2 ± 11.8%, EC$_{50}$ = 13.7 ± 8.2 µM). Additionally, increasing
hydrophobic bulk at position 1 through substitution with norleucine completely abolished
calcium signalling (**Nle**-YPGKF-NH$_2$, EC$_{50}$ = *n.d.*) and resulted in partial agonism of β-arrestin
recruitment (β-arr-1 40.2 ± 9.7%; β-arr-2 36.5 ± 8.8%). Together, data from these substitutions
indicate that the precise steric bulk of the side chain in position 1 is important for PAR4
agonism.
277
Next, we evaluated the impact of steric bulk relocation in position 1. Relocation of the methyl
group in alanine to the N-terminus and loss of side chain through substitution with sarcosine
(**Sar**-YPGKF-NH$_2$) resulted in poor agonism at both calcium signalling (EC$_{50}$ = *n.d.*) and β-
arrestin recruitment (β-arr-1 35.6 ± 2.0%, β-arr-2 39.8 ± 1.8%). Further, the addition of an α-
methyl group through substitution of alanine[1] with 2-aminoisobutyric acid (**Aib**-YPGKF-NH$_2$)
resulted in a significantly less potent partial agonist for calcium signalling (EC$_{50}$ = 50.2 ± 15.3
µM) and also resulted in a significant reduction in β-arrestin recruitment (β-arr-1 38.6 ± 3.9%;
β-arr-2 37.3 ± 3.6%). Similarly, replacement with *N*-methyl-serine [**(N-Me-S)**-YPGKF-NH$_2$]
also caused a significant attenuation of calcium signalling (EC$_{50}$ = *n.d.*), while also revealing a
decrease in β-arrestin recruitment compared to PAR4-AP (β-arr-1 78.1 ± 6.9%, β-arr-2 51.7 ±
6.3%). Thus, it is apparent that the precise position of the side chain in position 1 is important
for agonism.
290
To further probe the importance of charge localization at the N-terminal region of the peptide,
we examined the effect of changing the localization of the positive charge in position 1.
Substitution of alanine with either beta-alanine or isonipecotic acid significantly reduced the
potency and activity of the PAR4-AP, similar to the results with PAR4-AP N-terminal
acetylation (**betaAla**-YPGKF-NH2 calcium EC$_{50}$ = *n.d.*, β-arr-1 25.7 ± 4.8%, β-arr-2 31.8 ±
4.4%; **Inp**-YPGKF-NH$_2$ calcium EC$_{50}$ = *n.d.*, β-arr-1 32.0 ± 3.3%, β-arr-2 26.9 ± 3.0%).
Together these data suggest a role for N-terminal charge localization in agonist activity.
298
Effect of additional substitutions in positions 2-4 on PAR4-AP-mediated signalling.
Previously structure activity studies with PAR4 activating peptides revealed an apparent
requirement for an aromatic residue in position 2 (tyrosine) and found that alterations to
position 3 (proline) and position 4 (glycine) were not well tolerated (Faruqi et al., 2000). In the
present study, we observed that substitution of either tyrosine, proline, or glycine with alanine
reduced calcium signalling potency and β-arrestin recruitment, however none of these

substitutions completely abolished activity of the peptide (Fig. 1). Interestingly, substitution of either L-tyrosine or L-proline with the respective D-isomer significantly reduced the activity of the PAR4-AP (Fig. 2, Table 1). Together, these data suggest that while no individual residue is strictly required for agonist activity, the side chains of these residues and their positioning are important for agonist function. To further probe the contribution of these residues to the PAR4-AP, we generated peptides with alterations in the aromatic side chain of position 2, the backbone conformation of position 3, or position 4 (Fig. 5, Table 1).

Peptides with substitution of tyrosine[2] with either 4-fluoro-L-phenylalanine [A-**F(4-fluoro)**-PGKF-NH2] or *O*-methyl-tyrosine [A-**Tyr(Me)**-PGKF-NH$_2$] retained calcium signalling and β-arrestin recruitment with no significant changes in potency compared to the PAR4-AP. These data suggest that the side chain of tyrosine may be a site in which further modifications can be made to the peptide without affecting agonist potential; however, by comparing A**A**PGKF-NH$_2$ the presence of an aromatic ring in position 2 is of importance. Given the results obtained with alanine or D-proline substitution of proline[3] (Fig. 1 & 2, respectively), we furthered probed conformational contributions of this residue to the parental PAR4-AP. Proline was substituted with pipecolic acid (2-piperidinecarboxylic acid; Pip) which is structurally similar to proline but has an increased steric bulk and backbone elongation provided by a six-membered ring compared to the five-membered ring of proline. We observed partial agonism of calcium signalling in response to AY-**Pip**-GKF-NH$_2$ stimulation with no significant difference in agonist potency ($EC_{50} = 30.7 \pm 4.9$ μM) with a correlating partial agonism of β-arrestin recruitment (β-arr-1 38.8 ± 3.0%; β-arr-2 44.0 ± 2.7%). Thus, it appears that increasing steric bulk/backbone length may affect efficacy but still results in agonism. To further investigate the role of backbone conformation provided at position 3, proline was substituted with nipecotic acid (3-piperidinecarboxylic acid; Nip) which is similar to pipecotic acid, however, results in change of the peptide backbone direction (AY-**Nip**-GKF-NH$_2$). With this modification, both calcium signalling ($EC_{50} = $ *n.d.*) and β-arrestin recruitment (β-arr-1 32.9 ± 4.9%; β-arr-2 30.9 ± 4.5%) were significantly decreased in response to PAR4 stimulation. Elongation of the backbone in position four (glycine) by substitution with beta-alanine (AYP-**betaAla**-KF-NH$_2$) was also significantly detrimental to both calcium signalling ($EC_{50} = $ *n.d.*) and β-arrestin recruitment (β-arr-1 42.5 ± 1.8%, β-arr-2 24.9 ± 1.6%). Together these data suggest an important role for backbone conformation contributed by proline[3] and glycine[4] compactness in the PAR4-AP. As well, these data suggest peptide backbone length at proline[3] and glycine[4] has an important role; consistent with what we have observed with backbone elongation modifications at position 1 (**Inp**-YPGKF-NH$_2$ and **betaAla**-YPGKF-NH$_2$; Fig. 4, Table 1).

Effect of additional substitutions in position 5 on PAR4-AP-mediated signalling responses. Previously, investigations into glutamine substitutions in the native tethered ligand mimicking peptide, GYPGQV, revealed that agonism is maintained when this position is substituted with a charged residue such as arginine or ornithine (Faruqi et al., 2000). In the present study, we have identified that substitution of position 5 (lysine) of the PAR4-AP (AYPGKF-NH$_2$) with alanine does not significantly affect the agonist capacity of the peptide

(Fig. 1). Further, we identified that substitution of L-lysine with D-lysine significantly decreases agonist stimulation of calcium signalling and β-arrestin recruitment (Fig. 2). To further probe the effect of positively charged residues in position 5, we generated several PAR4-AP analogues with lysine^5 substitutions by either arginine or ornithine (Fig. 6, Table 1). Substitution of lysine^5 with arginine resulted in a partial agonist with respect to calcium signalling but increased β-arrestin recruitment compared to the parental peptide (AYPG**R**F-NH$_2$ calcium EC$_{50}$ = 4.0 ± 1.1 µM, β-arr-1 154.2 ± 10.3%, β-arr-2 140.1 ± 9.4%). This substitution delocalizes the positive charge of the side chain compared to lysine and increases steric bulk. Interestingly, substitution with ornithine, which is similarly charged but has a shortened distance between the charge and the peptide backbone compared to lysine, also resulted in partial agonism of PAR4-mediated calcium signalling accompanied by a modest increase in the recruitment of β-arrestins (AYPG**O**F-NH$_2$ calcium EC$_{50}$ = 4.2 ± 1.1 µM, β-arr-1 121.7 ± 8.5%, β-arr-2 112.1 ± 7.7%). To further probe if steric bulk in the side chain of position 5 affects PAR4-AP potency, we increased steric bulk through methylation of the epsilon (ε) amine of lysine [AYPG-(**N$_\varepsilon$-Me-K**)-F-NH$_2$]. This substitution yielded a full agonist for calcium signalling (EC$_{50}$ = 6.7 ± 1.9 µM) and a modestly more efficacious agonist for the recruitment of β-arrestins (β-arr-1 122.5 ± 12.4%; β-arr-2 116.7 ± 11.2%). All of these substitutions maintain a positive charge and are able to act as agonists for PAR4. To determine if charge in position 5 is required, we substituted lysine with citrulline, which maintains similar structure to arginine and lysine while removing a positive charge (AYPG-**Cit**-F-NH$_2$). Interestingly, we observed partial agonism with respect to calcium signalling (EC$_{50}$ = 5.4 ± 0.8 µM) with equipotent β-arrestin recruitment compared to the PAR4-AP, consistent with the alanine scan data for this position. Thus, positively charged residues in position 5 seem to contribute to agonism. Interestingly, we observed similar increases in β-arrestin recruitment and equipotent calcium signalling when stericity was either increased (AYPG**R**F-NH$_2$) or decreased (AYPG**O**F-NH$_2$), suggesting that the positive charge in this position is more important than the size of the side chain. Finally, although both the charge and size of the size chain have an effect, all modifications to position 5, with the exception of D-lysine (Fig. 2, Table 1) and N_α-methyl-lysine (Fig. 3, Table 1) modifications, were generally well-tolerated and there appears to be modest freedom in the side-chain of this position. This makes the side chain a good candidate for further investigations of structure-activity relationships as well as a good region to label with an imagine modality.

Effect of additional substitutions in position 6 on PAR4-AP-mediated signalling responses. Through our initial investigations into position 6 substitutions, we observed that replacement of phenylalanine^6 with either alanine, D-phenylalanine, or *N*-methyl-phenylalanine had varied effects on PAR4-AP potency and efficacy. Interestingly, while signalling was modulated, we did not observe a loss of activity with any of these substitutions. Previous investigations into position 6 substitutions in the native tethered ligand (GYPGQV) revealed the requirement for an aromatic residue, as substitution with non-aromatic residues such as lysine or ornithine significantly decreased or abolished the activity of the peptide (Faruqi et al., 2000). In addition to being an aromatic residue, phenylalanine is a hydrophobic

390   residue so we investigated the impact of altering the hydrophobicity/hydrophilicity and stericity
391   while maintaining an aromatic residue (Fig. 7, Table 1).
392
First, we investigated the effect of polar changes that decrease hydrophobicity. Substitution of
phenylalanine[6] with tyrosine results in the addition of a hydroxyl group to position 6
(AYPGK**Y**-NH$_2$). We observed partial agonism of calcium signalling with no significant
change in potency compared to the PAR4-AP (AYPGK**Y**-NH$_2$, EC$_{50}$ = 5.2 ± 1.0 μM).
Interestingly, the peptide was able to recruit β-arrestins at least as well as the PAR4-AP (β-arr-
1 120.5 ± 12.5%; β-arr-2 109.9 ± 11.4%). Further increasing polarity through substitution with
4-fluoro-L-phenylalanine, caused significant impairment of agonist activity, both decreasing
calcium signalling and significantly decreasing β-arrestin recruitment [AYPGK-**F(4-fluoro)**-
NH$_2$, calcium EC$_{50}$ = *n.d.,* β-arr-1 25.2 ± 6.5%; β-arr-2 41.3 ± 5.9%]. The results of these two
substitutions suggest that altering the polarity, while maintaining aromaticity, in position 6
reduces efficacy (Hoffmann et al., 2019). Since decreasing hydrophobicity did not increase
PAR4-AP activity, we next investigated alterations that increase stericity and hydrophobicity.
Substitution with 4-methyl-L-phenylalanine reduced the efficacy of the PAR4-AP to a partial
agonist for both calcium signalling and β-arrestin recruitment [AYPGK-**F(4-Me)**-NH$_2$ calcium
EC$_{50}$ = 9.4 ±3.0 μM, β-arr-1 56.4 ± 1.2%; β-arr-2 63.2 ± 1.1%]. Given these results, we wanted
to assess the pharmacological impact of substituting phenylalanine with (1-naphthyl)-L-alanine
(1Nal) and (2-naphthyl)-L-alanine (2Nal) which not only increase hydrophobicity but also have
a modest increase in steric bulk and π-stacking potential. Substitution of phenylalanine[6] with
(1-naphthyl)-L-alanine (1Nal; AYPGK-**1Nal**-NH$_2$) generated a partial agonist for both calcium
signalling (EC$_{50}$ = 9.6 ±1.9 μM) and β-arrestin recruitment (β-arr-1 77.5 ± 2.3%; β-arr-2 64.1
± 2.0%) compared to PAR4-AP. Notably, substitution to (2-naphthyl)-L-alanine (2Nal;
AYPGK-**2Nal**-NH$_2$), which differs in orientation compared to (1-naphthyl)-L-alanine such that
the alkyl chain is connected to position 2 of naphthyl group (instead of position one as in 1Nal),
restored potency of the peptide to the levels observed with PAR4-AP. AYPGK-**2Nal**-NH$_2$ was
an equipotent agonist for calcium signalling (EC$_{50}$ = 11.1 ± 2.3 μM) and β-arrestin recruitment
(β-arr-1 77.0 ± 7.0%; β-arr-2 91.6 ± 6.4%). Together, these results indicate that there is a
balance needed between hydrophobicity, stericity, and side chain positioning in the C-terminal
residue.
421
Probing antagonism of PAR4-AP-stimulated calcium signalling by calcium signalling null
peptides. The failure of some substituted peptides to stimulate calcium signalling could reflect
either a failure to bind or a loss of efficacy. To determine if such peptides can act as antagonists
of PAR4-AP stimulated calcium signalling, we pre-incubated HEK293 cells stably expressing
PAR4-YFP prior to stimulation with PAR4-AP. Pre-treatment and competition with 100 μM
of calcium null peptides were largely unable to affect the calcium response elicited by 30 μM
PAR4-AP (AYPGKF-NH$_2$) compared to vehicle treated controls. Interestingly, two calcium
null peptides with substitution in position 3 or 4 were able to decrease calcium signals elicited
by PAR4-AP. Calcium signalling reached a mean of 20.8 ± 1.3% of A23187 (3 μM) in response
to 30 μM AYPGKF-NH$_2$. Signalling in response to AYPGKF-NH$_2$ thus was significantly

reduced following pre-treatment with AY-**Nip**-GKF-NH$_2$ (10.6 ± 0.9% of A23187; p<0.05) or
AYP-**Sar**-KF-NH$_2$ (13.0 ± 1.5% of A23187; p<0.05) (Fig. 8).
434
PAR4-dependent calcium signalling is Gα$_{q/11}$-dependent. In order to verify that PAR4
dependent calcium signalling was Gα$_{q/11}$-dependent, HEK293 cells stably expressing PAR4-
YFP were stimulated with 100 µM AYPGKF-NH$_2$ following pre-incubation with selective
Gα$_{q/11}$-inhibitor YM254890. YM254890 inhibited AYPGKF-NH$_2$-stimulated calcium
signalling in a concentration-dependent manner, with complete inhibition observed at 100 nM
YM254890. The control vehicle (0.001% DMSO) had no effect on calcium signalling (Fig. 9).
441
Calcium signalling-null peptides do not activate MAPK signalling. Activation of G-protein
and β-arrestin pathways downstream of GPCRs often converge on the activation of the p44/42
MAP kinase-signalling pathway (Jean-Charles et al., 2017). We therefore monitored
phosphorylation of p44/42 MAPK in response to the 37 AYPGKF-NH$_2$ derivative peptides.
HEK293 cells, stably expressing PAR4-YFP, were incubated with control vehicle (0.001%
DMSO), 100 µM of AYPGKF-NH$_2$, or derivative peptides (100 µM) for 10 minutes. Data were
normalized as a ratio of phosphorylated-p44/42 (p-p44/42) to total-p44/42 (p44/42).
Interestingly, we find that most peptides that were unable to stimulate calcium signalling were
also unable to activate p44/42 MAPK signalling (*i.e.* A**y**PGKF-NH$_2$, AY**p**GKF-NH$_2$, (***N*-Me-**
S)-YPGKF-NH$_2$, **Inp**-YPGKF-NH$_2$, **Sar**-YPGKF-NH$_2$, AY-**Nip**-GKF-NH$_2$, AYP-**Sar**-KF-
NH$_2$). Similarly, peptides that perform as partial agonists of Gα$_{q/11}$-mediated calcium signalling
were also partial agonists for p44/42 phosphorylation. Several peptides were able to cause a
significantly greater increase in phosphorylation compared to the parental peptide (*i.e.* **Nle**-
YPGKF-NH$_2$, AYPG**O**F-NH$_2$, AYPG-(***N$_\varepsilon$*-Me-K**)-F-NH$_2$, AYPG-**Cit**-F-NH$_2$, AYPGK**Y**-
NH$_2$, AYPGK-**1Nal**-NH$_2$,) (Fig. 10, Table 1).
457
PAR4-mediated MAPK signalling is Gα$_{q/11}$-dependent and β-arrestin independent.
In order to further probe the contribution of different pathways downstream of PAR4 to p44/42
phosphorylation, we examined the effective of blocking Gα$_{q/11}$-signalling with the specific
antagonist YM254890 and examined the role of β-arrestins using a β-arrestin-1/-2 double
knockout HEK293 cell derived by CRISPR/Cas9 mediated targeting (Supp. Fig. 3). Pre-
treatment of PAR4-YFP HEK293 cells with 100 nM YM254890 for 20 minutes abolished
PAR4-AP (30 µM) stimulated activation of MAP kinase signalling and subsequent
phosphorylation of p-p44/42 at five minutes time (Fig. 11 A.). In contrast, there was no
significant difference in the phosphorylation of p44/42 MAPK between wild-type HEK293 and
β-arrestin-1/-2 knockout HEK293 cells stimulated with 30 µM PAR4-AP for 0-90 minutes
(Fig. 11 B.).
469
470 **Homology modelling of PAR4 and *in silico* docking of AYPGKF-NH$_2$.** In order to gain a
471 better understanding of the ligand-binding pocket in PAR4 we turned to homology modelling
472 and *in silico* docking experiments. A homology model of PAR4 was generated using known
473 coordinates of thermostabilized human PAR2 crystal structure in complex with AZ8838 (2.8Å
474 resolution, 5NDD.pdb, Cheng et al., 2017). Sequences of human PAR4 were aligned with the

structure-resolved residues of PAR2 using Clustal Omega (Madeira et al., 2019; Sievers et al., 2011). We generated a homology model of wild-type PAR2 on the PAR2 structure, which contained modifications necessary for crystallization. Twenty human PAR4 homology models, using the wild-type human PAR2 model as a template, were generated in MODELLER (Webb and Sali, 2014). The PAR4 model with the lowest discrete optimized protein energy (DOPE) score was utilized for *in silico* analyses.

Docking AYPGKF-NH$_2$ to the PAR4 homology model (GalaxyPepDock; Lee et al., 2015) revealed a number of highly predicted peptide-receptor interaction sites (Fig. 12 A.; Ser[67], Tyr[157], His[229], Asp[230], Leu[232], Leu[234], Asp[235], Ala[238], Gln[242], His[306], Tyr[307], Pro[310], Ser[311], Ala[314], Gly[316], Tyr[319], and Tyr[322]). Of these, the most frequently predicted interactions were receptor residues His[229], Asp[230], and Gln[242]. In peptide-docked models, Gln[242] made contact with alanine, tyrosine, and lysine, with the most frequent prediction being with phenylalanine. His[229] was predicted to interact with lysine and phenylalanine. Asp[230] was frequently predicted to interact with lysine and phenylalanine and contacts with lysine appeared in 60% of predictions (Fig. 12 B.). Interestingly, when visualizing receptor residues predicted to interact with the peptide, we found that a large number of these residues were localized on extracellular loop 2 (ECL2) (Fig. 12 A. and B.).

Calcium signalling and β-arrestin recruitment to predicted peptide binding-site mutant-PAR4. To experimentally examine *in silico* predictions, site-directed mutagenesis was performed introducing alanine substitutions at the three most highly predicted interacting residues, His[229], Asp[230], and Gln[242]. Additionally, we generated a double mutation, PAR4[H229A, D230A], to determine if combined loss of His[229] and Asp[230] would result in additive detriment to receptor activation. We evaluated the functional consequences of these mutations by recording calcium signalling in response to AYPGKF-NH$_2$ with all mutants. We observed no calcium response in cells expressing PAR4[D230A]-YFP or PAR4[H229A, D230A]-YFP mutant receptors up to 300 μM AYPGKF-NH$_2$ (Fig. 13 A.). Similarly, peptide induced β-arrestin-1/-2 recruitment to PAR4[D230A]-YFP and PAR4[H229A, D230A]-YFP was significantly reduced compared to the wild-type PAR4. Both PAR4[H229A] and PAR4[Q242A] mutations were fully able to recruit β-arrestin-1/-2 to the same levels observed with wild-type PAR4-YFP (Fig. 13 B. and C.).

To investigate whether these sites are important for tethered ligand activation of the receptor, we stimulated cells expressing wildtype and mutant constructs with thrombin. Thrombin cleavage of the receptor reveals the native tethered ligand, GYPGQV (Xu et al., 1998). Comparable to activation with AYPGKF-NH$_2$, we observed significantly diminished calcium signalling in HEK293 cells expressing PAR4[D230A]-YFP and PAR4[H229A, D230A]-YFP mutants, even up to 10 units/mL thrombin (Fig. 13 D.). Additionally, β-arrestin-1/-2 recruitment was also impaired to PAR4[D230A] and PAR4[H229A, D230A] mutants (Fig. 13 E and F). We observed no appreciable difference in calcium signalling or β-arrestin recruitment when PAR4[H229A]-YFP and PAR4[Q242A]-YFP mutants were stimulated with thrombin, with the exception of a modest decrease in β-arrestin-1 recruitment to PAR4[Q242A]-YFP mutant receptor.

Platelet aggregation response to differential activation of PAR4 signalling. To determine the effects of targeted signalling observed with several peptides we investigated PAR4-mediated platelet aggregation in response to the β-arrestin biased, calcium signalling-null peptide, A**y**PGKF-NH2, as well as a modestly more potent calcium agonist AYPG**R**F-NH2 (compared to AYPGKF-NH2). Thrombin (1 unit/mL) and 100 μM AYPGKF-NH2 were used as controls to ensure successfully washed platelet preparations. In response to thrombin stimulation, we observed a characteristic aggregation trace resulting in 100% aggregation at 1 unit/mL. In agreement with previously published data, AYPGKF-NH2 (100 μM) stimulation resulted in approximately 50% platelet aggregation (Ramachandran et al., 2017). We observed that platelet aggregation was not stimulated by 100 μM A**y**PGKF-NH2. Interestingly, when platelet preparations were stimulated with AYPG**R**F-NH2, with which we observed a modest increase in potency in calcium signalling assays, enhanced β-arrestin recruitment, and an increased phosphorylation of p44/42, we observed platelet aggregation response comparable to levels observed with thrombin (1 unit/mL) (Fig. 14).

Discussion:

PAR4 is emerging as a novel anti-platelet drug target. Here we examine structure activity requirements for PAR4 activation of calcium signalling, β-arrestin recruitment and MAP kinase activation. PAR cleavage by a number of different enzymes has been reported. Canonically, thrombin has been described as a PAR1 and PAR4 activator while trypsin is described as a PAR2 and PAR4 activator. While PAR3 can be cleaved by thrombin, its ability to act as a signalling receptor remains ambiguous. In each case, cleavage by serine proteases occurs at an arginine-serine site which reveals a unique tethered ligand specific to each PAR (Hollenberg, 2003) - human PAR1- SFLLRNP… (Liu et al., 1991; Vu et al., 1991) , human PAR2- SLIGKV… (Al-Ani et al., 2002; Nystedt et al., 1994) and human PAR4- GYPGQV… (Kahn et al., 1998; Xu et al., 1998). While these tethered ligand sequences are diverse, they retain some highly conserved properties including a highly conserved charged residue in position five of the revealed tethered-ligand-mimicking peptide (arginine in PAR1; lysine in PAR2). Additionally, PAR2 and PAR4 have a conserved glycine in position three and valine in position six. The similarities of the proteolytically-revealed tethered ligands is observed in the optimized tethered-ligand-mimicking synthetic agonist peptides. Hydrophobic residues leucine, isoleucine, and proline are preferred in position three of PAR1, PAR2, and PAR4 agonist peptides, respectively. Additionally, there is a basic residue in position five of the tethered ligand peptide in all three receptors (arginine in PAR1 and PAR2 and lysine in PAR4 activating peptides). These conserved residue properties highlight some of the key requirements that a PAR4 agonist peptide must meet. Previous structure-activity work starting with the proteolytically revealed sequences of human (GYPGQV) and murine (GYPGKF) PAR4 identified the hexapeptide AYPGKF-NH2 or SYPGKF-NH2 to be potent and selective agonists for PAR4 (Faruqi et al., 2000). In this study we started with the hexapeptide AYPGKF-NH2, as our parental peptide agonist, and incorporated various modifications at each of the six positions in this peptide to investigate determinants of agonist-induced PAR4-coupling to Gα$_{q/11}$, MAP kinase and β-arrestin-1/-2 recruitment to guide development of biased agonists of PAR4.

Previously, it was demonstrated that substitution of alanine or serine in position one of the
GYPGKF-NH$_2$ activating peptide was sufficiently able to trigger tritiated inositol 1,4,5-
triphosphate release comparable to levels observed with thrombin activation and much higher
than release caused by the GYPGKF-NH$_2$ (Faruqi et al., 2000). Interestingly, substitution of
position 1 with threonine was reported to significantly reduce the potency of the activating
peptide (Faruqi et al., 2000). In keeping with results from previous studies, we find a
requirement for small aliphatic side chains in position 1 of PAR4 activating peptides, with
substitutions that increased bulk in this position consistently resulting in poor agonists for both
calcium signalling and β-arrestin recruitment (N-Me-A Fig. 3; N-Me-S, Aib, Inp, V, Nle – Fig
4; Table 1). Additionally, we observed that both stereochemical inversion of L-alanine to D-
alanine (D-alanine, Fig. 2) and backbone elongation (betaAla., Fig. 4) were detrimental to both
calcium signalling and β-arrestin recruitment. Additionally, we observed that capping of N-
terminal charge via acetylation significantly decreases agonist potency. Together these results
indicate that positive charge and the localization of that charge in addition to precise side-chain
positioning at position 1 are critical to effective PAR4-agonism.

The aromatic sidechain, and to a lesser extent the hydroxyl group, of the tyrosine[2] residue of
the parental peptide also proved to be important for peptide activity with alanine or D-isomer
substitutions resulting in decreased activity of all pathways investigated. These results are
consistent with previous findings that replacement of tyrosine with other aromatic residues,
such as phenylalanine, significantly reduced both specificity and potency of the activating
peptide (Faruqi et al., 2000). Interestingly, substituting L-tyrosine[2] for either D-tyrosine (Fig.
2 A., Fig. 10) or N-methyl-tyrosine (Fig. 3 A. & 10) resulted in loss of calcium and p44/42
signalling, but had more modest effects on β-arrestin recruitment (Fig, 2 B. & C., Fig. 3 B. &
C., respectively; Table 1). Also, substitution of tyrosine with either 4-fluoro-phenylalanine or
O-methyl-tyrosine did not significantly affect calcium, p44/42 phosphorylation or β-arrestin
recruitment in comparison to AYPGKF-NH$_2$ (Fig. 5 & 10; Table 1). Our findings, together
with previous reported data (Faruqi et al., 2000), suggest the necessity of an aromatic residue
in position 2 to maintain agonist potency. Further, our data reveal that backbone orientation in
position 2 is a key determinant of agonist potency. Future experiments fine-tuning the aromatic
substituents may be beneficial for peptide analogue development.

Our data reveal that alterations affecting proline conformation or glycine residue compactness
were not well-tolerated resulting in decreased calcium signalling, p44/42 MAP kinase
signalling, and β-arrestin recruitment to PAR4. Calcium signalling has previously been
reported to be negatively impacted by alterations to either of these two residues (Faruqi et al.,
2000). Both proline and glycine residues are frequently found in β-turns and these data support
the notion that the conformation (proline[3] and glycine[4]) and rigidity (proline[3]) of PAR4
agonists enabled by these residues is important for activity. To test this hypothesis further, we
made several substitutions in position 3 (proline) aimed at evaluating favoured backbone
conformations. We observed that a return to a predominantly trans-amide conformation by
substitution of proline with alanine (Fig. 1), results in decreased potency in all pathways

studied. Interestingly, this substitution did not abolish signalling but did significantly decrease
agonist potency, which suggests that the backbone conformation of proline may be necessary
for potency but not efficacy. Substitution of proline to pipecolic acid (Pip., Fig. 5), which has
a similar peptide backbone orientation to proline with a six-membered aromatic ring, decreased
β-arrestin recruitment with only a modest decrease in calcium signalling activation.
Interestingly, substitution of proline with nipecotic acid (Nip., Fig. 5) abolished calcium
signalling and p44/42 phosphorylation and also decreased β-arrestin recruitment. Nipecotic
acid is similar to pipecolic acid with the exception that nipecotic acid changes the direction of
the peptide backbone compared to that adopted by proline, while pipecolic acid favours a
similar backbone directionality and conformation similar to proline. Interestingly, we observed
a similar pharmacological profile observed with AY-**Nip**-GKF-NH$_2$ when we substitute D-
proline in position 3, which also changes the backbone orientation of the parental peptide.
These data demonstrate that the backbone conformation provided by proline in position 3 of
the parental peptide or by pipecolic acid substitution is necessary for agonism of PAR4
signalling pathways. Further, substitution of a proline[3] with alanine (AY**A**GKF-NH$_2$, Fig. 2)
significantly and detrimentally altered the pharmacological profile compared to the parental
peptide PAR4-AP, perhaps indicating that agonist competency contributed by position 3 is
largely conformational not chemical. Substitution of sarcosine or beta-alanine at position 4 also
negatively affected all signalling pathways, while substitution of glycine[4] with alanine resulted
in a rightward shift in potency with calcium signalling and a reduction of β-arrestin recruitment.
Together these substitutions reveal the necessity of a small flexible residue in position 4,
consistent with previous reports (Faruqi et al., 2000).

It has been previously demonstrated that substitution of position 5 glutamine in the native
tethered ligand (GYPGQV), with arginine or ornithine, resulted in decreased potency in the
PAR4 agonist peptide for calcium signalling pathways (Faruqi et al., 2000). However, there is
a basic residue found in position 5 of PAR1 (arginine, TFLL**R**), PAR2 (arginine, SLIG**R**L),
and PAR4 (lysine, AYPG**K**F) tethered-ligand mimicking peptides. Given the conservation of
charged residues in position 5 of PAR tethered-ligand mimicking peptides, we generated
PAR4-AP derivative peptides with lysine[5] substitutions that maintained or altered charge in
this position to determine if this is a requirement for activity at PAR4. Substitution of lysine[5]
with alanine had no appreciable effect on calcium or MAP kinase signalling, while resulting in
decreased β-arrestin recruitment (Fig. 1). Substitution of L-lysine[5] with D-lysine, however,
significantly reduced potency and efficacy in calcium signalling and β-arrestin recruitment and
did not stimulate MAP kinase signalling (Fig. 2). We continued our exploration of the position
with N_α-methyl-lysine substitution, which increases steric bulk through the addition of a methyl
group to the backbone, while also removing a hydrogen bond donor and increasing the amount
of peptide in the cis-transformation of the amide bond between position 4 and 5. We found this
substitution detrimental to calcium signalling but advantageous for β-arrestin recruitment,
resulting in saturated and comparable levels of recruitment than those observed with PAR4-AP
but at lower concentrations than PAR4-AP (Fig. 3). Thus, backbone conformation and bulk at
position 5 is important for agonism of calcium signalling and MAP kinase activation, while
perhaps being an important site for modification in the search for β-arrestin biased ligands.

We next turned our attention to analogues that resemble the side chain of lysine[5] with a specific
interest on structure-activity as well as ascertaining the importance of charge in this residue. In
contrast to what was observed with substitutions to the tethered-ligand that mimicked the native
tethered ligand, we find that both calcium signalling and β-arrestin recruitment were not
significantly impacted by substitution to ornithine, which decreases steric bulk and shortens
the distance between the backbone and the positively charged amine compared to lysine
(AYPG**O**F-NH$_2$; Fig. 6). Interestingly, however, we observed significantly increased p44/42
phosphorylation in response to AYPG**O**F-NH$_2$ stimulation (Fig. 10). This is somewhat
puzzling since we find p44/42 phosphorylation to be G$\alpha_{q/11}$-dependent (Fig. 11); however, it
is important to note that MAP kinase and other cellular signalling pathways downstream of
receptor activation are subject to amplification/deamplification processes within the cell
(Chidiac, 2016) and these observations may point to additional differences in signalling
stimulated by this peptide that needs to be examined. We further probed stericity in this position
through increasing bulk with substitution of lysine[5] with arginine (AYPG**R**F-NH$_2$).
Interestingly, this substitution also results in a delocalization of charge compared to the charge
on the lysine[5] side chain. This modification provided a similar calcium signalling profile
compared to that observed with ornithine substitution, however, resulted in a significant
increase of β-arrestin recruitment compared to the parental PAR4-AP. Interestingly, there was
no significant change in p44/42 phosphorylation like that observed with ornithine substitution.
Another peptide with increased steric bulk, N_ε-methyl-lysine substitution in position 5,
improved calcium signalling potency, modestly improved β-arrestin recruitment, and
significantly increased p44/42 phosphorylation [AYPG-(**N_ε-Me-K**)-F-NH$_2$, Fig. 6]. We further
investigated whether charge was a significant determinant of agonism by substituting lysine[5]
with citrulline (AYPG-**Cit**-F-NH$_2$), which lacks charge but retains a similar structure compared
to arginine and, to a lesser extent, lysine. We observed decreased calcium, equivalent β-arrestin
recruitment, and significantly increased p44/42 phosphorylation compared to the parental
peptide (Fig. 6 & 10). Together these data reveal that changes to side chain charge, bond
donors/acceptors, and steric bulk result in varied pharmacological responses that may be
exploited to bias signalling responses in favour of β-arrestin recruitment (AYPG**R**F-NH$_2$) or
p44/42 phosphorylation (AYPG**O**F-NH$_2$; AYPG-(**N_ε-Me-K**)-F-NH$_2$) with modest
improvements to potency in calcium signalling. Interestingly, we observed that charge is not
required for activation of some pathways as alanine and citrulline substitutions resulted in
agonism of PAR4.

Finally, our structure-activity investigation turned to position 6, phenylalanine, of the parental
peptide. Given the significant impact that alterations to position 2 (tyrosine) had on agonist
potency, we predicted that there may be a pi-pi (π-π) stacking interaction between tyrosine and
phenylalanine which is supported by the backbone kink provided by proline in position 3.
Substitution of position 6 with alanine or D-phenylalanine reduced calcium signalling potency
while reducing β-arrestin recruitment and p44/42 phosphorylation. These alterations suggest
that phenylalanine in position 6 is not essential for PAR4 activation by agonist peptide but is
important for agonist potency and full activation of all signalling pathways. Methylation of the

backbone nitrogen in phenylalanine (AYPGK-(*N*-Me-F)-NH$_2$) resulted in improved activation of calcium signalling, β-arrestin recruitment, and p44/42 phosphorylation compared to AYPGKF-NH$_2$ (Fig. 3). Addition of either methyl (AYPGK-F(4-Me)-NH$_2$; Fig. 7), fluorine (AYPGK-F(4-fluoro)-NH$_2$; Fig. 7), or hydroxyl (AYPGKY-NH$_2$; Fig. 7) off of the *para*-position of the aromatic side chain of phenylalanine was deleterious to all signalling pathways studied. Substitution of unnatural residues that increase the size of steric bulk and π-stacking potential yielded mixed results based on the orientation of the naphthyl. Substitution of phenylalanine with (1-naphthyl)-L-alanine (1Nal) increased p44/42 phosphorylation but decreased both calcium signalling and β-arrestin recruitment (Fig. 7). Changing the alkyl chain connection from position 1 to position 2 of the naphthyl (AYPGK-2Nal-NH$_2$; Fig 7) resulted in activation similar to AYPGKF-NH$_2$. With (2-naphthyl)-L-alanine showing some flexibility in the steric bulk of position 6, it would be interesting for future work to investigate increasing the π-stacking potential at position 2 while simultaneously increasing the π-stacking potential at position 6, since we hypothesize that these two residues' aromatic regions of the PAR4-AP may interact.

To determine if any of the calcium signalling-null peptides were able to act as antagonists, cells were pre-treated with calcium null peptides and subsequently stimulated with AYPGKF-NH$_2$ to see if the parental peptide was able to still elicit a signal. We observed that both AY-**Nip**-GKF-NH$_2$ and AYP-**Sar**-KF-NH$_2$ were both able to significantly decrease the calcium signal elicited by PAR4-AP (Fig. 8). Interestingly, both substitutions resulted in ablation of both calcium and p44/42 signalling as well as significantly decreasing β-arrestin recruitment. These results may indicate that targeting PAR4 with small molecules mimicking position 3 and 4 dipeptides may provide a starting place for the design of novel PAR4 antagonists. Additionally, peptides that further explore modifications to position 3 and 4 could be investigated for potent PAR4 antagonists.

In platelets, it has previously been demonstrated that PAR4-mediated calcium mobilization induces activation and aggregation (Edelstein et al., 2014; Holinstat et al., 2006). Both thrombin and AYPGKF-NH$_2$ initiated PAR4-dependent platelet aggregation to levels comparable to those previously reported in the literature in the same rat platelet system (Ramachandran et al., 2017). As was hypothesized, we observed that stimulation of washed platelet samples with a calcium signalling null, β-arrestin biased peptide, A**y**PGKF-NH$_2$ (100 μM) did not stimulate platelet aggregation (Fig. 14). Further, stimulation of washed platelets with the agonist peptide, AYPG**R**F-NH$_2$ (Fig. 6), enhanced platelet aggregation to levels comparable to those observed with thrombin activation of PAR4 (Fig. 14). AYPG**R**F-NH$_2$ stimulates calcium signalling to levels comparable to that seen with AYPGKF-NH$_2$ but triggers more β-arrestin-1 recruitment than the parental peptide. These data suggest that calcium signalling is essential for triggering platelet activation, but arrestin recruitment may also play a role.

Having ascertained some of the governing peptide residue characteristics enabling PAR4 activation and validated their signalling consequences in an *ex vivo* system, we turned our

attention to investigating the binding site of the receptor. The extracellular loops of many GPCRs have been demonstrated to be important facilitators of ligand entry into the orthosteric site, with ECL2 possessing the most diversity in sequence and structure (Venkatakrishnan et al., 2014; Woolley and Conner, 2017). Analysis of common mechanisms of GPCR ligand interaction derived from crystal structure and biochemical experiments reveal a possible role of ECL2 secondary loop in binding and facilitating entry, while the diversity of residue sequences across class A GPCRs is thought to confer their ligand specificity (Venkatakrishnan et al., 2014; Woolley and Conner, 2017). In many class A GPCRs, a disulfide bond between ECL2 and a transmembrane 3 cysteine ($Cys^{3.25}$, Ballesteros-Weinstein numbering) is thought to stabilize the extracellular topology such that ligand recognition at ECL2 and subsequent orthosteric binding can occur (Wheatley et al., 2012). All PAR receptors possess this canonical TM3 $Cys^{3.25}$ (PAR1 $Cys175^{3.25}$, PAR2 $Cys148^{3.25}$, PAR3 $Cys166^{3.25}$, PAR4 $Cys149^{3.25}$) and residues adjacent to this conserved cysteine are important for ligand binding and receptor activation. Investigations into the role of PAR ECL2 domains have previously demonstrated that all PAR receptors possess a homologous CHDxL motif (…CHDVL… - $PAR1^{C254-L258}$ and $PAR2^{C226-L230}$; …CHDVH… $PAR3^{C245-L249}$ …CHDAL; … $PAR4^{C226-L230}$) in their ECL2 (Al-Ani et al., 1999; Xu et al., 1998). Given that the activating peptides of PAR1 and PAR2 (Al-Ani et al., 1999; Lerner et al., 1996) are similar (SFLLRN, PAR1; SLIGRL, PAR2), additionally studies were conducted that revealed that ligand specificity is conferred by acidic residues distal to the CHDxL motif (PAR1, $N^{259}ETL^{261}$; PAR2 $P^{231}EQL^{234}$; (Gerszten et al., 1994; Lerner et al., 1996; Nanevicz et al., 1995). Interestingly, the equivalent human PAR4 ECL2 residues distal to the CHDAL domain, $P^{233}LDA^{236}$, share sequence homology with those distal to the human PAR2 CHDVL domain ($P^{231}EQL^{234}$), both of which contain an acidic residue. Al-Ani *et al.* (1999) demonstrated that glutamic acid residues in the rat PAR2 ECL2 (PEEVL) conferred specificity to PAR2 ligands. In previous studies, a series of aspartic acids in the PAR4 extracellular loop 2 (ECL2) and amino terminal (N)-tail were proposed to be key for thrombin binding to PAR4 (Nieman, 2008; Nieman and Schmaier, 2007; Sánchez Centellas et al., 2017). Mutations of ECL2 aspartic acid residues $D^{224}A$, $D^{230/235}A$ (double mutation), and $D^{224/230/235}A$ mutants were all sufficiently able to decrease thrombin cleavage of PAR4 and activation in yeast models (Sánchez Centellas et al., 2017). Further, the activity of PAR4 agonist peptide AYPGKF-NH$_2$ was also abolished in PAR4 mutants lacking ECL2 single mutant D224A, double mutant $D^{230/235}A$, or triple mutant $D^{224/230/235}A$ aspartic acid mutations to alanine which the authors posit indicates that these aspartic acid residues may be important for both agonist peptide and tethered ligand bindings and activation of PAR4. The $D^{230/235}A$ mutation to alanine removed two acidic residues in the highly conserved …$CHD^{230}AL$ domain (PAR1 Asp^{256}; PAR2 Asp^{228}) as well as the homologous $PLD^{235}A$… specificity conferring domain observed in PAR1 (Glu^{260}) and PAR2 (Glu^{232}) making it difficult to assess whether the loss of cleavage and activation is due to the $CHD^{230}AL$ or $PLD^{235}A$ aspartic acid.

In the present study, *in silico* docking to a homology model of PAR4 revealed 17 predicted sites of interaction with AYPGKF-NH$_2$ (Ser^{67}, Tyr^{157}, His^{229}, Asp^{230}, Leu^{232}, Leu^{234}, Asp^{235}, Ala^{238}, Gln^{242}, His^{306}, Tyr^{307}, Pro^{310}, Ser^{311}, Ala^{314}, Gly^{316}, Tyr^{319}, Tyr^{322}; Fig. 12 A.). Interestingly, these predicted sites reveal residues that have been shown to or predicted to have a role in PAR4 activation by thrombin or agonist peptide including aspartic acid residue Asp^{230}

found in the highly conserved CHDxL domain in PARs as well as Asp^{235} in the downstream region that is homologous to the specificity conferring motif found in PAR1 and PAR2 (Sánchez Centellas et al., 2017). Additionally, a previous *in silico* docking experiment revealed a role for aspartic acid and histidine in this motif in binding of AYPGKF-NH$_2$ to PAR4 (Moschonas et al., 2017). We generated mutations of the most frequently predicted interaction sites (His^{229}, Asp^{230}, Gln^{242}, His^{229}/Asp^{230}) to alanine which revealed that Asp^{230} is essential for interaction of the peptide with the receptor and activation as mutation to alanine abolished calcium signal potentiation as well as β-arrestin recruitment. Interestingly, these results were consistent with both peptide agonist AYPGKF-NH$_2$ as well as with thrombin stimulated activation of PAR4, which agrees with previous reports of the importance of these sites while also elucidating the importance of the conserved $CHD^{230}AL$ motif in PAR4 activation, consistent with the role of this domain in PAR1 and PAR2 (Al-Ani et al., 1999; Blackhart et al., 1996; Gerszten et al., 1994; Lerner et al., 1996; Nanevicz et al., 1995; Sánchez Centellas et al., 2017). Another highly predicted residue belonging to the CHDAL domain, His^{229}, when mutated to alanine did not alter the ability of either AYPGKF-NH$_2$ or the tethered ligand as revealed by thrombin to activate calcium signalling or β-arrestin recruitment pathways. Further, we observe that a double mutation of both of these sites to alanine (H229A, D230A) did not result in any additive detriment to PAR4 activation; therefore, we conclude that Asp^{230} interaction with PAR4 peptide and tethered ligand agonist performs an essential role in activation of PAR4.

Finally, in the context of platelet activation we find that our modified peptides that are able to cause a greater level of calcium signalling and β-arrestin recruitment than PAR4-AP also trigger more platelet aggregation. In our hands, PAR4-AP was never able to elicit greater than 50% of the aggregation seen with 1 unit/mL of thrombin, while the peptide AYPG**R**F-NH$_2$ was able to trigger aggregation comparable to that seen with thrombin. A role for PAR4 dependent calcium signalling is well established (Edelstein et al., 2014; French and Hamilton, 2016; Tourdot et al., 2018), and consistently our calcium signalling null peptide A**y**PGKF-NH$_2$ failed to activate platelets. A role for β-arrestin mediated PAR4 signalling in platelets has also been demonstrated (Khan et al., 2014; Ramachandran et al., 2017). Together these data suggest that calcium signalling is critical for initiating PAR4 mediated platelet aggregation, but the precise role hierarchy of the different PAR4 signalling pathways in platelet responses remains to be fully established.

In summary, these studies identify key mechanisms involved in PAR4 activation and signalling. We also present and characterize a novel toolkit of PAR4 agonist peptides that could be used to study biased signalling through PAR4 in platelets as well as other physiological systems. We further show that platelet activation by PAR4 critically depends on the $G\alpha_{q/11}$ pathway, and selective targeting of this pathway might yield a useful anti-platelet agent. Overall, our work advances our understanding of agonist binding to PAR4 and will support future efforts aimed at defining signalling contribution in homeostatic signalling and the discovery of novel therapeutics targeting PAR4.

Materials and Methods:

All chemicals were purchased from Millipore-Sigma (Burlington, Massachusetts, United States), Thermo Fisher Scientific (Hampton, New Hampshire, United States), or BioShop Canada Inc. (Burlington, Ontario, Canada) unless otherwise stated. AYPGKF-NH$_2$ was purchased from and synthesized by Genscript (Piscataway, New Jersey, United States). Thrombin from human plasma was obtained from Millipore-Sigma (Cat. # 605195, 5000 units; Burlington, Massachusetts, United States).

Molecular Cloning and Constructs. The plasmid encoding PAR4-YFP was described previously (Ramachandran et al., 2017). Plasmid DNA mutations in the predicted orthosteric binding site of PAR4 were generated using the QuikChange XL Multi Site-Directed Mutagenesis Kit (Agilent Technologies, Mississauga, ON, Canada) to generate all mutants described in this study. All constructs were verified by sanger sequencing (Robarts DNA sequencing facility, University of Western Ontario). β-arrestin-1/-2-rluc constructs were a kind gift from Michel Bouvier (Université de Montréal). PX458 vector for CRISPR/Cas9 targeting was a kind gift from Feng Zhang (Massachusetts Institute of Technology; Addgene # 48138) *Site-directed mutagenesis of predicted PAR4 receptor-peptide interaction sites.* Site-directed mutagenesis using Agilent QuikChange-XL was undertaken to generate mutations in highly predicted receptor residues to validate GalaxyPepDock model and determine key binding site residues for the native PAR4 tethered ligand and PAR4 agonist peptide AYPGKF-NH$_2$. Residues were mutated by single-nucleotide substitutions replacing native residues with alanine. Successful mutations were confirmed by DNA sequencing (London Regional Genomics Centre, London, ON).

Cell Lines and Culture Conditions. All media and cell culture reagents were purchased from Thermo Fisher Scientific (Hampton, New Hampshire, United States). HEK-293 cells (ATCC; Manassas, Virginia, United States) and CRISPR/Cas9 HEK-293 β-arrestin double knockout (β-arr knockout) cell lines were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, 1% sodium pyruvate, and 1% streptomycin penicillin. Cells stably transfected with PAR4-YFP vector was routinely cultured in the above media supplemented with 600 µg/mL G418 sulfate (Geneticin; ThermoFisher Scientific Waltham, Massachusetts, United States). Since trypsin activates PAR4, cells were routinely sub-cultured using enzyme-free isotonic phosphate-buffered saline (PBS), pH 7.4, containing 1 mM EDTA followed by centrifugation to remove PBS-EDTA and plated in appropriate culture plates for further experimentation. Cells were transiently transfected with appropriate vectors using a modified calcium phosphate transfection (Nuclease free water, 2.5M CaCl$_2$, and 2x HEPES Buffered Saline). Transient transfections were rinsed with phosphate buffered saline (PBS) 24 hours post-transfection and replaced with growth media. Transiently transfected cells were assayed or imaged at 48 hours post-transfection. *Generation of β-arrestin-1/-2 double knockout HEK-293 cells using CRISPR/Cas9.* A β-arrestin-1/-2 double knockout HEK-293 cells cell line (β-arr. knockout) was generated using CRISPR/Cas9 mediated gene targeting. Guides sequences targeted to β-arrestin-1 or β-arrestin-2 were designed using the design tool at http://crispr.mit.edu. Gene targeting guides (β-arrestin 1;

TGTGGACCACATCGACCTCG, β-arrestin 2; GCGGGACTTCGTAGATCACC, were
ligated into the PX458 vector (Kind gift from Dr. Feng Zhang, MIT, Addgene plasmid #48138),
verified by direct sequencing and transfected in HEK-293 cells via calcium phosphate
transfection (Ferguson and Caron, 2004; Graham and van der Eb, 1973). GFP expressing cells
were single-cell selected by flow cytometry (FACSAria III; London Regional Flow Cytometry
Facility) into wells of a 96-well plate. Clonal cell populations were expanded and screened by
western blot to identify successful knockout (anti-β-arrestin-1/-2 antibody, rabbit mAb D24H9;
Cell Signalling Technologies, Danvers, Massachusetts, United States; anti-β-actin antibody,
mouse mAb 8H10D10; Cell Signalling Technologies; Supplemental Figure 3).
871

General Methods for Peptide Synthesis
All reagents were purchased from Sigma-Aldrich, ChemImpex, or Thermo Fisher Scientific
and used without further purification. Peptides were synthesized using standard Fmoc-solid
phase peptide synthesis (SPPS) on Rink amide MBHA resin (256 mg, 0.1 mmol, 0.39 mmol/g)
with a Biotage® Syrowave^{TM} automated peptide synthesizer (0.4 mmol of HCTU, 0.4 mmol
of Fmoc-amino acids, 0.6 mmol of DIPEA, 1 h coupling). Some peptides also involved
additional manual synthesis (see procedures below). Following automated and manual
synthesis, peptides were cleaved off the resin by being shaken in a cleavage cocktail solution
of 95% TFA, 2.5% TIPS, and 2.5% H_2O (4 mL, 5 h), precipitated in ice cold *tert*-butyl methyl
ether, lyophilized, purified by preparative RP-HPLC, and further lyophilized to obtain a dry
powder. Purity was assessed by analytical RP-HPLC and characterized by mass spectrometry
(MS). The analytical RP-HPLC was performed on a system consisting of an analytical Agilent
Zorbax SB-C8 column (4.6 x 150 mm, 5 μm), Waters 600 controller, Waters in-line degasser,
and Waters Masslynx software (version 4.1). Two mobile phases were used; eluent A (0.1%
TFA in acetonitrile) and eluent B (0.1% TFA in MilliQ water). The flow rate was set at 1.5
mLmin^{-1} over 10 minutes with an additional 5-minute wash (95% solvent A in solvent B). A
Waters 2998 Photodiode array detector (200-800 nm) and an ESI-MS (Waters Quattro Micro
API mass spectrometer) were used to monitor the column eluate. The preparative RP-HPLC
used the same system, eluents, and detection method as mentioned above for the analytical RP-
HPLC, except that a preparative Agilent Zorbax SB-C8 column (21.2 x 150 mm, 5 μm) at a
flow rate of 20 mLmin^{-1} was used. The mass spectra for all peptides were determined in positive
mode using an ESI ion source on either a Bruker micrOToF II mass spectrometer or a Xevo
QToF mass spectrometer (Table S1). All peptides had purity >95% as determined by analytical
HPLC (see supplemental information, Figure S1).
896

Synthesis of Ac-AYPGKF-NH₂. The hexapeptide (AYPGKF) was synthesized using the
standard automated Fmoc-SPPS discussed in the general methods for peptide synthesis. While
still on resin, with all of the side chains protected, and with the N-terminal Fmoc-deprotection
completed, the N-terminus of this peptide was acetylated by being shaken in a solution of acetic
anhydride and DMF (v/v 1:4, 5 mL, 30 min). Following this, the peptide was cleaved, purified,
and analyzed as mentioned in the general methods for peptide synthesis.
903

Synthesis of Peptides Involving N-methylation of the Peptide Backbone Peptides [(*N*-Me-A)-YPGKF-NH$_2$, (*N*-Me-S)-YPGKF-NH2, A-(*N*-Me-Y)-PGKF-NH2, AYPG-(*N*$_\alpha$-Me-K)-F-NH2, and AYPGK-(*N*-Me-F)-NH2] were synthesized using abovementioned automated Fmoc-SPPS up to the desired amino acid that was *N*-methylated (e.g. KF-resin for AYPG-(*N*$_\alpha$-Me-K)-F-NH$_2$). Following Fmoc-deprotection of the resin-bound N-terminal amino acid (e.g. lysine for AYPG-(*N*$_\alpha$-Me-K)-F-NH$_2$), site selective *N*-methylation of the N-terminal amino acid was completed as follows: First, protection of the primary amine was completed to favour monomethylation by adding 2-nitrobenzenesulfonyl chloride (110.8 mg, 0.5 mmol) and 2,4,6-trimethylpyridine (132 µL, 1.0 mmol) in NMP (4 mL) to the on-resin peptide (0.1 mmol) and shaken (15 min). Following completion of the protection reaction, the resin was washed thrice with NMP (3X 5 mL) and dry THF (3X 5 mL). Second, *N*-methylation was completed through Mitsunobu conditions by adding triphenylphosphine (131.2 mg, 0.5 mmol), diisopropyl azodicarboxylate (98 µL, 0.5 mmol), and methanol (41 µL, 1.0 mmol) in dry THF (4 mL) to the on-resin peptide and shaken under N$_2$ (10 min). Following completion of the *N*-methylation reaction, the resin was washed thrice with dry THF (3X 5mL) and NMP (3X 5 mL). Third, the methylated N-terminal amine was then deprotected by adding 2-mercaptoethanol (70 µL, 1.0 mmol) and 1,8-diazabicyclo(5.4.0)undec-7-ene (75 µL, 0.5 mmol) in NMP (4 mL) to the on-resin peptide and shaken (5 min). The solution turned bright yellow as expected. The solution was removed and then this third step was repeated, followed by being washed five times with NMP (5X 5 mL) and thrice with DMF (3X 5 mL). Addition of the next sequential amino acid (except for inapplicable peptides i.e. (*N*-Me-A)-YPGKF-NH$_2$ and (*N*-Me-S)-YPGKF-NH$_2$) was added manually [0.3 mmol of HATU, 0.3 mmol of Fmoc-amino acid (e.g. Fmoc-Gly-OH for AYPG-(*N*$_\alpha$-Me-K)-F-NH$_2$), 0.6 mmol of DIPEA, 4 mL of DMF, 3 h coupling]. Following completion of the sequential amino acid reaction, the resin was washed 5 times with DMF (5X 5mL) and then the next sequential amino acid was added (on applicable peptides i.e AYPG-(*N*$_\alpha$-Me-K)-F-NH$_2$ and AYPGK-(*N*-Me-F)-NH$_2$) using the automated procedure mentioned in the general methods for peptide synthesis. Following completion of the synthesis, the peptide was cleaved, purified, and analyzed as mentioned in the general methods for peptide synthesis. The amino acid *N*-methylglycine, commonly known as sarcosine, was obtained from a commercial source, and therefore, *N*-methylglycine containing peptides (i.e. Sar-YPGKF-NH$_2$ and AYP-Sar-KF-NH$_2$) were synthesized solely by the standard automated Fmoc-SPPS discussed in the general methods for peptide synthesis.

Synthesis of AYPG-(N$_\varepsilon$-Me-K)-F-NH$_2$. The hexapeptide (AYPGKF) was synthesized using the standard automated Fmoc-SPPS discussed in the general methods for peptide synthesis, except that Fmoc-Lys(Alloc)-OH was used in place of standard Fmoc-Lys(Boc)-OH. While still on resin, with all of the side chains protected, and with the N-terminal Fmoc-deprotection completed, the N-terminus of this peptide was Boc protected by adding Boc-ON (98.5 mg, 0.4 mmol) and DIPEA (70 µL, 0.4 mmol) in DMF (4 mL) and shaken (4 hours). The solution was removed and this Boc protection was repeated overnight (20 hours) to ensure reaction completion. The resin was washed thrice with DMF (3X 5 mL) and dry DCM (3X 5 mL) and was placed under N$_2$ for site selective Alloc deprotection of the on-resin lysine side chain. Phenylsilane (296 µL, 2.4 mmol) in dry DCM (2 mL) was added to the resin. Tetrakis(triphenylphosphine) palladium(0) (23.1 mg, 20 µmol) was dissolved in dry DCM (1

mL) and added to the resin. The peptide column was flushed with N_2 (2 min) followed by being
shaken (5 min). The resin was washed thrice with dry DCM (3X 5 mL). The procedure was
repeated and shaken again (30 min). The resin was washed four times with DCM (4X 5 mL)
and DMF (4X 5 mL). Following this N-terminal Boc protection and subsequent side chain
Alloc deprotection, *N*-methylation of the on-resin lysine side chain was completed through the
site selective *N*-methylation procedure mentioned above. Following completion of the
synthesis, the peptide was cleaved, purified, and analyzed as discussed in the general methods
for peptide synthesis.

SPPS Reaction Monitoring. Three methods were used to monitor SPPS reactions. Two
methods (the Kaiser Test and the small-scale resin cleavage) were used to monitor general
peptide synthesis. Three methods (the Kaiser Test, the small-scale resin cleavage, and the
Chloranil Test) were used to monitor synthesis of peptides involving *N*-methylation. For the
Kaiser Test, several resin beads were placed in a test tube followed by the addition of 42.5 mM
phenol in ethanol (50 µL), 20 µM potassium cyanide in pyridine (50 µL), and 280.7 mM
ninhydrin in ethanol (50 µL). The mixture was then heated (100 °C, 5 min); where a positive
test indicates the presence of a free primary amine. For the small-scale resin cleavage method,
several beads and cleavage cocktail (500 µL) are shaken, worked up as per the full cleavage of
the resin procedure described above, and the desired peptide was confirmed through HPLC-
MS. For the Chloranil Test, several resin beads were placed in a test tube followed by the
addition of 357.8 mM acetaldehyde in DMF (50 µL) and 81.3 mM *p*-chloranil in DMF (50 µL)
and allowed to stand at room temperature (5 min); where a positive test indicates the presence
of a free secondary amine.

Calcium Signalling Assay. HEK293 cells or HEK293 cells stably expressing PAR4-YFP were
loaded with calcium sensitive dye (Fluo-4 NW, F36206, Life Technologies, Thermo Fisher)
and aliquoted into cuvettes with Hanks buffered salt solution (HBSS containing Ca^{2+} and
Mg^{2+}). Samples were excited at 480nm and emission recorded at 530nm. Baseline fluorescence
from PAR4-YFP excitation remains constant throughout experiment. Following baseline
recording, agonist induced changes in fluorescence was recorded. Change in fluorescence from
baseline to maximum response at a given agonist concentration (delta) were expressed as a
percentage of maximum cellular response elicited by calcium ionophore (A23187, Sigma-
Aldrich). *Calcium Signalling Assay with PAR4 orthosteric site-directed mutants*. HEK293 cells
were transiently transfected with 2µg of wild-type YFP tagged PAR4 or PAR4 mutant
receptors (described above) using the calcium phosphate transfection protocol as described
above. Calcium signalling was investigated as described above for the wild type receptor.

β-arrestin Recruitment - Bioluminescence Resonance Energy Transfer (BRET). HEK293
cells were co-transfected with PAR4-YFP (2µg) or mutant PAR4-YFP and β-arrestin-1 or -2-
renilla luciferase (rluc, 0.2µg) using modified calcium phosphate transfection. Cells were re-
plated (24 hours post-transfection) into white 96-well plates (Corning). At 48-hours post-
transfection, agonist-stimulated β-arrestin recruitment to PAR4 was detected by measuring
BRET following 20 minutes of agonist stimulation. Coelenterazine (5µM; Nanolight

Technology, Pinetop, AZ) was added ten minutes prior to BRET collection using a Berthold
Mithras LB 940 plate reader.
993
Testing Peptide library specificity for PAR4 Previous studies have pointed to the possibility
of PAR agonist peptides cross activating other members of this family. For example, the PAR1
tethered ligand peptide SFLLRN is able to activate PAR2 and substitution of a phenylalanine
residue in certain PAR4 agonist peptides confers PAR1 activity (Boitano et al., 2011; Faruqi
et al., 2000; Lerner et al., 1996). Activating peptides typically require an aromatic sidechain in
position 2 (F in PAR1) and we wondered if substitution of tyrosine for phenylalanine may
decrease selectivity at PAR4 and increase crosstalk with PAR1. We examined whether
substitutions made to AYPGKF-NH_2 affected their selectivity using a calcium signalling assay
in HEK-293 cells that endogenously express PAR1 and PAR2. None of the peptides tested
elicited a signalling greater than 5% of the calcium ionophore A23187 showing that they retain
selectivity towards PAR4 (Supp. Fig. 4). Additionally, we investigated specificity using BRET
for β-arrestin-2 recruitment to cells expressing PAR4-YFP or PAR2-YFP. None of the peptides
were able to induce β-arrestin recruitment in PAR2-YFP expressing HEK-293 cells (NET
BRET of 0.05 and over) with the exception of AYPGK-1Nal-NH_2 (Supp. Fig. 5).
1008
PAR4-stimulated Mitogen-activated Protein Kinase (MAPK) signalling <u>***HEK-293 and***</u>
<u>***CRISPR/Cas9 β-arrestin-1/-2 double knockout HEK-293.***</u> Agonist-stimulated mitogen-
activated protein kinase (MAPK) signalling in CRISPR/Cas9 HEK-293 β-arrestin double
knockout cells (β-arr. knockout) and wild-type HEK-293 cells transfected with PAR4-YFP
were analysed by western blot. Cells were placed in serum free media (serum-free DMEM,
Gibco) for 3 hours before beginning experiment. Cells were then stimulated with 30μM
AYPGKF-NH_2 for varying time periods (0, 2, 5, 10, 20, 30, 60, 90 minutes) at 37 ˚C. Treated
cells were placed on ice following stimulating period. Total protein was extracted adding kinase
lysis buffer with phosphatase and protease inhibitors (20 mM Tris-HCl, pH 7.5, 100 mM NaCl,
10 mM $MgCl_2$, 1 mM EDTA, 1 mM EGTA, 0.5% NP-40, 2.5 mM sodium pyrophosphate, 1
mM b-glycerophosphate, 1 mM Na_3VO_4, 25 mM NaF, 1 mg/mL leupeptin, 1 mg/m aprotinin,
1 mM phenylmethylsulfonyl fluoride, and 1mM dithiothreitol) for 20 minutes and cell
membranes were cleared by centrifugation (13,300g for 3 minutes). Protein concentration<u>s</u>
were measured using the Pierce BCA Protein Assay (Thermo Fisher Scientific). Protein
samples were heat-denatured at 98°C for 8 minutes in denaturing Laemmli buffer (containing
2-mercaptoethanol) and resolved on 4–12% gradient Invitrogen Bolt Bis-Tris Plus gels
(Thermo Fisher Scientific). The resolved proteins were transferred via wet transfer in buffer
(1X Tris-glycine, 20% methanol) to a polyvinylidene difluoride membrane and blocked in
TBST buffer (TBS with 0.1% (v/v) Tween-20) supplemented with 1% ECL Advance Blocking
Agent (GE Healthcare, Waukesha, WI) for 40 minutes at room temperature. p44/42
(Thr202/Tyr204) phosphorylation was detected with specific antibodies (Cell Signalling
Technology, Danvers, MA; diluted 1:1000 in TBST) overnight at 4°C. phospho-p44/42
immunoreactivity was detected using the horseradish peroxidase-conjugated anti-rabbit
secondary antibody (Cell Signalling Technology; 1:10,000 in TBST for 1 hour). After washing
the membrane with TBST (3 x 10-minute washes), the peroxidase activity was detected with

the chemiluminescence reagent ECL Advance (GE Healthcare) on iBright image Station (Invitrogen, Burlington, ON). Membranes were then stripped with blot stripping buffer (Thermo Fisher Scientific) at room temperature and blocked in TBST with 1% ECL Advance Blocking Agent before incubation with the p44/42 (1:1000 in TBST with 1% ECL Advance Blocking Agent) overnight at 4°C. Following incubation with primary antibody, blots were rinsed in TBST and incubated with appropriate horseradish peroxidase -conjugated anti-rabbit or anti-mouse secondary antibody (1:10,000 in TBST for 1 hour). Membranes were washed and imaged as previously described. Band intensities representing activated MAPK were quantified using FIJI (Schindelin et al., 2012). Phospho-kinase levels were normalized for differences in protein loading by expressing the data as a ratio of the corresponding total-p44/42 signal. Data are expressed as a fold increase over unstimulated baseline control. *PAR4-stimulated MAPK signalling in HEK-293 +/- YM254890.* Agonist-stimulated MAPK signalling in wild-type HEK-293 cells transfected with PAR4-YFP were analysed by western blot. Cells were placed in serum free media (serum-free DMEM, Gibco) for 3 hours before beginning experiment. Prior to agonist stimulation cells were treated with either control (0.001% DMSO) or YM254890 (100nM) for 20 minutes. Cells were then stimulated with 30μM AYPGKF-NH$_2$ for varying time periods (0, 2, 5, 10, 20, 30, 60, 90 minutes) at 37 degrees Celsius. Cells were lysed, protein quantified, and samples run and imaged as previously described. *MAPK signalling in PAR4-YFP stable cells with peptide library.* Agonist-stimulated mitogen-activated protein kinase (MAPK) signalling in HEK-293 cells stably expressing PAR4-YFP were analysed by western blot following stimulation with PAR4 peptide agonists. Cells were placed in serum free media (serum-free DMEM, Gibco) for 3 hours before beginning experiment. Cells were stimulated with either control (no agonist), AYPGKF-NH$_2$ or agonist peptides described at 100μM concentration. Cells were stimulated for 10 minutes with agonist and lysed, quantified, and blot as previously described.

Preparation of washed platelets and light-transmission aggregometry. Blood was drawn from the abdominal aorta or heart of Sprague-Dawley rats into trisodium citrate (3.4% w.v). The blood was centrifuged at 200g for 15 minutes at 22 degrees Celsius to isolate platelet-rich plasma (PRP). PRP was transferred to a 14 mL polypropylene round-bottom tube (Falcon) and centrifuged at 1800 rotations per minute (rpm.; 594g) for 10 minutes at 22˚C to pellet platelets and obtain platelet-poor plasma (PPP). PPP was decanted and used for 100% light transmission control. To obtain washed platelet sample, platelets were washed twice in Tyrode's buffer (12 mM NaHCO$_3$, 127 mM NaCl, 5 mM KCl, 0.5 mM NaH$_2$PO$_4$, 1 mM MgCl$_2$, 5 mM glucose, and 10 mM HEPES; Ramachandran et al., 2017) and resuspended to a volume equivalent to that of the initial blood draw. Washed platelet sample was equilibrated for one hour at ambient temperature following the addition of 1M CaCl$_2$ (1 μl/mL). 400 μl of this washed platelet suspension was used for light transmission aggregometry on a platelet aggregometer (Model 700; Chrono-Log Corp, Havertown, PA).

Structural homology modelling of PAR4. Residues resolved in the PAR2 crystal (5NDD.pdb) were used to generate a homology model of wild-type human PAR2. Homology model of wild-type human PAR2 was then aligned with the wild-type human PAR4 amino acid sequence [National Center for Biotechnology Information (NCBI) accession number NM_003950) using

Clustal Omega (Madeira et al., 2019; Sievers et al., 2011). Only residues resolved in the crystal
structure of PAR2 were used as template coordinates for PAR4 homology modelling. Twenty
human PAR4 homology models were generated using the human PAR2 crystal structure
coordinates (5NDD.pdb; Cheng et al., 2017) as a template in MODELLER (version 9.16, Webb
and Sali, 2017). The model with the lowest DOPE score was taken as the best structure and
visualized using PyMOL (version 1.7.4, Schrödinger, LLC). Electrostatic surface potential was
calculated after assignment of partial charges using PDB2PQR and using the Adaptive Poisson-
Boltzmann Solver plugin in PyMOL (Dolinsky et al., 2004).

1086

In silico docking of AYPGKF-NH$_2$ to the PAR4 homology model. *in silico* docking of PAR4
agonist peptide, AYPGKF-NH$_2$, to PAR4 homology model was undertaken using
GalaxyPepDock (Lee et al., 2015). The lowest energy PAR4 homology model PDB file was
loaded into GalaxyPepDock along with a text file of the PAR4 agonist peptide sequence which
GalaxyPepDock uses to predict a peptide structure for docking analyses (Ala-Tyr-Pro-Gly-
Lys-Phe; amidation is not included in the structure prediction). Ten models of PAR4-AP
docking were generated. Frequency of receptor-peptide residue interactions, taken as atom-
atom distances of less than 3 Angstroms, was quantified manually to determine highly recurrent
interactions.

1096

Data analysis and statistical tests. Data are shown as mean ± SEM. Calcium signalling and β-
arrestin-1/-2 recruitment concentration effect curves were calculated using nonlinear regression
curve fitting (three parameter; Prism7). EC$_{50}$ values are reported (Table 1) where signal has
saturated. Where signalling is not saturated EC$_{50}$ is reported for calcium as not determined
(*n.d.*). The F-statistic calculated to compare EC$_{50}$ values was calculated using the extra sum-
of-squares analysis (Kim et al., 2019). For table 1, values comparing percentage of signalling
or recruitment at 100 μM to values obtained from the parental PAR4-AP, AYPGKF-NH$_2$, are
shown. Significance was determined for comparisons with 100 μM AYPGKF-NH$_2$ and p44/42
phosphorylation by one-way analysis of variance (ANOVA). Western blot protein
densitometry was quantified using FIJI. Transformation and analysis were conducted on raw
data and significance is indicated by '*' ($p < 0.05$).

1108

1109 ***Acknowledgements and Funding.*** These studies were funded by a Canadian Institutes of
1110 Health Research (CIHR) grant (RR) and grants from the Natural Sciences and Engineering
1111 Research Council of Canada (LL and PS). PT is the recipient of a doctoral QEII Graduate
1112 Scholarships in Science and Technology. JCL is the recipient of a Canada Graduate
1113 Scholarship and an Ontario Graduate Scholarship.

1114

1115 ***Competing interests.*** The authors declare that no competing interests exist.

1116

	Peptide (alterations shown in red)	Ca^{2+} EC$_{50}$ (μM)	Ca^{2+} Max. (% of A23187)	% of AYPGKF-NH$_2$ Response (100μM)			p-p44/42 (fold over baseline; 100 μM)
				Ca^{2+} signaling	β-Arr-1 recruitment	β-Arr-2 recruitment	
PAR4-AP	**AYPGKF-NH$_2$**	12.1 ± 3.1	27.4 ± 1.7	100.0 ± 8.2	100.0 ± 3.9	100 ± 3.5	5.9 ± 1.5
Alanine	A**A**PGKF-NH$_2$	*n.d.*	51.8 ± 25.4	52.9 ± 12.4*	47.3 ± 3.6*	46.4 ± 3.2*	0.8 ± 0.1
	AY**A**GKF-NH$_2$	63.7 ± 20.8*	29.7 ± 3.4	75.4 ± 11.1	48.5 ± 7.9*	43.4 ± 7.1*	1.2 ± 0.3
	AYP**A**KF-NH$_2$	21.2 ± 6.6	22.4 ± 1.9	72.2 ± 6.3	56.4 ± 5.6*	54.6 ± 5.1*	1.0 ± 0.2
	AYPG**A**F-NH$_2$	9.8 ± 2.9	25.7 ± 1.8	96.7 ± 17.8	59.3 ± 7.2*	70.8 ± 6.5	6.5 ± 2.7
	AYPGK**A**-NH$_2$	17.3 ± 5.0	20.0 ± 1.5	69.4 ± 4.1	47.9 ± 3.8*	52.6 ± 3.5*	3.8 ± 2.2
D-isomer	**a**YPGKF-NH$_2$	*n.d.*	19.2 ± 4.8	24.0 ± 4.4*	48.7 ± 2.6*	47.5 ± 2.4*	0.8 ± 0.2
	A**y**PGKF-NH$_2$	*n.d.*	1.1 ± 0.5	3.9 ± 2.8*	50.7 ± 4.6*	46.4 ± 4.2*	0.6 ± 0.1
	AY**p**GKF-NH$_2$	*n.d.*	1.2 ± 0.4	6.1 ± 3.4*	45.2 ± 10.5*	38.1 ± 9.5*	0.7 ± 0.2
	AYP**g**kF-NH$_2$	*n.d.*	16.9 ± 5.3	25.2 ± 5.2*	47.4 ± 1.0*	36.5 ± 0.9*	1.0 ± 0.3
	AYPGK**f**-NH$_2$	22.2 ± 4.7	21.9 ± 1.2	76.7 ± 5.2	80.3 ± 6.6	74.1 ± 6.0	6.5 ± 2.9
***N*-methyl.**	**(*N*-Me-A)**-YPGKF-NH$_2$	*n.d.*	15.0 ± 7.5	13.3 ± 1.3*	119.2 ± 11.9	80.6 ± 10.8	1.3 ± 0.3
	A-**(*N*-Me-Y)**-PGKF-NH$_2$	*n.d.*	2.7 ± 0.2	9.8 ± 2.9*	71.4 ± 10.8	61.5 ± 9.8*	2.2 ± 0.2
	AYP-**Sar**-KF- NH$_2$	*n.d.*	2.4 ± 0.3	8.8 ± 1.5*	35.9 ± 4.3*	36.1 ± 3.9*	1.4 ± 0.2
	AYPG-**(*N*α-Me-K)**-F-NH$_2$	*n.d.*	9.6 ± 1.9	18.3 ± 1.5*	126.7 ± 18.6	77.1 ± 16.9	1.8 ± 0.9
	AYPGK-**(*N*-Me-F)**-NH$_2$	7.6 ± 1.9	26.3 ± 1.4	99.4 ± 7.8	136.9 ± 18.0*	96.6 ± 16.9	15.7 ± 4.6
N-terminal	**Ac**-AYPGKF-NH$_2$	*n.d.*	11.0 ± 3.3	49.9 ± 16.5*	47.3 ± 3.1*	45.5 ± 2.8*	1.2 ± 0.4
Position 1	**V**YPGKF-NH$_2$	22.9 ± 8.3	16.8 ± 1.7	48.5 ± 4.6*	73.6 ± 13.0	55.2 ± 11.8*	2.3 ± 0.7
	Nle-YPGKF-NH$_2$	*n.d.*	1.1 ± 0.1	4.0 ± 1.1*	40.2 ± 9.7*	36.5 ± 8.8*	21.4 ± 7.0*
	Sar-YPGKF- NH$_2$	*n.d.*	1.6 ± 0.6	9.7 ± 8.5*	35.6 ± 2.0*	39.8 ± 1.8*	1.6 ± 1.1
	Aib-YPGKF-NH$_2$	50.2 ± 15.3*	25.1 ± 2.5	72.9 ± 12.0	38.6 ± 3.9*	37.3 ± 3.6*	7.8 ± 3.5
	(*N*-Me-S)-YPGKF-NH$_2$	*n.d.*	2.2 ± 0.5	8.8 ± 1.8*	78.1 ± 6.9	51.7 ± 6.3*	1.8 ± 0.4
	betaAla-YPGKF- NH$_2$	*n.d.*	1.9 ± 0.4	10.3 ± 3.9*	25.7 ± 4.8*	31.8 ± 4.4*	9.1 ± 3.4
	Inp-YPGKF-NH$_2$	*n.d.*	3.4 ± 0.8	18.9 ± 7.3*	32.0 ± 3.3*	26.9 ± 3.0*	1.3 ± 0.3
Position 2	A-**Tyr(Me)**-PGKF-NH$_2$	15.7 ± 4.0	25.7 ± 1.6	91.1 ± 10.1	77.9 ± 3.8	85.8 ± 3.4	12.7 ± 5.8
	A-**F-(4-fluoro)**-PGKF-NH$_2$	19.9 ± 5.4	35.1 ± 2.5	105.4 ± 9.0	72.1 ± 9.0	86.9 ± 8.1	10.1 ± 3.9
Position 3	AY-**Pip**-GKF- NH$_2$	30.7 ± 4.9	23.3 ± 1.1	74.2 ± 4.8	38.8 ± 3.0*	44.0 ± 2.7*	1.3 ± 0.3
	AY-**Nip**-GKF- NH$_2$	*n.d.*	2.5 ± 0.3	10.0 ± 1.8*	32.9 ± 4.9*	30.9 ± 4.5*	1.0 ± 0.3
Position 4	AYP-**betaAla**-KF- NH$_2$	*n.d.*	9.7 ± 2.4	19.8 ± 5.8*	42.5 ± 1.8*	24.9 ± 1.6*	1.8 ± 0.6
Position 5	AYPG**O**F- NH$_2$	4.2 ± 1.1	20.7 ± 1.1	77.3 ± 7.9	121.7 ± 8.5	112.1 ± 7.7	23.4 ± 7.2*
	AYPG**R**F-NH$_2$	4.0 ± 1.1	21.3 ± 1.2	82.0 ± 8.3	154.2 ± 10.3*	140.1 ± 9.4*	3.0 ± 0.3
	AYPG-**(*N*ε-Me-K)**-F-NH$_2$	6.7 ± 1.9	23.8 ± 1.5	88.9 ± 9.3	122.5 ± 12.4	116.7 ± 11.2	20.5 ± 5.3*
	AYPG-**Cit**-F-NH$_2$	5.4 ± 0.8	18.2 ± 0.6	65.5 ± 1.3	94.1 ± 10.1	87.7 ± 9.2	25.1 ± 7.5*
Position 6	AYPGK**Y**-NH$_2$	5.2 ± 1.0	16.9 ± 0.7	61.2 ± 4.3*	120.5 ± 12.5	109.9 ± 11.4	20.2 ± 9.7*
	AYPGK-**F(4-Me)**-NH$_2$	9.4 ± 3.0	17.8 ± 1.3	62.0 ± 5.5*	56.4 ± 1.2*	63.2 ± 1.1*	0.9 ± 0.4
	AYPGK-**F(4-fluoro)**-NH$_2$	*n.d.*	*n.d.*	16.5 ± 10.2*	25.2 ± 6.5*	41.3 ± 5.9*	1.1 ± 0.5
	AYPGK-**1Nal**-NH$_2$	9.6 ± 1.9	18.7 ± 0.9	65.4 ± 7.4	77.5 ± 2.3	64.1 ± 2.0*	21.3 ± 10.8*
	AYPGK-**2Nal**-NH$_2$	11.1 ± 2.3	25.3 ± 1.3	93.9 ± 4.2	77.0 ± 7.0	91.6 ± 6.4	4.3 ± 1.7

Table 1. Summary table of all calcium, β-arrestin, and p44/42 data. Calcium data is displayed as EC$_{50}$ or not determined (*n.d.*) if EC$_{50}$ could not be calculated. Maximum calcium response achieved with each peptide is shown as a percentage of the maximum calcium response achievable in a given experiment revealed by calcium ionophore (% of A23187 calcium ionophore). Statistically significant shifts in EC$_{50}$ compared to PAR4-AP, AYPGKF-NH$_2$, were found using the F-statistic (*$p < 0.05$). Given that saturation did not always occur, all peptides are compared to the parental PAR4-AP, AYPGKF-NH$_2$, and shown as a percent of

the response obtained with PAR4-AP at a given concentration (100 µM). Finally, fold increase

over unstimulated baseline is shown for phosphorylation of p44/42 MAP kinase activation.

Statistical significance for comparisons at 100 µM and p44/42 phosphorylation are calculated

using one-way ANOVA of response compared to PAR4-AP ($*p < 0.05$).

Figure 1. Alanine
substitutions in positions 2-6
significantly decreases β-
arrestin recruitment and
differentially affect calcium
signalling compared to PAR4-
AP. (A) Agonist-stimulated
calcium signalling. Calcium
signalling is shown as a
percentage of the maximum
calcium signalling obtained by
application of calcium
ionophore (A23187, 3 μM). (*n*
= *3-7*) **Agonist-stimulated**
recruitment of β-arrestin-1
(B) and β-arrestin-2 (C) to
PAR4. Concentration effect
curves of BRET values from
cells treated increasing
concentrations of peptide are
presented as normalized net
BRET ratio. (*n = 3*).

Figure 2. D-isomer amino acid
substitutions in positions 1, 2,
3, and 5 of the PAR4-AP,
significantly reduces potency
in calcium signalling and β-
arrestin recruitment assays.
(A) Agonist-stimulated
calcium signalling. Calcium
signalling is shown as a
percentage of the maximum
calcium signalling obtained by
application of calcium
ionophore (A23187, 3 μM). (*n =*
3-4) **Agonist-stimulated**
recruitment of β-arrestin-1 (B)
and β-arrestin-2 (C) to PAR4.
Concentration effect curves of
BRET values from cells treated
increasing concentrations of
peptide are presented as
normalized net BRET ratio. (*n =*
3).
1172

Figure 3. Peptides with *N*-methylation of either alanine[1] or lysine[5] do not initiate calcium signalling but enhance β-arrestin recruitment compared to PAR4-AP. *N*-methylation of phenylalanine[6] enhances β-arrestin recruitment while signalling calcium equipotently in comparison to PAR4-AP. (A) Agonist-stimulated calcium signalling. Calcium signalling is shown as a percentage of the maximum calcium signalling obtained by application of calcium ionophore (A23187, 3 μM). (*n = 3-7*) **Agonist-stimulated recruitment of β-arrestin-1 (B) and β-arrestin-2 (C) to PAR4.** Concentration effect curves of BRET values from cells treated increasing concentrations of peptide are presented as normalized net BRET ratio. (*n = 3*).

Figure 4. Calcium signalling and
β-arrestin recruitment to PAR4 in
response to AYPGKF-NH₂ or
peptides with N-terminal
acetylation or additional
substitutions in position 1
(alanine). (A) Agonist-stimulated
calcium signalling. Calcium
signalling is shown as a percentage
of the maximum calcium signalling
obtained by application of calcium
ionophore (A23187, 3 μM). (*n = 3-*
7) **Agonist-stimulated recruitment**
of β-arrestin-1 (B) and β-arrestin-
2 (C) to PAR4. Concentration effect
curves of BRET values from cells
treated increasing concentrations of
peptide are presented as normalized
net BRET ratio. (*n = 3*).
1215

Figure 5. Calcium and β-arrestin recruitment to PAR4 in response to AYPGKF-NH$_2$ or peptides with additional substitutions in tyrosine[2], proline[3], or glycine[4]. (A) Agonist-stimulated calcium signalling. Calcium signalling is shown as a percentage of the maximum calcium signalling obtained by application of calcium ionophore (A23187, 3 μM). (*n = 3-7*) **Agonist-stimulated recruitment of β-arrestin-1 (B) and β-arrestin-2 (C) to PAR4.** Concentration effect curves of BRET values from cells treated increasing concentrations of peptide are presented as normalized net BRET ratio. (*n = 3*).

Figure 6. Calcium and β-arrestin
recruitment to PAR4 in response
to AYPGKF-NH$_2$ or peptides with
additional substitutions in lysine[5].
(A) Agonist-stimulated calcium
signalling. Calcium signalling is
shown as a percentage of the
maximum calcium signalling
obtained by application of calcium
ionophore (A23187, 3 μM). (*n = 3-*
7) **Agonist-stimulated recruitment**
of β-arrestin-1 (B) and β-arrestin-
2 (C) to PAR4. Concentration effect
curves of BRET values from cells
treated increasing concentrations of
peptide are presented as normalized
net BRET ratio. (*n = 3*).
1251

Figure 7. Calcium and β-arrestin
recruitment to PAR4 in response
to AYPGKF-NH₂ or peptides with
additional substitutions in
phenylalanine[6]. (A) Agonist-
stimulated calcium signalling.
Calcium signalling is shown as a
percentage of the maximum calcium
signalling obtained by application of
calcium ionophore (A23187, 3 μM).
(*n = 3-7)* **Agonist-stimulated**
recruitment of β-arrestin-1 (B)
and β-arrestin-2 (C) to PAR4.
Concentration effect curves of
BRET values from cells treated
increasing concentrations of peptide
are presented as normalized net
BRET ratio. (*n = 3*).

Figure 8. PAR4-AP (AYPGKF-NH₂) stimulated calcium signalling responses following pre-treatment with calcium signalling null peptides. Calcium signalling (Fluorescence Em: 530) in HEK293 cells stably expressing PAR4-YFP were recorded following 30 μM AYPGKF-NH₂ addition after 30-minute pre-treatment incubation with 100 μM of calcium signalling-null peptides. Data are shown as a percentage of maximum response obtained in cells treated with the calcium ionophore A23187 (3 μM). Agonist peptide concentrations are in M. Pre-treatment with calcium-null peptides did not affect AYPGKF-NH₂-stimulated calcium signalling in most cases with the exception of AY-Nip-GKF-NH₂ and AYP-Sar-KF-NH₂, which significantly decreased AYPGKF-NH₂ stimulated calcium signalling compared to vehicle treated control. (*p > 0.05; *n = 3*)

Figure 9. PAR4-mediated calcium signalling is Gα$_{q/11}$-dependent. HEK293 cells stably expressing PAR4 were pre-treated with either DMSO vehicle control (0.001%) or increasing concentration of the selective Gα$_{q/11}$ inhibitor YM254890. Calcium signalling following stimulation with AYPGKF-NH$_2$ (100μM) was recorded. PAR4-mediated calcium signalling was blocked by YM254890 at concentrations greater than 100 nM. (*n = 4*)

Figure 10. PAR4-mediated phosphorylation of p44/42(ERK) is attenuated upon
stimulation of PAR4 with calcium-null peptide. HEK293 cells stably expressing PAR4-YFP
were stimulated with AYPGKF-NH$_2$ (30 µM) or peptide (30 µM) from library of compounds
and Western blot for p44/42 phosphorylation (p-p44/42). Data are shown as normalized fold
increase over unstimulated control cells. Dashed baseline shows mean of unstimulated baseline
for comparison. Dotted baseline shows mean response achieved following AYPGKF-NH$_2$
stimulation. (# $p > 0.05$ AYPGKF-NH$_2$ compared to untreated control; *$p > 0.05$ AYPGKF-
NH$_2$ compared to other peptide agonists; one-way ANOVA). Peptides that were unable to
stimulate calcium signalling were also generally unable to stimulate phosphorylation of p44/42.
Interestingly, several peptides were able to significantly increase phosphorylation of p44/42
compared to AYPGKF-NH$_2$. (Representative blots shown in supplemental figure S2; $n = 4$-5)

Figure 11. PAR4-mediated phosphorylation of p44/42 (ERK) is Gα$_{q/11}$-dependent and β-arrestin-independent. To evaluate contribution of Gα$_{q/11}$ and β-arrestin to p44/42 phosphorylation, p-p44/42 was monitored with western blot following stimulation with AYPGKF-NH$_2$ (30 μM) for various time points. To evaluate contribution of Gα$_{q/11}$-signalling cells were incubated with vehicle control (0.001% DMSO) or Gα$_{q/11}$-inhibitor, YM254890 (100 nM) prior to agonist stimulation (A). To determine contribution of β-arrestin-signalling to p44/42 phosphorylation, HEK293 and β-arrestin-1/2 CRISPR knockouts were assayed following agonist stimulation (B). Values are expressed as fold increase over unstimulated 0-minute time [(p-p44/22 / total p44/42) / (baseline p-p44/42 / total p44/42)]. Data are analysed with two-way ANOVA (*$p > 0.05$). We find that Gα$_{q/11}$ inhibition with YM254890 results in a significant reduction of p44/42 phosphorylation. We observe no statistical difference in ERK activation between HEK293 and β-arrestin-1/2 knockouts HEK293 cells. (Representative blots shown; *n = 4*)

Figure 12. (A) Frequency table of PAR4/AYPGKF-NH$_2$ interactions as predicted by *in silico* peptide docking (GalaxyPepDock). Numbers represent the frequency with which an atom in a PAR4-AP residue was predicted to interact with an atom of a PAR4 residue. **(B) Representation of the most highly predicted AYPGKF-NH$_2$ binding site (Asp^{230}, shown as sticks) interacting (shown as yellow dashed lines) with position 5 (Lys^5; shown as ball and sticks) of AYPGKF-NH$_2$.** Interaction of Lys^5 and Asp^{230} was highly predicted in 7/10 models (B). Alignment of the receptor backbone (shown as transparent cartoon) reveals similar binding poses adopted by the side chain position of Asp^{230} and the position of Lys^5 of AYPGKF-NH$_2$.

Figure 13. AYPGKF-NH$_2$-stimulated calcium signalling and β-arrestin-1/-2 recruitment in PAR4 extracellular loop 2 mutants. Agonist-stimulated Calcium signalling Calcium signalling in response to AYPGKF-NH$_2$ (A) or thrombin (D) was recorded in HEK293 cells transiently expressing PAR4-YFP, single site-directed mutant PAR4-YFP (PAR4^{H229A}-YFP, PAR4^{D230A}-YFP, PAR4^{Q242A}-YFP), or double site-directed mutant PAR4-YFP (PAR4$^{H229A, D230A}$-YFP, PAR4$^{H229D, A231D}$-YFP). Data are shown as a percentage of maximum response obtained in cells treated with the calcium ionophore A23187 (3 μM) for each transient cell line, per experimental day. AYPGKF-NH$_2$ concentration is in M and thrombin concentration is shown as units. (N=3-5) **Agonist-stimulated β-arrestin-1/-2 recruitment** HEK293 cells were transiently transfected with BRET pair of either β-arrestin-1-rluc or -2-rluc and PAR4-YFP or site-directed mutant PAR4-YFP (described above) and agonist-stimulated arrestin recruitment to PAR4 was monitored. Cells were incubated with increasing concentration of either

1331   AYPGKF-NH$_2$ (B, C) or thrombin (E, F) for 20 minutes and BRET values were recorded.

1332   BRET values from cells treated with HBSS were also collected to normalize the data. Renilla-

1333   luciferase substrate coelenterazine-h (5 µM) was added 10 minutes prior to data collection.

1334   Data are presented as the normalized net BRET ratio [eYFP/rluc - baseline (HBSS control)].

1335   AYPGKF-NH$_2$ concentration is in M and thrombin concentration is shown as units. ($n = 4$).

Figure 14. PAR4 agonist-mediated platelet aggregation. Both thrombin (1 unit/mL; shown in black) and AYPGKF-NH$_2$ (100 μM; shown in grey) initiated PAR4-dependent rat platelet aggregation. We observed that stimulation of washed platelet samples with a calcium signalling null, β-arrestin biased peptide, AyPGKF-NH$_2$ (100 μM; shown in yellow) did not stimulate platelet aggregation. Further, stimulation of washed platelets with the peptide agonist AYPGRF-NH$_2$ (100 μM; shown in purple), caused a robust platelet aggregation comparable to the response with 1unit/mL thrombin response (*n = 2-4*).

References.

Adams MN, Ramachandran R, Yau M-K, Suen JY, Fairlie DP, Hollenberg MD, Hooper JD. 2011. Structure, function and pathophysiology of protease activated receptors. *Pharmacology & therapeutics* **130**:248–282.

Al-Ani B, Saifeddine M, Kawabata A, Hollenberg MD. 1999. Proteinase activated receptor 2: Role of extracellular loop 2 for ligand-mediated activation. *Br J Pharmacol* **128**:1105–1113.

Al-Ani B, Wijesuriya SJ, Hollenberg MD. 2002. Proteinase-activated receptor 2: differential activation of the receptor by tethered ligand and soluble peptide analogs. *The Journal of pharmacology and experimental therapeutics* **302**:1046–1054.

Bar-Shavit R, Maoz M, Kancharla A, Jaber M, Agranovich D, Grisaru-Granovsky S, Uziely B. 2016. Protease-activated receptors (PARs) in cancer: Novel biased signaling and targets for therapy. *Methods in cell biology* **132**:341–358. doi:10.1016/bs.mcb.2015.11.006

Blackhart BD, Emilsson K, Nguyen D, Teng W, Martelli AJ, Nystedt S, Sundelin J, Scarborough RM. 1996. Ligand cross-reactivity within the protease-activated receptor family. *The Journal of biological chemistry* **271**:16466–16471.

Boitano S, Flynn AN, Schulz SM, Hoffman J, Price TJ, Vagner J. 2011. Potent Agonists of the Protease Activated Receptor 2 (PAR2). *J Med Chem* **54**:1308–1313. doi:10.1021/jm1013049

Cheng RKY, Fiez-Vandal C, Schlenker O, Edman K, Aggeler B, Brown DG, Brown GA, Cooke RM, Dumelin CE, Doré AS, Geschwindner S, Grebner C, Hermansson N-O, Jazayeri A, Johansson P, Leong L, Prihandoko R, Rappas M, Soutter H, Snijder A, Sundström L, Tehan B, Thornton P, Troast D, Wiggin G, Zhukov A, Marshall FH, Dekker N. 2017. Structural insight into allosteric modulation of protease-activated receptor 2. *Nature* **545**:112–115. doi:10.1038/nature22309

Chidiac P. 2016. RGS proteins destroy spare receptors: Effects of GPCR-interacting proteins and signal deamplification on measurements of GPCR agonist potency. *Methods* **92**:87–93. doi:10.1016/j.ymeth.2015.08.011

Coughlin SR. 2005. Protease-activated receptors in hemostasis, thrombosis and vascular biology. *J Thromb Haemost* **3**:1800–1814.

Coughlin SR. 1999. How the protease thrombin talks to cells. *Proceedings of the National Academy of Sciences of the United States of America* **96**:11023–11027.

Covic L, Misra M, Badar J, Singh C, Kuliopulos A. 2002. Pepducin-based intervention of thrombin-receptor signaling and systemic platelet activation. *Nature Medicine* **8**:1161–1165.

Dolinsky TJ, Nielsen JE, McCammon JA, Baker NA. 2004. PDB2PQR: an automated pipeline for the setup of Poisson-Boltzmann electrostatics calculations. *Nucleic Acids Res* **32**:W665-667. doi:10.1093/nar/gkh381

Edelstein LC, Simon LM, Lindsay CR, Kong X, Teruel-Montoya R, Tourdot BE, Chen ES, Ma L, Coughlin S, Nieman M, Holinstat M, Shaw CA, Bray PF. 2014. Common variants in the human platelet PAR4 thrombin receptor alter platelet function and differ by race. *Blood* **124**:3450–3458. doi:10.1182/blood-2014-04-572479

Erlandson SC, McMahon C, Kruse AC. 2018. Structural Basis for G Protein–Coupled Receptor Signaling. *doi.org* **47**:annurev–biophys–070317–032931.

Faruqi TR, Weiss EJ, Shapiro MJ, Huang W, Coughlin SR. 2000. Structure-function analysis of protease-activated receptor 4 tethered ligand peptides. Determinants of specificity and utility in assays of receptor function. *The Journal of biological chemistry* **275**:19728–19734.

Ferguson SSG, Caron MG. 2004. Green Fluorescent Protein-Tagged β-Arrestin Translocation as a Measure of G Protein-Coupled Receptor Activation In: Smrcka AV, editor. G Protein Signaling: Methods and Protocols, Methods in Molecular Biology™. Totowa, NJ: Humana Press. pp. 121–126. doi:10.1385/1-59259-430-1:121

Fredriksson R, Schioth HB. 2005. The repertoire of G-protein-coupled receptors in fully sequenced genomes. *Molecular pharmacology* **67**:1414–1425. doi:10.1124/mol.104.009001

French SL, Hamilton JR. 2016. Protease-activated receptor 4: from structure to function and back again. *Br J Pharmacol* **173**:2952–2965. doi:10.1111/bph.13455

Gerszten RE, Chen J, Ishii M, Ishii K, Wang L, Nanevicz T, Turck CW, Vu TK, Coughlin SR. 1994. Specificity of the thrombin receptor for agonist peptide is defined by its extracellular surface. *Nature* **368**:648–651.

Graham FL, van der Eb AJ. 1973. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* **52**:456–467.

Hilger D, Masureel M, Kobilka BK. 2018. Structure and dynamics of GPCR signaling complexes. *Nature structural & molecular biology* **25**:4–12. doi:10.1038/s41594-017-0011-7

Hoffmann W, Langenhan J, Huhmann S, Moschner J, Chang R, Accorsi M, Seo J, Rademann J, Meijer G, Koksch B, Bowers MT, von Helden G, Pagel K. 2019. An Intrinsic Hydrophobicity Scale for Amino Acids and Its Application to Fluorinated Compounds. *Angewandte Chemie International Edition* **58**:8216–8220. doi:10.1002/anie.201813954

Holinstat M, Voss B, Bilodeau ML, McLaughlin JN, Cleator J, Hamm HE. 2006. PAR4, but not PAR1, signals human platelet aggregation via Ca2+ mobilization and synergistic P2Y12 receptor activation. *The Journal of biological chemistry* **281**:26665–26674.

Hollenberg MD. 2003. Proteinase-activated receptors: Tethered ligands and receptor-activating peptides. *Drug Dev Res* **59**:336–343. doi:10.1002/ddr.10301

Hollenberg MD, Mihara K, Polley D, Suen JY, Han A, Fairlie DP, Ramachandran R. 2014. Biased signalling and proteinase-activated receptors (PARs): targeting inflammatory disease. *British journal of pharmacology* **171**:1180–1194.

Ishihara H, Connolly AJ, Zeng D, Kahn ML, Zheng YW, Timmons C, Tram T, Coughlin SR. 1997. Protease-activated receptor 3 is a second thrombin receptor in humans. *Nature* **386**:502–506.

Jacques SL, Kuliopulos A. 2003. Protease-activated receptor-4 uses dual prolines and an anionic retention motif for thrombin recognition and cleavage. *The Biochemical journal* **376**:733–740.

Jacques SL, LeMasurier M, Sheridan PJ, Seeley SK, Kuliopulos A. 2000. Substrate-assisted catalysis of the PAR1 thrombin receptor. Enhancement of macromolecular association and cleavage. *The Journal of biological chemistry* **275**:40671–40678.

Jean-Charles P-Y, Kaur S, Shenoy SK. 2017. GPCR signaling via β-arrestin-dependent mechanisms. *J Cardiovasc Pharmacol* **70**:142–158. doi:10.1097/FJC.0000000000000482

Kahn ML, Zheng YW, Huang W, Bigornia V, Zeng D, Moff S, Farese RVJ, Tam C, Coughlin SR. 1998. A dual thrombin receptor system for platelet activation. *Nature* **394**:690–694.

Khan A, Li D, Ibrahim S, Smyth E, Woulfe DS. 2014. The Physical Association of the P2Y12 Receptor with PAR4 Regulates Arrestin-Mediated Akt Activation. *Molecular pharmacology* **86**:1–11.

Kim BH, Wang FI, Pereverzev A, Chidiac P, Dixon SJ. 2019. Toward Defining the Pharmacophore for Positive Allosteric Modulation of PTH1 Receptor Signaling by Extracellular Nucleotides. *ACS Pharmacol Transl Sci* **2**:155–167. doi:10.1021/acsptsci.8b00053

Lee H, Heo L, Lee MS, Seok C. 2015. GalaxyPepDock: a protein–peptide docking tool based on interaction similarity and energy optimization. *Nucleic Acids Res* **43**:W431–W435. doi:10.1093/nar/gkv495

Lerner DJ, Chen M, Tram T, Coughlin SR. 1996. Agonist recognition by proteinase-activated receptor 2 and thrombin receptor. Importance of extracellular loop interactions for receptor function. *The Journal of biological chemistry* **271**:13943–13947.

Li D, D'Angelo L, Chavez M, Woulfe DS. 2011. Arrestin-2 differentially regulates PAR4 and ADP receptor signaling in platelets. *The Journal of biological chemistry* **286**:3805–3814.

Liu LW, Vu TK, Esmon CT, Coughlin SR. 1991. The region of the thrombin receptor resembling hirudin binds to thrombin and alters enzyme specificity. *The Journal of biological chemistry* **266**:16977–16980.

Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN, Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res* **47**:W636–W641. doi:10.1093/nar/gkz268

Miller MM, Banville J, Friends T, Gagnon M, Hangeland JJ, Lavalee J-F, Martel A, O'Grady H, Remillard R, Ruediger EH, Tremblay F, Posy S, Allegretto N, Guarino VR, Harden D, Harper TW, Hartl K, Josephs J, Malmstrom S, Watson CA, Yang Y, Zhang G, Wong PC, Yang J, Bouvier M, Seiffert DA, Wexler RR, Lawrence RM, Priestley ES, Marinier A. 2019. Discovery of Potent Protease-Activated Receptor 4 Antagonists with in vivo Antithrombotic Efficacy. *J Med Chem*. doi:10.1021/acs.jmedchem.9b00186

Morrow DA, Braunwald E, Bonaca MP, Ameriso SF, Dalby AJ, Fish MP, Fox KA, Lipka LJ, Liu X, Nicolau JC, Ophuis AJ, Paolasso E, Scirica BM, Spinar J, Theroux P, Wiviott SD, Strony J, Murphy SA. 2012. Vorapaxar in the secondary prevention of atherothrombotic events. *The New England journal of medicine* **366**:1404–1413.

Morrow DA, Scirica BM, Fox KA, Berman G, Strony J, Veltri E, Bonaca MP, Fish P, McCabe CH, Braunwald E. 2009. Evaluation of a novel antiplatelet agent for secondary prevention in patients with a history of atherosclerotic disease: design and rationale for the Thrombin-Receptor Antagonist in Secondary Prevention of Atherothrombotic Ischemic Events (TRA 2 degrees P)-TIMI 50 trial. *Am Heart J* **158**:335–341 e3.

Moschonas IC, Kellici TF, Mavromoustakos T, Stathopoulos P, Tsikaris V, Magafa V, Tzakos AG, Tselepis AD. 2017. Molecular requirements involving the human platelet protease-activated receptor-4 mechanism of activation by peptide analogues of its tethered-ligand. *Platelets* **28**:812–821. doi:10.1080/09537104.2017.1282607

Nanevicz T, Ishii M, Wang L, Chen M, Chen J, Turck CW, Cohen FE, Coughlin SR. 1995. Mechanisms of thrombin receptor agonist specificity. Chimeric receptors and complementary mutations identify an agonist recognition site. *The Journal of biological chemistry* **270**:21619–21625.

Nieman MT. 2008. Protease-Activated Receptor 4 Uses Anionic Residues To Interact with α-Thrombin in the Absence or Presence of Protease-Activated Receptor 1. *Biochemistry* **47**:13279–13286. doi:10.1021/bi801334s

Nieman MT, Schmaier AH. 2007. Interaction of Thrombin with PAR1 and PAR4 at the Thrombin Cleavage Site. *Biochemistry* **46**:8603–8610. doi:10.1021/bi700597p

Nystedt S, Emilsson K, Wahlestedt C, Sundelin J. 1994. Molecular cloning of a potential proteinase activated receptor. *Proceedings of the National Academy of Sciences of the United States of America* **91**:9208–9212.

Pierce KL, Premont RT, Lefkowitz RJ. 2002. Seven-transmembrane receptors. *Nature Reviews Molecular Cell Biology* **3**:639–650.

Ramachandran R, Mihara K, Thibeault P, Vanderboor CM, Petri B, Saifeddine M, Bouvier M, Hollenberg MD. 2017. Targeting a Proteinase-Activated Receptor 4 (PAR4) Carboxyl Terminal Motif to Regulate Platelet Function. *Molecular pharmacology* **91**:287–295.

Rwibasira Rudinga G, Khan GJ, Kong Y. 2018. Protease-Activated Receptor 4 (PAR4): A Promising Target for Antiplatelet Therapy. *Int J Mol Sci* **19**. doi:10.3390/ijms19020573

Sánchez Centellas D, Gudlur S, Vicente-Carrillo A, Ramström S, Lindahl TL. 2017. A cluster of aspartic residues in the extracellular loop II of PAR 4 is important for thrombin interaction and activation of platelets. *Thrombosis Research* **154**:84–92. doi:10.1016/j.thromres.2017.04.006

Scarborough RM, Naughton MA, Teng W, Hung DT, Rose J, Vu TK, Wheaton VI, Turck CW, Coughlin SR. 1992. Tethered ligand agonist peptides. Structural requirements for

thrombin receptor activation reveal mechanism of proteolytic unmasking of agonist function. *The Journal of biological chemistry* **267**:13146–13149.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**:676–682. doi:10.1038/nmeth.2019

Scirica BM, Bonaca MP, Braunwald E, De Ferrari GM, Isaza D, Lewis BS, Mehrhof F, Merlini PA, Murphy SA, Sabatine MS, Tendera M, Van de Werf F, Wilcox R, Morrow DA. 2012. Vorapaxar for secondary prevention of thrombotic events for patients with previous myocardial infarction: a prespecified subgroup analysis of the TRA 2°P-TIMI 50 trial. *The Lancet* **380**:1317–1324. doi:10.1016/S0140-6736(12)61269-0

Seyedabadi M, Ghahremani MH, Albert PR. 2019. Biased signaling of G protein coupled receptors (GPCRs): Molecular determinants of GPCR/transducer selectivity and therapeutic potential. *Pharmacol Ther* **200**:148–178. doi:10.1016/j.pharmthera.2019.05.006

Shapiro MJ, Weiss EJ, Faruqi TR, Coughlin SR. 2000. Protease-activated receptors 1 and 4 are shut off with distinct kinetics after activation by thrombin. *The Journal of biological chemistry* **275**:25216–25221.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* **7**:539. doi:10.1038/msb.2011.75

Sriram K, Insel PA. 2018. G Protein-Coupled Receptors as Targets for Approved Drugs: How Many Targets and How Many Drugs? *Molecular pharmacology* **93**:251–258.

Tourdot BE, Stoveken H, Trumbo D, Yeung J, Kanthi Y, Edelstein LC, Bray PF, Tall GG, Holinstat M. 2018. Genetic Variant in Human PAR (Protease-Activated Receptor) 4 Enhances Thrombus Formation Resulting in Resistance to Antiplatelet Therapeutics. *Arterioscler Thromb Vasc Biol* **38**:1632–1643. doi:10.1161/ATVBAHA.118.311112

Venkatakrishnan AJ, Deupi X, Lebon G, Tate CG, Schertler GF, Babu MM. 2014. Molecular signatures of G-protein-coupled receptors. *Nature* **494**:185–194. doi:10.1038/nature11896

Vu TK, Hung DT, Wheaton VI, Coughlin SR. 1991. Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* **64**:1057–1068.

Webb B, Sali A. 2017. Protein Structure Modeling with MODELLERFunctional Genomics. New York, NY: Humana Press, New York, NY. pp. 39–54. doi:10.1007/978-1-4939-7231-9_4

Wheatley M, Wootten D, Conner MT, Simms J, Kendrick R, Logan RT, Poyner DR, Barwell J. 2012. Lifting the lid on GPCRs: the role of extracellular loops. *Br J Pharmacol* **165**:1688–1703. doi:10.1111/j.1476-5381.2011.01629.x

Wilson SJ, Ismat FA, Wang Z, Cerra M, Narayan H, Raftis J, Gray TJ, Connell S, Garonzik S, Ma X, Yang J, Newby DE. 2017. PAR4 (Protease-Activated Receptor 4) Antagonism With BMS-986120 Inhibits Human Ex Vivo Thrombus Formation. *Arteriosclerosis, Thrombosis, and Vascular Biology* ATVBAHA.117.310104. doi:10.1161/ATVBAHA.117.310104

Wong PC, Seiffert D, Bird JE, Watson CA, Bostwick JS, Giancarli M, Allegretto N, Hua J, Harden D, Guay J, Callejo M, Miller MM, Lawrence RM, Banville J, Guy J, Maxwell BD, Priestley ES, Marinier A, Wexler RR, Bouvier M, Gordon DA, Schumacher WA, Yang J. 2017. Blockade of protease-activated receptor-4 (PAR4) provides robust antithrombotic activity with low bleeding. *Science translational medicine* **9**:eaaf5294. doi:10.1126/scitranslmed.aaf5294

Woolley MJ, Conner AC. 2017. Understanding the common themes and diverse roles of the second extracellular loop (ECL2) of the GPCR super-family. *Molecular and Cellular Endocrinology*, Pleiotropic GPCR signaling in health and disease **449**:3–11. doi:10.1016/j.mce.2016.11.023

Wu CC, Huang SW, Hwang TL, Kuo SC, Lee FY, Teng CM. 2000. YD-3, a novel inhibitor of
protease-induced platelet activation. *Br J Pharmacol* **130**:1289–1296.
Xu WF, Andersen H, Whitmore TE, Presnell SR, Yee DP, Ching A, Gilbert T, Davie EW,
Foster DC. 1998. Cloning and characterization of human protease-activated receptor
4. *Proceedings of the National Academy of Sciences of the United States of America*
95:6642–6646.
Young SE, Duvernay MT, Schulte ML, Nance KD, Melancon BJ, Engers J, Wood MR,
Hamm HE, Lindsley CW. 2010. A Novel and Selective PAR4 Antagonist:
ML354Probe Reports from the NIH Molecular Libraries Program. Bethesda (MD):
National Center for Biotechnology Information (US).
1570

Supplemental tables and figures:

Table S1: MS data of synthesized peptides.

Compound	Molecular Formula (M)	Calc. [M+H]$^+$ (m/z)	Found [M+H]$^+$ (m/z)
Ac-AYPGKF-NH$_2$	$C_{36}H_{50}N_8O_8$	723.38	723.43
AAPGKF-NH$_2$	$C_{28}H_{44}N_8O_6$	589.35	589.30
AYAGKF-NH$_2$	$C_{32}H_{46}N_8O_7$	655.36	655.43
AYPAKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.46
AYPGAF-NH$_2$	$C_{31}H_{41}N_7O_7$	624.31	625.36
AYPGKA-NH$_2$	$C_{28}H_{44}N_8O_7$	605.34	606.42
aYPGKF-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.44
AyPGKF-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.44
AYPGkF-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.46
AYPGKf-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.48
Aib-YPGKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.49
Sar-YPGKF-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.38
Inp-YPGKF-NH$_2$	$C_{37}H_{52}N_8O_7$	721.40	721.50
betaAla-YPGKF-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.46
A-Tyr(Me)-PGKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.48
A-F(4-fluoro)-PGKF-NH$_2$	$C_{34}H_{47}N_8O_6F$	683.37	683.36
AY-Pip-GKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
AY-Nip-GKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
AYP-Sar-KF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
AYP-betaAla-KF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.48
AYPGK-1Nal-NH$_2$	$C_{38}H_{50}N_8O_7$	731.38	731.48
AYPGK-2Nal-NH$_2$	$C_{38}H_{50}N_8O_7$	731.38	731.50
AYPGK-F(4-fluoro)-NH$_2$	$C_{34}H_{47}N_8O_7F$	699.36	699.38
AYPGK-F(4-Me)-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
AYpGKF-NH$_2$	$C_{34}H_{48}N_8O_7$	681.37	681.37
AYPGK-(*N*-Me-F)-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
AYPG-(N_α-Me-K)-F-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
A-(*N*-Me-Y)-PGKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
(*N*-Me-A)-YPGKF-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
VYPGKF-NH$_2$	$C_{36}H_{52}N_8O_7$	709.40	709.40
(*N*-Me-S)-YPGKF-NH$_2$	$C_{35}H_{50}N_8O_8$	711.38	711.38
Nle-YPGKF-NH$_2$	$C_{37}H_{54}N_8O_7$	723.42	723.42
AYPGRF-NH$_2$	$C_{34}H_{48}N_{10}O_7$	709.38	709.38
AYPGOF-NH$_2$	$C_{33}H_{46}N_8O_7$	667.36	667.36
AYPG-Cit-F-NH$_2$	$C_{34}H_{47}N_9O_8$	710.36	710.36
AYPG-(N_ε-Me-K)-F-NH$_2$	$C_{35}H_{50}N_8O_7$	695.39	695.39
AYPGKY-NH$_2$	$C_{34}H_{48}N_8O_8$	697.37	697.37

1576 **Figure S1: Analytical RP-HPLC UV detection chromatograms of synthesized peptides.**
1577 Purity of all peptides are > 95% as determined by analytical RP-HPLC UV detection (see traces below).
1578

1579 Ac-AYPGKF-NH$_2$:
1580 t_R = 10.50 min; gradient = 10% to 50% (solvent A in solvent B, 0 to 10 min)

1581
1582

1583 AAPGKF-NH2:
1584 t_R = 8.08 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1585
1586

1587 AYAGKF-NH$_2$:
1588 t_R = 9.14 min; gradient = 10% to 60% (solvent A in solvent B, 0 to 10 min)

1597 AYPAKF-NH$_2$:

1598 t_R = 10.53 min; gradient = 10% to 50% (solvent A in solvent B, 0 to 10 min)

1599

1600

1601 AYPGAF-NH$_2$:

1602 t_R = 10.83 min; gradient = 10% to 50% (solvent A in solvent B, 0 to 10 min)

1603

1604

1605 AYPGKA-NH$_2$:

1606 t_R = 9.71 min; gradient = 5% to 30% (solvent A in solvent B, 0 to 10 min)

1607

1608

1609

1610

1611

1612

1613

1614

1615

1616

1617

aYPGKF-NH$_2$:

t_R = 10.06 min; gradient = 10% to 50% (solvent A in solvent B, 0 to 10 min)

AyPGKF-NH$_2$:

t_R = 9.49 min; gradient = 15% to 50% (solvent A in solvent B, 0 to 10 min)

AYPGkF-NH$_2$:

t_R = 9.17 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1639   AYPGKf-NH$_2$:

1640 t_R = 9.04 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1641

1642

1643   Aib-YPGKF-NH$_2$:

1644 t_R = 9.35 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1645

1646

1647   Sar-YPGKF-NH$_2$:

1648 t_R = 9.23 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1649

1650

1651

1652

1653

1654

1655

1656

1657

1658

1659

1660   Inp-YPGKF-NH$_2$:

1661 t_R = 9.26 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1662

1663

1664   betaAla-YPGKF-NH$_2$:

1665 t_R = 9.34 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1666

1667

1668   A-Tyr(Me)-PGKF-NH$_2$:

1669 t_R = 9.57 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1670

1671

1672

1673

1674

1675

1676

1677

1678   A-F(4-fluoro)-PGKF-NH$_2$:

1679 t_R = 9.63 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1680

1681

1682   AY-Pip-GKF-NH$_2$:

1683 t_R = 9.25 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1684

1685

1686   AY-Nip-GKF-NH$_2$:

1687 t_R = 8.77 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1688

1689

1690

1691

1692

1693

1694

1695

1696

1697

1698

1699   AYP-Sar-KF-NH$_2$:

1700 t_R = 9.56 min; gradient = 10% to 60% (solvent A in solvent B, 0 to 10 min)

1701

1702

1703   AYP-betaAla-KF-NH$_2$:

1704 t_R = 9.38 min; gradient = 10% to 70% (solvent A in solvent B, 0 to 10 min)

1705

1706

1707   AYPGK-1Nal-NH$_2$:

1708 t_R = 9.71 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1709

1710

1711

1712

1713

1714

1715

1716

1717

1718

1719   AYPGK-2Nal-NH$_2$:

1720 t_R = 9.72 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1721

1722

1723   AYPGK-F(4-fluoro)-NH$_2$:

1724 t_R = 8.91 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1725

1726

1727   AYPGK-F(4-Me)-NH$_2$:

1728 t_R = 9.32 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1729

1730

1731

1732

1733

1734

1735

1736

1737

1738

1739

1740   AYpGKF-NH$_2$:

1741   t$_R$ = 9.02 min; gradient = 12% to 75% (solvent A in solvent B, 0 to 10 min)

1742
1743
1744   AYPGK-(*N*-Me-F)-NH$_2$:

1745   t$_R$ = 8.73 min; gradient = 12% to 80% (solvent A in solvent B, 0 to 10 min)

1746
1747
1748   AYPG-(*N*$_\alpha$-Me-K)-F-NH$_2$:

1749   t$_R$ = 8.33 min; gradient = 15% to 70% (solvent A in solvent B, 0 to 10 min)

1756   A-(N-Me-Y)-PGKF-NH$_2$:

1757 t_R = 8.74 min; gradient = 12% to 70% (solvent A in solvent B, 0 to 10 min)

1758
1759
1760   (N-Me-A)-YPGKF-NH$_2$:

1761 t_R = 8.88 min; gradient = 12% to 70% (solvent A in solvent B, 0 to 10 min)

1762
1763
1764   VYPGKF-NH$_2$:

1765 t_R = 8.90 min; gradient = 12% to 70% (solvent A in solvent B, 0 to 10 min)

1772   (*N*-Me-S)-YPGKF-NH$_2$:

1773 t_R = 9.00 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1774
1775
1776   Nle-YPGKF-NH$_2$:

1777 t_R = 8.88 min; gradient = 15% to 70% (solvent A in solvent B, 0 to 10 min)

1778
1779
1780   AYPGRF-NH$_2$:

1781 t_R = 8.73 min; gradient = 13% to 70% (solvent A in solvent B, 0 to 10 min)

1788   AYPGOF-NH$_2$:

1789 t_R = 9.16 min; gradient = 12% to 80% (solvent A in solvent B, 0 to 10 min)

1790
1791

1792   AYPG-Cit-F-NH$_2$:

1793 t_R = 8.82 min; gradient = 13% to 80% (solvent A in solvent B, 0 to 10 min)

1794
1795

1796   AYPG-(N_ε-Me-K)-F-NH$_2$:

1797 t_R = 8.74 min; gradient = 12% to 70% (solvent A in solvent B, 0 to 10 min)

1804   AYPGKY-NH$_2$:

1805 t_R = 8.88 min; gradient = 8% to 80% (solvent A in solvent B, 0 to 10 min)

1806

Figure S2. Representative blots of ERK phosphorylation in response to 100 µM of AYPGKF-NH₂ and derivative peptides. Data are summarized in figure 10.

A. β-Arrestin-1 TGTGGACCACATCGACCTCG
    β-Arrestin-2 GCGGGACTTCGTAGATCACC

B.

Figure S3. (A) Guide RNA sequences designed to target either β-arrestin-1 or -2. (B) Representative blots of β-arrestin-1/-2 protein from HEK-293 and β-arrestin-1/-2 double knockout HEK-293 (β-arr. Knockout) cells. We observe no β-arrestin-1/-2 protein in the β-arr. knockout HEK-293 cell line.

Figure S4. PAR4 peptide library is specific for PAR4 as shown by lack of calcium
signalling in HEK293 cells which do not endogenously express PAR4. Peptide AAPGKF-
NH$_2$ was found to have basal fluorescence even in the absence of cells. (*n = 3*)

Figure S5. PAR4 peptide library is specific for PAR4 and does not cause recruitment of
β-arrestin-2 to PAR2 (shown as black bars) at 100 μM, with the exception of AYPGK-
1Nal-NH$_2$ which shows some β-arrestin recruitment to PAR2. BRET ratios obtained with
the parental peptides for both PAR4 (AYPGKF-NH$_2$) and PAR2 (SLIGRL-NH$_2$) are
shown. (*n = 3*)