Early vegetation recovery after the 2008-2009 explosive eruption of the Chaitén Volcano.

Ricardo Moreno-González[1*], Iván A. Díaz[2], Duncan A. Christie[2,3], Rafael E. Coopman[2,4], Antonio Lara[2, 3]

1. Department of Palynology and Climate Dynamic, University of Göttingen, Germany.

2. Instituto de Conservación, Biodiversidad y Territorio, Universidad Austral de Chile, Valdivia, Chile.

3. Center for Climate and Resilience Research (CR)²

4. Marilyn Ball's Lab, Plant Science Division, Research school of Biology, The Australian National University

* Corresponding author: Ricardo Moreno González

E-mail: rmoreno@uni-goettingen.de

Abstract

In May 2008, Chaitén volcano entered in eruptive process, one of the world largest eruptions in the last decades. The catastrophic event left different type of disturbance and caused diverse environmental damage distributed heterogeneously in the surrounding areas of the volcano. We went to the field to assess the early vegetation responses a year after the eruption, in September 2009. Particularly, we evaluated the lateral-blast disturbance zone. We distributed a set of plots in three disturbed sites, and one in an undisturbed site. In each of these sites, in a plot of 1000m^2 we marked all stand tree, recording whether they were alive, resprouting or dead. Additionally, in each site 80 small-plots (~4m^2) we tallied the plants regeneration, its coverage, and the log-volume. We described whether the plant regeneration was growing on mineral or organic substrate. In the blast-zone the eruption created a gradient of disturbance. Close to the crater we found high devastation marked by no surviving species, scarce standing-dead trees and logs, as well as no tree regeneration. On the other extreme of the disturbance gradient, the trees with damaged crown were resprouting, small-plants were regrowing and seedlings were more dispersed. The main regeneration strategy was the resprouting of trunks or buried roots, while few seedlings were observed in the small plots and elsewhere in disturbed areas. However, the assessment was too soon after the eruption and updated monitoring is required to confirm observed patterns. The main findings of this study are: i) a mosaic of pioneering-wind dispersed species, scattered survivors regrowing and spreading from biological legacies, and plant species dispersed by frugivorous birds, likely favored by the biological legacies; (ii) the early succession is influenced by the interaction of the species-specific life history, altitudinal gradient and the different intensity of disturbance.

41 Keywords: 2008-Chaiten Volcano, Volcanic disturbance, vegetation dynamics, *Nothofagus*

42 dynamics, South American temperate rainforest, early succession.

1. Introduction

Volcanic eruptions are large-scale disturbance that modify the composition, the structure and

ecosystems processes (Swanson et al., 2013; Turner and Dale, 1998; Turner et al., 1997). As

volcanoes enter in eruptive phase complex mechanism involve the events in a gradient of

multiple type of disturbances such as explosive blasts, thermal and toxic chemical waves,

landslides, glowing avalanche deposits, debris flows, lava flows, and air falls of volcanic tephra

(Dale et al., 2005; Swanson et al., 2013). A volcanic eruption could have effects at very large

spatial and temporal scales (Peet, 1992; Robock and Oppenheimer, 2003), yet directly on

surrounded forest throughout burning, burying and/or blowing down trees or other plants (Dale

et al., 2005). However, since this kind of large disturbance are infrequent (Turner et al. 1998)

or have been seldom monitored (Chrisafully and Dale, 2018), our understanding of vegetation

responses is still limited. Despite the importance of volcanic eruptions, few studies have

analyzed the effect of eruptions short after the event (Dale et al., 2005; Chrisafully and Dale

2018). On the other hand, the effect of volcanism has been studied by comparing different

stands in chronosequences assuming that different forest patches would reflect the successional

stage after the disturbance (e.g. Drury and Nisbet 1973; Franklin and Swanson, 2010; Pickett

1988; Walker et al 2010).

Forest recovery is considerably slow following disturbances that heavily impact soils as well

as aboveground vegetation (Chazdon, 2003). Despite volcanoes are considered as catastrophic

disturbance event, it is possible that diverse biological legacies can remain after the event. The

persistence and heterogeneous distribution of these biological legacies might modulate and help

the recolonization of the plants (Foster et al., 1998; Franklin et al., 1985; Turner et al., 1998).

For example, the life-history traits of the pre-disturbed community will result in particular

survivors perhaps able to re-colonize and expand over the new conditions on the disturbed

environment (Foster et al., 1998; Turner et al., 1998). Therefore, the arrange and the amount of

the biological legacies would drive the rate and the pathways of the new colonizers.

One of the most active tectonic and volcanic areas in the world correspond to the subduction

area of the Nazca and the South American plate (e.g., Stern, 2004), which also originated the

longest and the second highest mountain range in the world, the Andes. The volcanoes in the

Chilean Andes are quite active. The Southern and Austral Volcanic Zones comprise 74 active

volcanoes since the Post-Glacial up to the present and at least 21 have had one large explosive

eruption (Fontijn et al., 2014). In the southern portion of the Andes, it is located the South

American Temperate rainforest (SATR) (Armesto et al., 1998). This cordillera plays a critical

role in SATR dynamics, where current successional models after volcanic disturbance suggest

the *Nothofagus* are dominant pioneer trees that benefit from disturbances to persist (Veblen et

al., 2016). The SATR represent a biogeographical island, rich in endemic species and

dominated by evergreen, broad-leaved species (Armesto et al., 1996; Villagran and Hinojosa,

1997, Arroyo et al., 2004). South of 40°S, extensive areas of old-growth forest dominate the

mountainous landscape. These forest are not directly disturbed by modern human activity, and

represent one of the last pristine areas with pre-industrial biogeochemical conditions (Perakis

and Hedin 2002)

On May 1st 2008 started the last eruption of the Chaiten Volcano (42° 50'S), near the small

town of Chaiten, in southern Chile. This eruption was the largest rhyolite eruption since the

great eruption of Katmai Volcano in 1912, being the first rhyolite eruption in which some

attributes of its dynamics and impacts have been monitored (Major and Lara, 2013). The

eruption consisted of an approximately 2-week-long explosive phase that generated as much as

1 km^3 bulk volume tephra (~0.3 km^3 dense rock equivalent) followed by an approximately 20-

month-long effusive phase that erupted about 0.8 km^3 of high-silica rhyolite lava that formed a

new dome within the volcano's caldera (Major and Lara, 2013). Chaiten volcano was

surrounded by extensive old-growth undisturbed forest by humans, providing a unique

opportunity to evaluate the effects of volcanism over undisturbed landscape, then providing an

opportunity to understand how volcanism may have affected the forest, under pre-industrial

conditions.

In this study, we characterized the early establishment of the vegetation after the eruption of

the Chaitén volcano along a disturbance gradient one year after the eruption started, from

nearby the crater to the closest old-growth forest with vegetation completely alive. We predict

that plant species diversity and the number of living trees decrease close to the crater, and

pioneering *Nothofagus* species are colonizing the areas affected by the eruption. Our objectives

included the creation of a baseline for further long-term monitoring of vegetation recovering,

and to identify and analyze the importance of biological legacies for forest recovery.

2. Material and methods

2.1 Study site

The Chaitén volcano is located in the west side of the Andean Range in southern Chile, at 42º

50' S and 72º 39' W (Fig 1) and is part of the so-called Southern Volcanic Zones. The Chaitén

Volcano is a dome of 1100 m height, located at 10 km NW of the town of Chaitén, populated

by approximately 5000 inhabitants before the eruption. The climate of the area corresponds to

a humid temperate with a strong oceanic influence (Luebert and Pliscoff, 2006). The landscape

is characterized by sharp mountains shaped by glaciers and valleys originated by fluvial and

glacio-fluvial outwash (Swanson et al., 2013). Vegetation within the influence area of the

Chaiten volcano is dominated by a dense old-growth temperate rainforest of northern

Patagonian and Valdivian types (Luebert and Pliscoff, 2006; Swanson et al., 2013; Veblen et

al., 1983; Veblen et al., 1996). These forests composition is dominated by broad-leaved

evergreen trees such as *Nothofagus dombeyi* (Mirb.) Oerst. (Nothofagaceae), *Laureliopsis*

philippiana (Looser) Schodde, *Gevuina avellana* (Molina), *Amomyrtus luma* (Molina) D.

Legrand and Kausel, *Luma apiculata* (DC.) Burret, *Drymis winteri* J.R. Forst. and G. Forst.,

Eucryphia cordifolia Cav. and *Weinmannia trichosperma* Cav. (both Cunoniaceae). These

species are normally arranged in multi-layered canopy strata and old, large-emergent trees (up

to c. 40 m height) and abundant snags and logs. The understory is densely covered by ferns

such as *Lophosoria quadripinnata* (J.F.Gmel.) C.Chr. and *Blechnum magellanicum* (Desv.)

Mett., seedlings, saplings and bamboo tickets (*Chusquea* spp.).

Figure 1. Study area.

2.2 Study design

We carried out the field work in September 2009. We first conducted a general inspection of

the study area (del Moral, 1981). Later, we selected four sites with different degrees of volcanic

impact following similar classifications of previous studies (del Moral, 1981). The first site

corresponds to a forest with low volcanic impact that only received few millimeters of tephra

deposition. This site was called "Old-growth forest". It was located 5 km from the center of the

new dome at an elevation of 600 m a.s.l. (Figure 1b). The second site was characterized by the

presence of abundant dead standing trees, and higher tephra deposition. This site was called

"Standing-dead" located at 3.5 km from the center of the new dome, at 160 m. The third site

was more affected by the eruption, and was characterized by high tephra deposition, where most

trees were broken and throw down by the eruption. This site was called "Blow down" and was

located at 2.3 km from the center of the new dome at an elevation of 300 m a.s.l. Finally, the

fourth and most affected site was in the border of the crater, at 1.5 km of the center of the new

dome and at 750 m a.s.l. This site presented the higher tephra deposition and all trees were blow

down by the eruption and heavily buried under volcanic ash. This site was called "Total

destruction".

In each one of these four areas was installed a 50 x 20 m permanent plot where we

measured the diameter at the breast height (dbh) of all trees $\geq$ 5cm. Every tree was identified at

the species level when possible and marked. We registered if the stems were alive, resprouting

after the eruption, or snag (standing dead). To estimate the presence and abundance of tree

regeneration, we traced four transects of 120 m long. Transects were distributed parallel,

separated 20 m from each other. In each transect, we mounted five circular plots of 1.5m radius

($4.71m^2$) every 30 m. On these plots we counted all seedlings and saplings present, and

recording the substrate where they rooted (forest soil, logs, or volcanic tephra). We registered

other non-tree plant species in the circular plots, estimated its abundance as the relative cover,

and we also recorded the substrate where they rooted were organic or mineral (e.g., forest soil,

logs, or volcanic tephra). We assessed the volume of all exposed logs in the plot, until each

border within the plot. Assuming a cylindrical form of each log, we calculated the volume of

logs present inside the plot and standardizing its measurements to m^3 of logs per m^2 of plot

surface.

2.3 Data Analysis

Trees seedlings, herbs and shrubs species richness was compared among sites using rarefaction

analysis (Colwell 2006). Rarefaction curves allow for comparing species richness among

environments avoiding the bias in the number of species detected due to the increased

probability of species detection in areas where individuals were more abundant, by plotting the

number of species detected in function of the number of individuals observed (Colwell and

Coddington, 1994, Gotelli & Colwell 2011, Colwell et al., 2012). This procedure includes

Monte Carlo simulations delivering an average value of species richness with a confidence

interval of 95% (Colwell 2006). Rarefaction analyses were conducted using the software

EstimateS Win 8 (Colwell 2006). Non-vascular species were excluded from the rarefaction

analyses since not all of them were classified up to the species level, and one genus could

include several species. Plant composition among sites was compared using Sørensen's

similarity index based on the presence/absence of plant species. Rarefaction and similitude

analyses were both carried out with the software Estimates 8.2 (Colwell 2006). In attempting

to define if the plant composition of one site was a subset of the composition of other site, we

conducted a Nested Analysis in NestCalc free access software (Atmar and Patterson, 1995).

3. Results

3.1 General description of the effects of the eruption

The eruption of the Chaitén volcano has an impressive effect on the forest. The eruption in the

study area produced a gradient of destruction. Around the crater all the trees were blown down

and heavily buried by tephra. Other trees were partially buried, broken in half, with the trunks

covered with impacts of many little stones looking similar to the fire of gun shots (Fig x).

Moving away and down from the crater, trees were blown down and densely covered by ash(Fig

x), while further away trees were standing dead presenting progressively finer branches(Fig x).

Many small patches of the organic litter were exposed and sparsely distributed and protected

from the direct effect of the eruption by root discs and large trunks, or were exposed by small

ravines that flush away the volcanic ash. These small patches looked like clumps of organic

soil, with roots of ferns and other small plants that were not buried by the tephra. These clumps

were covered by plants that survived in the area affected by the eruption, and their frequency

increases when moving away from the crater towards the undisturbed forests.

In the surrounding, we did not observed evidence of fire caused by the eruption, but the trunk

and bark of several trees looked charred within the influence area. Trees were dead and dry by

the eruptive explosions, but some trees in Blow-down and Standing-dead areas maintained the

bark in the opposite side of the crater, perhaps with possibilities to resprouting hereafter. Tephra

buried the fallen trees in the whole study area, but principally in Blow-down and Total-

destruction areas. Tephra was very compacted, seeming as pavement that covered the fallen

forests. Heterogeneous tephra deposition exposed some trunks, clumps and remnants with

organic-forest soil likely from the original forest. Shrubs and epiphytes were spread everywhere

in the Standing-dead area. Ferns, bamboo and tree seedlings were growing in remnants of

exposed organic soil, usually below or at one side of trunks or unearth roots. Surviving trees

also increases when moving away and down from the crater.

3.2 Species composition

We found 10 species of trees in the area, and 50 species of smaller plants, including seedlings,

herbs and ferns (Table 1). Rarefaction analysis for tree species showed that the Standing-dead

plot holds one more species than the Old-growth forest plot (Fig. 3). This same analysis showed

the Blow-down plot holds very few individuals, but the initial slope and shape of the curve was

similar to the other two curves. It indicates that at similar number of individuals, all curves

showed a similar number of tree species. Rarefaction analysis of small vascular plants showed

the Old-growth forest site had the richest species composition, while decreasing in the Standing

dead and Blow-down plots (Fig. 3). The Total-destruction plot showed practically no species.

Table 1. List of plant species present in the Chaitén volcano and their frequency in the big plot and in 20 plots of 3.14m² in the different studied areas.

Families	Species	Old-growth	Standing dead	Blow-down	Near crater
SEEDLINGS					
Cunoneaceae	*Weinmannia trichosperma*		2		
	Caldcluvia paniculata	4	4		
Eucryphiaceae	*Eucryphia cordifolia*	5			
Flacourtiaceae	*Azara lanceolata*			1	
Monimiacae	*Laureliopsis philippiana*	6			
Podocarpaceae	*Podocarpus saligna*	1			
Winteraceae	*Drimys winteri*	3			
Proteaceae	*Embothrium coccineum*		1	4	
	Lomatia ferruginea	5			
	Gevuina avellana	1			
Myrtaceae	*Myrceugenia parviflora*	3			
	Myrceugenia planipes	12			
	Amomyrtus luma	16	4		
Araliacae	*Raukaua laetevirens*	3			
HERBS AND SMALL SHRUBS					
Gesneraceae	*Mitraria coccinea*	8			
	Asteranthera ovata	1	1		
	Campsidium valdivianum	5			
Poaceae	*Chusquea uliginosa*	7	1	1	
Apocinaceae	*Elytropus chilensis*	6	1		
Fitolaceae	*Ercilla syncarpellata*	3			
Saxifragaceae	*Ribes magellanicum*		1		
Bromeliaceae	*Greigia landbeckii*	4			
Griselineaceae	*Griselinia racemosa*	1			
	Griselinia ruscifolia	4	1	1	
Guneraceae	*Gunnera tinctorea*			1	

Family	Species				
Rubiaceae	*Nertera granadensis*	1			
Hydrangeaceae	*Hydrangea serratifolia*	12			
Luzuriagaceae	*Luzuriaga poliphylla*	8	1	1	
	Luzuriaga radicans	14			

FERNS

Family	Species				
Dicksoniaceae	*Lophosoria quadripinnata*	6	4	3	
Hymenophylaceae	*Hymenophyllum tortuosum*			2	
	Hymenophyllum magellanico	2			
	Hymenoglossum cruentum	6			
	Hymenophyllum caudiculatum	3			
	Hymenophyllum cuneatum	2			
	Hymenophyllum dentatum	5			
	Hymenophyllum dicranotrichium	12			
	Hymenophyllum krauseanum	5			
	Hymenophyllum pectinatum	3			
	Hymenophyllum plicatum	3			
	Serpillopsis caespitosa	5			
Aspleniaceae	*Asplenium daeroides*	4			
	Pleurosorus papaverifolius	4		3	
Dryopteridaceae	*Megalastrum spectabile*	7			
Blechnaceae	*Blechnum mochaenum*	1			
	Blechnum arcuatum	1			
	Blechnum chilense	2	1	1	
	Blechnum magellanicum	1			
Dennstaedtiaceae	*Hypolepis rugosula*			1	
Total general		**201**	**22**	**20**	**0**

Figure 3. Rarefaction analysis showing the relationship between number of tree seedlings species (a) and small vascular plant species (b) as function of the number of individuals. Both panel the shaded area is indicating one standard error). All sites affected by the eruption of Chaitén Old-growth, Standing-dead and Blow-down are compared, except Total-destruction was not included because of the absence of plants.

Plant composition among Old-growth and Standing-dead plots showed some similarities (34%),

but less related with Blow-down (20%) and the Total-destruction plot (0%). Standing-dead and

Blow-down plots showed the highest similarities in species composition (Table 2). In the

disturbed plots, the plant species recorded were a subgroup of the species present in the Old-

growth forest plot (Nestedness calculator T= 13.85º). The most frequent plant was the fern *L.*

quadripinnata that survived the direct impact of the eruption in small clumps of organic matter

protected by trunks or root disc of trees, and were not completely covered by tephra. Bamboo

and ferns were sprouting from roots in the remnants of exposed forest soil.

Table 2. Sørensen similarity index for the plant composition among the different sites in the Chaitén influence area.

Site	Old-growth	Standing dead	Blow down	Total destruction
Old-growth	1			
Standing dead	0.342	1		
Blow down	0.2	0.526	1	
Total destruction	0	0	0.1	1

3.3. Forest structure
The forest structure of the Old-growth site showed individuals of all sizes, with several large

emergent trees (Fig. 4). Most trees in the Old-growth plot were alive and the snags most

probably represent trees that died before the eruption, while in the other plots most trees were

killed by the eruption. The Standing-dead plot showed similar dbh distribution than the Old-

growth plot, with higher frequency of small dbh. The Blow-down plot showed a very small

number of standing trees, concentrated in intermediate size class, while the Total-destruction

plot practically had no individual standing trees (Fig. 4).

Figure 4. Diameter at the breast height (dbh) distribution for living and standing dead trees in the study sites influenced by the eruption of the Chaitén volcano.

The basal area of the Old-growth plot was dominated by *L. philippiana,* followed by *E.*

cordifolia and *C. paniculata* (Table 3). In the Standing-dead and Blow-down plots *W.*

trichosperma and *N. dombeyi* were the dominant components of the basal area, but also with a

high number of unidentified trees (Table 3). We were not able to identify the species of the

standing dead trees in the Total destruction plot. The volume of fallen trees increased in the

Blow-down and in Standing-dead plots, but it was much lower in the Old-growth and the Total-

destruction plots (Fig. 5). In the Total-destruction plot most logs were heavily buried by a dense

tephra layer.

Table 3. Total basal area in 1000 m^2 of tree species of the study plots in the Chaitén volcano area, classified as alive, resprouting after the eruption, or snag

	Total-destruction	Blow-down		Standing-dead		Old-growth	
	Snag	Snag	Resprouting	Snag	Resprousting	Snag	Alive
Amomyrtus luma				0,02	0,34		0,47
Amomyrtus meli					0,06		
Caldcluvia paniculata							2,54
Drimys winterii				0,14			
Lomatia ferruginea				0,,04	0,01		0,02
Myrceugenia planipes							0,84
Eucryphia cordifolia		0,55		0,59	0,61	0,26	6,37
Laureliopsis philippiana		0,25		0,87			9,64
Nothofagus dombeyi				3,46			
Weinmannia trichosperma		0,95	0,41		2,38		0,71
not identified	1,41	0,94		1,45			
Total	**1,41**	**2,69**	**0,41**	**6,57**	**3,39**	**0,26**	**20,59**

Figure 5. Volume of fallen dead trees (logs) in the study sites influenced by the eruption of the Chaitén volcano in the study sites.

3.4. Seedlings and small plants

We found a total of 90 seedlings from 5 trees and shrubs species in the disturbed plots, while

we registered 174 seedlings of 12 species in Old-growth plots. The Standing-dead plots had 83

seedlings from five species, while the Blow-down plots presented 7 seedlings from two species.

We did not find any seedling in the total destruction plot. In the Old-growth plots, seedlings

were largely dominated by the shade tolerant *A. luma*, *M. parviflora* and *M. planipes*. In the

Standing-dead plots, seedlings were dominated by *C. paniculata* (48% of the total), but also

present the shade-tolerant tree *A. luma* (24%)*,* the pioneer tree species *W. trichosperma* and *E.*

coccineum (18% and 8% respectively). In the Blow-down plot, *E. coccineum,* a pioneer species,

and *A. lanceolata* were the only recorded species.

We found different mechanism of forest regeneration in the plots. In the Old-growth plot most

regeneration was from seeds, while in the Standing-dead plots most regeneration become from

resprout and few tree seedlings from seeds (Table 4). In the Old-growth plots, most of the tree

seedlings corresponded to species with fleshy fruits dispersed by birds, while in the Standing-

dead plots most seeds were wind-dispersed. In the Blow-down plots, all seedlings came from

seeds (Table 4), but a half of the seedlings corresponded to species dispersed by birds (e.g.

Ribes magellanicum) and the other half were species dispersed by wind (e.g. *E. coccineum*).

Second, in all plots most seedlings were growing on organic substrata (Table 4), such as organic

soil on the ground, root discs and logs, while few seedlings were growing directly either on the

tephra or on the rocks. In the Blow-down plot we found 7 seedlings of *E. coccineum*, growing

just in remnants of organic litter; in Standing-dead plot 11 seedlings growing directly in the

tephra, while the other 72 seedlings were resprouting directly from the standing tree's trunks.

In the Old-growth plots we found just one species growing in an exposed rock while the rest of

the seedlings were growing in organic substrates (Table 4).

Table 4. Number of seedlings per regeneration type and substratum type in the plots.

	Regeneration type		Substratum type	
	Resprouting	Seed	Mineral	Organic
Total-destruction	-	-	-	-
Blow-down	-	7	-	7
Standing-dead	74	9	11	72
Old-growth	31	127	1	157

Other small plant species in the studied plots (Table 1) showed differences along the disturbance

gradient (Table 2). For example, the Old-growth plot has species in all categories of plant cover,

with few species covering over 75% of the ground, and many species covering between 1-5%

(Fig. 6). In contrast, Standing-dead and Blow-down plots were covered by fewer species, and

concentrated in the lower ranges of coverage (Fig. 6). In the Total destruction plot we only

found a non-vascular plant (*Marchantia* sp.) with a cover between 5-25 % (Fig. 6) growing on

a log exposed to the surface. The dominant understory species in the Old-growth forest were

the bamboo *Chusquea* spp. and the fern *L. quadripinnata.* These species were also present in

the Standing-dead and Blow-down plots, although scattered and with a low percent of coverage

in the disturbed plots.

Figure 6. Plant cover (including vascular and non-vascular species) at the ground level of the study areas affected by the Chaitén volcano.

4. Discussion

4.1 Early vegetation establishment: the relevance of biological legacies

Few studies have been conducted short after an eruption in the southern Andes, however successional model predict that disturbed areas in low and mid altitude (<1000 m a.s.l.) forests would be dominated by *Nothofagus* species due to the shade-intolerance (e.g. Veblen and Ashton, 1978). However, in this study after the first year we did not register *Nothofagus* species despite were abundant in the surroundings non disturbed areas. It might be due to cyclic seed production with not constant seed rain along the years (Donoso, 1993; Veblen, 1985), and the short-term and limited spatial extension we were able to monitor the vegetation. However, our results pointed that regeneration occurs mainly due to an *in situ* development of plants associated to biological legacies such as logs, soil clumps and individuals that survive to the eruption. Other areas disturbed by Chaiten volcano presented similar patterns of types and abundance of biological structures (Swanson et al., 2013).

The initial establishment of the tree species, beside of the disturbance gradient, may be influenced by the interaction between life history of the species and the altitudinal gradient. Life history attributes such as seed dispersal, flowering phenology, growth form, resprouting ability or light-temperature stress resistance are a mechanism that influence the recovery response of the post-disturbance vegetation (Eriksson 2000; McIntyre et al., 1994; McLachlan and Bazely, 2001; Pausas and Lavorel, 2003; Tilman, 1997). In the disturbed plots, the principal mechanism of regeneration was the resprouting capability of some species (Tables 1 and 4). This is a frequent strategy in many plant species in response to disturbances of different types and intensities (Bellingham and Sparrow, 2000; Bond and Midgley, 2001). The seed dispersion syndrome could be also an important mechanism that influence the colonization after disturbance. Chilean forests have one of the highest proportions of plant dispersed by

frugivorous birds compared to other temperate rainforests (Armesto et al. 1987, Willson et al.,

1996). Birds consume a high proportion of fruits and may disperse the seed to specific places,

such as to a perch (Armesto et al., 2001; Lindenmayer and Franklin, 2002). The seed availability

might increase in sites where remnant snags, shrubs or perching trees where available (e.g.,

Bustamante-Sánchez, 2011). Finally, among the wind dispersed seeds, we only found seedlings

of *E. coccineum* in disturbed plots. While this tree is adapted to stressful condition, the seed is

able to disperse for long-distances (Donoso, 2006).

For non-tree species, Old-growth plot had more species than Standing-dead plot. On the

contrary, for tree species Standing-dead plot is richest than Old-growth, revealing the effect of

the eruption on the small plant species. On the other hand, in Blow-down and in Total-

destruction plot, i.e. from mid to high disturbance intensity at mid to high altitude, and without

or sparse biological legacies, we found low species richness and coverage, and sparse tree

regeneration. This mosaic of legacies in the landscape offers opportunities for the

recolonization of many different species in the affected area. The different species composition

would promote different pathways to early succession with different communities, but also

different recovery rates. We propose that in low to mid elevation areas, the vegetation recovery

will be faster, dominated by species shade-tolerant to semi-tolerant and dispersed by birds,

mostly due to biological legacies. On the other hand, at higher altitude, the vegetation recovery

would be slower, recolonization made by shade-intolerant and seed dispersed by winds because

the most stressful conditions of temperature at higher altitudes.

In synthesis, the results point to an early succession with (i) a mosaic of pioneering-wind

dispersed species, scattered survivors regrowing and spreading from biological legacies, and

plant species dispersed by frugivorous birds, likely favored by the biological legacies; (ii) the

early succession is influenced by the interaction of the species-specific life history (including

invasion by exotic species), altitudinal gradient and the different intensity of disturbance in a

broad area.

Acknowledgements

350 Field work was financed by CONICYT PDA-24. A.L. and D.C. acknowledge support from

351 grant CONICYT/FONDAP/15110009. RM acknowledge the scholarship DAAD/Becas Chile

352 during late stages of this manuscript.

REFERENCES

Armesto, J. J., Rozzi, R., Miranda, P., & Sabag, C. (1987). Plant/frugivore interactions in South American temperate forests. Revista Chilena de Historia Natural, 60(2), 321-336.

Aravena, J.C., Carmona, M., Perez, C.A., Armesto, J.J., 2002. Changes in tree species richness, stand structure and soil properties in a successional chronosequences in northern Chiloé Island, Chile. Revista Chilena de Historia Natural. 75, 339-360.

Armesto, J.J., Arroyo, M.T.K., Villagran, C., 1996. Ecología de los Bosques Nativos de Chile. Editorial Universitaria, Chile. 470pp.

Armesto, J.J., Díaz, I.A., Papic, C., Willson, M.F., 2001. Seed rains of fleshy and dry propagules in different habitats in the temperate rainforest of Chiloé Island, Chile. Austral Ecology 26, 311-320.

Armesto, J.J., Pickett, S.T.A., 1991. Foreword. Revista Chilena de Historia Natural 644, 389.

Arroyo, M.T.K., P. Marquet, C. Marticorena, J. Simonetii, L. Cavieres, F.A. Squeo & R. Rozzi, 2004. Chilean winter rainfall-valdivian forests. In (Mittermeier, R.A., P. Robles Gil, M. Hoffmann, J. Pilgrim, T. Brooks, C. Goettsch, J. Lamoreux & G.A.B. da Fonseca, eds), "Hotspots Revisited", pp. 99-103. CEMEX, México.

Atmar, W., Patterson, B.D., 1993. The measure of order and disorder in the distribution of species in fragmented habitat. Oecologia 96, 373-382.

Atmar, W., Patterson. B.D., 1995. The nestedness temperature calculator: a visual basic program, including 294 presence-absence matrices. AICS Research, Inc., University Park, NM, and The Field Museum, Chicago.

Bellingham, P., Sparrow. A.D., 2000. Resprouting as a life-history strategy in woody plant communities. Oikos 89(2), 409-416.

Bond, W.J., Midgley, J.J., 2001. Ecology of sprouting in woody plants: the persistence niche. Trends in Ecology & Evolution 16(1), 45-51.

Bustamante-Sánchez, M.A., Armesto, J.J., Halpern, C.B., 2011. Biotic and abiotic controls on tree colonization in three early successional communities of Chiloé Island, Chile. Journal of Ecology 99(1), 288-299.

Chazdon, R.L., 2003. Tropical forest recovery: legacies of human impact and natural disturbances. Perspective in Plant Ecology, Evolution and Systematics 6, 51-71.

Colwell, R.K., Chao, A., Gotelli, N.J., Lin, S., Mao, C., Chazdon, R.L., Longnio, J.T., 2012. Models and estimators linking individual-based and sample-based rarefaction, extrapolation and comparison of assemblages. Journal of Plant Ecology 5, 3-21.

Colwell, R.K., and Coddington, J.A., 1994. Estimating terrestrial biodiversity through
extrapolation. Phil. Trans. R. Soc. Lond. B 345: 101-118.

Crews, T., Kitayana, K., Fownes, J., Riley, R., Herber, D., Mueller-Dombois, D., Vitousek,
P., 1995. Changes in soil phosphorusfractions and ecosystem dynamics across a long
chronosequence in Hawaii. Ecology 76: 1407-1424

Dale V.H., Swanson, F.J., Crisafulli, C.M., 2005. Ecological responses to the 1980 eruption
of Mount St. Helens. Springer, 342pp.

del Moral, R. 1981. Initial life returns to Mount St. Helens. Natural History 90, 36-49.

del Moral, R., Grishim, S., 1999. Volcanic Disturbance and Ecosystem recovery, in: L.R.
Walker (Ed.) Ecosystem of disturbed Ground. Elsevier, pp. 137-160.

Dinerstein, E., Olson, D., Graham, D., Webster, A., Primm, S., Bookbinder, M., Ledec, G.,
1995. A conservation assessment of the terrestrial ecoregions of Latin America and the
Caribbean. The International Bank for Reconstruccion and Development. Waschington, DC:
The World Bank.

Donoso, C., 1993. Bosques templados de Chile y Argentina, variación, estructura y dinámica.
Tercera edición. Editorial Universitaria, Santiago-Chile, 484p.

Donoso, C., 2006. Las Especies arbóreas de los Bosques Templados de Chile y Argentina.
Autoecología. Marisa Cúneo Ediciones, Valdivia-Chile, 678p.

Donoso, C., Lara, A., 1999. Silvicultura de los Bosques Nativos de Chile. Editorial
Universitaria, Santiago-Chile, 421p.
420
Donoso PJ, Promis A (2013) Silvicultura en Bosques Nativos. Avances en la investigación en
Chile, Argentina y Nueva Zelandia. Estudios en Silvicultura de Bosques Nativos, Vol. 1. Ed.
Marisa Cuneo, Valdivia, Chile. p 253

Eriksson, O., 2000. Funtional roles of remnants plant population in communities and
ecosystem. Global Ecology and Biogeography 9, 443-449.

Foster, D., Knight, D., Franklin, J.F., 1998. Landscape patterns and legacies resulting from
Large, Infrequent forest disturbance. Ecosystems 1, 497-510.

Franklin, J.F., 1990. Biological Legacies: a critical Management concept from Mount St.
Helens. Transactions 55th North American Wildlife and Natural Resources Conference, 216-219.

Franklin, J.F., Armesto, J.J., 1996. La retención de elementos estructurales del bosque durante
la cosecha: Una alternativa de manejo para los bosques nativos chilenos. Ambiente y
Desarrollo 12 (2), 69-79.

Franklin, J. F., MacMahon, J. A., Swanson, F. J., Sedell, J. R., 1985. Ecosystem responses to
the eruption of Mount St. Helens. National Geographic Research 1:198-216.

Franklin, J.F., Mitchell, R.J., Palik, B.J., 2007. Natural disturbance and Stand development
principles for ecological forestry. United States Department of agriculture, Forest Service,
Northern Research Station. General Report NRS-19. 48p.

Franklin, J.F., Lindenmayer, D., MacMahon, J.A., McKee, A., Magnuson, J., Perry, D.A.,
Waide, R., Foster, D., 2000. Threads of continuity: ecosystem disturbance, recovery, and the
theory of biological legacies. Conservation Biology in Practice 1, 8-16.

Franklin, J.F., Spies, T.A., van Pelt, R., Carey, A., Thornburgh, D.A., Rae Berg, D.,
Lindenmayer, D.B., Harmon, M.E.,Keeton, W.S., Shaw, D.B., Bible, K., Chen, J., 2002.
Disturbance and structural development of natural forest ecosystem with silvicultural
implications, using Douglas-fir forest as an example. Forest Ecology and Management 155,
399-423.

Franklin, J.R., Swanson, M.E., 2010. Long-term ecological research in the forest of the
United States: key lessons for its application in Chile and around the world. Revista Chilena
de Hitoria Natural 83, 185-194.

Gallardo, M.B., Pérez, C., Nuñez-Avila, M., Armesto, J.J. 2012. Desacoplamiento del
desarrollo del suelo y la sucesión vegetal a lo largo de una cronosecuencia de 60 mil años en
el volcán Llaima, Chile. Revista Chilena de Historia Natural 85: 291-306

Gotelli, N.J., Entsminger G.L., 2005. EcoSim: Null models software for ecology. Version 5.0.
Acquired Intelligence Inc. & Kesey-Bear.
http://homepages.together.net/~gentsmin/ecosim.htm.
455
Grosse, H. y Quiroz, I. 1999. Silvicultura de los bosques de segundo crecimiento de roble, raulí
y coigüe en la región centro-sur de Chile. p 95-125. Silvicultura de los bosques nativos de Chile.
Editorial Universitaria. 421p

Haberle, S.G., Szeicz, J.M., Bennett, K.D., 2000. Late Holocene vegetation dynamics and
lake geochemistry at Laguna Miranda, XI Region, Chile. Revista Chilena de Historia Natural
73, 655-669.

Johnson, E.A., Miyanishi, K., 2008. Testing the assumptions of chronosequences in
succesion. Ecology Letters 11: 419-431.

Kitzberger, T., Veblen, T.T, Villalba, R., 1995. Tectonic influences on tree growth in northern
Patagonia, Argentina: the roles of substrate stability and climatic variation. Canadian Journal
of Forest Research 25, 1684-1696

MacArthur, R.H., Wilson, E.O., 1967. The theory of island biogeography. Princeton
University Press, Princeton

Major, J., Lara, L.E., 2013. Overview of Chaiten Volcano, Chile, and its 2008-2009 eruption.
Andean Geology 40(2), 196-215.

McIntyre, S., Lavorel, S., Tremont, M., 1994. Plant Life-History2 Attributes: Their
relationship to disturbance response in herbaceous vegetation. Journal of Ecology 83(1), 31-
44.

McLachalan, S.M., Bazely, D.R., 2001. Recovery patterns of understory herbs and their use
as indicators of deciduos forest regeneration. *Conservation Biology* 15, 98-110.

Müller-Dombois, D., Ellenberg, H., 1974. Aims and methods in vegetation ecology. The
Blackburn Press. 580p.

Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, A.B., Kent, J., 2000.
Biodfiversity hotspot for conservation priorities. Nature, 402, 853-58.

Lara, L.E., 2009. The 2008 eruption of the Chaitén Volcano, Chile: a preliminary report.
Andean Geology 36(1), 125-129.

Lindenmayer, D.B., Franklin, J.F., 2002. Conserving Forest Biodiversity: A Comprehensive
Multiscaled Approach. Island Press, Washington, 351p.

Luebert, F., Pliscoff, P., 2006. Sinopsis bioclimática y vegetacional de Chile. Editorial
Universitaria, Santiago-Chile, 316p.

Pausas, J.G., Lavorel, S., 2003. A hierchical deductive approach for functional types in
disturbed ecosystem. Journal of Vegetation Sciences 14(3): 409-416.

Peet, R., 1992. Community structure and ecosystem function, in: Glenn-Lewin D., Peet R.,
Veblen, T. (Eds.), Plant succession theory and prediction. Chapman & Hall, London, pp. 103-
151.

Perakis, S.S., Hedin L., 2002. Nitrogen loss from unpolluted South American forest mainly
via dissolved organic compounds. Nature 415, 416-419.

Pickett, S.T.A., Collins, S.L., Armesto, J.J., 1987. Models, mechanism and pathways of
succession. The Botanical Review 53(3), 335-371.

Pickett, S.T.A., 1988. Space-for-time substitution as an alternative to long term studies, in:
Likens, G.E. (Ed.), Long-term Studies in Ecology. Springer, New York, NY, pp. 110–135.

Robock, A., Oppenheimer, C., 2003. Volcanoes and Earth's Atmosphere. Geophysical
monograph 139.

Runkle, J.R., 1985. Disturbance regimes in temperate forests, in: Pickett, S.T.A., White, P.S.
(Eds.), The Ecology of Natural Disturbance and Patch Dynamics. Academic Press, Orlando,
FL, pp. 17-33

Stern, C.R., 2004. Active Andean volcanism: its geologic and tectonic setting. Andean
Geology. 31(2), 161-206

Swanson, F., Franklin, J., 1992. New Forestry principles from ecosystem analysis of Pacif
Northwest Forest. Ecological applications 2(3), 262-274

Swanson, F., Jones, J., Crisafulli, Ch., Lara A., 2013. Effects of volcanic and hydrology
processes on forest vegetation: Chaiten Volcano, Chile. Andean Geology 40(2), 359-391

Swanson, M., Franklin, J., Beschta, R., Crisafulli, C., DellaSala, D., Hutto, R., Lindenmaver,
D., Swanson, F., 2011. The Forgotten stage of of forest succession: Early-successional
ecosystems on forest site. Biological Sciences Faculty Publications. Paper 278.
http//scholarworks.s.umt.edu/biosci_pubs/278

Tilman D., 1997. Community invasibility, recruitment limitation, and grassland biodiversity.
Ecology 78, 81–92
Turner, M., Dale, V., 1998. Comparing large, infrequent disturbance: what have we learned?.
Ecosystem 1, 493-496.

Turner, M., Dale, M., Everham, E., 1997. Fires, hurricanes, and volcanoes: Comparing large
disturbances. BioSciences 47(11), 758-768.

Veblen, T.T., 1985. Stand dynamics in Chilean *Nothofagus* forest, in: Pickett, S.T.A., White,
P.S. (Eds.), The ecology of natural disturbance and patch dynamics. Academic Press,
Orlando, FL, pp 35-51.

Veblen, T.T., Ashton, D.H., 1978. Catastrophic influences on the vegetation of the Valdivian
Andes, Chile. Vegetatio 36(3): 149-167.

Veblen, T.T., Ashton, D.H., Schlegel, F., Veblen, A., 1977. Plant succession in a timberline
depressed by vulcanism in South-Central Chile. Journal of Biogeography 4: 275-294.

Veblen, T.T., Donoso, C., 1987. Alteración natural y dinámica regenerativa de las especies
chilenas de *Nothofagus* de la Región de Los Lagos. Bosque 8(2), 132-142.

Veblen, T.T., Donoso, C., Kitzberger, T., Rebertus, A.J., 1996. Ecology of southern Chilean
and Argentinean Nothofagus forests, in: Veblen T.T., Hill, R.S., Read, J., (Eds.) The ecology
and biogeography oí Nothofagus forests. Yale University Press, New Haven, Connecticut,
USA, pp. 293-353

Veblen, T.T., Donoso, C., Schlegel, F., Escobar, B., 1981. Forest dynamics in South-Central
Chile. Journal of Biogeography 8, 211-247.

Veblen, T.T., Schlegel, F., Oltremari, J., 1983. Temperate broad-leaved evergreen forest of
South America, in: Ovington, J.D., (Ed.) Temperate broad-leaved evergreen forest. Elsevier,
Amsterdam, pp 5-31.

Villagran, C., Hinojosa, F., 1997. Historia de los Bosques Templados de Sudamerica, II:
Análisis fitogeográfico. Revista Chilena de Historia Natural, 70, 241-267.

Willson, M.F., De Santo,T. L., Sabag, C. E., Armesto, J.J., 1996. Avian communities in
temperate rainforest of North and South America. In: Lawford, R., Alaback, P. B., Fuentes,
E., (Eds.) High-latitude rainforests and associated ecosystems of the west coast of the
Americas. Springer, New York. pp. 228–247