Spooky interaction at a distance in cave and surface dwelling electric fishes

Eric S. Fortune[1,*]**, Nicole Andanar**[1]**, Manu Madhav**[2]**, Ravi Jayakumar**[2]**, Noah J. Cowan**[2]**, Maria Elina Bichuette**[3]**, and Daphne Soares**[1]

[1]*New Jersey Institute of Technology, Biological Sciences, Newark, 07102, USA*
[2]*Johns Hopkins University, Whiting School of Engineering, Baltimore, 21218, USA*
[3]*Universidade Federal de São Carlos, Departamento de Ecologia e Biologia Evolutiva, São Carlos, 13565-905, Brasil*

Correspondence*:
Eric Fortune
eric.fortune@njit.edu

2 ABSTRACT

Glass knifefish (*Eigenmannia*) are a group of weakly electric fishes found throughout the Amazon basin. We made recordings of the electric fields of two populations of freely behaving *Eigenmannia* in their natural habitats: a troglobitic population of blind cavefish (*Eigenmannia vicentespelaea*) and a nearby epigean (surface) population (*Eigenmannia trilineata*). These recordings were made using a grid of electrodes to determine the movements of individual fish in relation to their electrosensory behaviors. The strengths of electric discharges in cavefish were larger than in surface fish, which may be a correlate of increased reliance on electrosensory perception and larger size. Both movement and social signals were found to affect the electrosensory signaling of individual *Eigenmannia*. Surface fish were recorded while feeding at night and did not show evidence of territoriality. In contrast, cavefish appeared to maintain territories. Surprisingly, we routinely found both surface and cavefish with sustained differences in electric field frequencies that were below 10 Hz despite being within close proximity of less than one meter. A half century of analysis of electrosocial interactions in laboratory tanks suggest that these small differences in electric field frequencies should have triggered the jamming avoidance response. Fish also showed significant interactions between their electric field frequencies and relative movements at large distances, over 1.5 meters, and at high differences in frequencies, often greater than 50 Hz. These interactions are likely envelope responses in which fish alter their EOD frequency in relation to changes in the depth of modulation of electrosocial signals.

Keywords: Gymnotiformes, weakly electric fish, troglobitic, epigean, envelope, cavefish, jamming avoidance response, diceCT

1 INTRODUCTION

Gymnotiformes are a group of nocturnal fishes characterized by a suite of adaptations which allow them to localize conspecifics (Davis and Hopkins, 1988) and capture prey (Nelson and MacIver, 1999) in complete darkness. These fishes produce a weak electric field, typically less than 100 mV/cm (Assad et al., 1998), using an electric organ located along the sides of the animal and in its tail (Heiligenberg, 1991). This electric field, known as the electric organ discharge or EOD, is detected using specialized electroreceptors embedded in the skin. These receptors encode modulations generated through interactions with the electric

fields of conspecifics and by nearby objects. This system provides a mechanism for communication among
conspecifics and for the detection and characterization of prey and other salient environmental features
(Nelson and MacIver, 1999; Crampton, 2019; Pedraja et al., 2018; Yu et al., 2019).

The nocturnal life histories of Gymnotiform species make them well suited for life in caves. Interestingly,
a single species of Gymnotiform fish, *Eigenmannia vicentespelaea*, has been discovered in a cave system
in central Brazil (Triques, 1996; Bichuette and Trajano, 2006, 2017). These fish exhibit features commonly
found in species adapted to life in caves, including reduced pigmentation and reduction and/or elimination
of the eyes (Culver and Pipan, 2019). The population is estimated to be around only 300 individuals
(Bichuette and Trajano, 2015).

Fish in the genus *Eigenmannia* produce quasi-sinusoidal electric signals that are maintained at frequencies
between about 200 and 700 Hz (Heiligenberg, 1991). Individual *Eigenmannia* change their electric field
frequencies in response to electrosensory signals produced by nearby conspecifics (Watanabe and Takeda,
1963). In the Jamming Avoidance Response (JAR), individuals raise or lower their electric field frequency
to avoid differences of less than about 10 Hz (Heiligenberg, 1991). Fish also change their electric field
frequencies in relation to the relative movement between individuals that are encoded in the amplitude
envelope of their summed electric fields (Stamper et al., 2013; Metzen and Chacron, 2013; Thomas et al.,
2018; Huang and Chacron, 2016). Envelope responses can also occur in groups of three or more fish when
the pairwise differences of their electric field frequencies are close, with 1-8 Hz, of one another (Stamper
et al., 2012).

To discover the potential consequences of adaptation for troglobitic life on electrosensory behavior, we
compared the electric behavior and movement of the cavefish *Eigenmannia vicentespelea* to nearby epigean
(surface) relatives, *Eigenmannia trilineata*, that live in the same river system (Fig. 1A). We used a recently
developed approach for characterizing electric behaviors and locomotor movements of weakly electric
fishes in their natural habitats (Madhav et al., 2018; Henninger et al., 2020). This approach, which uses a
grid of electrodes placed in the water, permits an estimation of the electric field of each fish and an analysis
of their concurrent movement.

2 METHODS

54 These observational studies were reviewed and approved by the Rutgers Institutional Animal Care and
55 Use Committee and follow guidelines established by the National Research Council and the Society for
56 Neuroscience.

2.1 Study sites

The study sites were located in Terra Ronca State Park (46° 10'–46° 30' S, 13° 30'–13° 50' W), in the
Upper Tocantins river basin, state of Goiás, central Brazil (Fig. 1A). We measured the electric behavior
of the cavefish *Eigenmannia vicentespelaea* in the São Vicente II cave (13° 58'37" S, 46° 40'04" W) in
October of 2016 (Fig. 1B). The electric behavior of the epigean species, *Eigenmannia trilineata*, was
measured in the Rio da Lapa at the mouth of the Terra Ronca cave (13° 38'44" S; 46° 38' 08" W) in April
of 2016 (Fig. 1B). These streams have moderate water currents, clear water with conductivity below 20 μS,
and the substrate is composed of sand, rocks, and boulders.

2.2 Anatomy

Four alcohol fixed specimens from the collection at the Universidade Federal de São Carlos (Dr. Bichuette)
were submerged in 11.25 Lugol's iodine (I2KI) solution for up to 36 hours prior to diffusable iodine based
contrast enhanced computer tomography (DiceCT). Stained specimens were removed from Lugol's solution,

rinsed in water to remove excess stain and sealed in rubber sleeves to prevent dehydration. Samples were
then loaded into 50 mL polypropylene centrifuge tubes for scanning.

Stained and unstained specimens were scanned at the Core Imaging Facility of the American Museum
of Natural History (New York, NY), using a 2010 GE Phoenix v-tome-x s 240CT high-resolution
microfocus computed tomography system (General Electric, Fairfield, CT, USA). DiceCT scanning
permits visualization of soft tissue details of the head and the body. Scans were made at 125 kV, with an
exposure time of 60 seconds. Voxel sizes were 20.0-25.9μm. Volume reconstruction of raw X-ray images
were achieved using a GE Phoenix datos—x.

2.3 Recordings of electric behavior at field sites

Eigenmannia were initially identified and located using hand-held single-electrode probes with a custom
amplifier/speaker system. This river system also permits direct visualization of the animals - the water
is sufficiently clear and free of debris to see and photograph the fish (Hero Cam, GoPro 3, USA) when
swimming in open water (see Supplemental videos S1, S2). Electric recordings were made using an array
of amplified electrodes (50 cm between electrodes) (Madhav et al., 2018). For measurements of the epigean
fish, an array of 8 electrodes was placed along the edges of the Rio da Lapa stream after sundown when the
fish were active. In the São Vicente II cave, an array of 16 electrodes was placed in eddies and side pools
along the primary stream. The flow at the center of the stream was too strong for grid array. Unfortunately,
we were unable to use the larger grid on our second visit to the surface site due to a concurrent religious
festival. As a result, the maximum XY range of the surface grid is about 100 cm diameter smaller than the
cave grid. In all other respects, the measurements from each grid are identical.

We used an algorithm (Madhav et al., 2018) (code available at doi:10.7281/T1/XTSKOW) to identify
each *Eigenmannia* using its time-varying fundamental frequency. We calculated an estimate of the strength
of the electric field for each fish. Fish were considered to be ideal current dipoles—a source-sink pair of
equal but time varying strength $I(t)$, separated by a small distance d. The electric current dipole moment
for the fish is defined as $p = Id$ which has the units of Ampere-meter or "Am." We calculated an estimate
of the position and pose of each fish. Position and pose were calculated in relation to the distribution of
power at each EOD frequency across the grid of electrodes. In these recordings, which were made in
shallow water of no more than 40 cm depth with a planar array of electrodes, the position estimates were
restricted to the XY-plane.

Continuous recording sessions using the grid were made both at the cave site ($N = 14$) and surface site
($N = 5$) of at least 600 seconds to over 1200 seconds duration. Intervals between recording sessions ranged
from 5 minutes to several hours. Because fish could not be tracked between recording sessions, it is likely
that some individual fish were measured across recording sessions.

2.4 Analysis of position and EOD frequency data

The position and EOD frequency data were analyzed, unless otherwise described, in 300 second epochs
with 50% overlap. All analyses were conducted using custom scripts in Matlab (Mathworks, Natick MA).
These data and scripts will be made freely available upon publication of this manuscript (URL). We
estimated the XY region of movement for each fish for each epoch using a minimum convex polygon fitted
to its positions. We then calculated the pairwise overlap between each convex polygon. Euclidean distances
between pairs of fish were computed as a function of time during each 300 second epoch.

To assess the relations between pairs of EOD frequencies and their relative movement, we calculated
Pearson correlations between 1) instantaneous distance between pairs of fish (distance) and 2) instantaneous
difference in EOD frequency (dF). Pearson correlations were measured over 300 second epochs with

150 second overlap between epochs. These "dF/distance correlations" ranged from -0.93 to 0.90. The values for dF/distance Pearson correlations matched our impressions of the strengths of relations between dF and distance by visual inspection of the data. To estimate the rate of spontaneous correlations between dF and distance, we shuffled (Matlab randperm) the epochs of dF and distance measurements.

3 RESULTS

3.1 General observations

Eigenmannia at both the surface and cave sites were found in clear water streams with rock and sand substrates (Bichuette and Trajano, 2003, 2006, 2015, 2017). At the surface site, there was a marked diurnal modulation of behavior. During the day, fish were found alone or in groups along the banks of the Rio da Lapa, typically below or around boulders and rocks. The grid system was not used to make recordings of the fish during the day due to a local festival. Nevertheless, the distribution of *Eigenmannia* during the day appeared to be similar to that described previously at sites in Ecuador (Tan et al., 2005). Unlike in previous measurements at other study sites (Stamper et al., 2010), no other Gymnotiform species were detected in our short survey. At night, we observed *Eigenmannia* swimming in open water in the center of the stream, typically near the bottom.

Surface fish foraged in sandy substrates. Foraging behavior included hovering with slow forward or backward swimming punctuated by strikes into the substrate (see Supplemental video S1). These strikes involved tilting of the head and body downwards with a rapid forward lunge to drive the mouth into the sand a few millimeters. These strikes differ from feeding behavior described in *Apteronotus albifrons* in which fish captured freely swimming *Daphnia*, typically from below the prey (Nelson and MacIver, 1999). We observed groups of *Eigenmannia* simultaneously foraging with inter-fish distances on the order of 10s of centimeters. These distances suggest that fish experience significant ongoing electrosensory interference from conspecifics. At other locations, we visually observed single fish foraging while swimming rather than hovering a particular location.

At the cave site, fish did not exhibit diurnal modulation of behavior. Cavefish were observed along the banks of the stream and in small eddies and pools. Fish were alone or in small groups of up to 10 individuals. Cavefish retreated to crevices and spaces within boulders and rocks when disturbed, forming temporary aggregates of individuals (see Supplemental video S2). Videos show eyeless cavefish orienting face-to-face during social interactions. Such movements, in the absence of visual cues, must be controlled, at least in part, using electrosensory signals. The distinctive substrate foraging behavior routinely seen in the surface fish was not observed in these cavefish.

3.2 Morphology

Surface fish (*Eigenmannia trilineata*) had large eyes that were circumferential and of the same size (Fig. 1C,E) whereas the cavefish (*Eigenmannia vicentespelaea*) had eyes in various states of degeneration, from microphthalmic to completely absent (Fig. 1D,F). Preliminary review of scans of four fish showed potential differences in the size of the electric organ (Fig. 1E,F), but additional material will be necessary for quantitative analysis of electric organ structure and physiology.

A prior study found that surface fish are smaller than the cavefish (Bichuette and Trajano, 2006): mean length of the snout to the end of the anal fin base was reported to be 8.45 cm (sd=2.67) in surface fish and 11.1 cm (sd=2.47) in cavefish. Size is important as it likely impacts the strength of electric fields: larger fish typically can generate larger currents in their electric organs.

3.3 Electric field amplitudes

The grid recording system permits an estimation of the strength of each individual's electric field using Fourier analysis. The strength of electric fields (Ampere-meter, "A-m") of surface fish were significantly lower than cavefish (Wilcoxon sign-rank, two-sided, z=2.98, p=0.0029) with mean strength of 6.5×10^{-4} A-m for surface fish and 9.66×10^{-4} A-m) for cavefish (Fig. 3B). The surface fish are generally shorter than the cavefish *Eigenmannia* and also appear to have relatively smaller electric organs. The increase in EOD amplitude may be an energetically costly adaptation (Markham et al., 2016) to life in this cave: larger EOD amplitudes would increase the ability to detect objects and capture prey (Nelson and MacIver, 1999) in the absence of visual cues.

3.4 Electric field frequencies

We used the grid recording system to calculate the EOD frequency of each fish within the grid (Fig. 2). EOD frequencies of surface fish were between 299.9 Hz and 435.6 Hz (N=110) whereas EOD frequencies of cavefish were between 230.0 Hz and 478.6 Hz (N=82) (Fig. 3A). The distribution of EOD frequencies of surface and cavefish are significantly different (Wilcoxon sign-rank, two-sided, z=2.57, p=0.0100). The mean and median frequencies of surface fish were 375.8 Hz and 382.0 Hz, and for cavefish were 356.6 Hz and 360.8 Hz. There was a dramatic diurnal modulation of behavior in surface fish, but grid data were collected only at night. The electric fields of some species of Gymnotiform fishes have been shown to have diurnal modulations and amplitude and frequency content (Stoddard et al., 2006; Markham et al., 2009; Migliaro et al., 2018; Sinnett and Markham, 2015). We have not observed diurnal modulation of EOD frequencies of *Eigenmannia* in previous field studies (Tan et al., 2005; Stamper et al., 2010).

We observed variations in EOD frequencies that likely include social signals such as chirps and envelope responses (Fig. 2C,D). The variance of EOD frequencies in freely swimming groups of surface fish (mean=0.64 Hz, n=184) and cavefish (mean=1.10 Hz, n=292) were significantly different (T-test, p<0.0001, df=467, tstat=6.1). We expect the variance of EOD frequencies in surface fish will differ during daylight hours, when the fish are hiding along the shores of the rivers, in relation to night hours when the fish are active. Additional recordings made during the day are needed.

3.5 Differences in EOD frequencies

The JAR is a behavior that, in laboratory settings, reduces the likelihood that pairs fish will have a difference in EOD frequency (dF) of less than 10 Hz (Heiligenberg, 1991). In surface fish the mean dF was 40.32 Hz (sd=26.90, N=2251) while the cavefish the mean dF was 78.38 Hz (sd=47.81, N=3908) (Fig. 4A). The distribution of mean dFs was significantly larger in cavefish than surface fish (T-test p<0.0001, df=6157, tstat=34.74). These findings are consistent with a role of the JAR in maintaining larger dFs between most individuals, but may also simply reflect the greater density of fish that we observed at the surface site.

Interestingly, we routinely found nearby fish with sustained, lasting for hundreds of seconds, differences in electric field frequencies that were below 10 Hz (Fig. 4B). In surface fish, there were 294 pairs with mean dFs of less than 10 Hz over 300 second duration epochs, and of those, 142 were less than 5 Hz (out of total 1947 pairwise interactions). In cavefish, there were 18 epochs with mean dFs that were less than 10 Hz, of which 4 were less than 5 Hz (out of total 435 pairwise interactions). Low dFs between pairs of fish were at distances within 50 cm (Fig. 4C-E).

3.6 Similarities and differences in movement patterns between surface and cavefish

During daylight hours, surface fish were found deep within crevices and other refugia along the sides of the Rio da Lapa, but emerged after dark for feeding and social behavior that occurred in open water

throughout the stream. In contrast, cavefish were found swimming with no apparent variation with respect to time of day along the sides of streams and in pools and eddies.

We estimated the position of each fish in and around the grid over time (Madhav et al., 2018). Cavefish appeared to swim within small regions or territories on the order of tens of centimeters in diameter. In contrast, surface fish appeared to have widely overlapping swimming trajectories. To examine the relative movements of fish, we divided the data into 300 second duration epochs and fitted a minimum convex polygon to each fish's positions.

Although there was no difference in the overall size of the convex polygons between surface and cavefish (T-test, p=0.5840, df= 499, tstat=0.5480), we found significantly more overlap in the trajectories of surface fish. The overlap between convex polygons of pairs surface fish (mean 13.89% overlap, std=9.57, n=1371) was significantly greater (T-test, p<0.0001, df=2219, tstat=13.22) than cavefish (mean 8.41%, std 9.08, n=950). Our impression is that the cavefish are territorial whereas surface fish were not territorial while feeding. The increased overlap in trajectories in surface fish may be a result of their lower amplitude electric fields, which could reduce the distance for detection of conspecifics, or simply due to the larger numbers of surface fish at the study sites. Further, we expect that the surface fish may defend territories during the day when hiding in refugia. The necessary grid recordings were not possible during the day—additional recordings are needed to better understand the diurnal modulation of social behavior in the surface fish.

3.7 Social stimuli and movement interact to modulate EOD signaling

Movement affected social signaling. Three surface fish were placed in tubes that restricted their movement during grid recordings. The variation of EOD frequencies in these fish (mean=0.39 Hz, std=0.36, n=29) was significantly lower than in freely swimming fish (T-test, p=0.0013, df=211, tstat=3.26). This is particularly interesting because the fish in tubes were surrounded by freely moving fish. These fish were exposed to conspecific electric fields that varied in amplitude as the other fish swam around them, producing envelopes (Stamper et al., 2013; Thomas et al., 2018). The dramatic reduction in the variation of their electric fields suggests that fish may modulate their electric signaling to conspecifics in relation to their own movement. Alternatively, other ancillary factors may have diminished overall sensorimotor responsiveness in restrained fish.

We also measured the EODs of three solitary cavefish—individuals that were freely moving but nevertheless isolated from conspecific electric fields. These fish also exhibited reduced variability of their EOD frequencies (mean=0.24 Hz, std=0.09, N=9) compared to cavefish in groups (T-test, p=0.0080, df=292, tstat=0.94). This variability of the electric field frequencies of solitary cavefish was not significantly different from the variability of surface fish in tubes (T-test, p=0.2554, df=36, tstat=1.16). These data are consistent with the theory that variability in EOD frequencies is driven by a combination of a fish's own movement and its electrical interactions with conspecifics.

3.8 Correlations between movement and EOD frequencies suggest envelope responses in the field

We examined the relations between relative movement and EOD frequency as fish interacted with conspecifics. We measured the dFs of all pairs of fish and their simultaneous pairwise distances (Fig. 5). In many pairs of fish, dF and distance appeared to be correlated—either changing in the same or opposite directions (Fig. 5A,B). Negative correlations (e.g. dF increases as distance decreases) may be driven by a JAR-like response known as 'envelope' responses (Stamper et al., 2012, 2013; Metzen and Chacron, 2013; Thomas et al., 2018; Huang and Chacron, 2016). As *Eigenmannia* move closer together, the relative amplitudes of each fish's electric field increases (similar to moving closer to a sound source). Fish respond

to these increases by shifting their electric field frequency either up or down (Stamper et al., 2013; Metzen
and Chacron, 2013; Thomas et al., 2018; Huang and Chacron, 2016).

Correlations between two independently varying measurements may, however, occur spontaneously.
To assess the rate of spontaneous correlations, we shuffled the distance and dF trajectories in time. We
compared distributions of Pearson correlations between dF and distance in the shuffled and original data
(Fig. 5C,D).

In surface fish, we found significantly more negative correlations but not positive correlations (Fisher's
exact test, less than 0.6: p=0.048, greater than 0.6: p=0.719)(Fig. 5C). In cavefish, we found both
significantly more negative and positive correlations (Fisher's exact test, less than 0.6: p<0.0001, greater
than 0.6: p<0.0001)(Fig. 5D). These results suggest that both surface and cavefish exhibit JAR-like
envelope responses—negative correlations between distance and dF. However, cavefish routinely exhibited
positive dF-distance correlations, which may be produced during aggressive behaviors, such as active
jamming (Tallarovic and Zakon, 2002).

Positive correlations were observed in fish with dFs of lower than 10 Hz (Fig. 5B). Such positive
correlations at low dFs are unexpected because of the impact of the JAR. The JAR is strongest at dFs of
approximately 2-8 Hz (Heiligenberg, 1991) and should increase dFs in nearby fish, generally resulting in
negative correlations between distance and dF. These unexpected positive correlations may be a result of
the production of social signals that 'override' the JAR, or may be driven by social interactions with other
nearby fish that have higher dFs.

Interestingly, we found examples of strong Pearson correlations between dF and distance at dFs over
50 Hz distances of over 100 cm and (Fig. 5A). This is interesting because these distances were previously
believed to be beyond the boundary of the animal's ability to detect such signals (Nelson et al., 1997;
Bastian, 1981; Henninger et al., 2018). These data are similar to evidence in other species of weakly
electric fish which show that individuals can detect each other via weak modulations of their electric signals
(Henninger et al., 2018; Raab et al., 2019).

4   DISCUSSION

263 We compared the electric behavior and locomotor movement in a population of epigean fish *Eigenmannia*
264 *trilineata* and a population of troglobitic fish *Eigemannia vicentespelea*. We observed that surface fish had
265 diurnal changes in behavior, hiding in refugia along the sides of streams during the day and swimming and
266 feeding along the bottom of sandy and rocky streams during the night. In contrast, cavefish showed no
267 apparent diurnal modulation of behavior.

4.1   Energetics

The EOD amplitudes of the troglobitic *Eigenmannia* were, on average, higher than the nearby surface
fish. This may be in part due to body size: a previous report (Bichuette and Trajano, 2006) showed that
the cavefish are generally larger than the surface fish. Further, our preliminary anatomical evidence from
diceCT scans suggest that the electric organs of cavefish may be relatively larger than in the surface fish.
Irrespective of size, the energetic cost of generating electric fields is high, consuming up to one quarter of
an individuals energy budget (Salazar et al., 2013; Markham et al., 2016). The fact that cave *Eigenmannia*
produce such energetically costly electric fields suggests that sufficient food resources are available and
accessible. The loss of eyes and pigment in these cavefish, therefore, is likely not under strong selection for
energetic costs, but rather neutral selection (Jeffery, 2009).

Indeed, Gymnotiform fishes throughout the Amazon basin have relatively small eyes. Over the years, we have encountered many individual fish with missing or damaged eyes. Further, we routinely observe dense infestations of nematode parasites in the eyes of individuals from a related genus, *Apteronotus*. These anecdotal observations are consistent with the theory that Gymnotiform fishes rely more heavily on electric sensing than vision for survival and reproduction. Gymnotiform fishes, including the troglobitic weakly electric cavefish *Eigenmannia vicentespelea*, represent a unique opportunity to study evolutionary changes related sensory perception and behavioral control.

4.2 EOD frequencies

The distribution of EOD frequencies in both of these groups of *Eigenmannia* were uni-modal, as has been previously observed (Tan et al., 2005). Further, we were not able to identify any frequency-dependent signalling that might be correlated with sex. Sex differences in EOD frequencies are well known in *Apteronotus* (Fugère et al., 2010; Raab et al., 2019) and *Sternopygus* (Hopkins, 1972), and there may be sex differences in EOD frequencies in *Eigenmannia* (Dunlap and Zakon, 1998). There was a difference in the distributions of EOD frequencies between the cavefish and surface fish, but the meaning of these differences remains unclear. The larger range of frequencies seen in the cavefish may be due to sustained interactions related to territoriality—fish in adjacent territories may increase their dFs over time.

Unexpectedly, we observed sustained, low-frequency dFs in both nearby surface and cavefish. Such interactions are not captured by models of the JAR (Madhav et al., 2013). Low-frequency dFs potentially impair electrolocation in *Eigenmannia* (Ramcharitar et al., 2005), and can be avoided by performing a JAR (Heiligenberg, 1991) or moving further apart (Tan et al., 2005). It seems unlikely that these fish were experiencing significant impairment of sensory function, as they were engaged in feeding on prey under the substrate in complete darkness. Studies in *Apteronotus albifrons* demonstrates that fish may use ampullary receptors to detect exogenously generated electric fields for prey capture (Nelson and MacIver, 1999). In addition, elasmobranchs have also been shown to detect and discriminate signals from substrate-bound prey using passive electroreception mediated by ampullary receptors (Kalmijn, 1971, 1982).

4.3 Territoriality

Territoriality is a form of space-related dominance (Kaufmann, 1983). The most prominent function of having a territory is to provide the holder with an secured supply of resources. In the epigean streams outside of the Terra Ronca cave, food resources for *Eigenmannia* appear to be widely distributed in sandy substrates. Our guess is that the size and distribution of prey items precludes territorial defense of food resources. On the other hand, we expect to find evidence of territoriality during the day, as refugia likely vary in quality and are relatively small.

Why cavefish exhibit evidence of territoriality is unclear. There are no known predators of *Eigenmannia* in the cave, eliminating the value of protective refugia. Territoriality may occur as a result of uneven distribution of food resources in the cave, due to other physical features that impact the fish, or a consequence of plesiomorphic social and/or reproductive behaviors. Territoriality has been described in other genera, such as *Gymnotus* (Zubizarreta et al., 2020).

4.4 Spooky interactions at a distance

The strength of electric fields in water decay at a rate of approximately distance cubed (Henninger et al., 2018). As a result, the distances between fish determine the strength of the interaction of their electric fields: nearby fish will experience higher EOD voltages than from those of distant fish. Because the EODs of *Eigenmannia* are nearly sinusoidal, distance will have an effect on the amplitude of modulations caused by the summation of EODs: nearby fish will have large amplitude modulations near 100%, whereas distant

fish will have far lower depths of modulation, often below 10%. The relative movement between fish will
cause concomitant changes in the strengths of EODs and depths of modulation that are proportional to
distance. The changes are known as 'envelopes'—the modulation of amplitude modulations. Envelope
stimuli can elicit changes in electric field frequencies in *Eigenmannia* and other Gymnotiform species
(Stamper et al., 2013; Thomas et al., 2018).

We found strong correlations between distance and pairwise differences in EOD frequencies at large
distances of over 1.5 meters and dFs of over 50 Hz in both surface and cavefish. These results are similar to
reports from field studies that examined other Gymnotiform species (Henninger et al., 2018; Raab et al.,
2019). These field studies suggest that electric fish are far more sensitive to electrosocial stimuli than
previously appreciated, requiring a reexamination of the neural systems for their perception.

Although many of the correlations between distance and dF were spontaneous, both surface and cavefish
exhibited significantly more negative correlations between distance and dF than random. These negative
correlations are likely driven by the amplitude envelope of electrical interference patterns, producing JAR-
like behavioral responses (Stamper et al., 2012). These findings show that laboratory studies of envelope
responses are ecologically relevant (Stamper et al., 2012, 2013; Metzen and Chacron, 2013; Thomas et al.,
2018; Huang and Chacron, 2016). Cavefish also exhibited significantly more positive correlations between
distance and dF. We believe that these may be aggressive signals in which fish actively jam each other
(Tallarovic and Zakon, 2002).

4.4.1   Permission to Reuse and Copyright

340 Figures, tables, and images will be published under a Creative Commons CC-BY licence and
341 permission must be obtained for use of copyrighted material from other sources (including re-
342 published/adapted/modified/partial figures and images from the internet). It is the responsibility of the
343 authors to acquire the licenses, to follow any citation instructions requested by third-party rights holders,
344 and cover any supplementary charges.

CONFLICT OF INTEREST STATEMENT

345 The authors declare that the research was conducted in the absence of any commercial or financial
346 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

347 Bichuette, Soares, and Fortune conceived and executed the field research. Andanar, Madhav, Jayakumar, and
348 Fortune analyzed the electrical data. Soares collected and analyzed the diceCT data. Madhav, Jayakumar,
349 and Cowan contributed to the design, use, and analysis of data from the grid system. All authors contributed
350 to the preparation of the manuscript.

FUNDING

351 This work was supported by a collaborative National Science Foundation (NSF) Award to Noah J Cowan
352 (1557895) and Eric S Fortune (1557858). The field work was supported by startup funds provided by the
353 New Jersey Institute of Technology to Daphne Soares. Maria Elina Bichuette received funding from the
354 CNPq (Awards 308557/2014-0).

ACKNOWLEDGMENTS

355 Field research permits were granted by the ICMBio and SEMARH/SECIMA. We would like to
356 thank DF Torres, MJ Rosendo and CCP de Paula for help in the field. Thanks to Dr. Jessica

357 Ware for the use of the CT scanner. This manuscript has been released as a pre-print at
358 [https://www.biorxiv.org/content/10.1101/747154v1], (Fortune et al., 2019).

SUPPLEMENTAL DATA

359 Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures,
360 please include the caption in the same file as the figure. LaTeX Supplementary Material templates can be
361 found in the Frontiers LaTeX folder.

DATA AVAILABILITY STATEMENT

362 The data and the Matlab scripts used to generate the figures will be made available on publication of this
363 study on github (URL).

REFERENCES

Assad, C., Rasnow, B., Stoddard, P., and Bower, J. (1998). The electric organ discharges of the gymnotiform
fishes: II. *Eigenmannia*. *Journal of Comparative Physiology A: Sensory, Neural, and Behavioral*
Physiology 183, 419–432. doi:10.1007/s003590050268
Bastian, J. (1981). Electrolocation I. how electroreceptors of *Apteronotus albifrons* code for moving objects
and other electrical stimuli. *Journal of Comparative Physiology A: Sensory, Neural, and Behavioral*
Physiology 144, 465–479. doi:10.1007/bf01326832
Bichuette, M. and Trajano, E. (2003). Epigean and subterranean ichthyofauna from the São Domingos
karst area, Upper Tocantins River basin, Central Brazil. *Journal of Fish Biology* 63, 1100–1121
Bichuette, M. and Trajano, E. (2006). Morphology and distribution of the cave knifefish *Eigenmannia*
vicentespelaea triques, 1996 (gymnotiformes: Sternopygidae) from Central Brazil, with an expanded
diagnosis and comments on subterranean evolution. *Neotropical Ichthyology* 4, 99–105. doi:10.1590/
s1679-62252006000100011
Bichuette, M. and Trajano, E. (2015). Population density and habitat of an endangered cave fish
Eigenmannia vicentespelaea Triques, 1996 (Ostariophysi: Gymnotiformes) from a karst area in central
Brazil. *Neotropical Ichthyology* 13, 113–122
Bichuette, M. and Trajano, E. (2017). Biology and behavior of *Eigenmannia vicentespelaea*, a troglobitic
electric fish from Brazil (Teleostei: Gymnotiformes: Sternopygidae): a comparison to the epigean
species, *E. trilineata*, and the consequences of cave life. *Tropical Zoology* 30, 68–82. doi:10.1080/
03946975.2017.1301141
Crampton, W. (2019). Electroreception, electrogenesis and electric signal evolution. *Journal of Fish*
Biology 95, 92–134. doi:10.1111/jfb.13922
Culver, D. and Pipan, T. (2019). *The Biology of Caves and Other Subterranean Habitats* (Oxford University
Press)
Davis, E. and Hopkins, C. (1988). Behavioural analysis of electric signal localization in the electric fish,
Gymnotus carapo (Gymnotiformes). *Animal Behaviour* 36, 1658–1671. doi:10.1016/s0003-3472(88)
80106-4
Dunlap, K. and Zakon, H. (1998). Behavioral actions of androgens and androgen receptor expression in
the electrocommunication system of an electric fish, *Eigenmannia virescens*. *Hormones and Behavior*
34, 30–38. doi:10.1006/hbeh.1998.1460
Fortune, E., Andanar, N., Madha, M., Jayakumar, R., Cowan, N., Bichuette, M., et al. (2019). Spooky
interaction at a distance in cave and surface dwelling electric fishes. *bioRxiv* doi:10.1101/747154

Fugère, V., Ortega, H., and Krahe, R. (2010). Electrical signalling of dominance in a wild population of
electric fish. _Biology Letters_ 7, 197–200. doi:10.1098/rsbl.2010.0804

Heiligenberg, W. (1991). _Neural Nets in Electric Fish (Computational Neuroscience)_ (Bradford Books)

Henninger, J., Krahe, R., Kirschbaum, F., Grewe, J., and Benda, J. (2018). Statistics of natural
communication signals observed in the wild identify important yet neglected stimulus regimes in
weakly electric fish. _The Journal of Neuroscience_ 38, 5456–5465. doi:10.1523/jneurosci.0350-18.2018

Henninger, J., Krahe, R., Sinz, F., and Benda, J. (2020). Tracking activity patterns of a multispecies
community of gymnotiform weakly electric fish in their neotropical habitat without tagging. _The Journal_
of Experimental Biology 223, jeb206342. doi:10.1242/jeb.206342

Hopkins, C. (1972). Sex differences in electric signaling in an electric fish. _Science_ 176, 1035–1037.
doi:10.1126/science.176.4038.1035

Huang, C. and Chacron, M. (2016). Optimized parallel coding of second-order stimulus features by
heterogeneous neural populations. _Journal of Neuroscience_ 36, 9859–9872. doi:10.1523/jneurosci.
1433-16.2016

Jeffery, W. (2009). Regressive evolution in _Astyanax_ cavefish. _Annual Review of Genetics_ 43, 25–47.
doi:10.1146/annurev-genet-102108-134216

Kalmijn, A. (1971). The electric sense of sharks and rays. _The Journal of Experimental Biology_ 55,
371–383

Kalmijn, A. (1982). Electric and magnetic field detection in elasmobranch fishes. _Science_ 218, 916–918.
doi:10.1126/science.7134985

Kaufmann, J. (1983). On the definitions and functions of dominance and territoriality. _Biological reviews_
58, 1–20

Madhav, M., Jayakumar, R., Demir, A., Stamper, S., Fortune, E., and Cowan, N. (2018). High-resolution
behavioral mapping of electric fishes in Amazonian habitats. _Scientific Reports_ 8. doi:10.1038/
s41598-018-24035-5

Madhav, M., Stamper, S., Fortune, E., and Cowan, N. (2013). Closed-loop stabilization of the jamming
avoidance response reveals its locally unstable and globally nonlinear dynamics. _Journal of Experimental_
Biology 216, 4272–4284. doi:10.1242/jeb.088922

Markham, M., Ban, Y., McCauley, A., and Maltby, R. (2016). Energetics of sensing and communication
in electric fish: A blessing and a curse in the anthropocene? _Integrative and Comparative Biology_ 56,
889–900. doi:10.1093/icb/icw104

Markham, M., McAnelly, L., Stoddard, P., and Zakon, H. (2009). Circadian and social cues regulate ion
channel trafficking. _PLoS Biology_ 7, e1000203. doi:10.1371/journal.pbio.1000203

Metzen, M. and Chacron, M. (2013). Weakly electric fish display behavioral responses to envelopes
naturally occurring during movement: implications for neural processing. _The Journal of Experimental_
Biology 217, 1381–1391. doi:10.1242/jeb.098574

Migliaro, A., Moreno, V., Marchal, P., and Silva, A. (2018). Daily changes in the electric behavior of
weakly electric fish naturally persist in constant darkness and are socially synchronized. _Biology Open_ 7,
bio036319. doi:10.1242/bio.036319

Nelson, M. and MacIver, M. (1999). Prey capture in the weakly electric fish _Apteronotus albifrons_:
sensory acquisition strategies and electrosensory consequences. _Journal of Experimental Biology_ 202,
1195–1203

Nelson, M., Xu, Z., and Payne, J. (1997). Characterization and modeling of P-type electrosensory afferent
responses to amplitude modulations in a wave-type electric fish. _Journal of Comparative Physiology A:_
Sensory, Neural, and Behavioral Physiology 181, 532–544. doi:10.1007/s003590050137

Pedraja, F., Hofmann, V., Lucas, K., Young, C., Engelmann, J., and Lewis, J. (2018). Motion parallax in electric sensing. *Proceedings of the National Academy of Sciences* 115, 573–577. doi:10.1073/pnas.1712380115

Raab, T., Linhart, L., Wurm, A., and Benda, J. (2019). Dominance in habitat preference and diurnal explorative behavior of the weakly electric fish *Apteronotus leptorhynchus*. *Frontiers in Integrative Neuroscience* 13. doi:10.3389/fnint.2019.00021

Ramcharitar, J., Tan, E., and Fortune, E. (2005). Effects of global electrosensory signals on motion processing in the midbrain of *Eigenmannia*. *Journal of Comparative Physiology A: Sensory, Neural, and Behavioral Physiology* 191, 865–872. doi:10.1007/s00359-005-0008-2

Salazar, V., Krahe, R., and Lewis, J. (2013). The energetics of electric organ discharge generation in gymnotiform weakly electric fish. *The Journal of Experimental Biology* 216, 2459–2468. doi:10.1242/jeb.082735

Sinnett, P. and Markham, M. (2015). Food deprivation reduces and leptin increases the amplitude of an active sensory and communication signal in a weakly electric fish. *Hormones and Behavior* 71, 31–40. doi:10.1016/j.yhbeh.2015.03.010

Stamper, S., Carrera-G, E., Tan, E., Fugere, V., Krahe, K., and Fortune, E. (2010). Species differences in group size and electrosensory interference in weakly electric fishes: Implications for electrosensory processing. *Behavioural Brain Research* 207, 368–376. doi:10.1016/j.bbr.2009.10.023

Stamper, S., Fortune, E., and Chacron, M. (2013). Perception and coding of envelopes in weakly electric fishes. *The Journal of Experimental Biology* 216, 2393–2402. doi:10.1242/jeb.082321

Stamper, S., Madhav, M., Cowan, N., and Fortune, E. (2012). Beyond the jamming avoidance response: weakly electric fish respond to the envelope of social electrosensory signals. *The Journal of Experimental Biology* 215, 4196–4207. doi:10.1242/jeb.076513

Stoddard, P., Zakon, H., Markham, M., and McAnelly, L. (2006). Regulation and modulation of electric waveforms in gymnotiform electric fish. *Journal of Comparative Physiology A: Sensory, Neural, and Behavioral Physiology* 192, 613–624. doi:10.1007/s00359-006-0101-1

Tallarovic, S. and Zakon, H. (2002). Electrocommunication signals in female brown ghost electric knifefish, *Apteronotus leptorhynchus*. *Journal of Comparative Physiology A: Sensory, Neural, and Behavioral Physiology* 188, 649–657. doi:10.1007/s00359-002-0344-4

Tan, E., Nizar, J., Carrera-G, E., and Fortune, E. (2005). Electrosensory interference in naturally occurring aggregates of a species of weakly electric fish, *Eigenmannia virescens*. *Behavioural Brain Research* 164, 83–92. doi:10.1016/j.bbr.2005.06.014

Thomas, R., Metzen, M., and Chacron, M. (2018). Weakly electric fish distinguish between envelope stimuli arising from different behavioral contexts. *The Journal of Experimental Biology* 221, jeb178244. doi:10.1242/jeb.178244

Triques, M. (1996). *Eigenmannia vicentespelaea*, a new species of cave dwelling electrogenic neotropical fish (Ostariophysi: Gymnotiformes: Sternopygidae). *Revue Francaise de Aquariologi* 23

Watanabe, A. and Takeda, K. (1963). The change of discharge frequency by A.C. stimulus in a weak electric fish. *The Journal of Experimental Biology* 40, 57–66

Yu, N., Hupe, G., Longtin, A., and Lewis, J. (2019). Electrosensory contrast signals for interacting weakly electric fish. *Frontiers in Integrative Neuroscience* 13. doi:10.3389/fnint.2019.00036

Zubizarreta, L., Quintana, L., Hernández, D., Teixeira de Mello, F., M, M., Honji, R., et al. (2020). Seasonal and social factors associated with spacing in a wild territorial electric fish. *bioRxiv* doi:10.1101/2020.01.29.924761

FIGURE CAPTIONS

Figure 1. Surface and cave *Eigenmannia*. A) Study sites are in a clear water system in the Rio da Lapa karst region in Goiás, Brazil. B) Top, the surface *Eigenmannia* site was in the entrance of the Rio da Lapa cave. B) bottom, the cave *Eigenmannia* were found in the São Vicente II cave. C) Surface *Eigenmannia* have well-developed eyes and distinctive markings. D) Cave *Eigenmannia* have poorly developed or missing eyes and lack pigmented features. diceCT imaging reveals the differences in eye sizes and a potential difference in the relative size of electric organs. E) Coronal sections through the head and mid-body of a surface fish and F) two cavefish (dorsal is up). Large, bright cells in the caudoventral coronal sections appear relatively larger in the cave versus surface species.

Figure 2. Electrical and locomotor behavior of *Eigenmannia*. A-B) Positions of surface and cave *Eigenmannia* over a period of 120 seconds. C-D) EOD frequencies of these fish. Each color represents a unique fish in each recording.

Figure 3. Differences in the distribution of EOD frequencies and amplitudes in surface fish (blue, up) and cavefish (red, down). A) Percent observations of EOD frequencies for surface fish and cavefish. B) Distribution of EOD amplitudes.

Figure 4. Differences in EOD frequencies. A) Percent observations of dFs for surface fish (blue, up) and cavefish (red, down). Grey bar highlights the low frequency region shown in B. C) Example epoch of two fish with sustained low-frequency dF. Top, EOD frequencies each fish; bottom Euclidian distance over time. D-E) X-Y positions of each fish.

Figure 5. Correlations between EOD frequency and movement. A) Example of a strong negative correlation between distance (purple, left) and dF (orange, right) of a pair of *Eigenmannia* over a period of 5 minutes. Dots are measurements, lines are low-pass fits of the data. B) Example of a strong positive correlation. C) Distribution of Pearson correlations between distance and dF for surface fish (blue) and shuffled data (black). D) Distribution of Pearson correlations between distance and dF for cavefish (red) and shuffled data (black). There were significantly more strong negative Pearson correlations, below -0.6 (marked in grey) in both surface and cavefish. Unlike in surface fish, there were significantly more examples of high Pearson correlations (above 0.6) in cavefish than in shuffled data.