Single cell transcriptomics reveal temporal dynamics of critical regulators of germ

cell fate during mouse sex determination

Authors: Chloé Mayère[1,2], Yasmine Neirijnck[1,3], Pauline Sararols[1], Chris M Rands[1],

Isabelle Stévant[1,2], Françoise Kühne[1], Anne-Amandine Chassot[3], Marie-Christine

Chaboissier[3], Emmanouil T. Dermitzakis[1,2], Serge Nef[1,2,]*.

Affiliations:

[1]Department of Genetic Medicine and Development, University of Geneva, 1211 Geneva,

Switzerland;

[2]iGE3, Institute of Genetics and Genomics of Geneva, University of Geneva, 1211

Geneva, Switzerland;

[3]Université Côte d'Azur, CNRS, Inserm, iBV, France;

Lead Contact:

*Corresponding Author:** Serge Nef, 1 rue Michel-Servet CH-1211 Genève 4,

Serge.Nef@unige.ch. + 41 (0)22 379 51 93

Running Title: Single cell transcriptomics of germ cells

Abbreviations;

AGC: Adrenal Germ Cell

GC: Germ cell

OGC: Ovarian Germ Cell

TGC: Testicular Germ Cell

scRNA-seq: Single-cell RNA-Sequencing

DEG: Differentially Expressed Gene

Keywords:

Single-cell RNA-Sequencing (scRNA-seq), sex determination, ovary, testis, gonocytes,

oocytes, prospermatogonia, meiosis, gene regulatory network, germ cells, development,

RNA splicing

Summary

Despite the importance of germ cell (GC) differentiation for sexual reproduction, the gene networks underlying their fate remain unclear. Here, we comprehensively characterize the gene expression dynamics during sex determination based on single-cell RNA sequencing of 14,914 XX and XY mouse GCs between embryonic days (E) 9.0 and 16.5. We found that XX and XY GCs diverge transcriptionally as early as E11.5 with upregulation of genes downstream of the Bone morphogenic protein (BMP) and Nodal/Activin pathways in XY and XX GCs, respectively. We also identified a sex-specific upregulation of genes associated with negative regulation of mRNA processing and an increase in intron retention consistent with a reduction in mRNA splicing in XY testicular GCs by E13.5. Using computational gene regulation network inference analysis, we identified sex-specific, sequential waves of putative key regulator genes during GC differentiation and revealed that the meiotic genes are regulated by positive and negative master modules acting in an antagonistic fashion. Finally, we found that rare adrenal GCs enter meiosis similarly to ovarian GCs but display altered expression of master genes controlling the female and male genetic programs, indicating that the somatic environment is important for GC function. Our data is available on a web platform and provides a molecular roadmap of GC sex determination at single-cell resolution, which will serve as a valuable resource for future studies of gonad development, function and disease.

Introduction

Germ cells are the precursors of the gametes, either sperm or eggs, required for sexual reproduction. GCs are initially bipotential, and their sex-specific fate depends on somatic cues provided by the ovarian and testicular environments, rather than the chromosomal sex they carry (1-3). In mice, primordial germ cells (PGCs) arise in the posterior proximal epiblast around embryonic day (E) 6.25. PGCs rapidly proliferate and colonize the gonads at around E10.5 (4, 5).

In fetal ovaries, GCs enter meiosis asynchronously in a wave from anterior to posterior over about two days, between E12.5 and E14.5 (6-8). This entry into meiosis is considered a hallmark of commitment to oogenesis (9). It is triggered by the expression of the pre-meiotic gene *Stra8* and the meiosis-associated gene *Rec8* and, at the same time, by the downregulation of pluripotency markers such as *Oct4* (*Pou5f1*), *Sox2* and *Nanog* (6, 7, 10). The up-regulation of the *Stra8* and *Meiosin* genes (10, 11) depends on the expression of the transcriptional regulator ZGLP1 that triggers the oogenic fate and stimulates the meiotic program (12). All-trans retinoic acid (ATRA) signaling has been long considered as the inducer of *Stra8* expression and germ cell decision to enter meiosis (13, 14). However, ATRA signaling has recently been excluded as the meiosis-instructing substance in oogonia (15, 16).

In contrast, GCs in fetal testes differentiate into prospermatogonia through a number of distinct, potentially interrelated events that occur asynchronously over a period of several days, but does not involve entry into meiosis (for a review see (17) and (18)). GCs

adopting the male fate up-regulate cell-cycle inhibitors such as *Cdkn2b* (19) and are

mitotically arrested from E12.5 onward (20). They transiently activate the

NODAL/CRIPTO signaling pathway (21-23) and down-regulate pluripotency genes such

as *Nanog*, *Sox2* and *Pou5f1* (24). From E13.5 onward, they begin to express male-

specific genes including *Nanos2* (25), *Dnmt3l* (26) and *Piwil4* (27), which ensure normal

male gametogenesis by regulating spermatogonial stem cell properties.

Although the cellular origins of oogonia and spermatogonia are well documented (28),

numerous questions related to the molecular mechanisms underlying their sex fate and

differentiation remain. For instance, the transcriptional programs mediating GC sex

determination are incompletely understood, and the essential genes and transcriptional

regulators orchestrating such specifications remain poorly defined.

To date, most transcriptional analyses relevant for mouse or human GC sex determination

have been conducted using traditional methods such as microarrays or bulk RNA-seq on

either whole gonads or isolated GC populations at few selected time points (29-41).

These studies, although informative, provided only an average transcriptional summary,

masking the inherent variability of individual cells and lineage types and thereby limiting

their capacity to reveal the precise dynamics of gene expression during GC sex

determination.

To determine the sequence of transcriptional events that are associated with GC

commitment and differentiation toward oogenesis and spermatogenesis, we performed

time-series single-cell RNA sequencing (scRNA-seq) on developing mouse gonads. We

recovered 14,914 GCs from XX and XY gonads across six developmental time points

from E9.0 to E16.5, encompassing the entire developmental process of gonadal sex

determination and differentiation. We reconstructed the developmental trajectories of

male and female GCs, characterized the associated genetic programs, and predicted gene

regulatory networks that regulate GC commitment and differentiation.

Material & Methods

Transgenic Mice

All work on animals has been carried out in accordance with the ethical guidelines of the

Direction Générale de la Santé of the Canton de Genève (experimentation ID GE/57/18).

Tg(Nr5a1-GFP) mouse strain was described previously (42) and has been maintained on

a CD1 genetic background.

Mouse urogenital ridges, testes, ovaries and adrenal glands collection

CD-1 female mice were bred with heterozygous *Tg(Nr5a1-GFP)* transgenic male mice.

Adult females were time-mated and checked for the presence of vaginal plugs the next

morning (E0.5). E9.0 (18±2 somites (ts)), E10.5 (8±2 tail somites), E11.5 (19±4 ts),

E12.5, E13.5, E16.5 and E18.5 embryos were collected and the presence of the *Nr5a1-*

GFP transgene was assessed under UV light. Sexing of E9.0, E10.5 and E11.5 embryos

was performed by PCR with a modified protocol from (43). Posterior trunks (E9.0) and

urogenital ridges (E10.5, E11.5) from each sex, XY adrenal glands, testes or ovaries were

pooled for tissue dissociation. Posterior trunks, urogenital ridges or adrenal glands were

enzymatically dissociated at 37ºC using the Papain dissociation system (Worthington

#LK003150). Cells were resuspended in DMEM 2%FBS, filtered through a 70 μm cell

strainer and stained with the dead cell marker Draq7™ (Beckman Coulter, #B25595).

Viable single cells were collected on a BD FACS Aria II by excluding debris (side scatter

vs. forward scatter), dead cells (side scatter vs. Draq7 staining), and doublets (height vs.

width). Testes and ovaries (from E12.5 to E16.5) were enzymatically dissociated at 37ºC

during 15 minutes in Trypsin-EDTA 0.05% (Gibco #25300054), resuspended in DMEM

2%FBS and filtered through a 70 μm cell strainer. After counting, 3000 to 7000 single

cells were loaded on a 10x Chromium instrument (10x Genomics). Single-cell RNA-Seq

libraries were prepared using the Chromium Single Cell 3′ v2 Reagent Kit (10x

Genomics) according to manufacturer's protocol. Each condition (organ, sex and

developmental stage) was performed in two biological independent replicates.

Sequencing

Library quantification was performed using the Qubit fluorometric assay with dsDNA HS

Assay Kit (Invitrogen). Library quality assessment was performed using a Bioanalyzer

Agilent 2100 with a High Sensitivity DNA chip (Agilent Genomics). Libraries were

diluted, pooled and sequenced on an Illumina HiSeq4000 using paired-end 26□+□98 +

8□bp as the sequencing mode. Libraries were sequenced at a targeted depth of 100 000 to

150 000 total reads per cell. Sequencing was performed at the Health 2030 Genome

Center, Geneva.

RNA scope and DAZL immunohistochemistry imaging

For RNAscope experiments, E14.5 XX and XY embryos were dissected in Phosphate

Buffered Saline (PBS). Samples were fixed in 4% paraformaldehyde (PFA) overnight at

room temperature. Five µm sections were processed for RNA *in situ* detection using the

RNAscope 2.0 High Definition-RED Kit according to the manufacturer's instructions

(ACDBio). *Rbm38, Supt6 and Tbrg4* probes were designed by ACDBio. Slides were

counterstained with DAPI diluted in the mounting medium at 10 µg/ml (Vectashield,

Vector laboratories) to detect nuclei, and with DAZL antibody (cat # GTX89448,

Genetex; 1:200) to detect GCs. RNAscope results were examined under a confocal

microscope Leica DM5500 TCS SPE on an upright stand (Leica Microsystems,

Mannheim, Germany), using an ACS APO 63X oil 1.3 NA objective. The lasers used

were diodes laser (405 nm, 488 nm and 532 nm). The microscope was equipped with a

galvanometric stage in order to do z-acquisitions.

Bioinformatic Analysis

Data processing with the Cell Ranger package, cell selection and quality controls

Computations were performed at the Vital-IT Center for high-performance computing of

the SIB (Swiss Institute of Bioinformatics) (http://www.vital-it.ch). Demultiplexing,

alignment, barcode filtering and UMI counting were performed with the Cell Ranger v2.1

pipeline (10x Genomics). Algorithms and versions used are listed in the key resources

table. Data were mapped to the mouse reference genome GRCm38.p5 in which the eGFP

(NC_011521.1) combined with the bovine GH 3′-splice/polyadenylation signals

(Stallings et al., 2002) (NM_180996.1) sequences have been added.

For cell-associated barcode selection, we computed the knee point and the inflection

point of the ranked barcodes distribution plot. The local minimum between these points

on the density curve (density base R version 3.6.1 function, bw=500, n=4096) of the UMI

number per barcode was detected using the quantmod package. This local minimum was

used as a threshold for cell-associated barcode selections. When no local minimum could

be detected between the two points, the nearest local minimum was used. Quality controls

regarding abnormal mitochondrial or ribosomal content, UMI number, detected gene

numbers, unmapped reads and putative doublet identification (Scrublet `0.2.1-0`) were

performed, but no data filtering was applied as no important anomalies were detected. In

total, for gonadal samples, we obtained 92,267 cells. It included 13,060 cells from E9.0,

14,962 cells from E10.5, 16,670 cells from E11.5, 20,285 cells from E12.5, 25,794 cells

from E13.5 and 16,994 cells from E16.5. In total, we obtained 61,755 XY cells and

46,010 XX cells. Adrenal glands were collected only in XY embyos and 26,639 cells

were collected in total with 7,296, 11,255, 8,822 and 7,609 cells collected at E12.5,

E13.5, E16.5 and E18.5 respectively.

Gene expression normalization

UMI counts per gene per cell were divided by the total UMI detected in the cell,

multiplied by a scale factor of 10,000 and log transformed.

Batch correction, clustering, UMAP visualization and germ cells selection

To classify the cell populations present in the developing testis and ovary, we selected all

genes detected in more than 50 cells (21,103 genes), log normalized their expression and

ran Independent Component Analysis (ICA, Seurat package v2.3.0) on these (100

components). ICA extracts non-Gaussian components from data, allowing a better

discrimination of small cell populations in heterogeneous dataset (44). We set the number

of ICs to 100 as we expected to have less than 100 different populations in our total

dataset. Another advantage of ICA is its good performance with nearly no gene filtering,

such as highly variable genes selection.

A neighbor graph corrected for batch between replicates was computed using BBKNN

(1.3.5) (45) with default values and n_neighbors_within_batch = 3 and the ICA as input

(all components).

Clustering was performed using the Scanpy Leiden method with resolution 1 and UMAP

were generated using the Scanpy UMAP method with default parameters. We selected

clusters with a strong expression of 10 well-known GCs markers (**Figure S1**).

 Pseudotime ordering of the cells

To order the cells along a pseudotime, we took advantage of the discrete prior knowledge

we have about the embryonic day at which each cell was harvested and generated an

ordinal regression model (adapted from (46)) to obtain a continuous pseudotime score

reflecting the differentiation status of the cells (Bmrm package v4.1). We trained the

model using a tenfold cross-validation schema on the whole expression matrix with a

correction for unequal numbers of XX and XY cells. The trained model outputs a weight

for each gene according to its ability to predict the embryonic time of each cell in an

ordered manner. We then predicted a continuous score for all cells using the weights of

the 100 top weighted genes to avoid over fitting of the model.

To determine which genes were important for sex specific progression of the cells along

time, we trained two distinct models on XX and XY cells and selected the genes with a

significantly higher absolute weight for classification (compared to the complete set of

genes). We merged the two lists of genes and obtained 688 unique genes, of which 128

were common to both XX and XY lineages, 266 were specific to XX germ cells, and 294

were specific to XY germ cells (**Fig. S2A** and **B**)

Gene regulatory network generation

GRN was generated using pyScenic package (v0.9.14) (47). First adjacencies were

generated using the grnboost function (48) with the whole expression matrix of all the

germ cells as input and the transcription factor list provided with the package. Next,

modules were generated, associating groups of genes to a putative regulator using

modules_from_adjacencies function. Finally, the obtained modules were pruned on the

basis of the presence or absence of the master regulator binding site footprint in the

neighborhood of the TSS of each regulated gene. AUCell was then ran on the obtained

regulons to infer its activity within each cell.

Regulons and genes hierarchical clustering

Regulons or genes were ordered using ward.d hierarchical clustering on the AUC or

expression matrix with Spearman correlation distance. Modules were determined using

cutree with k=10 for negative regulons and k=30 for positive regulons (packages

dendextend v1.13.4, factoextra v1.0.7, tidyr v1.1.0).

Heatmaps and expression curves

Heatmaps were generated using R (packages pheatmap v1.0.12 and heatmaps v1.8.0),

smoothing was performed using smoothheatmap function on log-normalized expression

levels for genes or AUC enrichment values for regulons. Expression curves were

generated using ggplot2 (v3.2.0) geom_smooth function with method "gam" on log

normalized expression levels.

Spliced and unspliced counts analysis

To generate spliced and unspliced counts data, the velocyto.py script from velocyto

(v0.17.8) package was called on each bam file with aforementioned reference genome

annotation. Spliced and unspliced counts were normalized using scvelo package (v0.1.19)

by dividing all UMI counts of a cell by the total counts in the cell, multiplying the result

by the median count number in all cells and applying log1p function. Scatter plots of each

specific gene with spliced and unspliced counts were generated using Ms and Mu layers

(kNN pooled expression) respectively.

Ectopic adrenal germ cells analysis

The analysis was performed using aforementioned steps: log normalization, ICA,

neighbor graph, and clustering with the same parameters. GC clusters were selected with

the same 10 GC marker genes (see **Supplementary information** for details). For

pseudotime ordering of the cells, a model was trained on gonadal cells only and the

pseudotime for adrenal GCs was predicted from it.

Data and source code availability

GCs single-cell RNA-seq data is available on GEO (accession number GSE136220).

Both adrenal and gonadal gene expression data are included in ReproGenomics Viewer

(49, 50).

 Web interface to browse single cell data

 To allow biologists and other researchers to explore the GC data we created an interactive

 website: http://neflabdata.info/. This allows anyone to easily display the expression of

 any mouse gene in our matrix on a UMAP representation of the GC data. Other metadata

 (such as sex, time point, or cluster) can also be shown on the UMAP.

 Our website uses the Cellxgene data explorer application, which has further

 functionalities, such as allowing the comparison of the expression profiles of multiple

 genes, as described in their documentation:

 https://chanzuckerberg.github.io/cellxgene/posts/gallery. We packaged Cellxgene in a

 Docker hosted on Google Cloud Platform (GCP) with our GC data H5AD output file

 from Scanpy. On GCP, we run a virtual machine instance of 2-standard machine type

 with 2 CPUs and 4-8 GB of memory.

 Results

 Generation of a single-cell transcriptional atlas of germ cell sex determination and

 differentiation

 To generate a gene expression atlas of GCs over sex determination, we used droplet-

 based 3' end scRNA-seq (10x Genomics Chromium) of XX and XY gonads from mouse

 embryos at six developmental stages (E9.0, E10.5, E11.5, E12.5, E13.5, and E16.5) (**Fig.**

 1A and **B**). The selected time points cover the entire process of gonadal sex determination

and span the emergence and differentiation of the major testicular and ovarian lineages

including the gonocytes. For each of the 12 conditions, we generated two independent

replicates from different pregnancies and sequenced an average of 10,000 cells. The

transcriptomes of the individual cells were sequenced at the depth of ~100,000-150,000

reads/cell. Using ten well-established GC markers, namely *Ddx4*, *Dazl*, *Mael*, *Dppa3*,

Sycp3, *Pecam*, *Pou5f1*, *Stra8*, *Dmrt1,* and *Dnmt1*, we identified 14,914 GCs among a

total of 107,765 cells (**Fig. S1A-C**) (see also **Supplementary information**). Among

GCs, 8,329 were XX (55.9%) and 6,585 were XY (44.1%). This included 56 cells from

E9.0, 71 cells from E10.5, 946 cells from E11.5, 4,395 cells from E12.5, 6,633 cells from

E13.5 and 2,813 cells from E16.5. The median number of detected expressed genes was

higher in GCs than in somatic cells with 5,536 and 4,502 respectively (**Fig. S1D**),

consistent with data indicating that GCs undergo pervasive and diverse transcription (51).

The generation of this single-cell transcriptional atlas allowed us to further analyze the

dynamic, the sex-specific features and the response to the somatic environment during

germ cell development.

Pseudotime ordering of the cells and differential expression analyses reveals the

dynamics of gene expression across XX and XY germ cell sex determination

UMAP projection of the 14,914 GCs permits visualization of the transcriptional

landscape (**Fig. 1C-E** and **Material & Methods**). To allow researchers to explore GC

expression data, we have created an interactive and user-friendly web site:

http://neflabdata.info/ to easily display the expression of any gene of our dataset on a

UMAP representation. Our analysis revealed that, at early stages (E9.0, E10.5 and

E11.5), the transcriptomes of XX and XY cells globally overlapped, while more

differentiated cells from later stages formed two sex-specific branches (**Fig. 1C** and **D**).

We then analyzed how the transcriptomes of XX and XY cells progress during the

process of sex determination by ordering the cells along a pseudotime that captures the

transcriptional state of each cell. To construct a pseudotime, we used ordinal regression

modeling (46, 52) with prior knowledge about the developmental stage of each cell

capture (see **Fig. 1E**, **Fig. S1E**, **Material & Methods** and **Supplementary information**).

We then represented the smoothed expression of 688 significant genes for two

independent sex specific ordinal regression models (**Table S1**) with a double heatmap

(mirror heatmap) in which the center represents the earliest cells (pseudotime 0, E9.5

cells) and the extremities represent the lineage endpoints (pseudotime 100, E16.5) of XX

GCs (left) and XY GCs (right). The mirror heatmap revealed that XX and XY GCs

diverged as early as E11.5, exhibiting dynamic and sex-specific differentiation programs

mediated by hundreds of genes (**Fig. S2**). In addition, using 63 well-known genes

involved in mouse GC pluripotency, sexual development and differentiation (18, 53), we

confirmed that our single-cell data accurately recapitulated known male and female GC

specification patterns (**Fig. 1F**).

To assess when sex-specific genetic programs are initiated in XX and XY GCs, we

investigated which genes where differentially expressed between these two populations

of GCs at each stage of development, retaining only genes with a mean log fold change

(LogFC) > 0.25 and an FDR-adjusted p-value < 0.05. Since we had a relatively small

number of GCs at E9.0 and E10.5 (56 and 71, respectively), we combined them into a

single group called pre-sex determination (pre-SD). As expected, few genes (20) were

differentially expressed prior to sex determination (**Fig. 1G**). We then observed a sharp

increase in differentially expressed genes (DEGs) with 101 genes at E11.5, whose

functional annotations are linked to chaperonine complex and proteasome/proteolysis

(see **Tables S2** and **S3** for more detail about DEGs and GO biological process enrichment

analysis). Then, 399, 1572 and 3,236 DEGs were identified at E12.5, E13.5 and E16.5,

respectively with the XX DEGs associated with meiotic cell cycle, female gamete

generation and synaptonemal complex assembly, while the XY DEGs were enriched for

RNA splicing and posttranscriptional regulation functions. In XX GCs, a substantial

proportion of the DEGs were located on the X chromosome, in particular at E11.5 (**Fig.**

S1F). These results are consistent with the gradual reactivation of X-linked genes after

E10.5 in XX GCs (54).

Interestingly, E12.5 XY DEGs were also associated to terms such as "axis specification",

"anterior/posterior axis specification", and "somite rostral/caudal axis specification".

Detected genes associated to this category are *Pitx2, Lhx1, Lefty1, Otx2, Wnt3, Ski,*

Tdrd5, Tmed2 and include multiple Nodal/Smad2/Smad3 targets (55). Among these

genes *Lefty1, Lefty2* and *Pycr2* were upregulated from E11.5 to 13.5 in XY GCs.

Conversely, the BMP pathway target gene *Id1* was found differentially expressed and

upregulated in XX compared to XY cells from E11.5 to E16.5. This is consistent with the

described activity of Nodal and BMP pathways in XY and XX GCs respectively (21, 56).

Taken together, these results indicate that XX and XY GCs diverge transcriptionally as

early as E11.5. At this stage, a small set of genes is differentially expressed between XX

and XY GCs, but some of them- mostly involved in BMP and Nodal/Activin pathways-

mark the first steps of transcriptional differences that will increase throughout cell

differentiation (discussed later). Afterwards, the number of differentially expressed genes

increases drastically and involves meiosis in XX GCs and multiple biological processes

in XY GCs, including RNA processing and pluripotency maintenance.

Reconstructing Gene Regulatory Networks mediating germ cell sex determination

reveals the sequential and modular architecture of germ cell development

The transcriptional state of a cell emerges from an underlying gene regulatory network

(GRN) in which the activities of a small number of key transcription factors and co-

factors regulate each other and their downstream target genes (47). To comprehensively

infer the GRNs at play during XX and XY GC sex determination, we applied the

pySCENIC pipeline (47) to our single-cell data. In brief, SCENIC links cis-regulatory

sequence information together with scRNA-seq data. It first catalogs coexpression (or

repression) modules between transcription factors and candidate target genes and then

identifies modules for which the regulator's binding motif is significantly enriched across

target genes; the identified modules are called regulons. Finally, the activity of each

regulon is attributed to each cell, using the AUCell method (47). For GCs, we identified

950 regulons (487 positive and 463 negative regulons) containing 11,336 genes (**Table**

S2). To compare how XX and XY GCs acquire their sex-specific identity, we selected the

406 positive regulons with binarized AUCell activity in more than 100 cells and

containing more than 5 genes, and classified them according to their expression pattern

into 30 modules (M1 to M30) along pseudotime (**Fig. 2**). We represented the smoothed

regulon expression level of XX and XY GCs with a mirror heatmap. Each module was

also functionally annotated via enrichment analyses using GO biological process and

Reactome Pathway terms. The most relevant enriched categories are displayed on **Figure**

2. Strikingly, the expression patterns revealed numerous stage and sex-specific regulon

profiles, mostly at late developmental stages (late E13.5 and E16.5). Initially, three

modules common to both XX and XY gonocytes (M1, M4 and M6, which were cell cycle

and splicing associated) were observed at early developmental stages, M1 and M6 persist

longer in XY and XX gonocytes respectively. These common modules were superseded

sequentially by a handful of transient and overlapping sex-specific regulon modules, M5

(associated to response to BMP, cell cycle arrest and FGF response), M26 (Wnt signaling

associated), M27, M29 (associated to meiosis I) in XX GCs, and M2 (RNA splicing and

cell cycle associated), M10 (cell junction, cell cycle arrest), M16 and M17 (mRNA

processing) in XY GCs. By late E13.5 and E16.5, we noted numerous oogonia-specific

like M21, M22 (Calmodulin induced events), and M30 (associated to corticosteroid

signaling pathway, Chromatin modifying enzymes and SUMOylation)   and

spermatogonia-specific modules such as M11 (response to FGF, signaling by RTK),

M15, as well as late common modules like M8 (associated to meiosis I, male gamete

generation and cell cycle arrest), M9, M13, M14, and M19 (associated with DNA

methylation).

We also selected 248 negative regulons with the same criteria and classified them

according to their expression pattern into 10 modules (M1 to M10) (**Figure S3**). Negative

regulons (-) all displayed activity in opposition to their repressing master regulator and

most of them showed a pattern with high activity at early times points, and a progressive

downregulation/repression over time (M5 to M10 containing 222 regulons). Among

them, regulons of M4(-), M5(-), M9(-) and M10(-) could be distinguished by the onset of

their sex-specific repression: regulons from M4(-), M5(-) and M9(-) showed a repression

starting earlier (E12.5) in XX cells than in XY cells. We also detected 4 modules formed

by 26 regulons with sequential transient expression: M1(-) (associated with meiosis,

DNA methylation and response to BMP), M2(-) (positive regulation of cell cycle arrest),

M3(-) (DNA methylation) and M10(-) (DNA replication). Among these, target genes

belonging to regulons of modules 1 and 10 were transiently repressed specifically in XY

GCs, in accordance with the male-specific expression of their 11 master regulators. This

implies a possible active repression of the oogonial program in XY GCs by a few master

regulators.

Overall, we identified and functionally annotated 654 regulons whose activities were

grouped under 40 modules based on their sex-specific and temporal expression. The

transient, sequential, and often sex-specific profiles likely represent a

sequential/hierarchical organization of regulatory modules required for oogonia and

spermatogonia differentiation. The regulons provide a good summary and new insights

on the regulation of the events occurring during GC differentiation, for instance, the

negative regulation of oogonial fate and the strong presence of mRNA processing

associated transcripts in XY GCs.

RNA splicing is differentially regulated in XY compared to XX germ cells

As mentioned above, we found a strong enrichment of RNA splicing and mRNA

processing terms in XY over-expressed E13.5 DEGs. Messenger RNA splicing represents

a powerful mechanism to modulate gene expression and is known to contribute to the

fine-tuning of cell differentiation programs (57). 51 DEGs are associated with mRNA

processing term and their XY-specific or XY-enriched expression profiles between E12.5

and E16.5 are illustrated in **Figure 3A**. The specific expression of genes associated with

mRNA processing in XY GCs was confirmed by *in situ* hybridization at E14.5 for the

Tbrg4, *Rbm38* and *Supt6* genes (**Fig. 3B** and **C**). *Supt6* codes for a transcription

elongation factor which also has a role in mRNAs splicing (58). *Tbrg4* gene encodes a

known mitochondrial RNA binding protein that may also have a role in cell cycle

regulation (59), while *Rbm38* is coding for an RNA binding protein involved in cell cycle

arrest and mRNA splicing (60). RNA-Scope® *in situ* hybridization analysis of these three

genes, followed by immunostaining for the DAZL GC marker, reveals that these

transcripts were colocalized with the DAZL protein and were specifically enriched in XY

GCs, consistent with our scRNA-seq data (**Figure 3C**).

The testis specific expression of RNA processing-associated genes as well as its splicing

pattern have already been investigated during adulthood in many species and is known to

play a crucial for multiple cellular decision such as mitosis-meiosis transition and stem

cell population maintenance (For reviews see (61) and (62)). However, to our knowledge,

the presence of a sex specific splicing pattern during embryonic development has not

been thoroughly investigated. DEGs overexpressed in XY GCs at E13.5 were associated

with terms such as negative regulation of mRNA processing, negative regulation of

mRNA splicing and other terms figuring a decreased splicing process. The term "negative

regulation of translation" was enriched as well, indicating most likely a global post-

transcriptional silencing of gene expression. We therefore formulated the hypothesis that

more transcripts should be unspliced in XY GCs at E13.5 compared to XX GCs.

Consequently, when looking at each gene in XY and XX GCs, the ratio of intronic/exonic

reads should be higher in XY GCs. To test this hypothesis, we counted intronic reads in

our dataset and calculated the ratio of intronic counts over total counts (exonic+intronics)

for each gene. If the ratio is high, then this indicates that more introns are retained and

consequently a reduction in mRNA splicing. We observed a small but highly significant

increase in the intronic/exonic ratio in XY GCs (p-value$<10^{-16}$) (**Fig. 3D**) consistent with

a reduction in mRNA splicing. This was confirmed at the level of individual genes.

Indeed, we have identified genes known for their importance in spermatogenesis, such as

Bnc2, *Dmrt1* and *Tet1*, or cell cycle control such as *Ccnd3* whose transcripts have a

significantly higher unspliced ratio in XY compared to XX GCs (**Fig. 3E**) (63-66).

Overall, we observed that an important set of RNA processing protein associated

transcripts are present in developing XY GCs compared to XX GCs. Moreover, we could

infer that RNA splicing is reduced in XY GCs compared to XX GCs, suggesting that this

type of post-transcriptional regulation plays an important role in the sex-specific

differentiation of the GCs.

The Nodal/Activin and BMP pathways are important early players in the sexual

differentiation of GCs.

BMP signaling and the BMP target Zinc Finger GATA Like Protein 1 (ZGLP1) have

been reported to be essential for the oogenic fate determination and meiosis entry (12,

56). It has also been shown that the BMP pathway can be repressed at the ligand-receptor

level by Nodal/Activin (NA) activity under certain conditions, notably in stem cells (67).

In addition, the NA pathway is specifically active in XY GCs from E11.5 to E13.5 and its

inhibition allows meiosis of TGC in cultured gonads (21). We therefore investigated the

expression of NA and BMP downstream genes in our dataset. As illustrated on **Figure**

4A, we found that NA and BMP target genes are specifically expressed as early as E11.5

in XY GCs and XX GCs respectively and are reciprocally silent in the other sex.

In order to identify factors secreted by supporting cells potentially activating or inhibiting

the NA and BMP pathways in GCs, we used single-cell RNA sequencing data (scRNA-

seq) from somatic XX and XY NR5A1$^+$ cells during their differentiation process into

granulosa and Sertoli cells (68). We found that *Bmp2* was specifically expressed in pre-

granulosa cells as early as E11.5 and not in Sertoli cells (**Fig. 4B**) as described previously

(33). We also observed that the gene coding for follistatin (*Fst*), a potent inhibitor of

Activin pathway (69), was also overexpressed in pre-granulosa cells from E12.5 onward

(**Fig. 4C**). Conversely, we observed a strong sexual dimorphism and overexpression of

the beta subunit of activin (*Inhbb*) and the BMP inhibitor *Grem2* in Sertoli cells from

E12.5 (**Fig. 4D** and **E**).

Taken together, these findings indicate that GC fate determination in mice occurs as early

as E11.5 with the activation of the NA and BMP pathways in XX and XX GCs,

respectively. Interestingly, our data suggest that XX and XY supporting cells express and

secrete early on factors such as *Bmp2*, *Fst*, *Inhbb* and *Grem2* that either promote one

pathway or prevent the activation of the other (**Fig. 4F**).

Error! Bookmark not defined.

Ectopic adrenal germ cells also enter into meiosis, but numerous major

transcriptional regulators of oocyte differentiation are absent or downregulated

While the majority of PGCs migrate toward the gonadal ridges, a small fraction of GCs

are lost along the way and end up in extragonadal organs such as the nearby adrenal

glands and mesonephroi (70-72). These adrenal GCs (AGCs), irrespective of their genetic

sex, have been reported to undergo meiosis, differentiate into oocytes and display

morphological characteristics identical to those of young oocytes in primordial follicles

before disappearing around 3 weeks of age (71, 72). In the developing adrenal, AGCs are

not surrounded by supporting cells such as granulosa cells in the ovary and Sertoli cells in

the testis. So, to evaluate how an extragonadal somatic environment affects GC fate, we

investigated at the transcriptional level how ectopic adrenal GCs enter into meiosis and

commit toward the female fate.

We conducted time-series 3' single-cell RNA sequencing of developing mouse adrenal

cells from E12.5, E13.5, E16.5, and E18.5 XY embryos and identified 312 adrenal GCs

based on the expression of the classical GC markers *Dazl*, *Ddx4* and *Mael* (see **Material**

& Methods). Overall, we captured 187 AGCs at E12.5, 92 cells at E13.5, 18 cells at

E16.5, and 15 cells at E18.5. The relatively low number of GCs at later stages may reflect

the smaller proportion of GCs in the growing adrenal glands or their premature death in

this environment. UMAP representation of these 312 XY AGCs combined with our

14,718 gonadal GCs revealed that the transcriptome of XY AGCs partially overlapped

with the transcriptome of XX ovarian germ cells (OGCs), suggesting that XY AGCs

enter into meiosis and differentiate into oocytes in synchrony with gonadal oocytes (**Fig.**

5A).

To investigate further the similarities and differences between AGCs, testicular germ

cells (TGCs) and OGCs, we ran differential expression analysis between these three cell

populations at meiosis onset (E12.5 and E13.5) (**Table S4**). We found that the

comparison between AGCs and TGCs yielded more differentially expressed genes than

the comparison between OGCs and AGCs, in particular at E13.5 (**Fig. 5B**). Enrichment

analysis of GO terms revealed that terms associated with meiosis are enriched in genes

over-expressed in AGCs compared to TGCs while terms associated with splicing and

mRNA processing are enriched in TGCs over-expressed genes (**Table S5**). Likewise,

splicing ratios were very similar between AGCs and OGCs, confirming that AGCs have

overall transcriptional characteristics comparable to OGCs (**Fig. 5C**).

Despite these strong transcriptomic similarities, some significant differences could

nevertheless be observed between AGCs and OGCs. In general, meiosis-related genes

showed similar patterns and levels of expression between AGCs and OGCs. This is the

case for *Stra8*, *Sycp1*, *Sycp3, Sync3, Spo11*, *Ccdc155*, *Dmc1*, *Mei1*, *Mei4*, *Meioc*,

Hormad1, *Hormad2*, *Msh5*, *Tex11*, *Prdm9, Zglp1* and *Smc1b* (**Fig. 4D** and **Fig. S5**).

However, a notable exception was *Rec8*, whose expression is reduced in the adrenals

compared to ovarian GCs (**Fig. S4A**). These results confirmed published data that

meiosis is not significantly affected in ectopic AGCs (71, 72). However, we found

numerous key female genes exhibiting either downregulation or a testis-like profile, for

example, genes involved in the WNT-β−catenin pathway (*Axin2, Lef1,* and *Sp5*) and

BMP target genes transcription factor genes such as *Gata2*, *Id1*, *Id2*, *Id3*, *Cdx2*, *Smad6*,

Dlx3, *Msx1* and *Msx2*, the cell cycle gene *E2f1*, and the oocyte-specific basic helix-loop-

helix transcription factor gene *Figla* (**Fig. 4D** and **Fig. S4A and B**). Finally, we found

that most genes involved in male GC fate were not upregulated in AGCs with few notable

531 exceptions including the retinoic receptor gene *Rara* and the male GC regulator gene

532 *Ctcfl.* Some other genes such as the NODAL target genes *Lefty1, Lefty2, Pitx2, Lhx1* (a

533 spermatogonial stem cell self-renewal gene) and *Otx2* and the male fate marker *Nanos2*

534 showed an intermediate expression between OGC and TGC level. (**Fig. 4D** and **Fig.**

535 **S4B**). Overall, these results indicated that ectopic adrenal XY GCs enter into meiosis at

536 around the same time as ovarian XX GCs (and unlike their XY counterparts in the testis),

537 but numerous genes related to both the female and male genetic programs were mis-

538 regulated.

540 **Discussion**

541 This study represents the first large-scale scRNA-seq analysis of XX and XY GCs

542 throughout the gonadal sex determination and differentiation process. The large number

543 of individual GCs profiled, allowed us to reconstruct a near continuous representation of

544 changes in gene expression occurring during the process of gonadal differentiation,

545 including the mitosis-to-meiosis switch in GCs in developing ovaries, and spermatogonial

546 commitment and differentiation in fetal testes. This represents a major advance beyond

547 previous work and has broad implications for studies in GC development, sex

548 determination and the etiology of human GC diseases. We also provide a user-friendly

549 website to display the expression of any gene expressed in XX and XY GCs during

550 gonadal sex determination using a UMAP representation at **http://neflabdata.info/.**

How do GCs commit to and acquire sex-specific fates during differentiation of the gonad

into a testis or an ovary? From our time course scRNA sequencing profiling of six

developmental stages encompassing gonadal sex differentiation, we could observe sex

specific transcriptional patterns as early as E11.5. At this stage, part of this differential

expression can be explained by the X chromosome reactivation occurring in XX GCs.

However, we found that the Nodal/Activin and BMP pathways are important early actors

in the sex-specific differentiation of GCs with the early and sex-specific expression of

multiple target genes of the Nodal/Activin pathway expressed from E11.5 onward in XY

GCs as well as numerous target genes of the BMP pathway upregulated in XX GCs. In

addition, we found that RNA splicing is differentially regulated with increased unspliced

mRNAs in XY GCs around E13.5 potentially promoting male GC fate, regulating mitotic

arrest, inhibiting meiotic entry and/or antagonizing the female GC fate.

Transient, sequential – and often sex-specific – profiles of regulatory modules

mediating oogonia and spermatogonia differentiation

Here we uncover the gene regulatory networks (GRNs) driving sexual differentiation of

the GCs in a comparative way. We have comprehensively constructed the GRNs for XX

and XY GCs during the process of sex differentiation. We found that specific

combinations of transcription factors drive sex-specific transcriptomes in GCs and that

the gene regulatory circuitry mediating GC sex determination is composed of 654

positive regulons that can be grouped into 40 modules, each of them exhibiting transient,

sequential and often sex-specific profiles. The fact that regulons are grouped into

modules displaying transient sex-specific profiles suggests a sequential organization of

regulatory modules that work together to fine-tune the interrelated cellular events that

lead to XX and XY GC differentiation. The master regulator genes present in each

specific module may not be controlling a single cellular event, but instead a combination

of overlapping sex-specific developmental processes including mitotic arrest,

prospermatogonia commitment and *de novo* methylation for XY GCs, as well as entry

into meiosis, and suppression of pluripotency genes for XX GCs.

The GRN inference analysis also provides an opportunity to identify new critical master

regulators of GC sex determination. While various master regulators have already been

implicated in playing a key role in pluripotency and GC sex-specific commitment and

differentiation, such as *Dazl*, *Pou5f1*, *Dmc1*, *Rec8*, *Stra8*, *Nodal*, *Nanos2*, and *Dnmt3l*,

our analysis predicted more than 800 positive and negative regulons (**Fig. 2, Fig. S3** and

Table S2), including a large set of new potentially critical regulators of GC commitment

and differentiation, for example RAD21, YBX1, EGR4, ETV4, ETV5, KDM5A,

KDM5B, NR3C1 and PHF8. These predictions provide an important framework and

guide for future experimental investigation.

Messenger RNA processing and intron retention as post-transcriptional regulation

in developing mammalian germ cells

Another level of control of gene expression and function lies in RNA maturation,

stabilization and degradation (57). We observed a global post-transcriptional silencing of

gene expression in XY GCs around E13.5 potentially involved in repressing premature

meiosis entry, regulating mitotic arrest and promoting male GC fate. Not only we found

numerous DEGs overexpressed in XY GCs associated with terms such as negative

regulation of mRNA processing and splicing but also determined that RNA splicing is

globally reduced in XY GCs compared to XX GCs. We found a large variation in the

ratios of unspliced/spliced transcripts between XX and XY GCs suggesting that intron

retention is another differential feature of XY GCs, allowing them to adopt the

appropriate sexual fate through sex-specific differential expression. Finally we provide

example of genes known for their importance in spermatogenesis, such as *Bnc2*, *Dmrt1*

and *Tet1* (63-65), or cell cycle control such as *Ccnd3* (66) which all displayed increased

levels of unspliced mRNAs in XY GCs. Intron retention has been shown to be a

prominent feature of the meiotic transcriptome of mouse spermatocytes (73). It can either

favor accumulation, storage, and timely usage of specific transcripts during highly

organized cell differentiation programs or cause transcript instability at specific

developmental transitions (73-77). The temporal or sex-specific variation in intronic

retention appears to be surprisingly frequent during the process of GC sex determination

and may ensure proper and timely expression of selected transcripts. Full length isoform

single cell sequencing with long reads would further illuminate and validate our splicing

results and the importance for the male specific differentiation of GCs (78, 79).

Nodal/Activin and BMP show opposite activity patterns in XX and XY GCs and may be a kick-starter of germ cell sexual differentiation.

Historically, STRA8 has been described as the only gatekeeper that engages the meiotic program in developing female GCs (10). The onset of *Stra8* expression in GCs of the developing ovary and its lack of expression in GCs of the developing testis led to a search for the presence of the female "meiosis-initiating substance" (MIS) or male "meiosis-preventing substance" (MPS) (17, 80). While RA has emerged as a potential MIS (13, 14), recent reports revealed that female GCs enter meiosis normally even in the absence of RA signaling (15, 16, 81). Recently Nagaoka et al. showed that BMP signaling and its downstream transcriptional regulator ZGLP1 were essential for the oogenic fate, with RA signaling assisting in the maturation of the oogenic program as well as to the repression of the PGC program (12). On the other hand, it is interesting to note that many BMP target genes such as *Gata2*, *Id1*, *Id2*, *Id3*, *Msx1*, *Msx2*, *Cdx2* were not over-regulated in AGCs and have an expression profile closer to TGCs (**Fig. S5**). This suggests that a strong BMP signalling is dispensable for meiosis initiation and specification of female GC fate, at least in the adrenal gland (**Fig. 5A**). It also suggests that upregulation of *Zglp1* in AGCs requires little or no BMP activity. However, as AGCs do not survive later in adrenal gland, we can't exclude BMP signaling may be important for cell survival and complete oogenic differentiation.

There are clear evidence that the somatic environment of the embryonic gonad, in

particular the supporting lineage, influence GC sex determination through the secretion of

factors that act either as promoting or inhibiting factors (17). Investigating the expression

profile of developing Sertoli cells and granulosa cells, we identified several sex

determining cues secreted by these supporting cells that promote or inhibit the BMP and

Nodal/Activin signaling pathways. On one side, the granulosa cells promote the BMP

pathway through the expression of *Bmp2* and repress Activin activity via the expression

of *Fst*. Conversely, Smad2/3 activity, which is intrinsically triggered by GCs by Nodal

expression is reinforced by *Inhbb* which is expressed by Sertoli cells, which also repress

the BMP pathway through the expression of *Grem2*. GREM1 and GREM2 are members

of the DAN family of BMP inhibitor that bind BMPs and prevent them from associating

with their receptors (82, 83). Both *Grem1* and *Grem2* have been shown to be expressed in

the postnatal ovary and to have roles in regulating follicular development (84-86).

Whether GREM2 has or not an MPS activity in the developing testis requires further

investigations.

Comparing adrenal and gonadal germ cells development provides a tool to

investigate the importance of the somatic environment in the process of germ cells

differentiation.

By comparing the transcriptome of adrenal and gonadal GCs, we have been able to

investigate how GCs respond to three different somatic environments: the adrenal,

ovarian and testicular environments. It allowed us to also investigate whether the gene

regulatory circuitries mediating GC sex determination, composed of 950 regulons, are

interconnected or act independently. The dynamic expression pattern of key marker genes

of meiosis is strikingly similar in both AGCs and OGCs, suggesting that the initiation and

maintenance of meiosis proceeds relatively normally in AGCs, despite their XY

karyotype. However, we also observed a significant alteration in the expression of some

key female master regulator genes as well as upregulation of some male specific genes,

indicating that the somatic environment in the adrenal gland cannot completely support a

female fate for these gonocytes. In particular, we observed a lack of upregulation of

genes involved in the canonical WNT/β−catenin signaling pathway (*Axin2, Lef1,* and

Sp5), suggesting that GCs in this environment are unable to respond to WNT signals, or

to express their receptors (87-89). The expression of other genes downstream of the BMP

or NA signalling pathways such as transcription factors *Id1*, *Id2*, *Id3*, *Dlx3*, *Gata2*, *Msx1*

and *Msx2* and *Cdx2* (90) as well as the oocyte-specific basic helix-loop-helix

transcription factor gene *Figla* also displayed significant alteration. The findings that

downstream genes for both the WNT and BMP signaling pathways are not turned on in

AGCs (with the notable exception of *Zglp1*) suggest that these two pathways may be

dispensable to initiate meiosis and the female GC fate. Interestingly, the absence of *Msx1*

and *Msx2* expression may explain why *Rec8* expression, but not other meiotic genes such

as *Stra8*, is blunted in adrenal GCs (**Fig. S4B**). Based on our GRN analysis, we found

that both MSX1 and MSX2 are predicted to positively regulate *Rec8* expression.

The large majority of male-specific master regulators are not expressed in XY AGCs with

few notable exceptions such as the NA responsive genes *Lefty1*, *Lefty2* and *Lhx1*

suggesting partial suppression of the NA activity in XY AGCs. Other genes normally

expressed in TGCs such as *Nanos2, Rara* and *Ctcfl* are also expressed in AGCs

supporting the idea that overall, the adrenal environment does not provide all the

necessary signal(s) required to commit GCs to oogenesis, resulting in AGCs

characterized by an altered ovarian identity despite meiosis entry.

Compiling single cell transcriptomes from mouse GCs at six developmental stages during

the process of sex determination, both in gonadal and extragonadal tissues, allowed us to

provide a comprehensive insight into the sex-specific genetic programs and gene

regulatory networks that regulate GC commitment toward a male or female fate. As such,

we have created a valuable and rich resource for the broad biology community that

should provide insights into how this fundamental decision impacts the etiology of

infertility and human gonadal GC tumors, two of the main clinical consequences of

defects in GC sex determination.

Acknowledgments:

This work was supported by grants from the Swiss National Science Foundation (grants

31003A_173070 and 51PHI0-141994) and by the Département de l'Instruction Publique

of the State of Geneva (to S.N.). We thank Luciana Romano and Deborah Penet for the

sequencing, Christelle Borel (GEDEV department, University of Geneva) for her advice

and help with 10X technology, Cécile Gameiro and Gregory Schneiter (Flow Cytometry

Facility, University of Geneva), the team of the Animal Facility (Faculty of Medicine,

University of Geneva), Julien Prados (Basic Neuroscience, University of Geneva) for his

help for pseudotime computation and Valentin Durand Graphic Design for help with

artwork. We thank also Andy Greenfield, the members of the Nef and Dermitzakis

laboratories for helpful discussion and critical reading of the manuscript.

Author Contributions

704   Conceptualization, S.N.; data generation and investigation, C.M., Y.N., P.S., A.A.C., I.S.

705   and F.K.; Formal Analysis and Data Curation, C.M, C.M.R. and P.S.; Writing – Original

706   Draft, S.N., C.M., and M.-C.C.; Funding Acquisition, S.N., and E.T.D.; Resources, S.N.,

707   and E.T.D.; Supervision, S.N., and E.T.D.

Declaration of Interests

709   The authors declare no competing interests.

References

References

1.   Byskov, A. G., and Saxen, L. (1976) Induction of meiosis in fetal mouse testis in vitro.
Dev Biol **52**, 193-200
2.   Evans, E. P., Ford, C. E., and Lyon, M. F. (1977) Direct evidence of the capacity of the
XY germ cell in the mouse to become an oocyte. *Nature* **267**, 430-431
3.   McLaren, A. (1983) Sex reversal in the mouse. *Differentiation* **23 Suppl**, S93-98
4.   Saitou, M., and Yamaji, M. (2012) Primordial germ cells in mice. *Cold Spring Harb*
Perspect Biol **4**
5.   Tam, P. P., and Snow, M. H. (1981) Proliferation and migration of primordial germ cells
during compensatory growth in mouse embryos. *J Embryol Exp Morphol* **64**, 133-147
6.   Bullejos, M., and Koopman, P. (2004) Germ cells enter meiosis in a rostro-caudal wave
during development of the mouse ovary. *Mol Reprod Dev* **68**, 422-428
7.   Menke, D. B., Koubova, J., and Page, D. C. (2003) Sexual differentiation of germ cells in
XX mouse gonads occurs in an anterior-to-posterior wave. *Dev Biol* **262**, 303-312
8.   Yao, H. H., DiNapoli, L., and Capel, B. (2003) Meiotic germ cells antagonize
mesonephric cell migration and testis cord formation in mouse gonads. *Development* **130**,
5895-5902
9.   Adams, I. R., and McLaren, A. (2002) Sexually dimorphic development of mouse
primordial germ cells: switching from oogenesis to spermatogenesis. *Development* **129**,
1155-1164
10. Baltus, A. E., Menke, D. B., Hu, Y. C., Goodheart, M. L., Carpenter, A. E., de Rooij, D.
G., and Page, D. C. (2006) In germ cells of mouse embryonic ovaries, the decision to
enter meiosis precedes premeiotic DNA replication. *Nat Genet* **38**, 1430-1434
11. Ishiguro, K. I., Matsuura, K., Tani, N., Takeda, N., Usuki, S., Yamane, M., Sugimoto,
M., Fujimura, S., Hosokawa, M., Chuma, S., Ko, M. S. H., Araki, K., and Niwa, H.
(2020) MEIOSIN Directs the Switch from Mitosis to Meiosis in Mammalian Germ Cells.
Dev Cell **52**, 429-445 e410
12. Nagaoka, S. I., Nakaki, F., Miyauchi, H., Nosaka, Y., Ohta, H., Yabuta, Y., Kurimoto,
K., Hayashi, K., Nakamura, T., Yamamoto, T., and Saitou, M. (2020) ZGLP1 is a
determinant for the oogenic fate in mice. *Science* **367**
13. Bowles, J., Knight, D., Smith, C., Wilhelm, D., Richman, J., Mamiya, S., Yashiro, K.,
Chawengsaksophak, K., Wilson, M. J., Rossant, J., Hamada, H., and Koopman, P. (2006)
Retinoid signaling determines germ cell fate in mice. *Science* **312**, 596-600
14. Koubova, J., Menke, D. B., Zhou, Q., Capel, B., Griswold, M. D., and Page, D. C. (2006)
Retinoic acid regulates sex-specific timing of meiotic initiation in mice. *Proc Natl Acad*
Sci U S A **103**, 2474-2479
15. Vernet, N., Condrea, D., Mayere, C., Feret, B., Klopfenstein, M., Magnant, W., Alunni,
V., Teletin, M., Souali-Crespo, S., Nef, S., Mark, M., and Ghyselinck, N. B. (2020)
Meiosis occurs normally in the fetal ovary of mice lacking all retinoic acid receptors. *Sci*
Adv **6**
16. Chassot, A. A., Le Rolle, M., Jolivet, G., Stevant, I., Guigonis, J. M., Da Silva, F., Nef,
S., Pailhoux, E., Schedl, A., Ghyselinck, N. B., and Chaboissier, M. C. (2020) Retinoic
acid synthesis by ALDH1A proteins is dispensable for meiosis initiation in the mouse
fetal ovary. *Sci Adv* **6**, eaaz1261

17. Kocer, A., Reichmann, J., Best, D., and Adams, I. R. (2009) Germ cell sex determination
in mammals. *Mol Hum Reprod* **15**, 205-213
18. Spiller, C., and Bowles, J. (2019) Sexually dimorphic germ cell identity in mammals.
Curr Top Dev Biol **134**, 253-288
19. Western, P. S., Miles, D. C., van den Bergen, J. A., Burton, M., and Sinclair, A. H.
(2008) Dynamic regulation of mitotic arrest in fetal male germ cells. *Stem Cells* **26**, 339-
347
20. McLaren, A. (1984) Meiosis and differentiation of mouse germ cells. *Symp Soc Exp Biol*
38, 7-23
21. Souquet, B., Tourpin, S., Messiaen, S., Moison, D., Habert, R., and Livera, G. (2012)
Nodal signaling regulates the entry into meiosis in fetal germ cells. *Endocrinology* **153**,
2466-2473
22. Spiller, C. M., Bowles, J., and Koopman, P. (2012) Regulation of germ cell meiosis in the
fetal ovary. *Int J Dev Biol* **56**, 779-787
23. Spiller, C. M., Feng, C. W., Jackson, A., Gillis, A. J., Rolland, A. D., Looijenga, L. H.,
Koopman, P., and Bowles, J. (2012) Endogenous Nodal signaling regulates germ cell
potency during mammalian testis development. *Development* **139**, 4123-4132
24. Western, P. S., van den Bergen, J. A., Miles, D. C., and Sinclair, A. H. (2010) Male fetal
germ cell differentiation involves complex repression of the regulatory network
controlling pluripotency. *FASEB J* **24**, 3026-3035
25. Suzuki, A., and Saga, Y. (2008) Nanos2 suppresses meiosis and promotes male germ cell
differentiation. *Genes Dev* **22**, 430-435
26. La Salle, S., Mertineit, C., Taketo, T., Moens, P. B., Bestor, T. H., and Trasler, J. M.
(2004) Windows for sex-specific methylation marked by DNA methyltransferase
expression profiles in mouse germ cells. *Dev Biol* **268**, 403-415
27. Aravin, A. A., Sachidanandam, R., Bourc'his, D., Schaefer, C., Pezic, D., Toth, K. F.,
Bestor, T., and Hannon, G. J. (2008) A piRNA pathway primed by individual transposons
is linked to de novo DNA methylation in mice. *Mol Cell* **31**, 785-799
28. Ginsburg, M., Snow, M. H., and McLaren, A. (1990) Primordial germ cells in the mouse
embryo during gastrulation. *Development* **110**, 521-528
29. Gkountela, S., Zhang, K. X., Shafiq, T. A., Liao, W. W., Hargan-Calvopina, J., Chen, P.
Y., and Clark, A. T. (2015) DNA Demethylation Dynamics in the Human Prenatal
Germline. *Cell* **161**, 1425-1436
30. Guo, F., Yan, L., Guo, H., Li, L., Hu, B., Zhao, Y., Yong, J., Hu, Y., Wang, X., Wei, Y.,
Wang, W., Li, R., Yan, J., Zhi, X., Zhang, Y., Jin, H., Zhang, W., Hou, Y., Zhu, P., Li, J.,
Zhang, L., Liu, S., Ren, Y., Zhu, X., Wen, L., Gao, Y. Q., Tang, F., and Qiao, J. (2015)
The Transcriptome and DNA Methylome Landscapes of Human Primordial Germ Cells.
Cell **161**, 1437-1452
31. Houmard, B., Small, C., Yang, L., Naluai-Cecchini, T., Cheng, E., Hassold, T., and
Griswold, M. (2009) Global gene expression in the human fetal testis and ovary. *Biol*
Reprod **81**, 438-443
32. Irie, N., Weinberger, L., Tang, W. W., Kobayashi, T., Viukov, S., Manor, Y. S.,
Dietmann, S., Hanna, J. H., and Surani, M. A. (2015) SOX17 is a critical specifier of
human primordial germ cell fate. *Cell* **160**, 253-268
33. Jameson, S. A., Natarajan, A., Cool, J., DeFalco, T., Maatouk, D. M., Mork, L., Munger,
S. C., and Capel, B. (2012) Temporal transcriptional profiling of somatic and germ cells

reveals biased lineage priming of sexual fate in the fetal mouse gonad. *PLoS Genet* **8**, e1002575

34. Lesch, B. J., Dokshin, G. A., Young, R. A., McCarrey, J. R., and Page, D. C. (2013) A set of genes critical to development is epigenetically poised in mouse germ cells from fetal stages through completion of meiosis. *Proc Natl Acad Sci U S A* **110**, 16061-16066

35. Molyneaux, K. A., Wang, Y., Schaible, K., and Wylie, C. (2004) Transcriptional profiling identifies genes differentially expressed during and after migration in murine primordial germ cells. *Gene Expr Patterns* **4**, 167-181

36. Rolland, A. D., Jegou, B., and Pineau, C. (2008) Testicular development and spermatogenesis: harvesting the postgenomics bounty. *Adv Exp Med Biol* **636**, 16-41

37. Rolland, A. D., Lehmann, K. P., Johnson, K. J., Gaido, K. W., and Koopman, P. (2011) Uncovering gene regulatory networks during mouse fetal germ cell development. *Biol Reprod* **84**, 790-800

38. Small, C. L., Shima, J. E., Uzumcu, M., Skinner, M. K., and Griswold, M. D. (2005) Profiling gene expression during the differentiation and development of the murine embryonic gonad. *Biol Reprod* **72**, 492-501

39. Soh, Y. Q., Junker, J. P., Gill, M. E., Mueller, J. L., van Oudenaarden, A., and Page, D. C. (2015) A Gene Regulatory Program for Meiotic Prophase in the Fetal Ovary. *PLoS Genet* **11**, e1005531

40. Tang, W. W., Dietmann, S., Irie, N., Leitch, H. G., Floros, V. I., Bradshaw, C. R., Hackett, J. A., Chinnery, P. F., and Surani, M. A. (2015) A Unique Gene Regulatory Network Resets the Human Germline Epigenome for Development. *Cell* **161**, 1453-1467

41. Niu, W., and Spradling, A. C. (2020) Two distinct pathways of pregranulosa cell differentiation support follicle formation in the mouse ovary. *Proc Natl Acad Sci U S A* **117**, 20015-20026

42. Stallings, N. R., Hanley, N. A., Majdic, G., Zhao, L., Bakke, M., and Parker, K. L. (2002) Development of a transgenic green fluorescent protein lineage marker for steroidogenic factor 1. *Mol Endocrinol* **16**, 2360-2370

43. McFarlane, L., Truong, V., Palmer, J. S., and Wilhelm, D. (2013) Novel PCR assay for determining the genetic sex of mice. *Sex Dev* **7**, 207-211

44. Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N. J., Livak, K. J., Mikkelsen, T. S., and Rinn, J. L. (2014) The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386

45. Park, J.-E., Polański, K., Meyer, K., and S.A., T. (2018) Fast Batch Alignment of Single Cell Transcriptomes Unifies Multiple Mouse Cell Atlases into an Integrated Landscape. *BioRxiv*

46. Teo, C. H., Vishwanthan, S. V. N., Smola, A. J., and Le, Q. V. (2010) Bundle Methods for Regularized Risk Minimization. *Journal of Machine Learning Research* **11**, 311–336

47. Aibar, S., Gonzalez-Blas, C. B., Moerman, T., Huynh-Thu, V. A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J. C., Geurts, P., Aerts, J., van den Oord, J., Atak, Z. K., Wouters, J., and Aerts, S. (2017) SCENIC: single-cell regulatory network inference and clustering. *Nat Methods* **14**, 1083-1086

48. Moerman, T., Aibar Santos, S., Bravo Gonzalez-Blas, C., Simm, J., Moreau, Y., Aerts, J., and Aerts, S. (2019) GRNBoost2 and Arboreto: efficient and scalable inference of gene regulatory networks. *Bioinformatics* **35**, 2159-2161

49.   Darde, T. A., Lecluze, E., Lardenois, A., Stevant, I., Alary, N., Tuttelmann, F., Collin, O.,
Nef, S., Jegou, B., Rolland, A. D., and Chalmel, F. (2019) The ReproGenomics Viewer: a
multi-omics and cross-species resource compatible with single-cell studies for the
reproductive science community. *Bioinformatics*
50.   Darde, T. A., Sallou, O., Becker, E., Evrard, B., Monjeaud, C., Le Bras, Y., Jegou, B.,
Collin, O., Rolland, A. D., and Chalmel, F. (2015) The ReproGenomics Viewer: an
integrative cross-species toolbox for the reproductive science community. *Nucleic Acids*
Res **43**, W109-116
51.   Soumillon, M., Necsulea, A., Weier, M., Brawand, D., Zhang, X., Gu, H., Barthes, P.,
Kokkinaki, M., Nef, S., Gnirke, A., Dym, M., de Massy, B., Mikkelsen, T. S., and
Kaessmann, H. (2013) Cellular source and mechanisms of high transcriptome complexity
in the mammalian testis. *Cell Rep* **3**, 2179-2190
52.   Telley, L., Agirman, G., Prados, J., Fièvre, S., Oberst, P., Vitali, I., Nguyen, L., Dayer,
A., and D., J. (2018) Single-cell transcriptional dynamics and origins of neuronal
diversity in the developing mouse neocortex. *BioRxiv*
53.   Hill, P. W. S., Leitch, H. G., Requena, C. E., Sun, Z., Amouroux, R., Roman-Trufero, M.,
Borkowska, M., Terragni, J., Vaisvila, R., Linnett, S., Bagci, H., Dharmalingham, G.,
Haberle, V., Lenhard, B., Zheng, Y., Pradhan, S., and Hajkova, P. (2018) Epigenetic
reprogramming enables the transition from primordial germ cell to gonocyte. *Nature* **555**,
392-396
54.   Sugimoto, M., and Abe, K. (2007) X chromosome reactivation initiates in nascent
primordial germ cells in mice. *PLoS Genet* **3**, e116
55.   Guzman-Ayala, M., Lee, K. L., Mavrakis, K. J., Goggolidou, P., Norris, D. P., and
Episkopou, V. (2009) Graded Smad2/3 activation is converted directly into levels of
target gene expression in embryonic stem cells. *PLoS One* **4**, e4268
56.   Miyauchi, H., Ohta, H., Nagaoka, S., Nakaki, F., Sasaki, K., Hayashi, K., Yabuta, Y.,
Nakamura, T., Yamamoto, T., and Saitou, M. (2017) Bone morphogenetic protein and
retinoic acid synergistically specify female germ-cell fate in mice. *EMBO J* **36**, 3100-
3119
57.   Kalsotra, A., and Cooper, T. A. (2011) Functional consequences of developmentally
regulated alternative splicing. *Nat Rev Genet* **12**, 715-729
58.   Hubbard, T., Barker, D., Birney, E., Cameron, G., Chen, Y., Clark, L., Cox, T., Cuff, J.,
Curwen, V., Down, T., Durbin, R., Eyras, E., Gilbert, J., Hammond, M., Huminiecki, L.,
Kasprzyk, A., Lehvaslaiho, H., Lijnzaad, P., Melsopp, C., Mongin, E., Pettett, R.,
Pocock, M., Potter, S., Rust, A., Schmidt, E., Searle, S., Slater, G., Smith, J., Spooner,
W., Stabenau, A., Stalker, J., Stupka, E., Ureta-Vidal, A., Vastrik, I., and Clamp, M.
(2002) The Ensembl genome database project. *Nucleic Acids Res* **30**, 38-41
59.   Wolf, A. R., and Mootha, V. K. (2014) Functional genomic analysis of human
mitochondrial RNA processing. *Cell Rep* **7**, 918-931
60.   Cho, S. J., Zhang, J., and Chen, X. (2010) RNPC1 modulates the RNA-binding activity
of, and cooperates with, HuR to regulate p21 mRNA stability. *Nucleic Acids Res* **38**,
2256-2267
61.   Licatalosi, D. D. (2016) Roles of RNA-binding Proteins and Post-transcriptional
Regulation in Driving Male Germ Cell Development in the Mouse. *Adv Exp Med Biol*
907, 123-151

62.   Pushpa, K., Kumar, G. A., and Subramaniam, K. (2017) Translational Control of Germ
893   Cell Decisions. *Results Probl Cell Differ* **59**, 175-200

894   63.   Vanhoutteghem, A., Messiaen, S., Herve, F., Delhomme, B., Moison, D., Petit, J. M.,
895   Rouiller-Fabre, V., Livera, G., and Djian, P. (2014) The zinc-finger protein basonuclin 2
896   is required for proper mitotic arrest, prevention of premature meiotic initiation and
897   meiotic progression in mouse male germ cells. *Development* **141**, 4298-4310

898   64.   Matson, C. K., Murphy, M. W., Griswold, M. D., Yoshida, S., Bardwell, V. J., and
899   Zarkower, D. (2010) The mammalian doublesex homolog DMRT1 is a transcriptional
900   gatekeeper that controls the mitosis versus meiosis decision in male germ cells. *Dev Cell*
901   **19**, 612-624

902   65.   Vincent, J. J., Huang, Y., Chen, P. Y., Feng, S., Calvopina, J. H., Nee, K., Lee, S. A., Le,
903   T., Yoon, A. J., Faull, K., Fan, G., Rao, A., Jacobsen, S. E., Pellegrini, M., and Clark, A.
904   T. (2013) Stage-specific roles for tet1 and tet2 in DNA demethylation in primordial germ
905   cells. *Cell Stem Cell* **12**, 470-478

906   66.   Motokura, T., Keyomarsi, K., Kronenberg, H. M., and Arnold, A. (1992) Cloning and
907   characterization of human cyclin D3, a cDNA closely related in sequence to the
908   PRAD1/cyclin D1 proto-oncogene. *J Biol Chem* **267**, 20412-20415

909   67.   Galvin, K. E., Travis, E. D., Yee, D., Magnuson, T., and Vivian, J. L. (2010) Nodal
910   signaling regulates the bone morphogenic protein pluripotency pathway in mouse
911   embryonic stem cells. *J Biol Chem* **285**, 19747-19756

912   68.   Stevant, I., Kuhne, F., Greenfield, A., Chaboissier, M. C., Dermitzakis, E. T., and Nef, S.
913   (2019) Dissecting Cell Lineage Specification and Sex Fate Determination in Gonadal
914   Somatic Cells Using Single-Cell Transcriptomics. *Cell Rep* **26**, 3272-3283 e3273

915   69.   Pentek, J., Parker, L., Wu, A., and Arora, K. (2009) Follistatin preferentially antagonizes
916   activin rather than BMP signaling in Drosophila. *Genesis* **47**, 261-273

917   70.   Heeren, A. M., He, N., de Souza, A. F., Goercharn-Ramlal, A., van Iperen, L., Roost, M.
918   S., Gomes Fernandes, M. M., van der Westerlaken, L. A., and Chuva de Sousa Lopes, S.
919   M. (2016) On the development of extragonadal and gonadal human germ cells. *Biol Open*
920   **5**, 185-194

921   71.   Upadhyay, S., and Zamboni, L. (1982) Ectopic germ cells: natural model for the study of
922   germ cell sexual differentiation. *Proc Natl Acad Sci U S A* **79**, 6584-6588

923   72.   Zamboni, L., and Upadhyay, S. (1983) Germ cell differentiation in mouse adrenal glands.
924   *J Exp Zool* **228**, 173-193

925   73.   Naro, C., Jolly, A., Di Persio, S., Bielli, P., Setterblad, N., Alberdi, A. J., Vicini, E.,
926   Geremia, R., De la Grange, P., and Sette, C. (2017) An Orchestrated Intron Retention
927   Program in Meiosis Controls Timely Usage of Transcripts during Germ Cell
928   Differentiation. *Dev Cell* **41**, 82-93 e84

929   74.   Edwards, C. R., Ritchie, W., Wong, J. J., Schmitz, U., Middleton, R., An, X., Mohandas,
930   N., Rasko, J. E., and Blobel, G. A. (2016) A dynamic intron retention program in the
931   mammalian megakaryocyte and erythrocyte lineages. *Blood* **127**, e24-e34

932   75.   Pimentel, H., Parra, M., Gee, S. L., Mohandas, N., Pachter, L., and Conboy, J. G. (2016)
933   A dynamic intron retention program enriched in RNA processing genes regulates gene
934   expression during terminal erythropoiesis. *Nucleic Acids Res* **44**, 838-851

935   76.   Wong, J. J., Ritchie, W., Ebner, O. A., Selbach, M., Wong, J. W., Huang, Y., Gao, D.,
936   Pinello, N., Gonzalez, M., Baidya, K., Thoeng, A., Khoo, T. L., Bailey, C. G., Holst, J.,

937		and Rasko, J. E. (2013) Orchestrated intron retention regulates normal granulocyte
938		differentiation. *Cell* **154**, 583-595
939	77.	Yap, K., Lim, Z. Q., Khandelia, P., Friedman, B., and Makeyev, E. V. (2012)
940		Coordinated regulation of neuronal mRNA steady-state levels through developmentally
941		controlled intron retention. *Genes Dev* **26**, 1209-1223
942	78.	Tian, L., Jabbari, J. S., Thijssen, R., Gouil, Q., Amarasinghe, S. L., Kariyawasam, H., Su,
943		S., Dong, X., Law, C. W., Lucattini, A., Chung, J. D., Naim, T., Chan, A., Ly, C. H.,
944		Lynch, G. S., Ryall, J. G., Anttila, C. J. A., Peng, H., Anderson, M. A., Roberts, A. W.,
945		Huang, D. C. S., Clark, M. B., and Ritchie, M. E. (2020) Comprehensive characterization
946		of single cell full-length isoforms in human and mouse with long-read sequencing.
947		*bioRxiv*, 2020.2008.2010.243543
948	79.	Lebrigand, K., Magnone, V., Barbry, P., and Waldmann, R. (2020) High throughput error
949		corrected Nanopore single cell transcriptome sequencing. *Nat Commun* **11**, 4025
950	80.	McLaren, A., and Southee, D. (1997) Entry of mouse embryonic germ cells into meiosis.
951		*Dev Biol* **187**, 107-113
952	81.	Kumar, S., Chatzi, C., Brade, T., Cunningham, T. J., Zhao, X., and Duester, G. (2011)
953		Sex-specific timing of meiotic initiation is regulated by Cyp26b1 independent of retinoic
954		acid signalling. *Nat Commun* **2**, 151
955	82.	Kattamuri, C., Luedeke, D. M., Nolan, K., Rankin, S. A., Greis, K. D., Zorn, A. M., and
956		Thompson, T. B. (2012) Members of the DAN family are BMP antagonists that form
957		highly stable noncovalent dimers. *J Mol Biol* **424**, 313-327
958	83.	Avsian-Kretchmer, O., and Hsueh, A. J. (2004) Comparative genomic analysis of the
959		eight-membered ring cystine knot-containing bone morphogenetic protein antagonists.
960		*Mol Endocrinol* **18**, 1-12
961	84.	Nilsson, E. E., Larsen, G., and Skinner, M. K. (2014) Roles of Gremlin 1 and Gremlin 2
962		in regulating ovarian primordial to primary follicle transition. *Reproduction* **147**, 865-874
963	85.	Myers, M., Tripurani, S. K., Middlebrook, B., Economides, A. N., Canalis, E., and
964		Pangas, S. A. (2011) Loss of gremlin delays primordial follicle assembly but does not
965		affect female fertility in mice. *Biol Reprod* **85**, 1175-1182
966	86.	Sudo, S., Avsian-Kretchmer, O., Wang, L. S., and Hsueh, A. J. (2004) Protein related to
967		DAN and cerberus is a bone morphogenetic protein antagonist that participates in ovarian
968		paracrine regulation. *J Biol Chem* **279**, 23134-23141
969	87.	Chassot, A. A., Gregoire, E. P., Lavery, R., Taketo, M. M., de Rooij, D. G., Adams, I. R.,
970		and Chaboissier, M. C. (2011) RSPO1/beta-catenin signaling pathway regulates oogonia
971		differentiation and entry into meiosis in the mouse fetal ovary. *PLoS One* **6**, e25641
972	88.	Chassot, A. A., Ranc, F., Gregoire, E. P., Roepers-Gajadien, H. L., Taketo, M. M.,
973		Camerino, G., de Rooij, D. G., Schedl, A., and Chaboissier, M. C. (2008) Activation of
974		beta-catenin signaling by Rspo1 controls differentiation of the mammalian ovary. *Hum*
975		*Mol Genet* **17**, 1264-1277
976	89.	Naillat, F., Prunskaite-Hyyrylainen, R., Pietila, I., Sormunen, R., Jokela, T., Shan, J., and
977		Vainio, S. J. (2010) Wnt4/5a signalling coordinates cell adhesion and entry into meiosis
978		during presumptive ovarian follicle development. *Hum Mol Genet* **19**, 1539-1550
979	90.	Lee, K. L., Lim, S. K., Orlov, Y. L., Yit le, Y., Yang, H., Ang, L. T., Poellinger, L., and
980		Lim, B. (2011) Graded Nodal/Activin signaling titrates conversion of quantitative
981		phospho-Smad2 levels into qualitative embryonic stem cell fate decisions. *PLoS Genet* **7**,
982		e1002130

    Figure legends

    Figure 1. Generation of the germ cell sex determination atlas. (A) Schematic

    representation of developing testis and ovary highlighting the major events of male and

    female GC differentiation as well as the time points used in the study. (B) Illustration of

    the experimental workflow using the 10x Genomics Chromium platform. UMAP

    projection of 14,914 GCs colored by time (C), sex (D) and pseudotime (PT) going from 0

    (E9.5 cells) to 100 (E16.5 cells) (E). (F) Mirror heatmap of 63 known genes involved in

    GC differentiation. Cells were ordered along a pseudotime. Cells with lowest PT (E10.5)

    are in the center of the figure and those with highest PT (E16.5) are on the left side for

    XX cells and on the right side for XY cells. Gene expression was smoothed to reduce

    dropout effect and obtain a better visualization of expression tendencies (expression

    scale: log normalized counts normalized per gene). The relevant processes regulated by

    these genes (cell cycle, male sex determination, meiosis, pluripotency and others) are

    indicated on the left side of the heatmap. (G) Barplot representing the number of

    significantly over-expressed genes in XX or XY cells at each stage. The numbers above

   the bars indicate the number of genes.

   Figure 2. Gene regulation network analysis reveals transient patterns of

   transcription activation during germ cell sex determination. Mirror heatmap of the

   406 regulons with positive association with their master regulator. Regulons were

   clustered in 30 modules (M1-M30) using hierarchical clustering with Spearman

1006   correlation distance. Cells were ordered according to their pseudotime score with lowest

1007   score (E9.5) in the center of the figure and highest score (E16.5) on the left side for XX

1008   cells and on the right side for XY cells. Boxes on the right display examples of master

1009   regulators of interest that are colored by dominant activity in XX (pink) or XY (blue)

1010   germ cells. In parentheses is the number of target genes for each master regulator.

1011

Figure 3. Relative abundance of immature (unspliced) and mature (spliced)

transcripts reveals sex- or gene-specific differences in RNA splicing.

(A) Mirror heatmap showing the expression profiles of 50 genes associated to the GO

term RNA processing found overexpressed in E13.5 XY GCs. Cells and genes were

ordered in the same way as in Figure 1F. (B) Expression profiles of *Tbrg4*, *Rbm38* and

Supt6, three RNA processing associated genes along pseudotime. Expression levels were

smoothed using gam method and 95% confidence interval is figured by grey areas. (C)

RNA-Scope analysis followed by immunostaining for the germ cell marker DAZL in

E14.5 XX and XY gonads reveals that *Tbrg4*, *Rbm38* and *Supt6* are enriched in XY GCs.

(D) Boxplot showing the mean ratio of unspliced reads over total number of reads for

each gene in all cells at a given time. Stars above illustrate the significance level of a

Mann-Whitney Wilcoxon test for each comparison: $p < 0.05$:*, $p < 0.001$:**, $p < 0.0001$:***.

(E) Scatter plots illustrating a time and sex specific splicing ratio in four selected genes

Ccnd3, *Bnc2m Tet1* and *Dmrt1*. The X axis represents spliced reads and the Y axis

represents unspliced reads. For each plot, a cloud of E13.5 XY GCs show a higher

number of unspliced reads compared to all other cells. Expression is in Log normalized

counts.

1029

1030

Figure 4. The Nodal/Activin and BMP pathways are important players in the sexual

differentiation of GCs.

(A) Mirror heatmap representing the expression profiles of ten BMP (pink box) and seven

Nodal/Activin (blue box) known target genes. Cells and genes were ordered in the same

way as in Figure 1F. Expression profiles of *Bmp2* (B), *Fst* (C), *Inhbb* (D) and *Grem2* (E)

in pre-granulosa and (pre-)Sertoli cells at E11.5, E12.5, E13.5 and E16.5. Expression

levels are in log RPKM. Above each time point p-values from a Mann-Whitney test

comparing the mean of expression of XX and XY cells are displayed. (F) A model of

regulation of the BMP and Nodal/Activin pathways in XX and XY PGCs resulting in

primary sex differentiation of the GCs as well as meiosis priming for XX ovarian GCs.

1041

Figure 5. Altered identity and delayed meiosis in adrenal XY germ cells. (A) UMAP

projections of 14,718 gonadal cells and 312 ectopic XY GCs developing in the adrenal

colored by organ (left) and time (right). Adrenal GCs are represented in green, with larger

dots. (B) Number of differentially over-expressed genes for each comparison adrenal GC

(AGC) vs ovarian GC (OGC,) AGC vs testicular GC (TGC) and reciprocally at E12.5

and E13.5. (C) Boxplot representing splicing ratios observed in AGC, TGCs and OGCs.

(D) Expression profiles of selected genes involved in meiosis, oogonia and

spermatogonia differentiation process. The solid line represents the smoothed expression

curves of the gene in the cells ordered by pseudotime, and the fade band is the 95%

confidence interval of the model.

1052

Figure 1

Figure 4

Figure 5