Bacteria- and temperature-regulated peptides modulate beta-catenin signaling in *Hydra*

Taubenheim J[1,2], Willoweit-Ohl D[2], Knop M[2], Franzenburg S[3], He J[2], Bosch TCG[2], Fraune S[1,2]

[1]HHU Düsseldorf, Institute of Zoology and Organismic Interactions, 40225 Düsseldorf, Germany

[2]CAU Kiel, Zoological Institute, 24118 Kiel, Germany

[3]CAU Kiel, Institute of Clinical Molecular Biology, 24118 Kiel, Germany

Corresponding author: fraune@hhu.de

Abstract

Animal development has traditionally been viewed as an autonomous process directed by

the host genome. But in many animals biotic and abiotic cues, like temperature and bacterial

colonizers, provide signals for multiple developmental steps. *Hydra* offers unique features to

encode these complex interactions of developmental processes with biotic and abiotic

factors. Here, we used the model animal *Hydra* to investigate the impact of bacterial

colonizers and temperature on the pattern formation process. In *Hydra*, formation of the head

organizer involves the canonical Wnt pathway. Treatment with alsterpaullone (ALP) results in

acquiring characteristics of the head organizer in the body column. Intriguingly, germ-free

Hydra polyps are significantly more sensitive to ALP compared to control polyps. In addition

to microbes, β-catenin dependent pattern formation is also affected by temperature. Gene

expression analyses led to the identification of two small secreted peptides, named Eco1 and

Eco2, being upregulated in the response to both, *Curvibacter* sp, the main bacterial colonizer

of *Hydra*, and low temperatures. Loss-of function experiments revealed that Eco peptides are

involved in the regulation of pattern formation and have an antagonistic function to Wnt

signaling in *Hydra*.

Introduction

Organisms develop in a specific environment, which is recognized and integrated into developmental programs influencing the phenotype and fitness of an individual. Thereby, environmental factors influencing the developmental processes are very diverse like temperature (1), oxygen levels (2), social interaction (3, 4) or the associated microbiome (5). Adjusting the developmental programs to environmental conditions directly translate into the fitness of an individual and has impact on the evolution (6). The field of Ecological Evolutionary Developmental (Eco-Evo-Devo) Biology integrates these three factors into a theoretical framework (7, 8).

Numerous studies explore the effect of temperature on phenotypic differences and the impact on developmental processes (9–12). However, whether the effect is caused by a mere altered chemical reaction norm or whether temperature is actively sensed and developmental programs are adjusted accordingly, is still under debate. There are evidences for both scenarios (11, 13) and they might not exclude each other.

Similarly, the associated bacteria of an organism have been shown to affect developmental processes of the host (14). They can drive the first cleavage and determine the anterior-posterior orientation of the fertilized egg of nematodes (15), induce the morphogenesis and settlement of tubeworm larva (5), or impact the correct molting event in filarial nematodes (16). In the squid *Euprymna*, bacteria control the development of the ciliated appendages of the light organs (17) and in vertebrates they affect the maturation of the gut (18). How microbial signals or environmental cues are received and integrated into the developmental program of the host is only poorly understood. Sensory nerve cells have been shown to recognize several environmental triggers, like temperature in *C. elegans* (19, 20) and nutrients in *D. melanogaster* (21) and are able to alter phenotypic outcomes during development (22, 23). However, it is unclear whether developmental plasticity has common hubs which are triggered by several environmental cues.

The freshwater polyp *Hydra* harbors a stable microbiota within the glycocalyx of the ectodermal epithelium, which is dominated by a main colonizer *Curvibacter sp.* (24, 25). The microbiota is actively maintained by the host (26–28) and is involved in the protection against fungal infection (25). It appears likely that the microbiota has also an influence on the development of *Hydra*, as constantly occurring developmental processes such as regulation of body size are prone to environmental cues (29). *Hydra* belongs to the phylum of Cnidaria, the sister group of all bilateria. It has a radial symmetric body plan with only one body axis and two blastodermic layers, the endo- and the ectoderm (30, 31). While the stem cells reside in the body column , differentiated cells migrate into the head and foot region (32–35).

The constant proliferation and differentiation of stem cells and the migration of cells from the
body column into the extremities necessitates ongoing pattern formation processes. In
Hydra, pattern formation is mainly controlled by a Wnt signaling center in the very tip
(hypostome region) of the head (36–38). Transplantation of tissue containing the Wnt
organizer can induce a secondary axis in recipient polyps depending on the position of
excision and transplantation and follows a morphogenetic field model of diffusion reaction
(39–42) . The formation of the organizer integrates not only position information, but is also
dependent on the surrounding temperature (39). In addition, ectopic activation of Wnt-
signaling by the inhibition of the GSK3-β kinase with Alsterpoullone (ALP) is inducing stem
cell differentiation and secondary axis formations (43) by translocating and activating the
TCF transcription co-factor β-catenin into the nucleus (38). The unlimited stem cell capacity
and the constant pattern formation in *Hydra* endows the organism extremely plasticity in
terms of regeneration, body size adaption to environment and non-senescence (44–50) On
the molecular level temperature acts on the Wnt-TGF-β signaling axis influencing the
outcome of developmental decisions such as budding and size regulation of the adult polyp
(29). All these processes are reversible indicating a high degree of plasticity of the
developmental programs in *Hydra*. How the environmental cues are received and integrated
into the developmental program of the animal remains unknown.

Here, we describe the taxonomically restricted gene (TRG) *eco1* and its paralogue *eco2* to
be regulated by long-term temperature and microbiota changes in the freshwater polyp
Hydra. Changes in the expression of *eco* genes adjust the developmental decisions during
pattern formation by interference with the Wnt-signaling pathway, controlling axis stability
and continuous stem cell differentiation in *Hydra.*

Results

Temperature and bacteria modulate beta-catenin activity

To consolidate our previous finding that temperature interferes with the Wnt-dependent developmental program in *Hydra* (29) we treated polyps cultured continuously at 12 °C and 18 °C with Alsterpaullone (ALP) at 18°C (**Figure 1A**). ALP is an inhibitor for GSK3-β and causes an activation of the Wnt-signaling pathway leading to the formation of ectopic tentacles in *Hydra* (29, 43, 51). The number of ectopic tentacles can be used as a proxy to evaluate Wnt signaling strength in the animal, where higher levels of Wnt signaling leads to a higher number of tentacles. Animals reared at 18°C formed ~40% more ectopic tentacles than animals reared at 12°C (**Figure 1B**), indicating that lower temperatures decreased Wnt-signaling strength and thus plays a role in controlling axis formation and maintenance of the proliferation zone in *Hydra*.

Figure 1: Wnt-signaling is temperature dependent. (A) Animals reared at 12°C and 18°C were treated with ALP for 24 h, before assessment of ectopic tentacle formation after 96 h. (B) Lower temperature leads to fewer ectopic tentacle formation after ALP treatment. (t-test, n =12, * p<0.05) (C-F) Constitutive active Wnt signaling in β-catenin OE animals causes multiple heads/body axes at 18°C, while the severity of the phenotype was subdued when animals were reared at 12°C. (G) The number of heads produced by the β-catenin OE animals is reversible and depends on the rearing temperature, where surrounding temperatures of 12°C resulted in fewer heads per polyp (n = 10).

In order to understand how temperature is interfering with the Wnt signaling pathway, we
reared transgenic animals, carrying a constitutively active β-catenin over expression (OE)
(38) construct at 12°C and 18°C. *Hydra* polyps carrying this construct formed multiple
secondary axis and pattern formation is significantly disturbed in these animals (**Figure 1C-**
F) (38). Transferring these animals from 18°C to 12°C rescued this phenotype nearly
completely (**Figure 1C, D**) by reducing the number of heads in these animals over a course
of 26 d (**Figure 1G**). Interestingly, the effect of temperature on axis formation was reversible,
as animals with few axis reared at 12°C developed multiple axis within 26 d if cultured at
18°C (**Figure 1E, F, G**). Temperature thereby had neither an effect on the expression of
members of the Wnt-signaling pathway nor the β-catenin OE construct (**Supplementary**
Figure 1).

To test whether other environmental factors such as the associated microbiota also affects β-
catenin dependent development in *Hydra,* we performed the same ALP-treatment on animals
with and without associated bacteria (**Figure 2A**). Germfree animals responded nearly four
times more to the ectopic activation of Wnt signaling compared to control animals (**Figure**
2B-D). In a second experiment we tested, whether recolonization (conventionalizing) of the
germfree animals reduced the increased tentacle formation after ALP treatment and found a
significant mitigation of the ALP effect in these animals (**Supplementary Figure 2,**
Supplementary Table 1). The observations indicate that not only higher temperatures but
also loss of host-associated bacteria increase the Wnt signaling in *Hydra* and affect
maintenance of the proliferation zone along the body column.

Figure 2: Wnt-signaling is dependent on microbial colonization. (A) Animals were ALP treated
for 24 h and number of generated ectopic tentacles were assessed 96 h after treatment, comparing
the treatment outcome of germfree (GF) and normally colonized animals (control). (b-d) Ectopic
tentacle formation is increased when colonizing bacteria were removed, suggesting a role of microbial
colonization in pattern formation in Hydra (Mann Whitney U test, n =58, *** p<0.0001).

Both temperature and bacteria influence the expression of *eco1* and *eco2*

To elucidate the underlying molecular mechanism of the environment-development interaction, we compared differentially regulated genes of a previous microarray study comparing GF to control animals (52) with a recent RNA-seq data set (29), which compared the transcriptome of animals reared at 12°C and 18°C. Both sets of differentially regulated genes overlapped in 55 contigs (**Figure 3A, Supplementary Table 2**). Within this overlap, 18 genes were regulated in the same direction upon lower rearing temperature and removed bacterial colonization (of these 14 [25,45%] down, 4 [7.27%] up). The rest of the genes (37 in total, 67.27%) showed contrary regulation in both conditions (19 [34.55%] down at lower rearing temperature, 18 [32.73%] up at lower rearing temperature) (**Supplementary Table 2**). As low temperature and removal of bacterial colonizers resulted in contrary responses to ALP we searched for genes with contradicting gene expression in this data set. We ranked the genes according their mean expression, fold change and significance level and excluded metabolic genes for further analysis (see methods for details of candidate gene selection). The contig 18166 was ranked highest in this analysis and showed highest differential gene regulation in the temperature experiment and was on position 15 in the bacterial data set among the 55 contigs. Apart from contig 18166 a paralogue contig (14187) was found to be regulated by both temperature and bacteria (position 5 and 18 in the temperature and bacterial data set, respectively). Both paralogues show high sequence homology, encode for small peptides with a predicted signal for secretion, and contain four conserved cysteine residues (**Figure 3B**). For both paralogues no homologs were detectable by BLAST search outside the taxon *Hydra*, indicating that these genes represent taxonomically restricted genes (TRGs) (53, 54).

We tested the expression of both paralogues upon temperature (**Figure 3C**) and bacterial cues (**Figure 3D**) via qRT-PCR and observed a significant up-regulation for both genes at low temperature (**Figure 3C**) and in the presence of bacterial colonizers (**Figure 3D**), confirming the initial screening result. Notably, the expression response of both genes to temperature changes was much stronger compared to the bacterial response. The spatial expression pattern of both paralogues were analyzed by whole mount *in-situ* hybridizations. At 12°C, both paralogues are expressed in the ectodermal cell layer along the whole body column, while tentacle and foot tissue showed no expression (**Figure 3E, F**). The expression at 18°C is restricted to the foot in the case of 18166 (**Figure 3G**) and to the lower budding region in the case of 14187 (**Figure 3H**) or showed no detectable level of expression at all (89%) (**Supplementary Figure 3**). In all cases the expression domain of 18166 at 18°C was more expanded than the expression of 14187 (**Figure 3G; H, Supplementary Figure 3**). These results indicate that the expression of both genes is up-regulated due to the expansion of its expression domain from a foot-restricted expression at 18°C to the expression through the whole body column at 12°C.

Figure 3: Two TRGs, expressed in the ectoderm, are potential recognition signals for environmental changes. (A) Re-analysis of microarray data, comparing germfree animals with colonized ones, and transcriptomic data, comparing animals at different rearing temperatures, revealed two candidate genes with common regulation upon environmental changes: *eco1* and *eco2*. (B) Sequence analysis of the candidate genes revealed a paralogous relationship between the two genes and a secretion signal peptide (red square), without any domain structure. (C-D) qRT-PCR assessment of *eco* gene expression in germfree animals and upon temperature shift confirms down regulation of *eco1* and *eco2* expression at higher rearing temperatures (n =3, Two-way ANOVA, Bonferroni post tests, ** p<0.01, *** p<0.001) and in disturbed microbiome conditions (n =3, Two-way ANOVA, Bonferroni post tests, * p<0.05, ** p<0.01, *** p<0.001). (E-H) *eco* genes were expressed in the foot and lower third of the animals at 18 °C rearing temperature, but the expression domain expands to the body column and parts of the head at 12 °C. (i-j) Immunohistochemistry with a polyclonal antibody raised against a fragment of Eco1 (underlined in b) show the production of the peptides in the ectodermal epithelium and packaging in vesicles localized in the apical part of the cells, which suggests a secretion of the peptide. The images are displayed in pseudocolor, green, red and blue corresponding to Eco1 (AlexaFluor488), the actin cytoskeleton (rhodamine-phalloidin, a proxy for the mesoglea) and the cell nuclei (TO-PRO3).

Using a polyclonal antibody, which was generated against a specific peptide encoded by
contig 18166 (**Figure 3B**, underlined sequence), we could observe that the peptide is
expressed in the ectodermal epithelial layer and localized in small vesicles (**Figure 3I**)
accumulating at the apical side of the epithelial cells (**Figure 3J**). This cellular localization
suggest that the peptide is secreted at the apical side of the ectodermal cells. Considering

their ecological dependence, we termed the genes *eco1* (contig 18166) and *eco2* (contig
14187), respectively.

Expression of *eco1* and *eco2* response to environmental changes within two weeks

Having confirmed that both genes respond to changes in temperature and bacterial
colonization, we assessed the expression of *eco1* and *eco2* over time in germfree animals
and two controls, conventionalized (conv) animals and wildtype polyps (**Figure 4A**). While 8
days post recolonization (dpr) the expression level of *eco1* and *eco2* in conventionalized
animals were still equivalent to the levels in germfree (GF) animals, the expression level of
both genes recovered within the second week, with rising expression levels similar to control
animals (**Figure 4A**). Furthermore, we tested if recolonization with the main colonizer
Curvibacter sp. alone is sufficient for the regulation of *eco* genes. Analyzing the expression
two weeks after recolonization we observed a recovery of the expression levels of both
genes (**Figure 4B**), indicating that the specific crosstalk between *Curvibacter* and *Hydra* is
sufficient to regulate gene expression of *eco1* and *eco2*.

To get insights into the temporal expression dynamics of *eco1* and *eco2* after temperature
change, we transferred animals cultured at 12°C to 18°C and vice versa, and monitored the
expression over the course of 28 days (**Figure 4C**). Both genes responded to temperature
changes within days, reaching a new stable expression level after around two weeks (**Figure
4C**). Thereby, *eco2* showed a higher up regulation at 12°C compared to *eco1* (**Figure 4C**),
which might reflect the fact that *eco2* is expressed at a lower level at 18°C compared to *eco1*
(**Figure 3 G, H**).

The fact that both factors, temperature and bacteria, strongly influence the expression of
eco1 and *eco2* raised the question if both factors are interacting with each other. To
disentangle both factors, we generated GF animals and maintained them at 18°C or
transferred them to 12°C (**Figure 4D-F**). We observed an increase of gene expression in
animals transferred to 12°C independent of the microbiota state of the animals (**Figure 4D-
E**), while the absence of bacteria reduced gene expression independent of the temperature
(**Figure 4F**). The effect of temperature was highly significant in an ANOVA ($p = 8.16E-9$)
which was corrected for gene variation while the bacteria effect was only marginal significant,
due to the smaller effect size ($p = 0.0557$). The interaction term of temperature and
colonization was not significant ($p = 0.2$) indicating no interaction of temperature and
colonization in the regulation of *eco* genes (**Supplementary Table 3**). These results suggest
independent gene regulation by temperature and microbiota for *eco1* and *eco2*.

Figure 4: Expression dynamics of eco genes were a long term adaptation to changing microbial state or rearing temperatures. (A) eco gene expression was reestablished by recolonization with bacteria after 14 d, but did not reach normal levels after only 8 d recolonization (n =3, Two-way ANOVA, Bonferroni posttests, * $p<0.05$, ** $p<0.01$, *** $p<0.001$) (B) Recolonization with the main colonizer Curvibacter sp. alone was sufficient to alleviate the expression suppression of eco genes after 14 d of colonization. (n =4, Two-way ANOVA, Bonferroni posttests, * $p<0.05$, *** $p<0.001$) (C) eco gene expression changes were established within 14-16 d upon rearing temperature shifts from 18 °C to 12 °C and vice versa. (D) Microbial colonization state and rearing temperature were independent inputs for the expression regulation of eco genes. Gene expression is up regulated upon temperature shifts from 18 °C to 12 °C, independent of colonization state of the animals. (E, F) The graphic displays effect size cleaned of overlaying variances. (E) The expression of *eco* genes increases within 14 days after reduction of rearing temperature, independent of the colonization status of the genes (ANOVA, $p = 8.16E-9$ for temperature, $p = 6.99E-6$ for temperature-time interaction, see Supplementary Table 3). (F) Removing bacteria from *Hydra* reduces expression of *eco* genes independent of temperature (ANOVA, $p = 0.0557$ for colonization, $p = 0.20$ for colonization-temperature interaction, see Supplementary Table 3).

In summary, temperature and bacteria dependent regulation of the two genes are reversible
and reflect a long-term acclimation to both factors, rather than a short-term regulation and an
immediate stress response. The timing of expression changes correlates with the reduction
of secondary heads in the β-catenin OE animals reared at 12°C (**Figure 1**). Thus, *eco1* and
eco2 might act as effector genes controlling phenotypic plasticity relaying environmental
cues directly to developmental pathways.
214

Eco1 and Eco2 act as antagonists to Wnt-signaling
To functionally analyze the role of Eco1 peptides, we designed a hairpin (HP) construct
based on the sequence of *eco1* fused to GFP (**Figure 5A**). We generated two transgenic
lines (B5 and B8), which displayed constitutive expression of the HP in the ectodermal
epithelial cells. The mosaic nature of genetically modified hatchlings allows for the selection
of transgenic and non-transgenic lines, which served as genetically identical control lines
(except for the hairpin construct). On the level of *in-situ* hybridization the transgenic line B8
showed a dramatically reduced expression level of *eco1* in the whole body column (**Figure
5B**) in comparison to its control line (**Figure 5C**). Checking the knock-down rate of *eco1* by
qRT-PCR in both lines revealed a strong down regulation by hairpin-mediated RNAi. Due to
high sequence similarity, the hairpin-mediated RNAi targeted also *eco2*, leading to similar
down-regulation compared to *eco1* in both transgenic lines (**Figure 5D**). While at 12°C the
rate of reduction was between 95%-100% for both genes, the knock-down rate at 18°C was
between 20%-40% (**Figure 5D**).
To test the hypothesis that Eco peptides act as antagonist to Wnt-signaling we treated both
transgenic Eco-knockdown (KD) lines with ALP and compared the number of ectopic
tentacles to control animals (**Figure 5E**). We found a significant 3-fold (B5) and 2-fold (B8)
increase in number of ectopically formed tentacles in Eco-KD animals, respectively (**Figure
5F**).

Figure 5: Knockdown of eco genes results in increased Wnt signaling. (a) Eco–hairpin construct for generation of transgenic *Hydra* (as, antisense; s, sense; TAA, stop codon; P, promoter; T, terminator) (b, c) Whole mount *in-situ* hybridization of *eco1* expression in Eco-KD (b) and control (c) animals. (d) *eco1* and *eco2* expression in two transgenic lines (B8 and B5) at 12°C and 18°C (n =3, Two-way ANOVA, Bonferroni posttests, *** p<0.001) (e-f) Treating Eco1-KD animals with ALP revealed an increased potential to form ectopic tentacles, indicating a higher Wnt signaling activity. (f) The number of heads formed after transplantation of near head tissue into the body axis of an acceptor polyp serves as a readout for the head inhibition potential, which is governed by the inhibition of the Wnt signaling. (g-h) Control (unlabeled) and Eco-KD (GFP labeled) animals served as acceptor polyps for head tissues from wild type (dsRed labeled) animals to assess the head inhibition potential under disturbed *eco* expression. (i) Eco-KD polyps showed a reduced head inhibition potential, indicating an impaired Wnt inhibition in these animals. n = 75, p=0.0085, Fisher's exact test.

Since the Wnt signaling is instructive for head regeneration, we tested, whether Eco peptides

are involved in the regeneration process. We cut Eco-KD animals reared at 12°C and 18°C in

the middle of the body column, to regenerate a head or a foot (**Supplementary Figure 4A**).

During head regeneration we observed no difference neither in the timing (**Supplementary**

Figure 4B) nor in the number of tentacles which regenerated between the Eco-KD and
control animals (**Supplementary Figure 4C**). Similarly, foot regeneration was not disturbed
in Eco-KD animals (**Supplementary Figure 4D**). This result indicates no major role of *eco*
genes in the regeneration processes of *Hydra*.

To consolidate the notion of Eco peptides being an antagonist to Wnt-signaling, we
performed transplantation experiments to measure the head inhibition potential of Eco-KD
animals. In *Hydra,* head activation and inhibition is governed by a gradient of an activator
and inhibitor, following a model first described by Alan Turing and specified by Alfred Gierer
and Hans Meinhardt (55–57). The model describes a two component system of molecules,
which can explain the head forming properties of *Hydra's* patterning processes. The idea
was experimentally tested by Harry MacWilliams in the 1970s and 80s, using transplantation
techniques (58, 59). In his experiments he described properties of the head inhibitor and
head activator in the animals, using the fact, that head near pieces have organizer
functionality (40) and can induce a head in the body column of an acceptor polyp. He
described two main findings: first, the rate of head induction increased as the site of
transplantation was further away from the head of the donor (head inhibition gradient)(59).
Second, the potential to form a head decreases with the distance to the head of the excised
pieces (head activation gradient) (58). We used a similar approach and assessed properties
of the head inhibition gradient in the Eco-KD background, by comparing the fraction of heads
formed in Eco-KD and control animals. To this end, we transplanted head near pieces
(directly beneath the tentacle ring) from control donor animals into the body column
(approximately one-third from the head) of acceptor animals. We chose the site of
transplantation (one-third length from the head) as it was reported that head formation was
medium to low in this experimental setting (60). If our notion of decreased head inhibition for
the knockdown of the *eco* gene expression was true, we expected an increased fraction of
heads formed after transplantation of head near tissue into the body column of Eco-KD
animals.

We found, that the fraction forming heads after transplantation is doubled in Eco-KD animals
(31 of 75), compared to control animals (16 of 76), indicating a reduced head inhibition
potential (**Figure 5G-I**). Together with the ALP experiment, these results demonstrated that
Eco peptides antagonize the effects of Wnt-signaling. Lastly, we checked whether *eco* gene
expression is regulated by the Wnt pathway and can act as a feedback mechanism in the
head formation process. We treated animals with different concentrations of ALP (0.2, 1, and
5 µM) for 24 h and measured *eco* expression via qRT-PCR, but could not detect a regulation
of the genes (**Supplementary Figure 5**).

Taken together these results show that the *eco* genes are able to relay environmental
signals, like bacterial colonization and temperature, to the Wnt-signaling cascade and by that
modulate axis and head formation in *Hydra* according to environmental conditions.

Figure 6: Eco peptides are environmentally triggered inhibitors of Wnt signaling. Microbial colonization and low temperature are independent inputs for Eco peptides and are able to induce its transcription. Eco peptides are synthesized as small peptides, packed into vesicles and secreted into extracellular matrix of the ectoderm. By an unknown mechanism Eco peptides are recognized and the Wnt signaling is inhibited. The inhibition takes place downstream of the β-catenin stabilization, presumably at the expression of Wnt target genes as Eco expression were able to inhibit activation of the pathway at the GSK-3β level after ALP treatment (see text).

Discussion

Wnt-signaling: an evolutionary conserved signaling hub integrating environmental signals to stem cell behavior

The Wnt-signaling pathway most likely evolved in the common ancestor of multicellular animals. Members of the pathway are present in all metazoan animals, but not in fungi, plants, or unicellular eukaryotes, and have regulatory functions in embryogenesis and cell differentiation (61).

In *Hydra* the Wnt pathway is involved in head formation (36), control of bud formation (37, 62) and the differentiation of stem cells (63, 64). Most Wnt family member are expressed in the tip of the hypostome of *Hydra* (37) and the Wnt pathway has been shown to form the activating agent in the head organizer of *Hydra* (36, 37). Wnt corresponds to the head inducer in the Gierer-Meinhardt model, which describes a self-organizing system of head activation (HA) and head inhibition (HI) and is suitable to explain regeneration and axis

formation in *Hydra* (42). In this model the HI would correspond to a molecule/gene which is
activated by Wnt and is inhibiting the Wnt pathway. Thus, the expression of such a gene
would be expected to be expressed in a graded manner from head to foot. Here, we show
that Eco peptides antagonizes the Wnt-signaling pathway (**Figure 6**) by reducing both
ectopic tentacle formation after ALP treatment (**Figure 5F**) and by transplantation
experiments (**Figure 5H-J**). However, the expression domain of *eco* genes is restricted to the
foot at 18°C and expands to the head region only, if the animals were reared at 12°C for at
least 14 days (**Figure 3E-H, Figure 4C**). The expansion of the expression domain of *eco*
genes corresponds to a dramatic increase of the gene expression level, which seems not to
appear graded in any form (**Figure 3C, E-H**). Eco peptides do thus not equal the proposed
HI of the Gierer-Meinhardt model and therefore add another layer to the Wnt-pathway
regulation in *Hydra*.
The properties of the HI in *Hydra* were elaborately determined by transplantation
experiments by MacWilliams (39). Inhibitor level changes occurred during regeneration within
6-24 h after ablation, indicating that the here described inhibitor of the Wnt signaling acts on
completely different time scales, than the immediately responding HI. MacWilliams further
noted that temperature changes had an influence on the HI potential of *Hydra.* However,
these temperature changes were also applied only several hours before transplantation,
where *eco* expression changes have not occurred (**Figure 4C**). We therefore argue, that Eco
peptides do not contribute to the immediate organizing function of Wnt during head
formation, but is rather adjusting background levels of Wnt signals throughout the body. This
notion is confirmed by the undisturbed regeneration of Eco-KD animals (**Supplementary**
Figure 4) and the otherwise normal appearance of the polyps (including regular budding). A
regulation of the background Wnt signaling strength by Eco peptides seems reasonable
because Wnt signaling is important for size regulation in *Hydra* by means of bud initiation
(29*,* 62*)* and body size is regulated with temperature (29, 49). In addition, the variation of
multiple axes in the β-catenin OE animals at different rearing temperatures occurred in the
same time frame as *eco* gene regulation upon temperature shift (**Figure 1C-G**). Altogether,
this could imply a function of Eco peptides in adjustment of Wnt signaling strength as they
are regulated with temperature and antagonize Wnt signaling. However, Eco-KD animals
showed now obvious size differences in our cultures, which indicates, that the temperature-
Wnt regulation seems to be more complex and can not be reduced to Eco signaling alone.
Notably, the proposed HI of the Gierer-Meinhardt model remains enigmatic on the molecular
level till today, though various candidates have been proposed, for instance a Hydra dickkopf
homolog (65, 66), the transcription factor Sp5 (67), or the glycoprotein thrombospondin (68),
but none of them seem to resemble all assumed properties of the HI in the Gierer-Meinhardt
model. Further, several downstream genes of the Wnt signaling pathway have been shown
to be involved in specific aspects of Wnt signaling, for example the homeobox gene *rax* is

important for tentacle formation (69), PKC signaling seems to be important for head
regneration (70), but not in bud formation (71), while ERK signaling was essential for both
processes (70, 71). We thus argue, that several levels of Wnt-inhibition exist and that the
activity of Eco peptides provides one mechanism. Until now, it is unclear how Eco peptides
antagonizes Wnt pathway, but we assume that they act downstream of β-catenin given the
following arguments. First, we showed that Eco-KD animals have a higher probability of
forming tentacles after ALP treatment, but have normal regeneration capability, thus Rax
could be a target transcription factor inhibited upon Eco signaling (69). Secondly, we showed
that low temperature is antagonizing the effect of over expressed β-catenin. As β-catenin
was mutated in a way that it was stabilized against APC dependent degradation (38), we
argue that the antagonistic activity is most likely downstream of β-catenin and independent of
the APC pathway. Therefore, we predict the presence of an endogenous receptor that binds
Eco peptides and activates a signaling cascade antagonizing β-catenin activity.

To date the inputs of *eco* gene regulation on the molecular level are unknown. We can
exclude, that Wnt signaling actively regulates *eco* genes, which would be expected if Eco
signaling is involved in the immediate head formation process. However, one could speculate
on acyl-homoserine-lacton (AHL) as a possible recognition molecule. We previously showed
that AHLs produced by *Curvibacter* are quenched by the *Hydra* host to avoid detrimental
gene expression of *Curvibacter* on the host (24). This direct interaction indicates an active
recognition of AHLs by the host and would mediate a possible route of regulating *eco* gene
expression. From the microarray data we can exclude also a MyD88 mediated *eco*
regulation, as MyD88-KD animals showed no differences in expression level of *eco* genes
(52).

Predicting inputs for the temperature signal is unequally more difficult, as to date no definite
temperature sensing mechanism has been found in *Hydra*. TRPs are known to sense high
and low temperature and have been identified in *Hydra* (72). TRPM3 has also been linked to
high temperature sensation (heat shock) in Hydra (73) but if similar mechanisms exist for
cold temperature or whether they can mediate the long term effect of *eco* gene expression is
unclear to date. *Eco* expression might be regulated by other head inhibiting substances as
MacWilliams has suggested, that physical properties of the HI change at different
temperatures, resulting in a longer range of action at lower temperatures, which eventually
might translate in an activated *eco* gene expression (39).

Increased Wnt-signaling causes the stem cells in *Hydra* to differentiate and to lose their
potential to self-renew (63, 64). Interestingly, the expression of both Eco peptides is
regulated by the presence of *Curvibacter*, the main bacterial colonizer of *Hydra*, and low
temperature. Individual pathways may signal both external signals to the promotor of *eco*

genes, as both factors regulate the expression of both genes independently (**Figure 6,**
Figure 4D).

Our model suggests, that the ratio of recognized Wnt and Eco peptides by an individual cell determines the activation of Wnt-target genes. Thereby, the opposing signaling outcomes of Wnt and Eco would determine the degree of differentiation in the tissue, as increased Wnt signaling causes the differentiation of stem cells (63, 64). This process is mediated by Myc1 in interstitial cells, which is directly affected by the increase of Wnt signals due to ALP treatment (64). Given, that Eco peptides are able to counteract increased Wnt signals, we thus argue that the *eco* genes integrate environmental signals into the developmental program and thereby promote the stemness of the cells in the body column of *Hydra*.

There are several studies in vertebrates that showed the integration of environmental signals into the Wnt pathway. In zebrafish, the intestinal bacterium *Aeromonas veronii* enhances β-catenin stability after recognition by MyD88 resulting in higher cell proliferation in the intestine (74). In human epithelial cells the CagA peptide produced by *Helicobacter pylori*, activates β-catenin, leading to transcriptional upregulation of genes implicated in cancer (75, 76). In addition, activation of the aryl hydrocarbon receptor by natural ligands, which are converted from dietary tryptophan and glycosinolates by intestinal microbes (77, 78), results in the degradation of β-catenin (78) and the suppression of intestinal carcinogenesis. We showed another mechanism, how bacterial signals are integrated into the Wnt signaling pathway, via an orphan peptide not present outside the *Hydra* genus. All described mechanisms are highly specific to each of the different model systems (MyD88, CagA, AhR, Eco1) and thus most likely evolved independently, which highlights the necessity of individual adaptation dependent on the life styles to cope with changing environments. However, conserved developmental signaling like the Wnt pathway seems to form signaling hubs to integrate environmental cues. The Wnt signaling pathway thereby fulfills central tasks in pattern formation as well as stem cell regulation in all metazoans and is clearly an important target to mediate developmental decisions upon environmental changes.

Orphan genes as circuit of environmental cues to host development

This study adds another example to taxonomic restricted or orphan genes as important factors for adaptations and stresses the importance of these fast evolving genes for the interaction with the environment. Orphan genes have been found to play important roles by recruiting new energy resources (79, 80), and occupying new habitats (81). Orphan genes often display functions in developmental programs (81, 82), immunity (26, 83) and mediate interaction with the environment (54, 84, 85). Here we support this notion by showing that the two *eco* paralogues interact with the environment in a way that the current conditions for

402   temperature and microbiota association are sensed and signaled to the Wnt pathway.
403   Thereby, the developmental program is tuned by environmental factors via the regulation of
404   the two orphan peptides Eco1 and Eco2.
405

Environment matters
Animal development has traditionally been viewed as an autonomous process directed by
the host genome. In recent years, it got evident that biotic and abiotic cues provide a variety
of signals that are integrated into the developmental program. These observations resulted in
the Eco-Evo-Devo concept (7, 8). Our results provide new evidences for this concept on
different levels. Firstly, our study supports the idea that development is plastic and
responsive to abiotic (temperature) and biotic (microbiota) factors, as *eco* gene expression is
highly dependent on these factors. Secondly, we show that evolution of new traits does not
necessarily demand the change in core developmental pathways, but that newly obtained
genes can act as modulators of these pathways mediating a gain of function in
developmental programs. Thirdly, we show that evolutionary young genes, which modulate
conserved developmental signaling cascades, can mediate phenotypic plasticity and
increase robustness of patterning formation after disturbance.
419

Material and Methods

Animal culture. All experiments were carried out with either wild type or transgenic *Hydra vulgaris AEP (Hydra carnea* (86, 87)) which were cultured in Hydra medium (HM, 0.28 mM CaCl$_2$, 0.33 mM MgSO$_4$, 0.5 mM NaHCO$_3$ and 0.08 mM KCO$_3$) at 18 °C with a 12/12 h light-dark cycle. Animals were fed *ad libitum* 2-3 times the week.

Transgenic animals. The β-catenin over expression construct contained a truncated β-catenin (AA Δ1-138, containing the phosphorylation site important for Wnt signaling transduction) obtained from *Hydra magnipapillata* terminally fused to eGFP in a modified Hot-G Bluescript II SK + backbone (alias: pHyVec4, Steele, Genebank: DQ385853) (38). The Eco1 hairpin constructs were cloned into a pGEM-T backbone obtained from *Hydra vulgaris* AEP (*Hydra carnea)* sequences (Compagen: HAEP_celera_v1_contig18166) containing base pair 3-303 with a 307 bp linker. dsRED and eGFP labelling constructs were codon usage optimized for *Hydra* and cloned into pGEM-T expression vectors. All constructs were driven by constitutively expressed *Hydra magnipapillata* actin promoters (~1.5 kb upstream the actin gene) and terminated by an actin terminator (~1 kb downstream of the gene). Stable transgenic animals were generated by microinjection of genetic constructs into 2-8 cell blastomere of *Hydra*. Transgenic animals hatched with a mosaic expression of eGFP, which served for selection of transgenic cells. The asexual mode of reproduction was used to select for transgenic cells and generate full transgenic animals on the on hand, while on the other hand non transgenic cells were selected to generate a non-transgenic line, which served as a control, containing the same genetic background.

Antibiotics treatment and recolonization. To render animals germ-free, we treated *Hydra* with a cocktail of ampicillin, rifampicin, streptomycin, and neomycin (50 µg/ml each) in sterile HM for two weeks and daily exchange of the medium. To control for germ-freeness a randomly chosen polyp from each batch were homogenized and plated on R2A-Agar, missing CFU indicated removal of bacteria. Furthermore, whole DNA from a randomly chosen polyp of each batch were isolated (DNeasy Blood & Tissue Kit, *Quiagen*) and a PCR for bacterial DNA contamination was performed (Eub-27F: AGAGTTTGATCCTGGCTCAG, Eub-1492R: GGTTACCTTGTTACGACTT).

To control for the effect of antibiotics in our experiments, animals were conventionalized by recolonizing them with naturally hosted bacteria after antibiotic treatment. To this end, for each germ-free *Hydra* one untreated animal were homognized, resuspended in 100 µl autoclaved HM and added to ~4 ml HM containing the germ-free animals. The animals were washed 24 h after recolonization with sterile HM to remove excess bacterial load.

Alsterpaullone experiment. Animals were fed one day prior experiment and treated with 5 µM Alterpaullone (ALP, *Sigma-Aldrich*) for 24 h to inhibit the glycogen synthase kinase 3 β (GSK3-β) and activate the Wnt signaling pathway at 18°C, regardless of previous rearing conditions. After treatment, the polyps were washed and incubated in HM before assessment of ectopic tentacle formation under the binocular 4 d after treatment. To regard any form of the temperature reaction norm we assessed the number of tentacles per polyp after tentacles were clearly developed so that a further increase in number of tentacles could not be observed, even at later time points.

Temperature treatment of β-catenin animals. β-catenin over expression animals were reared at either 18 °C or 12 °C and standard conditions prior experiment. Animals were transferred to treatment temperature HM (18 °C → 12 °C; 12 °C → 18 °C) kept as single polyps in a 12-well plate. Head structures of the visually largest animal were assessed at day 0, 5, 9, 14, 19, 22, 26, and 30. Smaller animal fragments, which might have appeared during the experiment, were removed from the cavity.

RNA extraction, cDNA generation, qRT-PCR. Total RNA was isolated by dissolving 15-25 animals in 1 ml TRIzol (*Invitrogen*) by vortexing, after starvation for 2 days. After addition of 300 µL chloroform (p. a.), samples were centrifuged (12,000 g at 4°C), the resulting aqueous phase was transferred in a clean tube, and 400 µl ethanol (99.9 %) was added. The solution was cleaned and desalted using the silica column protocol from Ambion PureLink RNA Mini Kit (*Thermo Scientific*), including an on column DNAse digestion step. 650 ng RNA was used to generate cDNA applying the protocol of the First Strand cDNA Synthesis Kit (*Fermentas*). Resulting cDNA was equilibrated within each experimental setup by semi-quantitative PCR using primer for the house keeping gene ELF1a. Equal amounts (according to PCR quantification) were used with the GoTaq qPCR Master Mix (*Promega*) in a 7300 real-time PCR system (*ABI*). For data analysis the simplified method after Pfaffl (88) was used, assuming a primer efficiency of 100 %.

Gene selection from transcriptomic analysis. We defined a gene set from the overlap of differentially expressed genes from a microarray comparing GF vs. recolonized animals and in a RNA-seq experiment comparing 12°C vs. 18°C rearing temperature (55 genes in total) (29, 52). For technical details of RNA-seq and microarray analysis, see (29, 52). This gene set was associated with ranks according mean expression, p-value and fold change for both conditions (decreasing order for fold change and mean expression, increasing order for p-values). The ranks for each category were summed, to generate a score which was used to sort the genes. The genes Eco1 (contig 18166) and Eco2 (contig 14187) were at position 2 and 7 in this list. We did not consider the first gene in the table (a serine acetyltransferase,

496 contig 18585) as it was predicted to be associated with metabolic changes which was not the
497 focus of this study. Eco2 was considered only after further investigation of Eco1 and
498 appearing also in a high rank in this list.
499

In-situ hybridization. Whole open reading frame of *eco1* and *eco2* where cloned into
pGEM-T (*Promega*). DIG-labeled probes where generated using T7/SP6 transcriptions start
sites and the digoxigenin RNA Labeling Mix kit (*Roche*) after manufactures instructions.
Hybridization was performed as described previously (89), in short: animals were relaxed
with 2 % urethan for 2 min at 4 °C, fixed in 4 % PFA over night at 4 °C, washed three times á
10 min with PBT (PBS + 0.1 % Tween), and bleached in methanol (100 %) three times for
minimum 30 min. Methanol was washed out with four decreasing concentrations of ethanol in
PBT (100 %, 75 %, 50 %, 25 % v/v) followed by three times PBT washing steps, all 10 min or
longer on a shaker. Samples were digested with 10 mg ml^{-1} Proteinase K in PBT for 20 min
at room temperature and digestion was stopped by two subsequent washing steps with
4 mg ml^{-1} glycine in PBT, followed by three PBT washing steps each 10 min on a shaker. In
order to acetylate positve charges in the samples and reduce background we treated
samples twice with triethanolamine (0.1 M, pH 7.8) 5 min, once more with triethanolamine
adding 2.5 µl µm^{-1} acetic anhydrid for 5 min to add another 2.5 µl ml^{-1} acetic anhydrid for
another 5 min. The samples where washed three times in PBT, refixed in 4 % PFA over night
at 4 °C and washed another three times in PBT to remove fixative. Samples were treated
twice with 2x SCC (0.3 M NaCl, 30 mM tri-sodium citrate, pH 7) for 10 min at room
temperature and twice for 20 min at 70 °C, to inactivate endogenous phosphatases. To
transfer the samples into hybridization solution (5x SSC + 50 % formamide (v/v), 0.02 %
Ficoll (w/v), 0.02 % bovine serum albumin (w/v), 0.02 % polyvinylpyrolidone (w/v), 0.1 %
Tween20 (v/v), 0.1 % CHAPS (w/v), 100 µg ml^{-1} Heparin), the medium was exchanged to
prewarmed 50 % hybridization solution in 2x SSC for 10 min at 57 °C, followed by 10 min in
hybridization solution at 57 °C and 2 h blocking in hybridization solution + 100 µg ml^{-1} tRNA
at 57 °C. Around 20 ng µl^{-1} *in-situ* probes were denatured for 10 min at 70 °C in 10x SSC +
50 % formamid. The denatured probe was added to the hybridization solution + tRNA to a
final concentration of 2 ng µl^{-1} and incubated over night at 57 °C. Sense probes were used
equally and served as negative control. To remove the probes and prepare antibody staining,
the samples were washed in four steps of decreasing hybridization solution in 2x SSC
(100 %, 75 %, 50 %, 25 %) for 10 min at 57 °C each. Afterwards, samples were washed
twice in 2x SSC containing 0.1 % CHAPS for 30 min at 57 °C and twice in MAB-T (100 mM
maleic acid, 150 mM NaCl, 0.1 % Tween20 (v/v), pH 7.5) for 10 min at room temperature. To
block unspecific antibody binding sites, the samples were blocked with MAB-T + 0.1 %
bovine serum albumin for 1 h at room temperature and with MAB-T + 0.1 % bovine serum
albumin, 20 % heat inactivated sheep serum for 2 h at 4 °C. Anti-digoxigenin alkaline

phosphate linked fab fragments (*Roche*) were used to detect probes by incubating the samples with a 1:2000 solution in MAB-T + 0.1 % bovine serum albumin, 20 % heat inactivated sheep serum over night at 4 °C. Excess antibody solution was removed by 8 washes with MAB-T for 15 min at room temperature. Sample staining was prepared by one NTMT (100 mM NaCl, 100 mM TRIS-HCl, 50 mM MgCl2, 0.1 % Tween20 (v/v), pH 9.5) wash and incubation with NTMT + 1 mM Levamisol for 5 min to inhibit unspecific endogenous alkaline phosphatases. Staining was performed by incubation of the samples in NTMT + 2 % NBT/BCIP in dark until a clear staining was obtained. The reaction was stopped with three washes of a*qua dest.* before dehydration in with three steps increasing ethanol concentrations (50 %, 70 %, 100 % in water) was performed and samples were embedded on glas slides in Euparal. Expression profiles of *eco* genes under the different treatments were analyzed under the microscope.

Immunohistochemistry. Immunostaining was performed using standard procedures (90): animals were relaxed in 2 % urethan for 2 min on ice and fixed in 4 % PFA over night at 4 °C. Fixative was removed with four washes in PBT (PBS + 0.1 % Tween20) for 15 min, cell membranes opened with PBS + 0.5 % TritonX100 for 30 min, and unspecific binding sites blocked with PBT + 1 % bovine serum albumin for 1 h, everything done at room temperature. The primary antibody was a polyclonal rabbit antibody raised against a 14 AA fragment (NSIKENMENFYPVE, AA 52-65, *GeneScript, USA*) of Eco1 and was incubated in PBT + 1 % bovine serum albumin, 10 µg/ml rabbit-α-Eco1-AB over night at 4 °C. The primary antibody was removed with four washes in PBT + 1 % bovine serum albumin for 15 min at room temperature. Secondary anitbody (goat-α-rabbit Alexa Flour 488 coupled, #A11034; LOT 1670152, *Life Technologies*) was incubated at 1 µg/ml in PBT + 1 % bovine serum albumin and incubated for 2 h at room temperature, from this step on samples were kept in dark. Secondary antibody was removed by four washes in PBS + 0.5 % Tween20, 1 % bovine serum albumin for 15 min at room temperature. Actin cytoskeleton staining was performed with rhodamine-phalloidin (#P1951, *Sigma*) in PBT for 1 h at room temperature and afterwards removed with four washes of PBS + 0.5 % Tween20, 1 % bovine serum albumin. To stain the nucleus we incubated the samples with TO-PRO3 iodide AlexaFluor633 (#T3605, LOT 23929W, *Invitrogen*) in PBS + 0.5 % Tween20, 1 % bovine serum albumin for 10 min at room temperature. Afterwards samples were transferred onto glas slides, embedded in Moviol and stored at 4 °C until analysis using a confocal laser scanning microscope (TCS SP1, *Leica*)

Transplantation experiments. Rings of tissue right beneath the tentacle ring were excised from donor animals and grafted into the ~1/3 of the body axis from head to foot of the acceptor animal, using fishing strings and polypropylene tubes for fixation of the tissues.

572   Grafts were grown together after 2-3 h and fishing strings were removed. The induction of
573   heads was assessed 2-3 days post transplantation using the fluorescent markers and a
574   binocular. Every form of head structure was evaluated as head, including a single tentacle, a
575   hypostome or a complete hypostome with an tentacle ring.
576

Author contributions

578   J. Taubenheim, S. Franzenburg, T. C. G. Bosch and S. Fraune. designed the experiments;
579   J. Taubenheim , M. Knop , D. Willoweit-Ohl, S. Franzenburg, and J. He conducted
580   experiments. J. Taubenheim , M. Knop and S. Fraune performed data analysis.
581   J. Taubenheim , T. C. G. Bosch and S. Fraune wrote the paper, all authors corrected final
582   version of the paper.
583

Acknowledgement

585   We greatly appreciate provision and microinjection of the *Hydra* embryos by Jörg Wittlieb.
586   We further thank Alexander Klimovich for help in the design of the antibodies and help with
587   the confocal microscopy. This work was funded by the German Research Foundation (DFG,
588   CRC1182). T. C. G. B. gratefully appreciates support from the Canadian Institute for
589   Advanced Research (CIFAR).

References

1. McGlashan JK, Spencer R-J, Old JM (2012) Embryonic communication in the nest: metabolic responses of reptilian embryos to developmental rates of siblings. *Proc R Soc B Biol Sci* 279(1734):1709–1715.

2. Callier V, Nijhout HF (2011) Control of body size by oxygen supply reveals size-dependent and size-independent mechanisms of molting and metamorphosis. *Proc Natl Acad Sci* 108(35):14664–14669.

3. Zera AJ, Tiebel KC (1988) Brachypterizing effect of group rearing, juvenile hormone III and methoprene in the wing-dimorphic cricket, Gryllus rubens. *J Insect Physiol* 34(6):489–498.

4. Ogawa A, Streit A, Antebi A, Sommer RJ (2009) A Conserved Endocrine Mechanism Controls the Formation of Dauer and Infective Larvae in Nematodes. *Curr Biol* 19(1):67–71.

5. Shikuma NJ, Antoshechkin I, Medeiros JM, Pilhofer M, Newman DK (2016) Stepwise metamorphosis of the tubeworm Hydroides elegans is mediated by a bacterial inducer and MAPK signaling. *Proc Natl Acad Sci* 113(36):10097–10102.

6. West-Eberhard MJ (2003) *Developmental plasticity and evolution* (Oxford University Press) Available at: https://books.google.de/books?id=iBkQyA2PkxEC&dq=Developmental+Plasticity+and+Evolution&hl=de&source=gbs_navlinks_s [Accessed April 25, 2019].

7. Gilbert SF, Bosch TCG, Ledón-Rettig C (2015) Eco-Evo-Devo: Developmental symbiosis and developmental plasticity as evolutionary agents. *Nat Rev Genet* 16(10):611–622.

8. Abouheif E, Favé MJ, Ibarrarán-Viniegra AS, Lesoway MP, Rafiqi AM, Rajakumar R (2014) Eco-Evo-Devo: The time has come. *Adv Exp Med Biol* 781:107–125.

9. Ghosh SM, Testa ND, Shingleton AW (2013) Temperature-size rule is mediated by thermal plasticity of critical size in Drosophila melanogaster. *Proc R Soc B Biol Sci* 280(1760):20130174–20130174.

10. Atkinson D (1994) Temperature and Organism Size—A Biological Law for Ectotherms?, pp 1–58.

11. Gillooly JF, Charnov EL, West GB, Savage VM, Brown JH (2002) Effects of size and temperature on developmental time. *Nature* 417(6884):70–73.

12. Bergmann C (1848) Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse. Available at: https://books.google.de/books?hl=de&lr=&id=EHo-

AAAAcAAJ&oi=fnd&pg=PA3&dq=Bergmann+1848&ots=YkWTtFnaf5&sig=5w
W6LvN-JR7V7g5L_Ewjhh8Fjws [Accessed April 29, 2019].

13.   Li Q, Gong Z (2015) Cold-sensing regulates Drosophila growth through insulin-
producing cells. *Nat Commun* 6:1–11.

14.   Fraune S, Bosch TC (2010) Why bacteria matter in animal development and
evolution. *Bioessays* 32(7):571–580.

15.   Landmann F, Foster JM, Michalski ML, Slatko BE, Sullivan W (2014) Co-
evolution between an Endosymbiont and Its Nematode Host: Wolbachia
Asymmetric Posterior Localization and AP Polarity Establishment. *PLoS Negl*
Trop Dis 8(8):e3096.

16.   Landmann F, Voronin D, Sullivan W, Taylor MJ (2011) Anti-filarial Activity of
Antibiotic Therapy Is Due to Extensive Apoptosis after Wolbachia Depletion
from Filarial Nematodes. *PLoS Pathog* 7(11):e1002351.

17.   Montgomery MK, McFall-Ngai M (1994) Bacterial Symbionts Induce Host
Organ Morphogenesis during Early Postembryonic Development of the Squid
Euprymna Scolopes. *Development* 120(7):1719–1729.

18.   De Vadder F, Grasset E, Holm LM, Karsenty G, Macpherson AJ, Olofsson LE,
Bäckhed F (2018) Gut microbiota regulates maturation of the adult enteric
nervous system via enteric serotonin networks. *Proc Natl Acad Sci U S A*
115(25):6458–6463.

19.   Hedgecock EM, Russell RL (1975) Normal and mutant thermotaxis in the
nematode Caenorhabditis elegans. *Proc Natl Acad Sci* 72(10):4061–4065.

20.   Kammenga JE, Doroszuk A, Riksen JAG, Hazendonk E, Spiridon L, Petrescu
A-J, Tijsterman M, Plasterk RHA, Bakker J (2007) A Caenorhabditis elegans
wild type defies the temperature-size rule owing to a single nucleotide
polymorphism in tra-3. *PLoS Genet* 3(3):e34.

21.   Okamoto N, Nishimura T (2015) Signaling from Glia and Cholinergic Neurons
Controls Nutrient-Dependent Production of an Insulin-like Peptide for
Drosophila Body Growth. *Dev Cell* 35(3):295–310.

22.   Fujiwara M, Sengupta P, McIntire SL (2002) Regulation of body size and
behavioral state of C. elegans by sensory perception and the EGL-4 cGMP-
dependent protein kinase. *Neuron* 36(6):1091–102.

23.   Sawala A, Gould AP (2017) The sex of specific neurons controls female body
growth in Drosophila. *PLOS Biol* 15(10):e2002252.

24.   Pietschke C, Treitz C, Forêt S, Schultze A, Künzel S, Tholey A, Bosch TCG,
Fraune S (2017) Host modification of a bacterial quorum-sensing signal
induces a phenotypic switch in bacterial symbionts. *Proc Natl Acad Sci U S A*
114(40). doi:10.1073/pnas.1706879114.

25. Fraune S, Anton-Erxleben F, Augustin R, Franzenburg S, Knop M, Schröder K,
Willoweit-Ohl D, Bosch TCG (2015) Bacteria-bacteria interactions within the
microbiota of the ancestral metazoan Hydra contribute to fungal resistance.
ISME J 9(7). doi:10.1038/ismej.2014.239.

26. Franzenburg S, Walter J, Künzel S, Wang J, Baines JF, Bosch TCG, Fraune S
(2013) Distinct antimicrobial peptide expression determines host species-
specific bacterial associations. *Proc Natl Acad Sci U S A* 110(39).
doi:10.1073/pnas.1304960110.

27. Mortzfeld BM, Taubenheim J, Fraune S, Klimovich AV, Bosch TCG (2018)
Stem cell transcription factor FoxO controls microbiome resilience in hydra.
Front Microbiol 9(APR). doi:10.3389/fmicb.2018.00629.

28. Fraune S, Bosch TCG (2007) Long-term maintenance of species-specific
bacterial microbiota in the basal metazoan Hydra. *Proc Natl Acad Sci U S A*
104(32):13146–13151.

29. Mortzfeld BM, Taubenheim J, Klimovich A V., Fraune S, Rosenstiel P, Bosch
TCG (2019) Temperature and insulin signaling regulate body size in Hydra by
the Wnt and TGF-beta pathways. *Nat Commun* 10(1):3257.

30. Bosch TCG (2007) Why polyps regenerate and we don't: Towards a cellular
and molecular framework for Hydra regeneration. *Dev Biol* 303(2):421–433.

31. Field KG, Olsen GJ, Lane DJ, Giovannoni SJ, Ghiselin MT, Raff EC, Pace NR,
Raff RA (1988) Molecular Phylogeny of the Animal Kingdom. *Science (80-)*
239(2):748–753.

32. Campbell RD (1967) Tissue dynamics of steady state growth in Hydra littoralis.
Dev Biol 15(5):487–502.

33. Campbell RD (1967) Tissue dynamics of steady state growth inHydra littoralis.
II. Patterns of tissue movement. *J Morphol* 121(1):19–28.

34. Campbell RD (1967) Tissue dynamics of steady state growth inhydra littoralis.
III. Behavior of specific cell types during tissue movements. *J Exp Zool*
164(3):379–391.

35. Holstein TW, Hobmayer E, David CN (1991) Pattern of epithelial cell cycling in
hydra. *Dev Biol* 148(2):602–611.

36. Hobmayer B, Rentzsch F, Kuhn K, Happel CM, von Laue CC, Snyder P,
Rothbächer U, Holstein TW (2000) WNT signalling molecules act in axis
formation in the diploblastic metazoan Hydra. *Nature* 407(6801):186–9.

37. Lengfeld T, Watanabe H, Simakov O, Lindgens D, Gee L, Law L, Schmidt HA,
Özbek S, Bode H, Holstein TW (2009) Multiple Wnts are involved in Hydra
organizer formation and regeneration. *Dev Biol* 330(1):186–199.

38. Gee L, Hartig J, Law L, Wittlieb J, Khalturin K, Bosch TCG, Bode HR (2010)
beta-catenin plays a central role in setting up the head organizer in hydra. *Dev*
Biol 340(1):116–24.

39. MacWilliams HK (1983) Hydra transplantation phenomena and the mechanism
of Hydra head regeneration. *Dev Biol* 96(1):217–238.

40. Browne EN (1909) The production of new hydranths in Hydra by the insertion
of small grafts. *J Exp Zool* 7(1):1–23.

41. Turing A (1952) The chemical basis of morphogenesis. *Philos Trans R Soc*
Lond B Biol Sci 237(641):37–72.

42. Gierer A, Meinhardt H (1972) A theory of biological pattern formation.
Kybernetik 12(1):30–39.

43. Broun M, Gee L, Reinhardt B, Bode HR (2005) Formation of the head
organizer in hydra involves the canonical Wnt pathway. *Development*
132(12):2907–2916.

44. Klimovich A, Rehm A, Wittlieb J, Herbst E-M, Benavente R, Bosch TCG (2018)
Non-senescent Hydra tolerates severe disturbances in the nuclear lamina.
Aging (Albany NY) 10(5):951–972.

45. Bosch TCG, Adamska M, Augustin R, Domazet-Loso T, Foret S, Fraune S,
Funayama N, Grasis J, Hamada M, Hatta M, et al. (2014) How do
environmental factors influence life cycles and development? An experimental
framework for early-diverging metazoans. *Bioessays* 36(12):1185–94.

46. Boehm A-M, Rosenstiel P, Bosch TCG (2013) Stem cells and aging from a
quasi-immortal point of view. *BioEssays* 35(11):994–1003.

47. Galliot B, Buzgariu W, Schenkelaars Q, Wenger Y (2018) Non-developmental
dimensions of adult regeneration in Hydra. *Int J Dev Biol* 62(6-7–8):373–381.

48. Gufler S, Artes B, Bielen H, Krainer I, Eder M-K, Falschlunger J, Bollmann A,
Ostermann T, Valovka T, Hartl M, et al. (2018) β-Catenin acts in a position-
independent regeneration response in the simple eumetazoan Hydra. *Dev Biol*
433(2):310–323.

49. Bisbee JW (1973) Size determination in Hydra: The roles of growth and
budding. *Development* 30(1).

50. Otto JJ, Campbell RD (1977) Tissue economics of hydra: regulation of cell
cycle, animal size and development by controlled feeding rates. *J Cell Sci*
28(1).

51. Leost M, Schultz C, Link A, Wu YZ, Biernat J, Mandelkow EM, Bibb JA, Snyder
GL, Greengard P, Zaharevitz DW, et al. (2000) Paullones are potent inhibitors

of glycogen synthase kinase-3beta and cyclin-dependent kinase 5/p25. *Eur J*
Biochem 267(19):5983–94.

52. Franzenburg S, Fraune S, Kunzel S, Baines JF, Domazet-Loso T, Bosch TCG
(2012) MyD88-deficient Hydra reveal an ancient function of TLR signaling in
sensing bacterial colonizers. *Proc Natl Acad Sci* 109(47):19374–19379.

53. Khalturin K, Hemmrich G, Fraune S, Augustin R, Bosch TCG (2009) More than
just orphans: are taxonomically-restricted genes important in evolution? *Trends*
Genet 25(9):404–413.

54. Tautz D, Domazet-Lošo T (2011) The evolutionary origin of orphan genes. *Nat*
Rev Genet 12(10):692–702.

55. Turing AM (1952) The Chemical Basis of Morphogenesis. *Philos Trans R Soc*
B Biol Sci 237(641):37–72.

56. Meinhardt H, Gierer A (1974) Applications of a theory of biological pattern
formation based on lateral inhibition. *J Cell Sci* 15(2):321–46.

57. Greber MJ, David CN, Holstein TW (1992) A quantitative method for separation
of living Hydra cells. *Roux's Arch Dev Biol* 201(5):296–300.

58. MacWilliams HK (1983) Hydra transplantation phenomena and the mechanism
of Hydra head regeneration. II. Properties of the head activation. *Dev Biol*
96(1):239–57.

59. MacWilliams HK (1983) Hydra transplantation phenomena and the mechanism
of Hydra head regeneration. I. Properties of the head inhibition. *Dev Biol*
96(1):217–238.

60. Takano J, Sugiyama T (1983) Genetic analysis of developmental mechanisms
in hydra. VIII. Head-activation and head-inhibition potentials of a slow-budding
strain (L4). *J Embryol Exp Morphol* 78(2):141–68.

61. Holstein TW (2012) The evolution of the Wnt pathway. *Cold Spring Harb*
Perspect Biol 4(7):a007922.

62. Watanabe H, Schmidt HA, Kuhn A, Höger SK, Kocagöz Y, Laumann-Lipp N,
Özbek S, Holstein TW (2014) Nodal signalling determines biradial asymmetry
in Hydra. *Nature* 515(7525):112–115.

63. Khalturin K, Anton-Erxleben F, Milde S, Plötz C, Wittlieb J, Hemmrich G, Bosch
TCG (2007) Transgenic stem cells in Hydra reveal an early evolutionary origin
for key elements controlling self-renewal and differentiation. *Dev Biol*
309(1):32–44.

64. Hartl M, Glasauer S, Gufler S, Raffeiner A, Puglisi K, Breuker K, Bister K,
Hobmayer B (2019) Differential regulation of myc homologs by Wnt/β-Catenin
signaling in the early metazoan Hydra. *FEBS J* 286(12):2295–2310.

65.   Guder C, Pinho S, Nacak TG, Schmidt HA, Hobmayer B, Niehrs C, Holstein
776   TW (2006) An ancient Wnt-dickkopf antagonism in Hydra. *Development*
777   133(5):901–911.

778   66.   Augustin R, Franke A, Khalturin K, Kiko R, Siebert S, Hemmrich G, Bosch TCG
779   (2006) Dickkopf related genes are components of the positional value gradient
780   in Hydra. *Dev Biol* 296(1):62–70.

781   67.   Vogg MC, Beccari L, Iglesias Ollé L, Rampon C, Vriz S, Perruchoud C, Wenger
782   Y, Galliot B (2019) An evolutionarily-conserved Wnt3/β-catenin/Sp5 feedback
783   loop restricts head organizer activity in Hydra. *Nat Commun* 10(1):312.

784   68.   Lommel M, Strompen J, Hellewell AL, Balasubramanian GP, Christofidou ED,
785   Thomson AR, Boyle AL, Woolfson DN, Puglisi K, Hartl M, et al. (2018) Hydra
786   Mesoglea Proteome Identifies Thrombospondin as a Conserved Component
787   Active in Head Organizer Restriction. *Sci Rep* 8(1):1–18.

788   69.   Reddy PC, Gungi A, Ubhe S, Pradhan SJ, Kolte A, Galande S (2019)
789   Molecular signature of an ancient organizer regulated by Wnt/β-catenin
790   signalling during primary body axis patterning in Hydra. *Commun Biol* 2(1):1–
791   11.

792   70.   Cardenas M, Fabila Y V., Yum S, Cerbon J, Böhmer FD, Wetzker R, Fujisawa
793   T, Bosch TCG, Salgado LM (2000) Selective protein kinase inhibitors block
794   head-specific differentiation in hydra. *Cell Signal* 12(9–10):649–658.

795   71.   Fabila Y, Navarro L, Fujisawa T, Bode HR, Salgado LM (2002) Selective
796   inhibition of protein kinases blocks the formation of a new axis, the beginning of
797   budding, in Hydra. *Mech Dev* 119(2):157–164.

798   72.   Schüler A, Schmitz G, Reft A, Özbek S, Thurm U, Bornberg-Bauer E (2015)
799   The Rise and Fall of TRP-N, an Ancient Family of Mechanogated Ion
800   Channels, in Metazoa. *Genome Biol Evol* 7(6):1713–1727.

801   73.   Malafoglia V, Traversetti L, Del Grosso F, Scalici M, Lauro F, Russo V,
802   Persichini T, Salvemini D, Mollace V, Fini M, et al. (2016) Transient Receptor
803   Potential Melastatin-3 (TRPM3) Mediates Nociceptive-Like Responses in
804   Hydra vulgaris. *PLoS One* 11(3):e0151386.

805   74.   Cheesman SE, Neal JT, Mittge E, Seredick BM, Guillemin K (2011) Epithelial
806   cell proliferation in the developing zebrafish intestine is regulated by the Wnt
807   pathway and microbial signaling via Myd88. *Proc Natl Acad Sci*
808   108(Supplement_1):4570–4577.

809   75.   Franco AT, Israel DA, Washington MK, Krishna U, Fox JG, Rogers AB, Neish
810   AS, Collier-Hyams L, Perez-Perez GI, Hatakeyama M, et al. (2005) Activation
811   of β-catenin by carcinogenic Helicobacter pylori. *Proc Natl Acad Sci U S A*
812   102(30):10646–10651.

76. Murata-Kamiya N, Kurashima Y, Teishikata Y, Yamahashi Y, Saito Y, Higashi
H, Aburatani H, Akiyama T, Peek RM, Azuma T, et al. (2007) Helicobacter
pylori CagA interacts with E-cadherin and deregulates the β-catenin signal that
promotes intestinal transdifferentiation in gastric epithelial cells. *Oncogene*
26(32):4617–4626.

77. Heath-Pagliuso S, Rogers WJ, Tullis K, Seidel SD, Cenijn PH, Brouwer A,
Denison MS (1998) Activation of the Ah Receptor by Tryptophan and
Tryptophan Metabolites [†]. *Biochemistry* 37(33):11508–11515.

78. Kawajiri K, Kobayashi Y, Ohtake F, Ikuta T, Matsushima Y, Mimura J,
Pettersson S, Pollenz RS, Sakaki T, Hirokawa T, et al. (2009) Aryl hydrocarbon
receptor suppresses intestinal carcinogenesis in ApcMin/+ mice with natural
ligands. *Proc Natl Acad Sci* 106(32):13481–13486.

79. Li L, Zheng W, Zhu Y, Ye H, Tang B, Arendsee ZW, Jones D, Li R, Ortiz D,
Zhao X, et al. (2015) QQS orphan gene regulates carbon and nitrogen
partitioning across species via NF-YC interactions. *Proc Natl Acad Sci*
112(47):14734–14739.

80. Li L, Foster CM, Gan Q, Nettleton D, James MG, Myers AM, Wurtele ES (2009)
Identification of the novel protein QQS as a component of the starch metabolic
network in Arabidopsis leaves. *Plant J* 58(3):485–498.

81. Santos ME, Le Bouquin A, Crumière AJJ, Khila A (2017) Taxon-restricted
genes at the origin of a novel trait allowing access to a new environment.
Science (80-) 358(6361):386–390.

82. Khalturin K, Anton-Erxleben F, Sassmann S, Wittlieb J, Hemmrich G, Bosch
TC (2008) A novel gene family controls species-specific morphological traits in
Hydra. *PLoS Biol* 6(11):e278.

83. Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG
(2007) Dynamic evolution of the innate immune system in Drosophila. *Nat
Genet* 39(12):1461–1468.

84. Colbourne JK, Pfrender ME, Gilbert D, Thomas WK, Tucker A, Oakley TH,
Tokishita S, Aerts A, Arnold GJ, Basu MK, et al. (2011) The Ecoresponsive
Genome of Daphnia pulex. *Science (80-)* 331(6017):555–561.

85. Kuo C-H, Kissinger JC (2008) Consistent and contrasting properties of lineage-
specific genes in the apicomplexan parasites Plasmodium and Theileria. *BMC
Evol Biol* 8(1):108.

86. Schwentner M, Bosch TCG (2015) Revisiting the age, evolutionary history and
species level diversity of the genus Hydra (Cnidaria: Hydrozoa). *Mol
Phylogenet Evol* 91:41–55.

87. Hemmrich G, Anokhin B, Zacharias H, Bosch TCG (2007) Molecular
phylogenetics in Hydra, a classical model in evolutionary developmental
biology. *Mol Phylogenet Evol* 44(1):281–290.

88. Pfaffl MW (2001) A new mathematical model for relative quantification in real-
time RT-PCR. *Nucleic Acids Res* 29(9):e45.

89. Grens A, Gee L, Fisher DA, Bode HR (1996) CnNK-2,an NK-2 homeobox
gene, has a role in patterning the basal end of the axis in *Hydra*. *Dev Biol*
180(2):473–488.

90. Engel U, Ozbek S, Streitwolf-Engel R, Petri B, Lottspeich F, Holstein TW
(2002) Nowa, a novel protein with minicollagen Cys-rich domains, is involved in
nematocyst formation in Hydra. *J Cell Sci* 115(Pt 20):3923–3934.

 Supplements

Supplementary Figure 1: The temperature shift does not change the expression of Wnt pathway components. Temperature change has no impact on the expression of Wnt3a, Wnt5, Wnt8, TCF, Frizzled, head formation associated hox genes (brachyury 1 and 2) or the expression of the β-catenin OE construct, but is associated with an 20 to 30 fold up regulation of the eco genes (n =3, Two-way ANOVA, Bonferroni posttests, *** p<0.001).

Supplementary Figure 2: Ectopic tentacle formation after ALP is dependent on the bacterial colonization in *Hydra*. Treatment of germ-free *Hydra* polyps with ALP leads to an increase of ectopic tentacle formation, compared to control and conventionalized animals (recolonized for 4 days prior treatment). The conventionalized and the germ-free animals show some dynamics in tentacle formation, with most tentacles formed 96 h after ALP treatment. Conventionalized *Hydra* represent an intermediate state for tentacle formation after ALP treatment, but is able to mitigate the effect in germ-free animals. n = 15 for conv and GF, 20 for control, fitted generalized linear model with poisson log link function for each time point individually. Multiple comparison for the treatment factor with generalized linear hypothesis testing (R package multcomp) and p-values corrected with Benjamini-Hochberg procedure. Asterisks code for the highest p-value in the comparisons at the different time points: *** p < 0.001, ** p < 0.01. See Supplementary Table 1 for test statistics.

Supplementary_Figure 3: Missing expression signal for *eco2* gene expression after in-situ hybridization. Most of the animals showed no signal after in-situ hybridization with an anti-sense probe (40/45). The low fraction of stained animals is caused by the low *eco2* signal at 18°C rearing temperature.

Supplementary Figure 4: *eco* expression does not influence the regeneration processes in Hydra. (A) To test for a role of eco genes in pattern formation during regeneration, animals were cut in half and each piece was placed individually in a 24-well plate. The regenerating pieces were observed for 3-4 days, counting emerging tentacles and foots every day. (B) The timing of the head regeneration (first appearance of a tentacle) was not altered by the knock-down of the eco genes in 12°C as well as 18°C (Wilcoxon-test, with Benjamini-Hochberg correction, n = 11, 12, 13, 9, 13, 10 polyps for each group, following the order of the graphic left to right). (C) Number of tentacles at different time points of regeneration was indifferent between Eco-KD and control animals (t-test @dpa 3 and 4, with Benjamini-Hochberg correction, n=6-14 polyps for each group). (D) The fraction of polyps regenerating a foot after ablation was not changed in an Eco-KD background (Fisher's-exact test @dpa =3 (12°C) or 2 (18°C), n = 6-14 polyps for each group).

ALP effect on eco genes

Figure legends

Figure 1: Wnt-signaling is temperature dependent. (A) Animals reared at 12 °C and 18 °C
were treated with ALP for 24 h, before assessment of ectopic tentacle formation after 96 h.
(B) Lower temperature leads to fewer ectopic tentacle formation after ALP treatment. (t-test,
n =12, * p<0.05) (C-F) Constitutive active Wnt signaling in β-catenin OE animals causes
multiple heads/body axes at 18 °C, while the severity of the phenotype was subdued when
animals were reared at 12 °C. (G) The number of heads produced by the β-catenin OE
animals is reversible and depends on the rearing temperature, where surrounding
temperatures of 12 °C resulted in fewer heads per polyp (n = 10).

Figure 2: Wnt-signaling is dependent on microbial colonization. (A) Animals were ALP
treated for 24 h and number of generated ectopic tentacles were assessed 96 h after
treatment, comparing the treatment outcome of germfree (GF) and normally colonized
animals (control). (b-d) Ectopic tentacle formation is increased when colonizing bacteria were
removed, suggesting a role of microbial colonization in pattern formation in *Hydra* (Mann
Whitney U test, n =58, *** p<0.0001).

Figure 3: Two TRGs, expressed in the ectoderm, are potential recognition signals for
environmental changes. (a) Re-analysis of microarray data, comparing germfree animals
with colonized ones, and transcriptomic data, comparing animals at different rearing
temperatures, revealed two candidate genes with common regulation upon environmental
changes: eco1 and eco2. (b) Sequence analysis of the candidate genes revealed a
paralogous relationship between the two genes and a secretion signal peptide, without any
domain structure. (c-d) qRT-PCR assessment of eco gene expression in germfree animals
and upon temperature shift confirms down regulation of eco1 and eco2 expression at higher
rearing temperatures (n =3, Two-way ANOVA, Bonferroni posttests, ** p<0.01, *** p<0.001)
and in disturbed microbiome conditions (n =3, Two-way ANOVA, Bonferroni posttests, *
p<0.05, ** p<0.01, *** p<0.001). (e-h) eco genes were expressed in the foot and lower third
of the animals at 18 °C rearing temperature, but the expression domain expands to the body
column and parts of the head at 12 °C. (i-j) Immunohistochemistry with a polyclonal antibody

raised against a fragment of Eco1 (underlined in b) show the production of the peptides in
the ectodermal epithelium and packaging in vesicles localized in the apical part of the cells,
which suggests a secretion of the peptide.
902
Figure 4: Expression dynamics of eco genes were a long term adaptation to changing
microbial state or rearing temperatures. (A) eco gene expression was reestablished by
recolonization with bacteria after 14 d, but did not reach normal levels after only 8 d
recolonization (n =3, Two-way ANOVA, Bonferroni posttests, * $p<0.05$, ** $p<0.01$, ***
$p<0.001$) (B) Recolonization with the main colonizer Curvibacter sp. alone was sufficient to
alleviate the expression suppression of eco genes after 14 d of colonization. (n =4, Two-way
ANOVA, Bonferroni posttests, * $p<0.05$, *** $p<0.001$) (C) eco gene expression changes were
established within 14-16 d upon rearing temperature shifts from 18 °C to 12 °C and vice
versa. (D) Microbial colonization state and rearing temperature were independent inputs for
the expression regulation of eco genes. Gene expression is up regulated upon temperature
shifts from 18 °C to 12 °C, independent of colonization state of the animals. (E, F) The
graphic displays effect size cleaned of overlaying variances. (E) The expression of *eco*
genes increases within 14 days after reduction of rearing temperature, independent of the
colonization status of the genes (ANOVA, $p = 8.16E-9$ for temperature, $p = 6.99E-6$ for
temperature-time interaction, see Supplementary Table 3). (F) Removing bacteria from
Hydra reduces expression of *eco* genes independent of temperature (ANOVA, $p = 0.0557$ for
colonization, $p = 0.20$ for colonization-temperature interaction, see Supplementary Table 3).
920
Figure 5: Knockdown of eco genes results in increased Wnt signaling. (a) Eco–hairpin
construct for generation of transgenic *Hydra* (as, antisense; s, sense; TAA, stop codon; P,
promoter; T, terminator) (b, c) Whole mount in situ hybridization of eco1 expression in Eco-
KD (b) and control (c) animals. (d) eco1 and eco2 expression in two transgenic lines (B8 and
B5) at 12 °C and 18 °C (n =3, Two-way ANOVA, Bonferroni posttests, *** $p<0.001$) (e-f)
Treating Eco1-KD animals with ALP revealed an increased potential to form ectopic
tentacles, indicating a higher Wnt signaling activity. (f) The number of heads formed after
transplantation of near head tissue into the body axis of an acceptor polyp serves as a
readout for the head inhibition potential, which is governed by the inhibition of the Wnt
signaling. (g-h) Control (unlabeled) and Eco-KD (GFP labeled) animals served as acceptor
polyps for head tissues from wild type (dsRed labeled) animals to assess the head inhibition
potential under disturbed eco expression. (i) Eco-KD polyps showed a reduced head
inhibition potential, indicating an impaired Wnt inhibition in these animals. n = 75, $p=0.0085$,
Fisher's exact test.
935
Figure 6: Eco peptides are environmentally triggered inhibitors of Wnt signaling.
Microbial colonization and low temperature are independent inputs for Eco peptides and are
able to induce its transcription. Eco peptides are synthesized as small peptides, packed into
vesicles and secreted into extracellular matrix of the ectoderm. By an unknown mechanism
Eco peptides are recognized and the Wnt signaling is inhibited. The inhibition takes place at
the level of β-catenin stabilization or the expression of Wnt target genes as Eco expression
were able to inhibit activation of the pathway at the GSK-3β level.
943
944 **Supplementary Figure 1: The temperature shift does not change the expression of Wnt**
945 **pathway components.** Temperature change has no impact on the expression of Wnt3a,
946 Wnt5, Wnt8, TCF, Frizzled, head formation associated hox genes (brachyury 1 and 2) or the

947 expression of the β-catenin OE construct, but is associated with an 20 to 30 fold up
948 regulation of the eco genes (n =3, Two-way ANOVA, Bonferroni posttests, *** p<0.001).
949
Supplementary Figure 2: Ectopic tentacle formation after ALP is dependent on the
bacterial colonization in *Hydra*. Treatment of germ-free *Hydra* polyps with ALP leads to an
increase of ectopic tentacle formation, compared to control and conventionalized animals
(recolonized for 4 days prior treatment). The conventionalized and the germ-free animals
show some dynamics in tentacle formation, with most tentacles formed 96 h after ALP
treatment. Conventionalized *Hydra* represent an intermediate state for tentacle formation
after ALP treatment, but is able to mitigate the effect in germ-free animals. n = 15 for conv
and GF, 20 for control, fitted generalized linear model with poisson log link function for each
time point individually. Multiple comparison for treatment factor with generalized linear
hypothesis testing (R package multcomp) and p-values corrected with Benjamini-Hochberg
procedure. Asterisks code f or the highest p-value in the comparisons at the different
timepoints: *** p < 0.001, ** p < 0.01. See Supplementary Table 1 for test statistics.
962
963 **Supplementary Figure 3: Missing expression signal for *eco2* gene expression after in-**
964 **situ hybridization.** Most of the animals showed no signal after in-situ hybridization with an
965 anti-sense probe. The low fraction of stained animals is caused by the low *eco2* signal at
966 18°C rearing temperature.
967
Supplementary Figure 4: Eco peptides are not involved in the regeneration process of
Hydra. (A) To test for a role of *eco* genes in pattern formation during regeneration, animals
were, randomized, cut in half and each piece was placed individually in a 24-well plate. The
regenerating pieces were observed for 3-4 days, counting emerging tentacles and foots
every day. (B) The timing of the head regeneration (first appearance of a tentacle) was not
altered by the knock-down of the *eco* genes in 12°C as well as 18°C (Wilcoxon-test, with
Benjamini-Hochberg correction, n = 9-13 polyps for each group). (C) Number of tentacles at
different time points of regeneration was indifferent between Eco-KD and control animals (t-
test @dpa 3 and 4, with Benjamini-Hochberg correction, n=6-14 polyps for each group). (D)
The fraction of polyps regenerating a foot after ablation was not changed in an Eco-KD
background (Fisher's-exact test @dpa =3 (12°C) or 2 (18°C), n = 6-14 polyps for each
group).
980
981 **Supplementary Figure 5: ALP treatment does not result in expression changes of eco**
982 **genes.** Animals treated with different concentrations of ALP for 24 h showed now significant
983 difference in the gene expression for eco genes (ANOVA, gene-stratified p = 0.532, see
984 Supplementary Table 4).