Visual working memory representations in visual and parietal cortex do

not remap after eye movements

Tao He[1*], Matthias Ekman[1], Annelinde Vandenbroucke[1,2] and Floris P. de Lange[1]

[1] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Kapittelweg 29, 6525 EN Nijmegen, the
Netherlands
[2] Current Address: Dutch national research agenda, Laan van Nieuw Oost Indië 300, 2593 CE Den Haag, the Netherlands
[*] Correspondence: t.he@donders.ru.nl

Abbreviated title: No remapping of VWM after eye movements

Number of pages: 30

Number of figures: 3

Number of words: Abstract = 248, Sig. Statement = 102; Introduction = 549, Discussion = 1008

Acknowledgements: This work was supported by The Netherlands Organisation for Scientific

Research Vidi Grant 452-13-016 to F.P.d.L., the EC Horizon 2020 Program ERC Starting Grant

678286 "Contextvision" to F.P.d.L, the James S.McDonnell Foundation 220020373 to F.P.d.L

and the China Scholarship Council (CSC; 201608330264) to T.H.. We thank David Richter and

Chuyao Yan for helpful comments and discussions of the manuscript.

Competing Interest: The authors declare no competing financial interest.

Author contributions: T.H. wrote the first draft of the paper; T.H., M.E., and F.P.d.L. edited the

paper; T.H., M.E., A.V., and F.P.d.L. designed research; T.H. and A.V. performed research; T.H.,

M.E., and F.P.d.L. analyzed data.

ABSTRACT

Previous studies have suggested that stimulus representations in the visual system

are remapped across eye movements during perception. Interestingly, our visual

system does not only process stimuli that are directly available to our eyes, but may

also have a role in maintaining information in VWM over a period of seconds. It

remains unclear if saccadic remapping also occurs for stimuli that are held in working

memory and whether eye movements degrade VWM representations in visual

cortex. Here, we directly compared the content of VWM in early visual areas for

saccade and no-saccade conditions using MVPA of delay-related activity measured

with fMRI. Specifically, we probed item-related VWM representations of the

remembered grating orientation from contralateral and ipsilateral early visual areas

(V1 – V3) and the intraparietal sulcus (IPS). When participants did not make a

saccade, VWM representations were robustly present in contralateral early visual

cortex. Surprisingly, VWM representations disappeared from contralateral V1-V3

after the saccade shifted the location of the remembered grating to the opposite

visual field. In contrast to our saccadic remapping hypothesis, we found no evidence

for the representations of the remembered grating at the saccadic target location in

the opposite visual field, suggesting that there is no evidence for remapping of VWM

in early visual cortex. Interestingly, IPS showed persistent VWM representations in

both the saccade and no-saccade condition. Together, our results indicate that VWM

representations in early visual cortex are not remapped across eye movements,

potentially limiting the role of early visual cortex in VWM storage.

Significance Statement

It has been well documented that our visual system does not only process stimuli

that are directly available to our eyes, but may also have a role in maintaining

information in visual working memory over a few seconds. Previous research has

indicated that stimulus features are remapped across eye movement to aid visual

stability. In the current study, we show that visual working memory information does

not remap after eye movements. In addition, visual working memory representations

are degraded after eye movements in early visual cortex, but not in parietal cortex,

potentially limiting the role of early visual cortex in VWM storage.

Introduction

Humans perform about three saccadic eye movements per second in order to guide the fovea toward regions of interest in the visual field (Rayner, 1998). Previous research has indicated that both stimulus features and attentional pointers are remapped across eye movement to aid visual stability (Melcher, 2007; Rolfs et al., 2011; He et al., 2018). Furthermore, research in the last decade has suggested that our visual system may not only process incoming information, but also be relevant for maintaining internal representations of previously observed visual stimuli, i.e. visual working memory (VWM). The ability to maintain information that is no longer in view is critical for reasoning about and mentally manipulating visual information despite temporal discontinuous in visual inputs that occur for example during eye movements, occlusions, and object motion (Curtis and D'Esposito, 2003; Serences, 2016; Lorenc et al., 2018).

VWM information has been observed both in early sensory areas (Ester et al., 2009; Harrison and Tong, 2009; Albers et al., 2013), as well as higher-order parietal (Christophel et al., 2015; Ester et al., 2015; Bettencourt and Xu, 2016; Xu, 2017, 2018), and prefrontal regions (Durstewitz et al., 2000; Riley and Constantinidis, 2016). Moreover, VWM representations in early visual cortex have been found to be spatially specific, maintained in a retinotopic manner (Pratte and Tong, 2014). Crucially however, in these previous studies, participants were instructed to maintain fixation while remembering a visual stimulus during a retention period. While this is a common approach to investigate VWM in a lab environment, it is arguably quite different from real-world settings that are marked by multiple eye movements per second. In our daily life, VWM is typically dynamic and fragile. For instance, when looking for a key in an apartment, one needs to maintain a mental image of the key

while continuously moving one's eyes toward different search locations. It is largely

unknown how eye movements impact the mental VWM representations in early

visual areas and beyond.

One possibility is that VWM representations are dynamically remapped and

follow the shift of the retinotopic location caused by eye movements. In this case, the

cortical location representing the VWM would be updated after an eye-movement.

Alternatively, working memory information might be maintained across multiple brain

regions and not require remapping following eye movements. On this account, after

every eye movement, the up-to-date feature of the stimulus (e.g., the latest location

of the stimuli) would be integrated with the previous working memory information.

Here, we tested the potential remapping of VWM in the early visual system,

by presenting participants with an orientated grating in either the left or right visual

field. The grating orientation had to be maintained for a subsequent retention period,

while we measured blood oxygenation level-dependent (BOLD) signals with

functional magnetic resonance imaging (fMRI). Using multivariate pattern analysis

(MVPA), we attempted to decode the remembered orientation in both the

contralateral and the ipsilateral hemispheres during trials in which participants

performed either a saccade or maintained fixation during the working memory

period.

To preview, we found a contralateral VWM representation in the early visual

system. This VWM representation disappeared when participants made a saccade

and was not remapped to ipsilateral visual cortex. In contrast, VWM representations

in IPS persisted after saccades. These findings suggest a limited role of the early

visual system in VWM.

Materials and Methods

Data and code availability

All data and code used for stimulus presentation and analysis is freely available on

the Donders Repository (https://data.donders.ru.nl/login/reviewer-

77603154/wWWKvrhkeJS0w6mNBkWAHBvST_jPJyodeGVyxxYwpLc).

Participants

Thirty healthy participants with normal or corrected-to-normal vision were recruited

from the institute's subject pool in exchange for either monetary compensation or

study credits. All participants were naive with respect to the purposes of the study.

The experiments were approved by the Radboud University Institutional Review

Board and were carried out in accordance with the guidelines expressed in the

Declaration of Helsinki. Written informed consent was obtained from all participants

prior to the start of the study. All participants had to complete the entire experimental

protocol (i.e., the behavioral experiment and the two fMRI sessions). Six participants

were excluded before the final analysis: one participant didn't complete the fMRI

sessions due to discomfort, one participant failed to follow the instructions (i.e., fixate

and saccade) during experiment, four participants had excessive head motion

(motion cutoff = 1 mm). The remaining 24 participants were included in all analyses

(13 female, mean age 23.3 years, range from 19 to 31 years). The desired sample

size of N=24 included subjects was based on an a priori power calculation that

aimed to achieve 80% power to detect an effect size of at least Cohen's d > 0.6.

Stimuli

Stimuli were programmed in MATLAB (v2016a, The MathWorks, Inc., Natick, MA) using the Psychtoolbox (Brainard, 1997). The circular sinusoidal grating stimulus subtended 10° and was centered with a small jitter (0.3°) on the screen center. The grating was full contrast, with a spatial frequency of 1 cycles/degree, a random phase and an orientation of either 25° or 115° (with a small jitter of 3°) from the horizontal axis. The contrasts of the edges of the grating were linearly attenuated over the distance from 4.5° to 5.0° radius. Two filled dots (0.5°, one green, one black) were presented at the periphery of the screen, which were served as the fixation dot and saccade target in the task. In the behavioral training session, stimuli were presented on a 24-in. flat-panel display (BenQ XL2420T, 1,920 x 1,080 resolution, 60 Hz refresh rate). In the fMRI sessions, stimuli were displayed on a rear-projection screen using an EIKI LC-XL100L (EIKI, Rancho Santa Margarita, CA) multimedia projector (1,024 x 768 resolution, 60 Hz refresh rate).

Experimental design

Behavioral training procedure. Prior to the fMRI scan sessions, all participants completed a one-hour behavioral training session to familiarize themselves with the fMRI main task and to establish their individual orientation discrimination threshold, which served as an initial orientation difference of the gratings in the following fMRI sessions. The experimental design of the behavioral training task was exactly the same as during the fMRI main task, except that the delay period and the inter-trial interval (ITI) were shortened to 3 s to reduce the experimental time. Participants completed two to three blocks of 56 trials until a stable orientation discrimination threshold was obtained, during which the eye movements were continuously monitored by an Eyelink 1000 plus eye tracker. In addition, participants were

familiarized with the localizer tasks and the retinotopic mapping procedures that

were used in the fMRI session at the end of the behavior session.

fMRI main task.   The experimental design (Figure 1a) was adapted from a well-

known delayed orientation discrimination task used before (Harrison and Tong,

2009). Each trial began with two filled dots (one green, one black) that were

presented at the periphery of the screen (2 s). Participants were asked to fixate at

the green dot through the trial. Two oriented gratings were flashed sequentially at the

center of the screen (which could be in the left or right visual field of the participants,

depending on the location of the green fixation dot) for 200 ms, respectively, with an

inter-stimulus interval of 400 ms, and followed by a retro-cue (400 ms, "1" or "2") to

indicate which orientation of the grating should be remembered during the following

delay period. The sequence of the orientations (25° and 115°) were randomly

chosen on each trial. On one third of trials (No-saccade condition), the two dots did

not swap their locations, which indicated to participants that they needed to keep

their fixation at the initial fixation location during the 10 s delay period. On the

remaining two thirds of trials (Saccade condition), the green dot was shifted to the

opposite side of the screen at the beginning of the delay period (2 s after the offset of

the retro-cue). This instructed participants to move their eyes to the opposite location

of the screen and maintain their fixation at the new position for the remaining 8 s. A

probe grating was presented at the screen center after the delay period and

participants were required to make a judgement of whether the probe orientation was

rotated clockwise or countclockwise relative to the grating they memorized during the

delay period. Finally, feedback was provided and a black fixation dot instructed

participants to move their eyes to the center of the screen in preparation for the next

trial. Trials were separated by an ITI of 9.6 s. The central fixation dot changed its

color from black to gray at the last second of the ITI to indicate that the next trial was

going to start.

In order to rule out the potential confound of global representations of VWM,

four catch trials were included in each run. Stimuli and timing of the catch trials were

identical to the main trials, except for the following changes. During the presentation

of the probe in the catch trial, the probe was horizontally shifted 1° to the left or right

with respect to the location of the sample gratings. Participants were instructed to not

respond when the location of the probe grating did not match with the sample

gratings. The rationale of this was that it forced participants to memorize the

orientation stimulus at its original location in order to successfully perform the task,

resulting in a spatially specific VWM representation.

Each run comprised 28 trials, consisting of 8 no-saccade trials, 16 saccade

trials, and 4 catch trials. The order of trials was a pseudo-randomized within each

run. Each trial lasted 26.4 s, consisting of a 16.8 s task period and a 9.6 s ITI. Each

run lasted 12.32 min and started with 6 s of fixation that was discarded from the

analysis.

Staircase Procedure

The staircase procedure was used to ensure equal task difficulty across participants

and to equate task difficulty levels across no-saccade and saccade conditions within

participants. In the behavioral training session, immediately after the delay period,

participants were asked to compare the tilted orientation between the probe and the

internal remembered grating. The staircase procedure estimated the difference

between probe and remembered grating that ensured 75% performance, using

QUEST (Prins and Kingdom, 2018). A maximum orientation difference of 20°

between the probe and remembered grating was enforced. The staircase was

initiated with an orientation difference of 10° and dynamically adapted according to

participants' performance on previous trial until a stable threshold was acquired. This

threshold was used as a seed in the following fMRI sessions and the same staircase

procedure was also used during scanning.

Eye tracking

Eye position was monitored with an MR-compatible Eyelink 1000 (SR Research Ltd.,

Ottawa, Canada) eye tracker. Only the left eye was recorded in the scanner. Pupil

and corneal reflection were sampled at 1000 Hz and analyzed offline to ensure that

participants fixated at the correct location. The eye tracker was calibrated at the

beginning of each session and repeated between runs if necessary. In 10 out of 192

runs, the eye tracker signal was lost during scanning due to subjects' head motion or

technical problems. During these runs, the experimenter monitored the eye position

online via the live video feed from the camera. All participants were trained on the

fixation task and to perform saccades in the behavioral training session prior to the

scanning session.

fMRI parameters

Functional and anatomical images were carried out with a 3T Siemens Prisma fit

MRI system (Siemens, Erlangen, Germany), using a 32-channel headcoil. Functional

images were acquired using a whole brain T2*-weighted multiband-4 sequence

(TR/TE = 1,200/39 ms, voxel size 2.4x2.4x2.4 mm, 56 transversal slices, 65° flip

angle, A/P phase encoding direction, FOV = 210 mm, BW = 2030 Hz/Px).

Anatomical images were acquired using a T1-weighted magnetization prepared rapid

gradient echo (MP-RAGE) sequence (TR/TE = 2,300/3.03ms, voxel size 1x1x1 mm,

192 transversal slices, 8° flip angle).

fMRI Preprocessing

fMRI data were preprocessed using FSL (Oxford, UK) (Smith et al., 2004), including

motion correction (six-parameter affine transform), temporal high-pass filtering (100

s) and Savitzky–Golay low-pass filter (time window length = 11 TRs, polynomial

order = 3; Savitzky & Golay, 1964) for each run separately. No spatial smoothing

and slice timing correction was performed. All univariate and multivariate analyses

were performed in native subject space using custom python code.

Functional localizers

In addition to the main experiment, participants underwent two localizer runs, which

were used to select voxels that maximally responded to stimuli presented in the

contralateral hemifield and served as independent training data for the multivoxel

pattern analysis. The same grating stimulus parameter were used in localizer runs as

those in the main experiment. Participants fixated at the left or right green dot and

the grating was presented at the center of screen for 16 s, with a frequency of 4 Hz.

Throughout the localizer, participant had to fixate at the green dot and monitor a

sequence of rapidly changing letters just above fixation, to which they had to

respond by button press whenever a target letter ("X" or "Z") occurred in a stream of

non-target letters ("A", "H", "R", "N", "T", "V", "U", "Y"). Letters were presented at a

frequency of 2 Hz. When the green dot shifted from the left to the right side of the

screen (or vice versa) at the end of the trial, participants had to move their eyes to

follow the fixation dot. Each trial was separated by a 2 s ITI in which only the fixation

cross was presented, to give participants enough time to saccade and stabilize their

eyes in the new position. Two blocks of 20 trials (10 trials per visual field: left or right)

and orientation (25° or 115°) were collected.

Images of the localizer runs were also preprocessed following the same

procedure as the main tasks. Onsets and duration of the localizer blocks were

convolved with a single-gamma haemodynamic response function (HRF) and fitted

using a general linear model in left and right hemisphere separately. For each

participant a two-sided t-contrast was calculated contrasting between left and right

hemispheres. Resulting statistical maps were thresholded at $Z > 2.3$ and a corrected

cluster significance threshold of $p < 0.05$.

Retinotopic mapping of early visual areas (V1 – V3)

Early visual areas (V1, V2, and V3) for each participant were identified using two

retinotopic mapping blocks based on the standard traveling-wave method using

rotating wedges (DeYoe et al., 1996; Engel et al., 1997; Wandell et al., 2007),

consisting of a clockwise and a counterclockwise-rotating run. Participants were

instructed to fixate at the center of the screen and fulfill rapid letter detection task

(exact same as that in localizer runs). The BOLD responses to the wedges were

used to estimate the polar angle of the visual field representation.

Regions of interest (ROIs)

Early visual areas (V1 - V3) and the intraparietal sulcus (IPS0 – IPS5) within each

hemisphere were independently chosen as the ROIs in our analyses. For the ROI of

the early visual areas, we first used Freesurfer (http://surfer.nmr.mgh.harvard.edu/,

(Fischl et al., 2002)) to define the gray–white matter boundary and perform cortical

surface reconstruction. The borders of the early visual areas V1 - V3 were delineated

based on retinotopic maps. The ROI of the intraparietal sulcus was defined based on

anatomical probability maps of retinotopic areas in the intraparietal sulcus (IPS0 –

IPS5) from the Probabilistic atlases (Wang et al., 2015). Finally, all surface-based

ROIs were backward-transformed into participant's native space. Next, the visually

active voxels corresponding to the grating that was positioned at the left and right

visual field was identified based on statistical activation maps from the functional

localizer runs. Two types of ROI were defined in the decoding analyses: contralateral

ROIs and ipsilateral ROIs. For instance, when decoding the orientation of the grating

that was initially presented at participant's right visual field, regardless of the no-

saccade and saccade condition, the left hemisphere would be labeled as the

contralateral ROIs, while the right hemisphere would be labeled as the ipsilateral

ROIs (see Figure 1b) and vice versa. Finally, we selected the 200 most active voxels

across the entire early visual cortex (V1 -V3) and the entire intraparietal sulcus (IPS0

– IPS5) to do the decoding analyses, separately.

Univariate fMRI analyses

To estimate the BOLD response of each hemisphere in each condition, we selected

the same voxels as those used for decoding. We separately modeled the onset of

each trial for each hemisphere and for no-saccade and saccade condition

separately. To quantify BOLD activity during the trial, contrasts between left and right

hemisphere regressors for each condition were created. All fMRI data were

transformed from MRI signal intensity to units of percent signal change, calculated

relative to the average level of activity for each voxel across the first volume of each

trial within each run.

Multivariate fMRI analyses

We used multivariate pattern analyses (MVPA) to determine whether the pattern of

activity in each ROI and each hemisphere contained orientation information, as

implemented in Scikit-learn 0.20.3 (Pedregosa et al., 2011). To this end, linear

support vector machines (SVMs) were trained to discriminate between the two

grating orientations based on the pattern of BOLD activity over voxels. In this study

we used classification distance as an indication of the amount of orientation

information being maintained in each hemisphere. To calculate the classification

distance, we measured the distance of each sample to the separating hyperplane

that was trained by a linear SVM with a positive/negative sign (the sign indicates the

class). We then averaged the distance within each class and calculated the absolute

distance between these two classes. While this approach can give us a binary

predicted label (25° or 115°), which can be used to calculate classification accuracy,

it additionally allows one to review the confidence of this classification and provide a

continuous metric of the decoding performance. In each ROI, the larger the

classification distance, the more confident the classifier is in determining the stimulus

class based on the BOLD activity pattern, and hence the more orientation

information is contained within the pattern of BOLD activity.

When training and testing within the main experiment we averaged the BOLD activity

over time points 7.2 - 10.8 s after the onset of the delay period. This time period was

selected to be as far from the onset of the two gratings at the beginning of the trial as

possible, and therefore not reflect activity elicited by the stimuli, but also prior to the

onset of the probe stimulus. These time series were then normalized using a z-

scored transformation within each run. For generalization from perception to VWM,

the classifier was trained using a full time series (0 - 16 s) of the average BOLD

activity in the localizer and then applied to the averaged VWM period.

A leave-one-run-out cross-validation procedure was used to train the classifier where

we trained in 7 out of 8 runs and tested on the remaining run. We also used this

cross-validation procedure on the time-resolved decoding analysis, but within each

time point independently. BOLD activity and classification distance over time were

statistically tested using nonparametric cluster-based permutation t test (10,000

permutations; two-sided $p < 0.05$; cluster formation threshold $p < 0.05$) (Maris and

Oostenveld, 2007).

Bayesian analyses

In order to further evaluate non-significant tests, we performed the Bayesian

equivalents of the above outlined analyses. JASP 0.10.2 (JASP Team, 2019,

RRID:SCR_015823) was used to perform all Bayesian analyses, using default

settings. Thus, for Bayesian t-tests a Cauchy prior width of 0.707 was chosen.

Qualitative interpretations of Bayes Factors are based on criteria by Lee &

Wagenmakers (2014).

341
Results

Our primary goal was to investigate whether VWM representations in early visual

cortex (V1-V3) are remapped after eye movements. We used MVPA (Kamitani and

Tong, 2005; Haynes and Rees, 2006; Norman et al., 2006) to determine whether the

contents of VWM that was initially represented in the contralateral hemisphere

(corresponding to the stimulus location) was remapped to the opposite (ipsilateral)

hemisphere following an eye movement to the opposite visual field during the

retention period. Twenty-four participants performed a delayed orientation

discrimination task, in which they initially fixated at the peripheral fixation dot while

passively viewing two successive grating stimulus orientations (25° and 115°) in one

visual field, followed by a cue and a delay period. Importantly, in two third of the trials

(*saccade condition*), participants were asked to saccade to the opposite side of the

screen during the delay period, while in the remaining third of trials (*no-saccade*

condition), participants maintained fixation at the initial side through the entire trial. A

probe grating was presented at the end of the trial (Figure 1a, b).

In order to ensure that participants memorized the grating in its original

location and thereby rule out of the potential confound of spatially unspecific

representations of VWM in early visual cortex (Ester et al., 2009; Pratte and Tong,

2014), we also included catch trials in which the probe grating was horizontally

shifted 1.2° with respect to the location of the sample gratings. Participants were

instructed to discriminate the orientation between the probe and the memorized

grating only when they were presented at the same location (see Methods for

details). This experimental design was chosen to encourage participants to maintain

both the orientation and location of the presented grating. Indeed, behavioral data

showed that participants could successfully withhold their response at an accuracy of

66.67% (SD 12.25) when the probe was presented at a displaced location,

demonstrating that participants remembered the orientation in a spatially specific

way.

Behavioural data confirmed that participants could successfully discriminate

small differences between the cued grating and the probe grating (mean

discrimination threshold, no-saccade condition: 8.17°, SD 4.80; saccade condition:

8.69°, SD 4.61) at an accuracy of 81.7% (SD 4.96) and 81.4% (SD 4.90) in both no-

saccade and saccade trials. Notably, there was no difference in accuracy between

trials with and without a saccade ($t_{(23)} = 0.33$, $p = 0.7465$, Cohen's $d = 0.07$; $BF_{01} =$

4.426), and no differences in the mean discrimination threshold ($t_{(23)} = 1.04$, $p =$

0.3085, Cohen's $d = 0.21$; $BF_{01} = 2.866$), indicating that the VWM performance was

not impaired by the eye movement during the delay period.

A univariate BOLD analysis showed that the presentation of the grating stimuli

induced stronger BOLD activity in contralateral than ipsilateral early visual areas

(cluster permutation test, $p = 0.0001$). As expected, probe presentation at the end of

the trial led to a BOLD activity increase in the contralateral hemisphere during no-

saccade trials and a BOLD activity increase in the ipsilateral hemisphere,

corresponding to the updated stimulus location after the eye movement, in the

saccade condition (Figure 1c). Taken together, these results suggest that

participants engaged successfully in the task.

Figure 1. Experimental paradigm and BOLD activity in the early visual cortex. (a)
Participants performed a delayed orientation discrimination task. At the start of each trial, two
dots were presented at the periphery of the screen and participants were instructed to fixate
at the green dot. Two orientated gratings (25° and 115°) were successively flashed in the
center of the screen, followed by a retro-cue (1 or 2) that indicated which grating to remember
in the following delay period. Crucially, during a 10 s delay period, if the green dot shifted to
the opposite side of the screen (*saccade condition*), participants had to make an eye
movement to the opposite side and maintained at the new location through the following time
period (the dashed arrow is for illustration only, not present in the actual test). Conversely, in
the no-saccade condition, the green dot did not change its position and participants
maintained fixation at the initial side throughout the entire trial. After the delay period a probe
was presented, and participants indicated whether the probe was tilted clockwise or

counterclockwise relative to the remembered grating. **(b)** Illustration of the experimental

design probing remapping. In the no-saccade condition (left), the remembered grating

presented in right visual field is initially represented in the left (contralateral) early visual cortex.

In the saccade condition (right), after participants saccade from the left to the right side of the

screen, we set out to examine whether VWM contents that initially generated in contralateral

(left) early visual cortex transfer to the ipsilateral (right) visual cortex. **(c)** Group averaged

(N=24) BOLD time course in contralateral and ipsilateral early visual areas (V1 – V3). In the

no-saccade condition, the presentation of grating stimuli evoked a higher BOLD response in

contralateral than in ipsilateral early visual areas. In the saccade condition, while the BOLD

activity was still higher in contralateral relative to ipsilateral early visual areas after

presentation of the grating stimuli, the pattern inverted after presentation of the probe – BOLD

activity became higher in ipsilateral than in contralateral early visual areas. Shaded areas

denote SEM.

413
In order to evaluate the potential remapping of VWM in early visual cortex, we

assessed whether the patterns of activation in both the hemispheres, contralateral

and ipsilateral to the item location, contained the remembered orientation information

during the delay period (Figure 1c, gray region). We trained a pattern classifier to

discriminate the grating orientation (25° vs. 115°) during an independent, passive-

viewing localizer run. The classifier was then used to discriminate the gratings during

the retention period of the main experiment. We used classification distance to

measure the amount of orientation information present in the activity patterns

(Buitinck et al., 2013). This approach results in a binary prediction (25° or 115°), but

additionally yields a continuous metric of the decoding performance that can be seen

as the confidence of the classification (see Methods for additional details).

425
No remapping of VWM in early visual cortex following eye movements.

During the no-saccade condition, activity patterns in the contralateral early

visual areas (V1 -V3) contained information about the maintained orientation during

the delay period (7.2 s - 10.8 s after onset of maintenance) (Figure 2a, left green bar;

$t_{(23)}$ = 3.60, p = 0.0015, Cohen's d = 0.73), whereas there was no orientation

information in the ipsilateral hemisphere (Figure 2a, left purple bar; $t_{(23)}$ = -0.25, p =

0.8078, Cohen's d = 0.05), in line with a retinotopically specific VWM storage (Pratte

and Tong, 2014). Indeed, Bayesian analyses indicated moderate support (BF_{01} > 3)

for the null hypothesis of no VWM representations in ipsilateral early visual areas

(BF_{01} = 4.531). In the saccade trials, however, orientation information in the

contralateral early visual cortex could no longer be significantly discriminated by the

classifier (Figure 2a, right green bar; $t_{(23)}$ = 1.9, p = 0.0706, Cohen's d = 0.39),

suggesting that the VWM content that was initially maintained in contralateral regions

is degraded by an eye movement during the delay period. Interestingly, the VWM

representations in contralateral early visual areas did not remap to its ipsilateral

hemisphere, given that the classifier could not infer, above chance, the remembered

orientation from activity patterns in ipsilateral early visual areas (Figure 2a, right

purple bar; $t_{(23)}$ = 0.69, p = 0.495, Cohen's d = 0.14). Bayesian analyses also showed

moderate support for the absence of orientation information in ipsilateral early visual

areas in saccade condition (BF_{01} = 3.743).

Figure 2. VWM representations in ipsi- and contralateral early visual areas (V1 - V3). (a)
Orientation classification performance for gratings presented to the contralateral and
ipsilateral hemispheres during the delay period (7.2 s – 10.8 s after onset of maintenance). In
the no-saccade condition, only the contralateral early visual areas (V1 -V3) maintained the

orientation information during delay period, whereas following an eye movement from one side

to the opposite side during the delay period, the remembered orientation could no longer be

reliably decoded from either the contralateral or the ipsilateral early visual areas, suggesting

that no remapping of VWM occurred around the eye movement. **(b)** The time course of

orientation classification performance in both contralateral and the ipsilateral early visual areas

in the no-saccade condition. Information about the orientation of the remembered grating was

present in the contralateral early visual areas during delay period. No remembered orientation

information was present in the ipsilateral early visual areas. The horizontal bars indicate

statistically significant different from zero (cluster permutation test, 1^{st} cluster: $p = 0.0011$; 2^{nd}

cluster: $p = 0.0346$). **(c)** The time course of orientation classification performance in both

contralateral and ipsilateral early visual areas in the saccade condition, in which participants

had to make a saccade to the opposite side of the screen at the beginning of the delay period.

The remembered orientation information was degraded in the contralateral visual cortex but

did not remap to the ipsilateral hemisphere. The horizontal bars indicate statistically significant

different from zero (cluster permutation test, $p = 0.008$). Error bars and shaded areas denote

SEM. *$p < 0.05$; ** $p < 0.01$; n.s., $p > 0.05$.

Next, we examined the time course of VWM representations in the contralateral and

ipsilateral early visual areas separately. To this end, we applied the classifier that

was trained using the passive localizer session to each time point of the working

memory task, including no-saccade and saccade trials. During no-saccade trials, the

classifier could initially not discriminate between orientations of the remembered

grating in either the contralateral or the ipsilateral early visual areas. However, after

a few seconds following the stimulus presentation, the classifier was able to select

the remembered orientation in the contralateral early visual cortex, while there was

no orientation information in the ipsilateral early visual cortex (Figure 2b, cluster

permutation test, $p < 0.05$). Compared to the saccade trials, after the eye movement

during the delay period, we found that orientation information was strongly degraded

in contralateral early visual areas. Crucially, the remembered orientation information

could not be decoded at any time point during delay period from activity patterns in

the ipsilateral early visual areas (Figure 2c), suggesting that VWM contents were not

remapped after the eye movement.

Our primary approach to quantify orientation-specific activity patterns in early

visual cortex was based on a separate data set, during which participants viewed the

grating stimuli while their attention was engaged elsewhere. While this approach

allows for strong inference (i.e., patterns of activity during the maintenance period

resemble those elicited by the physical presentation of the same stimuli), it cannot

detect consistent orientation-specific activity patterns during VWM maintenance that

are only robustly present during the VWM task, and not during passive viewing.

Therefore, in an exploratory analysis, we used a leave-one-run-out cross-validation

approach to train the classifier within the working memory period and test on the left-

out run.

Unlike in the decoding analysis trained on the localizer, we found that in no-

saccade trials, orientation information was present in both the contralateral (Figure

3a, left green bar; $t_{(23)} = 3.29$, $p = 0.0032$, Cohen's $d = 0.67$) and the ipsilateral

(Figure 3a, left purple bar; $t_{(23)} = 3.18$, $p = 0.0042$, Cohen's $d = 0.65$) early visual

areas. This indicates that there were consistent differences during VWM

maintenance of the different orientations not only in the contralateral but also the

ipsilateral early visual areas during the delay period. However, also here, the

classifier was not able to distinguish between grating orientation in either the

contralateral (Figure 3a, right green bar; $t_{(23)} = 1.50$, $p = 0.1469$, Cohen's $d = 0.31$;

$BF_{01} = 1.738$) or the ipsilateral (Figure 3a, right purple bar; $t_{(23)} = 1.94$, $p = 0.0642$,

Cohen's $d = 0.40$; $BF_{01} = 0.936$) early visual cortex after the eye movement during

saccade trials.

Figure 3. Orientation classification performance during the delay period in early visual cortex (V1 – V3) and intraparietal sulcus (IPS) using a leave-one-run-out cross-validation approach. (a) In the no-saccade condition, both the contralateral and the ipsilateral early visual areas contained orientation information during the delay period. In the saccade condition, however, the orientation information degraded in both hemispheres. **(b)** In the IPS, the classifier could select the remembered orientation in both the contralateral and the ipsilateral hemisphere in the no-saccade trials, in line with the findings in early visual cortex. In the saccade condition, after the eye movement there was still reliable orientation information in the contralateral IPS. Error bars denote SEM. $*$ $p < 0.05$; $**$ $p < 0.01$; $***$ $p < 0.001$; n.s., $p > 0.05$.

Persistent VWM representations after an eye movement in the IPS.

Our findings revealed that VWM representations in early visual cortex during

maintenance were not remapped following eye movements. Bettencourt & Xu (2016)

have recently shown that while remembered orientation information was degraded

by irrelevant distractors in early visual cortex, VWM information remained available

in the superior intraparietal sulcus (IPS). We sought to test whether we could

observe similar results in our study. To this end, we applied the leave-one-run-out

cross-validation method to IPS. First, we found that both the contralateral (Figure 3b,

left green bar; $t_{(23)} = 4.24$, $p = 0.0003$, Cohen's $d = 0.86$) and the ipsilateral IPS

(Figure 3b, left purple bar; $t_{(23)} = 2.22$, $p = 0.0369$, Cohen's $d = 0.45$) contained

information about the remembered orientation in no-saccade trials during delay

period, as also observed in early visual cortex. However, in contrast to the early

visual cortex, the remembered grating orientation could also be decoded from the

contralateral IPS after an eye movement (Figure 3b, right green bar; $t_{(23)}$ = 3.33, p =

0.0029, Cohen's d = 0.68), while there was no reliable orientation information in the

ipsilateral IPS (Figure 3b, right purple bar; $t_{(23)}$ = 1.48, p = 0.1515; BF_{01} = 1.775).

Discussion

In the current study, we investigated whether VWM representations are remapped

following eye movements and whether VWM information persists in the early visual

and parietal cortex after the execution of a saccade. We found robust encoding of

maintained orientation information in the contralateral hemisphere of early visual

cortex. However, this information degraded upon making a saccade to the opposite

hemifield and did not remap to the ipsilateral hemisphere in early visual cortex. This

suggests that there is no remapping of VWM representations across eye movements

in the early visual cortex. Additionally, VWM representations in early visual cortex

were impaired by the saccade during the retention period. In contrast, there was

orientation information in the IPS contralateral to the remembered item, that

remained reliably present also after execution of a saccade. These findings show

that early visual cortex is susceptible to interference of eye movements during a

working memory period, while IPS appears less sensitive to this interference.

Retinotopically specific encoding of VWM information in early visual cortex

In contrast to previous studies that have shown that VWM in early visual cortex is

represented in a spatially global mode (Ester et al., 2009), our study indicates that

the orientation information can be restricted to the contralateral hemisphere when

the orientation and location of the stimulus need to be jointly maintained, in line with

several recent studies (Pratte and Tong, 2014; Lorenc et al., 2018). In Ester et al.

(2009), although the remembered grating was always presented in one of the

hemifields, participants could use a strategy in which they memorized the orientation

of the grating perifoveally instead of at its original location (Williams et al., 2008).

This strategy would lead to a bilateral VWM representation irrespective of where the

grating is presented. In order to encourage our participants to represent the grating

in its original location, we inserted catch trials, in which the probe grating and the

remembered gratings were horizontally offset one visual degree. Participants were

instructed to respond only when the probe location matched the location of the

initially presented grating. Therefore, they had to maintain the orientation of the

gratings at the presented location in order to be able to successfully complete the

task. Our results revealed that VWM representation during our task were indeed

retinotopically specific (see Figure 2a), in line with previous findings (Pratte and

Tong, 2014).

No remapping of VWM following eye movements.

Attentional or feature based remapping of visual input just prior to a saccade has

been well established from a variety of studies and is thought to maintain visual

stability across eye movements. For instance, Rolfs et al. (2011) found that attention

could be predictively remapped to the future retinotopic location of an upcoming

target, even before the execution of the eye movement. In the current experiment,

we asked whether memory traces would also be updated with eye movements.

However, we found no evidence that visual working memory representations were

remapped following eye movements. At least two reasons should be emphasized to

address the discrepancy in remapping between the attention and VWM. First, to

survive in a multifaceted and dynamically changing world, our attention needs to be

selectively shifted to the most important stimuli that are goal-relevant. In order to be

competent for such a complicated task, remapping of attention could be one of the

most crucial mechanism to prepare information in advance. In contrast, the main

function of working memory is to temporarily retain information available for

processing. While attention could be flexibly remapped to different locations to

extract information, storing working memory information in one or several stationary

regions could be potentially an efficient way for retrieval. Second, in line with the

arguments of 'activity-silent' models of WM when compared to the persistent activity

models of WM (Stokes, 2015; Wolff et al., 2017), remapping of VWM after each eye

movement could also be energetically expensive, especially if VWM is maintaining

an up-to-date information of the external environment. Therefore, a stationary

working memory representation could be an ideal way for ecologically maintaining

and manipulating information in the brain.

Parietal cortex may subserve VWM representations after saccades.

In recent years, the parietal cortex has been repeatedly implicated in VWM

maintenance (Todd and Marois, 2004; Christophel et al., 2015; Ester et al., 2015;

Bettencourt and Xu, 2016), alongside the involvement of sensory areas (Harrison

and Tong, 2009; Serences et al., 2009; Emrich et al., 2013). Our results indicate that

the early visual cortex is susceptible to the interference of eye movements during the

delay period, potentially limiting the role of the early visual cortex in VWM. Similarly,

Bettencourt and Xu (2016) observed that VWM representations in early visual cortex

were degraded by the presentation of distractors, while superior IPS exhibited

persistent activity that was not impaired by the presentation of intervening distractor

stimuli. In our experiment, the eye movements during the delay period potentially

may also generate distracting new input to the visual system, which could disturb the

VWM representations. Intriguingly, while VWM representations in early visual cortex

were degraded following the eye movement, these representations were still

maintained in IPS. Behaviorally, participants' memory performance was

indistinguishable between the saccade and no saccade trials. Therefore, it could be

that the preserved behavioral performance we observed is supported by these IPS

representations, rather than the early visual cortex representations. In a more recent

study, Lorenc et al. (2018) found that the VWM information could be reliably

maintained only in early visual cortex without distractors. However, these early visual

cortex representations seem to transfer to the IPS representations when distractors

were presented, which suggesting a flexible coding of VWM between the early visual

cortex and the IPS depending on the task demands. This shows that VWM

information maintained in visual cortex is susceptible to interference, in line with the

idea that the early visual cortex may only play a limited role in VWM storage.

In conclusion, our results indicate that VWM representations in early visual

cortex are degraded and not remapped after eye movements, limiting the role of

early visual cortex in VWM storage.

References

Albers AM, Kok P, Toni I, Dijkerman HC, de Lange FP (2013) Shared

Representations for Working Memory and Mental Imagery in Early Visual

Cortex. Curr Biol 23:1427–1431.

Bettencourt KC, Xu Y (2016) Decoding the content of visual short-term memory

under distraction in occipital and parietal areas. Nat Neurosci 19:150–157.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Buitinck L, Louppe G, Blondel M, Pedregosa F, Mueller A, Grisel O, Niculae V,

Prettenhofer P, Gramfort A, Grobler J, Layton R, Vanderplas J, Joly A, Holt B,

Varoquaux G (2013) API design for machine learning software: experiences

from the scikit-learn project. ArXiv Prepr ArXiv:1309.0238.

Christophel TB, Cichy RM, Hebart MN, Haynes J-D (2015) Parietal and early visual

cortices encode working memory content across mental transformations.

NeuroImage 106:198–206.

Curtis CE, D'Esposito M (2003) Persistent activity in the prefrontal cortex during

working memory. Trends Cogn Sci 7:415–423.

DeYoe EA, Carman GJ, Bandettini P, Glickman S, Wieser J, Cox R, Miller D, Neitz J

(1996) Mapping striate and extrastriate visual areas in human cerebral cortex.

Proc Natl Acad Sci 93:2382–2386.

Durstewitz D, Seamans JK, Sejnowski TJ (2000) Dopamine-Mediated Stabilization of

Delay-Period Activity in a Network Model of Prefrontal Cortex. J Neurophysiol

83:1733–1750.

Emrich SM, Riggall AC, LaRocque JJ, Postle BR (2013) Distributed Patterns of

Activity in Sensory Cortex Reflect the Precision of Multiple Items Maintained

in Visual Short-Term Memory. J Neurosci 33:6516–6523.

Engel SA, Glover GH, Wandell BA (1997) Retinotopic organization in human visual

cortex and the spatial precision of functional MRI. Cereb Cortex 7:181–192.

Ester EF, Serences JT, Awh E (2009) Spatially Global Representations in Human

Primary Visual Cortex during Working Memory Maintenance. J Neurosci

29:15258–15265.

Ester EF, Sprague TC, Serences JT (2015) Parietal and Frontal Cortex Encode

Stimulus-Specific Mnemonic Representations during Visual Working Memory.

Neuron 87:893–905.

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A,

Killiany R, Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, Dale AM

(2002) Whole Brain Segmentation: Automated Labeling of Neuroanatomical

Structures in the Human Brain. Neuron 33:341–355.

Harrison SA, Tong F (2009) Decoding reveals the contents of visual working memory

in early visual areas. Nature 458:632–635.

Haynes J-D, Rees G (2006) Neuroimaging: Decoding mental states from brain

activity in humans. Nat Rev Neurosci 7:523–534.

He T, Fritsche M, Lange FP de (2018) Predictive remapping of visual features

beyond saccadic targets. J Vis 18:20–20.

JASP Team (2019) JASP (Version 0.10.2)[Computer software].

Kamitani Y, Tong F (2005) Decoding the visual and subjective contents of the human

brain. Nat Neurosci 8:679–685.

Lee MD, Wagenmakers E-J (2014) Bayesian Cognitive Modeling: A Practical

Course. Cambridge University Press.

Lorenc ES, Sreenivasan KK, Nee DE, Vandenbroucke ARE, D'Esposito M (2018)

Flexible Coding of Visual Working Memory Representations during

Distraction. J Neurosci 38:5267–5276.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-

data. J Neurosci Methods 164:177–190.

Melcher D (2007) Predictive remapping of visual features precedes saccadic eye

movements. Nat Neurosci 10:903–907.

Norman KA, Polyn SM, Detre GJ, Haxby JV (2006) Beyond mind-reading: multi-

voxel pattern analysis of fMRI data. Trends Cogn Sci 10:424–430.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M,

Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D,

Brucher M, Perrot M, Duchesnay É (2011) Scikit-learn: Machine Learning in

Python. J Mach Learn Res 12:2825–2830.

Pratte MS, Tong F (2014) Spatial specificity of working memory representations in

the early visual cortex. J Vis 14:22.

Prins N, Kingdom FAA (2018) Applying the Model-Comparison Approach to Test

Specific Research Hypotheses in Psychophysical Research Using the

Palamedes Toolbox. Front Psychol 9:1250.

Rayner K (1998) Eye movements in reading and information processing: 20 years of

research. Psychol Bull 124:372–422.

Riley MR, Constantinidis C (2016) Role of Prefrontal Persistent Activity in Working

Memory. Front Syst Neurosci 9:181.

Rolfs M, Jonikaitis D, Deubel H, Cavanagh P (2011) Predictive remapping of

attention across eye movements. Nat Neurosci 14:252–256.

Savitzky Abraham, Golay MJE (1964) Smoothing and Differentiation of Data by

Simplified Least Squares Procedures. Anal Chem 36:1627–1639.

Serences JT (2016) Neural mechanisms of information storage in visual short-term

memory. Vision Res 128:53–67.

Serences JT, Ester EF, Vogel EK, Awh E (2009) Stimulus-Specific Delay Activity in

Human Primary Visual Cortex. Psychol Sci 20:207–214.

Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg

H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J,

Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM (2004) Advances

in functional and structural MR image analysis and implementation as FSL.

NeuroImage 23:S208–S219.

Stokes MG (2015) 'Activity-silent' working memory in prefrontal cortex: a dynamic

coding framework. Trends Cogn Sci 19:394–405.

Todd JJ, Marois R (2004) Capacity limit of visual short-term memory in human

posterior parietal cortex. Nature 428:751–754.

Wandell BA, Dumoulin SO, Brewer AA (2007) Visual Field Maps in Human Cortex.

Neuron 56:366–383.

Wang L, Mruczek REB, Arcaro MJ, Kastner S (2015) Probabilistic Maps of Visual

Topography in Human Cortex. Cereb Cortex 25:3911–3931.

Williams MA, Baker CI, Op de Beeck HP, Mok Shim W, Dang S, Triantafyllou C,

Kanwisher N (2008) Feedback of visual object information to foveal retinotopic

cortex. Nat Neurosci 11:1439–1445.

Wolff MJ, Jochim J, Akyürek EG, Stokes MG (2017) Dynamic hidden states

underlying working-memory-guided behavior. Nat Neurosci 20:864–871.

Xu Y (2017) Reevaluating the Sensory Account of Visual Working Memory Storage.

Trends Cogn Sci 21:794–815.

Xu Y (2018) Sensory Cortex Is Nonessential in Working Memory Storage. Trends

Cogn Sci 22:192–193.