Visual working memory representations in visual and parietal cortex do not remap after eye movements

Tao He[1*], Matthias Ekman[1], Annelinde R.E. Vandenbroucke[1,2] and Floris P. de Lange[1]

[1] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Kapittelweg 29, 6525 EN Nijmegen, the Netherlands
[2] Current Address: Dutch National Research Agenda, Laan van Nieuw Oost Indië 300, 2593 CE Den Haag, the Netherlands
[*] Correspondence: t.he@donders.ru.nl

Highlights

- Visual working memory (VWM) representations do not remap after making saccades

- Eye movement degrade VWM representations in early visual cortex, limiting the role of early visual cortex in VWM storage

- Parietal cortex shows persistent VWM representations across saccades

ABSTRACT

It has been suggested that our visual system does not only process stimuli that are

directly available to our eyes, but also has a role in maintaining information in VWM

over a period of seconds. It remains unclear however what happens to VWM

representations in the visual system when we make saccades. Here, we tested the

hypothesis that VWM representations are remapped within the visual system after

making saccades. We directly compared the content of VWM for saccade and no-

saccade conditions using MVPA of delay-related activity measured with fMRI. We

found that when participants did not make a saccade, VWM representations were

robustly present in contralateral early visual cortex. When making a saccade, VWM

representations degraded in contralateral V1-V3 after the saccade shifted the

location of the remembered grating to the opposite visual field. However, contrary to

our hypothesis we found no evidence for the representations of the remembered

grating at the saccadic target location in the opposite visual field, suggesting that

there is no evidence for remapping of VWM in early visual cortex. Interestingly, IPS

showed persistent VWM representations in both the saccade and no-saccade

condition. Together, our results indicate that VWM representations in early visual

cortex are not remapped across eye movements, potentially limiting the role of early

visual cortex in VWM storage.

Keywords: visual working memory; eye movements; fMRI; remapping; MVPA

Introduction

Research in the past decade has suggested that our visual system may not only

process incoming information, but also be relevant for maintaining internal

representations of previously observed visual stimuli, i.e. visual working memory

(VWM). The ability to maintain information that is no longer in view is critical for

reasoning about and mentally manipulating visual information despite temporal

discontinuous in visual inputs that occur for example during eye movements,

occlusions, and object motion (Curtis and D'Esposito 2003; Serences 2016; Lorenc

et al. 2018).

VWM information has been observed both in early sensory areas (Ester et al.

2009; Harrison and Tong 2009; Albers et al. 2013; Sreenivasan et al. 2014), as well

as higher-order parietal (Christophel et al. 2015; Ester et al. 2015; Bettencourt and

Xu 2016), and prefrontal regions (Goldman-Rakic 1995; Durstewitz et al. 2000;

reviewed in Riley and Constantinidis 2016). Moreover, VWM representations in early

visual cortex have also been found to be spatially specific, maintained in a

retinotopic manner, at least in V1 and V2 (Pratte and Tong 2014; although see Ester

et al. 2009). Crucially however, in these previous studies, participants were

instructed to maintain fixation while remembering a visual stimulus during a retention

period. While this is a common approach to investigate VWM in a lab environment, it

is arguably quite different from real-world settings that are marked by multiple eye

movements per second. Humans perform about three saccadic eye movements per

second in order to guide the fovea toward regions of interest in the visual field

(Rayner 1998). Previous research has indicated that both stimulus features and

attentional pointers are remapped across eye movement to aid visual stability

63   (Melcher 2007; Rolfs et al. 2011; He et al. 2018). It is currently unclear however,

64   whether VWM is also remapped within the visual system after making saccades.

65   One possibility is that VWM representations are dynamically remapped and

follow the shift of the retinotopic location caused by eye movements. In this case, the

cortical location representing the VWM would be updated after an eye-movement.

Alternatively, if VWM does not remap, working memory information might be

maintained across multiple brain regions following eye movements. On this account,

after every eye movement, the up-to-date feature of the stimulus (e.g., the latest

location of the stimuli) would be integrated with the previous working memory

information.

73   Here, we tested the potential remapping of VWM in the early visual system,

by presenting participants with an orientated grating in either the left or right visual

field. The grating orientation had to be maintained for a subsequent retention period,

while we measured blood oxygenation level-dependent (BOLD) signals with

functional magnetic resonance imaging (fMRI). Using multivariate pattern analysis

(MVPA), we attempted to decode the remembered orientation in both the

contralateral and the ipsilateral hemispheres during trials in which participants

performed either a saccade or maintained fixation during the working memory

period.

82   To preview, we found a contralateral VWM representation in the early visual

83   system. This VWM representation degraded when participants made a saccade and

84   was not remapped to ipsilateral visual cortex. In contrast, VWM representations in

85   IPS persisted after saccades. These findings suggest a limited role of the early visual

86   system in VWM.

Materials and Methods

Data and code availability

All data and code used for stimulus presentation and analysis is freely available on
the Donders Repository (https://data.donders.ru.nl/login/reviewer-
77603154/28KfrmSvRmvGnYdOkGqdQQfMufTI7W29jY686hMMaro).

Participants

Thirty-four healthy participants (18 females, mean age 24.3 years, ranging from 19 to
33 years) with normal or corrected-to-normal vision were recruited from the institute's
subject pool in exchange for either monetary compensation or study credits. This
sample size of N = 34 included subjects ensured 80% power to detect an effect size
of at least Cohen's $d \geq 0.5$. All participants were naive with respect to the purposes
of the study. The experiments were approved by the Radboud University Institutional
Review Board and were carried out in accordance with the guidelines expressed in
the Declaration of Helsinki. Written informed consent was obtained from all
participants prior to the start of the study. Only participants who completed the entire
experimental protocol (i.e., the behavioral experiment and the two fMRI sessions)
were included in the final analysis. Data from six participants were excluded: one
participant didn't complete the fMRI sessions due to discomfort, one participant failed
to follow the instructions (i.e., fixate and saccade) during experiment, four
participants had excessive head motion (motion cutoff = 1 mm).

Stimuli

Stimuli were programmed in MATLAB (v2016a, The MathWorks, Inc., Natick, MA)
using the Psychtoolbox (Brainard 1997). The circular sinusoidal grating stimulus

subtended 10° and was centered with a small jitter (0.3°) on the screen center. The

grating was full contrast, with a spatial frequency of 1 cycles/degree, a random

phase and an orientation of either 25° or 115° (with a small jitter of 3°) from the

horizontal axis. The contrasts of the edges of the grating were linearly attenuated

over the distance from 4.5° to 5.0° radius. Two filled dots (0.5°, one green, one

black) were presented at the periphery of the screen (6° left or right away from the

center of the screen), which were served as the fixation dot and saccade target in

the task. In the behavioral training session, stimuli were presented on a 24-inch flat-

panel display (BenQ XL2420T, 1,920 x 1,080 resolution, 60 Hz refresh rate). In the

fMRI sessions, stimuli were displayed on a rear-projection screen using an EIKI LC-

XL100L (EIKI, Rancho Santa Margarita, CA) multimedia projector (1,024 x 768

resolution, 60 Hz refresh rate).

Experimental design

Behavioral training procedure.   Prior to the fMRI scan sessions, all participants

completed a one-hour behavioral training session to familiarize themselves with the

fMRI main task and to establish their individual orientation discrimination threshold,

which served as an initial orientation difference of the gratings in the following fMRI

sessions. The experimental design of the behavioral training task was exactly the

same as during the fMRI main task, except that the delay period and the inter-trial

interval (ITI) were shortened to 3 s to reduce the experimental time. Participants

completed two to three blocks of 56 trials until a stable orientation discrimination

threshold was obtained, during which the eye movements were continuously

monitored by an Eyelink 1000 plus eye tracker. In addition, participants were

familiarized with the localizer tasks and the retinotopic mapping procedures that

were used in the fMRI session at the end of the behavior session.

fMRI main task.   The experimental design (Figure 1a) was adapted from a well-

known delayed orientation discrimination task used before (Harrison and Tong

2009). Each trial began with two filled dots (one green, one black) that were

presented at the periphery of the screen (2 s). Participants were asked to fixate at

the green dot through the trial. Two oriented gratings were flashed sequentially at the

center of the screen (which could be in the left or right visual field of the participants,

depending on the location of the green fixation dot) for 200 ms, respectively, with an

inter-stimulus interval of 400 ms, and followed by a retro-cue (400 ms, "1" or "2") to

indicate which orientation of the grating should be remembered during the following

delay period. The sequence of the orientations (25° and 115°) were randomly

chosen on each trial. On 10 out of 28 trials (No-saccade condition) in each run, the

two dots did not swap their locations, which indicated to participants that they

needed to keep their fixation at the initial fixation location during the 10 s delay

period. On the remaining 18 trials (Saccade condition) in the run, the green dot was

shifted to the opposite side of the screen at the beginning of the delay period (2 s

after the offset of the retro-cue). This instructed participants to move their eyes to the

opposite location of the screen and maintain their fixation at the new position for the

remaining 8 s. A probe grating was presented at the screen center after the delay

period and participants were required to make a judgement of whether the probe

orientation was rotated clockwise or count-clockwise relative to the grating they

memorized during the delay period. Finally, feedback was provided and a black

fixation dot instructed participants to move their eyes to the center of the screen in

preparation for the next trial. Trials were separated by an ITI of 9.6 s. The central

fixation dot changed its color from black to gray at the last second of the ITI to

indicate that the next trial was going to start.

In order to incentivize participants to form spatially specific visual working

memories, four catch trials were included in each run. Stimuli and timing of the catch

trials were identical to the main trials, except for the following changes. During the

presentation of the probe in the catch trial, the probe was horizontally shifted 1.2° to

the left or right with respect to the location of the sample gratings. Participants were

instructed to not respond when the location of the probe grating did not match with

the sample gratings. The rationale of this was that it forced participants to memorize

the orientation stimulus at its original location in order to successfully perform the

task, resulting in a spatially specific VWM representation.

Each run comprised 28 trials, consisting of 10 no-saccade trials (including 2

catch trials), 18 saccade trials (including 2 catch trials). We chose these trial

numbers because the saccade trials were of primary interest and we therefore

wanted to have maximal sensitivity for this condition. The order of trials was a

pseudo-randomized within each run. Each trial lasted 26.4 s, consisting of a 16.8 s

task period and a 9.6 s ITI. Each run lasted 12.32 min and started with 6 s of fixation

that was discarded from the analysis.

Staircase Procedure

The staircase procedure was used to ensure equal task difficulty across participants

and to equate task difficulty levels across no-saccade and saccade conditions within

participants. In the behavioral training session, immediately after the delay period,

participants were asked to compare the tilted orientation between the probe and the

internal remembered grating. The staircase procedure estimated the difference

between probe and remembered grating that ensured 75% performance, using

QUEST (Watson and Pelli 1983; Prins and Kingdom 2018). A maximum orientation

difference of 20° between the probe and remembered grating was enforced. The

staircase was initiated with an orientation difference of 10° and dynamically adapted

according to participants' performance on previous trial until a stable threshold was

acquired. This threshold was used as a seed in the following fMRI sessions and the

same staircase procedure was also used during scanning.

Eye tracking

Eye position was monitored with an MR-compatible Eyelink 1000 (SR Research Ltd.,

Ottawa, Canada) eye tracker. Only the left eye was recorded in the scanner. Pupil

and corneal reflection were sampled at 1000 Hz and analyzed offline to ensure that

participants fixated at the correct location. The eye tracker was calibrated at the

beginning of each session and repeated between runs if necessary. In 10 out of 192

runs, the eye tracker signal was lost during scanning due to subjects' head motion or

technical problems. During these runs, the experimenter monitored the eye position

online via the live video feed from the camera. All participants were trained on the

fixation task and to perform saccades in the behavioral training session prior to the

scanning session.

fMRI parameters

Functional and anatomical images were carried out with a 3T Siemens Prisma fit

MRI system (Siemens, Erlangen, Germany), using a 32-channel headcoil. Functional

images were acquired using a whole brain T2*-weighted multiband-4 sequence

(TR/TE = 1,200/39 ms, voxel size 2.4x2.4x2.4 mm, 56 transversal slices, 65° flip

angle, A/P phase encoding direction, FOV = 210 mm, BW = 2030 Hz/Px).

Anatomical images were acquired using a T1-weighted magnetization prepared rapid

gradient echo (MP-RAGE) sequence (TR/TE = 2,300/3.03ms, voxel size 1x1x1 mm,

192 transversal slices, 8° flip angle).

fMRI Preprocessing

fMRI data were preprocessed using FSL (FMRIB Software Library; Oxford, UK;

www.fmrib.ox.ac.uk/fsl; Smith et al. 2004, RRID:SCR_002823), including motion

correction (six-parameter affine transform), temporal high-pass filtering (100 s) and

Savitzky–Golay low-pass filter (time window length = 11 TRs, polynomial order = 3;

Savitzky & Golay, 1964) for each run separately. No spatial smoothing and slice

timing correction was performed. All univariate and multivariate analyses were

performed in native subject space with custom python code using Nibabel

[https://doi.org/10.5281/zenodo.1464282], Scipy (Oliphant 2007; Millman and Aivazis

2011) and scikit-learn (Pedregosa et al. 2011).

Functional localizers

In addition to the main experiment, participants underwent two localizer runs, which

were used to select voxels that maximally responded to stimuli presented in the

contralateral hemifield in both the univariate and multivariate analysis. The same

grating stimulus parameter were used in localizer runs as those in the main

experiment. Participants fixated at the left or right green dot and the grating was

presented at the center of screen for 16 s, with a frequency of 4 Hz. Throughout the

localizer, participant had to fixate at the green dot and monitor a sequence of rapidly

changing letters just above fixation, to which they had to respond by button press

whenever a target letter ("X" or "Z") occurred in a stream of non-target letters ("A",

"H", "R", "N", "T", "V", "U", "Y"). Letters were presented at a frequency of 2 Hz. When

the green dot shifted from the left to the right side of the screen (or vice versa) at the

end of the trial, participants had to move their eyes to follow the fixation dot. Each

trial was separated by a 2 s ITI in which only the fixation cross was presented, to

give participants enough time to saccade and stabilize their eyes in the new position.

Two blocks of 20 trials (10 trials per visual field: left or right) were collected.

Images of the localizer runs were also preprocessed following the same

procedure as the main tasks. Onsets and duration of the localizer blocks were

convolved with a double gamma haemodynamic response function (HRF) and fitted

using a general linear model in left and right hemisphere separately. For each

participant a two-sided t-contrast was calculated contrasting between left and right

hemispheres. Resulting statistical maps were thresholded at $Z > 3.1$ and a corrected

cluster significance threshold of $p < 0.05$.

Retinotopic mapping of early visual areas (V1 – V3)

Early visual areas (V1, V2, and V3) for each participant were identified using two

retinotopic mapping blocks based on the standard traveling-wave method using

rotating wedges (DeYoe et al. 1996; Engel et al. 1997; Wandell et al. 2007),

consisting of a clockwise and a counterclockwise-rotating run. Participants were

instructed to fixate at the center of the screen and fulfill rapid letter detection task

(exact same as that in localizer runs). The BOLD responses to the wedges were

used to estimate the polar angle of the visual field representation.

Regions of interest (ROIs)

Early visual areas (V1 - V3) and the intraparietal sulcus (IPS0 – IPS5) within each

hemisphere were independently chosen as the ROIs in our analyses. For the ROI of

the early visual areas, we first used Freesurfer (http://surfer.nmr.mgh.harvard.edu/,

Fischl et al. 2002) to define the gray–white matter boundary and perform cortical

surface reconstruction. The borders of the early visual areas V1 - V3 were delineated

based on retinotopic maps. The ROI of the intraparietal sulcus was defined based on

anatomical probability maps of retinotopic areas in the intraparietal sulcus (IPS0 –

IPS5) from the Probabilistic atlases (Wang et al. 2015). Finally, all surface-based

ROIs were backward-transformed into participant's native space. Next, the visually

active voxels corresponding to the grating that was positioned at the left and right

visual field was identified based on statistical activation maps from the functional

localizer runs. Two types of ROI were defined in the decoding analyses: contralateral

ROIs and ipsilateral ROIs. For instance, when decoding the orientation of the grating

that was initially presented at participant's right visual field, regardless of the no-

saccade and saccade condition, the left hemisphere would be labeled as the

contralateral ROIs, while the right hemisphere would be labeled as the ipsilateral

ROIs (see Figure 1b) and vice versa. Finally, we selected the 150 most active voxels

across visual areas V1, V2, V3, as well as 250 most active voxels across the

combined early visual cortex (V1 - V3) and the entire intraparietal sulcus (IPS0 –

IPS5) to perform the decoding analyses, separately.

Univariate fMRI analyses

To estimate the BOLD response of each hemisphere in each condition, we selected

the same voxels as those used for decoding. We separately modeled the onset of

each trial for each hemisphere and for no-saccade and saccade condition within

each run to fit voxel-wise general linear models (GLM) using FSL FEAT. An

additional nuisance regressor of 24 motion regressors (FSL's standard + extended

motion parameters) were also added to the GLM. To quantify BOLD activity during

the trial, contrasts between left and right hemisphere regressors for each condition

were created. Multiple-comparison correction was performed using Gaussian

random-field based cluster thresholding. The significance level was set at the

cluster-forming threshold of z > 3.1 (i.e., $p < 0.001$, two-sided) and a cluster

significance threshold of $p < 0.05$. All fMRI data were transformed from MRI signal

intensity to units of percent signal change, calculated relative to the average level of

activity for each voxel across the first volume of each trial within each run. BOLD

activity over time were statistically tested using nonparametric cluster-based

permutation t test (10,000 permutations; two-sided $p < 0.05$; cluster formation

threshold $p < 0.05$) (Maris and Oostenveld 2007).

Multivariate fMRI analyses

We used multivariate pattern analyses (MVPA) to determine whether the pattern of

activity in each ROI and each hemisphere contained orientation information, as

implemented in Scikit-learn 0.20.3 (Pedregosa et al. 2011). To this end, linear

support vector machines (SVMs) were trained to discriminate between the two

grating orientations based on the pattern of BOLD activity over voxels. In this study

we used classification distance as an indication of the amount of orientation

information being maintained in each hemisphere. To calculate the classification

distance, we measured the distance of each sample to the separating hyperplane

that was trained by a linear SVM with a positive/negative sign (the sign indicates the

class). We then averaged the distance within each class and calculated the absolute

distance between these two classes. While this approach can give us a binary

predicted label (25° or 115°), which can be used to calculate classification accuracy,

it additionally allows one to review the confidence of this classification and provide a

continuous metric of the decoding performance. In each ROI, the larger the

classification distance, the more confident the classifier is in determining the stimulus

class based on the BOLD activity pattern, and hence the more orientation

information is contained within the pattern of BOLD activity.

When training and testing within the main experiment we averaged the BOLD

activity over time points 7.2 - 10.8 s after the onset of the delay period. This time

period was selected to be as far from the onset of the two gratings at the beginning

of the trial as possible, and therefore not reflect activity elicited by the stimuli, but

also prior to the onset of the probe stimulus. These time series were then normalized

on a voxel-by-voxel and run-by-run basis for each voxel using a z-scored

transformation and sorted into one of sixteen bins based on four factors: hemisphere

(left or right hemisphere), orientation (25° or 115°), stimulus location (left or right

hemifield) and saccade condition (saccade or no-saccade). A leave-one-run-out

cross-validation procedure was used to train the classifier where we trained in 7

("training" dataset) out of 8 runs and tested on the remaining run ("test" dataset) for

each hemisphere, stimulus location and saccade condition pair, separately. Within

the independent "training" and "test" dataset, activation patterns comprising the

mean response of each voxel during 25° or 115° trials were calculated. Finally, the

decoding distances in each corresponding stimulus location and hemisphere pair

were collapsed into the contralateral (i.e., left hemifield and right hemisphere, right

hemifield and left hemisphere) or ipsilateral (i.e., left hemifield and left hemisphere,

right hemifield and right hemisphere) hemisphere condition.

Bayesian analyses

In order to further evaluate all the statistical tests, we performed the Bayesian

equivalents of the above outlined analyses. JASP 0.10.2 (JASP Team, 2019,

RRID:SCR_015823) was used to perform all Bayesian analyses, using default

settings. Thus, for Bayesian t-tests a Cauchy prior width of 0.707 was chosen.

Qualitative interpretations of Bayes Factors are based on criteria by Lee &

Wagenmakers (2014).

335
Results

Our primary goal was to investigate whether VWM representations in early visual

cortex (V1-V3) are remapped after eye movements. We used MVPA (Kamitani and

Tong 2005; Haynes and Rees 2006; Norman et al. 2006) to determine whether the

information in VWM that was initially represented in the contralateral hemisphere

(corresponding to the stimulus location) was remapped to the opposite (ipsilateral)

hemisphere following an eye movement to the opposite visual field during the

retention period. 34 participants performed a delayed orientation discrimination task,

in which they initially fixated at the peripheral fixation dot while passively viewing two

successive grating stimulus orientations (25° and 115°) in one visual field, followed

by a cue and a delay period. Importantly, in 18 out of 28 trials (*saccade condition*) in

each run, participants were asked to saccade to the opposite side of the screen

during the delay period, while in the remaining 10 trials (*no-saccade condition*) in the

same run, participants maintained fixation at the initial side through the entire trial. A

probe grating was presented at the end of the trial (**Figure 1a, b**).

In order to encourage participants to memorize the grating in its original

location, we also included catch trials in which the probe grating was horizontally

shifted 1.2° with respect to the location of the sample gratings. Participants were

instructed to discriminate the orientation between the probe and the memorized

grating only when they were presented at the same location (see Methods for

details). This experimental design was chosen to encourage participants to maintain

both the orientation and location of the presented grating. Indeed, behavioral data

showed that participants successfully withheld their response (i.e. correct rejection

rate) at a rate of 63.6% (SD 18.37) and 62.87% (SD 19.58) in both no-saccade and

saccade conditions when the probe was presented at a displaced location, and no

difference was found between the two conditions ($t_{(33)}$ = 0.257, *p* = 0.7989, Cohen's

$d = 0.039$; $BF_{10} = 0.189$), suggesting that participants remembered the orientation in

a spatially specific way.

Behavioural data further confirmed that participants could successfully

discriminate small differences between the cued grating and the probe grating

(**Figure 1c**, mean discrimination threshold, no-saccade condition: 8.63°, SD 5.07;

saccade condition: 8.89°, SD 4.55) at an accuracy of 82.1% (SD 5.05) and 81.1%

(SD 4.43) in both no-saccade and saccade trials. Notably, there was no difference in

accuracy between trials with and without a saccade ($t_{(33)} = 1.125$, $p = 0.2687$,

Cohen's $d = 0.193$; $BF_{10} = 0.328$), and no difference in the mean discrimination

threshold ($t_{(33)} = 0.66$, $p = 0.514$, Cohen's $d = 0.113$; $BF_{10} = 0.225$), indicating that the

VWM performance was not impaired by the eye movement during the delay period.

A univariate BOLD analysis showed that the presentation of the grating stimuli

induced stronger BOLD activity in contralateral than ipsilateral early visual areas

(cluster permutation test, $p = 0.0001$). As expected, probe presentation at the end of

the trial led to a BOLD activity increase in the contralateral hemisphere during no-

saccade trials and a BOLD activity increase in the ipsilateral hemisphere,

corresponding to the updated stimulus location after the eye movement, in the

saccade condition (**Figure 1d**). Taken together, these results suggest that

participants engaged successfully in the task.

Figure 1 Experimental paradigm, behavioral performance and BOLD activity in the early visual cortex. (a) Participants performed a delayed orientation discrimination task. At the start of each trial, two dots were presented at the periphery of the screen and participants were instructed to fixate at the green dot. Two orientated gratings (25° and 115°) were successively flashed in the center of the screen, followed by a retro-cue (1 or 2) that indicated which grating to remember in the following delay period. Crucially, during a 10 s delay period, if the green dot shifted to the opposite side of the screen (*saccade condition*), participants had to make an eye movement to the opposite side and maintained at the new location through the following time period (the dashed arrow is for illustration only, not present in the actual test). Conversely, in the no-saccade condition, the green dot did not change its position and participants maintained fixation at the initial side throughout the entire trial. After the delay period a probe was presented, and participants indicated whether the probe was tilted clockwise or counterclockwise relative to the remembered grating. **(b)** Illustration of the experimental design probing remapping. In the no-saccade condition (left), the remembered grating presented in right visual field is initially represented in the left (contralateral) early visual cortex.

In the saccade condition (right), after participants saccade from the left to the right side of the

screen, we set out to examine whether VWM contents that initially generated in contralateral

(left) early visual cortex transfer to the ipsilateral (right) visual cortex. **(c)** there was no

difference in the mean discrimination threshold between trials with and without a saccade,

demonstrating that the VWM performance was not impaired by the eye movement during the

delay period. Grey dots with connecting lines denote individual participants. Colors are

estimated densities, white dots are group medians, boxes are quartiles and whiskers are 1.5

interquartile range. **(d)** Group averaged (N = 34) BOLD time course in contralateral and

ipsilateral early visual areas (V1 – V3). In the no-saccade condition, the presentation of grating

stimuli evoked a higher BOLD response in contralateral than in ipsilateral early visual areas.

In the saccade condition, while the BOLD activity was still higher in contralateral relative to

ipsilateral early visual areas after presentation of the grating stimuli, the pattern inverted after

presentation of the probe – BOLD activity became higher in ipsilateral than in contralateral

early visual areas. The first vertical dashed gray line represents the onset of the gratings, while

the second vertical dashed gray line indicates the onset of the probe. The vertical dashed red

line in saccade condition indicates the onset of the saccade. The vertical gray bar represents

the delay period (7.2 s - 10.8 s after onset of maintenance) that selected for multivariate

analysis. BOLD activity data was interpolated and smoothed for display only, all statistical

tests were applied before data interpolation. Shaded areas denote SEM.

In order to evaluate the potential remapping of VWM in early visual cortex, we

assessed whether the patterns of activation in both hemispheres, contralateral and

ipsilateral to the item location, contained the remembered orientation information

during the delay period (**Figure 1d**, gray region). A leave-one-run-out cross-

validation approach was used to train the classifier to discriminate the grating

orientation (25° vs. 115°) within the working memory period and test on the left-out

run. We used classification distance to measure the amount of orientation

information present in the activity patterns (Dijkstra et al. 2019; Linde-Domingo et al.

2019). This approach results in a binary prediction (25° or 115°), but additionally

yields a continuous metric of the decoding performance that can be seen as the

confidence of the classification (see Methods for additional details).

Figure 2 Orientation classification performance during the delay period in early visual

cortex and intraparietal sulcus (IPS) using a leave-one-run-out cross-validation

approach. In the combined early visual cortex (V1 – V3), both the contralateral and the

ipsilateral early visual areas contained orientation information during the delay period in no-

saccade condition. In the saccade condition, however, the orientation information significantly

degraded in the contralateral hemispheres. Nevertheless, orientation information in the

ipsilateral hemisphere after an eye movement was not higher than that before the saccade,

suggesting no remapping of VWM in the early visual areas. In the IPS, the classifier could

select the remembered orientation in both the contralateral and the ipsilateral hemisphere in

the no-saccade trials, in line with the findings in early visual cortex. In the saccade condition,

after the eye movement there was still a reliable orientation information in both the

contralateral and ipsilateral IPS, suggesting a consistent VWM representation in the IPS. Error

bars denote SEM. *No saccade – Contra*: No saccade condition, contralateral hemisphere; *No*

saccade – Ipsi: No saccade condition, ipsilateral hemisphere; *Saccade – Contra*: Saccade

condition, contralateral hemisphere; *Saccade – Ipsi*: Saccade condition, ipsilateral

hemisphere. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

No remapping of VWM in early visual cortex following eye movements.

During the no-saccade condition, activity patterns in the early visual areas (V1

- V3) contralateral to the grating location contained information about the maintained

orientation during the delay period (7.2 s - 10.8 s after onset of maintenance)

(**Figure 2**, dark green bar; $t_{(33)} = 3.652$, $p = 0.0009$, Cohen's $d = 0.63$; $BF_{10} =$

35.369). There was also weak evidence for some orientation information in the

ipsilateral hemisphere of V1 - V3 (**Figure 2**, dark orange bar; $t_{(33)} = 2.192$, $p =$

0.0356, Cohen's $d = 0.38$; $BF_{10} = 1.501$), and no compelling evidence for a

difference between the contralateral and ipsilateral hemisphere of V1 - V3 ($t_{(33)} =$

1.669, $p = 0.1047$, Cohen's $d = 0.32$; $BF_{10} = 0.642$). The same pattern was also

visible when inspecting the early visual areas separately (**Supplementary Figure 1**).

However, in the saccade trials, in which participants made an eye movement that

shifted the location of the remembered grating to the opposite visual field during the

delay period, the classifier was not able to distinguish between grating orientations in

V1 – V3 contralateral to the item location (**Figure 2**, light green bar; $t_{(33)} = 0.03$, $p =$

0.9765, Cohen's $d = 0.005$; $BF_{10} = 0.184$), and similarly there was no compelling

evidence for orientation information in ipsilateral V1-V3 (**Figure 2**, light orange bar;

$t_{(33)} = 1.905$, $p = 0.0655$, Cohen's $d = 0.327$; $BF_{10} = 0.92$). Again, there was no

difference in orientation information between the contralateral and ipsilateral

hemisphere of V1 – V3 ($t_{(33)} = 1.722$, $p = 0.09438$, Cohen's $d = 0.286$; $BF_{10} = 0.694$).

Crucially, there was a significant degradation of orientation information in the

contralateral hemisphere of the early visual cortex after an eye movement ($t_{(33)}$ =

2.748, p = 0.0096, Cohen's d = 0.478; BF_{10} = 4.452).

Although orientation information in the contralateral hemisphere was

degraded after the saccade, the remapping hypothesis of VWM predicts that

orientation information should become stronger in the ipsilateral hemisphere after the

eye movement. However, in contrast to this hypothesis, there was no difference in

terms of orientation information in the ipsilateral hemisphere after the eye movement

compared to when no saccade was made ($t_{(33)}$ = 0.072, p = 0.9431, Cohen's d =

0.017; BF_{10} = 0.184). Therefore, these results indicate that there is no remapping of

VWM following eye movements in the early visual areas.

Persistent VWM representations after an eye movement in the IPS.

Our findings revealed that VWM representations in early visual cortex during

maintenance were not remapped following eye movements. Bettencourt & Xu (2016)

have recently shown that while remembered orientation information was degraded

by irrelevant distractors in early visual cortex, VWM information remained available

in the superior intraparietal sulcus (IPS). We sought to test whether we could

observe similar results in our study. To this end, we also applied the leave-one-run-

out cross-validation method to IPS. First, we found that both the contralateral

(**Figure 2**, dark green bar; $t_{(33)}$ = 4.451, p = 0.0001, Cohen's d = 0.775; BF_{10} =

272.079) and ipsilateral IPS (**Figure 2**, dark orange bar; $t_{(33)}$ = 5.075, p < 0.0001,

Cohen's d = 0.883; BF_{10} = 1444.41) contained information about the remembered

orientation in no-saccade trials during delay period, as also observed in early visual

cortex. However, in contrast to the early visual cortex, the remembered grating

orientation could still be decoded from both the contralateral (**Figure 2**, light green

bar; $t_{(33)}$ = 3.13, p = 0.0036, Cohen's d = 0.545; BF_{10} = 10.276) and ipsilateral

(**Figure 2,** light orange bar; $t_{(33)}$ = 2.6, p = 0.0138, Cohen's d = 0.453; BF_{10} = 3.281)

IPS after an eye movement, indicating a persistent VWM representations in IPS.

These results also provided a neural evidence for supporting the preserved

behavioral performance after a saccade.

Finally, in order to ensure that the results are not dependent on the a priori but

arbitrarily chosen mask sizes of the ROIs, we repeated the analyses for ROIs of

sizes ranging from 20 to 250 voxels in step of 10 voxels in the areas V1, V2 and V3,

or ranging from 50 to 500 voxels in step of 20 voxels in the combined V1 – V3 and

IPS (**Supplementary Figure 2**). Results were qualitatively identical to those

mentioned above (**Figure 2** and **Supplementary Figure 1**) for almost entire range of

ROI sizes, indicating that our results do not depend on an arbitrary ROI size but

represent a robust pattern within the areas.

Discussion

In the current study, we investigated whether VWM representations are remapped

following eye movements and whether VWM information persists in the early visual

cortex and parietal cortex after the execution of a saccade. We found robust

encoding of maintained orientation information in the contralateral hemisphere of

early visual cortex. However, this information significantly degraded upon making a

saccade to the opposite hemifield and did not remap to the ipsilateral hemisphere in

early visual cortex. This suggests that there is no robust representation of VWM

representations across eye movements in the early visual cortex nor remapping of

VWM representations. Additionally, although VWM representations in early visual

cortex were impaired by the saccade during the retention period, this orientation

information remained reliably present in the IPS even after execution of a saccade.

These findings show that early visual cortex is susceptible to interference of eye

movements during a working memory period, while IPS appears less sensitive to this

interference.

Spatially specific encoding of VWM information in early visual cortex?

The issue of how VWM information is stored in early visual cortex has been debated

previously. While some researchers observed a retinotopically specific encoding of

items into VWM (Pratte and Tong 2014), Ester and colleagues (2009) observed a

non-spatially specific coding of VWM in early visual cortex. In their study, they could

decode the orientation information in early visual areas both contralateral and

ipsilateral to the stimulus location when participants were instructed to remember the

orientation presented only in one visual hemifield. However, although the

remembered grating was always located in one of the hemifields, participants could

potentially use a strategy in which they memorized the orientation of the grating

perifoveally instead of at its original location (Williams et al. 2008). Thus, it is unclear

whether their result reflected the true effect of the spatially global representations or

was caused by the lack of relevance of the spatial dimension in their task. To

incentivize participants to form a spatially specific visual working memory, we

inserted catch trials, in which the probe grating and the remembered grating were

horizontally offset by 1.2 visual degree in each run. Participants were instructed to

respond only when the probe location matched the location of the initially presented

grating. Therefore, participants had to maintain the orientation of the gratings at the

presented location in order to be able to successfully complete the task. Behavioral

results confirmed that participants indeed remembered the orientation stimuli in a

spatially specific way. Decoding performance in the early visual cortex (i.e., V1 – V3)

showed a numerically larger amount of orientation information in contralateral than

ipsilateral cortex, echoing earlier results (Pratte and Tong 2014) and suggesting a

spatially specific code. However, this difference was not statistically reliable, and

Bayesian analyses showed anecdotal evidence for a lack of difference (BF_{10} =

0.642). Thus, while our results may tentatively suggest some degree of retinotopic

specificity of VWM encoding, our data preclude any strong conclusion.

No remapping of VWM following eye movements.

Attentional or feature based remapping of visual input just prior to a saccade has

been well established from a variety of studies and is thought to maintain visual

stability across eye movements. For instance, Rolfs et al. (2011) found that attention

is predictively remapped to the future retinotopic location of an upcoming target,

even before the execution of the eye movement. In the current experiment, we asked

whether memory traces would also be updated with eye movements. However, we

found no evidence that visual working memory representations were remapped

following eye movements. At least two reasons should be emphasized to address

the discrepancy in remapping between attention and VWM. First, to survive in a

multifaceted and dynamically changing world, our attention needs to be selectively

shifted to the most important stimuli that are goal-relevant. In order to be competent

for such a complicated task, remapping of attention may be a crucial mechanism to

prepare information in advance. In contrast, the main function of working memory is

to temporarily retain information available for processing. While attention could be

flexibly remapped to different locations to extract information, storing working

memory information in one or several stationary regions could be potentially an

efficient way for retrieval. Second, in line with the arguments of 'activity-silent'

models of WM when compared to the persistent activity models of WM (Stokes

2015; Wolff et al. 2017), remapping of VWM after each eye movement could also be

energetically expensive, especially if VWM is maintaining an up-to-date information

of the external environment. Therefore, a stationary working memory representation

could be an ideal way for ecologically maintaining and manipulating information in

the brain.

Parietal cortex may subserve VWM representations after saccades.

In recent years, the parietal cortex has been repeatedly implicated in VWM

maintenance (Todd and Marois 2004; Christophel et al. 2015; Ester et al. 2015;

Bettencourt and Xu 2016), alongside the involvement of sensory areas (Harrison and

Tong 2009; Serences et al. 2009; Emrich et al. 2013; Rademaker et al. 2019). Our

results indicate that the early visual cortex is susceptible to the interference of eye

movements during the delay period, potentially limiting the role of the early visual

cortex in VWM. Similarly, Bettencourt and Xu (2016) observed that VWM

representations in early visual cortex were destroyed by the presentation of

distractors, while superior IPS exhibited persistent activity that was not impaired by

the presentation of intervening distractor stimuli. In our experiment, the eye

movements during the delay period potentially may also generate distracting new

input to the visual system, which could disturb the VWM representations. Intriguingly,

while VWM representations in early visual cortex were degraded following the eye

movement, these representations were still maintained in IPS. Behaviorally,

participants' memory performance was indistinguishable between the saccade and

no saccade trials. Therefore, it could be that the preserved behavioral performance

we observed is supported by these IPS representations, rather than the early visual

cortex representations. In a more recent study, Lorenc et al. (2018) found that the

VWM information could be reliably maintained only in early visual cortex without

distractors. However, these early visual cortex representations seem to transfer to

the IPS representations when distractors were presented, which suggesting a

flexible coding of VWM between the early visual cortex and the IPS depending on

the task demands. This shows that VWM information maintained in visual cortex is

susceptible to interference, in line with the idea that the early visual cortex may only

play a limited role in VWM storage.

In conclusion, our results indicate that VWM representations in early visual

cortex are degraded and not remapped after eye movements, but stably represented

in parietal cortex. These results suggest a limited role of early visual cortex in VWM

storage.

Acknowledgements: This work was supported by The Netherlands Organisation for

Scientific Research Vidi Grant 452-13-016 to F.P.d.L., the EC Horizon 2020 Program

ERC Starting Grant 678286 "Contextvision" to F.P.d.L, the James S.McDonnell

Foundation 220020373 to F.P.d.L, the China Scholarship Council (CSC;

201608330264) to T.H., and The Netherlands Organization for Scientific Research

(Veni Grant No. 016.Veni.195.435) awarded to M.E.. We thank David Richter and

Chuyao Yan for helpful comments and discussions of the manuscript.

Competing Interest: The authors declare no competing financial interest.

Author contributions: T.H. wrote the first draft of the paper; T.H., M.E., and

F.P.d.L. edited the paper; T.H., M.E., A.V., and F.P.d.L. designed research; T.H. and

A.V. performed research; T.H., M.E., and F.P.d.L. analyzed data.

References

Albers AM, Kok P, Toni I, Dijkerman HC, de Lange FP. 2013. Shared
622   Representations for Working Memory and Mental Imagery in Early Visual
623   Cortex. Curr Biol. 23:1427–1431.

624   Bettencourt KC, Xu Y. 2016. Decoding the content of visual short-term memory
625   under distraction in occipital and parietal areas. Nat Neurosci. 19:150–157.

626   Brainard DH. 1997. The Psychophysics Toolbox. Spat Vis. 10:433–436.

627   Christophel TB, Cichy RM, Hebart MN, Haynes J-D. 2015. Parietal and early visual
628   cortices encode working memory content across mental transformations.
629   NeuroImage. 106:198–206.

630   Curtis CE, D'Esposito M. 2003. Persistent activity in the prefrontal cortex during
631   working memory. Trends Cogn Sci. 7:415–423.

632   DeYoe EA, Carman GJ, Bandettini P, Glickman S, Wieser J, Cox R, Miller D, Neitz J.
633   1996. Mapping striate and extrastriate visual areas in human cerebral cortex.
634   Proc Natl Acad Sci. 93:2382–2386.

635   Dijkstra N, Ambrogioni L, Gerven MAJ van. 2019. Neural dynamics of perceptual
636   inference and its reversal during imagery. bioRxiv. 781294.

637   Durstewitz D, Seamans JK, Sejnowski TJ. 2000. Dopamine-Mediated Stabilization of
638   Delay-Period Activity in a Network Model of Prefrontal Cortex. J Neurophysiol.
639   83:1733–1750.

640   Emrich SM, Riggall AC, LaRocque JJ, Postle BR. 2013. Distributed Patterns of
641   Activity in Sensory Cortex Reflect the Precision of Multiple Items Maintained
642   in Visual Short-Term Memory. J Neurosci. 33:6516–6523.

643   Engel SA, Glover GH, Wandell BA. 1997. Retinotopic organization in human visual
644   cortex and the spatial precision of functional MRI. Cereb Cortex. 7:181–192.

645   Ester EF, Serences JT, Awh E. 2009. Spatially Global Representations in Human
646   Primary Visual Cortex during Working Memory Maintenance. J Neurosci.
647   29:15258–15265.

648   Ester EF, Sprague TC, Serences JT. 2015. Parietal and Frontal Cortex Encode
649   Stimulus-Specific Mnemonic Representations during Visual Working Memory.
650   Neuron. 87:893–905.

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A,
Killiany R, Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, Dale AM.
2002. Whole Brain Segmentation: Automated Labeling of Neuroanatomical
Structures in the Human Brain. Neuron. 33:341–355.

Goldman-Rakic PS. 1995. Cellular basis of working memory. Neuron. 14:477–485.

Harrison SA, Tong F. 2009. Decoding reveals the contents of visual working memory
in early visual areas. Nature. 458:632–635.

Haynes J-D, Rees G. 2006. Neuroimaging: Decoding mental states from brain
activity in humans. Nat Rev Neurosci. 7:523–534.

He T, Fritsche M, Lange FP de. 2018. Predictive remapping of visual features
beyond saccadic targets. J Vis. 18:20–20.

JASP Team. 2019. JASP (Version 0.10.2)[Computer software].

Kamitani Y, Tong F. 2005. Decoding the visual and subjective contents of the human
brain. Nat Neurosci. 8:679–685.

Lee MD, Wagenmakers E-J. 2014. Bayesian Cognitive Modeling: A Practical
Course. Cambridge University Press.

Linde-Domingo J, Treder MS, Kerrén C, Wimber M. 2019. Evidence that neural
information flow is reversed between object perception and object
reconstruction from memory. Nat Commun. 10:1–13.

Lorenc ES, Sreenivasan KK, Nee DE, Vandenbroucke ARE, D'Esposito M. 2018.
Flexible Coding of Visual Working Memory Representations during
Distraction. J Neurosci. 38:5267–5276.

Maris E, Oostenveld R. 2007. Nonparametric statistical testing of EEG- and MEG-
data. J Neurosci Methods. 164:177–190.

Melcher D. 2007. Predictive remapping of visual features precedes saccadic eye
movements. Nat Neurosci. 10:903–907.

Millman KJ, Aivazis M. 2011. Python for Scientists and Engineers. Comput Sci Eng.
13:9–12.

Norman KA, Polyn SM, Detre GJ, Haxby JV. 2006. Beyond mind-reading: multi-voxel
pattern analysis of fMRI data. Trends Cogn Sci. 10:424–430.

Oliphant TE. 2007. Python for Scientific Computing. Comput Sci Eng. 9:10–20.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M,
Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D,
Brucher M, Perrot M, Duchesnay É. 2011. Scikit-learn: Machine Learning in
Python. J Mach Learn Res. 12:2825–2830.

Pratte MS, Tong F. 2014. Spatial specificity of working memory representations in
the early visual cortex. J Vis. 14:22.

Prins N, Kingdom FAA. 2018. Applying the Model-Comparison Approach to Test
Specific Research Hypotheses in Psychophysical Research Using the
Palamedes Toolbox. Front Psychol. 9:1250.

Rademaker RL, Chunharas C, Serences JT. 2019. Coexisting representations of
sensory and mnemonic information in human visual cortex. Nat Neurosci.
22:1336–1344.

Rayner K. 1998. Eye movements in reading and information processing: 20 years of
research. Psychol Bull. 124:372–422.

Riley MR, Constantinidis C. 2016. Role of Prefrontal Persistent Activity in Working
Memory. Front Syst Neurosci. 9:181.

Rolfs M, Jonikaitis D, Deubel H, Cavanagh P. 2011. Predictive remapping of
attention across eye movements. Nat Neurosci. 14:252–256.

Savitzky Abraham, Golay MJE. 1964. Smoothing and Differentiation of Data by
Simplified Least Squares Procedures. Anal Chem. 36:1627–1639.

Serences JT. 2016. Neural mechanisms of information storage in visual short-term
memory. Vision Res. 128:53–67.

Serences JT, Ester EF, Vogel EK, Awh E. 2009. Stimulus-Specific Delay Activity in
Human Primary Visual Cortex. Psychol Sci. 20:207–214.

Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg
H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J,
Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM. 2004. Advances
in functional and structural MR image analysis and implementation as FSL.
NeuroImage, Mathematics in Brain Imaging. 23:S208–S219.

Sreenivasan KK, Curtis CE, D'Esposito M. 2014. Revisiting the role of persistent
neural activity during working memory. Trends Cogn Sci. 18:82–89.

Stokes MG. 2015. 'Activity-silent' working memory in prefrontal cortex: a dynamic
coding framework. Trends Cogn Sci. 19:394–405.

Todd JJ, Marois R. 2004. Capacity limit of visual short-term memory in human
posterior parietal cortex. Nature. 428:751–754.

Wandell BA, Dumoulin SO, Brewer AA. 2007. Visual Field Maps in Human Cortex.
Neuron. 56:366–383.

Wang L, Mruczek REB, Arcaro MJ, Kastner S. 2015. Probabilistic Maps of Visual
Topography in Human Cortex. Cereb Cortex. 25:3911–3931.

Watson AB, Pelli DG. 1983. Quest: A Bayesian adaptive psychometric method.
Percept Psychophys. 33:113–120.

Williams MA, Baker CI, Op de Beeck HP, Mok Shim W, Dang S, Triantafyllou C,
Kanwisher N. 2008. Feedback of visual object information to foveal retinotopic
cortex. Nat Neurosci. 11:1439–1445.

Wolff MJ, Jochim J, Akyürek EG, Stokes MG. 2017. Dynamic hidden states
underlying working-memory-guided behavior. Nat Neurosci. 20:864–871.

Supplementary information

732
733 Visual working memory representations in visual and parietal cortex do not remap

734 after eye movements

736 Tao He, Matthias Ekman, Annelinde R.E. Vandenbroucke and Floris P. de Lange
737
738

Supplementary Figure 1 Orientation classification performance during the delay period in early visual areas separately. Same analysis as in Figure 2 but performed in V1, V2 and V3 individually. The same pattern was observed as in Figure 2. Specifically, in no-saccade trials, orientation information was present in the contralateral hemisphere of V1 ($t_{(33)}$ = 3.701, p = 0.0008, Cohen's d = 0.64; BF_{10} = 39.988), V2 ($t_{(33)}$ = 2.460, p = 0.0193, Cohen's d = 0.43; BF_{10} = 2.48), and V3 ($t_{(33)}$ = 1.892, p = 0.0673, Cohen's d = 0.33; BF_{10} = 0.901), and also in the ipsilateral hemisphere of V1($t_{(33)}$ = 2.883, p = 0.0069, Cohen's d = 0.502; BF_{10} = 5.945) and V3 ($t_{(33)}$ = 3.147, p = 0.0035, Cohen's d = 0.548; BF_{10} = 10.688), but not V2 ($t_{(33)}$ = 0.612, p = 0.5447, Cohen's d = 0.107; BF_{10} = 0.219). In saccade trials, however, the classifier was not able to distinguish between grating orientations in the contralateral hemisphere of the V1 ($t_{(33)}$ = -0.174, p = 0.8627, Cohen's d = 0.03; BF_{10} = 0.186) and V3 ($t_{(33)}$ = 0.313, p = 0.7562, Cohen's d = 0.054; BF_{10} = 0.192), but not V2 ($t_{(33)}$ = 2.542, p = 0.0159, Cohen's d = 0.443; BF_{10} = 2.917), while the orientation information remained stable in the ipsilateral hemisphere of the V2 ($t_{(33)}$ = 2.306, p = 0.0276, Cohen's d = 0.395; BF_{10} = 1.848) and V3 ($t_{(33)}$ = 2.028, p = 0.0507, Cohen's d = 0.348; BF_{10} = 1.128), but not V1 ($t_{(33)}$ = 0.132, p = 0.8959, Cohen's d = 0.023; BF_{10} = 0.185). The orientation information in the contralateral hemisphere of the V1 was significantly degraded by an eye movement ($t_{(33)}$ = 2.765, p = 0.0093, Cohen's d = 0.647; BF_{10} = 4.612). Error bars denote SEM. *No saccade – Contra*: No saccade condition, contralateral hemisphere; *No saccade – Ipsi*: No saccade condition, ipsilateral hemisphere; *Saccade – Contra*: Saccade condition, contralateral hemisphere; *Saccade – Ipsi*: Saccade condition, ipsilateral hemisphere. *p < 0.05; **p < 0.01; ***p < 0.001.

**Supplementary Figure 2 Orientation classification performance during the
delay period over a range of ROI sizes using a leave-one-run-out cross-
validation method.** Same analysis as in Figure 2 but performed over a wide range
of ROI sizes. In areas V1, V2, and V3, we rerun the analysis for ROIs of sizes
ranging from 20 to 250 voxels in step of 10 voxels, while in areas V1 – V3 and IPS,
the analysis was repeated for ROIs of sizes ranging from 50 to 500 voxels in step of
20 voxels. The dashed vertical line is at the predefined ROI of 150 voxels in areas
V1, V2, and V3, or 250 voxels in areas combined V1 – V3 and IPS. The same
pattern of effects was found for almost entire range of ROIs. Interestingly, in the
combined V1 – V3, while WM information in the contralateral hemisphere was
reduced after an eye movement (dark green line vs. light green line), it was
persistently presented in the ipsilateral hemisphere for trials with and without a
saccade (dark orange line vs. light orange line) over all range of ROI sizes. In IPS,
however, no matter in the saccade or no-saccade condition, both the contralateral
and ipsilateral hemispheres showed consistent orientation information over the range
of ROIs, suggesting a robust VWM representation in this region.

Area	No saccade		Saccade	
	Contralateral	Ipsilateral	Contralateral	Ipsilateral
V1	0.264*** (t = 3.701, p = 0.0008; BF_{10} = 39.988)	0.165** (t = 2.883, p = 0.0069; BF_{10} = 5.945)	-0.013 (t = -0.174, p = 0.8627; BF_{10} = 0.186)	0.011 (t =0.132, p =0.8959; BF_{10} =0.185)
V2	0.166* (t = 2.460, p = 0.0193; BF_{10} = 2.48)	0.037 (t = 0.612, p = 0.5447; BF_{10} = 0.219)	0.159* (t = 2.542, p = 0.0159; BF_{10} = 2.917)	0.135* (t = 2.306, p = 0.0276; BF_{10} = 1.848)
V3	0.158 (t = 1.892, p = 0.0673; BF_{10} = 0.901)	0.158** (t = 3.147, p = 0.0035; BF_{10} = 10.688)	0.022 (t = 0.313, p = 0.7562; BF_{10} = 0.192)	0.114 (t = 2.028, p = 0.0507; BF_{10} = 1.128)
V1 – V3	0.226*** (t = 3.652, p = 0.0009; BF_{10} = 35.369)	0.118* (t = 2.192, p = 0.0356; BF_{10} = 1.501)	0.002 (t = 0.03, p = 0.9765; BF_{10} = 0.184)	0.112 (t = 1.905, p = 0.0655; BF_{10} = 0.92)
IPS	0.239*** (t = 4.451, p = 0.0001; BF_{10} = 272.079)	0.293*** (t = 5.075, p = 0.0000; BF_{10} = 1444.41)	0.184** (t = 3.13, p = 0.0036; BF_{10} = 10.276)	0.177* (t = 2.6, p = 0.0138; BF_{10} = 3.281)

Supplementary Table 1 Decoding distance over 7.2 – 10.8 seconds after the onset
of the delay period for each ROI and post-hoc t-test of significant differences (see
Figure 2). Dark grey cells indicate that there was a significant effect of saccade and
hemisphere condition in that ROI, light grey cells represent the effect only reached a
marginal significance, while cells indicate that there was no significant effect in that
ROI. *p < 0.05; **p < 0.01; ***p < 0.001.