Dormancy-to-death transition in yeast spores occurs due to gradual loss of gene-expressing ability

Théo Maire[1,2], Tim Allertz[1,2], Max A. Betjes[1,2], and Hyun Youk[1,3,4,5*]

[1]Kavli Institute of Nanoscience, Delft, The Netherlands

[2]Department of Bionanoscience, Delft University of Technology, Delft, The Netherlands

[3]CIFAR, CIFAR Azrieli Global Scholars Program, Toronto, ON, Canada

[4]Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, MA, USA.

[5]Program in Systems Biology, University of Massachusetts Medical School, Worcester, MA, USA.

[*]Corresponding author. E-mail: hyun.youk@umassmed.edu

Note: Small figures with captions are embedded within the text. Larger, full page versions of the figures appear after the main text and are preceded by their captions.

Short summary (256 characters with space)

This study identifies systems-level quantities that decay during dormancy in *Saccharomyces cerevisiae* spores and thereby reveals the meaning of ageing for dormant yeast spores and shows that they die when their gene-expressing ability is irreversibly lost.

Highlights

- For a given glucose concentration, a dormant yeast spore has a well-defined probability of germinating ("germination ability").
- A spore's germination ability positively correlates with its RNAP I-III levels and the gene-expression (GFP) level it can realize when the expression is forced without nutrients.
- Ageing during dormancy means gradual decreases in germination ability, RNAP levels, and GFP-level realizable when expression is forced.
- Spores die after sufficiently losing gene-expressing ability and drugs that inhibit gene expression during dormancy shorten spores' lifespans (e.g., from months to a day).

ABSTRACT

Dormancy is colloquially considered as extending lifespan by being still. Starved yeasts form dormant spores that wake-up (germinate) when nutrients reappear but cannot germinate (die) after some time. What sets their lifespans and how they age are open questions because what processes occur - and by how much - within each dormant spore remains unclear. With single-cell-level measurements, we discovered how dormant yeast spores age and die: spores have a quantifiable gene-expressing ability during dormancy that decreases over days to months until it vanishes, causing death. Specifically, each spore has a different probability of germinating that decreases because its ability to - without nutrients - express genes decreases, as revealed by a synthetic circuit that forces GFP expression during dormancy. Decreasing amounts of molecules required for gene expression - including RNA polymerases - decreases gene-expressing ability which then decreases chances of germinating. Spores gradually lose these molecules because they are produced too slowly compared to their degradations, causing gene-expressing ability to eventually vanish and, thus, death. Our work provides a systems-level view of dormancy-to-death transition.

(175 words)

INTRODUCTION

When starved of nutrients, microbes can become dormant by forming spores (Rittershaus *et al,* 2013; Neiman *et al,* 2011; Stragier & Losick, 1996; Errington, 2003; Lennon & Jones, 2011; Nachman *et al,* 2007; Chu *et al,* 1998; Eldar *et al,* 2009; Suel *et al,* 2006; Suel *et al,* 2007). Dormancy is a state in which gene expression, metabolism, and all other cellular processes are thought to have nearly or completely ceased. But exactly to what extent each process of life has halted inside dormant spores remain ambiguous as attaching firm numbers to them has been challenging due to the barely detectable levels of activities inside dormant spores. Entering dormancy is a multi-step process that does not necessarily end with the spore formation. In fact, after sporulation, spores of some species are not dormant right away. For example, *Bacillus subtilis* spores - a model for prokaryotic spores - undergoes a week-long "ageing" process during which they are metabolically active, and they actively degrade or produce key transcripts to tailor their dormancy to the environmental temperature during this process (Segev *et al,* 2012). The spores of budding yeast, *Saccharomyces cerevisiae* - a model for eukaryotic spores - takes between a few hours to a few days to fully enter dormancy (Brengues *et al,* 2002; Thacker *et al,* 2011). During this time, the not-yet-dormant yeast spores express various genes - such as those involved in completing the spore wall - before gradually turning off gene expressions, at the end of which the spores are considered to be dormant (Brengues *et al,* 2002; Thacker *et al,* 2011). After entering dormancy, both bacterial and yeast spores are thought to have greatly reduced metabolism and vanishingly low, if any, genome-wide expression levels (Rittershaus *et al,* 2013; Segev *et al,* 2012; Brengues *et al,* 2002). Supporting this view for yeast spores are the recent discoveries of proteins and messenger RNPs (e.g., mRNAs bound to translational machineries) becoming inactive by aggregating into macromolecular structures inside dormant yeast spores and other fungal spores (Laporte *et al,* 2008; Petrovska *et al,* 2014). The resulting solid-like, glassy cytoplasm - being packed with these aggregates - greatly hinders proteins' movements and enzymatic activities (Ablett *et al,* 1999; Cowan *et al,* 2003; Dijksterhuis *et al,* 2007; Parry *et al,* 2014; Joyner *et al,* 2016; Munder *et al,* 2016). Taken together, previous findings support the widely accepted view that dormant yeast spores have nearly ceased all their activities (Rittershaus *et al,* 2013). But technical and conceptual difficulties lie in the phrase, "nearly ceased". Technically, the nearly ceased activities inside dormant yeast spores might, for instance, occur too slowly so that many measurement techniques would lack the required sensitivity to detect those activities, especially in individual dormant spores rather than in lysates of many spores. On the conceptual side: given that dormant yeast spores already appear still and that, as time passes without nutrients, they eventually die - meaning that nutrients can no longer revive

them - an intriguing and underexplored challenge is explaining how yeast spores age over time

to transition from being dormant (i.e., alive) to dead (i.e., what sets a spore's lifespan?).

Addressing this question would deepen our understanding of dormancy - which remains either

poorly understood or misunderstood for many species as the emerging discoveries show (Laporte

et al, 2008; Petrovska *et al,* 2014) - and the meaning of cellular death. We sought to address this

question in the context of yeast spores.

To address our question, we first reasoned that a dormant yeast spore's lifespan is likely

encoded by amounts of its stored intracellular factors (e.g., specific proteins and RNAs) which are

required for restarting replicative life. Their depletions below some values may cause spores to

die. For dormant spores of some species, researchers have identified several intracellular factors

that affect their revival (Segev *et al,* 2012; Dworkin & Shah, 2010; Mutlu *et al,* 2018; Donnini *et*

al, 1988; Geijer *et al,* 2012; Herman & Rine, 1997; Sturm & Dworkin, 2015; Sinai *et al,* 2015). For

example, alanine is more likely to revive *B. subtilis* spores that have more alanine dehydrogenase

stored in them than spores that stored less of it (Mutlu *et al,* 2018). But it is unclear how the known

and as-yet-unknown intracellular factors that affect revivability - perhaps the numbers of stored

ribosomes and RNA polymerases - collectively affect a dormant yeast or bacterial spore's

lifespan, which can be months or decades (Nagtzaam & Bollen, 1994) and thus vastly longer than

typical biomolecules' lifetimes. A brute-force way to address this question is identifying all

intracellular factors that affect spores' revival, then measuring how the amounts of all those factors

may change over time within a dormant spore, and then deducing when the spore dies, thereby

determining all combinations of intracellular factors and their amounts that are necessary for

sustaining dormancy. But this approach remains elusive because, for one, it requires using a

current snapshot of numbers (i.e., amounts of intracellular factors now) to predict a capacity for

achieving a complex future behavior - whether the dormant spore still has an ability to wake-up if

nutrients were to reappear. Addressing our question requires circumventing this difficulty.

As we will show, we found a single, readily measurable, systems-level metric - ability for

a dormant yeast spore to express genes - that reveals how a dormant yeast spore eventually

dies. In the process, we propose here a view of dormancy as a precisely measurable, continuous

quantity that a yeast spore loses over time as it approaches death. Our study begins by giving

various glucose concentrations to dormant yeast spores. From this, we discovered that not all

yeast spores wake-up (germinate) to re-enter a replicative life and that those that do not germinate

are primed - they undergo accelerated germinations if they encounter more glucose hours-to-

days later. These phenomena, which were previously unnoticed, led us to investigate why only a

fraction of genetically identical, dormant yeast spores in a population germinate despite having

ample glucose. To investigate this, we used a synthetic circuit to induce, in dormant yeast spores,

expression of a gene (*GFP*) that plays no role in germination. We found that one can induce GFP

expression in dormant yeast spores that are in water without any nutrients (no extracellular amino

acids and glucose) and that, surprisingly, the inducible GFP level in a dormant spore tells us the

probability of that spore germinating for a given concentration of glucose. Strikingly, we found that

dormant yeast spores, when induced, can express genes at levels that are as high as those of

vegetative yeasts. To our knowledge, we provide here one of the first evidences and

quantifications of active transcription in intact, dormant yeast spores at a single-cell resolution.

We found that the inducible GFP level in a dormant yeast spore is strongly and positively

correlated with the amounts of all three RNA polymerases - RNAP I-III - that the spore has but is

virtually uncorrelated with the amount of ribosomal RNA, which is a proxy for the amount of

ribosomes in spores. Having identified these links, our paper proceeds to show how key quantities

- gene-expression ability, amount of RNA polymerases, and ability to germinate - decrease over

time during dormancy, up until death. This led us to quantitatively determine when dormancy ends

(death occurs) for a yeast spore. Putting the results together, our paper ends by formulating a

new conceptual model that explains how yeast spores gradually lose dormancy and die: key

molecules that are required for gene expression, including RNA polymerases, are gradually lost

during dormancy because spores cannot produce them faster than they degrade - we measured

firm numbers (tens of days) for these extremely slow processes. These molecules eventually

reach below some threshold amounts after which the spore has permanently lost the ability to

express genes, meaning that even ample glucose cannot restore the gene-expressing ability.

Since, after receiving glucose, spores must express genes in order to build a new cell that will

bud off the spore (i.e., germinate), the permanent loss of gene expressing ability at the final

moments of dormancy marks the spore's death.

RESULTS

Not all genetically identical spores germinate despite encountering ample glucose

We began by re-examining the conventional test for determining whether a yeast spore is dormant

or dead. The test involves giving ample glucose to yeast spores and then observing whether they

germinate or not (Brengues *et al,* 2002; Joseph-Strauss *et al,* 2007) (Fig. 1A).

Figure 1 - Glucose germinates only a fraction of yeast spores.
(A) Conventional test to determine whether a spore bag (i.e., ascus) containing four haploid spores, in the absence of any external nutrients (blue box), is dormant or dead. Green box: outcome if dormant. Red box: outcome if dead. **(B)** Left: Filmstrip of a time-lapse movie in which a 2%-glucose was added at the beginning of the movie (10 minutes between frames). A spore bag is counted as having germinated at the moment that at least one budding cell emerges from the spore bag (white arrows). Right: Time taken by each "wild-type" spore bag to germinate in the time-lapse movie. n = 137 spore bags from a representative time-lapse movie. **(C)** Percentage of wild-type spore bags that germinated as a function of time after adding glucose to the minimal medium. Different colors represent different glucose concentrations (from 0.0002% to 2%). n = 3; error bars are s.e.m. **(D)** Average time taken to germinate (blue circles) and the total percentage of spore bags that germinated (red squares), both 16 hours (960 minutes) after receiving glucose (i.e., the plateau values for each color in Fig. 1C). Both are functions of glucose concentration. n = 3; error bars are s.e.m.

If the spore germinates, then it is considered to have been dormant whereas if it does not, then it

is considered to have been dead. This test, however, does not reveal why a spore that does not

germinate is dead in the first place and when it died. For instance, the test cannot distinguish

between a spore that died *while* it was being formed versus a spore that was alive (i.e., dormant)

after forming but died *during* its dormancy (and why and when the death occurred). Motivated by

this deficiency, we re-examined the test by asking whether all yeast spores can indeed germinate

after receiving ample glucose (Fig. 1A). Starving a laboratory-standard ("wild type") homozygous

diploid yeasts caused them to form genetically identical, haploid spores. Specifically, each diploid

cell formed a single "spore bag" (i.e., ascus) that contained four genetically identical haploid

spores (Fig. 1A). We incubated a population of these spore bags in a "minimal medium" - this has

all the essential amino acids but no glucose - and, as a modification to the conventional test,

supplemented the minimal medium with a less-than-saturating concentration of glucose (less than

2%) instead of the usual, 2%-glucose. We used a wide range of glucose concentrations that

spanned a 10,000-fold range, from 0.0002% to 2%. For each glucose concentration, we used a

wide-field microscope to track individual spore bags and count how many of them germinated -

that is, how many spore bags contained at least one spore that germinated (i.e., replicated) - as

a function of time after we added glucose (Fig. 1B and Appendix Fig S1). We focused on spore

bags instead of individual spores that are within each spore bag because we sought to assess

whether a diploid cell successfully formed at least one spore that germinates. With a saturating

glucose concentration (2%), nearly every spore bag in the population germinated (Fig. 1C). But

with lower glucose concentrations (i.e., less than ~0.01%), a noticeable percentage of spore bags

in the population - about 10% or more - did not germinate regardless of how many hours we

waited after adding the glucose (Fig. 1C). The percentage of spore bags that germinated changed

in a sharp, step-like (sigmoidal) manner as a function of the glucose concentration (Fig. 1D - red

points), with the change in step located at a glucose concentration of ~0.003% (i.e., at this

concentration, ~50% of spore bags germinate). In contrast, the average time taken to germinate

only weakly depended on the glucose concentration, increasing by at most 2-fold despite a

10,000-fold decrease in the glucose concentration from 2% to 0.0002% (Fig. 1D – blue points).

This indicates that glucose weakly affects the speed of germination. Importantly, the germinations

did not stop because the spores ran out of glucose for any of the glucose concentrations because

when we measured how much glucose was left in the minimal medium after all the germinations

stopped (i.e., ~600 minutes after adding glucose for all glucose concentrations), we always found

a large fraction of the original glucose remaining in the medium: the glucose concentration hardly

decreased even for the lowest starting glucose concentrations that we used (e.g., 0.002% and

0.001%) (Appendix Fig S2). Moreover, we observed that vegetative, diploid cells of the wild-type

strain - the same strain as the spores - could replicate multiple times even with the lowest glucose

concentration (i.e., 0.0002%) (Appendix Fig. S3). Hence, even the lowest glucose concentration

was ample enough for a vegetative cell to divide multiple times. These results establish that yeast
spores do not necessarily germinate even when there is ample glucose.

Spores that do not germinate after encountering ample glucose are not necessarily dead
To explain why some spores do not germinate even with ample glucose, we considered two
possibilities. One was that the spore bags that did not germinate (i.e., "un-germinated spore
bags") died while trying to germinate. The other possibility was that the un-germinated spore bags
were still able to germinate and thus were still alive (i.e., dormant). We distinguished these two
possibilities by repeating the above experiments but now by adding glucose in two steps (Fig.
2A). First, we gave the spores a relatively low concentration of glucose. We then waited, typically
~16 hours (~1000 minutes), after which no more germinations occurred. Then, we added more
glucose to increase the total glucose concentration and then observed if this led to any more
germinations. We found that some of the spore bags that did not germinate the first time were
germinating after receiving more glucose (Fig. 2B and Appendix Figs S4-S5). Yet, some of the
spore bags still did not germinate after receiving the second batch of glucose if the resulting, final
concentration was not the saturation value, 2% (Fig. 2C - top panel). For example, when the final
concentration was 0.002%, nearly 60% of the spore bags in the population remained un-
germinated. Intriguingly, this 60% is close to the percentage that would have remained as un-
germinated if we had given the 0.002%-glucose all at once instead of in two steps (compare Fig.
2C with Fig. 1C). This suggests that each spore bag may be pre-programmed to germinate for
certain glucose concentrations. Accordingly, nearly every spore bag eventually germinated if the
second batch of glucose increased the total glucose concentration to 2% (Fig. 2C - bottom panel).
These results establish that spores that do not germinate after encountering ample glucose are
not necessarily dead.

Spores that do not germinate after encountering low glucose concentrations are primed
so that they accelerate their germinations later
To better understand why only some spore bags germinated for a given glucose concentration,
we examined whether the un-germinated spore bags had any measurable response to the
glucose that they first encountered.

Figure 2 - Un-germinated spores, primed by a low glucose concentration, germinate faster upon encountering more glucose a few days later.

(A) Wild-type spores are first incubated in a low glucose concentration ("1st [glucose]") before we add more glucose later to increase the glucose concentration to "final [glucose]". **(B)** For experiment in (A), number of spore bags germinated within each binned time. First glucose concentration is 0.0005% (from 0 to 16 hours) (green bars) and the final concentration is 0.002% (from 16 to 32 hours) (orange bars). n = 143 spore bags (representative data). **(C)** For the experiment in (A), percentage of spore bags that germinated as a function of time since the first glucose addition (1st and final concentrations as indicated). More glucose added 1000 minutes after the first glucose (purple vertical line). n = 3; error bars are s.e.m. **(D)** For the experiment in (A), average time taken for a spore bag to germinate - denoted $\Delta\tau$ in bottom panel of (C) - after stepping up the glucose concentration. We varied the first glucose concentration, but the final concentration was always 2%. n = 3; error bars are s.e.m. **(E)** Spore bags that do not germinate after encountering the first glucose are "primed" to germinate faster upon encountering more glucose. **(F)** For the experiment in (A), average time ($\Delta\tau$) taken for a spore bag to germinate after stepping up the glucose concentration (final glucose concentration is 2%), plotted as a function of the first, low glucose concentration. Different colors represent different times at which we added more glucose after the first addition: 16 hours (yellow), 48 hours (blue) and 96 hours (red). "Relative $\Delta\tau$" is the average time $\Delta\tau$ divided by the $\Delta\tau$ of spore bags that were incubated in minimal media without any glucose (0%) for the same duration of time (after which they received a 2%-glucose). n = 3; error bars are s.e.m.

(G) Heat map showing transcriptome-wide changes in un-germinated spores at 0, 16, 48, and 96 hours after being primed by a 0.002%-glucose (via RNA-Seq). Appendix Table S1 lists the genes for each transcriptional module (each row in the heat map). Also see Appendix Fig S8. For each transcriptional module, we first divided the expression level of each gene in that module by its expression level at 0-hours - this yields "normalized expression level" for that gene for both primed and unprimed spores (unprimed spores were in minimal media without glucose for 0, 16, 48, and 96 hours). We averaged these values over all genes in a transcriptional module to obtain "normalized expression (primed)" and "normalized expression (unprimed)" for each transcriptional module. Colors represent ratio of these two values, averaged over three biological replicates ($n = 3$).

When we added glucose in two steps so that the final concentration was 2% (Fig. 2A), we found

that the spore bags took less time to germinate, in response to the second batch of glucose, than

they would have if they had received the entire 2%-glucose all at once (Fig. 2D). Specifically, if a

spore bag was in a minimal medium without any glucose for 16 hours and then encountered a

2%-glucose, it needed an average of ~200 minutes to germinate. But this time decreased by

about half (i.e., to ~120 minutes) if a spore bag was first in a minimal medium with a low glucose

concentration - ranging from 0.0002% to 0.002% - for 16 hours and then received more glucose

so that the final concentration was 2% (Fig. 2D). Thus, encountering a very low amount of glucose

"primes" some spores so that, upon encountering a saturating level of glucose at a later time, they

would germinate faster - up to two times faster on average - compared to spores that did not

previously encounter any glucose (Fig. 2E). Furthermore, when we primed the spores with a very

low glucose concentration and then waited between 16 hours to 4 days before increasing the

glucose concentration to 2%, the spores were still primed - they germinated faster compared to

spores that were kept in minimal medium without any glucose for the same amount of time. They

were primed for up to two days after we added the first glucose but no longer primed four days

after we added the first glucose (Fig. 2F and Appendix Fig S6). Thus, primed dormancy lasts for

and decays over days.

Transcriptome-wide view of primed dormancy

Before turning to the question of what causes only some spore bags to germinate for a given

glucose concentration, we sought to uncover gene expressions that underlie the primed

dormancy. We first primed the spores by incubating them with a low glucose concentration

(0.002%) for either 16 hours, 1 day, 2 days, or 4 days. We then used zymolyase, as is the standard

(Coluccio *et al,* 2004), to isolate the un-germinated spores from the surrounding vegetative cells

(Appendix Fig S7) and then analyzed their transcriptomes with RNA-seq. As a control, we also

analyzed the transcriptome of un-primed spores, which were incubated in minimal media without

glucose for the same amounts of time as the primed spores. Following an insightful previous study

(Joseph-Strauss *et al,* 2007) that analyzed the yeast spores' transcriptome over several hours

after they received a 2%-glucose, we grouped multiple genes together into a set, called a
"transcriptional module" (Joseph-Strauss *et al,* 2007; Ihmels *et al,* 2002), if those genes
participate in the same process (e.g., protein synthesis) (Appendix Table S1). For both the primed
and un-primed spores, we averaged the expression levels of all genes in a given module to obtain
one expression value for that module. For six of nine transcriptional modules, the primed spores
had higher expression values than the un-primed spores after 16 hours and 48 hours of
incubations. But both types of spores had nearly the same expression value after four days of
incubation (Fig. 2G - last six rows and Appendix Fig S8). This trend mirrors the trend followed by
the average time taken by primed spores to germinate (i.e., accelerated germinations up to 48
hours after being primed but no accelerated germinations after four days) (Fig. 2F). Two
transcriptional modules showed this trend in a particularly pronounced manner. One of them is
the module for mating - haploid cells mate after germinating (Fig. 2G - seventh row). The other is
the module for transitioning from cell cycle's G2-phase to mitosis - a crucial final step of
germination (Fig. 2G - last row and Appendix Fig S8). Intuitively, these results make sense since
one expects that getting ready for mitosis and mating would accelerate germinations. Together,
these results establish that very low glucose concentrations can trigger transcriptome-wide
changes in un-germinated spores to prime them, so that they can germinate faster when they
encounter more glucose hours-to-days later.

Hypothesis on why not every yeast spore germinates with low glucose concentrations
Although we now understand how un-germinated spores respond after receiving glucose, we
have not addressed our original question: what determines, in the first place, which spore bags
germinate? We hypothesized that each diploid, vegetative yeast forms a spore bag with a distinct
"internal state". Many intracellular factors may define an internal state, including amounts of ATPs
or ribosomes that are stored inside the spores or combinations of these and other molecules that
are stored inside spores. We then hypothesized that, for each glucose concentration, only some
of the spore bags have the "right" internal states that enable germination. A difficulty in testing
this hypothesis is that in our experiments thus far, we gave glucose to spores *and then* observed
their subsequent actions, including for discovering primed dormancy. But in such experiments,
due to all measurements occurring *after* the spores receive glucose, we cannot infer the spores'
internal states that existed *before* they encountered glucose since glucose likely changes the
internal states. Therefore, we sought to manipulate the internal states without giving glucose. In
particular, we reasoned that depleting any internal resources (e.g., ATPs or amino acids) that are

stored inside spores *before* adding glucose would either decrease - or alter in more complex ways
- the percentage of spore bags that germinate for each glucose concentration.
307
308
Synthetic circuit to induce *GFP* in dormant yeast spores without any nutrients
To test our hypothesis, we built a synthetic gene-circuit in vegetative diploid yeasts so that
doxycycline - a small inducer molecule that easily diffuses through the yeast's cell wall - would
cause the cells to produce the Green Fluorescent Protein (GFP). In vegetative yeasts, this
synthetic circuit functions in such a way that increasing the doxycycline concentration increases
the GFP expression. We formed spores out of these engineered diploid cells (Fig. 3A). We
reasoned that if doxycycline can induce GFP expression in these spores without any nutrients
(i.e., in water), then we might deplete the stored resources inside the spores and thereby alter the
percentage of spore bags that germinate for a given glucose concentration. But it was unclear
whether it was even possible to induce GFP expression or any arbitrary gene in dormant yeast
spores without nutrients. For one, if gene induction were possible, then it is unclear why,
apparently, almost all gene expressions are suppressed in dormant yeast spores (Rittershaus *et*
al, 2013) since being able to induce GFP expression would mean that there must be active RNA
polymerases and ribosomes, and chromosomal regions that are accessible to them. Moreover,
recent studies established that starved vegetative yeasts, dormant yeast spores, and other
dormant fungal spores have a solid-like, glassy cytoplasm that is packed with macroscopic
aggregates of proteins and mRNPs (mRNAs bound by proteins) that inhibit gene-expression
machineries and movement of molecules inside the cytoplasm (Laporte *et al,* 2008; Petrovska *et*
al, 2014). By aggregating, proteins needed for metabolism and gene expression would be
inactivated as well (Ablett *et al,* 1999; Cowan *et al,* 2003; Dijksterhuis *et al,* 2007; Parry *et al,*
2014; Joyner *et al,* 2016; Munder *et al,* 2016). Indeed, to our knowledge, there has not yet been
a direct, single-cell-level observation of gene expression occurring in yeast spores during
dormancy. A previous, bulk-level (population-level) study (Brengues *et al,* 2002) has shown, from
lysates of many yeast spores, that two genes - *PGK1* (involved in gluconeogenesis) and *SPS100*
(involved in forming spore walls during spore formation) - are expressed but that they turn off a
few days after sporulation (i.e., while the spores enter dormancy). But it remains unclear whether
their expressions completely cease or remain at low, non-zero levels during dormancy. This is
because these bulk-level measurements were based on finding ribosomes bound to mRNAs after
lysing populations of yeast spores, meaning that the measurements cannot clearly distinguish
between zero and very low expression levels.

Figure 3 - Inducing GFP expression during dormancy with synthetic circuit shows that spores that can realize higher GFP level (higher GFP inducibility) are more likely to germinate (higher germination ability) for any glucose concentration.
(A) Synthetic gene-circuit that constitutively expresses a transcription factor, rtTA (with ADH1-promoter) and an inducible promoter (TET-promoter) controlling *GFP* expression. Increasing doxycycline increases GFP expression. **(B)** Engineered spore bags (shown in (A)) transcribe and translate GFP in water (without any nutrients) with 25 μg/ml of doxycycline (top row) and in a saline solution (PBS) with 50 μg/ml of doxycycline (bottom row). Snapshots of GFP expression shown 22 hours after adding doxycycline. **(C)** GFP levels of individual spore bags (grey curves) over time (measured every 10 minutes with a wide-field epifluorescence microscope) after incubation in PBS with 10 μg/ml of doxycycline (top panel: *n* = 104 spore bags) and 100 μg/ml of doxycycline (bottom panel: *n* = 150 spore bags). **(D)** Engineered spore bags (shown in (A)) were first incubated for 22 hours in either PBS without any doxycycline or with 100 μg/ml of doxycycline. Next, we transferred them to minimal medium with various glucose concentrations. Plot shows the total percentage of the engineered spore bags that germinated (measured 20 hours after incubating with glucose) for spores pre-incubated in PBS without doxycycline (black points) and in PBS with 100 μg/ml of doxycycline (orange points). *n* = 3; error bars are s.e.m. **(E)** Top: Percentage of spore bags with the same GFP level ("GFP inducibility") from (D) that germinated after receiving a 0.001%-glucose. Percentages are averaged over all spore bags with the same binned GFP level (in the histogram below). Representative data shown (*n* = 145 spore bags in a population). **(F)** Germination landscape: colors represent the probability that a spore bag with a particular steady-state GFP level ("GFP inducibility") germinates for each glucose concentration (i.e., data in top panel of (E) represents a single row of this heat map). To measure each pixel, as in the experiment described in (D), we incubated spore bags in PBS with 100 μg/ml of doxycycline for 22 hours before adding glucose ([glucose] as indicated along the rows). Columns indicate GFP inducibility of a spore bag, measured at 22 hours after adding the doxycycline. Each pixel is an average over 3 replicate populations (*n* = 3). **(G)** Given a spore bag, its GFP inducibility is a read-out of the probability to

germinate ("germination ability") for each glucose concentration and the minimal glucose concentration required to guarantee germination. Spore bags with lesser gene-expressing abilities without nutrients require more glucose to germinate.

Moreover, the ribosomes bound to mRNAs may be from the macroscopic aggregates that formed
before the spores entered dormancy (Laporte *et al,* 2008; Petrovska *et al,* 2014), which may
disable translation of those ribosome-bound mRNAs (Ablett *et al,* 1999; Cowan *et al,* 2003;
Dijksterhuis *et al,* 2007; Parry *et al,* 2014; Joyner *et al,* 2016; Munder *et al,* 2016). Adding to the
ambiguity is the fact that a bulk-level study revealed a depletion of a minute fraction of radioactive
uracil and methionine from an extracellular medium by a dense population of yeast spores
(Brengues *et al,* 2002). This finding indicates that transcription (proxied by the depleted uracil)
and translation (proxied by the depleted methionine) may be occurring in dormant yeast spores,
though these are indirect measurements since they did not directly visualize gene-expression
dynamics inside individual, intact spores. Time-lapse imaging of gene expression in individual
yeast spores - as we will show with our synthetic circuit - would help in resolving these ambiguities
and subtleties of gene regulations in dormant yeast spores.
352
Before testing the hypothesis posed in the previous section, we investigated whether GFP
induction in dormant spores is even possible by incubating the engineered spores with
doxycycline in either water or saline solution (PBS). Surprisingly, we discovered that doxycycline
fully induced GFP expression in these spores (Fig. 3B and Appendix Figs S9-S10). Varying the
doxycycline concentration tuned the spores' GFP levels over the same, wide range of values as
in vegetative cells that also have the synthetic circuit (Appendix Fig. S9-S11). We found that both
the rate of GFP production and the final (steady state) level of GFP widely varied among spore
bags of the same population (Fig. 3C and Appendix Fig S12). The most striking feature was that
all spore bags expressed GFP very slowly – GFP levels plateaued at steady-state values after
~20 hours of doxycycline whereas the vegetative yeasts needed ~8 hours of doxycycline for the
GFP level to plateau (Appendix Fig S11). But a more puzzling discovery was that the spores' GFP
levels stabilized at steady-state values in the first place. This is puzzling because the spores were
not dividing and hence their GFP - a highly stable protein - could not be diluted away by cell
divisions. In replicating (vegetative) yeasts, highly stable proteins such as GFP reach steady-state
levels because their production rate matches the cell-division rate. After the spores' GFP levels
reached steady-state values, we incubated the spores in PBS without doxycycline for two days
during which their GFP levels remained virtually unchanged (Appendix Fig S13). This means that
GFP levels reached steady-state values in dormant spores because the spores stopped

producing GFP after a day, despite the saturating concentration of doxycycline (100 μg/ml) still

being present in the medium after a day, with barely detectable degradations (Appendix Fig S14).

Together, these results provide a direct proof, with intact spores, that one can fully activate - to

the same level as in vegetative yeasts - transcription and translation of an arbitrary gene such as

GFP in dormant yeast spores without any nutrients. These results also establish that gene

expression in dormant yeast spores can exhibit starkly different dynamics when compared to

vegetative yeasts, such as vastly slower timescales and gene regulations.

Inducing GFP expression in dormant yeast spores does not alter percentages of spore

bags that germinate

Returning to our hypothesis that depleting yeast spores' internal resources through GFP induction

hinders germination, we incubated spores in PBS with a saturating concentration of doxycycline

(100 μg/ml) for 24 hours, after which the GFP levels plateaued at steady-state values (Fig. 3C

and Appendix Fig S13). We then washed away the doxycycline and then incubated the spores in

a minimal medium with a fixed concentration of glucose. Then, with time-lapse microscopy, we

tracked individual spore bags to determine how many spore bags germinated. For comparison,

we used the same method to determine how many spore bags without GFP induction germinated

- these spores were incubated for 24 hours in PBS without doxycycline and then in a minimal

medium with the same concentration of glucose as the GFP-induced spores (Fig. 3D and

Appendix Fig S15). For every glucose concentration that we examined, we found that inducing

GFP expression did not appreciably alter the percentage of spore bags that germinated and that

it also did not appreciably alter the average times taken by the spores to germinate (Appendix Fig

S16). Thus, our hypothesis is incorrect: expressing *GFP* does not alter the spores' ability to

germinate. While it may now appear that we are "back to square one" - since we still have not yet

uncovered what causes only some of the spore bags to germinate for a given glucose

concentration - we discovered the answer, as we explain in the next section, when we examined

the GFP levels of individual spore bags rather than the average GFP level of the entire spore

population as we have just done.

GFP level that a spore can realize (GFP inducibility) encodes the probability of germinating

(germination ability) for each glucose concentration

In the previous experiment, we measured the steady-state GFP levels just before the spores

encountered glucose. This led to the finding that spore bags that produced more GFP - spore

bags with higher "GFP inducibility" - were more likely to germinate (Fig. 3E and Appendix Figs

S17-S18). For example, after encountering a 0.001%-glucose, nearly 100% of the spore bags

with the highest, steady-state GFP levels (highest GFP inducibility) germinated whereas, in the

same population, only ~10% of the spore bags with half the GFP inducibility germinated. In fact,

for every glucose concentration, we could precisely determine the probability of germinating for a

spore bag once we knew its GFP inducibility (Appendix Fig S18). We thereby established a

quantitative relationship between the inducible GFP level and the ability to germinate. Importantly,

this means that just because a spore bag can express any amount of GFP without nutrients does

not mean that it will germinate after receiving a certain amount of glucose. There is no binary

relationship between the GFP expression and germination. The quantitative relationship between

the GFP inducibility and the probability of germinating ("germination ability") for a given glucose

concentration establishes that the stochastic variability (Padovan-Merhar & Raj, 2013; Balazsi *et*

al, 2011) in the induced GFP expression among dormant spores is meaningful and predictive,

despite GFP not having any functional role in the complex, multi-step process that is germination.

Germination landscape visually represents germination ability as a function of glucose

concentration and GFP inducibility

To visualize our results, we plotted a "germination landscape" - a heat map whose color

represents a probability that a spore bag with a given GFP inducibility germinates (i.e.,

germination ability) for each glucose concentration (Fig. 3F and Appendix Fig S19). In the

germination landscape, yellow represents a near-certain germination (i.e., germination probability

of nearly 1), green represents a germination probability of ~0.5, and dark blue represents a

germination probability of nearly zero. The germination landscape shows a "coastline" of nearly-

yellow pixels moving up towards higher rows (i.e., towards higher glucose concentrations) as one

moves from right to left (i.e., as the GFP inducibility decreases), meaning that more glucose is

required to guarantee a germination for a spore bag with a lesser GFP inducibility. The blue-green

pixels are almost immediately below the coastline of yellow pixels, indicating that the probability

of germinating, for a fixed value of GFP inducibility, is a sharp step-like function of the glucose

concentration. We confirmed this by quantitatively extracting (by log-regression), from the

germination landscape, the minimum glucose-concentration required for a spore with a given

value of GFP inducibility to have a 99%-chance of germinating (Appendix Fig S20). We call this

concentration, given the sharpness of the nearly step-like probability function, the "minimum

glucose concentration required for germination". We determined that as a spore bag's GFP
inducibility decreases, the minimum glucose concentration required for germination increases
(Fig 3G and Appendix Fig S20). Importantly, since inducing GFP expression does not alter the
total percentage of spore bags that germinate for any glucose concentration (Fig. 3D), inducing
GFP expression in a spore bag does not change - neither increase nor decrease - the probability
of germinating. Hence, GFP inducibility merely indicates which spore bags are more likely to
germinate for a given glucose concentration but does not cause or hinder germination. Hence, for
the same glucose concentration, spores with higher GFP inducibility have higher germination
ability than spores with lesser GFP inducibility. Moreover, spores with a higher GFP inducibility
require less glucose to guarantee that they germinate (i.e., have 100% chance of germinating)
than spores with lesser GFP inducibility (Fig. 3G and Appendix Fig S20C).
450
451

Transcribing within a day during dormancy is crucial for spores to remain alive
GFP has no role in germination. So, it is puzzling why GFP inducibility so precisely predicts the
germination ability for each glucose concentration. To make sense of this, we reasoned that the
GFP inducibility represents the dormant spore's ability to express genes in general, not just *GFP*,
when they are forced to do so without any nutrients (e.g., inducers such as doxycycline force an
activation of GFP expression). This is because there is nothing special about GFP.
Notwithstanding the conventional view that gene expression has nearly ceased in dormant yeast
spores, we then reasoned that GFP inducibility is a proxy for gene expressions that are actually
occurring in dormant spores and which are required for germination. According to this reasoning,
inhibiting global (genome-wide) transcription and translation during dormancy would decrease the
percentages of spore bags that germinate (i.e., decrease germination ability) for each glucose
concentration. To test this idea, we incubated spores in PBS for one day with one of three drugs
- thiolutin, cycloheximide, and antimycin A (Fig. 4A). Each drug inhibits some key part of gene
expression. Thiolutin globally inhibits transcription (Grigull *et al,* 2004; Parker *et al,* 1991; Guan
et al, 2006; Lauinger *et al,* 2017; Pelechano & Perez-Ortin; 2008). Cycloheximide globally inhibits
translation (Belle *et al,* 2006; Buchanan *et al,* 2016). Antimycin A inhibits oxidative
phosphorylation and, in turn, synthesis of ATP - a molecule that is required for expressing *GFP*
and other genes (Ocampo *et al,* 2012). After the day-long drug treatment, we washed away the
drug and then added a 2%-glucose with minimal medium to check how many spores could still
germinate (i.e., were still alive) (Fig. 4A).

Figure 4 - More RNA polymerases I-III means higher germination ability and GFP inducibility
(A) Protocol for (B). See "Protocol for Fig. 4B" in Methods. Thiolutin inhibits transcription. Cycloheximide inhibits translation. Antimycin A inhibits ATP synthesis by inhibiting oxidative phosphorylation. **(B)** For experiment in (A), percentage of spore bags that germinated. n=3; error bar are s.e.m. **(C)** Method to detect all RNAs being made in yeast spores with a single-cell resolution. Spores were incubated for 24 hours in PBS with 1 mM of 5-Ethynyl Uridine (5-EU; denoted U*) that incorporates into freshly made RNAs. We fix the spores afterwards and let fluorophore (Alexa 488) enter the spores and bind U* (see Methods). "Total RNA level" of a spore bag is the total fluorescence from all the 5-EU labeled RNAs. **(D)** Microscope images show a spore bag (from strain "TS3") after either incorporating 5-EU as described in (C) (top two images) or, as a control, following the protocol in (C) but without the 5-EU (bottom two images) (also see Appendix Fig S23A). Scale bar: 2 μm. Top histogram: total RNA level per spore bag with 5-EU (i.e., Alexa 488 fluorescence; n = 103 spore bags). Bottom histogram: fluorescence per spore bag in control population (i.e., without 5-EU; n = 95 spore bags). Also see Appendix Fig. S22A. **(E)** Each dot is from a single spore bag ("TS3" strain) with GFP-inducing circuit (Fig. 3A). For each spore bag, we measured its GFP protein level ("GFP inducibility") and total RNA level (5-EU fluorescence) after incubating the spores for 24 hours in PBS with both 100-μg/ml doxycycline and 1 mM of 5-EU (see (C)). n = 245 spore bags. Alexa 594 fluorophore attached to 5-EU (see Methods). Green line: linear regression with R=0.24 and Pearson p-value = 0.00018. **(F)** See "Protocol for Fig. 4F" in Methods. GFP inducibility per spore bag (each dot, "TS8" strain) measured as in (E) but now with live time-lapse without the 5-EU. "RNAP II level" is the mCherry fluorescence per spore bag due to the mCherry protein fused to Rpb3, a subunit of RNA polymerase II. n = 182 spore bags; Red line: linear regression with R=0.64 and Pearson p-value = 3.02 x 10^{-22}. Scale bar = 2 μm. **(G)** Same protocol as in (F) but with "TS9" spores. "RNAP I & III level" is the mCherry fluorescence

per spore bag due to the mCherry protein fused to Rpc40, a subunit of both RNAP I and RNAP III. n = 185 spore bags; Red line: linear regression with R=0.63 and Pearson p-value = 6.6 x 10^{-22}. Scale bar = 2 μm. **(H)** Same protocol as in (F) but with "TT14" spores fixed after 24 hours of incubation in PBS with 100-μg/ml doxycycline. 18s rRNA level is from CAL Fluor Red 610 fluorescence emitted by single-molecule FISH probes bound to 18S rRNAs (see Methods). n = 213 spore bags; Red line: linear regression with R = 0.19 and Pearson p-value = 0.005. **(I)** Bottom: RNAP II levels of spore bags ("TS8" strain) in a population, measured as in (F). Top: As a function of the RNAP II level (binned in the histogram), percentage of spore bags that germinated after receiving a 0.0015%-glucose, averaged over all spore bags with the same binned RNAP II level. n = 80 spore bags. **(J)** Triangular relationship. Any pair of the following three are positively correlated: Germination ability for each glucose concentration (purple), GFP inducibility (green), and amounts of RNA polymerases I-III.

By convention, being a dead spore means that it cannot germinate even after receiving a 2%-
glucose. This definition of death is sensible in light of the germination landscape (Fig. 3F): since
spores are more likely to germinate with a higher glucose concentration than with a lower glucose
concentration, a dead spore is one that has no chance of germinating with the saturating
concentration of glucose and, thus, no chance of germinating for lesser glucose concentration
either. Hence measuring the percentage of spore bags that do not germinate after receiving a
2%-glucose tells us the percentage of spore bags that contain only dead spores.

We verified that all three drugs indeed inhibited GFP expression in yeast spores (Appendix
Fig. S21). Nearly 100% of the spores were still alive after we inhibited either ATP synthesis with
antimycin A or global translation with cycloheximide for 24 hours (Fig. 4B). In contrast, inhibiting
global transcription with thiolutin for 24 hours killed almost all spores: less than 5% of the spore
bags were still able to germinate with a 2%-glucose after the thiolutin treatment (Fig. 4B). These
results establish that transcription - and potentially RNA metabolism in general - is the major
process that keeps dormant spores alive.

Method devised for detecting RNA synthesis in dormant yeast spores at single-cell level
To further explore how transcribing during dormancy is connected to germination ability - and thus
to survival - for a given glucose concentration, we first developed a method to measure all the
RNA that is actively being made in dormant yeast spores with a single-cell resolution. We modified
an existing method for mammalian cells in which the cells uptake 5-Ethynyl Uridine (5-EU) and
incorporates it into RNA that is being synthesized, after which the cells are fixed and fluorophores
(e.g., Alexa 488) enter the fixed cells to bind the 5-EU-labeled RNAs (Jao & Salic, 2008) (Fig.
4C). Since 5-EU would only be found in RNAs that are synthesized while the extracellular 5-EU
was present, this method lets us determine the total amount of RNAs that are made during a
specified time window (i.e., only while the 5-EU was present). The method therefore distinguishes

these freshly made RNAs from the RNAs that were present before we added the 5-EU to the cell
culture medium. By adapting this method for yeast, we could visually verify, by fluorescence, that
yeast spores actively transcribe during dormancy and determine the amounts of RNA produced
by the dormant spores, relative to one another, during a 24-hour incubation in 5-EU (Fig. 4D). To
our knowledge, this is one of the first demonstrations of active transcription in dormant yeast
spores with a single-cell resolution, as previous studies used either indirect or bulk assays that
required lyses of many spores (Brengues *et al,* 2002). We used several methods to verify that our
method worked in yeast spores (Appendix Figs S22-S23).
508
509

Total transcription rate in dormant spores is virtually uncorrelated with germination ability
To apply the adapted method to the GFP-inducible spores, we incubated the spores in PBS with
both doxycycline and 5-EU for 24 hours. Hence, the total RNA level accumulated in a day is a
proxy for the total transcription rate of a dormant spore. We found a statistically significant but
weak, positive correlation (R=0.24) between the spores' total transcription rate (i.e., accumulated
RNA level) and their GFP inducibility (Fig. 4E). This, in turn, means that the total transcription rate
poorly correlates with the spore's chance of germinating. Thus, the total transcription rate cannot
explain why the GFP inducibility so precisely determines the germination ability for each glucose
concentration. Yet, the fact that a 24-hour exposure to thiolutin kills nearly all spores (Fig. 4B)
suggests that some aspect of transcription during dormancy - other than the total amount of
transcription - should strongly correlate with the GFP inducibility and, in turn, with the germination
ability for each glucose concentration. Since RNA polymerases I, II, and III are required and
central machineries for transcription, we next sought to address whether the amount of each RNA
polymerase in a dormant spore strongly correlates with its GFP inducibility.
524
525

Amount of RNA polymerases I-III are strong predictors of GFP inducibility and germination
ability
To measure the amounts of RNA polymerase (RNAP) I, II, and III in individual spores through live
time-lapse microscopy, we genetically engineered two yeast strains. One strain produced Rpb3 -
a subunit of RNAP II - as a protein fused to the fluorescent protein, mCherry (Fig. 4F). The other
strain produced Rpc40 - a subunit of both RNAP I and RNAP III - as a protein fused to mCherry
(Fig. 4G). Both strains also have the GFP-inducing synthetic circuit (Fig. 3A). For these strains,
the mCherry fluorescence was entirely confined to the nuclei of both spores and vegetative yeasts

(Appendix Fig S24). Hence, the mCherry levels accurately reflected the amounts of each RNAPs

in spores. After sporulating these two strains, we incubated the resulting spores in PBS with

doxycycline for 24 hours to induce their *GFP*. We then used a microscope to measure both the

GFP and mCherry (RNAP) levels in individual spore bags. In one of the strains, mCherry level

represents the level of RNAP II, which transcribes all the coding genes and also produces some

non-coding RNAs. In the other strain, mCherry level represents the amount of RNAP I and RNAP

III combined, both of which produce only non-coding RNAs such as ribosomal RNAs. Strikingly,

the amount of RNAP II strongly (R=0.64) and positively correlated with the GFP inducibility (Fig.

4F). The combined amount of RNAPs I and III also strongly (R=0.63) and positively correlated

with GFP inducibility (Fig. 4G). Hence the amounts of both classes of RNA polymerases - one for

transcribing mostly coding genes (RNAP II) and one for transcribing non-coding genes (RNAPs I

and III) - are strong predictors of GFP inducibility. We also used single-molecule RNA FISH (Youk

et al, 2010; Raj *et al,* 2008) to quantify the amount of 18S ribosomal RNA (18S rRNA) - a ribosomal

subunit - in dormant spores that we fixed after the 24 hours of doxycycline. We then determined

how the amount of 18S rRNA in spores correlated with the GFP inducibility. We found that the

amount of 18S rRNA in spores barely correlated with their GFP inducibility (R=0.19). Since the

amount of 18S rRNA is a proxy for the amount of ribosomes, this result suggests that the amounts

of ribosomes in a dormant yeast spore is virtually uncorrelated - and at most weakly correlated -

with GFP inducibility and thus with germination ability. This is consistent with transcription, rather

than translation, being the dominant process for surviving dormancy (Fig. 4B).

Since GFP inducibility strongly and positively correlates with the amounts of all three

RNAPs and germination ability strongly and positively correlates with GFP inducibility, we would

expect that the amount of RNAPs in a dormant spore should also strongly and positively correlate

with the spore's germination ability as well. To verify this for RNAP II - we will focus on RNAP II

from now on instead of RNAPs I and III because RNAP II transcribes all coding genes such as

GFP as well as some non-coding genes - we took the spores with Rpb3 fused to mCherry and

then incubated them in a low concentration of glucose (0.0015%). Then, using the same method

that we used for measuring the germination landscape (Fig. 3E), we determined how likely it is

that a spore bag with a given amount of RNAP II germinates after receiving the 0.0015%-glucose.

We found that spore bags with more RNAP II indeed were more likely to germinate (Fig. 4I).

Taken together, the results so far establish a triangular relationship in which a strong,

positive correlation exists between any two of the following three quantities in dormant yeast

spores (Fig. 4J): GFP inducibility without nutrients, amount of RNAPs (RNAP II level or level of
RNAPs I and III combined), and the germination ability for each glucose concentration.
570
571

Dormant spores gradually lose GFP inducibility as they age
With the triangular relationship, we now have a molecular view - through RNAP II level - and a
functional view - through GFP inducibility - for studying how dormant yeast spores age, which is
what we originally set out to study. To age the dormant spores, we incubated the wild-type and
GFP-inducible spores in either water or minimal medium without any glucose for days or weeks
at 30 ^{O}C. In this "ageing experiment", we found that the number of dormant (i.e., alive) spores -
the ones that we could germinate when we gave them a 2%-glucose - decreased by similar rates
over several weeks regardless of the strain type and regardless of whether we kept the spores in
water or minimal medium (Fig. 5A). Specifically, about half of the spores in the population died
after ~30 days without glucose and almost everyone died after ~60 days without glucose (Fig.
5A). Furthermore, we observed that spores needed more time to germinate if we kept them longer
without nutrients (Appendix Fig S25), suggesting that germinating is becoming more difficult as
the dormant spores age.
585

To measure the GFP inducibility in these ageing spores, we took some of the spores from
the ageing population, incubated them with doxycycline in PBS for 24 hours (Appendix Fig S26),
measured the resulting GFP levels in these spore bags with a microscope (Fig. 5B and Appendix
Fig S27), and thus determined how the GFP inducibility changed over the 80 days in PBS without
nutrients (Appendix Fig S27). Since we also observed dead spores in the ageing population, we
could also measure the dead spores' GFP inducibility with doxycycline on different days. The GFP
inducibility of dead spores remained at near zero (near background fluorescence value)
regardless of when the spores had died (Fig. 5B - grey points). The GFP inducibility of dormant
spores, when averaged over every dormant spore bag, gradually decreased and approached the
near-zero GFP inducibility of the dead spores that were in the same ageing population over tens
of days (Fig. 5B - blue points). The dormant spores' GFP inducibility eventually reached a barely
detectable level - matching the dead spores' GFP inducibility - after ~80 days. By this time, nearly
all spores had died (Appendix Fig S27). Together, these results show that dormant spores indeed
lose their GFP inducibility over time as they age and approach their deaths. Crucially, in
accordance with our revised hypothesis, we did not find any dead spores with a detectable GFP
inducibility. Hence dormant yeast spores lose their GFP inducibility *before* dying.

Figure 5 - Dormant spores gradually have less RNA polymerase II and GFP inducibility as they age, eventually losing all gene expressing ability just before dying.

(A) Percentage of wild-type and GFP-inducible spore-bags (Fig. 3A) that germinate due to a saturating glucose concentration (2%) that they encounter after ageing for 0 - 85 days in water (grey and black) or minimal medium with essential amino acids (red) at 30 °C. n = 3; error bars are s.e.m. **(B)** Mean GFP inducibility of dormant (blue) and dead (grey) spore bags that were incubated in water without nutrients at 30 °C over ~80 days (also see Appendix Figs S27-S28). Blue data points are the average GFP-levels from the histograms shown in Appendix Fig S27. Grey data points are averages of the dead spore bags' histograms shown in Appendix Fig S27. n=3, error bars show the average standard deviation from three biological replicates. n =3. **(C)** Average RNAP II level of spore bags incubated in water for 50 days at 30 °C. n=3, error bars represent the average standard deviation in a population, each population having at least 100 spore bags (see Appendix Figs S28-S29). **(D)** Top: Percentage of spore bags, having the same RNAP II level on 39th day of ageing in water (as in (A)), germinating after receiving a 2%-glucose. Percentages are averaged over all spore bags with the same binned RNAP II level (corresponding histogram shown below). n = 193 spore bags. **(E)** RNAP II levels of dormant (alive) spore bags in blue and of dead spores in grey; both from the same population on 39th day of ageing in water (as in (A)) before any glucose added. n = 111 spore bags (blue); n = 82 spore bags (grey). **(F)** Corresponds to histograms in (E). Blue is for dormant and grey is for dead spore bags. Line traces show, after receiving a 2%-glucose, the fold change in each spore bag's RNAP II level relative to the spore bag's initial RNAP II level (i.e., value just before we added the 2%-glucose). Black dots represent moments of germination. n = 38 for blue and n = 32 for grey. Data points in the rightmost plot show the RNAP II production rate in each spore bag profiled in the line traces. For each spore bag, we fitted an exponential function to its RNAP II level over time (one of the line traces). Average production rate of RNAP II in dormant spore bags (average over all blue points) is a positive value, 0.031 / hour and in dead spore bag (average over all grey points) is a negative value, -0.0030 / hour. For a t-test on the difference between the two values: p-value = 1.4 x 10^{-5}. Linear regression line for dormant (blue) is R = -0.19 and for dead (grey) is R = -0.23.

603   Moreover, the fact that the oldest dormant spores have barely detectable GFP inducibility, just
604   like the dead spores, suggest that GFP inducibility is nearly all lost moments before death.
605
606

Dormant spores gradually have less copies of RNA polymerase II as they age
To further validate that dormant spores lose nearly all GFP inducibility just before dying, we sought
to measure the RNAP II level in ageing spores since GFP inducibility strongly and positively
correlates with the RNAP II level in spores. We repeated the ageing experiment but now with the
spores that had the fluorescent RNAP II. By taking out aliquots of these spores from an ageing
population and then using a microscope to measure the RNAP II levels in each of these aged
spore bags, we found that spores' RNAP II level decreased as they aged. Moreover, after ~50
days of ageing, almost all the spores had barely detectable RNAP II levels which were near the
background fluorescence value (Fig. 5C and Appendix Figs S28-S29).
616
617

More RNA polymerase II means more likely to survive dormancy
After a month of ageing, more than ~50% of the spores were dead. This was the case, for
example, on day 39 of ageing in water. We aliquoted a subset of the aged spores on day 39.
Then, we used a microscope to measure the RNAP II levels in each of the aliquoted spore bags.
Afterwards, we incubated these spores in minimal medium with a 2%-glucose and then, with time-
lapse microscopy, determined which spore bags germinated (i.e., were alive) and which ones did
not germinate (i.e., were dead). Spores that had more RNAP II had a higher chance of being alive
than spores with less RNAP II (Fig. 5D). In particular, we found that nearly 50% or more spores
with the lowest observed RNAP II levels were dead. These spore bags had near background
fluorescence level of ~2000-3000 fluorescence units (Fig. 5D). Since GFP inducibility positively
correlates with the RNAP II level, this finding complements the observation that spores lose
almost completely their GFP inducibility just before dying, as also expected from the triangular
relationship (Fig. 4J).
631
632

633   **Aged spores at the edge of dormancy before dying have nearly depleted RNAP II and have**
634   **strained gene-expressing ability**
635   On the 39th day of ageing, many spores with barely detectable RNAP II levels (2000-3000
636   fluorescence units in Fig. 5D) were dead but many others also with the same barely detectable

RNAP II levels were alive, albeit with less than a 50% chance (Fig. 5D). This 39-days-old population is thus ideal for understanding the last moments of dormancy - dormancy just before death. With this population, we sought to understand what actually distinguishes the dead from dormant spores when both have comparable, barely detectable RNAP II levels (i.e., 2000-3000 fluorescence units in Fig. 5D). We examined two histograms of RNAP II levels: one for dead spores (Fig. 5E - lower, grey histogram) and the other for dormant spores (Fig. 5E - upper, blue histogram). The two histograms show many alive and dead spores having the nearly identical, low range of values for RNAP II level (Fig. 5E). After measuring the RNAP II levels in individual spore bags, we incubated both the dead and dormant spore bags together in a minimal medium with a 2%-glucose and then tracked the RNAP II levels in individual spore bags with time-lapse microscopy. We found that, after receiving the 2%-glucose, the dead spores did not make any detectable levels of new RNAP II (Fig. 5F - grey traces and points) whereas the dormant spores did make observable amounts of new RNAP II until they germinated (Fig. 5F - blue traces and points). Moreover, these dormant spores often started to make new RNAP II shortly before buds appeared (i.e., before they germinated) (Fig. 5F). They also, on average, took longer times to germinate (Appendix Fig S30). Strikingly, these aged dormant spores, near death, did not produce any detectable levels of new RNAP II for the first 5 to 10 hours after receiving the 2%-glucose (Appendix Fig S30). After this "lag time", they began producing new RNAP II and then germinated shortly afterwards.

These results paint a picture in which dormant spores with barely detectable amounts of RNAP II are struggling through gene-expression and germination after receiving a 2%-glucose, as expected intuitively from dormant spores that are near death. This picture also makes sense given that RNA polymerases, including RNAP II, are required to make more RNAPs I-III and that the aged, nearly dead spores have barely detectable amounts of RNAP II. Crucially, these results establish that dead spores, which have virtually undetectable GFP inducibilities during dormancy (Fig. 5B), also have virtually undetectable gene-expressing abilities after they receive a saturating amount (2%) of glucose. This is supported by the fact that RNAP II - required for expressing all coding and some non-coding genes - is at barely detectable levels in the aged spores that are near death (Fig. 5E) and that a 2%-glucose does not induce gene expressions that lead to the making of a new RNAP II (Fig. 5F).

Figure 6 - Stopping transcription for a day causes dormant spores with abundant RNA polymerases to lose all gene-expressing ability and thus die.
(A) Protocol for (B-C) (See "Protocol for Figs. 6(A-C)" in Methods). **(B)** For protocol in (A), mean GFP inducibility of dormant (blue) and dead (grey) spore bags as a function of amount of time spent in thiolutin. *n*=3, error bars represent the standard deviation among spore bags in a population, averaged over three populations. **(C)** RNAP II level of spore bags after 12 hours of thiolutin (green; protocol in (A)) or after 39 days of ageing in water without drugs (light yellow; data from Fig. 5C). **(D)** Protocol for (E) (See "Protocol for Fig. 6E" in Methods). **(E)** Corresponds to green histogram in (C). Blue is for dormant and grey is for dead spore bags. Line traces show, after receiving a 2%-glucose, the fold change in each spore bag's RNAP II level relative to the spore bag's initial RNAP II level (i.e., value just before we added the 2%-glucose). Black dots represent moments of germination. *n* = 12 for blue and *n* = 42 for grey. Data points in the rightmost plot show the RNAP II production rate in each spore bag profiled in the line traces. For each spore bag, we fitted an exponential function to its RNAP II level over time (one of the line traces). Average RNAP II production rate in dormant spore bag (average over all blue points) is a positive value, 0.044 / hour and in dead spore bag (average over all grey points) is a negative value, -0.0045 / hour. For a t-test on the difference between the two values: p-value = 6.5 x 10^{-3}.

Transiently inhibiting transcription during dormancy permanently decreases GFP inducibility

We found that ageing spores gradually lose the GFP inducibility. In other words, older dormant spores, when forced by doxycycline to express *GFP* without nutrients, yield less steady-state GFP levels. Although dormant spores have less RNAP II as they age (Fig. 5C), it is not obvious that less RNAP II should lead to less steady-state GFP level after 24 hours of doxycycline. This is all the more since we showed above that the GFP level reaches a steady-state value because the

spores stop producing GFP, despite the doxycycline not degrading and still being present at the

saturating concentration at the end of the 24-hour incubation (Appendix Figs S13-S14). To

understand why spores lose the GFP inducibility as they age, we sought a clearer understanding

of gene expression in general, which occurs during dormancy as we have shown with the 5-EU

labeling of RNAs (Fig. 4E).

As a start, we incubated the GFP-inducible spores in water with thiolutin for various hours

(between 0 and 24 hours). Afterwards, we washed away the thiolutin and then added doxycycline

for 24 hours to induce their GFP expression (Fig. 6A). At the end of the 24-hr incubation with

doxycycline, we measured the GFP inducibility of each spore bag. After measuring the GFP-

inducibility, we incubated the spores in a 2%-glucose to determine which ones were dormant

(alive) and which were dead. In this experiment, we found that the GFP inducibility of dead spore

bags always remained at the minimally observable values, as expected (Fig. 6B - grey points).

The GFP inducibility of dormant spore bags gradually decreased as the hours spent in thiolutin

increased, until the GFP inducibility reached barely detectable levels after ~15 hours in thiolutin

(Fig. 6B - blue points). Hence, removing thiolutin did not restore the spore's GFP inducibility back

to the level that it had before it encountered thiolutin. In other words, transiently inhibiting global

transcription for ~12 hours or more has permanently (irreversibly) reduced the GFP inducibility.

Along with this irreversible decrease in GFP inducibility, we observed that thiolutin accelerated

deaths, resulting in almost all spores being dead after ~24 hours with thiolutin as opposed to the

~2 months in the previous ageing experiment without thiolutin (Fig. 5A and Appendix Fig S31A).

These results establish that if dormant spores do not produce transcripts - and potentially proteins

from these transcripts - during a day, then they irreversibly and completely lose GFP inducibility

and die within a day. Thus transcribing - and potentially translating - during dormancy during a

day is required for surviving and directly determines how much GFP the spore can make when it

is forced to do so.

Spores that died due to transient inhibition of transcription do not express genes despite

having abundant RNAP II

To better understand how the transcription inhibition by thiolutin causes death, we focused on

spores that encountered thiolutin for 12 hours since many of them were either dead or nearly

dead, thereby affording us enough dead and nearly dead spores to compare. During the 12-hours

of thiolutin, the spores' production of RNAP II itself would be stopped. Indeed, these spores had

~30% less RNAP II after the thiolutin treatment than they had before the treatment. They also had

~30% less RNAP II after the thiolutin treatment than the spores that we kept in water for 12 hours

without thiolutin (Appendix Fig S31). But these thiolutin-treated spores still had abundant RNAP

IIs, nearly 2-3 times more than 39-day-old spores that we kept in water without thiolutin (Fig. 6C).

Surprisingly, however, ~50% of the spores were dead after the 12-hours of thiolutin, which we

discovered by incubating the spores in a 2%-glucose after washing away the thiolutin (Appendix

Fig. S31A and Fig. 6D). In comparison, without thiolutin, we would need to wait ~39 days for ~50%

of the ageing spores to be dead (Appendix Fig. S32).

To test whether the RNAP II-abundant spores that died during the 12 hours of thiolutin

could express genes after receiving a 2%-glucose, we used time-lapse microscopy to track the

RNAP II levels in both dead and dormant spores after washing away the thiolutin and then

incubating the spores in minimal medium with a 2%-glucose (Fig. 6E - left half). As with the spores

that aged for 39 days without thiolutin (Fig. 5F), the thiolutin-treated dormant spores made new

RNAP II after receiving the 2%-glucose, typically taking ~5-10 hours before producing detectable

amounts of new RNAP II (Fig. 6E - blue curves and points) just like the 39-day-old spores with

little RNAP II. As for the spores that died from the thiolutin treatment, we did not observe any of

them producing noticeable amounts of new RNAP II during the ~20 hours of incubation in the 2%-

glucose, despite all of them having abundant RNAP IIs (Fig. 6E - grey curves and data).

Taken together, the thiolutin experiments (Figs. 6B-E) establish that dead spores can have

many copies of RNAP II and yet a 2%-glucose may not induce gene expression in them, including

those necessary for producing RNAP II. These results support the idea that spores need to

produce some transcripts, within several hours (~12 hours as a conservative estimate), in order

to maintain GFP inducibility - and thus for a 2%-glucose to be able to induce gene expression in

them - and hence for the spores to survive. Moreover, the fact that transiently inhibiting

transcription permanently reduces GFP inducibility (Fig. 6B) strongly suggests that during the

transcription inhibition, spores lost copies of key molecules that are required for gene expression

and that the spores could not fully replenish these molecules after transcription became

permissible again. In other words, the experiments suggest that the production rates for these

molecules is lower than their degradation rates during dormancy. This is also consistent with the

fact that dormant yeast spores eventually die if left they are left alone in water without any drugs

(Fig. 5A) - there is a net loss of the molecules because, during dormancy, the production rates for

the molecules are lower than the degradation rates for the molecules.

Figure 7 - Gradual loss of molecules that are required for gene expression underlies ageing and eventual death of dormant yeast spores.

(A) Protocol for (B-E). See "Protocol for Fig. 7(B-E) in Methods". Two drugs: Thiolutin inhibits transcription and cycloheximide inhibits translation. **(B)** Experiment in (A) with cycloheximide. For one representative population with cycloheximide (other replicates and data for thiolutin are in Appendix Fig S32). RNAP II level relative to its day 0 value, averaged over all dormant spore bags in a population. We used a linear interpolation determine the time taken for the halving of RNAP II level (8.4 days). **(C)** Experiment in (A). Time taken for the RNAP II level to decrease by half in dormant spores without drugs (white), with cycloheximide (red), and with thiolutin (green). For spores with thiolutin, we extrapolated this value from the measurements because, even when every spore had died after 2 days of thiolutin, average RNAP II level was still more than half its initial level (Appendix Fig S31). n=3; error bar are s.e.m. **(D)** Experiment in (A). For one representative population with cycloheximide (other replicates and data for thiolutin are in Appendix Fig S32). Time taken for half of the spore population to die ("half-life of dormancy"), estimated by linear interpolation (19.2 days). **(E)** Experiment in (A). Time taken for half of the spore population to die for populations without drugs (white), with cycloheximide (red), and with thiolutin (green). n=3; error bar are s.e.m. **(F)** Putting all our data together leads to a conceptual model of how spores age during dormancy and eventually die: key molecules required for gene expression, including RNA polymerases, are gradually lost during dormancy because spores cannot produce them faster than they degrade. These molecules eventually reach below some threshold amounts after which the spore has permanently lost the ability to express genes, meaning that even ample glucose cannot restore the gene-expressing ability (inset at bottom right). Since, after receiving glucose, spores must express genes in order to build a new cell that will bud off the spore (i.e., germinate), the permanent loss of gene expressing ability at the final moments of dormancy marks the spore's death (grey parts of the curves in the top plot).

Globally inhibiting gene expression shortens spores' lifespans

To end our study, we tested the idea proposed above by estimating the production and degradation rates of key molecules that are necessary for gene expression during dormancy. To achieve this at a systems-level - without having to identify all such molecules and then measure their production and degradation rates - we continuously inhibited transcription and translation while ageing the spores that have the fluorescent RNAP II for ~2 months in water with either a saturating concentration of cycloheximide or thiolutin. On various days, we took out an aliquot of spores from the ageing population, washed away the drug from the aliquoted population, measured the RNAP II level in these spores, and then incubated them in minimal medium with a 2%-glucose to check how many were still alive (i.e., dormant) (Fig. 7A). From these measurements, we determined the rate at which a spore's ability to express genes during dormancy decreases - due to the net loss of key molecules required for gene expression - by measuring the net loss rate of RNAP II as a proxy (Figs. 7B-C). The RNAP II level averaged over all dormant spores decreased by half after ~8.4 days of ageing in cycloheximide (translation inhibitor) (Fig. 7B and Fig. 7C - red bar). As a comparison, the RNAP II level averaged over all dormant spores decreased by half after ~14 days if the spores were left to age in water without any drugs (Fig. 7C - white bar). Hence, inhibiting translation accelerates the net loss of RNAP II by ~1.5 times compared to natural (drug-free) dormancy. Moreover, with cycloheximide, 50% of the spore population were dead after ~19 days (Fig. 7D and Fig. 7E - red bar) whereas, without any drugs, half of the spore population were dead after ~38 days (Fig. 7E - white bar). The approximately 2-fold difference caused by cycloheximide in both the rate of RNAP II loss and the half-life of dormancy supports the idea that dormant spores age and eventually die because they gradually lose enough of the molecules that are required for gene expression, including RNAP II , and that spores cannot replenish these after receiving a 2%-glucose.

The fact that cycloheximide accelerates the net loss of RNAP II proves that RNAP II production, which requires transcription and translation, occurs during dormancy. Cycloheximide prevents spores from producing RNAP II and thus accelerates the RNAP II loss. By how much it is accelerated tells us the production rate of RNAP II during dormancy. We found that dormant spores in water without any drugs produced RNAP II at an extremely slow pace, having a characteristic production time of ~26 days (Appendix Fig S33). In comparison, after receiving a 2%-glucose but before germinating, spores produced new RNAP II 10-50 times faster than they did during dormancy (i.e., before receiving the 2%-glucose) (Fig. 5E and Appendix Fig S33). The extremely slow production of RNAP II during dormancy supports the idea that spores do not

replenish the molecules required for production, such as RNAP II, fast enough for preventing their

net losses. As RNAP II is certainly required for expressing coding genes and thus for germination

(i.e., for building a new cell after receiving a 2%-glucose), the net loss of RNAP II ensures that

dormant spores will eventually die if left alone. Our measurements on RNAP II also strongly

suggest that the molecules required for gene expression, which are lost on the order of at most

several hours during the thiolutin treatment (Fig. 6B), require transcription for them to be made

and are relatively unstable compared to macromolecular machineries such as RNAP II that are

also required for gene expression. In fact, since the RNAP II fluorescence arises from a single

protein, Rpb3, fused to mCherry, the slow degradation of RNAP II also means that smaller

proteins likely also degrade over days, not hours, during dormancy. Taken together, these results

suggest that key transcripts, which are required for gene expression, are lost on the order of at

most several hours during the transcription inhibition. In accordance with this view, spores

retained more than half of their original RNAP II level after a day thiolutin (Fig. 7C) yet the number

of surviving spores were halved after just ~12 hours (Fig. 7E).

Model for dormancy-to-death transition

Taken together, our experiments suggest a model of how dormant spores gradually approach

their deaths (Fig. 7F). In this model, when yeast spores enter dormancy sometime after

sporulation (Brengues *et al,* 2002; Thacker *et al,* 2011), they initially possess high amounts of key

molecules that are necessary for gene expression which include, but are not limited to, RNA

polymerases. Spore bags would vary in the initial amounts of these key molecules - leading to

different germination abilities among them for each glucose - but all of them initially have

sufficiently high amounts of the molecules to let them germinate to a 2%-glucose. During

dormancy, retaining the ability to express genes is crucial for being able to germinate and thus

for survival because a spore must build a new cell after glucose appears (and thus express genes

that code for the building blocks of the new cell). The molecules required for gene expression

become less abundant as time passes without any nutrients because spores cannot replenish

these molecules faster than the rate at which they degrade. The resulting net loss of these

molecules over time leads to the spores gradually losing the ability to express genes such as

those involved in making RNAP II. The spores also become less able to germinate as days pass

by, requiring more glucose to guarantee that they germinate as they age (Fig. 3G). An aged spore

that has lost the gene-expressing ability can neither build a new cell when a 2%-glucose appears

- so it cannot germinate - nor can it make the lost molecules that are crucial for regaining the

gene-expression ability when a 2%-glucose appears. Accordingly, dead spores cannot express

genes including *GFP* (Fig. 5B) and those involved in the making of RNAP II when a 2%-glucose

appears (Fig. 5F). Once the amounts of the molecules required for gene expression decrease

below certain threshold values - note that different molecules can have different threshold values

- the spore cannot germinate even with a saturating amount of glucose and is thus considered

dead. Our results suggest that among the many molecules that are required for gene expression,

the key molecules - the ones that degrade the fastest during dormancy - are RNA polymerases

and, likely, certain crucial transcripts which may be non-coding transcripts. Further studies,

however, are required to definitively show that such transcripts play a role and identify them.

DISCUSSION

Our study began by posing two broad, related questions: how an organism whose life has almost

ceased - including dormant organisms - remain alive and how we can monitor its approach to

death given that it has faint signatures of intracellular dynamics that are difficult to discern. We

addressed key aspects of these two questions in the case of dormant yeast spores while also

raising new questions that invite future investigations. A conceptual difficulty in addressing these

questions lies in the phrase, "almost ceased". Prior to our work, how slowly intracellular processes

such as gene expression occur in dormant yeast spores has been unclear as firm numbers were

missing. With single-cell level measurements, we identified key quantities that are relevant for

studying ageing during dormancy: probability of germinating (which we called "germination

ability"), the ability to express genes during dormancy when forced to so (proxied by the "GFP

inducibility"), and the amount of RNAP polymerases inside the spores (Figs. 1-4). After identifying

these, we put firm numbers on them and showed how their decays over time collectively represent

the process of ageing during dormancy for yeast spores (Figs. 5-7). We found that the RNAP II

production and degradation take days whereas they take minutes in vegetative yeasts. We

rigorously quantified the germination ability for every glucose concentration and found that it

decays over days. With the GFP-inducing synthetic circuit, we quantified the ability of a dormant

spore to express genes without nutrients when forced to do so and found that it also decays over

days. Finally, we measured the lifespan of dormancy and found that one can shorten it from

months - for spores kept in water without any drugs at 30 $^{\circ}$C - to either hours (by inhibiting

transcription during dormancy) or days (by inhibiting translation during dormancy). We revealed,

with a single-cell resolution, how these quantities decayed over time up to the last moments of

dormancy (i.e., just before the yeast spores die), and then determined when a yeast spore is no

longer dormant but dead. Crucially, we found that these quantities are linked to one another and
therefore we can study the dormancy-to-death transition by monitoring several different quantities.
Of these, the key relationship that we uncovered is the ability to express genes being directly
linked to the ability to germinate: a yeast spore cannot germinate when glucose appears after it
has sufficiently lost the ability to express genes. After all, a yeast spore would need to express
genes to build a new cell that will bud off the spore (i.e., germinate). By linking the different
quantities and measuring the extremely slow production rates - compared to degradation rates -
of gene expression machineries such as RNAP II, we showed that a complete loss of gene
expressing ability is inevitable for a yeast spore. These measurements led us to a new conceptual
model for how yeast spores gradually lose dormancy (Fig. 7F). One way to monitor the gradual
loss of dormancy is by measuring the gradually decreasing ability of the yeast spore to express a
generic gene when forced to do so without nutrients (e.g., GFP inducibility). For this reason, we
propose here a concept of "amount of dormancy". The amount of dormancy gradually decreases
in a yeast spore and once it goes below a certain amount, the spore dies because it cannot
express genes even when glucose appears.
864
There are several proxies for the amount of dormancy, such as the GFP inducibility and
the amount of RNAP II. But one class of molecules, which we did not directly identify and measure
here, are key transcripts that we argue are likely important for maintaining gene-expressing ability
in yeast spores. While RNA polymerases are, as expected, required and important for maintaining
gene-expression ability during dormancy, we have shown that ample RNA polymerases and
ribosomes themselves are insufficient for yeast spores to maintain gene-expressing ability during
dormancy. In fact, we found that after several hours of transcription inhibition, yeast spores with
ample RNAP II and ribosomes cannot express coding genes both during dormancy and after
receiving a saturating concentration of glucose - these spores died. By comparing the timescale
of protein degradations in spores with the timescale of deaths that occur during the transcription
inhibitions, we argue that certain transcripts - perhaps non-coding transcripts - that degrade on a
timescale of at most hours are required for maintaining gene-expression ability and, in turn, for
keeping spores alive. We expect that future studies will investigate this interpretation. Such future
studies may benefit from the method that we have, for the first time, adapted for yeast spores
here: 5-Ethynyl Uridine (5-EU) labeling of new RNAs being made during dormancy.
880
On a related topic, our work motivates an avenue of research that is underexplored:
understanding the central dogma of molecular biology in eukaryotic spores by using dormant

yeast spore as a model system. With a synthetic gene-circuit, we showed that inducing

transcription and translation of a generic gene is possible in dormant yeast spores without any

external energy source. We provided, to our knowledge, one of the first evidences of active gene

expression inside intact, dormant yeast spores at a single-cell resolution. This approach - building

synthetic circuits in dormant yeast spores - may enable us to reveal fundamental principles of

eukaryotic transcription and translation in dormancy. As a starting point, our work raises a

fundamental question: how does a dormant yeast spore globally repress gene expression given

that its total transcription rate appears to be nearly uncorrelated with the amount of RNA

polymerases that it has? We believe that a good starting point for addressing this question is our

finding that *GFP* expression stops after a day because the production - either transcription or

translation - stops despite the inducer (doxycycline) still being present. Moreover, we may gain

additional insights into how the central dogma of molecular biology functions in dormant

eukaryotic spores by further investigating the glucose-triggered gene expressions in the primed,

dormant yeast spores (Fig. 2G).

While much is known about bacterial spores, particularly *B. subtilis* spores (Suel *et al,*

2007; Segev *et al,* 2012; Ablett *et al,* 1999; Parry *et al,* 2014; Sinai *et al,* 2015; Dworkin & Shah,

2010), the dormant state of yeast spores - a model for eukaryotic spores - has remained

comparatively underexplored. Our discoveries on yeast spores may not apply to bacterial spores

such as *B. subtilis* spores and determining the similarities and differences between dormant state

of yeast spores with dormant states of bacterial spores is challenging because dormancy of both

remain incompletely understood. To bridge the gap, one may adapt the approach that we

introduced here: building synthetic gene circuits in bacterial spores (e.g., *B. subtilis* spores) and

then studying their dormancy-to-death transition. Since eukaryotic gene regulations have many

features that are distinct from those of prokaryotes, we expect that studies of gene regulation in

dormant bacterial spores - another underexplored topic - along with further studies of gene

regulation in dormant yeast spores will deepen our understanding of both forms of dormancy and

help us elucidate the similarities and difference between the two.

We hope that our study motivates future studies that find systems-level metrics for

monitoring how dormancy in a cell or multicellular organism gradually transitions to death.

Collectively, such studies will likely deepen our understanding of dormancy-to-death transitions.

Materials and Methods

Protocol for Fig. 4B. We first incubated spores (strain "TT14") for 24 hours in PBS with either 10 μg/ml of thiolutin (inhibiting transcription) or 200 μg/ml of cycloheximide (inhibiting translation), or 100 μg/ml of antimycin A (inhibiting ATP production by inhibiting oxidative phosphorylation), or without any drugs. Afterwards, spores were washed 4 times with PBS and then incubated in minimal medium with 2% glucose for 24 hours.

Protocol for Fig. 4F. We simultaneously measured the GFP inducibility and RNAP II level in individual spore bags after 24 hours of incubation in PBS with 100-μg/ml doxycycline (also see "Microscope Data Analysis" section). "RNAP II level" is the mCherry fluorescence per spore bag due to the mCherry protein fused to Rpb3, a subunit of RNA polymerase II.

Protocol for Figs. 6(A-C). We incubated spores for different amounts of time in PBS with 10 μg/ml of thiolutin. Then, after washing away the thiolutin four times with PBS, we incubated the spores for 24 hours in PBS with 100 μg/ml of doxycycline. We then measured the GFP levels ("GFP inducibility") of each spore bag on a microscope. Then, with the spores still under observation in the microscope, we washed away the doxycycline three times with PBS and then incubated the spores for 24 hours in minimal medium with a 2%-glucose to assign a state (dormant or dead) to each spore bag.

Protocol for Fig. 6E. After incubating spores for 12 hours with 10 μg/ml of thiolutin, we washed away the thiolutin four times with PBS. We then place the spores under a microscope and incubated them for 24 hours in minimal medium with a 2%-glucose. Spore bags that germinated within the 24 hours of glucose were counted as dormant (alive). The ones that did not germinate during this time were counted as dead.

Protocol for Fig. 7(B-E). We incubated a population of spores ("TS8" strain) in water with either 10 μg/ml of thiolutin or 200 μg/ml of cycloheximide, or without any drugs. At various time points during the incubation, we aliquoted some of the spores from the population and then, with a microscope, measured the mCherry fluorescence of each spore bag (representing RNAP II level since these spores had mCherry protein fused to Rpb3, a subunit of RNAP II). The "RNAP II level" for each spore bag is the maximal mCherry fluorescence obtained from 36 z-stacks with a z-step

0.2 μm. After measuring the RNAP II level, we kept these spores under the microscope and

washed away the drug four times with PBS. Then, while still keeping the spore bags under the

microscope, we incubated them for 24 hours in minimal medium with 2%-glucose. We measured

the percentage of spore bags that germinated during the 24 hours (these are counted as dormant

(alive)).

Drugs used in Fig. 4A. We used thiolutin from Sigma Aldrich (CAS Number 87-11-6). For

storage, we aliquoted thiolutin into DMSO at a final concentration of 1 mM. We used

cycloheximide from Sigma Aldrich (CAS Number 66-81-9) which we maintained in DMSO at a

final concentration of 100 mg/mL. We used antimycin A from Sigma Aldrich (CAS Number 1397-

94-0) which we diluted into ethanol at 10 mg/mL.

Strains. The "wild-type", homozygous diploid yeast strain that we used is from Euroscarf with the

official strain name "20000D" and genotype as follows: *MATa/MATα his3-11_15/his3-11_15;*

leu2-3_112/leu2-3_112; ura3-1/ura3-1; trp1Δ2/ trp1Δ2; ade2-1/ade2-1; can1-100/can1-100. This

strain generated four genetically identical, haploid spores for each spore bag. For engineering the

GFP-inducible spores, we started from the haploid versions of "20000D", which were also from

Euroscarf. These haploid strains were "20000A" (isogenic to another standard laboratory strain

called "W303" with mating-type "a") and "20000B" (isogenic to W303 with mating-type "alpha").

The 20000A's genotype is as follows: *MATa his3-11_15; leu2-3_112; ura3-1; trp1Δ2; ade2-1;*

can1-100. The 20000B's genotype is exactly the same as 20000A's, except that it is of the

opposite mating type (mating type "alpha"). The GFP-expressing diploid strain, called "TT14", is

nearly identical to the wild-type's except for the addition of selection-marker genes that we

introduced during the construction of the strain. TT14 has the following genotype: (*MATa/MATα*

his3-11_15/his3-11_15; leu2-3_112/leu2-3_112; ura3-1/ura3-1; ADE2/ADE2; can1-100/can1-

100; HygB/HygB; *trp1Δ2/TRP1; URA3/ura3-1; pADH1-rtTA/pADH1-rtTA; pTET07-GFP/pTET07-*

GFP). Here, *pADH1* is the constitutive promoter for *ADH1* gene in yeast (631 bases upstream of

ADH1's ORF), *rtTA* is the reverse tetracycline-controlled transactivator whose transcription-

activation domain is from the yeast's Msn2, and *pTET07* is the promoter with 7 binding sites for

rtTA. We sporulated TT14 to form the "GFP-inducible spores" (Fig. 3A). These constructs are

more fully described in a previous publication (Youk & Lim, 2014). For experiments involving the

metabolic labelling of RNA with 5-Ethynyl-Uridine (5-EU) (Fig. 4C), we had to obtain spores that

were able uptake the extracellular 5-EU for incorporation into intracellular RNA. To achieve this,

we engineered a diploid yeast strain, "TS3", which is the same as TT14 - TS3 also has the same

GFP-inducing synthetic circuit (Fig. 3A) as TT14 - but now with additional genotype modifications

(*HIS3/HIS3; hENT1-ADH1/hENT1-ADH1; HSV-TKGDP/HSV-TKGDP*) for which we used a

construct that was previously shown to let yeast import BrdU (Viggiani & Aparicio, 2006). *hENT1*

is the "human Equilibrative Nucleoside Transporter", under the control of the promotor for *ADH1,*

HSV is the Herpes Simplex Virus thymidine kinase under control of the promotor for *GPD1*, and

HIS3 is a selection marker gene used for the cloning. To obtain the spores whose subunits of

RNA polymerases were fused to mCherry (Rpb3 for RNAP II and Rpc40 for RNAP I and III) (Figs.

4F and 4G), we engineered two diploid strains, "TS8" and "TS9", which were the same as "TS3"

- TS8 and TS9 both have the GFP-inducing synthetic circuit and allow 5-EU uptake) - but now

with additional genotype modifications: (*RPB3-mCherry-NatR / RPB3-mCherry-NatR*) or (*RPC40-*

mCherry-NatR / RPC40-mCherry-NatR). Here "*RPB3-mCherry-NatR*" is the *RPB3* gene on the

yeast chromosome IX (YIL021W) fused at its 3'-end to the *mCherry* gene, using the protein-linker

sequence from (Sheff & Thorn, 2004). *NatR* is the Nourseothricin resistance marker inserted

behind the stop codon of the *RPB3-mCherry* fusion gene. "*RPC40-mCherry-NatR*" is the same

as the "*RPB3-mCherry-NatR"* but now with the *RPC40* gene that is on the yeast chromosome

XVI (YPR110C) instead of *RPB3.*

Strain construction. For each yeast transformation, we integrated a single copy of an

appropriate, linearized yeast-integrating plasmid at a desired genomic locus through a

homologous recombination. We first introduced a promoter of *ADH1* controlling rtTA expression

pADH1-rtTA into 20000A (wild-type haploid, mating type "a") and 20000B (wild-type haploid,

mating type "alpha") at the HO locus, by inserting a linearized, singly integrating, yeast-integrating

plasmid with *pADH1-rtTA* and Hygromycin-resistance gene as a selection marker. This yielded

two strains, "W303r1" (From 20000A) and "W304r1" (from 20000B). We next replaced the *ade2-*

1 "ochre mutation" in W303r1 and W304r1 with a functional *ADE2* gene by a homologous

recombination of *ADE2* that we obtained by PCR from the S288C reference genome (100-bp

homology on both flanking sites of the PCR product). This yielded two strains, "TT2" (from

W303r1) and "TT8" (from W304r1). We then inserted a *pTET07-GFP* at the *LEU2* locus in TT2

and TT8 by linearizing a yeast-integration plasmid that contains *pTET07-GFP*. This yielded two

strains, "TT7" (from TT2) and "TT9" (from TT8). We then introduced two constitutively expressed

selection-marker genes: *URA3* into TT7 to create "TT10" and *TRP1* into TT9 marker to create

"TT13". These selection markers allowed us to select the diploid strain ("TT14") that resulted from

mating TT10 with TT13 by using a double drop-out medium (lacking *-ura* and *-trp*). TT14 is

homozygous for *ADE2, pADH1-rtTA*, and *pTET07-GFP*, thus all four haploids in a TT14 spore-

bag has an inducible *GFP*. For the construction of TS3 strain (EU-incorporating diploids), we
transformed two haploid strains, "TT10" and "TT13", as stated in (Viggiani & Aparicio, 2006) with
a linearized fragment (digested with NheI) of the plasmid "p403-BrdU-Inc" (from (Viggiani &
Aparicio, 2006) and obtained from Addgene). This plasmid contains *hENT1-ADH1*, *HSV-TKGDP*,
and a constitutively expressed selection marker, *HIS3*. We linearized and then integrated the
plasmid, as a single-copy, into the *HIS3* locus in the yeast genome. This resulted in two strains,
"TS1" (from TT10) and "TS2" (from TT13), of opposing mating types. We mated these two to
obtain a diploid strain "TS3". To construct the strain, "TS8" (which has Rpb3 protein fused to
mCherry), we transformed the haploids strains, TS1 and TS2, by integrating mCherry-NatR linker
at the *RPB3* locus in the yeast genome. We obtained the mCherry-NatR linker taking a plasmid
"HypI" (used in (Laman Trip & Youk, 2020) and derived from "pkt150" (Sheff & Thorn, 2004)) and
then amplifying the mCherry-NatR linker through PCR. This resulted in two strains, "TS4" (from
TS1) and "TS5" (from TS2), which had opposite mating types. We mated them to obtain the diploid
strain, "TS8". For constructing the diploid strain, "TS9" (which has Rpc40 protein fused to
mCherry), we transformed the haploids strains "TS1" and "TS2" to obtained two strains, "TS6"
(from TS1) and "TS7" (from TS2), which had opposite mating types. We mated them to obtain the
diploid strain, "TS9".

1034

Yeast transformations and mating. We transformed yeasts with the standard, lithium-acetate-
based method. In short, log-phase cells were resuspended in 0.1M of lithium acetate in 1x TE,
together with the DNA to be inserted by homologous recombination at a desired location within
the yeast genome and herring ssDNA. We then added PEG3350, 10X TE and 1 M of lithium
acetate to the yeast culture so that we had final concentrations of 40% PEG 3350, 1X TE, and
0.1 M of LiOAc. We then incubated the culture at 42 °C for 30 minutes. We pelleted the resulting
cells by centrifuging and then washed them with water. Afterwards, we plated the cells on agar
plates with dropout media to select from transformed yeasts. We used a standard method of
mating yeasts. Namely, we mated yeasts of opposite sexes by inoculating a 500 μL of YPD
medium with two colonies - one from each sex. We then incubated the culture overnight in 30 °C
on a benchtop shaker (Eppendorf Mixmate) that agitated the culture at 300 rpm. We then spread
100 μL of the culture on double-dropout agar plates which selected for the diploids that resulted
from successful mating.

1048

Spore formation (sporulation). We used a standard protocol for sporulating yeasts. In short,
we first grew diploid yeasts (homozygous diploid wild-type or GFP-inducible strains) to saturation

overnight. We then transferred these cells to a "pre-sporulation media" (i.e., YPAc: consists of

Yeast Peptone media with a 2% potassium acetate) which we then incubated for 8 hours at 30°C.

We subsequently transferred the diploid yeasts to a "sporulation medium" (i.e., 2% potassium

acetate) and left them to sporulate for 5 days at 20 °C, while rotating as a liquid culture in a tube.

Afterwards, we transferred the resulting spores to water and stored them at 4°C. Through

measurements, we found that we could store these spores for several months without loss of

viability (i.e., 2%-glucose still germinates ~100% of the spore bags).

1058

Microscope sample preparation. We performed all microscope imaging with 96-well glass-

bottom plates (5242-20, Zell-Kontakt). Prior to each microscope imaging, we pre-treated the

glass-bottom by incubating it for 20 minutes with 0.1 mM of concanavalin A (ConA, C2010, Sigma-

Aldrich). We washed the ConA and then typically added 1 µL of spores in 200 µL of minimal

medium per well. The plates were then centrifuged at 1000 rpm for 1 minute to sediment and

attach all spore bags to the glass-bottom. We then performed the microscope imaging.

1065

Microscope data acquisition. We used Olympus IX81 inverted, epifluorescence, wide-field

microscope. For each time-lapse movie, we collected images once every 10 minutes for every

field of view. The temperature during microscope imaging was maintained by an incubator cage

(OKO Lab) that enclosed the microscope. We acquired each image with an EM-CCD Luca R

camera (Andor) and IQ3 software (Andor). We used a wide-spectrum lamp (AMH-600-F6S,

Andor) for exciting fluorescent proteins.

1072

Microscope data analysis. We processed the microscope images with ImageJ and MATLAB

(Mathworks). To measure the times taken to germinate (and to count spore bags that germinated),

we looked for the first haploid spore that formed a bud for each ascus (spore bag). We segmented

the spore bags by using the Sobel filtering of the brightfield images to create a mask. We extracted

fluorescence values inside the mask. We corrected for the background fluorescence, for each

ascus one by one, by subtracting the average background in a 50-pixel area that surrounded each

ascus. For GFP fluorescence (measuring GFP inducibility) and mCherry fluorescence (measuring

RNAP I-III levels through Rpb3-mCherry or Rpc40-mCherry), we computed the fluorescence per

spore by taking the maximal intensity from 36 z-stacks with a z-step size of 0.2 μm per spore bag.

1082

Hexokinase-based assay to measure glucose concentrations. 10 hours after adding a low

concentration of glucose to germinate some of the spores in a 1-mL minimal medium in a 24-well

microscopy plate (5242-20, Zell-Kontakt), we took 800 µL of the supernatant to measure the

concentration of glucose in it. We determined the glucose concentration by using a hexokinase-

based glucose assay kit (Glucose (HK) Assay, G3293, Sigma-Aldrich) that is based on converting

glucose through hexokinase and $NADP^+$ dependent glucose-6-phosphate-dehydrogenase.

1089

RNA-seq on un-germinated spores. We collected un-germinated spores by first collecting 1 mL

samples of spores that were incubated in a 0.002%-glucose at 0, 16, 48, and 96 hours after the

incubation began. To isolate the un-germinated spores, we treated the 1-mL samples with

zymolyase (786-036, G-Biosciences). Zymolyase lysed vegetative cells that formed from

germinated spores, thus ensuring that we only collected RNA from un-germinated spores for

sequencing. We then extracted the RNA from the leftover, un-germinated spores RiboPure Yeast

Kit (Ambion, Life Technologies) as described by its protocol. Next, we prepared the cDNA library

with the 3' mRNA-Seq library preparation kit (Quant-Seq, Lexogen) as described by its protocol.

Afterwards, we loaded the cDNA library on an Illumina MiSeq with the MiSeq Reagent Kit c2

(Illumina) as described by its protocol. We analyzed the resulting RNA-Seq data as previously

described (Trapnell *et al,* 2012): We performed the read alignment with TopHat, read assembly

with Cufflinks, and analyses of differential gene-expressions with Cuffdiff. We used the reference

genome for *S. cerevisiae* from ensembl. We used the transcriptional modules listed in Appendix

Table S1 for grouping the relevant genes into transcriptional modules.

1104

Fixing and permeabilizing spores. Spores were fixed and made permeable - to let fluorophores

into the spores for 5-EU labeling of RNAs and single-molecule RNA FISH - according to the

standard protocols for vegetative *S. cerevisiae*. An important modification to these protocols is

that to digest the spore wall, we extended the typical incubation in zymolyase to 2 hours during

which the spores were constantly agitated at 300 rpm in 30 ^{O}C. We used this fixation and

permeabilization treatment for performing single-molecule RNA FISH (Fig. 4H) and 5-EU labeling

of RNAs (Fig. 4C-D).

1112

1113   **FISH probes.** We designed single-molecule FISH probes for detecting 18S rRNA (Fig. 4H) with

1114   the Stellaris® FISH Probe Designer (Biosearch Technologies, Inc., Petaluma, CA):

1115   www.biosearchtech.com/stellarisdesigner. The probes were coupled to CAL Fluor Red 610

1116   (Biosearch Technologies, Inc.).

1117

1118   **Single-molecule RNA FISH.** We used the standard protocol for single-molecule RNA FISH in

1119   yeast, detailed in "Protocol for S.cerevisiae from Stellaris RNA FISH" (Biosearch Technologies,

1120   Inc., Petaluma, CA) and also in (Youk *et al,* 2010; Raj *et al,* 2008).

1121

Metabolic (5-Ethynyl Uridine) labelling of freshly synthesized RNA with click reaction. We

used click chemistry to bind fluorophores to the 5-Ethynyl Uridine (5-EU) labeled RNAs in spores.

To do so, we followed the protocol for mammalian cells, from "Click-iT® Plus Alexa Fluor® picolyl

azide toolkits" (Thermo Fisher) but with the following modifications so that the protocol would work

in yeast spores: (1) all reactions and washing steps were done in a 1.5-mL Eppendorf tube with

a centrifugation speed of 800 x G; (2) spore fixation and permeabilization were performed as

mentioned above; (3) total volume for the click reaction was 50 μL; (3) for the click reaction, fixed

spores were at OD ~ 0.02; (4) the click-reaction cocktail was incubated for 1 hour and 30 minutes.

All components (Ethynyl-Uridine, Fluorophore-Azide, and reagents for the click reaction) were

from the "Click-iT® Plus Alexa Fluor® picolyl azide toolkits" (Thermo Fisher). For simultaneously

imaging 5-EU bound RNAs with GFP proteins inside spores (Fig. 4E), we used Picolyl Azide

Alexa 594 from JenaBioscience. Here, the ratio of CuSO4 to the copper protectant was set to

1:20 in order to simultaneously detect the fluorescence from both GFP and Alexa 594.

Supporting Information

Appendix Figures S1-S33

Appendix Table S1

Acknowledgements:

We thank Sophie van de Gevel for her help with the 5-EU labelling of RNA. We thank Amir Mitchell for insightful discussions and critical comments on our manuscript. We thank the members of the Youk laboratory for fruitful discussions. We thank The Kavli Nanolab Imaging Centre (KNIC) and Jeremie Capoulade for support and help with microscopy. H.Y. was supported by the European Research Council (ERC) Starting Grant (MultiCellSysBio, #677972), Netherlands Organisation for Scientific Research (NWO) Vidi Award (#680-47-544), CIFAR Azrieli Global Scholars Program, and EMBO Young Investigator Award.

Author Contributions:

T.M. and H.Y. conceived the project and designed the experiments. T.A. and M.A.B. performed the experiments and data analyses for Fig. 1. T.M., T.A. and M.A.B, performed the experiments and data analyses for Fig. 2. T.M. performed all the experiments and data analyses for Figs. 3 to 7. H.Y. provided overall guidance. T.M. and H.Y. wrote the manuscript.

Conflict of interest:

The authors declare that they have no competing interests.

Data availability:

The authors declare that all data supporting the findings of this study are available within the paper and its supplementary information files. The data that support the findings of this study are available from the corresponding author upon reasonable request. The RNA-Seq data in Fig. 2G is available on NCBI GEO with accession number #.

References

Ablett S, Darke AH, Lillford PJ, Martin DR (1999) Glass formation and dormancy in bacterial spores. *Int. Journal of Food Sci Tech.* **34**: 59-69.

Balazsi B, Van Oudenaarden A, Collins JJ (2011) Cellular decision making and biological noise: From microbes to mammals. *Cell* **144**: 910-925.

Belle A, Tanay A, Bitincka L, Shamir R, O'Shea EK (2006) Quantification of protein half-lives in the budding yeast proteome. *Proc Natl Acad Sci USA* **103**: 13004-13009

Brengues M, Pintard L, Lapeyre B (2002) mRNA decay is rapidly induced after spore germination of Saccharomyces cerevisiae. *J Biol Chem* **277**, 40505-40512.

Buchanan BW, Lloyd ME, Engle SM, Rubenstein EM (2016) Cycloheximide chase analysis of protein degradation in Saccharomyces cerevisiae. *J Vis Exp* 110:53975.

Chu S, DeRisi J, Eisen M, Mulholland J, Botstein D, Brown PO, Herskowitz I (1998) The transcriptional program of sporulation in budding yeast. *Science* **282**: 699-705.

Coluccio A, Bogengruber E, Conrad MN, Dresser ME, Briza P, Neiman AM (2004) Morphogenetic pathway of spore wall assembly in Saccharomyces cerevisiae. *Eukaryot Cell* **3**: 1464-1475.

Cowan AE, Koppel DE, Setlow B, Setlow P (2003) A soluble protein is immobile in dormant spores of bacillus subtilis but is mobile in germinated spores: implications for spore dormancy. *Proc. Natl. Acad. Sci. USA* **100**: 4209-4214.

Dijksterhuis J, Nijsse J, Hoekstra FA, Golovina EA (2007) High viscosity and anisotropy characterize the cytoplasm of fungal dormant stress-resistant spores. *Eukaryotic Cell* **6**: 157-170.

1194 Donnini C, Puglisi PP, Vecli A, Marmiroli N (1988) Germination of Saccharomyces cerevisiae
1195   ascospores without trehalose mobilization as revealed by in vivo 13C nuclear magnetic
1196   resonance spectroscopy. *J. Bacteriol.* **170**, 3789-3791.

1197

1198 Dworkin J, Shah IM (2010) Exit from dormancy in microbial organisms. *Nat Rev Microbiol* **8**:
1199   890-896.

1200

1201 Eldar A, Chary VK, Xenopoulos P, Fontes ME, Loson OC, Dworkin J, Piggot PJ, Elowitz MB
1202   (2009) Partial penetrance facilitates developmental evolution in bacteria. *Nature* **460**: 510-
1203   514.

1204

1205 Errington J (2003) Regulation of endospore formation in Bacillus subtilis. *Nat Rev Microbiol*
1206   **1**:117-126.

1207

1208 Geijer C, Pirkov I, Vongsangnak W, Ericsson A, Nielsen J, Krantz M, Hohmann S (2012) Time
1209   course gene expression profiling of yeast spore germination reveals a network of transcription
1210   factors orchestrating the global response. *BMC Genomics* **13**:554.

1211

1212 Grigull J, Mnaimneh S, Pootoolal J, Robinson MD, Hughes TR (2004) Genome-wide analysis of
1213   mRNA stability using transcription inhibitors and microarrays reveals posttranscriptional
1214   control of ribosome biogenesis factors. *Mol Cell Biol.* **24**: 5534-5547.

1215

1216 Guan Q, Zheng W, Tang S, Liu X, Zinkel RA, Tsui KW, Yandell BS, Culbertson MR (2006) Impact
1217   of nonsense-mediated mRNA decay on the global expression profile of budding yeast. *PLoS*
1218   *Genet* **2**: e203

1219

1220 Herman PK, Rine J (1997) Yeast spore germination: a requirement for Ras protein activity during
1221   re-entry into the cell cycle. *EMBO J.* **16**: 6171-6181.

1222

1223 Ihmels J, Friedlander G, Bergmann S, Sarig O, Ziv Y, Barkai N (2002) Revealing modular
1224   organization in the yeast transcriptional network. *Nat. Genet.* **31**: 370-377
1225

1226 Jao CY, Salic A (2008) Exploring RNA transcription and turnover in vivo by using click chemistry.
1227   ***Proc Natl Acad Sci USA*** **105:** 15779–15784.

1228

1229 Joseph-Strauss D, Zenvirth D, Simchen G, Barkai, N (2007) Spore germination in Saccharomyces
1230   cerevisiae: global gene expression patterns and cell cycle landmarks. ***Genome Biol.*** **8:**
1231   R241.

1232

1233 Joyner RP, Tang JH, Helenius J, Dultz E, Brune C, Holt LJ, Huet S, Muller DJ, Weis K (2016) A
1234   glucose-starvation response regulates the diffusion of macromolecules. ***eLife*** **5:** e09376.

1235

1236 Laman Trip DS, Youk H (2020) Yeasts collectively extend the limits of habitable temperatures by
1237   secreting glutathione. ***Nat Microbiol*** **5:** 943-953.

1238

1239 Laporte D, Salin B, Daignan-Fornier B, Sagot I (2008) Reversible cytoplasmic localization of the
1240   proteasome in quiescent yeast cells. ***J Cell Biol*** **181:** 737-745.

1241

1242 Lauinger L, Li J, Shostak A, Cemel IA, Ha N, Zhang Y, Merkl PE, Obermeyer S, Stanovic-Valentin
1243   N, Schafmeier T, Wever WJ, Bowers AA, Carter KP, Palmer AE, Tschochner H, Melchior F,
1244   Deshaies RJ, Brunner M, Diernfellner A (2017) Thiolutin is a zinc chelator that inhibits the
1245   Rpn11 and other JAMM metalloproteases. ***Nat Chem Bio*** **13:** 709-714

1246

1247 Lennon JT, Jones SE (2011) Microbial seed banks: the ecological and evolutionary implications
1248   of dormancy. ***Nat Rev MIcrobiol*** **9:**119-130.

1249

1250 Munder MC, Midtvedt D, Franzmann T, Nuske E, Otto O, Herbig M, Ulbricht E, Muller P,
1251   Taubenberger A (2016) A pH-driven transition of the cytoplasm from a fluid- to a solid-like
1252   state promotes entry into dormancy. ***eLife*** **5:** e09347.

1253

1254 Mutlu A, Trauth S, Ziesack M, Naler K, Bergest JP, Rohr K, Becker N, Hofer T, Bischofs IB (2018)
1255   Phenotypic memory in Bacillus subtilis links dormancy entry and exit by a spore quantity-
1256   quality tradeoff. ***Nat Comm*** **9:**69.

1257

1258   Nachman I, Regev A, Ramanathan S (2007) Dissecting timing variability in yeast spores. *Cell*
1259   **131**: 544-556.

1260

1261   Nagtzaam MPM, Bollen GJ (1994) Long shelf life of Talaromyces flavus in coating material of
1262   pelleted seeds. *Eur. J. Plant Pathol.* **100**: 279-282.

1263

1264   Neiman AM (2011) Sporulation in the budding yeast Saccharomyces cerevisiae. *Genetics* **189**:
1265   737-765.

1266

1267   Ocampo A, Liu J, Schroeder EA, Shadel GS, Barrientos A (2012) Mitochondrial respiratory
1268   thresholds regulate yeast chronological lifespan and its extension by caloric restriction. *Cell*
1269   *Metab* **16**: 55-67.

1270

1271   Padovan-Merhar O, Raj A (2013) Using variability in gene expression as a tool for studying gene
1272   regulation. *Wiley Interdiscip Rev Syst Biol Med.* **5**: 751-759.

1273

1274   Parker R, Herrick D, Peltz SW, Jacobson A (1991) Measurement of mRNA decay rates in
1275   *Saccharomyces cerevisiae. Methods in Enzymology* **194:** 415-423

1276

1277   Parry BR, Surovtsev IV, Cabeen MT, O'Hern CS, Dufresne ER, Jacobs-Wagner C (2014) The
1278   bacterial cytoplasm has glass-like properties and is fluidized by metabolic activity. *Cell* **156**:
1279   183-194.

1280

1281   Pelechano V, Perez-Ortin JE (2008) The transcriptional inhibitor thiolutin blocks mRNA
1282   degradation in yeast. *Yeast* **25**: 85-92.

1283

1284   Petrovska I, Nuske E, Munder MC, Kulasegaran G, Malinovska L, Kroschwald S, Richter D,
1285   Fahmy K, Gibson K, Verbavatz JM, Alberti S (2014) Filament formation by metabolic
1286   enzymes is a specific adaptation to an advanced state of cellular starvation. *eLife* **3**: e02409.
1287

1288 Raj A, Van den Boogard P, Rifkin SA, Van Oudenaarden A, Tyagi S (2008) Imaging individual
1289   mRNA molecules using multiple singly labeled probes. **Nat Methods 5**: 877-879.
1290

1291 Rittershaus ESC, Baek S.-H, Sassetti CM (2013) The normalcy of dormancy: common themes in
1292   microbial quiescence. **Cell Host Microbe 13**: 643-651.
1293

1294 Segev E, Smith Y, Ben-Yehuda S (2012) RNA dynamics in aging bacterial spores. **Cell 148**: 139-
1295   149.
1296

1297 Sheff M, Thorn K (2004) Optimized cassettes for fluorescent protein tagging in Saccharomyces
1298   cerevisiae. **Yeast 21**: 661-670.
1299

1300 Sinai L, Rosenberg A, Smith Y, Segev E, Ben-Yehuda S (2015) The molecular timeline of a
1301   reviving bacterial spore. **Mol Cell. 57**: 695-707.
1302

1303 Stragier P, Losick R (1996) Molecular genetics of sporulation in Bacillus subtilis. **Annu. Rev.**
1304   **Genet. 30**, 297-341.
1305

1306 Sturm A, Dworkin J (2015) Phenotypic diversity as a mechanism to exit cellular dormancy.
1307   **Current Biology 25**: 2272-2277.
1308

1309 Suel G, Garcia-Ojalvo J, Liberman L, Elowitz MB (2006) An excitable gene regulatory circuit
1310   induces transient cellular differentiation. **Nature 440**: 545-550.
1311

1312 Suel G, Kulkarni RP, Dworkin J, Garcia-Ojalvo J, Elowitz MB (2007) Tunability and noise
1313   dependence in differentiation dynamics. **Science 315**: 1716-1719.
1314

1315 Thacker D, Lam I, Knop M, Keeney S (2011) Exploiting spore-autonomous fluorescent protein
1316   expression to quantify meiotic chromosome behaviors in Saccharomyces cerevisiae.
1317   **Genetics 189**: 423-439.
1318

1319   Trapnell C, Roberts A, Goff L, Pertea G, Kim Daehwan, Kelley DR, Pimentel H, Salzberg SL, Rinn

1320   JL, Pachter L (2012) Differential gene and transcript expression analysis of RNA-seq

1321   experiments with TopHat and Cufflinks. *Nat. Protoc.* **7**: 562-578.

1322

1323   Viggiani CJ, Aparicio OM (2006) New vectors for simplified construction of BrdU Incorporating

1324   strains of Saccharomyces cerevisiae. *Yeast* **23:** 1045–1051.

1325

1326   Youk H, Lim WA (2014) Secreting and sensing the same molecule allows cells to achieve versatile

1327   social behaviors. *Science* **343**: 1242782.

1328

1329   Youk H, Raj A, Van Oudenaarden, A (2010) Imaging single mRNA molecules in yeast. *Methods*

1330   *in Enzymology* **470**: 429-446

1331

Figure captions

Figure 1 - Glucose germinates only a fraction of yeast spores.

(A) Conventional test to determine whether a spore bag (i.e., ascus) containing four haploid spores, in the absence of any external nutrients (blue box), is dormant or dead. Green box: outcome if dormant. Red box: outcome if dead.

(B) Left: Filmstrip of a time-lapse movie in which a 2%-glucose was added at the beginning of the movie (10 minutes between frames). A spore bag is counted as having germinated at the moment that at least one budding cell emerges from the spore bag (white arrows). Right: Time taken by each "wild-type" spore bag to germinate in the time-lapse movie. n = 137 spore bags from a representative time-lapse movie.

(**C)** Percentage of wild-type spore bags that germinated as a function of time after adding glucose to the minimal medium. Different colors represent different glucose concentrations (from 0.0002% to 2%). n = 3; error bars are s.e.m.

(D) Average time taken to germinate (blue circles) and the total percentage of spore bags that germinated (red squares), both 16 hours (960 minutes) after receiving glucose (i.e., the plateau values for each color in Fig. 1C). Both are functions of glucose concentration. n = 3; error bars are s.e.m.

Figure 2 - Un-germinated spores, primed by a low glucose concentration, germinate faster upon encountering more glucose a few days later.

(A) Wild-type spores are first incubated in a low glucose concentration ("1st [glucose]") before we add more glucose later to increase the glucose concentration to "final [glucose]".

(B) For experiment in (A), number of spore bags germinated within each binned time. First glucose concentration is 0.0005% (from 0 to 16 hours) (green bars) and the final concentration is 0.002% (from 16 to 32 hours) (orange bars). n = 143 spore bags (representative data).

(C) For the experiment in (A), percentage of spore bags that germinated as a function of time since the first glucose addition (1st and final concentrations as indicated). More glucose added 1000 minutes after the first glucose (purple vertical line). n = 3; error bars are s.e.m.

(D) For the experiment in (A), average time taken for a spore bag to germinate - denoted $\Delta\tau$ in bottom panel of (C) - after stepping up the glucose concentration. We varied the first glucose concentration, but the final concentration was always 2%. n = 3; error bars are s.e.m.

(E) Spore bags that do not germinate after encountering the first glucose are "primed" to
germinate faster upon encountering more glucose.

(F) For the experiment in (A), average time ($\Delta\tau$) taken for a spore bag to germinate after stepping
up the glucose concentration (final glucose concentration is 2%), plotted as a function of the first,
low glucose concentration. Different colors represent different times at which we added more
glucose after the first addition: 16 hours (yellow), 48 hours (blue) and 96 hours (red). "Relative
$\Delta\tau$" is the average time $\Delta\tau$ divided by the $\Delta\tau$ of spore bags that were incubated in minimal media
without any glucose (0%) for the same duration of time (after which they received a 2%-glucose).
n = 3; error bars are s.e.m.

(G) Heat map showing transcriptome-wide changes in un-germinated spores at 0, 16, 48, and 96
hours after being primed by a 0.002%-glucose (via RNA-Seq). Appendix Table S1 lists the genes
for each transcriptional module (each row in the heat map). Also see Appendix Fig S8. For each
transcriptional module, we first divided the expression level of each gene in that module by its
expression level at 0-hours - this yields "normalized expression level" for that gene for both primed
and unprimed spores (unprimed spores were in minimal media without glucose for 0, 16, 48, and
96 hours). We averaged these values over all genes in a transcriptional module to obtain
"normalized expression (primed)" and "normalized expression (unprimed)" for each transcriptional
module. Colors represent ratio of these two values, averaged over three biological replicates (n =
3).

1384

1385

Figure 3 - Inducing GFP expression during dormancy with synthetic circuit shows that
spores that can realize higher GFP level (higher GFP inducibility) are more likely to
germinate (higher germination ability) for any glucose concentration.
(A) Synthetic gene-circuit that constitutively expresses a transcription factor, rtTA (with ADH1-
promoter) and an inducible promoter *(*TET-promoter) controlling *GFP* expression. Increasing
doxycycline increases GFP expression.

(B) Engineered spore bags (shown in (A)) transcribe and translate GFP in water (without any
nutrients) with 25 μg/ml of doxycycline (top row) and in a saline solution (PBS) with 50 μg/ml of
doxycycline (bottom row). Snapshots of GFP expression shown 22 hours after adding
doxycycline**.**

(C) GFP levels of individual spore bags (grey curves) over time (measured every 10 minutes with
a wide-field epifluorescence microscope) after incubation in PBS with 10 μg/ml of doxycycline
(top panel: n = 104 spore bags) and 100 μg/ml of doxycycline (bottom panel: n = 150 spore bags).

(D) Engineered spore bags (shown in (A)) were first incubated for 22 hours in either PBS without any doxycycline or with 100 μg/ml of doxycycline. Next, we transferred them to minimal medium with various glucose concentrations. Plot shows the total percentage of the engineered spore bags that germinated (measured 20 hours after incubating with glucose) for spores pre-incubated in PBS without doxycycline (black points) and in PBS with 100 μg/ml of doxycycline (orange points). n = 3; error bars are s.e.m.

(E) Top: Percentage of spore bags with the same GFP level ("GFP inducibility") from (D) that germinated after receiving a 0.001%-glucose. Percentages are averaged over all spore bags with the same binned GFP level (in the histogram below). Representative data shown (n = 145 spore bags in a population).

(F) Germination landscape: colors represent the probability that a spore bag with a particular steady-state GFP level ("GFP inducibility") germinates for each glucose concentration (i.e., data in top panel of (E) represents a single row of this heat map). To measure each pixel, as in the experiment described in (D), we incubated spore bags in PBS with 100 μg/ml of doxycycline for 22 hours before adding glucose ([glucose] as indicated along the rows). Columns indicate GFP inducibility of a spore bag, measured at 22 hours after adding the doxycycline. Each pixel is an average over 3 replicate populations (n = 3).

(G) Given a spore bag, its GFP inducibility is a read-out of the probability to germinate ("germination ability") for each glucose concentration and the minimal glucose concentration required to guarantee germination. Spore bags with lesser gene-expressing abilities without nutrients require more glucose to germinate.

Figure 4 - More RNA polymerases I-III means higher germination ability and GFP inducibility

(A) Protocol for (B). See "Protocol for Fig. 4B" in Methods. Thiolutin inhibits transcription. Cycloheximide inhibits translation. Antimycin A inhibits ATP synthesis by inhibiting oxidative phosphorylation.

(B) For experiment in (A), percentage of spore bags that germinated. n=3; error bar are s.e.m.

(C) Method to detect all RNAs being made in yeast spores with a single-cell resolution. Spores were incubated for 24 hours in PBS with 1 mM of 5-Ethynyl Uridine (5-EU; denoted U*) that incorporates into freshly made RNAs. We fix the spores afterwards and let fluorophore (Alexa 488) enter the spores and bind U* (see Methods). "Total RNA level" of a spore bag is the total fluorescence from all the 5-EU labeled RNAs.

(D) Microscope images show a spore bag (from strain "TS3") after either incorporating 5-EU as described in (C) (top two images) or, as a control, following the protocol in (C) but without the 5-EU (bottom two images) (also see Appendix Fig S23A). Scale bar: 2 μm. Top histogram: total RNA level per spore bag with 5-EU (i.e., Alexa 488 fluorescence; n = 103 spore bags). Bottom histogram: fluorescence per spore bag in control population (i.e., without 5-EU; n = 95 spore bags). Also see Appendix Fig. S22A.

(E) Each dot is from a single spore bag ("TS3" strain) with GFP-inducing circuit (Fig. 3A). For each spore bag, we measured its GFP protein level ("GFP inducibility") and total RNA level (5-EU fluorescence) after incubating the spores for 24 hours in PBS with both 100-μg/ml doxycycline and 1 mM of 5-EU (see (C)). n = 245 spore bags. Alexa 594 fluorophore attached to 5-EU (see Methods). Green line: linear regression with R=0.24 and Pearson p-value = 0.00018.

(F) See "Protocol for Fig. 4F" in Methods. GFP inducibility per spore bag (each dot, "TS8" strain) measured as in (E) but now with live time-lapse without the 5-EU. "RNAP II level" is the mCherry fluorescence per spore bag due to the mCherry protein fused to Rpb3, a subunit of RNA polymerase II. n = 182 spore bags; Red line: linear regression with R=0.64 and Pearson p-value = 3.02 x 10^{-22}. Scale bar = 2 μm.

(G) Same protocol as in (F) but with "TS9" spores. "RNAP I & III level" is the mCherry fluorescence per spore bag due to the mCherry protein fused to Rpc40, a subunit of both RNAP I and RNAP III. n = 185 spore bags; Red line: linear regression with R=0.63 and Pearson p-value = 6.6 x 10^{-22}. Scale bar = 2 μm.

(H) Same protocol as in (F) but with "TT14" spores fixed after 24 hours of incubation in PBS with 100-μg/ml doxycycline. 18s rRNA level is from CAL Fluor Red 610 fluorescence emitted by single-molecule FISH probes bound to 18S rRNAs (see Methods). n = 213 spore bags; Red line: linear regression with R = 0.19 and Pearson p-value = 0.005.

(I) Bottom: RNAP II levels of spore bags ("TS8" strain) in a population, measured as in (F). Top: As a function of the RNAP II level (binned in the histogram), percentage of spore bags that germinated after receiving a 0.0015%-glucose, averaged over all spore bags with the same binned RNAP II level. n = 80 spore bags.

(J) Triangular relationship. Any pair of the following three are positively correlated: Germination ability for each glucose concentration (purple), GFP inducibility (green), and amounts of RNA polymerases I-III.

Figure 5 - Dormant spores gradually have less RNA polymerase II and GFP inducibility as they age, eventually losing all gene expressing ability just before dying.

(A) Percentage of wild-type and GFP-inducible spore-bags (Fig. 3A) that germinate due to a saturating glucose concentration (2%) that they encounter after ageing for 0 - 85 days in water (grey and black) or minimal medium with essential amino acids (red) at 30 ^{O}C. n = 3; error bars are s.e.m.

(B) Mean GFP inducibility of dormant (blue) and dead (grey) spore bags that were incubated in water without nutrients at 30 ^{O}C over ~80 days (also see Appendix Figs S27-S28). Blue data points are the average GFP-levels from the histograms shown in Appendix Fig S27. Grey data points are averages of the dead spore bags' histograms shown in Appendix Fig S27. n=3, error bars show the average standard deviation from three biological replicates. n =3.

(C) Average RNAP II level of spore bags incubated in water for 50 days at 30 ^{O}C. n=3, error bars represent the average standard deviation in a population, each population having at least 100 spore bags (see Appendix Figs S28-S29).

(D) Top: Percentage of spore bags, having the same RNAP II level on 39th day of ageing in water (as in (A)), germinating after receiving a 2%-glucose. Percentages are averaged over all spore bags with the same binned RNAP II level (corresponding histogram shown below). n = *193* spore bags.

(E) RNAP II levels of dormant (alive) spore bags in blue and of dead spores in grey; both from the same population on 39th day of ageing in water (as in (A)) before any glucose added. n = 111 spore bags (blue); n = 82 spore bags (grey).

(F) Corresponds to histograms in (E). Blue is for dormant and grey is for dead spore bags. Line traces show, after receiving a 2%-glucose, the fold change in each spore bag's RNAP II level relative to the spore bag's initial RNAP II level (i.e., value just before we added the 2%-glucose). Black dots represent moments of germination. n = 38 for blue and n = 32 for grey. Data points in the rightmost plot show the RNAP II production rate in each spore bag profiled in the line traces. For each spore bag, we fitted an exponential function to its RNAP II level over time (one of the line traces). Average production rate of RNAP II in dormant spore bags (average over all blue points) is a positive value, 0.031 / hour and in dead spore bag (average over all grey points) is a negative value, -0.0030 / hour. For a t-test on the difference between the two values: p-value = 1.4 x 10^{-5}. Linear regression line for dormant (blue) is R = -0.19 and for dead (grey) is R = -0.23.

Figure 6 - Stopping transcription for a day causes dormant spores with abundant RNA
polymerases to lose all gene-expressing ability and thus die.
(A) Protocol for (B-C) (See "Protocol for Figs. 6(A-C)" in Methods).
(B) For protocol in (A), mean GFP inducibility of dormant (blue) and dead (grey) spore bags as a
function of amount of time spent in thiolutin. $n=3$, error bars represent the standard deviation
among spore bags in a population, averaged over three populations.
(C) RNAP II level of spore bags after 12 hours of thiolutin (green; protocol in (A)) or after 39 days
of ageing in water without drugs (light yellow; data from Fig. 5C).
(D) Protocol for (E) (See "Protocol for Fig. 6E" in Methods).
(E) Corresponds to green histogram in (C). Blue is for dormant and grey is for dead spore bags.
Line traces show, after receiving a 2%-glucose, the fold change in each spore bag's RNAP II level
relative to the spore bag's initial RNAP II level (i.e., value just before we added the 2%-glucose).
Black dots represent moments of germination. $n = 12$ for blue and $n = 42$ for grey. Data points in
the rightmost plot show the RNAP II production rate in each spore bag profiled in the line traces.
For each spore bag, we fitted an exponential function to its RNAP II level over time (one of the
line traces). Average RNAP II production rate in dormant spore bag (average over all blue points)
is a positive value, 0.044 / hour and in dead spore bag (average over all grey points) is a negative
value, -0.0045 / hour. For a t-test on the difference between the two values: p-value = 6.5×10^{-3}.
1517
1518
Figure 7 - Gradual loss of molecules that are required for gene expression underlies ageing
and eventual death of dormant yeast spores.
(A) Protocol for (B-E). See "Protocol for Fig. 7(B-E) in Methods". Two drugs: Thiolutin inhibits
transcription and cycloheximide inhibits translation.
(B) Experiment in (A) with cycloheximide. For one representative population with cycloheximide
(other replicates and data for thiolutin are in Appendix Fig S32). RNAP II level relative to its day
0 value, averaged over all dormant spore bags in a population. We used a linear interpolation
determine the time taken for the halving of RNAP II level (8.4 days).
(C) Experiment in (A). Time taken for the RNAP II level to decrease by half in dormant spores
without drugs (white), with cycloheximide (red), and with thiolutin (green). For spores with
thiolutin, we extrapolated this value from the measurements because, even when every spore
had died after 2 days of thiolutin, average RNAP II level was still more than half its initial level
(Appendix Fig S31). $n=3$; error bar are s.e.m.

(D) Experiment in (A). For one representative population with cycloheximide (other replicates and

data for thiolutin are in Appendix Fig S32). Time taken for half of the spore population to die ("half-

life of dormancy"), estimated by linear interpolation (19.2 days).

(E) Experiment in (A). Time taken for half of the spore population to die for populations without

drugs (white), with cycloheximide (red), and with thiolutin (green). n=3; error bars are s.e.m.

(F) Putting all our data together leads to a conceptual model of how spores age during dormancy

and eventually die: key molecules required for gene expression, including RNA polymerases, are

gradually lost during dormancy because spores cannot produce them faster than they degrade.

These molecules eventually reach below some threshold amounts after which the spore has

permanently lost the ability to express genes, meaning that even ample glucose cannot restore

the gene-expressing ability (inset at bottom right). Since, after receiving glucose, spores must

express genes in order to build a new cell that will bud off the spore (i.e., germinate), the

permanent loss of gene expressing ability at the final moments of dormancy marks the spore's

death (grey parts of the curves in the top plot).

A

How to distinguish dormant from dead?

Without energy source (kept in water)

With energy source

Yeast spore | Ascus (Spore bag)

No signs of life

Dormant or dead?

add sugar

If dormant

Replicating (Germinating)

alive

If dead?

No replication Ungerminated

why dead?

B

Added glucose

Time

Germinating

Germinating

[Glucose] = 2%

C

Only some spores germinate due to glucose

[Glucose] (%)
- 2
- 0.2
- 0.02
- 0.01
- 0.005
- 0.002
- 0.001
- 0.0005
- 0.0002

% of spore bags germinated

Time after adding glucose (minutes)

D

Speed of germination & total % germinated

Average time taken to germinate (minutes)

Total % of spores bags germinated

[Glucose] (%)

Fig. 1

Fig. 2

A Synthetic circuit to test arbitrary gene induction in spores

B Gene expression is possible in spores without any external energy source

C Spore bags vary in ability to produce GFP

(In PBS without nutrients)

Average
Single spore bag

Low [dox.]

Average

High [dox.]

GFP (a.u.)

Time after dox added (hours)

D Inducing GFP does not change % of spore bags that germinate

% of spore bags germinated

Without dox.
With dox.

[Glucose] (%)

E Spore bags with higher GFP inducibility are more likely to germinate

[Glucose] = 0.001%

% of spore bags germinated

Number of spore bags

In PBS 22h after dox

GFP before glucose addition (a.u.) ("GFP inducibility")

F Germination landscape

[Glucose] (in x10⁻³ %)

Probability of germination

GFP before glucose addition (a.u.) ("GFP inducibility")

G GFP inducibility during dormancy quantifies ability to germinate

Very low   Low   Medium   High

GFP inducibility

Minimum [glucose] required for germination

V. low   Low   Med.   High

No germination

Saturating (2%)

Fig. 3

A Inhibit gene expression in dormant spores

Add drug   Add glucose   Dead?

Time

Remove drug

Thiolutin   Cyclohex.   Antimycin A

DNA → RNA → Protein   ATP synthesis

B Transcription during dormancy is essential for survival

% germinating after drug removal

None   Transcription   Translation   ATP synth.

Process inhibited with drug for 24 hours

C Detect all RNAs being made in each dormant spore

DNA
AGTAGA

5-Ethynyl Uridine (U*)

Alexa 488

ŮCAŮCŮ

Newly made RNA

D Quantify all RNAs being made in each dormant spore

Alexa 488   DAPI

With U*

No U*

Fraction of spores

With U*

No U*

Total RNA level (a.u.)

E Total transcription rate poorly correlates with GFP inducibility

dox.

Inducible GFP protein level (a.u.)

R=0.24

Total RNA level (a.u.)

Amount of RNA polymerases highly correlates with GFP inducibility

F RNAP II   RPB3   mCherry

Inducible GFP protein level (a.u.)

R=0.64

RNAP II level (a.u.)

G RNAP I & III   RPC40   mCherry

Inducible GFP protein level (a.u.)

R=0.63

RNAP I & III level (a.u.)

H Ribosomal RNA level poorly correlates with GFP inducibility

rRNA

Inducible GFP protein level (a.u.)

R=0.19

18S rRNA level (a.u.)

I More RNA polymerase means more likely to germinate

% germinated

[Glucose] = 0.0015%

Number of spore bags

RNAP II level before receiving glucose (a.u.)

J Ability to germinate

Amount of RNA polymerases

GFP inducibility w/o nutrients

dox.

Fig. 4

A Spores lose ability to germinate (die) as days pass without glucose

- GFP-strain in amino acids
- GFP-strain kept in water
- Wild-type kept in water

% of spore bags germinated with 2%-glucose

Time without glucose (days)

B Dormant spores gradually lose GFP inducibility

dox.

- Can germinate (dormant) — Sub-populations
- Cannot germinte (dead)

Inducible GFP protein level (folds)

Time in water without glucose (days)

C Amount of RNA polymerase II gradually decreases during dormancy

Day 0   Day 20

BF

RNAP II (RPB3)

RNAP II level (a.u.)

Time in water without nutrients (days)

D More RNA polymerase II means more likely to survive dormancy

Day 39 in water without nutrients

% alive

After receiving 2% glucose

Number of spore bags

RNAP II level before receiving glucose (a.u.)

At edge of dormancy, near death (on day 39 in water without nutrients):

E Dormant spores have same near-zero RNAP II levels as dead spores

Dormant

Dead

Fraction of spores

RNAP II level (a.u.)

F After receiving glucose, dormant spores make new RNAP II but dead spores do not

• Germinates

Dormant

Dead

Fold change in RNAP II level relative to time = 0

Time after glucose (hrs)

2%-glucose added

Production rate of RNAP II (1 / hr)

Dormant

Dead

RNAP II level before receiving glucose (a.u.)

Fig. 5

A
Transient encounter with thiolutin

Thiolutin
DNA → RNA

Add thiolutin | Add dox | Measure GFP
Time
Remove thiolutin

B
Transient encounter with thiolutin permanently removes GFP inducibility

Inducible GFP protein level (a.u.)
- Dormant
- Dead
Time spent in thiolutin (hours)

After transient encounter with thiolutin for 12 hours:

C
RNAP II is still abundant

Fraction of spores
- 12 hrs in thiolutin
- 39 days in water (w/o thiolutin)
RNAP II level (a.u.)

D
Add thiolutin | Add glucose
12 h
Remove thiolutin
Time
Dead?

E
Dead spores have abundant RNAP II but they cannot make more RNAP II

Germinates
Dormant
Dead

Fold change in RNAP II level relative to time = 0

Time after glucose (hrs)
2%-glucose added

Production rate of RNAP II (1 / hr)
Dormant
Dead
RNAP II level before receiving glucose (a.u.)

Fig. 6

Inhibiting transcription or translation shortens lifespan of dormant spores

A

Measure RNAP II

Add drug
Check viability
Days
Dead?

Thiolutin   Cyclohex.
DNA   RNA   Protein

B

RNAP II level (relative to day 0)

8.4 days

Days in water with cyclohex.

C

Days taken for RNAP II level to be halved

☐ No drugs
■ Cyclohex.
■ Thiolutin

D

% alive

19.2 days

Days in water with cyclohex.

E

Half life of dormancy (days)

☐ No drugs
■ Cyclohex.
■ Thiolutin

F

Model for dormancy-to-death transition

Gene-expressing ability during dormancy:
High

— Dormant
— Dormant with translation inhibition
— Dormant with transcription inhibition

Molecules required for gene expression

Low

Glucose cannot restore gene-expressing ability beyond this point

Very low

Alive
Alive
Alive
Dead

Days without nutrients

Gene products   Germinates
Time
Glucose added

Gene products   Germinates
Time
Glucose added

Gene products   Dead
Time
Glucose added

Fig. 7