Integrating the metabolic processes of *Escherichia coli* with virulence by decreasing glucose
availability, inhibiting the acetyl-CoA carboxylase gene *accA* with asRNA, and through the
quantification of the *luxs* gene

Tatiana Hillman[1]
[1] Biological Sciences, TheLAB, Inc., Los Angeles, California, United States
Corresponding author:
Tatianna Hillman[1]
Email address: thillman@usc.edu

ABSTRACT

The study aims to demonstrate a possible link between bacterial cell metabolism and virulence

by combining the metabolic mechanisms of the gram-negative bacteria, *Escherichia coli*.

Glucose increases the proliferation of intestinal microflora, which augments the output of the

short-chain fatty acids. Bacteria ferment glucose into short-chain fatty acids, which help regulate

many biochemical processes and pathways. Each short-chain fatty acid maintains colonic pH,

promotes cell differentiation, and the apoptosis of colonocytes. The long-chain fatty acids are

also synthesized for plasma membrane and biofilm formation. To increase the synthesis of

acetyl-CoA carboxylase, an enzyme that catabolizes glucose into short- and long-chain fatty

acids, *Escherichia coli* was cultured in Luria broth enhanced with a high to a low concentration

of glucose. The 15mM, a high concentration of glucose, yielded qPCR products measured for the

target gene *accA*, which was 4,210ng/µL. The 7.5mM sample produced a concentration equal to

375 ng/µL, and the control sample measured an *accA* concentration of 196 ng/µL. The gene

accA, one of four subunits for the acetyl-CoA carboxylase enzyme, was suppressed by asRNA,

producing a qPCR product of 63 gene copies. Antisense RNA for *accA* reduced the amount of

luxs, a vital gene needed for propagating quorum-sensing signal molecules. The *luxs* gene, which

is responsible for releasing autoinducer 2 for cell-to-cell quorum sensing, was reduced by the

gene inhibition of *accA* with asRNA. The increase in *luxs* transcription augments biofilm

production in support of spreading virulence. The implications of the study advocate for

designing antibiotics that target bacterial cell metabolic processes to block bacterial antibiotic

resistance.

INTRODUCTION

Alexander Fleming discovered penicillin in 1928. His discovery catalyzed the use of

antibiotics that saved many lives, however, with the overuse of antibiotics the issue of antibiotic

resistance has occurred in parallel. Between the 1960s and the 1980s a profound formulation of

many new and novel antibiotics were discovered [1]. The new and novel antibiotics were

superseded by a vigorous evolution of bacterial antibiotic resistance. The three phases that occur

after bacterial horizontal gene transfer for proliferating antibiotic resistance include: 1) the

antibiotic genes received during conjugation code for the creation of efflux pumps; 2) the

antibiotic-resistant genes are enclosed within the pro-antibiotic genes, which can impede

antibiotic drugs attaching to the antibiotic gene target sites; and 3) the antibiotic-resistant genes

can direct the ubiquitous destruction of antibiotic drugs through hydrolysis [11] .

Only a few of the novel antibiotics were available to treat many of the antibiotic resistant

bacteria infections. Therefore, there has been a reduction in exertion and effort within the field of

pharmaceutics to design more new and novel antibiotics. Therefore, future research can

investigate the processes, mechanisms, and the dynamics of antibiotic resistance to increase the

development of novel antibiotics that can target those specific resistant processes. Antimicrobial

resistance or AMR has led to the deaths of 700,000 annually including an additional 10 million

people who is approximated to die by 2050 [1]. Generating novel antibiotics has not solved the

problem of AMR since bacterial resistance has reduced the efficacy of every antibiotic

administered in the clinical setting. There is a demand for more creative and advanced antibiotics

to deal with this tremendous public health issue.

The bacterial colonies within biofilms are profusely resistant to antibiotics and the

biofilm induces the continuance of chronic infections. Because the biofilm extracellular matrix

prevents penetration of antibiotics, is largely anaerobic or lacking oxygen, bacteria within the

biofilm are more tolerant of antibiotics [2]. Biofilms are an organization of bacterial groups

attached to a biotic or abiotic extracellular matrix. However, there is insufficient knowledge

about the effect of antibiotic resistant processes linked to bacterial biofilms. Bacteria use the

formation of biofilm as an advantageous survival strategy.

Biofilm allows bacterial cells to develop a high tolerance for antibiotics through

resistance. The biofilm formed consists of proteins, extracellular DNA, sugars, and is strongly

resistant to proration by antibiotics. When antibiotics are given, the antibiotics eradicate most of

the bacteria cells on the surface of the biofilm, but the bacteria embedded closer to the base of

the matrix remain [3]. Biofilm can endure and persist after antibiotic exposure due to many

physical, anatomical, and innate genetic factors. The pathogenic bacteria create many of their

virulent actions and biofilm production through quorum sensing or cell to cell communication.

Due to the cell-cell communication for biofilm formation, designing small molecules for

disturbing the quorum sensing signals can inhibit biofilm development.

After researchers exposed bacterial cells within biofilm to the antibiotics to rifampicin-

and kanamycin-resistant mutants within immature biofilm communities, the results displayed

that each mutant amplified in growth without the presence of antibiotics. The mutant cultures of

biofilm that were treated with kanamycin showed less antibiotic resistance, however, after 250

generations the resistant mutant bacteria increased in growth at the surface and at the base of the

biofilm matrix [4]. *Pseudomonas aeruginosa* can produce redox reactive pigments. These

pigments are called phenazines, which affect the expression of genes, cause metabolic changes,

and can alter bacterial cells tolerance of antibiotics [5]. Phenazines induce the survival of *P.*

aeruginosa through its ability to modify biofilm physiology.

Phenazines increase the formation of the initial matrix, composed of exopolysaccharide,

which is needed for bacteria to attach and form a mature biofilm. The Phenazines cause the

production of efflux pumps and affect the downward cascade of the central metabolic processes

of *P. aeruginosa*. Researchers measured the bacterial mutant-survivors of biofilm after exposure

to antibiotics and a carbon compound. They found that after biofilms of *P. aeruginosa* were

cultured with succinate, a carbon component that enters the mid-central metabolism of glucose,

the phenazine could no longer protect bacterial cells from antibiotic penetration and eradication.

A bacteria's increased tendency to develop biofilm bacterial populations can be a

significant factor of virulence. For example, most patients suffering with a skin infection known

as *S. epidermidis* produce a strong biofilm in vitro [6]. The exopolysaccharide matrix can lessen

the penetration of the deeper strata of the biofilm by antibiotics. When bacteria are subjected to

antibiotic exposure at deeper layers of the biofilm, resistant mutants of those bacteria can form

[**7**]. Although biofilm communities of bacteria can be exposed to large amounts of antibiotics for

longer periods of time, active and live bacterial cells can persist and survive.

Lux-S is a S-ribosylhomocysteine lyase that produces autoinducer-2. Lux-S is necessary

for the motility of bacteria and for forming bacterial biofilm. Autoinducer-2 is a signaling

molecule released that induces the cell to cell communication called quorum sensing. Bacterial

cells communicate through an intercellular process called quorum sensing. Amplified Lux-S

expression increases the release of autoinducer-2, intensifies quorum sensing, and produces more

biofilm for spreading pathogenic bacteria. The expression of the gene *luxs* induces the release of

autoinducer-2. Quorum sensing is a type of communication between bacterial cells that occurs

through the release of small signaling molecules called autoinducers. Releasing the autoinducers

regulates the aggregation of cells and monitors genetic expression. The autoinducers of gram-

negative bacteria consist of homoserine lactones versus gram-positive bacteria that release

oligopeptides [7].

In addition, the microflora can communicate with the host cells of the gastrointestinal

tract through a variety of chemical idioms [16]. A group of chemical molecules for bacterial

language expression and transfer includes homoserine lactones that are the most heavily studied.

When a bacterial infection occurs, homoserine lactones form and interrelate with the immune

system [17]. An increased release of quorum-sensing signals produces more biofilm, which is

needed for rapid motility and the spreading of infection. The production of biofilm induces

resistance to antimicrobial peptides. Biofilm maintains an intensive resistance to antibiotics as

multi-drug resistant pumps utilize biofilms for excreting antibiotics from bacteria. Mutations

present in genes that are a part of the metabolic pathway cause bacteria to become more

susceptible to antibiotic exposure.

However, the metabolism of glucose by bacteria significantly affects the output of

autoinducers and the production of biofilm. The acetyl-CoA carboxylase enzyme catalyzes the

biosynthesis of long-chain fatty acids and hydrolyzes glucose into short-chain fatty acids. The

LCFAs are synthesized after the acetyl-CoA transferase, the alpha subunit of ACCase or the

ACCA subunit transfers a carboxyl group from pyruvate to an acetyl-CoA, producing malonyl-

CoA, which catalyzes the elongation of the LCFAs. The LCFAs become structured into the

bacterial cell-matrix and the plasma membrane. The bacteria cells grow into large populations,

forcing biofilm formation and induces a quorum-sensing response. The *luxs* gene is essential for

synthesizing and releasing autoinducer-2.

Lux-S is required for quorum sensing, biofilm formation, and virulence. Inhibiting the

fatty-acid synthesis of bacterial cell metabolism can reduce Lux-S expression and biofilm

production. Increased metabolic fatty-acid synthesis can amplify Lux-S expression as a virulent

producer of biofilm, which can cause the persistence of antibiotic resistant bacteria. We do not

completely understand the exact mechanism that connects cell metabolism with the bacterial

pathogenesis of antibiotic resistance. The author suggests that blocking the enzymatic activity of

acetyl-CoA carboxylase may represent a potential aim for overcoming antibiotic resistance.

The metabolic processes of bacteria may provide much potential for generating and

designing novel therapeutics as antibiotics that can reduce biofilm formation. For example,

CbrAB is a histidine kinase that is membrane-bound, and a part of the metabolic regulatory

system called the CbrAB/Crc system [41]. The CbrAB/Crc system shows a connection of

bacterial cell metabolic processes with antibiotic resistance [10]. A process called catabolite

repression control or Crc inhibits translation of catabolite activities by targeting specific mRNAs.

The Crc protein targets mRNAs adjacent to ribosome binding sites [42]. When sources of carbon

availability are limited the CbrAB/Crc system is actuated. The CbrAB protein initiates the

expression of a small RNA termed *crcz*. The crcz alleviates the repression of the Crc-targeted

mRNAs. Inhibiting the *crc* gene lowers the growth rate of bacteria along with providing a higher

susceptibility to antibiotics [10].

Nutrient availability can determine the expression of metabolic genes and remove the

bacterial response of antibiotic resistance. In a study by Peng et al. (2015), bacteria cultured

without glucose or the non-polar amino acid of alanine were less susceptible to kanamycin, an

antibiotic. Only 9.5 ng/mL of the kanamycin was absorbed; however, adding alanine and glucose

to the cultures increased the uptake of kanamycin to 65-123 ng/mL and then to 113-231 ng/mL

[12]. Yu Bin Shu et al. (2018) performed a Michaelis-Menten kinetics experiment using multi-

drug efflux pumps while adding glutamate and acetate to growths of *Escherichia coli* [13]. They

confirmed that the *Escherichia coli* cells were more sensitive to gentamicin, which is an

antibiotic. Each of the previously mentioned studies displays a significant link between bacterial

metabolism and antibiotic resistance.

For that reason, the effects of extracellular nutrient availability on bacterial gene

expression were tested in this study. The concentration of glucose in Luria-Bertani broth media

was altered. RNA was extracted to test the yield of the gene expression of *Escherichia coli*. To

study the intracellular process, the gene expression of *accA*, a subunit of the acetyl-CoA

carboxylase enzyme, was quantified via real-time PCR. The ACCase complex is an important

enzyme for the first step of bacterial fatty-acid synthesis. The cell metabolism within *Escherichia*

coli cells induces the expression of the four genes, *accA*, *accB*, *accC*, and *accD*, which code for

the translation of acetyl-CoA carboxylase. The four genes for acetyl-CoA carboxylase modify

the composition of fatty acids and monitor the rate of production. The four genes also monitor

the overexpression of ACCase by inserting one of the four subunits, activating antisense RNA

expression [9]. To quantify the activity of ACCase, a part of the intracellular metabolism, the

concentration of the gene *accA* was measured after exposure to glucose conjoined with antisense

RNA inhibition.

To measure the effects of genetic inhibition of the intracellular metabolic processes, *accA*

was suppressed and the RNA concentrations recorded. The original bacterial protocell constrains

foreign and movable DNA through the transcription of antisense RNA, which is complementary

to its specific DNA target sequence [14]. The asRNAs bind to the sequences flanking the

ribosome-binding site and the start codon of the target mRNAs [15]. The antisense RNA blocks

the ribosomal detection by the RBS inhibiting translation [15]. In this study, antisense RNA

obtained amplification through the PCR amplification of an antisense DNA sequence. The DNA

sequence converts to an antisense sequence when the DNA is flanked with Xho1 and Nco1

restriction sites. The PCR product was implanted into the pHN1257 plasmid through ligation

(Fig 1). In addition, *Escherichia coli* cells were converted into antisense expressing bacteria after

transforming the competent bacteria cells with the recombinant IPTG-PT-asRNA plasmid called

pHN1257. Real-Time PCR analysis was performed to determine the number of gene copies for

accA.

Figure 1 Plasmid Assembly and the Gene Silencing of *accA*

Plasmid Assembly and Gene Silencing the PCR products and the PTasRNA expression vector

of the plasmid pHN1257 were digested with the restriction enzymes XhoI (upstream) and NcoI

(downstream). After the restriction digestion, the antisense PCR product was ligated into the

IPTG-PTasRNA inducible vector of pHN1257 at the multiple cloning site.

doi: 10.3390/ijms15022773 This article is an open access article distributed under the terms and

conditions of the Creative Commons Attribution license

(http://creativecommons.org/licenses/by/3.0/).

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3958881/figure/f6-ijms-15-02773/

In addition, suppressing the *accA* gene can model the mechanism of the pathogenic

bacteria repression of gene transcription based on the availability of nutrients and any other

physiological change [18]. In order to measure the effects of extracellular and intracellular

factors upon the intercellular communication between cells, the gene copy number of *luxs* will be

quantified. To test the correlation between glucose, *accA*, and *luxs*, the concentration of glucose

was decreased while inhibiting *accA*. The author hypothesizes that inhibiting *accA* while

lessening glucose availability will decrease the *luxs* gene copy number as quantified through

qPCR. The author expects to determine an integration between extracellular glucose,

intracellular metabolic gene expression, the effect of the two preceding stages of metabolism on

intercellular cell-to-cell signaling, and pathogenesis. The ultimate purpose of the study is to

demonstrate the immense potential of targeting the metabolic gene, *accA*, for overcoming

antibiotic resistance.

MATERIALS AND METHODS

BACTERIAL CELL CULTURE AND RNA EXTRACTION

The *Escherichia coli* TOP10 strain of cells were cultured and plated onto 25 mL agar

plates. The bacterial cells were inoculated into Luria-Bertani broth enhanced with glucose

concentrations of 200µM, 50µM, and 0µM. The cultures of 200, 50, and 0 micromolar (n=7) of

glucose were measured for the dsDNA and the RNA OD260 concentrations and repeated as

seven replicates. Varied concentrations of glucose including, 15mM, 7.5mM, 5mM, and control

of glucose were added to LB liquid with *Escherichia coli* cells. The plasmid pHN1257 was

utilized for assembly of the antisense expression vectors. The recombinant plasmid of pHN1257

for the expression of antisense RNA were plated with the kanamycin antibiotic and incubated at

37°C for 24 hrs. Cells grown on agar plates were inoculated into 4 mL of LB liquid media with

(+) and (-) PTasRNA bacterial cells expression of asRNA for *accA* cultured with kanamycin. A

sum of 20 RNA extractions were completed. A total of 3 mL of bacterial cell culture from the

cells grown in the varying concentrations of glucose, the (+) asRNA, and the (-) asRNA cells

were centrifuged at 4,000 x g for 10 minutes at 4°C. The medium was discarded, and the cells

were resuspended in 100μL of lysozyme/TE buffer. The solution was vortexed for 30 seconds.

Incubation occurred at 30°C for 10 minutes in a shaker-incubator. A lysis buffer of 350μL with

25mg of glass beads were added. The lysates were centrifuged for 5 minutes at maximum speed.

RNA was extracted using HiBind RNA mini columns through RNA Wash Buffers I and II. The

RNA was eluted with 50μL of DEPC water.

CDNA PREPARATION

The author used the OneScript Reverse Transcriptase cDNA Synthesis kit by Applied

Biological Materials Inc. or ABM. Approximately, 1μg of RNA from each sample, high glucose,

medium glucose, low glucose, and (-) glucose were added to 0.5μM oligonucleotides(dT)

(10μM). For the asRNA and (-) RNA, 1 microgram of each total RNA extract was added to the

initial primer/RNA and reaction mixture. The cDNA was also prepared for *luxs* quantification.

The initial primer/RNA mix consisted of 1μg of total RNA, 1μL of oligo(dt), 1μL of dNTP

(10mM) mix, and nuclease-free water for a 20μL reaction solution. The mixture was heated to

65°C for 5 minutes and then incubated on ice for 1 minute. The reaction mixture of 4μL 5X RT

buffer, 0.5μL of RNaseOFF, and 1μL OneScript RTase was added to the initial primer/RNA mix

in a 2μL microcentrifuge tube. The cDNA was synthesized by incubating the tube for 50 minutes

at 42°C. The reaction was stopped at 85°C for 5 minutes and chilled on ice. The cDNA was

stored at -20°C.

PCR

A Promega PCR MasterMix volume of 25µL was added to the upstream and downstream

primers each of 0.5µL specific for the *accA* gene target. PCR amplification was repeated as a

triplicate. The concentrations of the two cDNA samples added to the PCR master mix were

190ng and 230ng plus nuclease-free water. A 2X and 1X upstream primer, 10µM 0.5–5.0µl0.1–

1.0µM, and a downstream primer, 10µM 0.5–5.0µl 0.1–1.0µM, DNA template 1–5µl less than

250ng, and nuclease-free water were mixed into a 50µl PCR reaction mixture. The thermocycler

was set to denaturation of a 2-minute initial denaturation step at 95°C. Other subsequent

denaturation steps were between 30 seconds and 1 minute. The annealing step was optimized

with annealing conditions by performing the reaction starting approximately 5°C below the

calculated melting temperature of the primers and increasing the temperature in increments of

1°C to the annealing temperature. The annealing step was set at 30 seconds to 1 minute in 52°C.

For the extension reaction with Taq polymerase, there was 1 minute allowed for DNA to be

amplified at 72°C with a final extension of 5 minutes at 72–74°C. The PCR thermocycler

completed 40 cycles of amplification instead of 35 cycles.

AGAROSE GEL ELECTROPHORESIS AND PLASMID ASSEMBLY

The condition of the PCR products, without primer dimers and contamination, for the

target gene *accA*, were detected through agarose gel electrophoresis. The PCR products were

confirmed without primer dimers and contamination, the PCR products, and the PTasRNA

expression vector of the plasmid pHN1257 were digested with the restriction enzymes XhoI

(upstream) and NcoI (downstream) (New England Biolabs XhoI- catalog R0146S- 1,000 units

and NcoI- catalog R0193S- 1,000 units) (Fig. 1). Each microcentrifuge tube was placed in the

incubator for 30 minutes at 37°C. A heat block was heated to 90°C. The PCR products of the

accA gene were ligated into the pHN1257 plasmid by mixing 1µL of the DNA insert with 2µL of

the plasmids, adding 5µL of ligation mix, and then placing the tubes into a heat block of 90°C

for 15 minutes (Takara Ligation Kit 6023). Competent bacterial cells were transformed with the

pHN1257plasmid plus the antisense DNA insert of *accA*, for expressing (+) asRNA and (-)

asRNA, which is the control. The competent cells were incubated with the recombinant DNA for

45 minutes at 37oC.

REAL-TIME PCR

The author enhanced each bacterial sample with the glucose concentrations of 15mM,

7.5mM, 5mM, and 0mM (n=3). A triplicate of real-time PCR experiments was completed. The

cDNA from H-glucose, M-Glucose, L-glucose, and the control (-) glucose were diluted into a 1:8

serial dilution. A 10-fold dilution of standards was prepared for bacterial samples transformed

with (+) asRNA inducible plasmids and exposed to high to a low volume of glucose. The amount

of gene expression for *accA* and the target gene *luxs* were measured through absolute

quantification methods of qPCR. The diluted cDNA was pipetted into a clear 96 well plate. A

0.4µl of each primer pair mixture was pipetted into a microcentrifuge tube with 10µL of

SensiMix SYBR MasterMix, and with 1.2µL of DEPC water. The SYBR reaction mixture of

12µL was then pipetted into the wells with the diluted and undiluted cDNA. Optical film sealed

the plate. To reposition all liquids to the bottom of the wells, the PCR plates were centrifuged for

2 minutes at 2500 rpm. The real-time PCR Amplification Protocol (Roche Lightcycler 480

machine) began with a 3 min step at 95°C for 40 cycles with a step of 10 sec at 95°C, 45 sec at

65°C, and 20 sec at 78°C with one step of 1 min at 95°C. One step of 1 min at 55°C for 10s each

started at 55°C with a 0.5°C increment at each step up to 95°C. For the Roche Lightcycler 480

machine, the 96-well clear Roche plates were used for SYBR green detection.

THE qPCR ABSOLUTE QUANTIFICATION OF LUX-S

Bacterial cells were transformed with the pHN1257 plasmid expressing asRNA to inhibit

the genetic expression of the *accA* gene. Three replications of qPCR were performed for

quantifying the gene copies of *luxs*. The author added 25µM and 5µM concentrations of glucose

to the bacterial samples expressing antisense RNA for the inhibition of the gene *accA*. RNA was

extracted from cells grown with 25µM glucose and 5µM of glucose. The cells grown with

glucose in the medium were also transformed with antisense expressing IPTG-PT-asRNA

inducible pHN1257plasmids. Cells were cultured without (-) glucose but the expression of *accA*

was also inhibited, (+) asRNA. A control, without (-) glucose and (-) asRNA, was compared to

each sample. The number of gene copies for *luxs* were measured through the qPCR amplification

protocol and absolute quantification (Roche Lightcycler 480 machine).

STATISTICAL ANALYSIS

Figures were constructed with The Sigma Plot 14[th] edition, Microsoft excel, and with

Graph Prism Edition 8. The mean and standard deviations were calculated by Microsoft excel

and GraphPad Prism. The p-values and the ANOVA results were produced through GraphPad

Prism. Standard Error bars for each figure were added through Sigma plot. The standard curves

for the absolute of quantification of the genes *accA* and *luxs* were generated through Microsoft

excel with linear regression coefficients added and calculated for each standard curve for the

absolute quantification of *accA* and *luxs*.

RESULTS AND DISCUSSION

THE EFFECT OF GLUCOSE ON THE RNA CONCENTRATION AND THE

EXPRESSION OF ACCA

Hydrolytic enzymes of *Escherichia coli* split one glucose molecule into two pyruvates. The acetyl-CoA carboxylase enzyme transfers a carboxyl group from the pyruvate to the acetyl-CoA, producing malonyl-CoA (Fig 2). The acetyl-CoA carboxylase enzyme consists of four subunits combined into the AccABCD complex. The *accA* gene codes for the ACCA subunit, which is the transferase domain of the ACCase enzyme. From the results, the genetic expression of *accA* was proportional to the increased glucose concentrations in *Escherichia coli* bacterial samples. After culturing *Escherichia coli* with different levels of glucose, the RNA concentrations for each sample measured were of 1392, 797, 608, and 199 ng/ μL for the H-Glucose, M-Glucose, the L-Glucose, and the control, respectively. The genes of *accA* and *accD* are entirely independent of the *accBC* operon.

When *accB* is overexpressed, *accBC* transcription is blocked, but the overexpression of the other gene products, *accA* and *accD*, don't affect the *accBC* operon transcript levels [21]. A high level of glucose increases the availability of *accB*, and the transcription of the *accBC* operon is greatly reduced even after exposure to a small amount of *accB*. Although a high glucose concentration can stymie the production of *accBC*, the over-activated *accBC* operon did not inhibit the production of *accA*, the target gene of study. The subunit ACCA encoded in the *accA* gene may act as a more intense target for blocking bacterial cell fatty acid synthesis and metabolism. The *Escherichia coli* TOP10 bacteria cells were grown in LB broth overnight at 37°C with 15mM (High-glucose), 7.5mM (Medium-glucose), 5mM (Low-glucose), and a control of glucose concentration (Fig 3). RNA was extracted from each sample and the concentrations

determined by the Implen Nanophotometer 250. The high-glucose sample produced the highest

amount of RNA compared to the medium, the low, and to the control. The RNA for each sample

was reverse transcribed into first-strand cDNA and absolute quantification with qPCR measured

the amount of the target gene, *accA*, of each sample.

Figure 2 Formation of Malonyl-CoA A glucose molecule is hydrolyzed into two pyruvate

molecules. The acetyl-CoA Carboxyl transferase or the ACCA alpha subunit of the acetyl-CoA

carboxylase complex, AccABCD, transfers a carboxyl group from the pyruvate to the acetyl-

CoA to form malonyl-CoA. The malonyl-CoA enters the fatty acid synthesis pathway,

elongating fatty acids into long-chain fatty acids of phospholipids necessary for structuring

plasma membranes and for forming the initial extracellular matrix.

A) **Figure 3 Glucose concentration versus RNA** Glucose concentrations of 15mM, 7.5mM,

5mM, and the control samples each yielded RNA concentrations, in ng/uL, 1392, 797, 608, 179,

respectively. A) The bacterial sample with 15mM glucose yielded the largest measure of RNA,

noting a proportional relation between glucose and genetic expression of *E. coli*. B) Includes

qPCR results for 15mM, 7.5mM, 5mM, and the control. The samples with 15mM and 5mM

glucose displayed the most genetic activity of *accA* transcription, measuring *accA* concentrations

of 4,210 ng/uL and 1,500 ng/uL respectively. The *accA* qPCR concentrations for 7.5mM *accA*

equaled 372 ng/uL and a 196 ng/uL quantity for the control. C) Displays the OD260 measure of

DNA results for samples of *E. coli* grown in medium enhanced with 200uM, 50uM, 0mM of

glucose.

Glucose mM and [RNA]

A)

Glucose mM and qPCR of accA

B)

Glucose 200uM, Glucose 50uM and Gluc...

C)

The RNA of each bacterial sample enhanced with glucose was extracted and the

concentrations determined (N=20, 335.4 ± 329.2). The high-glucose sample provided a crossing

point value or a Cp-value of 12.28 and the concentration of *accA* was 4.21E3 ng/µL. The Cp of

the sample of medium-glucose equaled 16.51 with a concentration of 3.75E2 and the low-

glucose Cp was 14.08 with a target gene concentration of 1.50E3 (Fig 3). The concentration of

accA by qPCR analysis, in ng/µL, included: 4210, 375, 1500, and 196 for H-glucose, M-glucose,

L-glucose, and for the control, respectively. The number of qPCR products ranged from 11-186,

5-18, and 0-79 gene copies for High-glucose, Medium-glucose, and the Low-glucose/control,

respectively (N=12, 48±38).

The control group produced a Cp value of 17.64 with a target gene concentration of

1.96E2. For the glucose concentrations of 200µM (n=7,173±72), 50 µM (n=7,116±105), and the

control (n=7, 157±64) compared to their dsDNA concentrations showed significant variability

(Fig 3).Comparing 200 µM to its control group, the difference between the samples was

statistically significant (p=0.038). The cells grown in 200µM yielded a more significant and

larger production of dsDNA than cells cultured as a control. After loading a 96-well plate for

qPCR with 8-fold dilution standards of each sample, the standards of each sample were used to

calculate a standard curve (Fig 3).

Vinals et al., (1999), found results that determined that a high glucose concentration

causes the absorption of glucose to be blocked as a result of its repression of glucose

phosphorylation [19]. Ferraris et al. (1990), measured the luminal glucose osmolarity and the

concentrations in the small intestines. They found that the luminal glucose concentrations in a

range of 50mM to 500mM exhibited errors [20]. The previous researchers measured glucose

concentrations but did not recognize how osmolality is affected by Na+ and K+ salts, amino

acids, and peptides [20]. Ferrais et al. (1990), discovered the average of SI luminal osmolarities

were approximately 100 mosmol/kg, which were mainly hypertonic results. In the Ferrais et al.

(1990) study, for an animal's diet, the SI glucose concentrations averaged 0.4-24 mM and ranged

with time over a large amount of an SI region from 0.2 to 48 mM. They also found that high

concentrations of glucose lessened the absorption of glucose in endothelial cells called ECV304

cell lines. There was a 60 percent reduction of 2-deoxyglucose uptake at 30 mM glucose from

their study [20]. The high glucose osmolality resulted in a reduced Vmax value for 2-

deoxyglucose uptake with a constant Km, calculated from the Michaelis-Menten kinetic enzyme

equation [20].

THE QPCR RESULTS OF THE ANTISENSE RNA INHIBITON OF THE TARGET

GENE ACCA

383   Ninety percent of the genome for Eukaryotes consists of non-coding regions or RNA that

will not translate into proteins. Antisense transcription is the purpose of the noncoding regions of

human cell DNA [22]. The non-coding transcripts are called natural antisense RNAs. Antisense

activity has been recorded in mice, *Saccharomyces cerevisiae*, *Drosophila melanogaster*, and

Arabidopsis thaliana [23]. Recombinant DNA was produced when the PCR product of the gene

insert, *accA*, was ligated into the plasmid pHN1257 (Fig. 1), which was engineered to amplify

antisense RNA. The PCR product and the IPTG-PT-asRNAs plasmid of pHN1257 were cut with

the restriction enzymes XhoI and NcoI. The primers (Table. 1) each contained an extra 3 to 4 bps

of nucleic acids to accompany the sequences for Xhol and Nco1. The primers flanked the target

DNA, which totaled to 150 base pairs. The antisense sequence was constructed in a specific

orientation in that the restriction sites, which NcoI normally flanks the forward primer, reversed

positions. The XhoI was repositioned to flank the forward primer, and the Nco1 flanked the

reverse primer, creating the antisense sequence when inserted into the plasmid for pHN1257.

The PCR product was amplified (n=3, 219 ± 171), and then ligated into the pHN1257-asRNA

inducible vector.

Table 1 Primers for amplifying the *luxs* gene, antisense RNA for *accA*, and the *accA* gene.
399

Primers	DNA Sequence
XhoI+*accA* Forward	GAGATGAGTCTGAATTTCCTTGATT
NcoI+*accA* Reverse	TGGCAGTTCATCGCTTTTTTCAC
luxs Forward	CATACCCTGGAACATCTGTTTGC
luxs Reverse	AGTTCCTGCACTTTCAGCACATC
accA Forward	TCATCACCTTTATCGACACCCC
accA Reverse	TTCACTTTATCGCCCACGCC

400
401
Primers for PCR Amplification and qPCR Analysis. To produce antisense RNA for *accA*

inhibition the restriction enzyme site for XhoI was sequenced with the forward primer for *accA*.

The restriction enzyme NcoI was added to the reverse primer for *accA*. Through PCR

amplification, the PCR products were composed of antisense DNA for the *accA* gene. The PCR

product would be inverted when inserted into the IPTG-PT-asRNA inducible pHN1257 plasmid,

forming recombinant DNA plasmids for asRNA transcription in vitro. The plasmids with

recombinant antisense DNA were transformed into bacterial competent cells for qPCR analysis

of the *accA* gene. The primers for the *luxs* gene were applied to quantify the number of gene

copies from qPCR analysis for cells grown in glucose enhanced samples. The primers for the

accA gene were used for the quantification of the *accA* gene from the samples cultured with an

increasing gradient of glucose solutions.

The total RNA concentration for the control, (-) asRNA, equaled 739.44 ng/µL. The

control, (-) asRNA, yielded a miRNA concentration of 334.98 ng/µL (Fig 4). The RNA

concentration for transformed bacterial cells with the asRNA of *accA* pHN1257plasmid was

279.28ng/µL. The miRNA concentration of the cells with asRNA measured to 240.90 ng/µL.

The gene of *accA* was successfully suppressed by asRNA in vitro with 63 gene copies, qPCR

measured for bacteria cells (n=3) transformed with the recombinant antisense pHN1257 plasmid

DNA (37.8 ± 35.6). The real-time PCR results for the bacterial cells without the antisense gene

target and insert yielded 421.69 ng/µL for *accA* (Fig 5). There was a 147 % difference between

cells not expressing asRNA versus cells transcribing the asRNA for *accA* (p=0.027). Inhibiting

or lessening ACCase activity limits the synthesis of lipids, causing many physical changes.

Antisense RNA expression can modify the consistency of fatty acids by upregulating the

production of the ACCase enzyme [24].

Figure 4 RNA Concentrations vs. Glucose Concentration A) *E. coli* bacteria cultures (n=3)

were enhanced with 200µM, 50µM, and a control of glucose (59±49). B) The 50µM of glucose

in a bacterial sample highest yield was 566ng/µL, for 200uM 117ng/µL of RNA was produced,

and the control measured to 76ng/µL of RNA.

A)

B)

Figure 5 The qPCR results for quantifying *accA* concentration and the gene copy number.

The RNA for each sample, 15mM, 7.5mM, 5mM, and the control, was reverse transcribed into

first strand cDNA and absolute quantification with qPCR was used to measure the amount of the

target gene, *accA*, produced by each sample. A) High glucose had a Cp of 12.28 and the

concentration of *accA* was 4.21E3 ng/uL. The Cp of sample medium-glucose equaled 16.51 with

a concentration of 3.75E2 and the low-glucose Cp was 14.08 with target gene concentration of

1.50E3. B) The gene of *accA* was successfully suppressed by asRNA in vitro with 63 gene

copies measured for bacteria cells transformed with the recombinant antisense pHN1257 plasmid

DNA. The bacterial cells with the pHN1257 plasmid but without the antisense gene target and

insert produced 421.69 ng/uL for *accA*. There was a 147% percent difference between cells not

expressing asRNA versus the cell transcription of the asRNA for *accA* (p=0.027).

A)

B)

During short-chain fatty acid synthesis, malonyl-CoA is formed from acetyl-CoA, using

energy from ATP and bicarbonate production [21]. Glucose is hydrolyzed into pyruvate, which

is made into acetyl-CoA, forming acetate (Table.2). Through the Wood-Ljugdahl pathway,

pyruvate loses two hydrogens, carbon dioxide loses oxygen, altered into carbon monoxide, and a

methyl group is added to the oxidation, synthesizing acetyl-CoA into Acetate [25]. Butyrate is

produced when two molecules of acetyl-CoA combine into acetoacetyl-CoA. The acetoacetyl-

CoA is transformed into butyryl-CoA, or butyrate [20]. Propionate is composed through the

succinate or acrylate pathway [26]. The phosphoenolpyruvate, or PEP, is broken into pyruvate

which is further metabolized with water into succinate. The succinate is reduced into propionyl-

CoA, forming propionate (Table. 2). Through the acrylate pathway, lactate loses oxygen,

forming propionate [25].

Table 2 Pathways for Biosynthesis of SCFAs from Carbohydrate Fermentation For

synthesis of each SCFA, glucose is hydrolyzed into 2 molecules of pyruvate, which then lose a

hydrogen, forming Acetyl Coenzyme A. For butyrate the Acetyl coenzyme A loses a water,

forming B-hydroxy butyryl CoA, where it loses a hydrogen and the CoA is replaced with a

Phosphate, the phosphate is removed. After the phosphate is removed, butyrate is formed.

Acetate forms when the CoA in Acetyl Coenzyme A is replaced with a phosphate group, and

then the phosphate is added to an ADP to make ATP. Propionate can be formed when lactate

loses an oxygen, giving Acryloyl-CoA, converted into Propionyl-CoA, which the CoA is

replaced with a phosphate, and then removed by a kinase, to produce propionate.

Glucose	Glucose	Glucose
2 Pyruvates Lose a Hydrogen	2 Pyruvates Lose a Hydrogen	2 Pyruvates Lose 1 Hydrogen Gain 1 Hydrogen
Acetyl CoEnzyme A	Acetyl Coenzyme A	LACTATE Loses 1 Oxygen
B-hydroxybutyryl CoA Lose H2O	Phosphate Replaces The CoA	Creates Acryloyl-CoA
Crotonyl CoA Gains a Hydrogen	Creates Acetyl-Phosphate	Forms Propionyl-CoA
Butyryl CoA Phosphate Replaces The CoA	Phosphate added to ADP to make ATP	CoA in Propionyl-CoA Replaced with a Phosphate
Butyryl Phosphate formed The Phosphate moves to an ADP to form ATP **BUTYRATE**	**ACETATE**	Phosphate removed **PROPIONATE**

The LCFAs are synthesized when the ACCA subunit of ACCase transfers a carboxyl

group to acetyl-CoA converting into malonyl-CoA, which then enters the fatty acid elongation

cycle (Fig 6). The fatty acid cycle synthesizes long chains of fatty acids for the formation of the

bacterial plasma membranes and a thin-layered matrix for initial bacterial attachment to a biotic

or abiotic surface. After bacteria attach to a surface, more growth occurs until a microcolony of

bacterial cells form. In addition, the gene *accA* codes for the subunit of the Acetyl-CoA

Carboxylase complex. Acetyl-CoA carboxylase (ACC) initiates the first step of the fatty acid

synthesis. An acyl-group or an ACP is added to the malonyl-CoA and converted into a ketoacyl-

ACP. The Beta-ketoacyl-ACP is reduced, gains a hydrogen, into Beta-hydroxyacyl-ACP. The

Beta-hydroxyacyl-ACP is dehydrated, losing a molecule of water, to alter into an Enoyl-ACP,

which forms into an Acyl-ACP by gaining 2 hydrogen protons.

Figure 6 Long-Chain Fatty Acid Synthesis An acyl-group or an ACP is added to the

malonyl-CoA and converted into a ketoacyl-ACP. The Beta-ketoacyl-ACP is reduced, gains a

hydrogen, into Beta-hydroxyacyl-ACP. The Beta-hydroxyacyl-ACP is dehydrated, losing a

molecule of water, to alter into an Enoyl-ACP, which forms into an Acyl-ACP by gaining 2

hydrogen protons.

QUANTIFICATION OF LUX-S GENE EXPRESSION

The gene *accA* is a component of the ACCA subunit of the ACCase complex, which functions as a transferase. The ACCA subdomain of ACCase is vital for catalyzing fatty acid synthesis. Lipid biosynthesis is responsible for increased quorum sensing and biofilm formation. After glucose is hydrolyzed into two pyruvate molecules, the alpha subunit, ACCA, transfers a carboxyl group to acetyl-CoA to produce malonyl-CoA. The malonyl-CoA enters the lipid biosynthesis pathway. The fatty acids produced assist with secreting a more fluid membrane, stretching the span of the biofilm matrix. As the biofilm matrix expands the microcolonies of bacteria grow proportional to the size of the matrix. The increase in the number of colonies induces the release of autoinducers, which accrue into a display of bioluminescence. The bioluminescence is a signal of quorum sensing. The gene *luxs* is needed to produce autoinducer-2. *luxs* monitors the amount of biofilm, flagellum movement, and regulates the factors of virulence. The ACCA subunit catalyzes the synthesis of lipids required for inducing the release of AI-2, which are signals produced during quorum sensing.

The high levels of autoinducers initiate a positive feedback loop switching all bacterial cells into producing quorum signals, which create bioluminescence [20]. As autoinducers compile in the surrounding environment of a bacterial cell, the bacterial population increases [5]. The bacteria sense the growth in the population and begin tracking the changes in the number of cells, which help to change the amount of genetic expression [28]. Quorum sensing between bacterial cells monitors the genes responsible for organizing the activities executed by different bacterial groups. Activities completed through quorum sensing include bioluminescence, spore formation, antibiotic excretion, creating a biofilm, and releasing factors for virulence [5]. Biofilm can increase the movement of pathogenic bacteria into the bloodstream [7].

Therefore, there can exist a significant link between bacterial cell metabolism and

virulence. By quantifying the measurement of gene copies for *luxs* while inhibiting *accA* with

asRNA, the connection of metabolism with pathogenesis could be demonstrated (Fig 7).

Escherichia coli cells were grown in 25μM and 5μM of glucose and transformed with antisense

pHN1257expressing IPTG-PT-asRNA plasmids. Cell samples with 5μM glucose- (+) asRNA

produced a high measure of the *luxs* gene [11]. The gene copy number of *luxs* for (+) glucose/

(+) asRNA, and for (+) asRNA were quantified. For the results, the (-) glucose-(+) asRNA

sample produced a gene copy number of 199 and 2511 *luxs* copies for (-) glucose- (-) asRNA

(n=3, 277 ± 37). For the 25μM glucose-(+) asRNA and 5μM glucose (+), asRNA produced

39,810 and 316,228 gene copies of *luxs*, respectively (Fig 8). The control samples, the (-)

glucose-(-) asRNA, (n=3, 658,114 ± 483,499) and the 5μM glucose- (+) asRNA (n=3,

171571±204574) exhibited the highest amount of expression for *luxs* with a $p<0.05$ (p=0.025)

(Fig 8). An ANOVA analysis of the qPCR gene copies for the *luxs* gene is displayed in Table 3.

Figure 7 Gene Copies for *luxs* determined through qPCR measurement

The qPCR Results for *luxs* the number of gene copies for *luxs* was measured through qPCR and

absolute quantification A) (-) glucose-(+)asRNA sample had a gene copy number of 199. B)

There were 2511 *luxs* copies for (-)glucose-(-)asRNA. The 5uM glucose-(+)asRNA yielded

316,228 gene copies of *luxs* D) For the 25μM glucose-(+)asRNA 39,810 gene copies were

produced.

549 **A)**   **B)**

551
552
553

554   **C)**   **D)**

555
556

Figure 8 Total Production of *luxs* versus the Control Sample. The polar plot compares the

control sample for the qPCR quantification of the gene *luxs* after exposure to glucose and

antisense RNA in the form of the pHN1257-IPTG-asRNA expression plasmid. A) The control

compared to the asRNA sample, to the 25micromolar-glu-asRNA sample, and versus the 5**μM**-

glu-asRNA sample equaled the following polar coordinates (radius, theta): (1×10^6, 180°), (5×10^5,

60°), and (6x10^5,68°), respectively. B) The control, 5, 25, and the asRNA micromolar samples

production of the *luxs* gene are displayed in the 3D contour graph as an intense red, purple/blue

for the 5micromolar-glu-asRNA, and purple for the asRNA including the 25micromolar-glu-

asRNA sample. The control sample yielded a higher measure of *luxs* with $1x10^6$ gene copies

(658,114 ± 483,499). The sample without glucose but with asRNA expression produced 199

gene copies of *luxs* (n=3, 277 ± 37).

569
570
571
572   **A)**

588 **B)**

LUX-S Gene Copy Number (qPCR)

591 **Table 3 ANOVA results for the total experimental sample for quantifying gene copies**
592 **through qPCR.**

ANOVA Results of qPCR Gene Copies

	luxs-glu/(+) asRNA	*luxs*-glu/(+) asRNA	(-)glu(-)asRNA	*accA* (-)asRNA (+)glu/ (-) glu	*accA* (+)asRNA (+)glu/ (-) glu
1	2511	316228	1000000	186	100
2	2000	26915.4	1000000	79	32
3	741	17783	39810	18	5
4	199	251	151286	7.4	3
n	4	4	4	4	4

X	1363.750	90294.350	547774.000	72.600	35.000
s	1074.010	151028.260	524165.009	81.920	45.306
X_{ave}	127907.940				
total	19	1798315304447.048			

source	df	SS	MS	F	P-value
treatments	4	905636341810.108	226409085452.527	3.8044	0.0250
error	15	892678962636.940	59511930842.463		
total	19	1798315304447.04			

The ANOVA results reveal an overall mean of 127907.940 for the total number of trials that

yielded a total and a sum of gene copies through Real-Time PCR absolute quantification. The

sample without glucose and without antisense RNA, (-)glucose-(-)asRNA, produced the highest

amount of gene expression, for *luxs* and *accA* with 547774.000 as a mean of the total number of

gene copies yielded from qPCR analysis p=0.025.

The 5µM glucose- (+) asRNA sample expressed more of the *luxs* gene than the 25µM

sample. Wang et al., (2005) found adding 0.8 percent glucose to their bacterial culture and

growth medium increased the activity at the promoter site of the *luxs* gene. A glucose

concentration of 5µM results in a normal exogenous osmolarity surrounding bacterial cells

within the microflora in the gut of the lower digestive tract [24]. Because cells were overly

saturated with a 25µM concentration of glucose, it was expected to increase the level of *luxs*

gene expression. However, the excess saturation of glucose at 25µM did not result in the

amplified gene expression of *luxs*. Jesudhasan et al. (2010) proved that higher levels of

autoinducer-2 did not lead to the increased gene expression of *luxs* in *Salmonella typhimurium*

when cultured in high concentrations of glucose [25].

Glucose affected the rate of transcription, increased the production of nucleic acids, but

the inhibition of *accA* with asRNA significantly suppressed gene expression. From the control to

the 25 µM glucose exposure, bacteria cells transcribed DNA at an amplified rate (Fig 9). As the

glucose concentration increased to 50 µM the rate of transcription reached a sustained

equilibrium with no new nucleic acid production or activity as the concentration approached 200

µM. The samples expressing asRNA inhibition of *accA* produced 16% of the total yield of

nucleic acid. Samples not expressing asRNA inhibition, yielded 83% of the total quantity of

nucleic acid.

Moreover, by transforming *Escherichia coli* bacterial cells with IPTG-PT-asRNA

inducible vectors, which targeted the gene *accA*, the gene expression of *luxs* gene also decreased.

Because glucose delays the movement of the *luxs* mutant strain with bacterial overpopulation but

does not inhibit bacterial growth, each (+) glucose-(+) asRNA bacterial colony appeared smaller

and defected. Osaki et al., (2006) showed that *luxs* mutants can be smaller in diameter with 8.0

mm versus 12.3 mm in its wildtype [29]. The bacterial colonies expressing antisense RNA

(5mm) were a smaller size versus cells without antisense RNA transcription (8mm) of *accA*. The

cell samples with (+) asRNA transcription were found by many preceding research studies to

appear smaller in size [21].

Bacterial samples transformed with IPTG-PT-asRNA inducible pHN1257 plasmids and

designed to inhibit the *accA* gene displayed the least measure of *luxs* expression. The (-) glucose-

(+) asRNA sample exhibited the lowest amount of *luxs* expression of 199 gene copies (225 ±

37). As a result, when the *luxs* gene is mutated with less expression, the biofilm is thinner,

looser, and displaying an appearance of bacterial cells with defects. Lux-S forms a thicker and

more viscous biofilm, which is a result of many nucleic acids released into the extracellular

matrix of the biofilm.

Figure 9 The effect of glucose on the rate of transcription, inhibition with asRNA, and the

frequency of Nucleic Acid production. A) From the control to the 25 µM glucose exposure,

bacteria cells transcribed DNA at an amplified rate. As the glucose concentration increased to 50

the rate of transcription reached a sustained equilibrium with no new Nucleic Acid production or

activity. B) The gene *accA* was inhibited with asRNA. The samples expressing asRNA inhibition

of *accA* produced 16% of the total yield of nucleic acid. Samples not expressing asRNA

inhibition, yielded 83% of the total quantity of nucleic acid.

A)

641
642
643   **B)**
644

The ample amounts of DNA are released to maintain a more solid macromolecular matrix

and consistency of the biofilm [30]. Lux-S is required for AI-2 production and for regulating

gene expression in the early-log-growth phase. Pathogenic bacteria depend on biofilm formation

to attach to epithelial cells and tissues, spreading infectious diseases. Pathogenic bacteria infect

host cells by accumulating AI-2, exogenously, which then increases the amount and consistency

of the formation of biofilm (31). The gene *luxs* codes for the flagella complex, biofilm creation,

and controls virulent factors. The *luxs* gene, when it is more expressed, causes the spread of

virulent bacteria through increasing the motility and movement of pathogenic bacteria. More

Lux-S expression, as a result of mutations in the *luxs* gene, produces increased amounts of

biofilm, assisting pathogenic bacteria to invade the mucosal layers and the epithelial cells of its

host, developing into a systemic infection.

The autoinducer-2 that accumulates, monitors cell growth by permitting bacterial cells to

sense the excess AI-2 and then terminate population growth. The excess AI-2 distributes quorum

signals between bacterial cells, which notifies cells of limited nutrients, surface area, or space.

Autoinducer-2 requires the genetic expression and translation of *luxs* (Fig 8). When Lux-S and

autoinducer-2 are less regulated and overexpressed, infectious bacteria can propagate in stressful

environments with high acidity and salinity [23]. Kendall et al., (2008) demonstrated the ability

of *S. pyogenes* to adapt to acidic conditions as a result of an unregulated Lux-S/AI-2 system [32].

The *Vibrio fischeri* process of quorum sensing differs drastically compared to the

Escherichia Coli. *Vibrio fischeri* synthesizes AI-2 through an enzyme called an autoinducer

synthase or the Lux I that produces a hormone class of acyl-homoserine lactones, AHL, which is

the autoinducer 3OC6-homoserine lactone. In *Vibrio fischeri* the AHL binds to the autoinducer

receptor called Lux-R for initiating a transcriptional activator of DNA [27]. After AHL is

produced, it passes through the cell's plasma membrane as the number of bacterial cells increase,

and then it stops diffusing after cell density reaches a threshold. When the AHL is attached to the

LuxR complex, the operon is activated, transcribing the genes that code for the enzyme

luciferase [27]. The LuxR-AHL bond releases, into the periphery of the cell, high levels of

quorum sensing molecules that stimulate the cell to cell communication between bacteria species

[24].

The *Vibrio fischeri* process of quorum sensing immensely differs in comparison to

Escherichia Coli. In addition*, Escherichia Coli* and many gram-negative bacteria also regulate

the influx and output of autoinducer-2. In *Escherichia Coli* the transporter complex called

LsrABCD regulates the uptake of autoinducer-2. The repressor of LsrABCD and the LsR is a

kinase that is a part of the quorum sensing system of AI-2. The cyclic AMP/cyclic AMP receptor

proteins regulates the lsr operon. The cyclic AMP/cyclic AMP receptor proteins are expressed

from *lsrk* and *lsrR* that is positioned upstream of lsr [8]. Lsrk that is a kinase located in the

cytoplasm phosphorylates AI-2 that then attaches to the *lsr* repressor. The activated and attached

AI-2 suppresses the *lsr* repressor LsrR. After the activity of the *lsr* repressor is inhibited, the

release of AI-2 increases via the autoinducer-activated expression of the Lux-S protein.

A glucose molecule is hydrolyzed into two pyruvate molecules. The acetyl-CoA

Carboxyl transferase or the ACCA alpha subunit of the acetyl-CoA carboxylase complex,

AccABCD, transfers a carboxyl group from the pyruvate to the acetyl-CoA to form malonyl-

CoA. The malonyl-CoA enters the fatty acid synthesis pathway, elongating fatty acids into long-

chain fatty acids of phospholipids necessary for structuring plasma membranes and for forming

the initial extracellular matrix. The microcolonies attach, aggregate, and accumulate in the

extracellular matrix. As the colonies accumulate into large populations a release of autoiducer-2

molecules is triggered, causing a certain bioluminescence through an exchange of cell- to-cell

quorum sensing signals. The release of autoinducer-2 signals cells to increase population growth

or to preempt it. The gene *luxs* is needed for releasing AI-2. The excess AI-2 distributes quorum

signals between bacterial cells, which notifies cells of limited nutrients, surface area, or space. If

bacterial cells continue to accrue, more biofilm is produced within the extracellular matrix,

resulting in the dispersion of bacterial cells. Biofilm is a potent virulent factor. Pathogenic

bacteria can disperse from and cause a systemic infection through excess biofilm production.

To form biofilm for infection, the *luxs* gene must be activated. Its excessive amplification

and expression form a denser texture of biofilm [30]. Lux-S monitors virulence through

regulation of the process for generating biofilm in the nasopharynx in mice with pneumonia,

Streptococcus pneumoniae. Flagella expression increases with *luxs* mutation in the lungs and

bloodstream. The *luxs* mutants can infect the lungs or the bloodstream more rapidly than its wild-

type strain [27]. Flagellin increases inflammation by activating Toll-Like Receptor Signaling or

TLR5 pathways, which translates pro-inflammatory genes within the Mitogen-Activated Protein

Kinase or the MAPK pathways. Flagella modulates virulence and pathogenesis by allowing more

rapid motility of bacteria, infecting the colonization of host cells, and assisting infectious

bacterial cells into entering the mucosal layers [33].

Pathogenic bacteria determine the level of autoinducer signals specific for their species.

Pathogenic bacteria require a specific order and assortment of virulent genes to cause infection.

For example, the different residues or R-groups of amino acids within autoinducers bind to the

LUXR protein receptors in a conformation-specific to the amino acid side chains within the

binding sites. The orientation of the AI to protein receptor binding sites produces varied types of

side-chain lengths and amino acid substitutions [34]. For example, pathogenic bacteria as

Enteropathogenic *Escherichia coli* EPEC release intraspecies-type of signals that help to

colonize the small intestines because the small intestines are void of much commensal

microflora. The Enterohemorrhagic *Escherichia coli* or EHEC pathogenic form of *Escherichia*

coli conducts signals with other diverse bacterial cell types and with host cells. The EHEC

communicates through quorum sensing with other colonic bacteria cells. The EHEC infect

epithelial cells through activating the genes called the Locus of Enterocyte Effacement or LEE.

The genes of LEE are required for A/E lesions to form when EHEC cells attach to and efface

from epithelial cells, amplifying the level of pathogenicity [27].

When the production of Lux-S is blocked, the intensity of disease-causing bacteria

changes. For example, the mutation of *luxs* in *Haemophilus influenzae* increases virulence that

causes middle ear infections [35]. The *A. pleuropneumoniae* and *M. haemolytica* bacteria have

autoinducer mutations that lessen virulence in mouse models. When the *luxs* gene is inhibited,

there is a lesser amount of pustule-creation than the parental wild type strain [35]. However,

autoinducer 3 is not dependent on the expression of the *luxs* gene. *luxs* genetic mutants impede

the production of AI-3 (21-22). As a result, the pathogenic Escherichia coli, EHEC, collects

signals from host cells in the form of hormones known as epinephrine. The AI-3 can

communicate with the transmembrane protein of the host cell called Qsec [36]. The two

components for quorum sensing also include Las and RhI that are quorum systems consisting of

activator proteins. The activator proteins direct the synthesis of the autoinducers. The proteins

called Las and RhI turn genes on or off based on the density of bacterial populations, sensing the

accumulation of autoinducers [37].

The bacteria called *P. aeruginosa* secretes two types of autoinducers called N-3-

oxododecanoyl, a homoserine lactone, (3-oxo C12-HSL) and N-butyryl-L-homoserine lactone

(C4-HSL). These autoinducers released can cause virulence in people infected by *P. aeruginosa*

by supplying virulent factors that help colonize and spread the infection [37]. Therefore, the

results present significant data for the effect of glucose and genetic inhibition of *accA* on the

production of the *luxs* gene, an important component of bacterial virulence. The data

demonstrates that the genetic changes, as inhibiting the *accA* gene with antisense RNA, and the

availability of nutrients such as glucose, affected the production of *luxs*. Inhibition of the *luxs*

gene can reduce biofilm formation and decelerate the motility of bacteria, which the lessened

motility can impede the spread of pathogenic bacteria.

CONCLUSION

The overall purpose of the study was to confirm and identify the gene *accA* as a possible

target for blocking bacterial antibiotic resistance. Through examining the role of *accA*, a link

between bacterial cell metabolism and virulence was attempted and demonstrated. A cumulative

perspective, integrating bacterial cell metabolism via the ACCA subunit with bacterial

pathogenicity, may provide possible metabolic targets for designing new and potent antibiotics.

It was previewed in this study, an ability to design antibiotics that target bacterial cell metabolic

genes as *accA*, the binding sites of the ACCase enzyme, and the fatty acid synthesis pathway.

Targeting all three areas of bacteria cell metabolism can reduce the formation of bacterial cell

biofilm needed for spreading pathogenic bacteria. The strong condition of a bacterial

population's metabolic state can prevent bacterial susceptibility to antibiotics [34]. The results of

decreased RNA expression and cell growth in the presence of antisense RNA inhibition of *accA*

may serve as a model for blocking microbial metabolic processes that contribute to antibiotic

resistance [38].

Escherichia coli cells were cultured in high glucose to a control concentration and

showed a decreasing range of gene expression for the target gene, *accA*. The observation of the

bacterial cells, grown in an increased amount of available glucose, showed the highest measure

of transcription for *accA*. The effect of glucose extracellular of the bacteria, on the concentration

of RNA, was measured by modulating the glucose concentration gradient. The transcription of

intracellular molecules as the *accA* gene was determined through qPCR. The *accA* gene was

inhibited with antisense RNA, using an IPTG-inducible vector, pHN1257. There was a 138

percent difference of gene expression between cells transformed with antisense pHN1257 versus

cells negative for antisense RNA during in vitro transcription. The recombinant DNA and

plasmids were assembled from the IPTG-PT-asRNA of the pHN1257 (Fig 1).

The constructed plasmid of PHN257 consisted of flanking inverted repeats for the target

DNA, rebuilding a paired double-stranded RNA terminus that inhibited the transcription of *accA*.

The inhibition of *accA* was confirmed through the qPCR results of 63 gene copies for *accA* (+)

asRNA versus 422 gene copies for *accA* with (-) asRNA. The sum volume of *luxs* was

determined through qPCR. By increasing the glucose concentration in increments, Wang et al.

(2005) found that glucose increased the levels of autoinducer-2 of the culture as well [39]. The

observations displayed the RNA concentration for bacterial cells, expressing asRNA of *accA*,

included a concentration of 279.28 ng/µL. The miRNA concentration of the cells with asRNA

measured to 240.90 ng/µL.

The results supported the hypothesis. Decreasing the glucose concentration, while

suppressing the *accA* gene, led to an observed decrease in *luxs* expression. The author

hypothesized that decreasing the sum of glucose and suppressing *accA* gene expression would

decrease the expression of the *luxs* gene. The *luxs* gene is vital for quorum sensing and releasing

autoinducer-2. Increased expression of Lux-S produces more AI-2, which induces and initiates

quorum sensing between bacterial cells. The increased AI-2 molecules released triggers quorum

sensing for the amplified production of biofilm. Biofilm formation allows more pathogenic

bacteria cells to attach, proliferate, detach, and spread into a systemic infection.

The *accA* gene translated into the acetyl-CoA carboxyl transferase enzyme transfers a

carboxyl group from pyruvate to the acetyl-CoA to form a malonyl-CoA. More glucose induces

more production of the acetyl-CoA carboxyl transferase. Increased glucose also causes more AI-

2 quorum-sensing signal molecules to accumulate. The malonyl-CoA enzyme is important for

catalyzing the biosynthesis of long-chain fatty acids. The long-chain fatty acids are the building

blocks for biofilm formation. Therefore, a decrease of biofilm production can occur due to the

limited enzymatic activity of the ACCA transferase subunit via *accA* gene inhibition.

However, in the presence of glucose with antisense RNA transcription, *Escherichia coli*

cells continued to produce amplified amounts of *luxs* with 316,228 copies measured through

qPCR when enhanced with 5µM of glucose (n=3, 171571± 204574). The limitations of this

study include the contradictory results of the 5µM glucose-(+) asRNA sample for *luxs*. Adding a

5µM concentration of glucose to the LB broth culture increased the gene expression of *luxs*

although the *accA* gene was inhibited. These contradictory and unexpected results may show

early signs of reduced efficacy of the inhibition of the *accA* gene in the presence of glucose.

However, it is probable that the 5uM of glucose triggered increased expression in the

accD gene and of the ACCD subunit, which was not inhibited by asRNA. The gene *accA* can

continue to act as a potent metabolic target for suppressing antibiotic resistance. This study

suggests that designing antibiotics could include targeting metabolic genes and enzymes. During

the past 30 years, only a few potent antibiotics have been formulated and discovered [40].

However, there is a new motivation for examining bacterial targets located in metabolic

pathways. For future research, studying the gene expression, structure, and the function of the

ACCase complex can increase the incentive for designing small-molecule inhibitors specific for

the binding sites of bacterial-metabolic enzymes.

ACKNOWLEDGMENTS

Special thanks are given to the staff from TheLAB in Los Angeles, CA. They were responsible

for ordering and providing all materials for this project. Many thanks are given for their

commitment to assisting me with tips and solutions for laboratory procedures. Thank you to their

staff for helping me to frame the structure of this project. They were always available for

answering questions and giving much guidance. Thank You!

REFERENCES

1.Calero-Cáceres, W., Ye, M., & Balcázar, J. L. (2019). Bacteriophages as environmental
reservoirs of antibiotic resistance. *Trends in microbiology*.

2.Rojo-Molinero, E., Macià, M. D., & Oliver, A. (2019). Social behavior of antibiotic resistant
mutants within Pseudomonas aeruginosa biofilm communities. *Frontiers in Microbiology*, *10*,
570.

3.Singh, N., Romero, M., Travanut, A., Monteiro, P. F., Jordana-Lluch, E., Hardie, K. R., ... &
Alexander, C. (2019). Dual bioresponsive antibiotic and quorum sensing inhibitor combination
nanoparticles for treatment of Pseudomonas aeruginosa biofilms in vitro and ex vivo.
Biomaterials science.

4.France, M. T., Cornea, A., Kehlet-Delgado, H., & Forney, L. J. (2019). Spatial structure
facilitates the accumulation and persistence of antibiotic-resistant mutants in biofilms.
Evolutionary Applications, *12*(3), 498-507.

5.Schiessl, K. T., Hu, F., Jo, J., Nazia, S. Z., Wang, B., Price-Whelan, A., ... & Dietrich, L. E.
(2019). Phenazine production promotes antibiotic tolerance and metabolic heterogeneity in
Pseudomonas aeruginosa biofilms. *Nature communications*, *10*(1), 762.

6.Ardon, C. B., Prens, E. P., Fuursted, K., Ejaz, R. N., Shailes, J., Jenssen, H., & Jemec, G. B. E.
(2019). Biofilm production and antibiotic susceptibility of Staphylococcus epidermidis strains
from Hidradenitis Suppurativa lesions. *Journal of the European Academy of Dermatology and*
Venereology, *33*(1), 170-177.

7.Soares, A., Caron, F., & Etienne, M. (2019). Commentary: tolerance and resistance of pseudomonas aeruginosa biofilms to antimicrobial agents–how p. Aeruginosa can escape antibiotics. *Frontiers in Microbiology*, *10*, 2164.

8.Li, J., Attila, C., Wang, L., Wood, T. K., Valdes, J. J., & Bentley, W. E. (2007). Quorum sensing in Escherichia coli is signaled by AI-2/LsrR: effects on small RNA and biofilm architecture. *Journal of bacteriology*, *189*(16), 6011-6020.

9.K. Ali, and T. Aruna. Recent advances in acetyl CoA carboxylase: a key enzyme of fatty acidbiosynthesis in plants. Intl JAppl Biol Pharmac Technol. 2016:264-277.

10.Martínez JL, Rojo F. Metabolic regulation of antibiotic resistance. FEMS microbiology reviews. 2011 Sep 1;35(5):768-89.

11.Hillman T. 2019. Current methods for inhibiting antibiotic resistant bacteria by targeting bacterial cell metabolism and disrupting antibiotic elimination through the AcrAB-Tolc efflux pump. PeerJ Preprints 7:e27840v1 https://doi.org/10.7287/peerj.preprints.27840v1

12.Su YB, Peng B, Li H, Cheng ZX, Zhang TT, Zhu JX, Li D, Li MY, Ye JZ, Du CC, Zhang S. Pyruvate cycle increases aminoglycoside efficacy and provides respiratory energy in bacteria. Proceedings of the National Academy of Sciences. 2018 Feb 13;115(7): E1578-87.

13.B.R. tenOever, The Evolution of Antiviral Defense Systems. Cell Host Microbe. 2016; 19:142–149. doi: 10.1016/j.chom.2016.01.006.

14.N. Nakashima, T. Tamura. Conditional gene silencing of multiple genes with anti-senseRNAs and generation of a mutator strain of Escherichia coli. Nucleic Acids Res. 2009;103 (37)

15.C.M. Waters, B.L. Bassler. Quorum sensing: cell-to-cell communication in bacteria. Annu. Rev. Cell Dev. Biol. 2005;21: 319-346.

16.Telford, G., Wheeler, D., Williams, P., Tomkins, P.T., Appleby, P., Sewell, H., Stewart, G.S., Bycroft, B.W. and Pritchard, D.I. The Pseudomonas aeruginosaquorum-sensing signal moleculen(3-oxododecanoyl)-l-homoserine lactone has immunomodulatory activity. Infection and immunity. 1998; 66(1):36-42.

17.Wilson, J. W., Schurr, M. J., LeBlanc, C. L., Ramamurthy, R., Buchanan, K. L., and Nickerson, C. A. Mechanisms of bacterial pathogenicity. Postgraduate medical journal. 2002;78(918):216-224.

18.Francesc Vinals, Aachim Gross, Xavier Testar, Manuel Palacin, Peter Rosen, ¨ Antonio Zorzano, Highglucose concentrations inhibit glucose phosphorylation, but not glucose transport, in human endothelialcells, Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 1999;1450(2)119-129, ISSN 0167-4889

19.R.P. Ferraris, S. Yasharpour, K.C.K. Lloid, R. Mirzayan, J.M. Diamond, (1990) Luminal glucose concentration in the gut under normal conditions. Am. J. Physiol. 1990;259, 820–837.

20. M.M. Kendall, D.A. Rasko, V. Sperandio. Global effects of the cell-to-cell signaling
moleculesautoinducer-2, autoinducer-3, and epinephrine in a luxS mutant of enterohemorrhagic
Escherichia coli." Infection and Immunity. 2007;75(10):4875-4884.

21.Siller, M., Janapatla, R.P., Pirzada, Z.A., Hassler, C., Zinkl, D. and Charpentier, E., 2008.
Functional analysis of the group A streptococcal luxS/AI-2 system in metabolism, adaptation to
stress andinteraction with host cells. BMC microbiology. 2008;8(1), p.188.

22.Labandeira-Rey, M., Janowicz, D.M., Blick, R.J., Fortney, K.R., Zwickl, B., Katz, B.P.,
Spinola, S.M. and Hansen, E.J. Inactivation of the Haemophilus ducreyi luxS gene affects the
virulence ofthis pathogen in human subjects. Journal of Infectious Diseases. 2009;200(3):409-
416.

23.Osaki, T., Hanawa, T., Manzoku, T., Fukuda, M., Kawakami, H., Suzuki, H., Yamaguchi, H.,
Yan,X., Taguchi, H., Kurata, S. and Kamiya, S. Mutation of luxS affects motility and infectivity
ofHelicobacter pylori in gastric mucosa of a Mongolian gerbil model. Journal of medical
microbiology. 2006;55(11):1477-1485.

24.J. Georg, R.H. Wolfgang. cis-antisense RNA, another level of gene regulation in bacteria.
Microbiology and Molecular Biology Reviews. 2011;75(2): 286-300.

25.U. Emilio. Shifts in metabolic hydrogen sinks in the methanogenesis-inhibited
ruminalfermentation: a meta-analysis. Frontiers in Microbiology. 2015;6(37).

26.Jesudhasan, P.R., Cepeda, M.L., Widmer, K., Dowd, S.E., Soni, K.A., Hume, M.E., Zhu, J.
and Pillai, S.D. Transcriptome analysis of genes controlled by luxS/autoinducer-2 in
Salmonellaenterica serovar Typhimurium. Foodborne pathogens and disease.2010;7(4):399-410.

27.Vendeville, A., Winzer, K., Heurlier, K., Tang, C.M. and Hardie, K.R.,
Making'sense'ofmetabolism: autoinducer-2, LuxS and pathogenic bacteria. Nature Reviews
Microbiology. 2005;3(5):383.

28.S.T. Freestone, B.L. Bassler. Bacterial Quorum Sensing: Its Role in Virulence and
Possibilities for Its Control. Cold Spring Harbor Perspectives in Medicine. 2012;2(11): a012427.
doi:10.1101/cshperspect. a012427.

29.Vidal, J.E., Howery, K.E., Ludewick, H.P., Nava, P. and Klugman, K.P., 2013. Quorum-
sensingsystems LuxS/autoinducer 2 and Com regulate Streptococcus pneumoniae biofilms in a
bioreactor withliving cultures of human respiratory cells. Infection and immunity. 2013;81(4):
1341-1353.

30.Trappetti, C., Potter, A.J., Paton, A.W., Oggioni, M.R. and Paton, J.C. LuxS mediatesiron-
dependent biofilm formation, competence, and fratricide in Streptococcus pneumoniae. Infection
and Immunity. 2011;79(11):4550-4558.

31. Peng B, Su YB, Li H, Han Y, Guo C, Tian YM, Peng XX. Exogenous alanine and/or glucose
plus kanamycin kills antibiotic-resistant bacteria. Cell metabolism. 2015 Feb 3;21(2):249-62.
930

931 32. M.M. Kendall, D.A. Rasko, V. Sperandio. Global effects of the cell-to-cell signaling
932 moleculesautoinducer-2, autoinducer-3, and epinephrine in a luxS mutant of enterohemorrhagic
933 Escherichia coli." Infection and Immunity. 2007;75(10):4875-4884.
934

935 33. Yang, Y., Zhou, M., Hou, H., Zhu, J., Yao, F., Zhang, X., Zhu, X., Hardwidge, P.R. and Zhu,
936 G. Quorum-sensing gene luxS regulates flagella expression and Shiga-like toxin production in
937 F18abEscherichia coli. Canadian journal of microbiology. 2014;60(6):355-361.
938

939 34. A. Jayaraman, K.W. Thomas, Bacterial quorum sensing: signals, circuits, and implications for
940 biofilms and disease. Annu. Rev. Biomed. Eng. 2008; 10:145-167.
941

942 35. Vidal, J.E., Ludewick, H.P., Kunkel, R.M., Zahner, ¨ D. and Klugman, K.P. The
943 LuxSdependent quorum-sensing system regulates early biofilm formation by Streptococcus
944 pneumoniae strainD39. Infection and immunity. 2011;79(10):4050-4060.
945

946 36. E.S. James, J.E. Cronan. Expression of two *Escherichia coli* acetyl-CoA carboxylase
947 subunits isautoregulated. Journal of Biological Chemistry. 2004;279(4): 2520-2527.
948

949 37. Tateda, K., Ishii, Y., Horikawa, M., Matsumoto, T., Miyairi, S., Pechere, J.C., Standiford,
950 T.J., Ishiguro, M. and Yamaguchi, K. The Pseudomonas aeruginosa autoinducer N-3-
951 oxododecanoylhomoserine lactone accelerates apoptosis in macrophages and neutrophils.
952 Infection and immunity. 2003;71(10), pp.5785-5793.
953

954 38. Baquero F, Martínez JL. Interventions on metabolism: making antibiotic-susceptible bacteria.
955 mBio. 2017 Dec 29;8(6):e01950-17.
956

957 39. Wang, L., Hashimoto, Y., Tsao, C.Y., Valdes, J.J. and Bentley, W.E. Cyclic AMP (cAMP)
958 and cAMP receptor protein influence both synthesis and uptake of extracellular autoinducer 2
959 inEscherichia coli. Journal of bacteriology. 2005;187(6):2066-2076.
960

961 40. Polyak SW, Abell AD, Wilce MC, Zhang L, Booker GW. Structure, function and selective
962 inhibition of bacterial acetyl-coa carboxylase. Applied microbiology and biotechnology. 2012
963 Feb 1;93(3):983-92.
964

965 41. Zhang, X. X., & Rainey, P. B. (2008). Dual involvement of CbrAB and NtrBC in the
966 regulation of histidine utilization in Pseudomonas fluorescens SBW25. *Genetics*, *178*(1), 185-
967 195.
968

969 42. Sonnleitner, E., Valentini, M., Wenner, N., el Zahar Haichar, F., Haas, D., & Lapouge, K.
970 (2012). Novel targets of the CbrAB/Crc carbon catabolite control system revealed by transcript
971 abundance in Pseudomonas aeruginosa. *PloS one*, *7*(10), e44637.
972
973

985