Single cell ATAC-seq in human pancreatic islets and deep learning upscaling of rare cells reveals cell-specific type 2 diabetes regulatory signatures

Vivek Rai[1,*], Daniel X. Quang[1,*], Michael R. Erdos[2], Darren A. Cusanovich[3,5], Riza M. Daza[3], Narisu Narisu[2], Luli S. Zou[2], John P. Didion[2], Yuanfang Guan[1], Jay Shendure[3,*], Stephen C.J. Parker[1,4,*,‡], Francis S. Collins[2,*,‡]

[1] Department of Computational Medicine & Bioinformatics, University of Michigan, Ann Arbor MI 48109,
[2] National Human Genome Research Institute, National Institutes of Health, Bethesda MD 20892,
[3] Department of Genome Sciences, University of Washington, Seattle WA 98109,
[4] Department of Human Genetics, University of Michigan, Ann Arbor MI 48109,
[5] Department of Cellular and Molecular Medicine, University of Arizona, Tucson AZ 85721

[*] Contributed equally to this work.
[‡] Corresponding authors; University of Michigan, Departments of Computational Medicine & Bioinformatics and Human Genetics, 100 Washtenaw Ave, Ann Arbor, MI 48109, USA, scjp@umich.edu; National Human Genome Research Institute, National Institutes of Health, Bethesda, MD 20892, USA, collinsf@od.nih.gov

ABSTRACT

Objective: *Type 2 diabetes (T2D) is a complex disease characterized by pancreatic islet dysfunction, insulin resistance, and disruption of blood-glucose levels. Genome-wide association studies (GWAS) have identified >400 independent signals that encode genetic predisposition. More than 90% of the associated single nucleotide polymorphisms (SNPs) localize to non-coding regions and are enriched in chromatin-defined islet enhancer elements, indicating a strong regulatory component to disease susceptibility. Pancreatic islets are a mixture of cell types expressing distinct hormonal programs, and so each cell type may contribute differentially to the underlying regulatory processes that modulate T2D-associated transcriptional circuits. Existing chromatin profiling methods such as ATAC-seq and DNase-seq, applied to islets in bulk, produce aggregate profiles that mask important cellular and regulatory heterogeneity.*

Methods: *We present genome-wide single cell chromatin accessibility profiles in >1,600 cells derived from a human pancreatic islet sample using single-cell-combinatorial-indexing ATAC-seq (sci-ATAC-seq). We also developed a deep learning model based on the U-Net architecture to accurately predict open chromatin peak calls in rare cell populations.*

Results: *We show that sci-ATAC-seq profiles allow us to deconvolve alpha, beta, and delta cell populations and identify cell-type-specific regulatory signatures underlying T2D. Particularly, we find that T2D GWAS SNPs are significantly enriched in beta cell-specific and cross cell-type shared islet open chromatin, but not in alpha or delta cell-specific open chromatin. We also demonstrate, using less abundant delta cells, that deep-learning models can improve signal recovery and feature reconstruction of rarer cell-populations. Finally, we use co-accessibility measures to nominate the cell-specific target genes at 104 non-coding T2D GWAS signals.*

Conclusions: *Collectively, we identify the islet cell-type of action across genetic signals of T2D predisposition and provide higher-resolution mechanistic insights into genetically encoded risk*

48 *pathways.*
49
50 *Keywords: Islet; Epigenomics; Deep Learning; Single cell; Chromatin; Type 2 diabetes*
51
52 *Abbreviations: Assay for Transposase Accessible Chromatin Sequencing (ATAC-seq), Genome wide*
53 *association study (GWAS), Expression quantitative trait loci (eQTL), Glucagon (GCG), Insulin (INS),*
54 *Somatostatin (SST).*
55

1. INTRODUCTION

Pancreatic islets consist of a cluster of at least five different endocrine cell-types (alpha, beta, delta,
gamma, and epsilon), each producing a unique hormone in a distinct but coordinated manner [1].
Collectively, these clusters of cells work together to maintain insulin production and glucose
homeostasis. Disruption of the complex interplay between the cell types, their organization, and their
underlying regulatory interaction is known to be associated with type-2-diabetes (T2D) pathophysiology
[2]. However, the exact cellular mechanisms through which different risk factors contribute to the
disease risk are not completely understood. Using GWAS and eQTL mapping approaches, recent
studies have discovered >400 independent signals (>240 loci) associated with T2D and T2D-
associated traits [3], although remarkably, more than 90% of them localize to non-protein-coding
regions of the genome [4]. Growing evidence suggests that many of these variants likely influence the
RNA expression and cellular function of human pancreatic islets by altering transcription factor binding,
critical components of a cellular regulatory network [5–9].
69
High-throughput epigenomic profiling methods such as ATAC-seq [10] and DNase-seq [11] have
enabled profiling of chromatin accessibility across samples in a tissue-wide manner, providing the
opportunity to identify millions of context-specific regulatory elements. However, these bulk-
measurements of chromatin accessibility limit the precise understanding of how tissue heterogeneity
and multiple cell-types in the population contribute to overall disease etiology [12]. Recent advances in
single-cell transcriptomic and epigenomic profiling methods have enabled an unbiased identification of
cell-type populations and regulatory elements in a heterogeneous biological sample. By mapping the
chromatin-regulatory landscape at a single-cell resolution, recent single-nuclei studies have
demonstrated the potential to discover complex cell populations, link regulatory elements to their target
genes, and map regulatory dynamics during complex cellular differentiation processes [13–16]. The
pancreatic islet gene expression landscape has been investigated at single-cell resolution in existing
studies [17,18], but chromatin accessibility studies have been limited to fluorescence-activated cell
sorting (FACS) methods for obtaining cell-type populations [19,20]. FACS based methods will miss
identification of unknown or rarer cell-populations and are unable to produce pure cell-type populations
due to reliance on the specificity of cell-surface markers [21,22].
85
Here, we present a genome-wide map of chromatin accessibility in >1,600 nuclei derived from a human
pancreatic islet sample using single-nucleus-combinatorial-indexing ATAC-seq (sci-ATAC-seq) [23].
sci-ATAC-seq enables us to deconvolve cell populations and identify cell-type-specific regulatory
signatures underlying T2D. Notably, we find that T2D GWAS SNPs are significantly enriched in beta
cell-specific and cross cell-type shared islet open chromatin, but not in alpha or delta cell-specific open
chromatin. We also demonstrate, using the less represented delta cell population (< 5% of total islet
population), that the use of deep learning can improve signal recovery and feature reconstruction for
less abundant cell-populations. We anticipate that our deep learning method will enable analysis of
heterogeneous tissues that may be harder to obtain in large numbers or contain rare sub-populations.

Collectively, these results identify the islet cell-type of action across genetic signals of T2D
predisposition and provide higher-resolution mechanistic insights into genetically encoded
pathophysiology.
98
2. MATERIALS AND METHODS
2.1 Bulk Islet ATAC-seq
Sample processing. The human pancreatic islet samples were procured and processed as described
by Varshney et al [8]. Briefly, the islets were obtained from the National Disease Research Interchange
(NDRI) and processed according to the NHGRI institutional review board-approved protocols. The islet
was shipped overnight from the distribution center. Upon receipt, we pre-warmed the islet to 37 degree
in shipping media for 1-2h before harvest. ~50-100 islet equivalents (IEQs) were harvested and
transposed in triplicate following the methods in Buenrostro et al [10]. The ATAC-seq library was
barcoded and sequenced 2 x 125bp on a HiSeq 2000.
108
ATAC-seq analysis. Sequencing adapters were trimmed using cta (v0.1.2) [24] and aligned to hg19
reference genome using BWA-MEM (v0.7.15-r1140, options: -I 200,200,5000) [25]. Picard
MarkDuplicates (v2.18.27) was used for duplicate removal and samtools [26] was used to filter for
autosomal, properly-paired and mapped read pairs with mapping quality >= 30 (v1.9, options: -f 3 -F
3340 -q 30). Replicates across each sample were merged into a single file using samtools merge. For
peak calling, each sample was downsampled to 25 million (M) reads and converted to BED file. We
then used MACS2 [27] to call broad peaks (v2.1.1.20160309; options: --nomodel --broad --shift -100 --
extsize 200 --keep-dup all --SPMR) and removed those with FDR >0.05 and overlapping with ENCODE
hg19 blacklists [28]. ATAC-seq coverage tracks were displayed using UCSC Genome Browser and
Integrative Genomics Viewer (IGV). Summary statistics were calculated using Ataqv (v1.0) [29] and are
available in interactive and downloadable format online (**Table S7**). For comparative purposes, we
performed the same read trimming, alignment, filtering, downsampling, and peak calling steps on
publicly available ATAC-seq data (**Table S1**). Peaks from each sample were merged to create a master
peak set and Spearman correlation was computed on the RPKM normalized read-count matrix.
123
124 *Determination of high-confidence peaks.* We randomly sampled 2.5 M reads from each sample using
125 samtools view and pooled them into one file so that each sample is equally represented. Peaks were
126 called on the pooled file as discussed in the previous paragraph. We then determined the number of
127 samples overlapping with each master peak using peaks called on individual samples.
128
129 *Overlap of reads with chromHMM states.* We tested for enrichment of ATAC-seq peaks across 13 islet-
130 specific chromatin states using Genomic Association Tester (GAT) [30]. We ran GAT (v1.3.5, options: --
131 number-samples 10,000) and filtered chromatin states with no significant enrichment (Bonferroni
132 adjusted p-value < 0.05) of peaks in them. The log2 fold enrichment values across chromatin states
133 were clustered using hierarchical clustering of the correlation matrix.
134
2.2. sci-ATAC-seq analysis
Sample processing. We used the combinatorial cellular indexing method to generate single-nuclei
chromatin accessibility data as previously described in Cusanovich et al [23]. Briefly, a suspension of
islet cells were obtained and pelleted 5 min at 4°C 500 x g. The media was aspirated and the cells were
washed once in 1 ml PBS. The cells were pelleted again for 5 min at 4 °C 500 x g and then
resuspended in 1 ml of cold lysis buffer (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.1%
IGEPAL CA-630, supplemented with 1X protease inhibitors (Sigma P8340)). Nuclei were maintained on

ice whenever possible after this point. 10 µl of 300 µM DAPI stain was added to 1 ml of lysed nuclei for sorting. To prepare for sorting, 19 µl of Freezing Buffer (50 mM Tris at pH 8.0, 25% glycerol, 5 mM MgOAc$_2$, 0.1 mM EDTA, supplemented with 5 mM DTT and 1X protease inhibitors (Sigma P8340)) was aliquot into each well of a 96-well Lo-Bind plate. 2,500 DAPI$^+$ nuclei (single cell sensitivity) were sorted into each well of the plate containing Freezing Buffer. The plate was sealed with a foil plate sealer and then snap frozen by placing in liquid nitrogen. The frozen plate was then transferred directly to a -80 °C freezer. Subsequently, the sample was shipped from NIH to UW overnight on dry ice. The plate was then thawed on ice and supplemented with 19 µl of Illumina TD buffer and 1 µl of custom indexed Tn5 (each well received a different Tn5 barcode). The nuclei were tagmented by incubating at 55 °C for 30 min. The reaction was then quenched in 20 mM EDTA and 1 mM spermidine for 15 min at 37 °C. The nuclei were then pooled and stained with DAPI again. 25 DAPI$^+$ nuclei were then sorted into each well of a 96-well Lo-bind plate containing 11.5 µl Qiagen EB buffer, 800 µg/µl BSA, and 0.04% SDS. 2.5 µl of 10 µM P7 primers were added to each sample and the plate was incubated at 55 °C for 15 min. 7.5 µl of NPM was then added to each well. Finally, 2.5 µl of 10 µM P5 primers were added to each well and the samples were PCR amplified with following cycles: 72 °C 3min, 98 °C 30s, then 20 cycles of 98 °C for 10 s, 63 °C for 30 s, 72 °C for 1 min. The exact number of cycles was determined by first doing a test run on 8 samples on a real-time cycler with SYBR green (0.5X final concentration). PCR products were then pooled and cleaned on Zymo Clean&Concentrator-5 columns (the plate was split across 4 columns) eluting in 25 µl Qiagen EB buffer and then all 4 fractions were combined and cleaned using a 1X Ampure bead cleanup before eluting in 25 µl Qiagen EB buffer again. The molar concentration of the library was then quantified on a Bioanalyzer 7500 chip (including only fragments in the 200-1000 bp range) and sequenced on an Illumina NextSeq at 1.5 pM concentration.

QC and pre-processing.

Step 1. Barcode correction and filtering. Each barcode consists of four 8-bp long indexes (i5, i7, p5, and p7). Reads with barcode combinations containing more than 3 edit distance for any index were removed. If a barcode was within 3 edits of an expected barcode and the next best matching barcode was at least 2 edits further away, we corrected the barcode to its best match. Otherwise, the barcode was classified as ambiguous or unknown.

Step 2. Adapter trimming and alignment. Adapters were removed using Trimmomatic [31] with NexteraPE adapters as input (ILLUMINACLIP:NexteraPE.fa:2:30:10:1:true TRAILING:3 SLIDINGWINDOW:4:10 MINLEN:20) and aligned to hg19 reference using BWA-MEM (v0.7.15-r1140, options: -I 200,200,5000) [25]. The final alignment was filtered using samtools to remove unmapped reads and reads mapping with quality < 10 (-f3 -F3340 -q10) as well as reads that were associated with ambiguous or unknown barcodes.

Step 3. Deduplication and nuclei detection. Duplicates from the pruned file were removed using a custom Python script on a per-nucleus basis. Using the distribution of reads per barcode, we applied bi-culstering, as implemented in the mclust [32] R package, to differentiate between background barcodes and barcodes that correspond to a nucleus. Using the list of non-background barcodes, we split the aggregate bam file into constituent bam files corresponding to each barcode representing a single nucleus using a custom Python script.

Step 4. Quality assessment of each single nucleus. For each single nucleus, we computed ATAC-seq quality metrics such as fragment length distribution, transcription start site (TSS) enrichment, short-to-

mononucleosomal reads ratio, total autosomal reads, and fraction of reads overlapping peaks. We
removed nuclei with a) total reads outside 5% to 95% range [34578, 226755] of all the nuclei, and b)
TSS enrichment of <2.7 (5%-tile) from further downstream analysis.

Step 5. Aggregate sci-ATAC-seq peaks. We pooled reads from filtered barcodes from the previous
steps to create an aggregate bam file. Peaks were called and filtered as described previously in the
Bulk Islet ATAC-seq analysis section.

2.3. Cluster analysis
Feature selection and clustering. We generated a list of TSS distal peaks (>5 kb away from the nearest
TSS based on RefSeq genes [33]) from the aggregate sci-ATAC-seq data. For each nucleus, we
counted the number of reads overlapping the peaks using the Rsubread package [34]. We then
adopted a logistic regression approach to remove peaks where binarized accessibility across nuclei
was significantly associated (Bonferroni corrected p-value < 0.05) with sequencing depth. This
approach should help to reduce the bias associated with sequencing depth, as the remaining peaks are
no longer associated with this technical factor, a strategy that has been successfully implemented in
single cell RNA-seq data analysis [35]. The resulting count matrix was RPKM-normalized and
reweighted using the term-frequency and inverse-document-frequency (TF-IDF) method [13]. To do
this, we first weighted all the sites for individual nuclei by the total number of sites accessible in that cell
(''term frequency''). We then multiplied these weighted values by log(1 + the inverse frequency of each
site across all cells), the "inverse document frequency." The TF-IDF transformed matrix was then
reduced to 30 principal components using Principal Component Analysis (PCA) and used as input to
generate a two-dimensional embedding using the Uniform Manifold Approximation Method (UMAP,
n_neighbors = 20) [36]. We identified clusters in the two-dimensional embedding in an unsupervised
manner using a density-based clustering method (hdbscan, minPts = 20) as implemented in dbscan R
package [37].

Cell identity assignment. The cell identities were assigned based on de-facto cell-type-specific hormone
markers: *INS-IGF2* (beta), *GCG* (alpha), *SST* (delta) etc. A marker gene was said to be present in a
nuclei if a read mapped within 5 kb of the GENCODE (v19) gene body annotation [38]. For additional
verification of cell-identity, we computed RPKM normalized aggregate ATAC-seq signal across cell-type
marker genes reported in an islet scRNA-seq study [17]. Finally, we evaluated the enrichment of cells
from each cell-type cluster relative to their expected population proportion using two-sided binomial test
across ten bins of sequencing depth (~145 cells/bin).

2.4. Deep learning signal and peak upscaling
Model. The U-Net model [39] takes input sequences and outputs prediction sequences. The goal of
model training is to reduce the error between the prediction output and a representation of ground truth.
For signal upscaling, the input sequence is base-wise scores of BAM pileups (read-depth)
corresponding to a subsample of n cells (randomly sampled from 600 cells) and the output sequence is
base-wise scores of BAM pileups using reads from all 600 cells. Peak upscaling not only uses the
subsampled BAM pileup scores as inputs, but it also uses the binary base-wise values from calling
peaks with MACS2 on the subsampled BAM alignments. Output sequences for peak upscaling are the
binary base-wise values from calling peaks with MACS2 on the data. We created two models, each
separately on data from alpha and beta cells. Because both had different number of constituent single-
nuclei, we matched the size of output dataset by randomly sampling 600 cells from each cell-type
cluster. The input datasets were created by sampling n cells from the set of 600 cells such that the total

number of reads is similar across both models.
237
The network architecture of the U-Net used in this study is illustrated in **Fig. S2A**. It consists of a
contracting, convolutional path (left side) and an expansive, deconvolutional path (right side). The
contracting path consists of repeated applications of two kernel size 11 convolutions (unpadded
convolutions) with rectified linear unit (ReLU) activation, and a kernel size 2 max pooling operation with
stride 2 for downsampling. Each downsampling step halves the length of the activation sequence while
doubling the number of feature channels. Every step in the expansive path consists of a kernel size 2
deconvolution layer with a linear activation function that halves the number of feature channels, a
concatenation with the correspondingly cropped feature map from the contracting path, and two kernel
size 11 convolution layers with ReLU activations. The cropping is necessary due to the loss of border
sequence steps in non-padded convolution. At the final layer a kernel size 1 convolution with either an
ReLU (for signal upscaling) or sigmoid (for peak upscaling) activation function generates the sequence
of predictions. Due to the use of unpadded convolutions, the prediction sequence is shorter than the
input sequence by a constant border width. Although the U-Net can accept arbitrary length input
sequences, we fix all training samples to be of length 6700, which results in output prediction
sequences of length 4820. In total, the network has five steps each in the contracting and expansive
paths for a total of 27 convolutional layers and 8,998,529 training parameters. The model was
implemented using Keras [40] with the Tensorflow [41] backend, and experiments were run using Titan
Xp and GTX 1080 Ti GPUs.
256
To reduce overfitting, we split chromosomes into training, validation, and testing sets. The model was fit
using the ADAM optimizer [42] with a learning rate of 1e-5 and a batch size of 128 for 50 epochs.
Separate loss functions, and hence models, were used to solve signal and peak upscaling. For signal
upscaling, we used the mean squared error base-wise loss function. For peak upscaling, the loss
function was the sum of the cross entropy base-wise loss and the Dice-coefficient loss, also known as
F1 score. We used mean average precision, a common evaluator for object detection, and Pearson
correlation as the output evaluation metrics for peak and signal upscaling, respectively. This
downscaling and model training were repeated for n=5, 10, 28, 50, 100, 200, 300, 400, and 500 cells.
265
266 *Generating upscaled peaks.* In order to select a subset of high-confidence peaks from the predicted
267 model output, we adopted a post-hoc approach where we compared the number of cell-type-specific
268 peaks for alpha, beta, and delta cells, and chose a threshold where they had a similar number. For
269 predicted delta cell peaks, we combined the results from alpha and beta models at the same threshold
270 using bedtools [43] intersect (v.2.27.1) after filtering for the chosen threshold.
271
272 **2.5. Cell-type-specific peaks analysis**
273 *Cell-type-specific peaks.* Peaks specific to each cell-type were obtained by comparing peaks in one
274 cell-type with all other cell-types using bedtools.
275
276 *T2D GWAS SNPs enrichment.* Enrichment of T2D associated GWAS SNPs from DIAMANTE [3] was
277 tested using GREGOR (v1.3.1) [44]. Specifically, we used the following parameters: r2 threshold (for
278 inclusion of SNPs in LD with the diabetes associated GWAS SNPs) = 0.80, LD window size = 1 Mb,
279 and minimum neighbor number = 500. P-values were adjusted according to Bonferroni threshold for
280 multiple testing burden.
281
fGWAS analysis. We used fGWAS [45] to model shared properties of loci affecting a trait. We ran

fGWAS (v0.3.6) with DIAMANTE T2D GWAS summary data and cell-type ATAC-seq peaks from three
cell types as input annotations. For each individual annotation, the output model provided maximum
likelihood enrichment parameters and annotations were considered as significantly enriched if the
parameter estimates and 95% confidence interval (CI) did not overlap zero. We then used fGWAS to
run a conditional analysis in a pair-wise manner where enrichment of one model was evaluated
conditional on the output models from other annotations.
289
2.6 Cicero co-accessibility analysis
In order to link TSS distal ATAC-seq peaks with target genes, we used Cicero [46], which identifies co-
accessible pairs of DNA elements using single-cell chromatin accessibility data. We used these results
to infer connections between regulatory elements and their target genes. We ran Cicero (v1.0.15,
default parameters) with cells from the alpha and beta cell clusters separately. To do this, we first called
peaks on each cluster and counted the number of reads per nuclei overlapping the peaks. The resulting
count matrix was used as input to Cicero along with the UMAP projection for each cluster. Finally, in
order to decide a threshold for filtering co-accessible peak pairs, we computed Fisher odds ratio for
enrichment of co-accessible peaks versus distance matched non co-accessible peaks (co-accessibility
< 0) with three different three-dimensional chromatin looping data sets: islet Hi-C [47], islet promoter
capture Hi-C (pcHi-C) [48], and EndoC Pol2 ChIA-PET anchors [49]. For overlap, we checked if both
the ends of the Cicero loops intersected with both the anchors from the experimental chromatin looping
data. Public epigenome browser session links have been included in **Table S7**.
303
T2D GWAS SNP overlap analysis. In order to link T2D GWAS SNPs with the target genes, we utilized
380 independent GWAS signals from DIAMANTE that were genetically fine-mapped to 99% credible
set. We filtered SNPs within each set to have >0.05 posterior probability of association (PPAg). We
then checked for each GWAS signal whether SNPs passing the criteria mapped within 1 kb of cell-type-
specific ATAC-seq peaks. To obtain Cicero target genes, we checked if an ATAC-seq peak was a)
within 1 kb of a variant, b) outside 1 kb range of a RefSeq TSS, and (c) linked to an ATAC-seq peaks
that was within 1 kb of a RefSeq TSS. The binary overlap matrix was clustered using hierarchical
clustering with binary distance method.
312
3. RESULTS
3.1. sci-ATAC-seq captures tissue relevant characteristics similar to bulk ATAC-seq
Pancreatic islets represent approximately 1-2% (by mass) of total pancreatic tissue [1] and therefore
requires specialized approaches to isolate in a manner that maintains viability. We obtained a highly
pure (>95% purity and >92% viability) sample of human pancreatic islet tissue from one individual
(cadaveric donor, female, 43 years old, and non-diabetic) and profiled chromatin-accessibility using sci-
ATAC-seq protocol [23] as described previously (**Fig. 1A, Table S2**). In total, we obtained 1,690 single-
cell ATAC-seq datasets with depth ranging from 17,667 to 415,237 (median: 79,482) reads per
nucleus, and TSS enrichment from 0.77 to 9.80 (median: 3.91) after removing background barcodes
(**Fig. S1A**). For quality assessment of each single nucleus, we reasoned that total reads and TSS
enrichment values are more suitable metrics for identifying nuclei with poor signal-to-noise ratio than
using fraction of reads in peaks as the latter may bias counts for under-represented cell-type
populations in the analysis (**Fig. S1B-C**). Based on these criteria, we obtained high-quality sci-ATAC-
seq data for 1,456 single-nuclei. In addition to sci-ATAC-seq data, we generated high-quality bulk
ATAC-seq data for ten islet samples with >47 M reads and >4.4 TSS enrichment per sample (**Table
S3**). Using our approach to identify high-confidence (master) peak calls across samples (see methods),
we obtained 106,460 bulk islet accessible chromatin peaks.

We then compared the aggregate islet sci-ATAC-seq data with bulk ATAC-seq samples from islets and
other tissues. For this, we called 156,311 peaks on the aggregate sci-ATAC-seq. We found that
aggregate sci-ATAC-seq profiles were concordant and clustered together with the other bulk islet
samples indicating that aggregate sci-ATAC-seq can capture chromatin accessibility in a manner
equivalent to bulk ATAC-seq assays (**Fig. 1B-C, S1D-E**). Further, to understand if the aggregate sci-
ATAC-seq peaks capture islet-specific regulatory features, we compared the distribution of peaks
across chromHMM chromatin state maps in eight tissues, including islets and the EndoC human beta
cell line [8]. We found that islet sci-ATAC-seq peaks overlap active TSS and active enhancer
segmentations in islet and EndoC (a beta cell line) chromatin state maps to a larger extent compared to
other tissues (**Fig. 1D**). Because chromHMM enhancer states are driven by H3K27ac marks and are
known to be associated with tissue-specific enhancer activity [5,50], our results indicate that sci-ATAC-
seq data capture the underlying islet-specific chromatin architecture similarly to bulk islet ATAC-seq
assays. Overall, these results indicate that our aggregate islet sci-ATAC-seq data are of high quality
and suggests that the underlying individual nuclei could reveal valuable cell-specific patterns of the
constituent cell-types.

3.2. sci-ATAC-seq reveals constituent cell-types in pancreatic islets
The aggregate sci-ATAC-seq profile of the islet is constituted of signal from distinct cell-types. For
identifying these cell-types, we leveraged the observation that TSS distal regions capture cell-type-
specific accessibility patterns and are effective at classifying constituent cell-types [51]. We adopted a
multi-step process to robustly detect and identify islet subpopulations (see methods, **Table S4**). This
approach produced four distinct clusters (**Fig. 2A**). In order to assign a cell-type identity to the clusters,
we merged nuclei in each cluster to create aggregate chromatin accessibility profiles and systematically
examined the patterns of accessibility at multiple cell-type marker loci. We found three clusters to have
distinct chromatin-accessibility patterns at *GCG, INS-IGF2,* and *SST* loci corresponding to three major
islet cell-types: alpha, beta, and delta cells (**Fig. 2B**). The fourth cluster (95 nuclei, ~7% of all nuclei)
showed a "mixed" cell-type appearance as shown by signal at multiple cell-specific markers. We
reasoned that these are likely to be nuclei doublets resulting from barcode collisions inherent to the
combinatorial indexing protocol, and thus should have skewed ATAC-seq read coverage. Indeed, we
observed that nuclei assigned to the mixed cell cluster were significantly (nominal P-value = 7.3e-7,
binomial test) enriched in the high sequencing depth bin relative to nuclei from other clusters (**Fig. 2C**).
As such, these nuclei were removed from further analyses yielding a total of 1,361 nuclei with 51%,
47%, and 2% assigned to beta, alpha, and delta cell-type respectively. These estimates agree with the
existing estimates of pancreatic islet cell-type proportions observed in confocal microscopy or single-
cell transcriptomics experiments [17,52–54]. As additional validation of our cell-type assignments, we
used cell-type signature genes from a published islet scRNA-seq study [17] and observed cluster-
specific chromatin accessibility consistent with our assigned cell identities (**Fig. 2D-E**).

We then analyzed the chromatin accessibility profile for each cell-type cluster. For this, we aggregated
nuclei within each cluster and identified peaks using MACS2. We identified 129,046 sites for alpha and
120,116 sites for beta cells. However, because the delta cluster had only 28 cells (corresponding to ~2
M reads), we reasoned that MACS2 would not perform ideally on data with such low depth. Indeed, we
only identified 49,293 peaks using MACS2 on the delta cell aggregate reads.

3.3. Deep learning enables robust peak calls on less abundant delta cells
To solve the challenge of learning cell-type-specific features from the sparse signal in low count delta
cell cluster, we used a deep learning approach based on U-Net architecture (**Fig. S2A**). U-Net was first
developed for biomedical image segmentation but has since been applied to many other problems

including audio and image super resolution. Its use in the super resolution problem served as the main
impetus for our choice of model to upscale genomic signals. We formulated our approach as a
classification problem where we use sparse signal and corresponding peak calls (equivalent to a low-
resolution image) to predict dense and high-quality peak calls (equivalent to a high-resolution image). In
order to avoid overfitting, we adopted a rigorous training scheme. We divided the chromosomes into
training, validation, and testing sets (**Fig. 3A**, **S2B**), and tested the performance of models within the
same cell-type and across different cell-types. We reasoned that our islet sci-ATAC-seq data is an ideal
fit for this problem as all the nuclei come from the same individual and processing batch and should,
therefore, contain no genetic or technical biases that would influence within or across cell-type
predictions. Since we had high-quality data from two cell-types, we trained two models: one model was
trained using 28-cell and 600-cell data from alpha cells (alpha trained model), while the second model
was trained similarly on data from beta cells (beta trained model). We then compared peak predictions
from both models to corresponding MACS2 peaks from the 600-cell data. We found that results from
cross cell-type predictions of both models outperformed MACS2 peak results as measured by mean
average precision **(Fig. 3B)**, suggesting that the U-Net was able to reconstruct peak calls from sparse
signal independent of the specific cell-type it was trained on. We highlight several examples where the
model was able to successfully predict peaks that were absent in sparse 28-cell data but present in
600-cell data of a cell-type **(Fig. 3C).** These predictions could not have transferred or "copied over"
from the training data because the training cell-type had no signal or peak call at the given locus. Based
on these results, we decided to use the U-Net models to predict peaks for the low-count delta cell
cluster. Since the U-net model provides a posterior probability score for each peak call prediction, we
sought to create a high-confidence set of predicted peak calls for each cell-type. We used a threshold
of 0.625 to filter predicted peaks for each cell-type. The choice of threshold was used to control for
potential false positives and final number of predicted cell-type peaks (**Fig S2C-D**).
399
Further, considering that delta peak predictions from both alpha and beta model were highly concordant
(Jaccard index of 0.85), we used the intersection of the results as the final predicted outcome. Now with
representative peak calls from each cluster, we compared them to bulk islet ATAC-seq master peak
calls (**Table S5**). We found that master peaks derived from bulk islets were highly reproducible across
samples, with >70% of peaks occurring in five or more of the ten samples (**Fig. 3D**). Predictably, we
also observed that the chromatin states corresponding to "Active TSS" and "Active Enhancer" showed
enrichment with increasing reproducibility of master peaks. Likewise, chromatin states such as
"Repressed polycomb", "Weak transcription", "Quiescent/low signal" showed a depletion with increasing
islet ATAC-seq peak reproducibility (**Fig. 3E**). Similarly, when we compared cell-type peaks to the
master peaks, we found that the proportion of peaks from each cell-type increased with the increasing
reproducibility of bulk peaks, suggesting that highly reproducible peaks are driven by all constituent
cell-types while peaks that occur in fewer samples might originate from underlying cell-population
variability (**Fig. 3F).**
413
While the primary model of our interest was trained using data from 28 cells to predict 600-cell
equivalent peaks, we asked if the model would perform similarly for a varying resolution of input data.
For this, we subsampled cells from alpha and beta cell clusters to sets of different cell counts, starting
with as few as five cells to 500 cells. We found that the performance of the model increased with the
increasing number of cells used in the input training data (**Fig. S2E**). There was up to five-fold gain in
coverage of T2D GWAS SNPs in beta predicted peaks compared to MACS2 peaks (**Fig. S2F**) even
when fewer cells were used as input training data (**Fig. S2G)**. These results suggest that the deep-

learning strategy is applicable to a range of input data typically seen in single-cell sequencing
experiments.
423
424 Overall, our results show that deep-learning driven feature prediction can help us recover tissue and
425 cell-type relevant chromatin accessibility patterns from sparse and noisy data. Using this approach can
426 enhance biological discoveries, which is challenging with rare cell populations.
427
3.4. T2D GWAS enrichment at cell-type-specific chromatin signatures
We computed the overlap enrichment of T2D GWAS loci in cell-type peak annotations from alpha, beta,
and delta cells using a Bayesian hierarchical model, as implemented in fGWAS [45]. fGWAS allows
calculation of marginal enrichment associations for one cell type conditioned on another by using not
only the subset of genome-wide significant loci but also the full genome-wide association summary
statistics. We observed that annotations from all three cell types were highly enriched for T2D GWAS
loci, with beta-cell annotations having the highest enrichment values. (**Fig. 4A**). However, when we
accounted for marginal associations using a joint model, we found that beta cells are the only cell type
to remain enriched after adjusting for the other two cell types. This result suggests that shared or beta
cell-specific chromatin accessibility peaks drive the association with T2D GWAS. More broadly, these
findings illustrate how single cell chromatin profiling results, when coupled with conditional statistical
enrichment analyses, can dissect specific cell types that drive enrichment in bulk tissue samples.
440
We next partitioned the peaks into exclusive sets based on the cell-types shared by each peak.
Because the delta cell cluster has fewer reads compared to alpha and beta cells, we did not utilize read
count-based approaches to determine cell-type-specific peaks. Instead, we used peak level metrics to
identify peaks exclusive to a combination of cell types. We found that a majority of peaks (47,209) were
shared across all cell-types and that each cell type had a set of unique accessible sites (29,884 beta,
39,353 alpha, 31,330 delta) (**Fig. 4B**). Consistent with our expectations, TSS-proximal shared peaks
mostly overlapped active TSS chromatin states compared to cell-type-specific peaks which had a larger
proportion of peaks in active enhancer states (**Fig. S3A**). We then used a complementary enrichment
approach with the GREGOR tool [44] to determine if T2D GWAS loci are enriched in each subclass of
peaks. We found that T2D GWAS loci were highly enriched in shared peaks (P-value 1.64e-16, fold
enrichment 2.03) and beta cell-specific peaks (P-value 6.42e-6, fold enrichment 1.91) (**Fig. S3B**). We
also observed moderate enrichment of T2D GWAS SNPs in other sets of cell-type-specific peaks, but
strikingly, there was little enrichment in delta cell-specific peaks (P-value 3.12e-3, fold enrichment 1.55)
and no significant enrichment in alpha cell-specific peaks (P-value 1.83e-01, fold enrichment 1.16). This
suggests that the role of alpha and delta cells in the mechanisms underlying genetic predisposition to
T2D pathophysiology might be limited compared to beta cells. Further, the enrichment of shared peaks
also suggests that these variants might be affecting regulatory pathways that are shared across all cell
types in the islets, suggesting that a narrow focus on studying beta cells only for T2D might not yield a
complete picture of the disease mechanisms. These independent enrichment findings from the
GREGOR tool are consistent with the results from the fGWAS analysis (**Fig. 4A**), indicating the robust
nature of these results.
462
3.5. Linking cell-type-specific chromatin accessibility to target genes
One of the primary challenges in understanding the underlying biological mechanisms at non-coding
T2D GWAS variants is the identification of their target genes. Risk variants occurring in enhancer
regions can often interact with their target genes that are not adjacent. Multiple studies have examined
the regulatory landscape of pancreatic islets and relevant cell lines using chromosome conformation

capture techniques to nominate target genes [47–49]. However, most of these studies were conducted
on bulk islet samples, thereby obscuring any cell-specific signatures of chromatin looping. Additionally,
chromatin looping studies tend to have noisy signals when two regions are close in linear space, which
leads to a bias towards detecting longer-range interactions. In order to mitigate these limitations, we
adopted a recently published approach, Cicero [46], which leverages profiles of chromatin co-
accessibility across single cells to infer pairs of chromatin peaks that are likely to be in close physical
proximity. For this analysis, we focused on alpha and beta cell-types as they were the clusters with the
most nuclei. In order to filter the Cicero co-accessible scores for those peak pairs that are more likely to
represent true looping, we compared our results to experimentally-defined loops from three
independent chromatin looping data sets: islet Hi-C [47], islet promoter capture Hi-C (pcHi-C) [48], and
EndoC Pol2 ChIA-PET [49] loops. We found that Cicero peak-pairs from our sci-ATAC-seq data with
score >0.05 were strongly enriched to be called as loops in each of the three reference data sets (**Fig.**
4C, S4A). With this threshold, we found 190,176 beta cell and 147,716 alpha cell co-accessible peak-
pairs.
482
Using our new catalog of Cicero-inferred chromatin loops, we next sought to link TSS-distal T2D GWAS
variants to target gene promoters. We focused on the latest T2D GWAS results and used SNPs in
association signals that were genetically fine-mapped to be in a 99% credible set and had >0.05
posterior probability of association (PPAg) [3]. For this mapping procedure, we required that the
credible set SNP was not within 1 kb of an annotated TSS and that the other end of the chromatin loop
occurs within 1 kb of an annotated TSS. Using this approach across both alpha and beta cells, we
found that of the 265 independent GWAS signals containing SNPs that met our criteria, we were able to
nominate target genes at 104 of them **(Fig. 4D)**. In a similar manner, we checked if the SNPs within
each locus overlapped a cell-type-specific peak (**Table S6**). We observed several notable examples. At
the *C2CD4A/B* locus, we found rs7163757 (PPAg 0.095) to be linked to *C2CD4B* in alpha cells (**Fig.**
4E). Using an islet gene expression and genetic integration approach to identify expression quantitative
trait loci (eQTL), we previously showed that rs7163757 is associated with *C2CD4B* expression [8], and
a subsequent functional study corroborated these findings [55]. At a different locus, we found
rs11708067 (PPAg 0.79), located in an islet enhancer within the *ADCY5* gene, to be linked to the TSS
of the corresponding gene **(Fig. 4F).** The risk allele of rs11708067 has been reported to be associated
with reduced expression of *ADCY5* [56] and functional validation experiments show association with
impaired insulin secretion [4,9]. As an example of a beta cell-specific connection, we found variant
rs13262861 (PPAg 0.97) within the *ANK1* locus to be linked to nearby *NKX6-3* (**Fig. 4G).** We have
previously used islet eQTL data to nominate *NKX6-3* as an islet target gene at this locus [4,8]. The
extensive support from previous publications for these three loci serves as positive controls for our
results and reinforces the quality of this sci-ATAC-seq data and analyses. Finally, we highlight
rs62059712 (PPAg 0.34) within the *ATP1B2* locus as an example of a variant linked to multiple gene
promoters across both beta and alpha cell-types **(Fig. 4H)**. Notably, of the 104 T2D GWAS signals for
which we were able to nominate target genes in either cell type, 60 (~58%) had more than one
nominated target gene.
508
4. DISCUSSION
Single-nuclei chromatin accessibility profiling provides a unique approach for mapping of cell-type-
specific regulatory signatures. Here, we utilized the sci-ATAC-seq protocol to generate and study
chromatin accessibility profiles for 1,456 high-quality nuclei from a purified pancreatic islet sample. Our
dataset and analyses provide high-quality maps of cell-type accessibility profiles and regulatory
architecture using an unbiased approach compared to prior maps from sorted cell-type populations.

However, it is essential to emphasize that single-cell data present unique challenges, and that our
study, which analyzed only one pancreatic islet sample, may be limited in how it can address some of
them.
518
First, de-novo identification of cell types from the sparse single-cell chromatin accessibility data
continues to be a challenge. We adopted several strategies to address potential biases in our analysis.
Our logistic regression approach to eliminate read depth as a confounding technical variable, combined
with the binomial counting strategy to infer doublet enrichment in clusters, enabled us to identify three
major cell-type populations corresponding to alpha, beta, and delta cells. In order to assign these cell
identities, we relied not only on classical hormone markers, but we also leveraged findings from an
independent islet single-cell RNA-seq study to validate our results. While islets have been reported to
contain other rarer cell-type populations (<5% of all islet cells) [52], our ability to observe them was
limited due to the size of our dataset.
528
Second, we faced the challenge of identifying reliable cell-type-specific accessibility patterns across cell
types due to the relatively low abundance of delta cells. The lower numbers of reads reduce the
statistical power to call peaks and use count-based methods to identify distinct categories of accessible
regions. As such, our U-Net-based deep learning approach presents a novel strategy for addressing
this particular problem. We demonstrated, using alpha and beta cells as reciprocal training and testing
datasets, that our model successfully learns to predict high-quality peak calls from low cell count data.
We observed, however, that there are diminishing returns from using deep learning models when 200
or more cells are used as input to the model, an observation consistent with the threshold of
experimental reproducibility highlighted in a recent large-scale single nuclei ATAC-seq study [15]. This
consistency with an independent study reinforces the value of our deep learning approach but also
highlights a limitation of our delta peak predictions which derive from a low cell count input dataset.
Nonetheless, we envision that our method will be useful in scenarios where it is challenging or cost-
prohibitive to obtain specific cell populations.
542
Overall, an important implication of our findings comes from our ability to generate cell-specific
chromatin accessibility maps and to infer looping connections from accessible regions to target genes
of T2D GWAS variants. A recent T2D GWAS [3] reported >400 independent association signals, but
the molecular mechanisms underlying these signals is known only for a subset of the variants. Single
nuclei resolution cell-specific regulatory signatures provide a unique opportunity to infer target gene
links with non-coding elements. Thus, we integrated cell-type co-accessibility links with T2D GWAS
SNPs that were genetically fine-mapped to 99% credible sets to create a higher resolution map of the
regulatory landscape underlying 104 distinct T2D GWAS signals. Focusing on the cell-specificity of the
chromatin accessibility peaks that anchor these target gene associations, we observed seven classes,
representing: i. peaks that are unique to a cell type (three classes), ii. peaks that are shared across all
three cell types (one class), iii. peaks that occur in a pair of cell types (three classes). Interestingly, the
class of peaks shared across all three cell types comprised 26 of the 104 (25%) T2D GWAS to target
gene links even though this class is only one of seven. These results paint a complex picture of disease
mechanisms where certain risk variants may mediate target effects through cell-type-specific pathways,
while others might affect multiple target genes shared across cell-type populations.
558
559 We noted specific examples at the *C2CD4A/B* and *ANK1* loci, where we were able to nominate specific
560 variants linked with islet gene expression and their role in T2D pathophysiology as compelling targets
561 for future mechanistic studies. As this manuscript was under preparation, another similar study

appeared as a preprint [57], and as such an important future topic will be to combine and meta-analyze multiple islet single-cell ATAC-seq datasets. Such an endeavor will increase statistical power to detect chromatin features, including loops, at GWAS loci, and eventually enable single-cell resolution chromatin QTL studies, which will help to further narrow in on functional SNPs. Overall, we believe that the data, results, and methodology from this study will be of value to the broader research community.

DATA AVAILABILITY

The data reported in this paper have been deposited in the dbGaP (accession no. phs001188.v1.p1; FUSION Tissue Biopsy Study—Islet Expression and Regulation by RNAseq and ATACseq). Code for deep learning analysis is available on GitHub at https://github.com/ParkerLab/PillowNet. Custom scripts are available at https://github.com/ParkerLab/islet_sci-ATAC-seq_2019. Additional URLs are provided in **Table S7** of Supplementary Data file.

ACKNOWLEDGEMENTS

We thank the members of the Parker, Collins, and Shendure labs for helpful suggestions and feedback. This work was supported by American Diabetes Association (ADA) Fellowship to JPD, National Institute of Diabetes and Digestive and Kidney Diseases grant R01 DK117960, National Heart, Lung, and Blood Institute grant U01 HL137182, and the ADA Pathway to Stop Diabetes Grant 1-14-INI-07 to SCJP, and National Institutes of Health Grants 1-ZIA-HG000024 to FSC. The medical art images were obtained from Servier Medical Art by Servier under CC-BY-3.0 license. We thank F. Steemers and L. Christiansen at Illumina for providing indexed Tn5 transposase. We also thank NVIDIA Corporation for donating the Titan Xp GPUs to SCJP.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

FSC, JS, and SCJP conceived the study; MRE, DC, and RMD generated the data; LSZ, JPD, and NN performed initial analysis; VR, DXQ, and SCJP analyzed the data and performed the research presented here; DXQ and YG implemented the U-Net strategy and wrote the code; FSC, JS, and SCJP jointly supervised the work; and VR, DXQ, and SCJP wrote the manuscript with feedback from all the authors.

REFERENCES

[1]   DeFronzo RA, Ferrannini E, Groop L, Henry RR, Herman WH, Holst JJ, et al. Type 2 diabetes mellitus. Nat Rev Dis Primer 2015;1:15019. doi:10.1038/nrdp.2015.19.

[2]   Spellman CW. Pathophysiology of Type 2 Diabetes: Targeting Islet Cell Dysfunction. J Am Osteopath Assoc 2010;110:S2–7.

[3]   Mahajan A, Taliun D, Thurner M, Robertson NR, Torres JM, Rayner NW, et al. Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. Nat Genet 2018:1. doi:10.1038/s41588-018-0241-6.

[4]   Viñuela A, Varshney A, Bunt M van de, Prasad RB, Asplund OB, Bennett A, et al. Influence of genetic variants on gene expression in human pancreatic islets – implications for type 2 diabetes. BioRxiv 2019:655670. doi:10.1101/655670.

[5]   Parker SCJ, Stitzel ML, Taylor DL, Orozco JM, Erdos MR, Akiyama JA, et al. Chromatin stretch enhancer states drive cell-specific gene regulation and harbor human disease risk variants. Proc Natl Acad Sci 2013;110:17921–6. doi:10.1073/pnas.1317023110.

[6]   Pasquali L, Gaulton KJ, Rodríguez-Seguí SA, Mularoni L, Miguel-Escalada I, Akerman ĺ, et al.

Pancreatic islet enhancer clusters enriched in type 2 diabetes risk-associated variants. Nat Genet
2014;46:136–43. doi:10.1038/ng.2870.

[7]   Thurner M, van de Bunt M, Torres JM, Mahajan A, Nylander V, Bennett AJ, et al. Integration of
human pancreatic islet genomic data refines regulatory mechanisms at Type 2 Diabetes
susceptibility loci. ELife 2018;7:e31977. doi:10.7554/eLife.31977.

[8]   Varshney A, Scott LJ, Welch RP, Erdos MR, Chines PS, Narisu N, et al. Genetic regulatory
signatures underlying islet gene expression and type 2 diabetes. Proc Natl Acad Sci
2017;114:2301–6. doi:10.1073/pnas.1621192114.

[9]   Bunt M van de, Fox JEM, Dai X, Barrett A, Grey C, Li L, et al. Transcript Expression Data from
Human Islets Links Regulatory Signals from Genome-Wide Association Studies for Type 2
Diabetes and Glycemic Traits to Their Downstream Effectors. PLOS Genet 2015;11:e1005694.
doi:10.1371/journal.pgen.1005694.

[10] Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin
for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and
nucleosome position. Nat Methods 2013;10:1213–8. doi:10.1038/nmeth.2688.

[11] Hesselberth JR, Chen X, Zhang Z, Sabo PJ, Sandstrom R, Reynolds AP, et al. Global mapping of
protein-DNA interactions *in vivo* by digital genomic footprinting. Nat Methods 2009;6:283–9.
doi:10.1038/nmeth.1313.

[12] Shema E, Bernstein BE, Buenrostro JD. Single-cell and single-molecule epigenomics to uncover
genome regulation at unprecedented resolution. Nat Genet 2018:1. doi:10.1038/s41588-018-
0290-x.

[13] Cusanovich DA, Reddington JP, Garfield DA, Daza RM, Aghamirzaie D, Marco-Ferreres R, et al.
The *cis*-regulatory dynamics of embryonic development at single-cell resolution. Nature
2018;555:538–42. doi:10.1038/nature25981.

[14] Preissl S, Fang R, Huang H, Zhao Y, Raviram R, Gorkin DU, et al. Single-nucleus analysis of
accessible chromatin in developing mouse forebrain reveals cell-type-specific transcriptional
regulation. Nat Neurosci 2018;21:432–9. doi:10.1038/s41593-018-0079-3.

[15] Satpathy AT, Granja JM, Yost KE, Qi Y, Meschi F, McDermott GP, et al. Massively parallel single-
cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion.
Nat Biotechnol 2019;37:925–36. doi:10.1038/s41587-019-0206-z.

[16] Cusanovich DA, Hill AJ, Aghamirzaie D, Daza RM, Pliner HA, Berletch JB, et al. A Single-Cell
Atlas of In Vivo Mammalian Chromatin Accessibility. Cell 2018. doi:10.1016/j.cell.2018.06.052.

[17] Lawlor N, George J, Bolisetty M, Kursawe R, Sun L, Sivakamasundari V, et al. Single-cell
transcriptomes identify human islet cell signatures and reveal cell-type–specific expression
changes in type 2 diabetes. Genome Res 2017;27:208–22. doi:10.1101/gr.212720.116.

[18] Muraro MJ, Dharmadhikari G, Grün D, Groen N, Dielen T, Jansen E, et al. A Single-Cell
Transcriptome Atlas of the Human Pancreas. Cell Syst 2016;3:385-394.e3.
doi:10.1016/j.cels.2016.09.002.

[19] Arda HE, Tsai J, Rosli YR, Giresi P, Bottino R, Greenleaf WJ, et al. A Chromatin Basis for Cell
Lineage and Disease Risk in the Human Pancreas. Cell Syst 2018;7:310-322.e4.
doi:10.1016/j.cels.2018.07.007.

[20] Ackermann AM, Wang Z, Schug J, Naji A, Kaestner KH. Integration of ATAC-seq and RNA-seq
identifies human alpha cell and beta cell signature genes. Mol Metab 2016;5:233–44.
doi:10.1016/j.molmet.2016.01.002.

[21] Nica AC, Ongen H, Irminger J-C, Bosco D, Berney T, Antonarakis SE, et al. Cell-type, allelic, and
genetic signatures in the human pancreatic beta cell transcriptome. Genome Res 2013;23:1554–
62. doi:10.1101/gr.150706.112.

[22] Dorrell C, Grompe MT, Pan FC, Zhong Y, Canaday PS, Shultz LD, et al. Isolation of mouse
pancreatic alpha, beta, duct and acinar populations with cell surface markers. Mol Cell Endocrinol
2011;339:144–50. doi:10.1016/j.mce.2011.04.008.

[23] Cusanovich DA, Daza R, Adey A, Pliner HA, Christiansen L, Gunderson KL, et al. Multiplex single-
cell profiling of chromatin accessibility by combinatorial cellular indexing. Science 2015;348:910–
4. doi:10.1126/science.aab1601.

[24] John Hensley. cta: C++ implementation of Buenrostro adapter trimming. The Parker Lab at the

University of Michigan; 2017.

[25] Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
ArXiv13033997 Q-Bio 2013.

[26] Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics 2009;25:2078–9. doi:10.1093/bioinformatics/btp352.

[27] Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based Analysis
of ChIP-Seq (MACS). Genome Biol 2008;9:R137. doi:10.1186/gb-2008-9-9-r137.

[28] The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human
genome. Nature 2012;489:57–74. doi:10.1038/nature11247.

[29] A toolkit for QC and visualization of ATAC-seq results.: ParkerLab/ataqv. The Parker Lab at the
University of Michigan; 2018.

[30] Heger A, Webber C, Goodson M, Ponting CP, Lunter G. GAT: a simulation framework for testing
the association of genomic intervals. Bioinformatics 2013;29:2046–8.
doi:10.1093/bioinformatics/btt343.

[31] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 2014;30:2114–20. doi:10.1093/bioinformatics/btu170.

[32] Fraley C, Raftery AE, Scrucca L, Murphy TB, Fop M. mclust: Gaussian Mixture Modelling for
Model-Based Clustering, Classification, and Density Estimation. 2019.

[33] O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference sequence
(RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation.
Nucleic Acids Res 2016;44:D733-745. doi:10.1093/nar/gkv1189.

[34] Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by
seed-and-vote. Nucleic Acids Res 2013;41:e108. doi:10.1093/nar/gkt214.

[35] Gierahn TM, Wadsworth Ii MH, Hughes TK, Bryson BD, Butler A, Satija R, et al. Seq-Well:
portable, low-cost RNA sequencing of single cells at high throughput. Nat Methods 2017;14:395–
8. doi:10.1038/nmeth.4179.

[36] UMAP: Uniform Manifold Approximation and Projection. J Open Source Softw 2018.
doi:10.21105/joss.00861.

[37] Hahsler M, Piekenbrock M, Arya S, Mount D. dbscan: Density Based Clustering of Applications
with Noise (DBSCAN) and Related Algorithms. 2019.

[38] Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, et al. GENCODE:
The reference human genome annotation for The ENCODE Project. Genome Res 2012;22:1760–
74. doi:10.1101/gr.135350.111.

[39] Ronneberger O, Fischer P, Brox T. U-Net: Convolutional Networks for Biomedical Image
Segmentation. ArXiv150504597 Cs 2015.

[40] François Chollet. Keras. GitHub; 2015.

[41] Abadi M, Barham P, Chen J, Chen Z, Davis A, Dean J, et al. TensorFlow: A system for large-scale
machine learning. 12th USENIX Symp. Oper. Syst. Des. Implement. OSDI 16, 2016, p. 265–283.

[42] Kingma DP, Ba J. Adam: A Method for Stochastic Optimization. ArXiv14126980 Cs 2014.

[43] Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics 2010;26:841–2. doi:10.1093/bioinformatics/btq033.

[44] Schmidt EM, Zhang J, Zhou W, Chen J, Mohlke KL, Chen YE, et al. GREGOR: evaluating global
enrichment of trait-associated variants in epigenomic features using a systematic, data-driven
approach. Bioinformatics 2015;31:2601–6. doi:10.1093/bioinformatics/btv201.

[45] Pickrell JK. Joint Analysis of Functional Genomic Data and Genome-wide Association Studies of
18 Human Traits. Am J Hum Genet 2014;94:559–73. doi:10.1016/j.ajhg.2014.03.004.

[46] Pliner HA, Packer JS, McFaline-Figueroa JL, Cusanovich DA, Daza RM, Aghamirzaie D, et al.
Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin Accessibility Data.
Mol Cell 2018. doi:10.1016/j.molcel.2018.06.044.

[47] Greenwald WW, Li H, Benaglio P, Jakubosky D, Matsui H, Schmitt A, et al. Subtle changes in
chromatin loop contact propensity are associated with differential gene regulation and expression.
Nat Commun 2019;10:1054. doi:10.1038/s41467-019-08940-5.

[48] Miguel-Escalada I, Bonàs-Guarch S, Cebola I, Ponsa-Cobas J, Mendieta-Esteban J, Atla G, et al.
Human pancreatic islet three-dimensional chromatin architecture provides insights into the

genetics of type 2 diabetes. Nat Genet 2019;51:1137. doi:10.1038/s41588-019-0457-0.

[49] Lawlor N, Márquez EJ, Orchard P, Narisu N, Shamim MS, Thibodeau A, et al. Multiomic Profiling
Identifies cis-Regulatory Networks Underlying Human Pancreatic β Cell Identity and Function. Cell
Rep 2019;26:788-801.e6. doi:10.1016/j.celrep.2018.12.083.

[50] Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, et al. Super-Enhancers in the
Control of Cell Identity and Disease. Cell 2013;155:934–47. doi:10.1016/j.cell.2013.09.053.

[51] Bulger M, Groudine M. Functional and Mechanistic Diversity of Distal Transcription Enhancers.
Cell 2011;144:327–39. doi:10.1016/j.cell.2011.01.024.

[52] Cabrera O, Berman DM, Kenyon NS, Ricordi C, Berggren P-O, Caicedo A. The unique
cytoarchitecture of human pancreatic islets has implications for islet cell function. Proc Natl Acad
Sci 2006;103:2334–9. doi:10.1073/pnas.0510790103.

[53] Brissova M, Fowler MJ, Nicholson WE, Chu A, Hirshberg B, Harlan DM, et al. Assessment of
Human Pancreatic Islet Architecture and Composition by Laser Scanning Confocal Microscopy. J
Histochem Cytochem 2005;53:1087–97. doi:10.1369/jhc.5C6684.2005.

[54] Segerstolpe Å, Palasantza A, Eliasson P, Andersson E-M, Andréasson A-C, Sun X, et al. Single-
Cell Transcriptome Profiling of Human Pancreatic Islets in Health and Type 2 Diabetes. Cell
Metab 2016;24:593–607. doi:10/f3vcbz.

[55] Kycia I, Wolford BN, Huyghe JR, Fuchsberger C, Vadlamudi S, Kursawe R, et al. A Common Type
2 Diabetes Risk Variant Potentiates Activity of an Evolutionarily Conserved Islet Stretch Enhancer
and Increases C2CD4A and C2CD4B Expression. Am J Hum Genet 2018;102:620–35.
doi:10.1016/j.ajhg.2018.02.020.

[56] Roman TS, Cannon ME, Vadlamudi S, Buchkovich ML, Wolford BN, Welch RP, et al. A Type 2
Diabetes-Associated Functional Regulatory Variant in a Pancreatic Islet Enhancer at the ADCY5
Locus. Diabetes 2017;66:2521–30. doi:10.2337/db17-0464.

[57] Chiou J, Zeng C, Cheng Z, Han JY, Schlichting M, Huang S, et al. Single cell chromatin
accessibility reveals pancreatic islet cell type- and state-specific regulatory programs of diabetes
risk. BioRxiv 2019:693671. doi:10.1101/693671.

FIGURE LEGENDS

Figure 1. Schematic of sci-ATAC-seq study. (A) sci-ATAC-seq protocol for generating single-nuclei ATAC-seq data from a pancreatic islet sample. The data is then used to identify constituent cell-types and use deep-learning model to predict peaks on the clusters with fewer nuclei count. (B) ATAC-seq signal tracks for 10 bulk islet samples and sci-ATAC-seq islet sample. Bottom tracks show the signal across a random subset of upto 400 single-nuclei. Signal tracks are normalized to one million reads and scaled between 0-2. (C) Spearman correlation between aggregate sci-ATAC-seq, 13 bulk islets, 3 adipose, 2 muscle, 2 CD4+ T-cells, and 1 GM12878 sample (see **Table S1**). (D) Distribution of aggregate sci-ATAC-seq TSS proximal and distal peaks across bulk islet derived ChromHMM segmentations.

Figure 2. Clustering and identification of cell-type clusters in sci-ATAC-seq data. (A) UMAP projection with clustering of 1,456 single-nuclei islets represented by each single point into four clusters as identified by density-based clustering. (B) Enrichment of cells from each cluster relative to their expected population proportion across different read sequencing depth bins. Sequencing depth increases with the bin number. (C) Genome browser tracks showing signal at different cell-type marker loci: alpha (*GCG*), beta (*INS-IGF2*), delta (*SST*), and a housekeeping gene *(GAPDH)*. Tracks are normalized to one million reads and scaled between 0-5. (D) Overview of independent cluster verification scheme utilizing cell-type signature genes as identified by an islet scRNA-seq study by Lawlor et al (2017). (E) Plot of aggregate ATAC-seq signal (normalized using RPKM) at scRNA-seq derived cell-type signature genes for alpha, beta, and delta cells. Number of signature genes for each cell type indicated in the legend.

Figure 3. Deep learning upscaling from sparse low-count nuclei clusters. (A) Schematic of U-Net training scheme. Two models are depicted in the illustration: one trained on alpha cells data as input and other trained on beta cells as input. Delta cell peak predictions from both models are combined to get final predictions (see Methods). (B) Precision-recall curve comparing peak calls from MACS2 on downscaled data (alpha cell-type) with predicted peak calls from 28 cell U-net model (trained on beta, predicted on alpha). (C) Example loci illustrating peak upscaling with the model. For each cell-type, four tracks are shown: full signal track, peak calls on full data, peak calls on subsampled data, and predicted peak calls. The predicted peak calls are obtained from a model trained on a different cell-type. For delta predicted peak calls, intersection of prediction from both alpha and beta models are shown. Signal tracks normalized to one million reads and scaled between 0-2. (D) Reproducibility of master peaks from bulk islet ATAC-seq across individual samples. (E) Fold enrichment ($\log_2$) of different sets of reproducible peaks from bulk islet ATAC-seq across 13 islet chromatin states. Genic Enhancer is not shown because of no enrichment. (F) Overlap of cell-type peaks (alpha, beta, predicted delta) with different sets of reproducible peaks from bulk islet ATAC-seq data.

Figure 4. Enrichment of T2D GWAS signals in cell-type-specific chromatin and linking them to target genes. (A) Fold enrichment ($\log_2$) of T2D GWAS SNPs in cell-type peaks in single and conditional analysis mode using fGWAS tool. For each cell-type, three enrichment values with 95% confidence intervals are shown: None (single annotation mode), Alpha *(*conditioned on alpha), Beta (conditioned on beta), and Delta (conditioned on delta). (B) Partitioning of alpha, beta, and predicted delta peaks in mutually exclusive sets of cell-type-specific peaks. The subplot (on right) shows the total number of peaks for each cell-type. (C) Distance-matched Fisher odds that beta cell co-accessibility links overlap islet Hi-C, islet pcHi-C, and ChIA-PET chromatin loops across different co-accessibility

threshold bins. (D) Overlap of T2D GWAS credible set SNPs with cell-type-specific peaks. Bin is
colored if there's at least one SNP (PPAg > 0.05) in the 99% genetic credible set of the T2D GWAS
signal located within 1 kb of an ATAC-seq peak. Cicero score columns are colored to indicate the score
of the highest scoring link to the target gene. (E) Viewpoint plot of alpha Cicero connections centered at
rs7163757 for *C2CD4A/B* locus, (F) alpha Cicero connections centered at rs11708067 for *ADCY5*
locus, (G) beta Cicero connections centered at rs13262861 for *ANK1* locus, and (H) Cicero connections
for both alpha and beta centered at rs62059712 for *ATP1B2* locus. The viewpoint region is +/- 1kb of
the region from the variant.
801
802

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. ATAC-seq metrics of nuclei from sci-ATAC-seq. (A) Distribution of reads per barcodes shown with the threshold chosen for filtering background barcodes. (B) Fraction of reads in peaks versus TSS Enrichment, and (C) Total autosomal reads versus TSS enrichment for all single nuclei. Density units are arbitrary. (D) TSS coverage of aggregate sci-ATAC-seq, and (E) Fragment length distribution of aggregate sci-ATAC-seq compared with ten bulk islet ATAC-seq samples.

Figure S2. Peak calling using deep learning approach. (A) Schematic of U-Net learning strategy. (B) The training, testing, and validation scheme used for training the models delineating which chromosomes were part of what dataset. (C) Number of predicted peaks (from 28-cell trained model) for each cell type with different output posterior probability thresholds. (D) Number of cell-type specific peaks for alpha, beta, and delta after partitioning into mutually exclusive sets (see methods) with different output posterior probability thresholds. (E) Average precision in predicting peaks compared for all four models (two training and two prediction datasets) with different sizes of input training data. (F) Enrichment of T2D GWAS SNPs (N=378) in predicted beta peak calls (from alpha-trained model) compared with peaks calls from MACS2 on the data with varying size of input training data. (G) Precision and recall curves comparing predicted beta peaks (from alpha-trained model) for varying size of input training data.

Figure S3. (A) Distribution of TSS proximal and distal peaks (>5kb from nearest Refseq TSS) in shared peaks and peaks assigned only to alpha, beta, and delta cell-type. (B) Enrichment of T2D GWAS SNPs (N=378) across all cell-type-specific sets of peaks.

Figure S4. (A) Fisher odds score for enrichment of alpha co-accessible sites in loop anchors from three different dataset: Islet Hi-C, Islet pcHi-C, and EndoC ChIA-PET.