Title: The current obesity "epidemic": segregation of familial genetic risk in NHANES

cohort supports a major role for large genetic effects

Running title: Segregation of familial genetic obesity risk

Authors: Arthur B. Jenkins[1,4]*, Marijka Batterham[2], Lesley V. Campbell[3,4]

[1] School of Medicine, University of Wollongong, Wollongong NSW 2522 Australia

[2] Statistical Consulting Centre, School of Mathematics and Applied Statistics, University of

Wollongong NSW 2522 Australia

[3] Diabetes Service, St Vincent's Hospital, Sydney, NSW 2010 Australia

[4] Diabetes and Metabolism Division, Garvan Institute of Medical Research, Darlinghurst, NSW 2010

Australia

* Corresponding author

E-mail: ajenkins@uow.edu.au

The authors declare no competing financial or non-financial interests.

Abstract

Background/Objectives: The continuing increase in many countries in adult body mass index (BMI

kg/m^2) and its dispersion is contributed to by interactions between genetic susceptibilities and an

increasingly obesogenic environment. Whether population susceptibility to obesogenic environments

is mainly determined by a subgroup with high genetic susceptibility or susceptibility is more evenly

distributed throughout the population is unresolved, due to uncertainties around relevant genetic and

environmental architecture. We aimed to test the predictions of a Mendelian genetic architecture

based on collectively common but individually rare large-effect variants and its ability to account for

current trends in a large population-based sample.

Subjects/Methods: We studied publicly available adult BMI data (n = 9102) from 3 cycles of

NHANES (1999, 2005, 2013) adjusted for age, gender and race/ethnicity. A first degree family

history of diabetes (FH) served as a binary marker (FH_0/FH_1) of genetic obesity susceptibility. We

tested for multi-modality in BMI non-parametrically using a runs tests in conditional quantile

regression (CQR) models of FH effects, obtained parametric fits to a Mendelian model in FH_1, and

estimated FH-environment interactions in CQR models and in ANCOVA models incorporating

secular time.

Results: A unimodal FH effect on BMI was excluded ($p \leq 0.0001$) in CQR models and parametric fits

to a Mendelian model in FH_1 identified 3 modes at 25.8 ± 1.0 (SEM), 32.1 ± 1.8 and 40.6 ± 2.5 kg/m^2.

Mode separation accounted for ~40% of BMI variance in FH_1 providing a lower bound for the

contribution of large effects. CQR analysis identified strong interactions between FH and

environmental factors ($p \leq 0.01$) and FH_1 accounted for ~60% of the secular trends in BMI and its SD

in ANCOVA models.

Conclusions: Multimodality in the FH effect is inconsistent with a predominantly polygenic small

effect architecture. We conclude that large genetic effects interacting with obesogenic environment

provide a better, quantitative explanation for current trends in BMI.

Introduction

The recent and continuing increase in the global mean adult BMI, first seen in high income countries, is now apparent in most countries across a wide range of ethnic composition and socio-economic conditions (1) and is accompanied by increases in measures of dispersion (2, 3). Although BMI is known from family-based studies to be under strong genetic influences (4) population genetic backgrounds have been effectively constant over this time, implying that BMI trends are driven by change in environmental factors (obesogenic environment, OE). Evidence from twin studies, which demonstrate increased genetic variance over time, supports an important role for interactions between OE and genetic susceptibility (G x OE) on both mean and dispersion of BMI (3, 5), but how large a role is not yet clear. Defining the role of G x OE in "epidemic" obesity, and hence of genetic susceptibility itself, is hindered by problems of measurement and modeling of interactions (6) and by uncertainty around both the genetic architecture (4) and the exact nature of the environmental drivers (7). Whether a population's susceptibility to OE is predominantly determined by a subgroup with high genetic susceptibility or is more evenly spread within populations is unresolved despite important implications for the management of obesity and related disorders at population and individual levels (2, 8–10).

The genetic variants responsible for obesity susceptibility remain largely unknown. Genome-wide association studies (GWAS) have identified significant associations with >200 markers with small effects on BMI (polygenes), together explaining only approximately 3-4 % of total variance compared to family-based heritabilities (h^2) of 50-75% (11). Few causative mechanisms responsible for these phenotypically weak associations are known (4). The sources of the h^2 unaccounted for by GWAS are uncertain; proposals include overestimation of h^2, large numbers of common genetic variants with small, statistically insignificant effects on phenotypes (12, 13) and importantly, candidates not tested in most GWAS. Among the latter are rare genetic variants with large phenotypic

effects and G x OE interactions (4). Recently significant G x OE interactions have been detected in

individual GWAS loci and in composite genetic risk scores, which explain little of the missing

component of h^2 (14, 15).

A family history of diabetes (FH) is a potent, predominantly genetic (16, 17) risk factor for diabetes

diagnosis (DM) and for obesity-related phenotypes (18–21) consistent with the strong association

between type 2 DM and overweight/obesity. Familial effects on obesity-related phenotypes in adults

are also predominantly genetic (3, 22), so to the extent that the DM generating FH is of type 2

(approximately 94% of DM in the US population (23)), FH is a prevalent and readily obtained marker

of genetic susceptibility both to diabetes and to the obesity commonly preceding it. We have reported

evidence from a small sample of a multi-modal effect of FH on a composite adiposity index

consistent with segregation in families of discrete obesity risk (21). Polygenic risk scores (PRS)

based on large numbers of small effects are expected to be, and appear to be, normally-distributed

(24, 25) and cannot account for familial segregation of risk. The present work is based on the

alternative hypothesis that individually rare, but collectively common, genetic variants with large

phenotypic effects are the source of most of the missing h^2 and of most of G x OE, and that their

effects can be detected through analyses of phenotypic segregation in high-risk families (26).

The Continuous National Health and Nutrition Examination Survey (NHANES) is a continuing

(1999-) large-scale population-based survey incorporating an index of adiposity (Body Mass Index,

BMI) and first-degree FH (FH_0/FH_1) together with potential covariates and confounders. Although

BMI has recognized limitations as an adiposity phenotype (26, 27) it is the basis for most large-scale

genetic studies and like other authors, we assume that a large enough scale and appropriate modeling

of covariates will reduce effects of imprecision and bias (11). We aimed to test in a large multi-cycle

NHANES sample for the presence of familial segregation of genetic risk and to estimate the

contribution of FH, and by extension all discrete genetic risk, to recent secular trends in adult BMI.

Our results support a predominant role for large genetic effects interacting with OE in the obesity

"epidemic".

96   Data and Methods

Data

We used data from the 1999-2000, 2005-2006 and 2013-2014 cycles of NHANES

(https://www.cdc.gov/nchs/nhanes/index.htm accessed 25 Aug 2017). We extracted records for

participants age 20-65 years with non-missing gender, race/ethnicity, BMI, diabetes family history

(FH) and diabetes diagnosis data, and current smoking status if available. The definitions of two

fields changed over the sampling period: 1) Diabetes family history was defined in terms of 1° and 2°

relatives in 1999-2000 but by 1° relatives only in subsequent cycles. We recoded the 1999-2000

diabetes family history data to conform to the later definition using the separately collected data for

affected parents and siblings. 2) The self-identified race/ethnicity field (RIDRETH1) code was used

excluding other races (OR) to maintain consistency across cycles (Supplementary Methods). We

excluded from the primary analyses subjects diagnosed with diabetes because of possible

confounding by effects of either diabetes or diabetes therapies on BMI. The resulting data set is

summarized in Table 1.

Statistical analyses

Approach

We treat the data as a convenience sample and took no account of the sampling weights provided by

NHANES to permit nationally representative estimates. Our results are not intended to be

representative of the US population.

Our primary analyses are based on non-parametric visualization (kernel-smoothing) and analyses

(conditional quantile regression, CQR) of distributions requiring no prior distributional assumptions.

Parametric fits to multimodal distributions were then used to quantify the contributions of the

predicted large genetic effects model. FH(0/1) is treated as a binary genetic risk marker and calendar

time as a continuous surrogate of OE. Effects of OE interacting with FH were assessed in CQR

models, and also in least-squares ANCOVA models using bootstrap resampling to minimize

distributional assumptions in the calculation of effect size estimates and errors. All analyses were

performed using R 3.3.1 (28).

Summary statistics

Heterogeneity of the samples across cycles was assessed by Chi square test for categorical variables

and by one-way ANOVA for age. Effects on BMI were assessed by ANCOVA against continuous

time (yr = calendar start year - 1999). Effects on phenotype SD's were assessed by Bartlett's test in

one-way ANOVA models.

Adjustment

Prior to analysis BMI was adjusted for effects of age, gender and race/ethnicity in a linear model (age

$+ age^2 +$ race x gender). The adjustment model accounted for 4 % of the total variance in BMI

(Supplementary Table S1).

Distributions

Visualization

The effect of FH on the distribution of adjusted BMI was visualized using kernel-smoothed density

estimates by FH status (R base function density). The degree of smoothing is controlled by the

bandwidth parameter (bw) which was obtained in the full non-diabetic data set (bw = 0.99) from a

measure of the dispersion of the data (29). This produces a continuous distribution function and is

widely used to visualise features of potential interest which may be obscured in histograms. The

credibility of apparent effects of FH on the shape of the distribution was assessed by post-hoc

analysis of density differences ($FH_1 - FH_0$) by quantiles (n=20) of the full sample. Mean density

differences with SE were obtained by quantile by bootstrap resampling with replacement (1000

draws, stratified by FH status with resample sizes = strata sizes) and tested against zero using a Z-

test.

Conditional Quantile Regression (CQR)

Conditional quantile regression is a powerful tool for analyzing the effect of covariates on

distributions without assumptions of distributional shape. In contrast to ordinary least-squares (OLS)

regression which characterizes effects on global features of a distribution, CQR analyses local effects

of covariates independently at any specified quantiles and can detect and test variations in covariate

effects across quantiles. Originating in econometrics (30) it is now used in other areas including

genetics (31). In the CQR framework developed by Abadi et al (14) for analysis of genomic markers,

trends in effect sizes across quantiles represent interactions between genetic effects and unobserved

environmental and/or genetic factors. We treat FH as a binary genetic risk marker (FH_0/FH_1) and a

linear trend in quantile regression coefficients (β_{1i}) across quantiles (τ_i) represents summed linear

interactions of FH with unobserved factors. We analysed the effects of FH on adjusted BMI by CQR

using the R package quantreg. The effect of all interactions on the FH effect was tested in in a 2

parameter linear model:

for each quantile τ_i in y,

$$y(\tau_i \mid FH=fh)= \beta_{0i} + \beta_{1i} * FH$$

163   where $y(\tau_i \mid FH=fh)$ = the ith quantile of adjusted BMI conditional on the value of FH (0,1), the

164   intercept β_{0i} is the ith quantile value in FH_0 and β_{1i} is the FH effect size in quantile i.

166   The interaction between FH and continuous calendar time was estimated in the ANCOVA model:

$$y(\tau_i \mid FH=fh)= \beta_{0i} + \beta_{1i} * FH + \beta_{2i}*yr + \beta_{3i}*FH*yr$$

where yr = cycle start year – 1999, β_{0i} is the ith quantile value in FH_0 at yr =0 and β_{2i} and β_{3i} are CQR

coefficients for time and time*FH interaction effects in quantile i.

Equality of CQR parameter effect sizes across quantiles was tested using a Khmaladze test (32) (R

package quantreg).

The shape of the interaction relationship on the BMI scale was assessed in an OLS relationship

between quantile coefficients β_1 and β_0, which represents a linear model of a transformed quantile-

quantile plot between FH_1 and FH_0 after adjustment for covariates, and models a unimodal effect.

Residuals from the relationship were tested against randomness using a Wald-Wolfowitz runs test (R

package randtests), a simple, general non-parametric test of randomness in ordered binary

observations. The number of runs is compared to expectations under a normal approximation to the

random sampling distribution. Randomness is rejected in a two-sided test if the number of observed

runs is too high indicating high frequency oscillations, or too low indicating longer structures. We

applied a left-sided runs test to residuals from regression relationships testing for large structures and

interpret significance as excluding a unimodal effect and therefore as evidence of multiple modes in

the effect.

Parametric fits

We obtained fits to a 3-component normal mixture distribution representing a simple Mendelian

model of fixed genetic effect using an expectation-maximization algorithm (normalmixEM function

in the R package mixtools) . The models are characterised by the fitted means (μ_i), standard

deviations (σ_i) and mixing proportions (λ_i) of the three component distributions. Full model fits

were obtained in FH_1 but were not obtainable in FH_0 or DM_1 groups and we constrained μ_i in those

groups to values estimated in FH_1 in order to obtain comparable estimates of σ_i and λ_i. Risk allele

frequencies (q) under an additive Mendelian model of large genetic effects were calculated from the

fitted λ_i:

$$q = 0.5*\lambda_2 + \lambda_3$$

where λ_2 and λ_3 represent the proportions of carriers of 1 and 2 risk alleles respectively. Within-

sample consistency of calculated q across the three groups analysed was assessed by comparing

fitted q_{FH_1} with the prediction from random mating of DM_1 into the full sample:

$$\text{predicted } q_{FH_1} = (q_{DM} + \text{n-weighted mean}(q_{DM}, q_{FH_1}, q_{FH_0}))/2$$

Secular trends

Effects of diabetes family history status (FH_0, FH_1) and continuous time (yr = calendar start year -

1999) on adjusted BMI and its standard deviation (SD) were assessed in ANCOVA models of the

form:

$$y = \beta_0 + \beta_1 * FH + \beta_2 * yr + \beta_3 * FH * yr$$

where y = adjusted BMI mean or SD by FH status (0/1) and cycle, and yr = cycle start year – 1999.

Each OLS fit estimated 4 parameters from the 6 data points. Mean parameter estimates with 95% CI

were obtained by bootstrap resampling with replacement (1000 draws stratified by FH status and

cycle with resample sizes = strata sizes).

Comparison of cross-sectional and secular trend effects

Effects of FH on BMI distribution and on secular trends in BMI were compared by calculating the

contribution (%) of FH_1 to the effect in the full non-diabetic sample for calculated risk allele

frequency (q%) and to the slope (ß%) of the relationships between time and BMI in ANCOVA model

described above. Mean (± SE where possible) q% and ß% were calculated in the relevant bootstrap

samples.

Results

Participant characteristics

Data from 9102 non-diabetic subjects met the inclusion criteria, approximately equally distributed

across the 3 cycles. Gender balance varied little but there was a cycle effect in race/ethnicity, most

obvious in the reduced representation of MA in the two later cycles. Average age varied across cycles

but not its SD, while adjusted BMI and its SD showed linear trends with cycle time. FH_1 prevalence

was higher in the two later cycles compared to 1999-2000 as was DM_1 prevalence. Current smoking

status was predominantly missing in the data (55%) and was not included in the BMI adjustment

model. However smoking status was not related to FH whether analysed in the full data ($X^2 = 2.80$, 2

df , p = 0.25) or in those with non-missing smoking status ($X^2 = 0.43$, 1 df , p = 0.51), hence is

unlikely to confound analyses of FH effects. The mean age at diagnosis of DM (43.6 yr) is consistent

with predominantly type 2 DM in the sample.

Distributions

Visualization

Adjusted BMI in the non-diabetic sample showed an apparently unimodal distribution, right-skewed

compared to a normal model and closer to a log-normal model (Fig 1A). When visualized by FH

status (Fig 1B) the predicted multimodality in FH_1 was indicated with modes in the normal weight,

overweight and obese regions of the BMI distribution. In contrast FH_0 showed an apparently

unimodal distribution and a difference in shape between the two groups was supported by the pattern

of formal significance in the post-hoc analysis of density differences between groups (Fig 1B). BMI

distribution in the diabetic sample appeared to be depleted in the lower mode and enriched in the

upper modes compared to FH_1 (Fig 1C).

CQR

Analysis of the effect of FH status on the shape of the BMI distribution using CQR demonstrated

increasing FH effect size at higher levels of BMI ($p \leq 0.01$, Fig 2A, main panel), indicating strong

interactions between FH_1 and other variables not included in the model. FH_1 effect size ranged from

< 1 kg/m^2 in the lower quantiles to ~ 3 kg/m^2 in the upper quantiles, substantially different in both

regions to the OLS estimate (1.7 kg/m^2). The approximate linearity of FH_1 effects by quantile seen in

main panel of Fig 2A supports simple 1st order interactions, with that linearity emphasised by an OLS

analysis on the BMI scale between quantile coefficients ß$_1$ (FH_1) and ß$_0$ (FH_0) (slope $= 0.148 \pm 0.003$

(SE), $R^2 = 0.98$). Adjustment for main and interaction effects of calendar time in the model

(Supplementary Fig S2) weakened the trend in ß$_1$ across quantiles ($p > 0.1$, Fig 2B main panel) and

the OLS relationship between ß$_1$ and ß$_0$ (slope $= 0.103 \pm 0.006$, $R^2 = 0.86$), supporting the conclusion

that calendar time is a strong surrogate of obesogenic environmental influences interacting with

genetic factors as represented by FH status.

While the overall OLS relationship between $ß_1$ and $ß_0$ was linear there was strong evidence for

additional non-linear structure in an analysis of residuals from the relationship (Fig 2A inset) which

found significant non-randomness (runs test, p = 7.4 x 10^{-5}) attributable to a small number of broad

excursions. Adjustment for time effects accentuated the pattern (Fig 2B inset, runs test p = 2.2 x 10^{-5}).

While these patterns in the conditional quantiles do not map directly onto the unconditional quantile

plots in Fig 1B, by rejecting the unimodal effect hypothesis they add strong support to the conclusion

that FH_1 has multimodal discrete effects on BMI. The pattern of residuals after adjustment for time

effects (Fig 2B inset) exhibits 3 broad peaks, with suggestions of finer structure particularly in the

central peak. The lower peak in the low-normal weight range may derive from the presence of type 1

diabetes family history in the sample, while the upper two are consistent with the predicted discrete

effects of FH_1 derived from type 2 diabetes.

Parametric analysis

The distribution of BMI in FH_1 appears consistent with a simple Mendelian model and fitting a 3-

component normal distribution model to the FH_1 data resulted in robust estimates of component

means and separations (Fig 3A), as well as mixing coefficients and SDs (Table 2). Approximately

50% of FH_1 occupied the upper two modes and separation between modes accounted for

approximately 40% of the total variance in adjusted BMI with the remainder assigned to dispersion

within modes (Fig 2B). Under a Mendelian model the variance due to mode separation represents a

lower bound on the contribution of large effects as some of the dispersion within modes represents

variance in effect sizes of individual contributing causal loci (see Discussion) which will contribute to

the ~60% of variance assigned to within-modes. Estimates of component SDs and mixing proportions

with component means, constrained for FH_0 and DM_1 to those identified in the FH_1 data, support

enrichment in the two upper components in FH_1 compared to FH_0 (48% vs. 33%) and more strongly

in DM_1 (72%). Predicted risk allele frequencies in FH_1 (q - Table 2) express these distributional

properties in Mendelian terms and show within-sample consistency in that q_{FH1} predicted from

random mating of DM_1 (0.37) is not different to the fitted estimate (0.30 ± 10).

Secular trends

Adjusted BMI mean (Fig 2A) and SD (Fig 2B) increased over the sampling period significantly faster

in FH_1 compared to FH_0 in the bootstrapped ANCOVA model, and estimates of ß and Δß in the mean

data were indistinguishable from the OLS estimates provided by the CQR analysis (Supplementary

Fig S2). Similar results were obtained with log-transformed BMI (Supplementary Fig S3). FH_1

accounted for 62% of the BMI mean trend and 60% of the BMI SD trend in this sample over the

period 1999-2014, effects similar in magnitude to the estimated FH_1 contribution to the sample risk

allele frequency (50%, Supplementary Table S2).

Discussion

Summary

We tested the prediction of segregation of discrete effects of FH on adult BMI (21), modeled as

modes of distribution, and estimated the contribution of FH_1 to recent trends in BMI mean and

dispersion in a large population-based sample. The results support a predominant role in the recent

obesity "epidemic" for rare genetic variants with large effects interacting with OE.

Segregation of genetic susceptibility

The non-parametric analysis provided evidence for a multi-modal distribution in the FH_1 group

consistent with the prediction of segregation of large genetic effects in families (21). Multi-modality

was supported by the analysis of density differences between FH_1 and FH_0 by unconditional quantiles

(Fig 1B) and by evidence of shape in the relationship between CQR coefficients on the BMI scale

(Fig 2A&B insets). Polygenic risk scores (PRS) in population-based samples are expected to be

normally-distributed, and appear to be so (24, 25). Any elevated polygenic obesity risk in DM_1 will

dilute into the mating population resulting in a right-shifted distribution in FH_1 compared to FH_0, not

multi-modality. Alternative explanations for multi-modality might be discrete stratification of OE

which seems unlikely, or un-modeled interactions between FH and other covariates. Un-modeled

interactions between FH and stratified residual confounders may exist and contribute but we found no

evidence for this in plots of distributions by gender and race/ethnicity (Supplementary Fig S1) or in

an analysis of smoking status against FH. Discrete inheritance of genetic variants with large effects

remains the most likely explanation for multi-modality in the BMI distribution.

Approximately 40% of the adjusted BMI variance in FH_1 was accounted for by between-modes

variance (Fig 3B) but this represents a lower bound since the identified modes are likely to be

synthetic ie composed of a range of effect sizes due to rare variants at different loci. Indications of

fine structure within the broad peaks (Fig 2B inset) are suggestive. Examples of rare variants with

large effects on BMI in adults (ß) are known from studies of candidate genes and monogenic obesity

loci (26) while more recently a common variant in Samoans (EAF = 0.26, ß $\approx$ 1.4 kg/m^2), very rare

in other populations (33), and an African-specific rare variant (EAF = 0.008, ß =4.6 kg/m^2)

undetected in Europeans and Asians (34) have been identified by GWAS. Overall, ß in these nine

examples ranges from 1.4- 9 kg/m^2 and a similar range in effect sizes in the NHANES sample would

contribute substantially to the within-mode variance estimated here. A combination of within-subject

variance (~5% (35)) with polygenic variance (~ 5% (4)) sets a lower bound for within-modes

variance and hence the upper bound for between modes, implying that between 40% and 90% of total

variance in FH_1 may be attributed to large genetic effects.

G x OE

FH_1 is a prevalent (36%) and powerful determinant of the rate of change of mean BMI and its

dispersion over time, accounting for 62% of the BMI trend and 60% of the BMI SD trend in this

sample over 1999-2014. Under a polygenic model the familial risk would be distributed normally

over FH_1 which would then be a marker of a large fraction of the at-risk population. However under

the Mendelian model supported here genetic risk would segregate in families and only approximately

50% of FH_1 would acquire the excess familial risk and only ~18% of the sample would then account

for ~60% of the trends. Individuals with DM_1 must represent a fraction of individuals with elevated

genetic obesity risk and it is likely that the remainder, particularly those with a family history of

obesity without DM, would substantially increase the genetic component of the trends consequent to

the high heritability of BMI (22). This Mendelian model is internally consistent in estimates of risk

allele frequencies (q) in FH_1, FH_0 and DM_1 (Table 2) and in comparisons of FH_1 effect sizes in cross

section (q, ~50%) and over time (ß, ~60%) (Supplementary Table S2). Our results support the

proposition that the largest part and perhaps all of recent trends in mean and dispersion of BMI are

due to a minor subset of individuals with elevated genetic susceptibility to OE.

Limitations

The design and interpretation of fits to parametric mixture distribution models involves choices

concerning the number of components, parameter starting values and algorithms, and fit to a specific

model cannot be taken in isolation as support for its structural validity. We base our choice and

structural interpretation of 3-component normal mixture model fits and parameters on the *a priori*

hypothesis of Mendelian segregation of obesity risk in families (21) supported by the non-parametric

distributional plots (Fig 1B,C) and CQR analysis (Fig 2A,B insets). Like Abadi et al (14) we

interpret interactions in the CQR analysis as predominantly G x OE although a contribution from G x

G interactions cannot be excluded. Our interpretation is supported by the effects on the interaction of

including calendar time in the CQR model (Fig 2B). Other limitations discussed above include our

inability to exclude discreet stratification of OE and the possible influence of unmeasured/unknown

confounders of FH.

Conclusions

We conclude that a Mendelian model of individually rare but collectively common genetic risk

variants with large effects interacting with OE provides a plausible quantitative explanation for recent

trends in obesity and should be favored over a predominantly polygenic model which does not. The

evidence for a predominant role for polygenes (13) can appear to be strong (eg "Polygenic obesity is

the most common form of obesity in modern societies..." (36)) but recent interpretations seek to

explain the still missing heritability in obesity in terms of unidentified large genetic effects and G x

OE (4, 37) and recommend a renewed focus on family-based designs and on specific populations in

which large effect variants may be enriched (33, 34). Our results strengthen that view by showing that

a model based on unidentified segregating variants with large effects interacting with OE can account

for the largest part of the secular trend in obesity and its dispersion in a large population-based

sample. We can't predict what positive clinical or public health changes might follow from these

findings but they should help refocus research away from the under-performing but entrenched big

idea (38) that is polygenic obesity.

Supplementary information is available at The International Journal of Obesity's website.

Supplementary Methods: Race/Ethnicity coding

Supplementary Table S1: BMI-adjustment model parameters

Supplementary Table S2: Comparison of effects of diabetes family history on secular trends in

adjusted BMI and its SD and on calculated risk allele frequencies (q) in non-diabetic participants.

Supplementary Fig S1: Kernel-smoothed adjusted BMI densities by FH, gender and race/ethnicity

Supplementary Fig S2: time and time x FH interaction effects in a Conditional quantile regression

of adjusted BMI in an ANCOVA model.

Supplementary Fig S3: Secular trends in log-transformed adjusted BMI mean and SD

Competing Interests

The authors declare no competing financial or non-financial interests.

References

1. Di Cesare M, Bentham J, Stevens GAA et al. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. Lancet. 2016;387:1377-1396.

2. Krishna A, Razak F, Lebel A, Davey Smith G, Subramanian SV. Trends in group inequalities and interindividual inequalities in BMI in the United States, 1993–2012. Am J Clin Nutr. 2015;101:598-605.

3. Silventoinen K, Jelenkovic A, Sund R et al. Differences in genetic and environmental variation in adult BMI by sex, age, time period, and region: an individual-based pooled analysis of 40 twin cohorts. Am J Clin Nutr. 2017;106:457-466.

4. Loos RJ. The genetics of adiposity. Curr Opin Genet Dev. 2018;50:86-95.

5. Rokholm B, Silventoinen K, Ängquist L, Skytthe A, Kyvik KO, Sørensen TI. Increased genetic variance of BMI with a higher prevalence of obesity. PLoS One. 2011;6:e20816.

6. Franks PW, McCarthy MI. Exposing the exposures responsible for type 2 diabetes and obesity. Science. 2016;354:69-73.

7. Hall KD. Did the Food Environment Cause the Obesity Epidemic? Obesity. 2018;26:11-13.

8. Kivimaki M, Stenholm S, Kawachi I. The widening BMI distribution in the United States[letter]. Am J Clin Nutr 2015;101(6):1307-1308.

9. Jenkins AB, Campbell LV. Variation in Genetic Susceptibility Drives Increasing Dispersion of Population BMI[letter]. Am J Clin Nutr 2015;101:1308.

10. Razak F, Smith GD, Krishna A, Lebel A, Subramanian SV. Variation in genetic susceptibility drives increasing dispersion of population BMI Reply[letter]. Am J Clin Nutr 2015;101(6):1308-1309.

11.   Speakman JR, Loos RJF, O'Rahilly S, Hirschhorn JN, Allison DB. GWAS for BMI: a treasure

trove of fundamental insights into the genetic basis of obesity. Int J Obes. 2018;42:1524-1531.

12.   Locke AE, Kahali B, Berndt SI et al. Genetic studies of body mass index yield new insights for

obesity biology. Nature. 2015;518:197-206.

13.   Khera AV, Chaffin M, Wade KH et al. Polygenic Prediction of Weight and Obesity

Trajectories from Birth to Adulthood. Cell. 2019;177:587-596.e9.

14.   Abadi A, Alyass A, Robiou du Pont S et al. Penetrance of Polygenic Obesity Susceptibility

Loci across the Body Mass Index Distribution. Am J Hum Gen. 2017;101:925-938.

15.   Nagpal S, Gibson G, Marigorta U. Pervasive Modulation of Obesity Risk by the Environment

and Genomic Background. Genes. 2018;9:doi: 10.3390/genes9080411.

16.   Hemminki K, Li X, Sundquist K, Sundquist J. Familial risks for type 2 diabetes in Sweden.

Diabetes Care. 2010;33:293-297.

17.   Willemsen G, Ward KJ, Bell CG et al. The Concordance and Heritability of Type 2 Diabetes in

34,166 Twin Pairs From International Twin Registers: The Discordant Twin (DISCOTWIN)

Consortium. Twin Res Hum Genet. 2015;18:762-771.

18.   Ghosh A, Liu T, Khoury MJ, Valdez R. Family History of Diabetes and Prevalence of the

Metabolic Syndrome in U.S. Adults without Diabetes: 6-Year Results from the National Health

and Nutrition Examination Survey (1999-2004). Public Health Genom. 2010;13:353-359.

19.   Tirosh A, Shai I, Afek A et al. Adolescent BMI trajectory and risk of diabetes versus coronary

disease. N Engl J Med. 2011;364:1315-1325.

20.   Scott RA, Consortium I. The link between family history and risk of type 2 diabetes is not

explained by anthropometric, lifestyle or genetic risk factors: the EPIC-InterAct study.

Diabetologia. 2013;56:60-69.

21.   Jenkins AB, Batterham M, Samocha-Bonet D, Tonks K, Greenfield JR, Campbell LV.

Segregation of a latent high adiposity phenotype in families with a history of type 2 diabetes

431  mellitus implicates rare obesity-susceptibility genetic variants with large effects in diabetes-

432  related obesity. PLoS One. 2013;8:e70435.

433 22. Stryjecki C, Alyass A, Meyre D. Ethnic and population differences in the genetic predisposition

434  to human obesity. Obes Rev. 2018;19:62-80.

435 23. Xu G, Liu B, Sun Y et al. Prevalence of diagnosed type 1 and type 2 diabetes among US adults

436  in 2016 and 2017: population based study. BMJ. 2018;362:k1497.

437 24. Llewellyn CH, Trzaskowski M, Plomin R, Wardle J. From modeling to measurement:

438  developmental trends in genetic influence on adiposity in childhood. Obesity. 2014;22:1756-

439  1761.

440 25. Rask-Andersen M, Karlsson T, Ek WE, Johansson A. Gene-environment interaction study for

441  BMI reveals interactions between genetic factors and physical activity, alcohol consumption

442  and socioeconomic status. PLoS Genet. 2017;13:e1006977.

443 26. Jenkins AB, Campbell LV. Future management of human obesity: understanding the meaning

444  of genetic susceptibility. Adv Genomics Genet. 2014;4:219-232.

445 27. Müller MJ, Geisler C, Blundell J et al. The case of GWAS of obesity: does body weight control

446  play by the rules. Int J Obes. 2018

447 28. R Development Core Team. R: A Language and Environment for Statistical Computing. R

448  Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/; 2016.

449 29. Sheather SJ, Jones MC. A Reliable Data-Based Bandwidth Selection Method for Kernel

450  Density Estimation. J Roy Stat Soc B Met. 1991;53:683-690.

451 30. Koenker R. Quantile Regression: 40 Years On. Ann Rev of Econ. 2017;9:155-176.

452 31. Briollais L, Durrieu G. Application of quantile regression to recent genetic and -omic studies.

453  Hum Genet. 2014;133:951-966.

454 32. Koenker R, Xiao Z. Inference on the quantile regression process. Econometrica. 2002;70:1583-

455  1612.

456   33.   Minster RL, Hawley NL, Su CT et al. A thrifty variant in CREBRF strongly influences body

457   mass index in Samoans. Nat Genet. 2016;48:1049-1054.

458   34.   Chen G, Doumatey AP, Zhou J et al. Genome-wide analysis identifies an African-specific

459   variant in SEMA4D associated with body mass index. Obesity. 2017;25:794-800.

460   35.   Wormser D, White IR, Thompson SG, Wood AM. Within-person variability in calculated risk

461   factors: Comparing the aetiological association of adiposity ratios with risk of coronary heart

462   disease. Int J Epidemiol. 2013;42:849-859.

463   36.   Albuquerque D, Nóbrega C, Manco L, Padez C. The contribution of genetics and environment

464   to obesity. Br Med Bull. 2017;123:159-173.

465   37.   Saeed S, Arslan M, Froguel P. Genetics of Obesity in Consanguineous Populations: Toward

466   Precision Medicine and the Discovery of Novel Obesity Genes: Obesity and Consanguinity.

467   Obesity. 2018;26:474-484.

468   38.   Joyner MJ, Paneth N, Ioannidis JP. What Happens When Underperforming Big Ideas in

469   Research Become Entrenched? JAMA. 2016;316:1355-1356.

Figure legends

<u>Fig 1:</u> Distribution of adjusted BMI in non-diabetic (DM_0) and diabetic (DM_1) participants in

combined NHANES 1999-2000, 2005-2006 and 2013-2014 cycles. A) Full non-diabetic sample

(n=9102) binned by quantiles (n=20) with superimposed kernel-smoothed and fitted densities in

normal and log-normal models. B) Kernel-smoothed adjusted BMI density by FH status. * significant

(p<0.05) difference in empirical density ($FH_1 - FH_0$) by quantiles of the full sample shown in panel

A. C) Kernel -smoothed adjusted BMI density in non-diabetic FH_1 (n=3297) compared to diabetic

participants (n=793).

<u>Fig 2:</u> Conditional quantile regression effects of FH on adjusted BMI in non-diabetic participants in

models consisting of FH alone (A) and in interaction with calendar time (B). The main panels show

the FH main effect size (β_1) by quantile with 95% CI (grey-shaded area), the OLS estimate of the

average effect (solid black line) with 95% CI (dotted black lines) and p-values associated with

Khmaladze tests for equality of β_1 across quantiles. The insets show the patterns of residuals ($\Delta\beta_1$)

from linear regressions of β_1 against conditional quantiles in FH_0 (β_0) , with 95% CI on the fits

(dotted lines) around the lines $\Delta\beta_1 = 0$ representing perfect fits.

<u>Fig 3:</u> A) Adjusted BMI density in FH_1 by quantile (grey bars) and kernel-smoothed (black line) with

fits to a three component normal mixture distribution. B) Estimated contributions in FH_1 of the

components of the mixture distribution to the prevalence (mixture coefficients, λ) and variance of

adjusted BMI.

<u>Fig 4:</u> Effects of diabetes family history ($FH_{0/1}$) on linear secular trends in age-, gender- and

race/ethnicity-adjusted BMI mean ± se (A) and standard deviation ± se (B). Parameter estimates

with 95% CI were obtained in ANCOVA models by stratified bootstrap resampling of all non-

diabetic individuals (see Methods). Dotted lines enclose 95% CI on fitted values at each point; β =

regression slope vs. time (kg/m^2 per year); $\Delta\beta = \beta_{FH1} - \beta_{FH0}$ (95%CI).

Table 1: Participants by diabetes status (DM_0/DM_1)

	All cycles	1999-2000	2005-2006	2013-2014	$p^{\dagger}$
		NHANES cycle			
		DM_0			
n	9102	2865	3076	3161	-
Gender (F%)	53	55	54	52	0.06
Race/Ethnicity (%) (MA/OH/NHW/NHB)[¥]	23/8/46/ 23	30/7/44/20	24/4/49/24	16/11/48/25	7.8×10^{-56}
Age (yr): Mean	40.5	40.7	39.4	41.5	4.5×10^{-9}
SD	13.2	13.2	13	13.2	0.63
BMI (kg/m^2)*: Mean	28.7	28.3	28.6	29.2	4.6×10^{-9}
SD	6.6	6.3	6.4	7.1	1.2×10^{-11}
Diabetes Family History (Y%)	36	29	42	38	4.1×10^{-25}
Current smoking (%, Y/N/missing)	25/20/55				-
		DM_1			
n (%)	793 (8.0)	211 (6.9)	252 (7.6)	330 (9.5)	0.0003
Age at Diagnosis (yr)	43.6	44.3	42.9	43.6	0.47

† Cycle effects (p) by ANOVA (age), ANCOVA (BMI), Bartlett's test (SD's) and Chi-squared test for categorical variables.

[¥] MA = Mexican American, OH = Other Hispanic, NHW = Non-Hispanic White, NHB = Non-Hispanic Black

* Adjusted for age, gender and race/ethnicity in a linear model (see Table S1).

Table 2: Three-component normal mixture distribution fits to adjusted BMI by FH and DM status[¥]

	Component	FH$_0$	DM$_0$ FH$_1$	DM$_1$
mean	1	*	25.8±1.0	*
	2	*	32.1±1.8	*
	3	*	40.6±2.5	*
sd	1	3.8	3.8±0.9	3.7
	2	5.7	5.1±1.2	4.5
	3	10	8.2±0.8	8.4
λ	1	0.67	0.52±0.16	0.28
	2	0.29	0.37±0.15	0.47
	3	0.04	0.11±0.06	0.25
q	-	0.18	0.30±0.10 (0.37)[§]	0.49

[¥] ± bootstrap standard error for FH$_1$ only

* Means in FH$_0$ and DM$_1$ constrained to fitted values in FH$_1$

[†] Calculated risk allele frequency in an additive Mendelian model of large effects

[§] Predicted from DM$_1$ mating randomly into the full sample.