Reduction in standard metabolic rate after multigenerational exposure to

elevated temperatures in the wild

Natalie Pilakouta[1], Shaun S. Killen[1], Bjarni K. Kristjánsson[2], Skúli Skúlason[2], Jan

Lindström[1], Neil B. Metcalfe[1], and Kevin J. Parsons[1]

[1] Institute of Biodiversity, Animal Health, and Comparative Medicine, University of

Glasgow, Glasgow, UK

[2] Department of Aquaculture and Fish Biology, Hólar University, Sauðárkrókur, Iceland

Corresponding authors: Natalie Pilakouta (natalie.pilakouta@glasgow.ac.uk); Kevin

Parsons (kevin.parsons@glasgow.ac.uk)

Running title: Thermal divergence in metabolic rate

Keywords: climate change, countergradient variation, Krogh's rule, metabolic cold

adaptation, physiology, *Gasterosteus aculeatus*

Abstract

In light of global climate change, there is a pressing need to understand and predict the

capacity of populations to respond to rising temperatures. Metabolic rate is a key trait that is

likely to influence the ability to cope with climate change. Yet, empirical and theoretical

work on metabolic rate responses to temperature changes has so far produced mixed results

and conflicting predictions. Our study addresses this issue using a novel approach of

comparing fish populations in geothermally warmed lakes and adjacent ambient-

temperature lakes in Iceland. This unique 'natural experiment' provides repeated and

independent examples of populations experiencing contrasting thermal environments for

many generations over a small geographic scale, thereby avoiding the confounding factors

associated with latitudinal or elevational comparisons. Using Icelandic sticklebacks from

three warm and three cold habitats, we measured individual metabolic rates across a range

of acclimation temperatures to obtain reaction norms for each population. We found a

general pattern for a lower standard metabolic rate in sticklebacks from warm habitats, as

predicted by Krogh's rule. Metabolic rate differences between warm- and cold-habitat

sticklebacks were more pronounced at more extreme acclimation temperatures, suggesting

the release of cryptic genetic variation upon exposure to novel conditions, which can reveal

hidden evolutionary potential. Lastly, we found a stronger divergence in metabolic rate

between thermal habitats in allopatry than sympatry, indicating that gene flow may

constrain physiological adaptation when dispersal between warm and cold habitats is

possible. In sum, our study suggests that fish may diverge toward a lower standard

metabolic rate in a warming world, but this might depend on connectivity and gene flow

between different thermal habitats.

Introduction

Climate change poses a substantial threat to biodiversity, as rising temperatures are altering abiotic and biotic environmental conditions and imposing novel selection pressures on organisms (Crozier & Hutchings 2014). In light of this, there is now a pressing need to understand the capacity of populations to respond and adapt to increasing temperatures. This knowledge is critical for our ability to predict the impacts of climate change and to inform management and conservation efforts (Donelson et al. 2011, Sinclair et al. 2016, Campbell et al. 2017). Ectothermic animals are particularly vulnerable to changes in ambient temperature, because this directly influences their body temperature (Zuo et al. 2012). Ectotherms are therefore expected to adapt to climate change through plastic and/or evolutionary changes in a wide range of physiological, morphological, and behavioural traits (Crozier & Hutchings 2014).

A key trait that is likely to influence the capacity of ectothermic animals to cope with increasing temperatures is metabolic rate (Donelson et al. 2012). Standard and maximum metabolic rates are among the most commonly measured physiological traits and can have important fitness implications for organisms (White and Kearney 2013, Pettersen et al. 2018). Standard metabolic rate (SMR) refers to the minimum rate of energy throughput needed to sustain an animal at rest, whereas maximum metabolic rate (MMR) sets the upper limit for the capacity to perform oxygen-consuming physiological activities (Killen et al. 2016). Another important physiological variable is absolute aerobic scope (AS), which is calculated as the difference between an individual's SMR and MMR. AS represents an individual's total capacity for simultaneous oxygen-consuming tasks above maintenance requirements (e.g., digestion, activity, growth; Pörtner & Farrell 2008). In ectotherms, AS

may increase with temperature up to an optimum, after which it declines with further

temperature increases. AS has therefore been suggested to play a key role in the response of

fish populations to changing temperatures (Farrell 2016, Sandblom et al. 2016). Notably,

however, many fish species appear to show little or no change in AS with changes in

temperature (Lefevre 2016, Nati et al. 2016, Jutfelt et al. 2018), so the degree to which AS

limits adaptive responses to warming remains an open question.

It is also still unclear how SMR and MMR will evolve in response to climate change.

According to Krogh's rule (also known as metabolic cold adaptation), ectotherms living in

cold environments should have higher metabolic rates than those in warm environments,

when the two are observed at the same temperature (Krogh 1916, Gaston et al. 2009). This

view is in line with the concept of countergradient variation, which is a result of stabilising

selection favouring similar phenotypes in different environments (Marcil et al. 2006) and

can arise when genetic differences in a trait counteract environmental effects (Conover &

Schultz 1995). An alternative hypothesis is that natural selection will favour a reduction in

metabolic rate in animals living at low temperatures if the energetic cost of cellular

respiration exceeds the benefit of additional ATP production (Clarke 1980, Clarke 1991,

Clark 1993, Chown and Gaston 1999). These opposing predictions have led to much

controversy and debate about the effects of temperature changes on metabolic rate

adaptation (Huey and Berrigan 1996, Chown and Gaston 1999, Addo-Bediako et al. 2002).

Until now, empirical studies aiming to address this issue have focused on interspecific

and intraspecific variation in metabolic rate across contrasting thermal habitats, such as

populations from different latitudes or elevations (e.g., Tsuji 1988, White et al. 2012,

Gaitán-Espitia and Nespolo 2014). However, there are many confounding factors associated

with these comparisons. For example, high-latitude habitats are not only colder on average

but also experience more extreme temporal fluctuations in temperature and photoperiod

than low-latitude habitats. These confounding factors may partly explain the mixed results

of previous research on the relationship between thermal environment and metabolic rate

(Clarke 2003, White and Kearney 2013, Alton et al. 2017).

Here, we use a novel approach by comparing Icelandic populations of threespine

sticklebacks (*Gasterosteus aculeatus*) found in geothermally warmed lakes and adjacent

ambient-temperature lakes. This unique 'natural experiment' provides repeated and

independent examples of populations experiencing contrasting thermal environments for

many generations over a small geographic scale, thereby avoiding the confounding factors

associated with latitudinal or elevational comparisons. In addition, there are no substantial

differences in water chemistry between geothermally warmed and ambient-temperature

lakes, allowing us to isolate the effects of temperature. We have previously shown that there

is strong morphological divergence between sticklebacks from these warm and cold habitats,

suggesting local adaptation (Pilakouta et al. 2019).

Using sticklebacks from three warm and three cold habitats, we measured individual

metabolic rates across a range of acclimation temperatures (10°C, 15°C, and 20°C) to obtain

reaction norms for each population. We examined whether living in a warm environment

over several generations results in a suppressed metabolic rate, providing a powerful test of

the controversial Krogh's rule. If metabolic rate responds in a similar way to thermal habitat

across multiple populations, this would suggest that metabolic rate evolution in response to

climate change may be predictable (Bolnick et al. 2018). We also investigated whether the

steepness of metabolic rate reaction norms differs between populations from warm vs cold

environments. Given the potential importance of physiological adaptation for population

persistence (Donelson et al. 2011, Sinclair et al. 2016), our study could provide valuable

insights into the capacity of ectotherms to cope with climate change.

Methods

Study animals

Using unbaited minnow traps, we collected adult threespine sticklebacks from six freshwater

populations in Iceland in May–June 2016 (Table 1). Two of these populations were

allopatric, meaning that the warm and cold habitats were in neighbouring but separate

water bodies with no potential for gene flow (Figure 1). We also sampled two sympatric

warm-cold population pairs, where the warm and cold habitats were in the same water

body, with no physical barriers between them (Figure 1).

The cold habitats have all existed for thousands of years, since the last glacial period

(Einarsson et al. 2004), but there is some variation in the age of the warm habitats (Table 1).

The 'Mývatn warm' and Grettislaug sites have been naturally heated by geothermal activity

for over 2000 years (Hight 1965, Einarsson 1982). In contrast, the 'Áshildarholtsvatn warm'

habitat originated only 50–70 years ago, fed by excess hot water runoff from nearby

residences using geothermal heating.

Animal husbandry

After being transported to the University of Glasgow, these fish were kept at densities of 10-

15 individuals per 10-litre tank in a common recirculation system at 15°C. The tanks

contained plastic plants as shelter. Fish were fed *ad libitum* twice a day with a mixture of

frozen bloodworms, *Mysis* shrimp, and *Daphnia*. Two months before the experiments, all

fish were anaesthetised using benzocaine and marked with visible implant elastomer tags

(Northwest Marine Technology Inc) to allow individual identification. They were kept at a

12h light:12h dark photoperiod throughout the experiment.

Fish were acclimated to 10°C, 15°C, or 20°C for at least one month before measuring

their metabolic rate. Different individuals were used at each of these temperatures. The

intermediate temperature (15°C) is close to the maximum temperature experienced by fish

in cold habitats in the summer and the minimum temperature experienced by fish in warm

habitats in the winter (Table 1). The lowest temperature in this range (10°C) is not generally

experienced by warm-habitat fish in the wild, and the highest temperature in this range

(20°C) is not experienced by cold-habitat fish (Table 1). Exposing fish to these unfamiliar

temperatures allowed us to examine the release of cryptic genetic variation in metabolic rate

and thus hidden evolutionary potential to respond to thermal changes (Paaby & Rockman

2014, Shama 2017).

Metabolic rate measurements

We used intermittent flow-through respirometry to estimate individual metabolic rates by

measuring oxygen uptake. Sixteen cylindrical, glass respirometry chambers (83 ml) were

submerged in a 93-L experimental tank (780 mm × 570 mm × 210 mm) containing air-

saturated water. The water temperature within the experimental tank was maintained at

10°C, 15°C, or 20°C depending on the treatment. This was done using a thermostated

reservoir connected to the experimental tank by a thermoregulator (TMP-REG system,

Loligo Systems, Denmark), which allowed us to maintain the water temperature within

0.2°C of our target temperature for the entire 24-h trial period. To maintain good water

mixing and avoid an oxygen gradient in the respirometry chambers, we used a peristaltic

pump (Masterflex, Cole-Parmer), which moved water through the chambers and around an

external circuit of gas-impermeable tubing (Masterflex, Cole-Parmer). Oxygen

concentration in the chambers was measured every two seconds using four Firesting

channel oxygen meters with sixteen associated sensors (PyroScience GmbH, Aachen,

Germany). To account for bacterial respiration during the trials, background bacterial

oxygen consumption was measured before and after each trial in the 16 respirometry

chambers (see *Data analysis*). A UV filter connected to the experimental tank was also used

to sterilise the water and minimise bacterial respiration.

We fasted fish for 48 h before the trials because metabolic rate increases during

digestion (Killen 2014). Trials started around 14:00 each day. Immediately before being

placed into a respirometry chamber, each fish was subjected to exhaustive exercise by being

chased in a circular tank; this allowed us to measure their maximum metabolic capacity

(Killen et al. 2012, Clark et al. 2013, Killen et al. 2017). After complete exhaustion, which

always occurred within 2–3 min of chasing, fish were placed into individual respirometers.

Rates of oxygen uptake were then measured in 3-min intervals over a 15-min period, during

which the respirometers were sealed and the decrease in oxygen content was used to

calculate rate of oxygen uptake (see *Data analysis*). The maximum rate of oxygen uptake

measured during these five 3-min intervals was used as a proxy for the maximum metabolic

rate (MMR).

The fish were left in the respirometers undisturbed until around 14:00 the following

day. Every 9 min, an automated water pump (Eheim GmbH & Co. KG, Germany) would

switch on for 2 min flushing the respirometers with aerated water. Based on the decrease in

oxygen concentration during the 7-min off-cycle of the pumps, we calculated the rate of

oxygen uptake. Standard metabolic rate (SMR) was estimated as the lowest 10th percentile

of measurements taken throughout the measurement period (Dupont-Prinet et al. 2010,

Killen 2014), excluding the first 5 h during which oxygen consumption was elevated due to

handling stress (Killen 2014). Absolute aerobic scope (AAS) was calculated as the difference

between SMR and MMR. Factorial aerobic scope (FAS) was calculated as the ratio of

MMR and SMR.

During the trial period, the experimental bath was covered with black plastic to avoid

external disturbances. We also covered the sides of the respirometry chambers with opaque

material to prevent visual stimuli from other individuals in the same trial. Fish were

removed from the respirometer after 24 h, at which point we weighed them and returned

them to their initial holding tank. Our sample sizes ranged between 15 and 33 per

population per temperature (Supplementary Tables 1 & 2). Due to equipment failure during

a small proportion of our trials, we have slightly more measurements of MMR than SMR

and AS (Supplementary Tables 1 & 2).

Data analysis

Oxygen concentration data derived from Firesting software were analysed in LabChart 7

(ADInstruments Pty Ltd, Australia). To calculate oxygen consumption rates (MO_2), we used

the average slope of each 7-min off-cycle measurement period derived from the linear

regressions between oxygen consumption over time (mg O_2 h^{-1}). All MO_2 data were

corrected for the volume of the respirometry chamber and body mass of the fish. Using the

bacterial respiration data collected for all chambers before and after trial measurements, we

assumed a linear increase in bacterial oxygen consumption over time. The estimated

bacterial background respiration at any given timepoint was then subtracted from the MO_2

data.

Statistical analyses were performed using R version 3.5.1 (R Core Team 2017), and

figures were generated using the ggplot2 package (Wickham 2009). SMR, MMR, AAS,

FAS, and body mass were log-transformed for use in models because of a non-linear

relationship between metabolic rate and mass. For each response variable (SMR, MMR,

AAS, and FAS), we used general linear models (GLM) with the following explanatory

variables: thermal habitat (warm or cold), population pair (allopatric, sympatric 1, or

sympatric 2), acclimation temperature (10°C, 15°C, or 20°C), and all possible interactions

between these three factors. A significant effect of the three-way interaction between thermal

habitat, population pair, and acclimation temperature would indicate non-parallel

divergence of metabolic rate reaction norms across population pairs (Bolnick et al. 2018).

After checking for homogeneity of slopes, body mass was also included as a covariate in all

models to correct for the effects of mass on metabolic rate. All model assumptions were

tested and verified, and the statistical results reported below are the values from the full

models including all interactions. Our results for AAS (Table 2, Figure 2C) versus FAS are

largely similar (Supplementary Table 3, Supplementary Figure 1), and we only present the

results for AAS, which is considered to be more informative and robust (Halsey et al. 2018).

Results

As expected, larger fish had higher absolute metabolic rates, and metabolic rate tended to

increase with acclimation temperature (Table 2). Acclimation temperature explained over

30% of the variation in SMR but only 7% and 3% of the variation in MMR and AAS,

respectively (Table 2). SMR was therefore more variable than MMR and AAS in response

to acclimation temperature. Fish from warm and cold habitats also differed in the steepness

of their metabolic rate reaction norms (Figure 2). In terms of SMR, warm-habitat fish had a

steeper metabolic rate reaction norm than cold-habitat fish in the allopatric population pair

but a less steep reaction norm in sympatric population 1 (Figure 2A).

We found a statistically significant three-way interaction between thermal habitat,

population pair, and acclimation temperature on SMR and AAS (Table 2). There was a

marginally non-significant effect of this three-way interaction on MMR (Table 2). This

interaction indicates that the divergence in metabolic rate reaction norms between warm-

and cold-habitat fish varied across the three population pairs (Figure 2). For example, there

was a stronger divergence in SMR and MMR between the warm- and cold-habitat fish in

the allopatric populations than in the sympatric populations (Figure 2A and 2B). Similarly,

the strong effect of population pair and acclimation temperature×population pair on SMR

(Table 2) indicates that local adaptation may be driving variation in this trait across different

sites.

Differences in SMR between fish from warm and cold habitats tended to be more

pronounced at more extreme acclimation temperatures: in the allopatric population pair,

thermal divergence in SMR decreased with increasing acclimation temperature, and in

sympatric population 1, thermal divergence in SMR increased with increasing acclimation

temperature (Figure 2A). However, in sympatric population 2, there was no divergence in

SMR between warm- and cold-habitat fish at any of the three acclimation temperatures

(Figure 2A).

Similar to SMR, differences in MMR between thermal habitats tended to be more

pronounced at more extreme acclimation temperatures. Warm-habitat fish from sympatric

population 2 had a lower MMR and AAS than cold-habitat fish when both were acclimated

to 20°C (Figure 2B and 2C). In the other two population pairs, warm-habitat fish had a

lower MMR and AAS than cold-habitat fish at 10°C (Figures 2B and 2C).

Discussion

We have taken advantage of a unique study system of geothermally heated and ambient-

temperature populations to test whether fish in warm environments show a suppressed or

elevated metabolic rate compared to those in cold environments. We found a general

pattern for a lower SMR in sticklebacks originating from warm habitats, although the extent

of this effect varied depending on the population pair and acclimation temperature.

Interestingly, the SMR of warm-habitat sticklebacks at their naturally experienced

temperatures (15-20°C) was similar to the SMR of cold-habitat sticklebacks at their naturally

experienced temperature (10-15°C). This results in similar metabolic costs in these

contrasting thermal environments and is consistent with the predictions of countergradient

variation and Krogh's rule. Our findings therefore suggest that fish may evolve a lower

metabolic rate as global temperatures increase in response to climate change.

In our study system, some populations of sticklebacks living in warm or cold habitats

are in separate water bodies (allopatry), while others are found in different parts of the same

water body (sympatry). Given the potential for gene flow between sympatric but not

allopatric population pairs, we might expect sympatric pairs to be less phenotypically

divergent than allopatric pairs (Hendry and Taylor 2004, Pinho and Hey 2010).

Alternatively, sympatric pairs might be more divergent because of character displacement,

whereby differences between morphs are more pronounced in areas where they co-occur

and minimised in areas where their distributions do not overlap (Brown and Wilson 1956,

Losos 2011). Our findings show a stronger divergence in metabolic rate between

sticklebacks from warm and cold habitats in allopatry than in sympatry. This suggests that

gene flow may constrain physiological adaptation in natural populations where physical

dispersal between thermal habitats is possible (Lenormand 2002, Hendry and Taylor 2004).

It also indicates that metabolic responses to thermal habitat vary across populations, making

it difficult to predict metabolic rate evolution in response to climate change.

The two sympatric populations differed in the extent of thermal divergence in their

metabolic rate reaction norms: sympatric population 1 showed a greater degree of

divergence than sympatric population 2. It is possible that this variation is related to

differences in the age of these warm habitats (Table 1). In populations that have been

diverging for longer, there is more scope for natural selection and genetic drift to introduce

adaptive or stochastic phenotypic differences (Ord and Summers 2015). In our study system,

we might thus expect the younger population pair (i.e., sympatric 1) to be less divergent

than the older populations pair (i.e., sympatric 2). However, we found the opposite pattern,

whereby the young sympatric population showed a greater degree of divergence in

metabolism between fish from warm and cold habitats. In this young sympatric population,

there was thermal divergence in both SMR and MMR reaction norms, whereas in the older

sympatric population, there was no divergence in SMR and only a small divergence in

MMR. These results are even more surprising considering that the warm and cold habitats

of the younger sympatric population are only tens of meters apart, whereas they are a few

kilometres apart for the older sympatric population (Table 1). One possible explanation is

that there are differences in food availability in the two sympatric populations (White and

Kearney 2013). Low food availability in the warm habitat of sympatric population 1, along

with elevated temperatures, may favour a lower SMR to allow organisms to resist starvation

(Alton et al. 2017). Correspondingly, if there is high food availability in the warm habitat of

sympatric population 2 (O'Gorman et al. 2016), selection due to temperature and food

availability might be operating in opposite directions, resulting in a similar SMR in the cold

and warm habitats (Figure 2A).

Another interesting finding was a higher degree of variability in SMR than MMR and

AS in response to acclimation temperature. Across the three warm-cold population pairs,

acclimation temperature explained over 30% of the variation in SMR but only 7% and 3%

of the variation in MMR and AS, respectively. If rising ambient temperatures cause a

greater increase in SMR relative to MMR, this will lead to a decrease in AS (Donelson et al.

2011). Our results support this prediction, as warm-habitat sticklebacks tended to have a

lower AS than cold-habitat sticklebacks at a high temperature (20°C). It has been proposed

that the capacity to meet increased oxygen demands at elevated temperatures may

determine the persistence of fish populations in a warming climate (Donelson et al. 2011;

but see Lefevre 2016, Jutfelt et al. 2018), so it is important to better understand and predict

changes in AS in response to temperature changes (Sinclair et al. 2016).

By measuring metabolic rates across a range of temperatures, rather than a single

temperature as is typically done (Bruneaux et al. 2014), we were able to compare the

steepness of the metabolic reaction norms of warm versus cold populations. In terms of

SMR, warm-habitat sticklebacks had a steeper metabolic rate reaction norm than cold-

habitat sticklebacks in the allopatric population pair but a less steep reaction norm in

sympatric population 1. Despite these contrasting trends, a common pattern emerging from

our results was that differences between warm- and cold-habitat sticklebacks tended to be

more pronounced at more extreme temperatures ($10\,^{\circ}$C or 20°C). This suggests that cryptic

genetic variation was released upon exposure to these novel conditions, thus revealing

hidden evolutionary potential (Paaby & Rockman 2014, Shama 2017). If we had only

measured metabolic rates at an intermediate temperature, we would have detected little or

no differences between warm- and cold-habitat sticklebacks. Our findings thus demonstrate

that when studying individuals originating from different thermal environments in a

common garden setting, it is much more informative to compare their reaction norms rather

than use a single acclimation temperature (Bruneaux et al. 2014).

Moreover, using populations exposed to contrasting thermal habitats for many

generations allowed us to examine long-term responses to elevated temperatures (Huss et al.

2019). This avoids the limitations of short-term laboratory experiments that expose animals

to high temperature for only one or a few generations. Nevertheless, the relative

contribution of genetic change and plasticity in driving metabolic rate differences in this

study system is still unknown. A recent review suggested that metabolic rate is generally

highly heritable, although active metabolic rate tends to be more heritable than resting

metabolic rate (Pettersen et al. 2018). Further work on the heritability of metabolic traits in

this system would allow us to assess their potential to respond to selection.

In summary, we have shown a general pattern for a lower metabolic rate in fish from

warm habitats, which provides a powerful test of and support for the controversial Krogh's

rule. We also found evidence for a stronger divergence in metabolic rate between warm and

cold habitats in allopatry than sympatry, suggesting that gene flow may constrain

physiological adaptation when dispersal between thermal habitats is possible. By focusing

on natural populations living in contrasting thermal environments over a small geographic

scale, our study offers valuable insights into how fishes and other ectotherms might

physiologically adapt to global climate change.

Acknowledgements

We thank Joseph Humble for his assistance with animal husbandry. We are also grateful to

Iain Hill, Tiffany Armstrong, Anna Persson, and Kári Heiðar Árnason for their help with

fieldwork in Iceland.

Ethical statement

Our study adheres to the ASAB/ABS Guidelines for the Use of Animals in Research, the

institutional guidelines at University of Glasgow, and the legal requirements of the UK

Home Office (Project License P89482164).

Funding

The study was funded by a Natural Environment Research Council Grant

(NE/N016734/1) awarded to KJP, NBM, SSK, and JL.

References

Addo-Bediako A, Chown SL, Gaston KJ. 2002. Metabolic cold adaptation in insects: a

large-scale perspective. Functional Ecology 16(3):332-8.

Alton LA, Condon C, White CR, Angilletta Jr MJ. 2017. Colder environments did not

select for a faster metabolism during experimental evolution of Drosophila

melanogaster. Evolution 71(1):145-52.

Bolnick DI, Barrett RD, Oke KB, Rennison DJ, Stuart YE. 2018. (Non) parallel evolution.

Annual Review of Ecology, Evolution, and Systematics 49:303-30.

Brown WL, Wilson EO. 1956. Character displacement. Systematic Zoology 5(2):49-64.

Bruneaux M, Nikinmaa M, Laine VN, Lindström K, Primmer CR, Vasemägi A. 2014.

Differences in the metabolic response to temperature acclimation in nine-spined

stickleback (*Pungitius pungitius*) populations from contrasting thermal environments.

Journal of Experimental Zoology Part A: Ecological Genetics and Physiology

321(10):550-65.

Campbell CS, Bean CW, Parsons KJ. 2017. Conservation evo-devo: preserving biodiversity

by understanding its origins. Trends in ecology & evolution 32(10):746-59.

Chown SL, Gaston KJ. 1999. Exploring links between physiology and ecology at macro-

scales: the role of respiratory metabolism in insects. Biological Reviews 74(1):87-120.

Clark TD, Sandblom E, Jutfelt F. 2013. Aerobic scope measurements of fishes in an era of

climate change: respirometry, relevance and recommendations. Journal of

Experimental Biology 216(15):2771-82.

Clarke A. 1980. A reappraisal of the concept of metabolic cold adaptation in polar marine

invertebrates. Biological journal of the Linnean Society 14(1):77-92.

Clarke A. 1991. What is cold adaptation and how should we measure it? American

Zoologist 31(1):81-92.

Clarke A. 1993. Seasonal acclimatization and latitudinal compensation in metabolism: do

they exist?. Functional Ecology 7(2):139-49.

Clarke A, Morris GJ, Fonseca F, Murray BJ, Acton E, Price HC. 2013. A low temperature

limit for life on Earth. PLoS One. 8(6):e66207.

Conover DO, Schultz ET. 1997. Natural selection and the evolution of growth rate in the

early life history: what are the trade-offs? In: Early life history and recruitment in fish

populations (pp. 305-332). Springer, Dordrecht.

Crozier LG, Hutchings JA. 2014. Plastic and evolutionary responses to climate change in

fish. Evolutionary Applications. 7(1):68-87.

Donelson JM, Munday PL, McCormick MI, Nilsson GE. 2011. Acclimation to predicted

ocean warming through developmental plasticity in a tropical reef fish. Global Change

Biology 17(4):1712-9.

Donelson JM, Munday PL, McCormick MI, Pitcher CR. 2012. Rapid transgenerational

acclimation of a tropical reef fish to climate change. Nature Climate Change 2(1):30.

Dupont-Prinet A, Chatain B, Grima L, Vandeputte M, Claireaux G, McKenzie DJ. 2010.

Physiological mechanisms underlying a trade-off between growth rate and tolerance of

feed deprivation in the European sea bass (*Dicentrarchus labrax*). Journal of

Experimental Biology 213(7):1143-52.

Einarsson A. 1982. The palaeolimnology of Lake Mývatn, northern Iceland: plant and

animal microfossils in the sediment. Freshwater Biology 12(1):63-82.

Einarsson A, Stefánsdóttir G, Jóhannesson H, Olafsson JS, Gíslason GM, Wakana I,

Gudbergsson G, Gardarsson A. 2004. The ecology of Lake Myvatn and the River

Laxá: variation in space and time. Aquatic Ecology 38(2):317-48.

Farrell AP. 2016. Pragmatic perspective on aerobic scope: peaking, plummeting, pejus and

apportioning. Journal of Fish Biology 88:322-343.

Gaitán-Espitia JD, Nespolo R. 2014. Is there metabolic cold adaptation in terrestrial

ectotherms? Exploring latitudinal compensation in the invasive snail *Cornu aspersum*.

Journal of Experimental Biology 217(13):2261-7.

Gaston KJ, Chown SL, Calosi P, Bernardo J, Bilton DT, Clarke A, Clusella-Trullas S,

Ghalambor CK, Konarzewski M, Peck LS, Porter WP. 2009. Macrophysiology: a

conceptual reunification. The American Naturalist 174(5):595-612.

Halsey LG, Killen SS, Clark TD, Norin T. 2018. Exploring key issues of aerobic scope

interpretation in ectotherms: absolute versus factorial. Reviews in Fish Biology and

Fisheries 28(2):405-15.

Hendry AP, Taylor EB. 2004. How much of the variation in adaptive divergence can be

explained by gene flow? An evaluation using lake-stream stickleback pairs. Evolution

58(10):2319-31.

Hight GA. 1965. The saga of Grettir the Strong. London, Dent.

Huey, R. B. and D. Berrigan. 1996. Testing evolutionary hypotheses of acclimation. In I. A.

Johnston and A. E Bennett (eds.), Phenotypic and evolutionary adaptation to

temperature, pp. 205-237. U. Cambridge Press, Cambridge, U.K.

Huss M, Lindmark M, Jacobson P, van Dorst RM, Gårdmark A. 2019. Experimental

evidence of gradual size-dependent shifts in body size and growth of fish in response to

warming. Global Change Biology 25:2285-2295.

Jutfelt F, Norin T, Ern R, Overgaard J, Wang T, McKenzie DJ, Lefevre S, Nilsson GE,

Metcalfe NB, Hickey AJ, Brijs J. 2018. Oxygen-and capacity-limited thermal

tolerance: blurring ecology and physiology. Journal of Experimental Biology

221(1):169615.

Killen SS, Marras S, Ryan MR, Domenici P, McKenzie DJ. 2012. A relationship between

metabolic rate and risk-taking behaviour is revealed during hypoxia in juvenile

European sea bass. Functional Ecology 26(1):134-43.

Killen SS. 2014. Growth trajectory influences temperature preference in fish through an

effect on metabolic rate. Journal of Animal Ecology 83(6):1513-22.

Killen SS, Glazier DS, Rezende EL, Clark TD, Atkinson D, Willener AS, Halsey LG. 2016.

Ecological influences and morphological correlates of resting and maximal metabolic

rates across teleost fish species. The American Naturalist 187(5):592-606.

Killen SS, Norin T, Halsey LG. 2017. Do method and species lifestyle affect measures of

maximum metabolic rate in fishes? Journal of Fish Biology 90:1037-1046.

Krogh A. 1916. The Respiratory Exchange of Animals and Man. London: Longmans,

Green and Co.

Lefevre S. 2016. Are global warming and ocean acidification conspiring against marine

ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high

CO_2 and their interaction. Conservation Physiology 4(1):cow009.

Lenormand T. 2002. Gene flow and the limits to natural selection. Trends in Ecology &

Evolution 17(4):183-9.

Losos JB. 2011. Convergence, adaptation, and constraint. Evolution: International Journal

of Organic Evolution 65(7):1827-40.

Marcil J, Swain DP, Hutchings JA. 2006. Genetic and environmental components of

phenotypic variation in body shape among populations of Atlantic cod (*Gadus morhua*

L.). Biological Journal of the Linnean Society 88(3):351-65.

Nati JJ, Lindström J, Halsey LG, Killen SS. 2016. Is there a trade-off between peak

performance and performance breadth across temperatures for aerobic scope in teleost

fishes? Biology Letters 12:20160191.

Ord TJ, Summers TC. 2015. Repeated evolution and the impact of evolutionary history on

adaptation. BMC Evolutionary Biology 15(1):137.

O'Gorman EJ, Ólafsson ÓP, Demars BO, Friberg N, Guðbergsson G, Hannesdóttir ER,

Jackson MC, Johansson LS, McLaughlin ÓB, Ólafsson JS, Woodward G. 2016.

Temperature effects on fish production across a natural thermal gradient. Global

Change Biology 22:3206-20.

Paaby AB, Rockman MV. 2014. Cryptic genetic variation: evolution's hidden substrate.

Nature Reviews Genetics 15(4):247.

Pettersen AK, Marshall DJ, White CR. 2018. Understanding variation in metabolic rate.

Journal of Experimental Biology 221(1):jeb166876.

Pilakouta N, Humble J, Hill I, Killen SS, Kristjánsson BK, Skúlason S, Lindström J,

Metcalfe NB, Parsons KJ. 2019. Testing the predictability of morphological evolution

in naturally warmed stickleback populations. *BioRxiv* [Preprint]. Available at

https://doi.org/10.1101/609933.

Pinho C, Hey J. 2010. Divergence with gene flow: models and data. Annual Review of

Ecology, Evolution, and Systematics 41:215-30.

Pörtner HO, Farrell AP. 2008. Physiology and climate change. Science 322(5902):690-2.

R Core Team (2017). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL: http://www.R-

project.org/

Sandblom E, Clark TD, Gräns A, Ekström A, Brijs J, Sundström LF, Odelström A, Adill

A, Aho T, Jutfelt F. 2016. Physiological constraints to climate warming in fish follow

principles of plastic floors and concrete ceilings. Nature Communications 7:11447.

Shama LNS. 2017. The mean and variance of climate change in the oceans: hidden

evolutionary potential under stochastic environmental variability in marine sticklebacks.

Scientific Reports 7:8889.

Sinclair BJ, Marshall KE, Sewell MA, Levesque DL, Willett CS, Slotsbo S, Dong Y, Harley

CD, Marshall DJ, Helmuth BS, Huey RB. 2016. Can we predict ectotherm responses

to climate change using thermal performance curves and body temperatures? Ecology

Letters 19(11):1372-85.

Tsuji JS. 1988. Thermal acclimation of metabolism in Sceloporus lizards from different

latitudes. Physiological Zoology 61(3):241-53.

White CR, Kearney MR. 2013. Determinants of inter-specific variation in basal metabolic

rate. Journal of Comparative Physiology B 183(1):1-26.

White CR, Frappell PB, Chown SL. 2012. An information-theoretic approach to evaluating

the size and temperature dependence of metabolic rate. Proceedings of the Royal

Society B: Biological Sciences 279:3616-21.

Wickham H. 2009. ggplot2: elegant graphics for data analysis. Springer New York.

Zuo W, Moses ME, West GB, Hou C, Brown JH. 2012. A general model for effects of

temperature on ectotherm ontogenetic growth and development. Proceedings of the

Royal Society B: Biological Sciences 279:1840-1846.

Table 1. Sampling locations of warm- and cold-habitat threespine sticklebacks collected in May–June 2016 in Iceland. Distance refers to how far apart the warm-habitat and cold-habitat sample sites are for each warm-cold pair. All cold habitats have existed since the last glacial period and are therefore approximately 10,000 years old, whereas warm habitats can be classified as either young (<80 years old) or old (>2,000 years old). The summer and winter temperatures listed are the average water temperatures recorded at each sampling location during the corresponding seasons.

Population pair	Water body	Thermal habitat	Age of warm habitat (years)	Distance (km)	Summer temperature (ºC)	Winter temperature (ºC)
allopatric populations	Grettislaug	warm	>2,000	21.04	24.9	13.5
	Garðsvatn	cold			14.6	2.2
sympatric population 1	Áshildarholtsvatn	warm	50–70	0.05	24.1	12.5
		cold			12.2	3.4
sympatric population 2	Mývatn	warm	>2,000	3.18	22.8	22.0
		cold			11.5	1.0

Table 2. Results of general linear models testing the effects of thermal habitat (warm or cold), population pair (allopatric, sympatric 1, or sympatric 2), acclimation temperature (10ºC, 15ºC, or 20ºC), and their interactions on SMR, MMR, and AS in threespine stickleback from six populations in Iceland. Df denotes degrees of freedom. Eta-squared (η^2) represents the percent variance explained by each factor, which was calculated by dividing the sum of squares for each factor by the total sum of squares and multiplying by 100.

	Standard metabolic rate (SMR)				Maximum metabolic rate (MMR)				Aerobic scope (AS)			
	η^2	df	*F*	*P*	η^2	Df	*F*	*P*	η^2	df	*F*	*P*
Thermal habitat	0.82	1	9.77	**0.002**	1.38	1	10.4	**0.001**	0.99	1	6.26	**0.013**
Population pair	1.38	2	8.25	**<0.001**	0.81	2	3.27	**0.039**	2.20	2	7.09	**<0.001**
Acclimation temperature	31.7	2	192	**<0.001**	6.92	2	26.9	**<0.001**	2.97	2	9.56	**<0.001**
Mass	29.3	1	355	**<0.001**	35.9	1	281	**<0.001**	28.6	1	186	**<0.001**
Thermal habitat × Population pair	0.94	2	5.65	**0.004**	0.35	2	1.56	0.21	0.22	2	0.89	0.41
Thermal habitat × Acclimation temperature	0.03	2	0.14	0.87	1.15	2	4.29	**0.014**	1.43	2	4.74	**0.009**
Population pair × Acclimation temperature	3.32	2	10.0	**<0.001**	0.46	2	0.98	0.42	1.21	2	2.05	0.087
Thermal habitat × Population pair × Acclimation temperature	0.82	4	2.49	**0.043**	1.15	4	2.25	0.063	2.97	4	4.79	**<0.001**
Error	31.7	385			51.9	406			59.4	385		

Figure 1. Map of Iceland showing the sampling locations of warm- and cold-habitat sticklebacks we collected for this study. Each of the three population pairs is indicated by a different colour.

Figure 1. Metabolic rate (mg O$_2$ hr^{-1}) of threespine sticklebacks from cold and warm habitats in Iceland that were acclimated to 10°C, 15°C, or 20°C. We show mass-corrected standard metabolic rate (A), maximum metabolic rate (B), and aerobic scope

(C). Error bars indicate standard errors, and small circles represent individual data points (blue=cold thermal habitat, red=warm thermal habitat). 'Allopatric' refers to Grettislaug and Garðsvatn, 'sympatric 1' refers to Áshildarholtsvatn, and 'sympatric 2' refers to Mývatn.