Contrasting signatures of introgression in North American box turtle (*Terrapene* spp.)

contact zones

Running title: Differential introgression among hybrid zones

Authors:

Bradley T. Martin[1,6,*], Marlis R. Douglas[1], Tyler K. Chafin[1], John S. Placyk, Jr.[2,5], Roger

D. Birkhead[3], Christopher A. Phillips[4], Michael E. Douglas[1]

[1]Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas 72701, USA;

Email: (BTM) btm002@uark.edu; (MRD): mrd1@uark.edu; (TKC): tkchafin@uark.edu; (MED):

med1@uark.edu.

[2]Department of Biology, University of Texas, Tyler, Texas, 75799, USA; Email:

japlacyk@gmail.com

[3]Alabama Science in Motion, Auburn University, Auburn, AL 36849, USA; Email:

birkhrd@auburn.edu

[4]Illinois Natural History Survey, Prairie Research Institute, University of Illinois, Champaign, IL

61820; Email: caphilli@illinois.edu

[5]Science Division, Trinity Valley Community College, Athens, TX 75751, USA.

Corresponding author:

[6,*]Global Campus, University of Arkansas, 2 E. Center St., Fayetteville, Arkansas 72701, USA;

Email: btm002@uark.edu

Disclosure statement: Authors have nothing to disclose

ABSTRACT

Hybridization occurs differentially across the genome in a balancing act between selection and migration. With the unprecedented resolution of contemporary sequencing technologies, selection and migration can now be effectively quantified such that researchers can identify genetic elements involved in introgression. Furthermore, genomic patterns can now be associated with ecologically relevant phenotypes, given availability of annotated reference genomes. We do so in North American box turtles (*Terrapene*) by deciphering how selection affects hybrid zones at the interface of species boundaries and identifying genetic regions potentially under selection that may relate to thermal adaptations. Such genes may impact physiological pathways involved in temperature-dependent sex determination, immune system functioning, and hypoxia tolerance. We contrasted these patterns across inter- and intra-specific hybrid zones that differ temporally and biogeographically. We demonstrate hybridization is broadly apparent in *Terrapene*, but with observed genomic cline patterns corresponding to species boundaries at loci potentially associated with thermal adaptation. These loci display signatures of directional introgression within intra-specific boundaries, despite a genome-wide selective trend against intergrades. In contrast, outlier loci for inter-specific comparisons exhibited evidence of being under selection against hybrids. Importantly, adaptations coinciding with species boundaries in *Terrapene* overlap with climatic boundaries and highlight the vulnerability of these terrestrial ectotherms to anthropogenic pressures.

Keywords: adaptive divergence; hybrid zone; ddRAD; temperature dependence; selection; genomic cline

1. INTRODUCTION

Hybrid zones are natural laboratories that allow the genetic architecture of local adaptation and/or reproductive isolation to be examined. They frequently juxtapose with underlying ecological gradients, allowing researchers to quantify the how selection impacts the genome (Barton & Hewitt 1985; Payseur 2010). Here, selection might prevent introgression at loci underpinning crucial adaptations while the rest of the genome essentially homogenizes (Via 2009; Feder *et al.* 2013). Once such loci are identified, the phenotypes inferred by adaptive divergence can then be inferred via a bottom-up, "reverse-ecology" approach (Li *et al.* 2008; Tiffin & Ross-Ibarra 2014). Hybrid zones effectively become unique "windows" into the speciation process by allowing functional loci to be associated with various aspects of ecology (Taylor *et al.* 2015).

Diminishing costs associated with genomic sequencing, coupled with an upsurge in genomic annotations, has facilitated the reverse-ecology approach. Examples include adaptive divergence in seasonal growth and variability in immune responses (Rödin⬚Mörch *et al.* 2019), proactive responses to environmental gradients (Keller & Seehausen 2012; Guo *et al.* 2016; Waterhouse *et al.* 2018; Teske *et al.* 2019), and an upsurge in contemporary effects such as anthropogenic modulation of reproductive boundaries (Garroway *et al.* 2010; Taylor *et al.* 2014; Grabenstein & Taylor 2018). Importantly, researchers can now effectively gauge the how the genome is impacted by ecological and climatic shifts, and the how species distributions are promoted by pre-existing adaptive gradients (Rosenzweig *et al.* 2008; Taylor *et al.* 2015; Ryan *et al.* 2018). Thus, genome-scale datasets can often refine our perspectives on two major evolutionary patterns: First, reproductive boundaries among historically co-existing species can

become blurred due to substantial environmental change, a situation directly analogous to the

imposition of contact among otherwise allopatric taxa (Rhymer & Simberloff 1996). Second,

selection and migration are balanced within hybrid zones (Key 1968; Parmesan *et al.* 1999), and

this balance can shift due to rapid and/or exceptional climate change (Seehausen *et al.* 2008;

Kearns *et al.* 2018). Genomic data extracted from hybrid zones may thus allow species

boundaries to be defined according to their phenotypic and genetic underpinnings.

Species boundaries can either be strengthened (Ryan *et al.* 2018) or eroded (Muhlfeld *et*

al. 2014) due to temperature shifts, a major component of climate change, potentially shifting

species distributions and/or hybrid zone dynamics. Furthermore the effects of temperature on

physiological and cellular mechanisms are well known (Kingsolver 2009), directly affecting

growth, development, reproduction, locomotion, and immune response (Keller & Seehausen

2012). As a result, the manner by which thermal gradients interact with species boundaries has

become a major focus (Qin *et al.* 2013). Herein, we attempt to clarify how species boundaries

reflect environmental processes by quantifying the geographic and ecological foundations of two

hybrid zones in the ectothermic North American box turtles (*Terrapene*).

1.1. *Hybridization in North American box turtles*

North American box turtles (Emydidae, *Terrapene*) are long-lived, omnivorous, and primarily

terrestrial ectotherms, with a rectangular appearance defined by a dome-shaped dorsal carapace

and a ventral plastron hinged to tightly close against the carapace (hence the common name)

(Dodd 2001). Their North American range is characterized by two well-known zones of

hybridization (Milstead 1969; Dodd 2001; Cureton *et al.* 2011) that provide excellent models

from which to contrast regional patterns of hybridization and introgression. To do so, we

evaluated four southeastern taxa [the Woodland (*T. carolina carolina), *Gulf Coast (*T. c. major*),

Three-toed (*T. carolina triunguis*), and Florida (*T. bauri*) box turtles (Auffenberg 1958, 1959;

Milstead & Tinkle 1967; Milstead 1969; Martin *et al.* 2013; Iverson *et al.* 2017)], and two

midwestern (the Ornate box turtle, *T. ornata ornata* and *T. c. carolina*; Cureton *et al.* 2011). Each

hybridizes regionally, and therefore we focus on inter- and intra-specific contacts within these

two regions.

One focal hybrid zone is nested within southeastern North America (Ricketts 1999), where

box turtles inhabit a biodiversity hotspot. Here, clear-cutting, invasive species, and altered fire

regimes are widespread (Stapanian *et al.* 1997, 1998; van Lear & Harlow 2002), and impact

numerous endemic species (Lydeard & Mayden 1995). The region also displays clinal

intergradation (i.e., interbreeding between subspecies), as well as hybridization across a variety of

taxa (Remington 1968; Swenson & Howard 2004), due largely to coincident ecological and

climatic transitions (Swenson & Howard 2005).

By contrast, contact zones in midwestern North America seemingly stem from secondary

contact (i.e., resumption of interbreeding following a geographic separation), as associated with

postglacial recolonization/ expansion (Swenson & Howard 2005). Here, prairie-grassland habitat

has been anthropogenically fragmented such that niche overlap now occurs between grassland

and woodland species (Johnson 1994; Samson & Knopf 1994; Rhymer & Simberloff 1996;

Samson *et al.* 2004). Furthermore, while overlapping forms in the Midwest represent distinct

species, southeastern forms are taxonomically in flux. Species status has varied for *T. m.*

triunguis, and *T. c. major* is now viewed as an intergrade population (Butler *et al.* 2011; Iverson

et al. 2017), despite recent molecular work suggesting specific status for *T. m. triunguis* and

phylogenetic structure in *T. c. major* (Martin *et al.* 2013, 2014, 2020). For the sake of clarity, we

follow Martin *et al.*, regarding *T. m. triunguis* as a full species and *T. c. major* as a recognized

subspecies.

We used ddRAD sequencing (Peterson *et al.* 2012) to contrast genome-wide patterns of

clinal introgression within each hybrid zone. We then identified/ quantified loci potentially under

selection by mapping them to an available genomic reference and accordingly interpreted their

potential ecological associations, which provide invaluable insights into how *Terrapene* has

responded to a fluctuating climate. As such, our results extend to a proactive management

paradigm that underscores the conservation of co-occurring forms.

2. MATERIALS AND METHODS

2.1. *Tissue and DNA collection*

Tissues for *T. carolina*, *T. ornata*, and *T. mexicana triunguis* were collected by volunteers

and agency collaborators (Table S1). Additional samples were provided by numerous museums

and organizations. Live animals were sampled non-invasively (e.g., blood, toenails, or toe-clips),

whereas road-kills were sampled indiscriminately. Isolation of genomic DNA was performed

using DNeasy Blood and Tissue Kits (QIAGEN), QIAamp Fast DNA Tissue Kit (QIAGEN), and

E.Z.N.A. Tissue DNA Kits (Omega Bio-tek). The presence of genomic DNA was confirmed via

gel electrophoresis using a 2% agarose gel.

2.2. *Library preparation*

In silico digests were carried out to optimize restriction enzyme selection, using available

genomic references [Painted turtle (*Chrysemys picta*), GenBank Accession #: GCA_000241765.2

(Shaffer *et al.* 2013); FRAGMATIC (Chafin *et al.* 2018); genome size=2.59 X 10^9 bp]. The

distribution of fragments (from N=24 samples) were first optimized then evaluated using an

Agilent 4200 TapeStation. Library preparation was conducted per standard protocol (Peterson *et*

al. 2012), using *PstI* (5'-CTGCAG-3') and *MspI* (5'-CCGG-3') restriction enzymes. We digested

~500-1,000ng of DNA/ sample at 37°C, with unique DNA barcode and sequencing adapters

subsequently ligated. Prior to sequencing, quality control checks were performed at the core

facility, to include fragment analysis for confirmation of correct size range and quantitative real-

time PCR. Individuals (N=96) were pooled per lane of single-end Illumina sequencing at the

University of Oregon Genomics and Cell Characterization Core Facility (Hi-Seq 4000, 1x100bp).

Populations were randomized across multiple lanes to mitigate batch effects.

2.3. *Assembly and quality control*

Read quality was quantified using FASTQC v. 0.11.5, then demultiplexed and aligned

using IPYRAD v. 0.7.28 (Eaton & Overcast 2020), with reads mapped to the scaffold-level *T.*

mexicana triunguis reference assembly (GenBank Accession #: GCA_002925995.2) at a distance

threshold of 0.15. Non-mapping reads were discarded. This alignment is herein termed the

"scaffold alignment" to differentiate it from a separate transcriptome-mapped alignment (see

section 2.6). Barcodes and adapters were trimmed, as were the last five base pair (bp) of each

read. Those exceeding five bases with low PHRED quality score (<33) were discarded, and

potential paralogs were filtered by excluding loci with high heterozygosity ($>75\%$) or >2 alleles

per individual. Loci with a sequencing depth of $<20X$ per individual or $<50\%$ presence across

individuals were also discarded. Our mapping and filtering steps above were conducted in

IPYRAD.

2.4. *Assessing Admixture and Population Structure*

ADMIXTURE (Alexander *et al.* 2009) was used to assess contemporary hybridization. It

employs a model-based ML approach that estimates the proportion of ancestry shared across the

genome-wide average of each sample. $K=1\text{-}13$ were used for datasets containing all sequenced

taxa and subsets from the Southeast and Midwest hybrid zones, with 20 independent replicates

per K (ADMIXPIPE; Mussmann *et al.* 2020). Hierarchical partitioning was done because

ADMIXTURE often underestimates K by detecting only the uppermost hierarchy of population

structure (Evanno *et al.* 2005). SNP data were pre-filtered using VCFTOOLS (Danecek *et al.*

2011), with SNPs randomly thinned to one per locus to alleviate linkage bias and filtered by

removing sites with a minor allele frequency (MAF)$<1.0\%$ to reduce bias associated with

erroneous genotypes and singletons (Linck & Battey 2019). Model support across K-values was

assessed using five-fold cross-validation (Alexander *et al.* 2009). ADMIXPIPE output was

summarized using the CLUMPAK server (Kopelman *et al.* 2015), with each individual

subsequently plotted as a stacked bar-chart (Rosenberg 2004).

We also performed Discriminate Analysis of Principal Components (DAPC) using the

adegenet R-package (v2.0-0) with identical filtering parameters (Jombart *et al.* 2010). The

find.clusters() function was utilized with 1,000,000 iterations to determine the optimal K with the

lowest Bayesian Information Criterion (BIC). DAPC cross-validation (100 replicates, 90%

training dataset) then evaluated which principle components and discriminant functions to retain,

with individuals plotted against the top three DAPC axes.

Finally, we also ran TESS3 (TESS3R R package; Caye *et al.* 2016) to estimate ancestry

coefficients (as with ADMIXTURE), but also incorporate spatial proximity into ancestral genotype

estimates. The TESS3 input alignments were subsets of those used in ADMIXTURE to include only

individuals with GPS coordinates. Cross-validation (with 10% sites randomly masked) was

performed for K=1-10 with twenty independent runs to assess optimal K. The output Q-matrix

was interpolated using spatial kriging (Jay *et al.* 2012).

2.5. *Identifying Hybrids*

NEWHYBRIDS (Anderson & Thompson 2002) was used to assign statistically-supported

hybrids to genotype frequency classes (i.e., Pure, F_1, F_2, and backcrosses between F_1 and parental

types). The *getTopLoc()* function in HYBRIDDETECTIVE (Wringe *et al.* 2017a) reduced the data to

300 loci containing the highest among-population differentiation (F_{ST}) and lowest linkage

disequilibrium correlation (r^2<0.2). Burn-in was 500,000 MCMC generations followed by

2,000,000 post burn-in sweeps. Seeds were randomized and the analysis employed the Jeffrey's

prior for θ and π. To train the data, individuals sampled outside the focal hybrid zones with

ADMIXTURE proportions=100% were pre-assigned as parentals. The following combinations of

taxa were employed: *Terrapene carolina carolina* X *T. c. major*, *T. c. carolina* X *T. m. triunguis*,

T. c. major X *T. m. triunguis*, and *T. c. carolina* X *T. o. ornata*. A posterior probability threshold

>0.8 was required for assignment into the genotype frequency classes, as determined using a

power analysis conducted with HYBRIDDETECTIVE and PARALLELNEWHYBRID pipelines (Wringe

et al. 2017b; a).

2.6. *Genomic Clines among Scaffolds and mRNA Mapping*

In addition to the "scaffold alignment," IPYRAD was rerun with reads mapped to the *T. m.*

triunguis reference transcriptome (GenBank Accession: GCA_002925995.2), with identical

filtering parameters. Three subsets of the resulting "transcriptomic alignment" were generated to

retain only individuals per each pairwise combination of southeastern taxa (*T. c. carolina*, *T. c.*

major, *T. m. triunguis*). The "scaffold" and "transcriptome" alignments were then independently

examined for patterns of differential introgression using INTROGRESS (Gompert & Buerkle 2010)

and Bayesian Genomic Clines (Gompert & Buerkle 2012). Both generate genomic clines, which

assess locus-specific ancestry to identify outliers *versus* the genome-wide average and can

identify outlier SNPs having cline shapes divergent from neutral expectations. Parental reference

populations were determined *a priori* via population structure and NEWHYBRIDS results. Parental

status was considered only for samples with ADMIXTURE ancestry coefficients=100%.

INTROGRESS derived neutral expectations from 1,000 parametric simulations (Gompert &

Buerkle 2010). Genomic clines were only generated for SNPs with a high allele frequency

differential (δ) between parental types (Andrés *et al.* 2013). Outlier SNPs were defined using a

Bonferroni-corrected α-significance threshold.

Prior to running BGC, sites with a MAF<5% were removed because an over-abundance

of uninformative loci inhibited parameter convergence. Five independent BGC runs were

conducted for both scaffold and transcriptomic alignments, which included 1,000,000 and

1,800,000 burn-in, respectively, each with 200,000 post-burn-in generations. Samples were

thinned with every 50 iterations retained to mitigate auto-correlation. Genotype uncertainty

corrections were applied to each locus, with the sequencing error rate prior computed in IPYRAD

(ranging from 0.1-0.2%). The BGC linkage model was tested but found to be computationally

intractable (i.e., >1Tb memory and unreasonable running times). Upon run completion, parameter

traces were visually inspected for convergence. Replicate runs were subsequently combined.

Two parameters [i.e., genomic cline center (α) and rate (ß)] represented BGC output

(Gompert & Buerkle 2011). The α-parameter indicates the direction of introgression, with

negative and positive outliers depicting excess P_1 and P_2 ancestry, respectively. ß characterizes

the rate of change, with negative values reflecting a wider genomic cline where loci more freely

introgress, and a steeper cline indicated by positive values with relatively sharp transitions from

P_1 to P_2 ancestry. BGC outliers were considered significant if they met either of two criteria: 1)

the 95% credible intervals for α or ß did not overlap zero, or 2) the median of the posterior

distribution exceeded the probability distribution's quantile interval $[(\frac{1-0.975}{2}, \frac{0.975}{2})$; (Gompert

& Buerkle 2011)].

2.7 *Mapping BGC Outliers to Chromosomes*

BGC parameters α and ß were plotted onto assembled chromosomes to visualize the

distribution of outliers across the genome. The *Terrapene* reference is a scaffold-level assembly,

so MINIMAP2 (Li 2018) and PAFSCAFF (https://github.com/slimsuite/pafscaff) were used to map

the *T. m. triunguis* reference to a closely related chromosome-level assembly (*Trachemys scripta*;

Simison *et al.* 2020; GenBank accession: GCA_013100865.1). The ASM20 MINIMAP2 preset was

chosen to accommodate the ~5-10% sequence divergence expected between *Trachemys* and

Terrapene (Feldman & Parham 2002). The mapping connected *Terrapene* BGC outliers with

Trachemys chromosomal positions, allowing the putative BGC outlier locations to be visualized

at the chromosome-level, with plots subsequently generated (RIDEOGRAM R package; Hao *et al.*

2020).

2.8 *Correlating Outliers with Environmental Variables*

We also independently identified SNP outliers by environmental association using

redundancy analysis (RDA). Outlier SNPs were correlated with the standard WorldClim v2

Bioclimatic variables that included 19 raster layers of temperature and precipitation, plus mean

annual solar radiation, mean annual wind speed, and elevation (Fick & Hijmans 2017). The finest

available scale (30 arc-seconds) was chosen for each raster. Layers were projected to WGS84 and

cropped to the sampling extent (RASTER R package; Hijmans & Van Etten 2016). Raster values at

each sampling location were then extracted.

Each predictor variable was scaled, centered, and tested for normality with a Shapiro-

Wilks test. Non-normal distributions were transformed using the BESTNORMALIZE R package

(Peterson & Cavanaugh 2019) per RDA's assumptions. The environmental layers were assayed

for predictive capabilities with uncorrelated variables retained (ADESPATIAL R package, forward

selection with 10,000 permutations). Predictors that failed forward selection were removed. To

account for underlying spatial influence, distance-based Moran's eigenvector maps (dbMEM)

were generated using sample coordinates (*quickMEM()* R function; Borcard *et al.* 2018). The

dbMEMs are a matrix of axes that capture spatial patterns from multiple angles rather than just a

latitudinal or longitudinal vector. To reduce overfitting, informative, non-redundant dbMEM axes

were subset, using forward selection with 1,000 permutations. Finally, the SNP matrix was

imported into R using ADEGENET, and missing data were imputed as the most frequent allele per

population, following RDA assumptions.

A partial RDA (pRDA) conditioned on the dbMEM spatial matrix was then conducted using

genotypes as the response variables, with 1,000 permutations (VEGAN R package; Oksanen *et al.*

2019). This approach "partialed out" spatial autocorrelative effects that could yield false negative

SNP-environment associations. Significant RDA axes were determined using VEGAN's

anova.cca() function, and SNPs with loadings +/- 3 standard deviations from the mean on a

significant axis were considered outliers. A full RDA and a pRDA conditioned on environment

were also conducted to estimate the contributions of spatial *versus* environmental predictors.

Each SNP was then correlated pairwise with all environmental variables using Pearson's

correlation coefficient (*r*), and those with the strongest correlations represented the best supported

SNP-environment association.

3. RESULTS

A total of 368 individuals (Tables S1, S2) were retained across 12,052 (combined alignment),

10,338 (Midwest-only), and 11,308 (Southeast-only) unlinked reference-mapped loci. This, a

result of quality control steps and post-alignment filters that eliminated individuals with >90%

missing data, and sites with MAF<1.0%. The scaffold alignment included 134,607 variable and

90,777 parsimoniously informative sites. The transcriptome-guided alignment contained 2,741 bi-

allelic SNPs across 247 individuals, with subsets generated for *T. c. carolina* X *T. c. major*

(EAxGU), *T. c. carolina* X *T. m. triunguis* (EAxTT), and *T. c. major* X *T. m. triunguis* (GUxTT).

3.1. *ADMIXTURE across the hybrid zones*

The combined east/ west ADMIXTURE CV (Fig. S1) supported *K*=6 (x⎕= 0.19279, SD =

0.00017), followed by *K*=4 (x⎕= 0.19490, SD = 0.0016) and *K*=5 (x⎕= 0.19738, SD = 0.00013).

The analysis indicated population structure for *T. c. carolina*, *T. m. triunguis*, two distinct *T. c.*

major subpopulations (Alabama/ Mississippi and Florida panhandles), and northern and southern

T. o. ornata subpopulations from Illinois+Wisconsin+Iowa and

Kansas+Texas+Colorado+Nebraska (Fig. S2). *Terrapene bauri* was excluded due to limited

sampling (N=4). Admixture occurred between *T. c. carolina* X *T. c. major* (EAxGU), *T. c.*

carolina X *T. m. triunguis* (EAxTT), *T. c. major* X *T. m. triunguis* (GUxTT), *T. o. ornata* X *T. c.*

carolina (EAxON), and the two *T. o. ornata* subpopulations.

The lowest CV score for the southeastern ADMIXTURE (N=259 individuals) was at K=4

($\bar{x}$= 0.21851, SD=0.00016), with K=3 ($\bar{x}$=0.22134, SD=0.00015) and K=5 ($\bar{x}$= 0.22519,

SD=0.00082) trailing (Fig. S3). Southeastern taxa included *T. c. carolina*, *T. c. major*, and *T. m.*

triunguis, and their analysis concurred with the all-taxa dataset in terms of both population

structure and admixed taxa (Figs. 1, S4). Admixture primarily occurred throughout Alabama and

the Florida panhandle (EAxGU), Georgia and South Carolina (EAxTT), and Mississippi/

southern Alabama (GUxTT). The two *T. c. major* subpopulations in the all-taxa analysis were

corroborated. The same four southeastern groups plus *T. bauri* were also produced by DAPC

(K=5; Fig. S5). We found *T. bauri* highly differentiated along axis 1 (71.9% variance explained),

whereas axes 2-3 delineated the remaining southeastern taxa (17.5% and 5.73%, respectively).

The midwestern analysis (Figs. 1, S6) included an optimal K=2 ($\bar{x}$= 0.24069, SD=

0.00023), followed by K=3 ($\bar{x}$=0.25703, SD=0.00454) and K=4 ($\bar{x}$=0.25861, SD=0.00374)

(Fig. S7). The K=2 groups consisted of *T. c. carolina* and *T. o. ornata*, with only a few

individuals indicating admixture. At K=3, *T. c. carolina* from Illinois split as a distinct group,

although only a few of the admixture proportions approached 100%. At K=4, the northern and

southern *T. o. ornata* subpopulations produced by the all-taxa analysis also materialized.

TESS3 corroborated both midwestern and southeastern ADMIXTURE analyses, with *T. o.*

ornata, *T. c. carolina T. m. triunguis*, and the two *T. c. major* subpopulations in Alabama/

Mississippi and Florida being delineated (Fig. 1). The Kriging interpolation also spatially

highlighted ancestry gradients consistent with ADMIXTURE (FIG. S8), with lower surface

prediction scores concordant with areas that contained frequently mixed ancestry.

3.2. *Genealogical hybrid classification*

HYBRIDDETECTIVE confirmed convergence for inter- and intra-simulation replicates (Fig.

S9) with 500,000 burn-in and 2,000,000 post-burn-in sweeps (the EAxON analysis required

4,000,000 sweeps with 1,000,000 burn-in). Our power analyses suggested 90% assignment

accuracy (+/- SD) for all genotype classes at a critical threshold of 0.8 (Figs. S10, S12, S14, S16).

Statistical power was also elevated ($\geq$0.8), although some genotype classes for EAxGU displayed

relatively lower power (<0.8) (Figs. S11, S13, S15, S17).

The southeastern hybrid zone consisted entirely of backcrosses (F_1 hybrids X parental

types), F_2-generation, and unassigned (>F_2) hybrids (Fig. 2, Table S3). Specifically, the EAxGU

analysis identified backcrosses with parental *T. c. major* and F_2 hybrids in the Florida panhandle

and southern Alabama (Fig. 2A). Similarly, all hybrid-generation EAxTT individuals from

Georgia were identified as backcrosses with both parental types, whereas South Carolina hybrids

were backcrosses with *T. c. carolina* (Fig. 2B).

Second-generation hybrids and backcrosses with both parental types were evident among

GUxTT (Fig. 2C). Mississippi contained individuals with all three hybrid genotype classes (F_2,

B_1, and B_2), but with backcrosses to parental *T. m. triunguis* at the greatest frequency. Alabama

and Florida GUxTT were only represented by *T. m. triunguis* backcrosses. Finally, *T. o. ornata*

and *T. c. carolina* in Illinois displayed relatively few hybrid genotypes (5%) but all were F_1, in

contrast to the southeastern analyses (Fig. 2D).

3.3. *Selective signatures at transcriptomic loci*

Using INTROGRESS, only SNPs with a high allelic frequency differential ($\delta>0.8$) were

retained (Andrés *et al.* 2013). One exception was the EAxGU analysis where $\delta>0.7$ was applied

because no loci at the higher threshold were recovered.

The INTROGRESS genomic cline analysis recovered three outlier mRNA loci for EAxGU,

and five each for EAxTT and GUxTT, with thirteen total and nine unique outliers among the

three pairwise comparisons (Table 1; Fig. 3). Clines were inconsistent in that some pairwise

taxon comparisons displayed rapid transitions from P_1 to P_2, whereas others demonstrated

patterns indicative of directional introgression.

For example, all three EAxGU outlier loci displayed an over-representation of EA alleles

in the hybrid zone, concomitant with an under-representation of heterozygotes and GU alleles

(Fig. 3A). The *SULT* locus was an extreme example, with excess EA alleles below a hybrid index

of ~0.8 (=80% assignment to GU at diagnostic loci). This pattern was replicated to a lesser degree

in *ZNF236*, whereas *TLR9* was more sigmoidal. These genotypic proportions, coupled with the

non-sigmoidal cline shape in *SULT* and *ZNF236*, suggest that introgression may be driven by a

directional shift towards homozygous P_1 genotypes. In contrast, a steep, sigmoidal cline

represented the genomic trend among scaffold assembly (and putatively non-functional) loci (Fig.

18). Taken together, these results suggest underlying directional introgression facilitating

exchange of EA alleles despite divergence being maintained at most loci.

Cline shape was inconsistent within the EAxTT hybrid zone (Fig. 3B). Three (of five)

outlier loci (*SASH3*, *SYPL2*, and *TLR9*) were significantly under-represented with regards to

heterozygotes. Their clines displayed steep slopes, suggesting rapid transition among parental

genotypes. An additional locus (*CITED4*) reflected an overrepresentation of P_2 (TT), and a fifth

(*FAM89B*) displayed three equally-represented genotypes. Of note, *FAM89B* did not differ

significantly from neutral expectations following Bonferroni correction (*P*=0.036).

By contrast, neutral expectations were rejected in all five GUxTT clines (*P*=0; α=0.007),

with four (i.e., *SASH3*, *SYPL2*, *ACAD11, FAM89B*) suggesting a pattern of restricted

introgression (Fig. 3C). A fifth (i.e., *TMEM214*) displayed directional introgression, with the

homozygous P_2 (TT) genotype being overrepresented. Both the GUxTT and EAxTT analyses

showed a ubiquitous signal of steep clines in non-transcriptomic loci (Fig. S18).

A greater number of SNP outliers (N=81) were identified with BGC rather than

INTROGRESS, likely reflecting the larger number of loci included in the former. All INTROGRESS

outliers were also corroborated by BGC (Fig. 4; Table 1). For EAxGU (Fig. 4A), all three

exhibited excess EA ancestry (negative α=excess P_1; positive α=excess P_2), but there were no

cline rate (ß) outliers in either direction. In contrast, four out of five EAxTT loci (*SYPL2*, *SASH3*,

TLR9, and *CITED4*) were positive ß outliers, indicating steep clines and thus restricted

introgression, with *SYPL2* also being an α outlier with excess EA (P_1) ancestry (Fig. 4B). The

fifth locus (*FAM89B*) was an α (not ß) outlier that favored EA alleles. Finally, GUxTT (Fig. 4C)

included two loci that were both α and ß outliers with steep clines and excess GU ancestry

(*SYPL2* and *FAM89B*). Two others were ß-only outliers with steep clines (*SASH3* and *ACAD11*),

and one (*TMEM214*) an α-only outlier with excess TT ancestry.

3.4 Environmental correlations with outliers

Shapiro-Wilks tests confirmed normality for all layers, following *OrderNorm* transformation.

Forward selection retained ten uncorrelated and predictive layers (Table 2), and pRDAs revealed

49.2% environmental and 40.9% spatial contributions to model inertia. After controlling for

spatial autocorrelation, the pRDA ANOVA identified four significant axes ($P<0.05$) explaining

21.5%, 10.6%, 10.3%, and 9.6% of the variance (cumulative 52.1%). The pRDA also identified

56 annotated outlier SNPs correlated with environmental predictors (Fig. 5). Twenty-eight pRDA

outliers overlapped with INTROGRESS and/or BGC analyses (Fig. 6). Many of the overlapping

SNPs, including all nine INTROGRESS loci that remained as outliers across all three analyses, were

most strongly correlated with temperature variables rather than precipitation or wind speed (Table

3).

4. DISCUSSION

Our analyses characterized introgression in two North American box turtle hybrid zones

(i.e., midwestern and southeastern North America). The midwestern hybrid zone showed no

evidence of introgression, with hybrids restricted at low frequency to F_1, whereas southeastern

hybridization was introgressive in nature, as evidenced by numerous backcrosses and F_2

individuals and a conspicuous lack of F_1 hybrids. Furthermore, contrasting intra- and inter-

specific southeastern hybrid zones revealed they not only varied in the genealogical composition

of their hybrids but also in the shapes and widths of locus-specific clines. We propose such a

contrast provides insight into the evolutionary histories of the taxa involved and serves to

delineate their appropriate taxonomic designations. Specifically, recent phylogenetic research

indicates that *T. m. triunguis* is a separate species from *T. c. carolina* and *T. c. major* (Martin *et*

al. 2013, 2014, 2020). The genomic clines herein are consistent with those phylogenetic results in

that transcriptomic loci with steep clines are only found in inter-specific comparisons (i.e., *T.*

mexicana versus T. carolina). The candidate genes at these transcriptomic loci may be targets of

selection and directional introgression, as they deviated from neutral expectations not only with

respect to genome-wide ancestry (i.e., genomic clines) but also multivariate environmental

associations. Below we consider the impact of these results on the evolutionary history, genomic

architecture, and species boundaries of *Terrapene*.

4.1.*Regional and Taxon-specific Perspectives*

Several of our conclusions have implications for *Terrapene* systematics. First, our

ADMIXTURE analyses substantiate the presence of discrete *T. c. major* populations in Florida and

Mississippi, with the Alabama and Apalachicola river drainages potentially serving as

biogeographic barriers (Fig. S8). This consideration was markedly absent in previous

morphological analyses (Butler *et al.* 2011) that concluded *T. c. major* merely represented an area

of admixture between other *Terrapene* in the region. We did indeed detect considerable

admixture between *T. c. major*, *T. c. carolina*, and *T. m. triunguis*, but the presence of two

apparently non-admixed *T. c. major* populations demonstrated the existence of cryptic genetic

variation (Douglas *et al.* 2009). We interpret these populations as representing distinct

evolutionary significant units (ESUs) or (at worst) management units (MUs).

Second, admixture is apparent among all southeastern *Terrapene* except *T. bauri*, which

absorbed the greatest amount of DAPC variation. This is likely attributed to Pliocene vicariance

in the Florida Peninsula, where the Okefenokee Trough divided northern from southern Florida

(Bert 1986; Douglas *et al.* 2009). Each of these aspects will require careful consideration when

conservation efforts are planned or implemented, particularly given that *Terrapene* are in decline

throughout their range (Dodd 2001).

On the other hand, *T. ornata* and *T. carolina* are separated by greater genetic distances

than are the southeastern taxa (Martin *et al.* 2013), which may suggest the presence of intrinsic

genetic incompatibilities (Barton 2001; Abbott *et al.* 2013) and is consistent with the lack of

hybrids beyond the F_1 generation. Furthermore, the low frequency of F_1 hybrids observed in the

Illinois ONxEA population may have resulted from recent degradation of the prairie grassland

habitat (Manning 2001; Mussmann *et al.* 2017), which subsequently initiated increased

heterospecific contact or otherwise disturbed reproductive boundaries by altering the fitness

consequences of hybridization (Chafin et al. 2019; Grabenstein and Taylor 2018). However, this

hypothesis cannot be explicitly tested herein. Although we did not identify contemporary back-

crossed individuals, Cureton *et al.* (2011) did depict potential introgressive hybridization between

ONxEA based on mitochondrial DNA and multiple microsatellite markers. These discrepancies

may reflect either historical admixture as a source of introgression, which we did not explicitly

test for, or back-crossed hybrids as a rarity not encountered in our sampling.

4.2. *Biogeography and hybrid zone formation in Terrapene*

The disparity in early and late-generation hybrids between the midwestern (ONxEA) and

southeastern (EAxGU, GUxTT, and EAxTT) hybrid zones suggests differences in the underlying

evolutionary processes. Such differences could involve regional variability in the extent or nature

of reproductive isolation, or simply their respective biogeographic histories. Pleistocene

glaciation precipitated numerous widespread distributional shifts across many taxa in the

Midwest (King 1981; Webb 1981). Subsequent postglacial "shuffling" has also been implicated

in as an historical driver of introgressive hybridization in populations of *Sistrurus* rattlesnakes

now allopatric in the same region (Sovic *et al.* 2016). The same process could explain evidence

for past introgression in the ONxEA hybrid zone (Cureton *et al* 2011b), despite a lack of

contemporary hybrids. Here, a rapid re-colonization [especially from northern refugia such as the

"Driftless Area" of Wisconsin, Iowa, and Illinois during the last glacial maximum (Holliday *et al.*

2002)] resulted in north-south contact zones near the glacial maximum (i.e., "leading-edge"

hypothesis; Hewitt 1996, 2000). However, our purported range overlap also represents a broad

interdigitation of "prairie" and "interior highland" habitats (Ennen *et al.* 2017), indicating that

later-generation hybrids simply requires a finer-scale sampling than ours to be detected.

The southeastern United States has long been known for the of co-occurrence of contact

zones (Remington 1968; Avise 2000), with migration from refugia in southern Florida and east

Texas/ west Louisiana as an hypothesized mechanism (Swenson & Howard 2005). The

divergence of these lineages prior to Pleistocene glaciation (Martin *et al.* 2013) seemingly

indicates postglacial expansion as a potential mechanism underlying southeastern hybrid zone

formation. Likewise, finer-scale phylogeographic breaks that corroborate those in *Terrapene* have

also been detected in numerous other turtles (Walker & Avise 1998). For example, *Sternotherus*

minor and *S. odoratus* show deep east-west phylogeographic breaks approximately centered on

Alabama, with unique lineages in peninsular Florida (Iverson 1977; Walker & Avise 1998).

Kinosternon subrubrum mirrors this pattern, but with an additional unique lineage in the

panhandle region (Walker *et al.* 1998). Similar breaks again appear in *Trachemys scripta,*

Macroclemys temminckii (Walker & Avise 1998; Roman *et al.* 1999), and *Gopherus polyphemus*

(Lamb *et al.* 1989). It is thus no surprise for the region to be identified as a hotspot for inter-

specific contact and phylogeographic concordance (Soltis *et al.* 2006; Rissler & Smith 2010), also

reflected in *Terrapene*. As always, it becomes inherently difficult to separate the relative

importance of historical processes from contemporary physiographic or ecological features.

4.3 *Functional Genomic Architecture in the Southeastern Hybrid Zone*

Regardless of the biogeographic scenarios invoked, a clear hotspot for hybridization exists

in the southeastern United States. The variance among locus-specific patterns of differentiation or

exchange allows for the potential interpretation of adaptive processes, at least in the context of

neutral expectations. SNPs located in several mRNA loci are implicated as potentially

contributing to selection in *Terrapene* from three southeastern hybrid zones. Among inter-

specific comparisons (Figs. 3B, 3C, 4B, 4C), the dominant pattern was a steep, sigmoidal cline

(GUxTT and EAxTT, but most clearly apparent in the latter). This accordingly points to selection

against interspecific heterozygotes (Fitzpatrick 2013). In contrast, the selective advantage of EA

alleles in the EAxGU hybrid zone (Figs. 3A, 4A) fails to agree with the general genome-wide

pattern of underdominance (Fig. S18), which suggests directional introgression within which EA

alleles are favored in hybrids under contemporary conditions. Mapping BGC outliers to

Trachemys chromosomes (Fig. 4) also revealed their ubiquitous rather than finely concentrated

distribution across the genome, and this in turn highlighted the differential nature of the observed

interspecific introgression. Inaccuracies regarding mapping a *Terrapene* assembly against a

Trachemys reference genome is also a possibility, although we attempted to minimize this by

resolving conflicts via PAFSCAFF.

Loci with steep genomic clines were most strongly correlated with temperature predictors,

suggesting the importance of thermal adaptations in maintaining species boundaries between

southeastern *Terrapene*. In contrast, neither precipitation nor wind-associated outliers followed

this pattern. Given the positive relationship between outlier genes and thermal predictors, a

natural extrapolation would be that a thermal gradient drives differential introgression. This has

multiple implications regarding the integrity of species boundaries if *Terrapene* are subjected to

future environmental changes. In one scenario, a shifting adaptive landscape may promote

hybridization by contravening long-term reproductive isolation (EAxGU), with subsequent

introgression at specific loci (as herein). Alternatively, rapid environmental change could simply

outpace the selective filtering of maladaptive variants, with a subsequent decrease in fitness

(Kokko *et al.* 2017). This would be particularly evident when effective population sizes are

already depressed following a population bottleneck (Chafin *et al.* 2019). Here, extreme rates of

change may also link with a genetic swamping effect (i.e., replacement of local genotypes by

hybrids; Todesco *et al.* 2016). Both scenarios implicate anthropogenic pressures as governing the

fates of diverse taxa across hybrid zones (Taylor *et al.* 2015).

The putative functions of the nine outlier loci supported by genomic clines and RDA

provide additional support for the strong impact of thermal selection. Two loci potentially relate

to TSD during embryonic development, while others seemingly associate with molecular

pathways in skeletal muscle and nervous tissues that involve tolerance to anoxia and hypoxia

(N=6), and immune response to pathogens [(N=2); see Table 1 for sources]. Anoxia/ hypoxia-

related genes have been associated with freeze tolerance in hibernating turtles (Storey 2006), thus

supporting an obvious association with thermal gradients. Here, three loci (*SYPL2, ACAD11,* and

TMEM214) may regulate brain function and metabolism by up-regulating Ca^{2+} concentrations

(Takeshima *et al.* 1998; Pamenter *et al.* 2016), inducing lipid metabolism (He *et al.* 2011; Gomez

& Richards 2018), and initiating stress-induced apoptosis (Kesaraju *et al.* 2009; Li *et al.* 2013).

Similarly, the *CITED4* gene (EAxTT) potentially inhibits hypoxia-related transcription factors

(Fox *et al.* 2004), whereas *FAM89B* (GUxTT) may become up-regulated when physiological

conditions turn hypoxic (Goyal & Longo 2014). The regulation of immune function is less clearly

associated with underlying thermal gradients, but may associate instead with behavioral

thermoregulation during infection, given that infection resistance increases at warmer

temperatures (Dodd 2001; Agha *et al.* 2017). However, we remain cautious in that these genes

have not been associated with specific functions in *Terrapene,* and to do so remains speculative.

Nevertheless, their potential connection with thermal adaptations is consistent with the RDA.

We emphasize that ectothermic vertebrates are exceptionally vulnerable to contemporary

pressures, and reflect an elevated extinction risk due to a strong reliance on environmental

thermoregulation and a dependence on suitable habitat (Gibbons *et al.* 2000; Sinervo *et al.* 2010;

Winter *et al.* 2016). Indeed, ectotherms can exhibit reduced fitness and growth-rates when

environmental conditions exceed their thermal optima (Deutsch *et al.* 2008; Martin & Huey 2008;

McCallum *et al.* 2009; Huey *et al.* 2012; Huey & Kingsolver 2019). Increased temperatures also

impact physiological pathways, and these are putatively comparable with the genes described

herein, such as increasing metabolic rates (Dillon *et al.* 2010), intensifying hypoxic stress despite

higher temperature-driven O_2 demands (Huey & Ward 2005), heightening disease transmission

(Pounds *et al.* 2006), or even the over-extension of thermal tolerances (Sinervo *et al.* 2010; Ceia-

Hasse *et al.* 2014). Going forward, climate change may facilitate evolutionary responses to

changing thermal conditions, potentially including local adaptation (Holt 1990; Norberg *et al.*

2012; Bush *et al.* 2016), physiological and behavioral mechanisms (e.g., thermoregulation,

phenology), plasticity (Urban *et al.* 2014; Sgrò *et al.* 2016), of shifts in species distributions

(Parmesan *et al.* 1999; Parmesan & Yohe 2003; Moreno-Rueda *et al.* 2012). Accordingly,

ectothermic species boundaries may be particularly susceptible if indeed governed by thermal

conditions, as seemingly exemplified herein with *Terrapene*.

4.3. *Conclusions*

Our study suggests that reproductive isolation in turtles involves numerous mechanisms

regulated by thermally induced selective pressures. Clearly, such selective pressures play a

prominent role in chelonian ecology. Many turtle species are at elevated risks from climate

change due to the imposition of TSD during embryonic development, as well as prolonged

generation times. Similarly, long generation times can also restrict the adaptive capacity of turtles

in a rapidly changing climate (Hoffmann *et al.* 2017). Here, climate change can shift sex-ratios

and promote demographic collapse, with warmer temperatures initiating male bias in biological

sex, and vice versa (Janzen 1994).

Two important evolutionary implications are evident in our data. First, we demonstrated

differential introgression along an ecological gradient in three taxa inhabiting a North American

hybrid zone. We then assessed scaffold-aligned and transcriptomic SNPs to identify several genes

whose functions are consistent with physiological processes related to thermal ecology, and as

such are capable of promoting adaptive divergence. In this sense, they potentially describe

ecological gradients related to TSD, anoxia/ hypoxia tolerance, and immune response in a hybrid

zone encompassing three taxa. While we acknowledge that the observed clinal patterns could

represent "molecular spandrels" reflecting an underlying neutral process, such as isolation-by-

distance, we sought environmental associations by actively controlling for spatial autocorrelation

as a corroboration of clinal loci with a putative adaptive role (Vasemägi 2006; Barrett & Hoekstra

2011). We also underscore specific loci displaying genomic cline patterns consistent with

directional introgression and selection, and these loci are potentially sustaining divergence across

species.

Second, we characterized a southeastern North American hybrid zone representing a

variety of biodiversity elements as being susceptible to anthropogenic and environmental changes

(Remington 1968; Swenson & Howard 2005; Rissler & Smith 2010). Our results demonstrate

that NGS population genomic methods can clearly identify population structure and detect

introgression, whereas traditional Sanger sequencing methods are inadequate to do so (Butler *et*

al. 2011; Martin *et al.* 2013). We also underscore specific loci prone to directional introgression

and selection that may potentially sustain divergence across species.

ACKNOWLEDGEMENTS

The research herein was conducted in partial fulfillment of the Ph.D. degree in Biological

Sciences at the University of Arkansas (BTM). We would like to extend our gratitude to the

countless volunteers and organizations who collected and/or provided tissue samples (Table S1).

We also thank the current and former members of the Douglas Lab and University of Arkansas

faculty who provided advice and support, especially A. Alverson, W. Anthonysamy, M. Bangs, J.

Koukl, S. Mussmann, J. Pummill, and Z. Zbinden. Finally, many thanks to the reviewers, whose

comments led to a greatly improved manuscript. Sample collections were approved under the

University of Texas-Tyler Animal Care and Use Committee (IACUC) permit #113 and

University of Illinois IACUC protocols 16160 and 18000. Funding was provided by the Lucille F.

Stickle Fund of the North American Box Turtle Committee, the American Turtle Observatory

(ATO), and the following endowments: The Bruker Professorship in Life Sciences (MRD) and

the Twenty-First Century Chair in Global Change Biology (MED). Analytical resources were

provided by the Arkansas High Performance Computing Cluster (AHPCC) and an NSF-XSEDE

Research Allocation (TG-BIO160065) that allowed access the Jetstream cloud service.

5. REFERENCES

Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJE, Bierne N, Boughman J, Brelsford A,
Buerkle CA, and Buggs R (2013) Hybridization and Speciation. *Journal of Evolutionary*
Biology, **26**, 229–246.

Agha M, Price S, Nowakowski A, Augustine B, and Todd B (2017) Mass mortality of eastern
box turtles with upper respiratory disease following atypical cold weather. *Diseases of*
Aquatic Organisms, **124**, 91–100.

Alexander DH, Novembre J, and Lange K (2009) Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research*, **19**, 1655–1664.

Anderson EC and Thompson EA (2002) A model-based method for identifying species hybrids
using multilocus genetic data. *Genetics*, **160**, 1217–1229.

Andrés JA, Larson EL, Bogdanowicz SM, and Harrison RG (2013) Patterns of transcriptome
divergence in the male accessory gland of two closely related species of field crickets.
Genetics, **193**, 501–13.

Auffenberg W (1958) Fossil turtles of the genus *Terrapene* in Florida. *Bulletin of the Florida
State Museum*, **3**, 53–92.

Auffenberg W (1959) A Pleistocene *Terrapene* hibernaculum, with remarks on a second
complete box turtle skull from Florida. *Quarterly Journal of the Florida Academy of
Science*, **22**, 49–53.

Avise JC (2000) *Phylogeography*. Harvard University Press, Cambridge, MA.

Babik W, Dudek K, Fijarczyk A, Pabijan M, Stuglik M, Szkotak R, and Zieliński P (2015)
Constraint and Adaptation in newt Toll-Like Receptor Genes. *Genome Biology and
Evolution*, **7**, 81–95.

Barrett RDH and Hoekstra HE (2011) Molecular spandrels: Tests of adaptation at the genetic
level. *Nature Reviews Genetics*, **12**, 767–780.

Barton NH (2001) The role of hybridization in evolution. *Molecular Ecology*, **10**, 551–568.

Barton NH and Hewitt GM (1985) Analysis of hybrid zones. *Annual Review of Ecology and
Systematics*, **16**, 113–148.

Bert T (1986) Speciation in western Atlantic stone crabs (genus *Menippe*): the role of geological
processes and climatic events in the formation and distribution of species. *Marine Biology*,
93, 157–170.

Borcard D, Gillet F, and Legendre P (2018) Spatial analysis of ecological data. In: *Numerical
ecology with R*, pp. 299–367. Springer.

Bush A, Mokany K, Catullo R, Hoffmann A, Kellermann V, Sgrò C, McEvey S, and Ferrier S
(2016) Incorporating evolutionary adaptation in species distribution modelling reduces
projected vulnerability to climate change. *Ecology Letters*, **19**, 1468–1478.

Butler JM, Dodd Jr. CK, Aresco M, and Austin JD (2011) Morphological and molecular evidence
indicates that the Gulf Coast box turtle (*Terrapene carolina major*) is not a distinct
evolutionary lineage in the Florida Panhandle. *Biological Journal of the Linnean Society*,
102, 889–901.

Caye K, Deist TM, Martins H, Michel O, and François O (2016) TESS3: Fast inference of spatial
population structure and genome scans for selection. *Molecular Ecology Resources*, **16**,
540–548.

Ceia-Hasse A, Sinervo B, Vicente L, and Pereira HM (2014) Integrating ecophysiological models into species distribution projections of European reptile range shifts in response to climate change. *Ecography*, **37**, 679–688.

Chafin TK, Douglas MR, Martin BT, and Douglas ME (2019) Hybridization drives genetic erosion in sympatric desert fishes of western North America. *Heredity*, **123**, 759–773.

Chafin TK, Martin BT, Mussmann SM, Douglas MR, and Douglas ME (2018) FRAGMATIC: in silico locus prediction and its utility in optimizing ddRADseq projects. *Conservation Genetics Resources*, **10**, 325–328.

Cureton JC, Buchman AB, Deaton R, and Lutterschmidt WI (2011) Molecular analysis of hybridization between the box turtles *Terrapene carolina* and *T. ornata*. *Copeia*, **2011**, 270–277.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G, Durbin R, and Group 1000 Genomes Project Analysis (2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–2158.

Deutsch CA, Tewksbury JJ, Huey RB, Sheldon KS, Ghalambor CK, Haak DC, and Martin PR (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*, **105**, 6668–6672.

Dillon ME, Wang G, and Huey RB (2010) Global metabolic impacts of recent climate warming. *Nature*, **467**, 704–706.

Dodd KC (2001) *North American Box Turtles, A Natural History*. University of Oklahoma Press, Norman, OK, USA.

Douglas ME, Douglas MR, Schuett GW, and Porras LW (2009) Climate change and evolution of the New World pitviper genus *Agkistrodon* (Viperidae). *Journal of Biogeography*, **36**, 1164–1180.

Eaton DAR and Overcast I (2020) ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics*, **36**, 2592–2594.

Ennen JR, Matamoros WA, Agha M, Lovich JE, Sweat SC, and Hoagstrom CW (2017) Hierarchical, Quantitative Biogeographic Provinces for All North American Turtles and Their Contribution to the Biogeography of Turtles and the Continent. *Herpetological Monographs*, **31**, 114–140.

Evanno G, Regnaut S, and Goudet J (2005) Detecting the number of clusters of individuals using the software structure: a simulation study. *Molecular Ecology*, **14**, 2611–2620.

Feder JL, Flaxman SM, Egan SP, and Nosil P (2013) Hybridization and the build-up of genomic divergence during speciation. *Journal of Evolutionary Biology*, **26**, 261–266.

Feldman CR and Parham JF (2002) Molecular phylogenetics of emydine turtles: Taxonomic revision and the evolution of shell kinesis. *Molecular Phylogenetics and Evolution*, **22**, 388–398.

Fick SE and Hijmans RJ (2017) WorldClim 2: new 1 km spatial resolution climate surfaces for
global land areas. *International journal of climatology*, **37**, 4302–4315.

Fitzpatrick BM (2013) Alternative forms for genomic clines. *Ecology and Evolution*, **3**, 1951–
1966.

Fox SB, Braganca J, Turley H, Campo L, Han C, Gatter KC, Bhattacharya S, and Harris AL
(2004) CITED4 inhibits hypoxia-activated transcription in cancer cells, and its cytoplasmic
location in breast cancer is associated with elevated expression of tumor cell hypoxia-
inducible factor 1α. *Cancer Research*, **64**, 6075–6081.

Garroway CJ, Bowman J, Cascaden TJ, Holloway GL, Mahan CG, Malcolm JR, Steele MA,
Turner G, and Wilson PJ (2010) Climate change induced hybridization in flying squirrels.
Global Change Biology, **16**, 113–121.

Gibbons JW, Scott DE, Ryan TJ, Buhlmann KA, Tuberville TD, Metts BS, Greene JL, Mills T,
Leiden Y, Poppy S, and Winne CT (2000) The global decline of reptiles, déjà vu
amphibians: reptile species are declining on a global scale. Six significant threats to reptile
populations are habitat loss and degradation, introduced invasive species, environmental
pollution, disease, unsustaina. *Bioscience*, **50**, 653–666.

Gomez CR and Richards JG (2018) Mitochondrial responses to anoxia exposure in red eared
sliders (*Trachemys scripta*). *Comparative Biochemistry and Physiology Part B:
Biochemistry and Molecular Biology*, **224**, 71–78.

Gompert Z and Buerkle CA (2010) INTROGRESS: a software package for mapping components
of isolation in hybrids. *Molecular Ecology Resources*, **10**, 378–384.

Gompert Z and Buerkle CA (2011) Bayesian estimation of genomic clines. *Molecular Ecology*,
20, 2111–2127.

Gompert Z and Buerkle CA (2012) BGC: Software for Bayesian estimation of genomic clines.
Molecular Ecology Resources, **12**, 1168–1176.

Goyal R and Longo LD (2014) Acclimatization to long-term hypoxia: gene expression in ovine
carotid arteries. *Physiological Genomics*, **46**, 725–734.

Grabenstein KC and Taylor SA (2018) Breaking barriers: causes, consequences, and
experimental utility of human-mediated hybridization. *Trends in Ecology and Evolution*, **33**,
198–212.

Guo B, Lu D, Liao WB, and Merilä J (2016) Genomewide scan for adaptive differentiation along
altitudinal gradient in the Andrew's toad *Bufo andrewsi*. *Molecular Ecology*, **25**, 3884–
3900.

Hao Z, Lv D, Ge Y, Shi J, Weijers D, Yu G, and Chen J (2020) RIdeogram: drawing SVG
graphics to visualize and map genome-wide data on the ideograms. *PeerJ Computer
Science*, **6**, e251.

He M, Pei Z, Mohsen A-W, Watkins P, Murdoch G, Van Veldhoven PP, Ensenauer R, and

Vockley J (2011) Identification and characterization of new long chain acyl-CoA dehydrogenases. *Molecular Genetics and Metabolism*, **102**, 418–429.

Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnaean Society*, **58**, 247–276.

Hewitt GM (2000) The genetic legacy of the Quaternary ice ages. *Nature*, **405**, 907–913.

Hijmans RJ and Van Etten J (2016) raster: Geographic Data Analysis and Modeling. R package version 2.5-8.

Hoffmann AA, Sgrò CM, and Kristensen TN (2017) Revisiting adaptive potential, population size, and conservation. *Trends in Ecology and Evolution*, **32**, 506–517.

Holliday VT, Knox JC, Running I V., Mandel RD, and Ferring CR (2002) The central lowlands. In: *The physical geography of North America* (ed Orme AR), pp. 335–362. Oxford University Press, New York.

Holt RD (1990) The microevolutionary consequences of climate change. *Trends in Ecology and Evolution*, **5**, 311–315.

Huey RB, Kearney MR, Krockenberger A, Holtum JAM, Jess M, and Williams SE (2012) Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **367**, 1665–1679.

Huey RB and Kingsolver JG (2019) Climate warming, resource availability, and the metabolic meltdown of ectotherms. *The American Naturalist*, **194**, E140–E150.

Huey RB and Ward PD (2005) Hypoxia, global warming, and terrestrial late Permian extinctions. *Science*, **308**, 398–401.

Iverson JB (1977) Geographic variation in the musk turtle, *Sternotherus minor*. *Copeia*, **1977**, 502.

Iverson JB, Meylan PA, and Seidel ME (2017) Testudines—Turtles. In: *Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico, with Comments Regarding Confidence in Our Understanding* (ed Crother BI), pp. 82-91. SSAR Herpetological Circular 43.

Janzen FJ (1994) Climate change and temperature-dependent sex determination in reptiles. *Proceedings of the National Academy of Sciences of the United States of America*, **91**, 7487–90.

Jay F, Manel S, Alvarez N, Durand EY, Thuiller W, Holderegger R, Taberlet P, and François O (2012) Forecasting changes in population genetic structure of alpine plants in response to global warming. *Molecular Ecology*, **21**, 2354–2368.

Johnson W (1994) Woodland expansion in the Platte River, Nebraska: patterns and causes. *Ecological Monographs*, **64**, 45–84.

Jombart T, Devillard S, and Balloux F (2010) Discriminant analysis of principal components: a
new method for the analysis of genetically structured populations. *BMC Genetics*, **11**, 94.

Kearns AM, Restani M, Szabo I, Schrøder-Nielsen A, Kim JA, Richardson HM, Marzluff JM,
Fleischer RC, Johnsen A, and Omland KE (2018) Genomic evidence of speciation reversal
in ravens. *Nature Communications*, **9**, 906.

Keller I and Seehausen O (2012) Thermal adaptation and ecological speciation. *Molecular
Ecology*, **21**, 782–799.

Kesaraju S, Schmidt-Kastner R, Prentice HM, and Milton SL (2009) Modulation of stress
proteins and apoptotic regulators in the anoxia tolerant turtle brain. *Journal of
Neurochemistry*, **109**, 1413–1426.

Key KHL (1968) The concept of stasipatric speciation. *Systematic Biology*, **17**, 14–22.

King JE (1981) Late Quaternary vegetational history of Illinois. *Ecological Monographs*, **51**, 43–
62.

Kingsolver JG (2009) The well-temperatured biologist. (American Society of Naturalists
Presidential Address). *The American Naturalist*, **174**, 755–68.

Kokko H, Chaturvedi A, Croll D, Fischer MC, Guillaume F, Karrenberg S, Kerr B, Rolshausen
G, and Stapley J (2017) Can evolution supply what ecology demands? *Trends in Ecology
and Evolution*, **32**, 187–197.

Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, and Mayrose I (2015) CLUMPAK: a
program for identifying clustering modes and packaging population structure inferences
across *K. Molecular Ecology Resources*, **15**, 1179–1191.

Kumaresan V, Pasupuleti M, Arasu MV, Al-Dhabi NA, Arshad A, Amin SMN, Yusoff FM, and
Arockiaraj J (2018) A comparative transcriptome approach for identification of molecular
changes in Aphanomyces invadans infected Channa striatus. *Molecular Biology Reports*, **45**,
2511–2523.

Lamb T, Avise JC, and Gibbons JW (1989) Phylogeographic patterns in mitochondrial-dna of the
desert tortoise (Xerobates-agassizi), and evolutionary relationships among the north-
american gopher tortoises. *Evolution*, **43**, 76–87.

van Lear DH and Harlow RF (2002) Fire in the eastern United States: influence on wildlife
habitat. In: *Proceedings: the role of fire for nongame wildlife management and community
restoration: traditional uses and new directions. General Technical Report 288* (eds Ford
W., Russell KR and Moorman CE), pp. 2–10. US Dept. of Agriculture, Forest Service,
Northeastern Research Station.

Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*, **34**, 3094–
3100.

Li YF, Costello JC, Holloway AK, and Hahn MW (2008) "Reverse ecology" and the power of
population genomics. *Evolution*, **62**, 2984–2994.

Li C, Wei J, Li Y, He X, Zhou Q, Yan J, Zhang J, Liu Y, Liu Y, and Shu H-B (2013) Transmembrane Protein 214 (TMEM214) mediates endoplasmic reticulum stress-induced caspase 4 enzyme activation and apoptosis. *The Journal of Biological Chemistry*, **288**, 17908–17.

Linck EB and Battey CJ (2019) Minor allele frequency thresholds strongly affect population structure inference with genomic datasets. *Molecular Ecology Resources*, **19**, 639–647.

Lydeard C and Mayden RL (1995) A diverse and endangered aquatic ecosystem of the southeast United States. *Conservation Biology*, **9**, 800–805.

Manning B (2001) *Critical Trends in Illinois Ecosystems*. Illinois Department of Natural Resources. Springfield, IL.

Martin BT, Bernstein NP, Birkhead RD, Koukl JF, Mussmann SM, and Placyk JS (2013) Sequence-based molecular phylogenetics and phylogeography of the American box turtles (*Terrapene* spp.) with support from DNA barcoding. *Molecular Phylogenetics and Evolution*, **68**, 119–134.

Martin BT, Bernstein NP, Birkhead RD, Koukl JF, Mussmann SM, and Placyk Jr JS (2014) On the reclassification of the *Terrapene* (Testudines: Emydidae): a response to Fritz & Havaš. *Zootaxa*, **3835**, 292–294.

Martin BT, Chafin TK, Douglas MR, Placyk JS, Birkhead RD, Phillips CA, and Douglas ME (2020) Machine learning substantiates biologically meaningful species delimitations in the phylogenetically complex North American box turtle genus *Terrapene*. *bioRxiv, doi: https://doi.org/10.1101/2020.05.19.103598*.

Martin TL and Huey RB (2008) Why suboptimal is optimal: Jensen's inequality and ectotherm thermal preferences. *The American Naturalist*, **171**, E102-18.

McCallum M, McCallum JL, and Trauth SE (2009) Predicted climate change may spark box turtle declines. *Amphibia-Replilia*, **30**, 259–264.

Milstead WW (1969) Studies on the evolution of the box turtles (genus *Terrapene*). *Bulletin of the Florida State Museum, Biological Science Series*, **14**, 1–113.

Milstead WW and Tinkle DW (1967) *Terrapene* of Western Mexico, with comments on species groups in the genus. *Copeia*, **1967**, 180–187.

Moreno-Rueda G, Pleguezuelos JM, Pizarro M, and Montori A (2012) Northward shifts of the distributions of Spanish reptiles in association with climate change. *Conservation Biology*, **26**, 278–283.

Muhlfeld CC, Kovach RP, Jones LA, Al-Chokhachy R, Boyer MC, Leary RF, Lowe WH, Luikart G, and Allendorf FW (2014) Invasive hybridization in a threatened species is accelerated by climate change. *Nature Climate Change*, **4**, 620–624.

Mussmann SM, Douglas MR, Anthonysamy WJB, Davis MA, Simpson SA, Louis W, and Douglas ME (2017) Genetic rescue, the greater prairie chicken and the problem of

conservation reliance in the Anthropocene. *Royal Society Open Science*, **4**, 160736.

Mussmann SM, Douglas MR, Chafin TK, and Douglas ME (2020) AdmixPipe: population
analyses in Admixture for non-model organisms. *BMC Bioinformatics*, **21**, 1–9.

Norberg J, Urban MC, Vellend M, Klausmeier CA, and Loeuille N (2012) Eco-evolutionary
responses of biodiversity to climate change. *Nature Climate Change*, **2**, 747–751.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara
RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, and Wagner H (2019) vegan:
Community Ecology Package.

Paitz RT and Bowden RM (2008) A proposed role of the sulfotransferase/sulfatase pathway in
modulating yolk steroid effects. *Integrative and Comparative Biology*, **48**, 419–427.

Pamenter ME, Gomez CR, Richards JG, and Milsom WK (2016) Mitochondrial responses to
prolonged anoxia in brain of red-eared slider turtles. *Biology Letters*, **12**, 20150797.

Parmesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, Descimon H, Huntley B, Kaila L,
Kullberg J, Tammaru T, Tennent WJ, Thomas JA, and Warren M (1999) Poleward shifts in
geographical ranges of butterfly species associated with regional warming. *Nature*, **399**,
579–583.

Parmesan C and Yohe G (2003) A globally coherent fingerprint of climate change impacts across
natural systems. *Nature*, **421**, 37–42.

Payseur BA (2010) Using differential introgression in hybrid zones to identify genomic regions
involved in speciation. *Molecular Ecology Resources*, **10**, 806–820.

Peterson RA and Cavanaugh JE (2019) Ordered quantile normalization: a semiparametric
transformation built for the cross-validation era. *Journal of Applied Statistics*, **2019**, 1–16.

Peterson BK, Weber JN, Kay EH, Fisher HS, and Hoekstra HE (2012) Double digest RADseq: an
inexpensive method for de novo SNP discovery and genotyping in model and non-model
species. *PLOS ONE*, **7**, e37135.

Pounds AJ, Bustamante MR, Coloma LA, Consuegra JA, Fogden MPL, Foster PN, La Marca E,
Masters KL, Merino-Viteri A, Puschendorf R, Ron SR, Sánchez-Azofeifa GA, Still CJ, and
Young BE (2006) Widespread amphibian extinctions from epidemic disease driven by
global warming. *Nature*, **439**, 161–167.

Qin D, Plattner G, Tignor M, Allen S, Boschung J, Nauels A, Xia Y, Bex V, and Midgley P
(2013) *Summary for policymakers. Climate change 2013: the physical science basis.*
Contribution of Working Group I to the fifth assessment report of the Intergovernmental
Panel on Climate Change, eds Stocker, TF et al. Cambridge University Press, Cambridge,
UK.

Remington CL (1968) Suture-zones of hybrid interaction between recently joined biotas. In:
Evolutionary Biology (ed Dobzhansky T), pp. 321–428. Springer, New York, NY.

Rhymer JM and Simberloff D (1996) Extinction by hybridization and introgression. *Annual*
Review of Ecology and Systematics, **27**, 83–109.

Ricketts TH (1999) *Terrestrial ecoregions of North America: a conservation assessment*. Island
Press, Washington, DC.

Rissler LJ and Smith WH (2010) Mapping amphibian contact zones and phylogeographical break
hotspots across the United States. *Molecular Ecology*, **19**, 5404–5416.

Rödin☐Mörch P, Luquet E, Meyer☐Lucht Y, Richter☐Boix A, Höglund J, and Laurila A (2019)
Latitudinal divergence in a wide☐spread amphibian: contrasting patterns of neutral and
adaptive genomic variation. *Molecular Ecology*, **28**, 2996–3011.

Roman J, Santhuff SD, Moler PE, and Bowen BW (1999) Population structure and cryptic
evolutionary units in the alligator snapping turtle. *Conservation Biology*, **13**, 135–142.

Rosenberg NA (2004) DISTRUCT: a program for the graphical display of population structure.
Molecular Ecology Notes, **4**, 137–138.

Rosenzweig C, Karoly D, Vicarelli M, Neofotis P, Wu Q, Casassa G, Menzel A, Root TL,
Estrella N, Seguin B, Tryjanowski P, Liu C, Rawlins S, and Imeson A (2008) Attributing
physical and biological impacts to anthropogenic climate change. *Nature*, **453**, 353–357.

Ryan SF, Deines JM, Scriber JM, Pfrender ME, Jones SE, Emrich SJ, and Hellmann JJ (2018)
Climate-mediated hybrid zone movement revealed with genomics, museum collection, and
simulation modeling. *Proceedings of the National Academy of Sciences*, 2017–14950.

Samson F and Knopf F (1994) Prairie conservation in North America. *Bioscience*, **44**, 418–421.

Samson FB, Knopf FL, and Ostlie WR (2004) Great Plains ecosystems: past, present, and future.
Wildlife Society Bulletin, **32**, 6–15.

Seehausen O, Takimoto G, Roy D, and Jokela J (2008) Speciation reversal and biodiversity
dynamics with hybridization in changing environments. *Molecular Ecology*, **17**, 30–44.

Sgrò CM, Terblanche JS, and Hoffmann AA (2016) What Can Plasticity Contribute to Insect
Responses to Climate Change? *Annual Review of Entomology*, **61**, 433–451.

Shaffer HB, Minx P, Warren DE, Shedlock AM, Thomson RC, Valenzuela N, Abramyan J,
Amemiya CT, Badenhorst D, and Biggar KK (2013) The western painted turtle genome, a
model for the evolution of extreme physiological adaptations in a slowly evolving lineage.
Genome Biology, **14**, R28.

Simison W, Parham J, Papenfuss T, Lam A, and Henderson J (2020) Annotated chromosome-
level reference genome of the red-eared slider turtle (*Trachemys scripta elegans*). *Genome*
Biology and Evolution, **12**, 456–462.

Sinervo B, Méndez-de-la-Cruz F, Miles DB, Heulin B, Bastiaans E, Villagrán-Santa Cruz M,
Lara-Resendiz R, Martínez-Méndez N, Calderón-Espinosa ML, Meza-Lázaro RN, Gadsden
H, Avila LJ, Morando M, De la Riva IJ, Victoriano Sepulveda P, Rocha CFD,

Ibargüengoytía N, Aguilar Puntriano C, Massot M *et al.* (2010) Erosion of lizard diversity by climate change and altered thermal niches. *Science*, **328**, 894–9.

Soltis DE, Morris AB, McLachlan JS, Manos PS, and Soltis PS (2006) Comparative phylogeography of unglaciated eastern North America. *Molecular Ecology*, **15**, 4261–4293.

Sovic MG, Fries AC, and Gibbs HL (2016) Origin of a cryptic lineage in a threatened reptile through isolation and historical hybridization. *Heredity*, **117**, 358–366.

Stapanian MA, Cassell DL, and Cline SP (1997) Regional patterns of local diversity of trees: associations with anthropogenic disturbance. *Forest Ecololgy and Management*, **93**, 33–44.

Stapanian MA, Sundberg SD, Baumgardner GA, and Liston A (1998) Alien plant species composition and associations with anthropogenic disturbance in North American forests. *Plant Ecology*, **139**, 49–62.

Storey KB (2006) Reptile freeze tolerance: metabolism and gene expression. *Cryobiology*, **52**, 1–16.

Swenson NG and Howard DJ (2004) Do suture zones exist? *Evolution*, **58**, 2391–2397.

Swenson NG and Howard DJ (2005) Clustering of contact zones, hybrid zones, and phylogeographic breaks in North America. *The American Naturalist*, **166**, 581–591.

Takeshima H, Shimuta M, Komazaki S, Ohmi K, Nishi M, Iino M, Miyata A, and Kangawa K (1998) Mitsugumin29, a novel synaptophysin family member from the triad junction in skeletal muscle. *The Biochemical Journal*, **331**, 317–22.

Taylor SA, Larson EL, and Harrison RG (2015) Hybrid zones: windows on climate change. *Trends in Ecology and Evolution*, **30**, 398–406.

Taylor SA, White TA, Hochachka WM, Ferretti V, Curry RL, and Lovette I (2014) Climate-mediated movement of an avian hybrid zone. *Current Biology*, **24**, 671–676.

Teske PR, Sandoval-Castillo J, Golla TR, Emami-Khoyi A, Tine M, von der Heyden S, and Beheregaray LB (2019) Thermal selection as a driver of marine ecological speciation. *Proceedings of the Royal Society B: Biological Sciences*, **286**, 20182023.

Tiffin P and Ross-Ibarra J (2014) Advances and limits of using population genetics to understand local adaptation. *Trends in Ecology and Evolution*, **29**, 673–680.

Todesco M, Pascual MA, Owens GL, Ostevik KL, Moyers BT, Hübner S, Heredia SM, Hahn MA, Caseys C, Bock DG, and Rieseberg LH (2016) Hybridization and extinction. *Evolutionary Applications*, **9**, 892–908.

Urban MC, Richardson JL, and Freidenfelds NA (2014) Plasticity and genetic adaptation mediate amphibian and reptile responses to climate change. *Evolutionary Applications*, **7**, 88–103.

Vasemägi A (2006) The adaptive hypothesis of clinal variation revisited: Single-locus clines as a result of spatially restricted gene flow. *Genetics*, **173**, 2411–2414.

Via S (2009) Natural selection in action during speciation. *Proceedings of the National Academy*
of Sciences, **106**, 9939–9946.

Walker DE and Avise JC (1998) Principles of phylogeography as illustrated by freshwater and
terrestrial turtles in the southeastern United States. *Annual Review of Ecology and*
Systematics, **29**, 23–58.

Walker DE, Moler PE, Buhlmann KA, and Avise JC (1998) Phylogeographic patterns in
Kinosternon subrubrum and *K. baurii* based on mitochondrial DNA restriction analyses.
Herpetologica, **54**, 174–184.

Waterhouse MD, Erb LP, Beever EA, and Russello MA (2018) Adaptive population divergence
and directional gene flow across steep elevational gradients in a climate-sensitive mammal.
Molecular Ecology, **27**, 2512–2528.

Webb T (1981) The past 11,000 years of vegetational change in eastern North America.
BioScience, **31**, 501–506.

Winter M, Fiedler W, Hochachka WM, Koehncke A, Meiri S, and De la Riva I (2016) Patterns
and biases in climate change research on amphibians and reptiles: a systematic review.
Royal Society Open Science, **3**, 160158.

Wringe BF, Stanley RRE, Jeffery NW, Anderson EC, and Bradbury IR (2017a)
HYBRIDDETECTIVE: a workflow and package to facilitate the detection of hybridization
using genomic data in R. *Molecular Ecology Resources*, **17**, e275–e284.

Wringe BF, Stanley RRE, Jeffery NW, Anderson EC, and Bradbury IR (2017b)
parallelnewhybrid: an R package for the parallelization of hybrid detection using
NEWHYBRIDS. *Molecular Ecology Resources*, **17**, 91–95.

DATA ACCESSIBILITY

The demultiplexed ddRADseq reads are deposited as FASTQ files to NCBI's sequence read

archive (https://www.ncbi.nlm.nih.gov/sra); Accessions: SAMN12668545-SAMN12668981

(BioProject ID: PRJNA563121). The R scripts, metadata, and input files for each analysis are

available from a Dryad Digital Repository (https://doi.org/10.5061/dryad.brv15dv7k).

AUTHOR CONTRIBUTIONS

BTM and TKC conceived the research, laboratory, and analytical tools, scripts, and approaches. BTM performed the lab work, analyzed the data, conducted bioinformatic analyses, and wrote the manuscript. MRD and MED were the study supervisors, guided the study design, and provided funding. JSP facilitated the collection of thousands of *Terrapene* tissues and provided expertise in methodological development. RDB collected hundreds of *Terrapene* tissues from southeastern North America and facilitated the collection of many additional individuals. CAP provided tissues from the midwestern hybrid zone as well as sample site expertise in Illinois. All authors contributed to editing and revising the manuscript

Table 1: Annotation information for overlapping INTROGRESS and BGC (Bayesian Genomic Cline) outliers (Figs. 3, 4). Each outlier was derived from an alignment mapped to the *Terrapene* transcriptome, with three pairwise taxa combinations performed. The significance threshold (α_B) was determined using a Bonferroni correction for multiple tests. Bold Gene abbreviations with an asterisk (*) differ significantly from neutral expectations. EA=Woodland (*T. c. carolina*), GU=Gulf Coast (*T. c. major*), TT=Three-toed (*T. m. triunguis*).

Gene Abbr. (P-value)	BGC (α/β)	Outlier (α/β)‡	Full Gene Name	Possible Function(s)	Source(s)
EAxGU (α_B=0.01)					
SULT (P=0)*	-1.59/-0.22	-/N	Amine Sulfotransferase-like	Regulates yolk steroids during TSD†	(Paitz & Bowden 2008)
TLR9 (P=0)*	-0.74/1.14	-/N	Toll-like Receptor 9	Immune Response to Pathogens	(Babik *et al.* 2015)
ZNF236 (P=0)*	-0.94/0.14	-/N	Zinc Finger Protein 236	Unknown	
EAxTT (α_B=0.007)					
SASH3 (P=0)*	-0.49/1.35	N/+	SAM and SH3 Domain Containing 3	Immune response	(Kumaresan *et al.* 2018)
SYPL2 (P=0)*	-0.68/1.65	-/+	Synaptophysin-like Protein 2 (AKA Mitsugumin 29)	Maintenance of [Ca^{2+}] during anoxia	(Takeshima *et al.* 1998; Pamenter *et al.* 2016)
FAM89B (P=0.036)	-0.59/0.23	-/N	Family with Sequence Similarity 89, member B	Upregulated in hypoxic conditions	(Goyal & Longo 2014)
CITED4 (P=0)*	0.26/0.97	N/+	Cbp/p300 Interacting Transactivator, Domain 4	Inhibits hypoxia-activated transcription	(Fox *et al.* 2004)
TLR9 (P=0)*	-0.38/1.63	N/+	Toll-like Receptor 9	Immune Response to Pathogens	(Babik *et al.* 2015)
GUxTT (α_B=0.007)					
SASH3 (P=0)*	-0.22/1.86	N/+	SAM and SH3 Domain Containing 3	Immune response	(Kumaresan *et al.* 2018)
SYPL2 (P=0)*	-0.48/2.23	-/+	Synaptophysin-like Protein 2	Maintenance of [Ca^{2+}] during anoxia	(Takeshima *et al.* 1998; Pamenter *et al.* 2016)
FAM89B (P=0)*	-0.46/0.66	-/+	Family with Sequence Similarity 89, member B	Upregulated in hypoxic conditions	(Goyal & Longo 2014)
ACAD11 (P=0)*	-0.24/1.16	N/+	Acyl-CoA Dehydrogenase, family member 11-like	Lipid metabolism	(He *et al.* 2011; Gomez & Richards 2018)
TMEM214 (P=0)*	0.77/0.11	+/N	Transmembrane Protein 214	Stress-induced apoptosis during anoxia	(Kesaraju *et al.* 2009; Li *et al.* 2013)

†TSD=Temperature-dependent sex determination

‡α outliers=excess P$_1$ (-) or P$_2$ (+) ancestry; ß outliers=slow (-) or rapid (+) genomic cline rates; N=neutral

Table 2: WorldClim environmental predictor abbreviations for the redundancy analysis (RDA). The raster layers were obtained from https://worldclim.org. Additional variables were excluded via forward selection due to low predictive capacity or correlation with the remaining layers.

Abbr.	Full Name	BioClim No.
tmDR2	Mean diurnal range [Monthly (Max temp - Min temp)]	2
tS4	Temperature seasonality (standard deviation X 100)	4
mxtWM5	Max temperature of warmest month	5
mintCM6	Min temperature coldest month	6
mtWQ8	Mean temperature of wettest quarter	8
pAM12	Annual precipitation	12
pDM14	Precipitation of driest month	14
pWQ16	Precipitation of wettest quarter	16
pCQ19	Precipitation of coldest quarter	19
Wind	Mean annual wind speed	N/A

Table 3: *Terrapene* ddRAD SNP outliers (N=118) most strongly associated with ten predictive and uncorrelated WorldClim variables (Table 2), collapsed into temperature, precipitation, or wind speed categories. Percentages include the total for only redundancy analysis (% RDA) and for all three outlier methods (% All): INTROGRESS, Bayesian genomic cline (BGC), and RDA.

Environment Type	No. Outliers	% (RDA)	% (All)
Temperature	44	78.6	34.7
Precipitation	10	17.9	8.5
Wind	2	3.6	1.7
Total	56	100.0	44.9

Figure 1: ADMIXTURE (barplots) and TESS3 (map) analyses for 11,308 *Terrapene* ddRADseq loci. All plots include individuals (N=320; black circles on map) from the Midwest and Southeast hybrid zones, with the optimal number of clusters (*K*) determined via cross-validation across 20 independent runs. Labels above ADMIXTURE plots indicate subspecies (when available) as identified in the field: EA=Woodland (*T. carolina carolina*), ON=Ornate (*T. o. ornata*), TC=*T. carolina* (subspecies identification unavailable), GU=Gulf Coast (*T. c. major*), TT=Three-toed (*T. mexicana triunguis*). Bottom labels show hybrid zone localities by U.S. state: IL=Illinois, AL=Alabama, GA=Georgia, FL=Florida, MS=Mississippi, LA=Louisiana. A * represents a group of "pure" individuals from multiple localities outside the hybrid zones. TESS3 ancestry coefficients (Q) are predicted across the spatial surface via Kriging interpolation (θ=10) and are color-coded with the ADMIXTURE plots. Lighter/ darker gradient shades depict lower/ higher Q.

Figure 2: Population-level NEWHYBRIDS plots for four pairs of southeastern and midwestern *Terrapene* taxa. Individuals were collapsed into populations based on field identification at the subspecific level. The first two characters represent: GU=Gulf Coast, *T. c. major*; EA=Woodland, *T. c. carolina*; TT=Three-toed, *T. m. triunguis*; ON=Ornate, *T. o. ornata*; TC=*T. carolina* (subspecies-level field identification unavailable). The last two characters represent U.S. state: AL=Alabama, FL=Florida, MS=Mississippi, SC=South Carolina, GA=Georgia, LA=Louisiana, IL=Illinois). Each plot corresponds to tests between parental groups (A) EAxGU (N=109), (B) EAxTT (N=135), (C) GUxTT (N=139), and (D) EAxON (N=112). A posterior probability threshold >0.8 was required for genotype frequency class assignments, which included P_1 and P_2 (parental types), F_1 and F_2 (first and second-generation hybrids), backcrosses (B_1 and B_2), and F_N (unclassified).

Figure 3: *Terrapene* Genomic clines depicting outlier SNPs found in transcriptome-aligned ddRAD loci. Pairwise comparisons are between EA=Woodland (*T. c. carolina*), GU=Gulf Coast (*T. c. major*), and TT=Three-toed (*T. mexicana triunguis*) box turtles. The gray area represents neutral expectations based on 2,660 (EAxGU), 2,623 (EAxTT), and 2,622 (GUxTT) transcriptome-aligned SNPs, and each line is a genomic cline for one outlier locus (abbreviations defined in Table 1).

Figure 4: Bayesian genomic cline (BGC) outliers for *Terrapene* ddRADseq SNPs, plotted as heatmaps mapped to *Trachemys scripta* chromosomes. Each chromosome repeats to display significant outliers from the genomic cline center (α; left) and rate (ß; right) BGC parameters. Thinner and thicker bands represent SNPs from unknown scaffolds and annotated genes. BGC was run pairwise for three taxa: EA=*T. carolina carolina*, GU=*T. c. major*, and TT=*T. mexicana triunguis.* Outliers were significant if they had a 95% CI excluding zero or exceeding the probability distribution's quantile interval ($\frac{1-0.975}{2}$, $\frac{0.975}{2}$). The □ plots depict transcriptome-aligned BGC genomic clines with hybrid index histograms above, and each line represents a genomic cline for one locus. The αXß plots illustrate the BGC parameters as a function of density. Polygons define density space for significant α (blue), ß (orange), and both (purple) outliers.

Figure 5: Redundancy analysis (RDA) representing outlier *Terrapene* SNPs correlated with ten predictive, non-redundant BioClim environmental variables (see Table 2 for predictor abbreviations). Significant outliers were designated as being +/- 3 standard deviations from the RDA axis loading means. Pairwise Pearson's correlations between each outlier and environmental variable were performed, and the strongest correlation coefficient (r) determined the best-supported predictor.

Figure 6: Venn Diagram depicting overlap between *Terrapene* transcriptomic outliers identified in INTROGRESS, Bayesian genomic cline (BGC), and redundancy analysis (RDA). Each value includes raw counts (top) and percentages (bottom).