Social bonding induces changes in brain functional connectivity in male and female monogamous voles: a longitudinal fMRI study

M. Fernanda López-Gutiérrez[1], Juan J. Ortiz[1], Francisco J. Camacho[1], Larry J. Young[2], Raúl G. Paredes[1,3] Nestor F. Diaz[4], Wendy Portillo[1] and Sarael Alcauter[1]

[1]Instituto de Neurobiología, Universidad Nacional Autónoma de México, Querétaro, México.
[2]Silvio O Conte Center for Oxytocin and Social Cognition, Center for Translational Social Neuroscience, Yerkes National Primate Research Center, Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, Georgia 30329, United States of America.
[3]Escuela Nacional de Estudios Superiores, Unidad Juriquilla. Querétaro, México.
[4]Instituto Nacional de Perinatología, Isidro Espinosa de los Reyes, México.

Abstract

Microtus ochrogaster, the prairie vole, has become a valuable rodent model to study complex social behaviors, including enduring social bonding. Previous studies have related pair bonding in the prairie vole with plastic changes in several brain regions. However, the network interaction between these regions and their relevance to such socio-sexual behaviors has yet to be described. In this study, we used resting state functional magnetic resonance imaging (rsfMRI) to explore the longitudinal changes in the functional interactions of a set of brain regions previously associated with pair bonding in prairie voles. Specifically, 32 prairie voles were scanned at baseline, after 24 hours and two weeks of cohabitation and mating with a sexual partner (16 male-female pairs). Between 48 and 72 hours of cohabitation voles showed significant preference for the partner over a stranger, confirming pair bond formation; however, individual variability was observed. Network based statistics showed no sex differences in functional connectivity, but revealed a common network with significant longitudinal changes between sessions, including the anterior cingulate cortex (ACC), anterior olfactory nucleus (AON), dentate gyrus (DG), dorsal hippocampus (dHIP), lateral septum (LS), medial prefrontal cortex (mPFC), retrosplenial cortex (RSC), ventral hippocampus (vHIP), and ventral tegmental area (VTA). Furthermore, baseline functional connectivity in three sub-networks including the ACC, nucleus accumbens (NAcc), basolateral amygdala, and DG (ACC-NAcc-BLA-DG), the medial amygdala (MeA)-ventral pallidum (VP), and RSC-VTA, predicted the onset of affiliative behavior (huddling latency) in the first hours of cohabitation, even before sexual experience. Finally, a relationship was found between the strength of the pair bond (partner preference index) with long-term changes in the functional connectivity between the ventral pallidum and medial amygdala (MeA-VP). Overall, our findings revealed the association between network-level changes and the process of social bonding, and provide a novel approach to further investigate the neurophysiology of complex social behaviors displayed in the prairie vole.

Introduction

The prairie vole (*Microtus ochrogaster*) is a rodent native of North America whose natural behavior involves pair bonding, which can be defined as forming an enduring, preferential association between a sexually mature male and female, characterized by selective contact, affiliation, and copulation with the partner over a stranger (Gubernick, 1994). Pair bonded voles will usually display selective aggression towards unfamiliar conspecifics; biparental care, including paternal behavior and alloparenting; incest avoidance and reproductive suppression of adult individuals within a family group (Carter et al., 1995). These behaviors make the prairie vole a valuable model to investigate behaviors associated with a socially monogamous reproductive strategy (K. A. Young et al., 2011), social bond disruption, social isolation and social buffering (Bosch et al., 2016 Lieberwirth & Wang, 2016). Being comparable to human-like social interactions, studying the neurobiology of social behavior in the prairie vole may allow further understanding of human social bonding, its alterations in psychological disorders and overall impact in health (Kiecolt-Glaser et al., 2010).

Pair bonding-related behaviors in the prairie vole depend on hormonal mechanisms and the activation of emotional, reward and sensory brain circuits (Broad et al., 2006; Numan & Young, 2016), integrating large functional networks (Johnson et al., 2016, Kelly & Goodson, 2014). Among these, the mesolimbic reward system and the social decision-making network (SDM) (Lauren A. O'Connell & Hofmann, 2011) are the proposed networks to be involved in pair bonding, being modulated by steroid hormones, dopamine (DA), oxytocin (OXT), arginine vasopressin (AVP), γ-aminobutyric acid (GABA), glutamate, and

corticotropin-releasing factor (CRF), among others (Johnson et al., 2017, Lieberwirth & Wang, 2014; K. A. Young et al., 2011). A current hypothesis suggests that pair bonding consists of two different plastic processes: the formation of a neural representation of the partner, and a selective preference for the partner, i.e. the maintenance of the pair bond (Walum & Young, 2018). In this process, an association has to be created between the reinforcing properties of sex (mating) and the olfactory signature from the partner (Ulloa et al., 2018; L. J. Young & Wang, 2004; L. J. Young et al., 2005). In broad terms, both OXT and AVP are necessary and sufficient for the formation of the pair bond (Cho et al., 1999; Johnson & Young, 2015; L. J. Young & Wang, 2004), and their release during social and sexual interactions are the likely triggers and critical modulators of the mentioned network, since most regions included in the SDM network express their corresponding receptor binding (Johnson et al., 2017; Owen et al., 2013). DA would also be released in concert with OXT and AVP in specific regions to modulate the adequate display of behavior and formation of the pair bond (Wang & Aragona, 2004), including the NAcc, which has substantially more D1-like receptor binding two weeks after female exposure, relative to non–pair bonded males (Aragona et al., 2006). However, the interplay of these broad networks has not been directly explored *in vivo*.

Recently, novel electrophysiologic and optogenetic techniques have been employed to demonstrate that the functional connectivity between the nucleus accumbens (NAcc) and the medial prefrontal cortex (mPFC) during initial cohabitation in female prairie voles modulates the affiliative behavior with their potential partner (Amadei et al., 2017), providing exciting data of the relevance of

such corticostriatal interactions for social bonding. However, this approach does not allow the study of the interaction of multiple brain regions, i.e. networks, and their relevance of such interactions in the process of pair bonding. Neuroimaging methods may provide the alternative to explore such networks in a longitudinal fashion, but only two studies have made use of positron-emission tomography (PET) to explore limited aspects of such longitudinal changes (Bales et al., 2007; Hostetler., 2017), providing the first longitudinal evidence of neurophysiological changes associated to pair bonding. Potentially, functional magnetic resonance imaging (fMRI) may be the ideal tool to explore the longitudinal changes in the functional brain networks (Biswal et al., 1995; Damoiseaux et al., 2006; Smith et al., 2009), providing high spatial resolution, wide brain coverage and being minimally invasive to explore longitudinal changes. In particular, resting state functional magnetic resonance imaging (rsfMRI) explores the low frequency fluctuations (<0.1 Hz) of the blood oxygen level dependent (BOLD) signal, which has proven to remain highly synchronized within the sensory, motor and associative networks in the brain of humans (Biswal et al., 1995; Damoiseaux et al., 2006; Smith et al., 2009), non-human primates (Rilling et al., 2007; Vincent et al., 2007) and rodents (Gozzi & Schwarz, 2016; Jonckers et al., 2011), including the prairie vole (Ortiz et al., 2018), turning this technique into a promising tool for translational research. Indeed, functional connectivity explored by means of the correlation of rsfMRI signals of anatomically separated brain regions (Friston et al., 1993; Biswal et al., 1995) has demonstrated to be correlated with neuronal activity (Mateo et al., 2017), and it is subject to change in plastic processes such as learning and memory in both humans (Jolles et al., 2013) and rodents (Nasrallah et

al., 2016). Here, we make use of this non-invasive technique to explore the longitudinal changes in the brain functional connectivity associated with pair bonding in prairie voles.

Methods

Animals. Thirty-two three-month-old sexually naïve female (*N*=16) and male (*N*=16) prairie voles (*Microtus ochrogaster*) were used in the study. The animals were housed in a temperature (23°C) and light (14:10 light-dark cycle) controlled room and provided with rabbit diet HF-5326 (LabDiet, St. Louis, MO, USA) oat, sunflower seeds, and water *ad libitum*. These voles were previously weaned at 21 days, housed in same-sex cages, and were descendants of voles generously donated by Dr. Larry J. Young from his colony at Emory University. Fourteen days before the experimental protocol, female voles were bilaterally ovariectomized. After recovery, silastic capsules (Dow Corning™ Silastic™ Laboratory Tubing; Thermo Fisher Scientific, Pittsburg, USA) containing estradiol benzoate (B2E; 17β-estradiol; Sigma Aldrich, Missouri, E.E.U.U) dissolved in corn oil (0.5 mg/mL of B2E) were implanted via s.c. to induce sexual receptivity 4 days before cohabitation protocol and remained implanted during the entire experimental protocol. The corresponding dose and procedure reliably induced sexual receptivity (Fowler et al., 2005; Ingberg et al., 2012).

Animals were anesthetized to avoid stress and excessive movement during scanning sessions. Isoflurane at 3% concentration in an oxygen mixture was used for induction and positioning in the scanner bed, in which the head was immobilized with a bite bar and the coil head holder. Once voles were securely

placed in the scanner bed, isoflurane anesthesia was adjusted at a 2% concentration and a single bolus of 0.05 mg/kg of dexmedetomidine (Dexdomitor; Zoetis, Mexico) was administered subcutaneously. Five minutes after the bolus injection, isoflurane anesthesia was lowered and maintained at 0.5%. MRI acquisition started when physiological readings were stabilized (~15 minutes after bolus injection). The use of both anesthetics has been reported as optimal for rsfMRI acquisition in rodents and closely resemble an awake condition (Grandjean et al., 2014, Paasonen et al., 2018), yet the mentioned combination of dose and administration route were previously standardized for this specific protocol in prairie voles (non-published data). Body temperature was maintained with a circulating water heating pad within the scanner bed, respiration rate was monitored with an MR-compatible pneumatic pillow sensor, and blood oxygen saturation was measured with an MR-compatible infrared pulse-oximeter (SA instruments Inc, Stony Brook NY). After the scanning sessions, animals were monitored until fully recovered and transferred back to their housing. All surgical, experimental and maintenance procedures were carried out in accordance with the "Reglamento de la Ley General de Salud en Materia de Investigación para la Salud" (Health General Law on Health Research Regulation) of the Mexican Health Ministry which follows the National Institutes of Health's "Guide for the Care and Use of Laboratory Animals" (NIH Publications No. 8023, revised 1978). The animal research protocols were approved by the bioethics committee of the Instituto de Neurobiología, UNAM.

Figure 1. Sequence of experiments during a 30-day period: Female voles were bilaterally ovariectomized before MR and behavioral protocols. After being allowed to recover from surgery for 10 days, silastic capsules containing E2B (estradiol benzoate) were implanted via s.c. 4 days for sexual receptivity induction before cohabitation. Once couples went under cohabitation, they were housed together for the remaining of the experiment and were only separated for PPT and MR scanning sessions. OVX: ovariectomy surgery. MR: magnetic resonance imaging scanning session. PPT: Partner preference test.

Image acquisition. Prairie voles underwent three MRI acquisition sessions: a baseline scan before cohabitation, a second scan after 24 hours of cohabitation, and a third scan after two weeks of cohabitation. The timeline of the experiment is described in Figure 1. MRI acquisition was conducted with a Bruker Pharmascan 70/16US, 7 Tesla magnetic resonance scanner (Bruker, Ettlingen, Germany), using an MRI CryoProbe transmit/receive surface coil (Bruker, Ettlingen, Germany). Paravision-6 (Bruker, Ettlingen, Germany) was used to perform all imaging protocols. Before running the fMRI sequence, local field homogeneity was optimized within an ellipsoid covering the whole brain and skull using previously acquired field maps. RsfMRI was acquired using a spin-echo echo planar imaging (SE-EPI) sequence: repetition time (TR) = 2000 ms, echo time (TE) = 19 ms, flip angle (FA) = 90°, field of view (FOV) = 18 × 16 mm^2, matrix dimensions = 108 × 96, yielding an in-plane voxel dimensions of 0.167 × 0.167 mm^2, and slice thickness of 0.7 mm, total volumes acquired = 305 (10 minutes and 10 seconds). After the rsfMRI sequence, an anatomical scan was obtained using a spin-echo rapid acquisition with refocused echoes (Turbo-RARE) sequence with the following

parameters: TR = 1800 ms, TE = 38 ms, RARE factor = 16, number of averages (NA) = 2, FOV = 18 × 20 mm^2, matrix dimensions = 144 × 160, slice thickness = 0.125 mm, resulting in isometric voxels of size 0.125 × 0.125 × 0.125 mm^3.

Behavioral testing. *Cohabitation:* the day after the baseline scanning session, female and male voles unrelated to each other were randomly assigned as couples and placed for cohabitation in a new home cage with fresh bedding to promote *ad libitum* mating and social interaction. The first 6 hours of cohabitation were video recorded for subsequent analysis of social and mating behavior. Specifically, mounting, intromission and ejaculation latencies from male voles; lordosis latency from females; and huddling latencies for each male and female pair. Huddling behavior was considered when the pair was sitting or lying in bodily contact for at least 10 seconds. Voles were housed in couples for the remaining of the experiment and were only separated for MRI scanning sessions and behavioral tests.

Partner preference test (PPT): Each subject underwent a 3-hour partner preference test to evaluate pair bond formation. This protocol was based on a previously described test (Williams et al., 1992) and was performed in custom built, 3-chambered clear plastic arenas divided by perforated clear plastic barriers that allowed visual, auditory and olfactory contact, but not physical interaction or mating behavior between subjects. In the central chamber, the vole being tested could roam freely, and time spent in the incentive areas at opposite sides of the chamber was recorded. The incentive areas were defined as the proximal space next to the chambers with its "partner" or with an opposite-sex, novel "stranger" vole. All

stranger voles were unrelated to subjects in the test and had the same age and hormonal condition than the sexual partner. On each test, partner or stranger voles were randomly and alternately positioned on the opposite chambers of the arena. Male and female partners were tested in alternate time periods (48 or 72h), assigned at random, to enable rest between tests and avoid excessive stress. PPT data analysis was performed with *UMATracker* software (Yamanaka & Takeuchi, 2018), which allowed quantification of the proportion of time spent with each of the stimulus voles. A partner preference index was calculated for each subject, consisting on the time with partner divided by the total time with any of the stimulus voles (time with partner plus time with strange). Shapiro-Wilk normality tests revealed that such behavioral data was not normally distributed, therefore, statistical differences were explored with Mann Whitney U tests.

Imaging data preprocessing. Imaging data preprocessing was performed with FMRIB's Software Libraries (FSL, v5.0.9; Jenkinson et al., 2012). To avoid initial signal instability, the first 5 volumes of each functional series were discarded. The, slice-timing correction and motion correction were applied, using the first non-discarded volume as reference. The reference volume was also taken to determine the rigid-body transformation to the corresponding anatomical image. The resulting rigid-body transformation was combined with a non-linear transformation to a prairie vole brain template obtained from previous published work (Ortiz et al., 2018). Functional images were later warped to the brain template and resampled to a resolution of $0.4 \times 0.4 \times 0.4$ mm^3. To minimize physiological confounds, the first 5 eigen-vectors (time-series) within the combined non-grey matter mask were

obtained (Behzadi et al., 2007). These eigen-vectors and the 6 motion parameters (3 rotations, 3 displacements) were regressed out from each subject's functional series. Datasets were band-pass filtered to retain frequencies between 0.01 and 0.1 Hz (Gorges et al., 2017). Finally, smoothing was applied with a gaussian kernel with a full width at half maximum of 0.8 mm.

Functional connectivity analysis. To explore changes in functional connectivity in regions associated with socio-sexual behavior before and after cohabitation, sixteen brain regions of interest (ROI) were defined. Specifically, ROIs were manually defined on the anatomical prairie vole brain template, visually guided with the Allen Mouse Brain Atlas (Lein et al., 2007) to select the minimum possible number of voxels that included/covered each of the following regions: the anterior cingulate cortex (ACC), anterior olfactory nucleus (AON), basolateral amygdala (BLA), bed nucleus of the stria terminalis (BNST), lateral septum (LS), medial amygdala (MeA), main olfactory bulb (MOB), medial prefrontal cortex (mPFC), nucleus accumbens (NAcc), retrosplenial cortex (RSC), paraventricular nucleus (PVN), ventral pallidum (VP), ventral tegmental area (VTA), dentate gyrus (DG), dorsal hippocampus (dHIP) and ventral hippocampus (vHIP) (Johnson & Young, 2017; Lieberwirth & Wang, 2016; Walum & Young, 2018) (Figure 2). For each subject and session, connectivity matrices were calculated as follows: first, the average time series for each region were extracted from the preprocessed fMRI datasets and then, partial correlation estimates were obtained for all possible pairs of ROIs, partialling out the remaining time series. Finally, correlation values were Fisher's z-transformed and further tested to identify connected nodes (ROIs) with

significant longitudinal changes. Due to technical problems, two subjects missed the baseline MRI acquisition (session 1), and two subjects missed the 2 week-cohabitation MRI acquisition (session 3). In order to obtain a balanced sample, missing connectivity data (4 out of 96 expected datasets) were substituted with the average connectivity matrices of the corresponding sex and scanning session. Repeated measures ANOVA tests were performed for two groups (male, female), three sessions (before, 24 hours and two weeks of cohabitation) and sex-group interactions. Sets of connected nodes, i.e. networks, with significant sex, session or interaction differences were identified based on the *Network Based Statistics* (NBS) analysis (Zalesky et al., 2010), which estimates the statistical significance of the identified networks by comparing it with a null distribution obtained with 5000 permutations of the original data (naturally controlling for the size of the original matrices in the permutation tests). In order to identify potential relationships between behavior and functional connectivity, huddling latencies and partner preference indexes were tested as linear covariates of the functional connectivity data in subsequent NBS analyses.

Figure 2. Regions of interest (ROIs) for network functional connectivity analysis. (a) Antero-posterior coronal slices of the prairie vole template overlayed with ROI masks with the resolution used in the analysis. Each color represents a different ROI. ACC: anterior cingulate cortex. AON: anterior olfactory nucleus. BLA: basolateral amygdala. BNST: bed nucleus of the stria terminalis. DG: dentate gyrus. dHIP: dorsal hippocampus. MeA: medial amygdala. MOB: main olfactory bulb. LS: lateral septum. mPFC: medial prefrontal cortex. NAcc: nucleus *accumbens*. PVN: paraventricular nucleus. RSC: retrosplenial cortex. VP: ventral pallidum. vHIP: ventral hippocampus. VTA: ventral tegmental area. **b)** 3D views of ROI maks embedded within the prairie vole template.

Results

Mount (*M±SD*: 65.4±31.7 min), intromission (116±35.3 min), and ejaculation (125±34.4 min) latencies were obtained for male voles (*N*=16). Lordosis latency (22.3±13.3 min) was also measured on females (*N*=16), and huddling latencies (69.5±15.8 min) were obtained for each male and female pair. Three of the sixteen couples did not mate during the recorded period, but all voles displayed huddling and licking/grooming behavior with their sexual partner.

Between 48 and 72 hours of cohabitation, partner preference was evaluated on each subject (see methods). A significant difference was found

between the percentage of time spent on the incentive area related to the partner (*median* = 55.9 %) with the area related to the stranger vole (median = 37.1 %) for all subjects (*U* = 378, *p* = 0.036, *r* = 0.32) (Figure 3). No significant differences were found between males and females in their preference for the partner (*U* = 121, *p* = 0.81, *r* = 0.05) or the stranger voles (*U* = 118, *p* = 0.72, *r* = 0.07), and there were also no significant differences in partner preference between time periods of PPT (48 and 72 hours) (*U* = 119, *p* = 0.75, *r* = 0.06).

Figure 3. PPT analysis: Between 48 and 72h of cohabitation, partner preference was evaluated on each subject (*N* = 32). A significant difference was found between the time spent on the incentive area related to the partner, with the incentive area related to the stranger vole. Boxplot graphs show whiskers with 10-90 percentiles, horizontal line inside the box shows data median and "+" represents data mean. (*) denotes significance at p < 0.05.

Male and female voles share network-level changes related to social bonding

Network Based Statistics analysis yielded significant changes between baseline, 24h and 2 weeks of cohabitation MR sessions (*p* = 0.005) in a network consisting of nine regions: ACC, AON, DG, dHIP, LS, mPFC, RSC, vHIP, and VTA (Figure 4a). No significant differences were found between male or female voles. The nodes with more connections changed were the LS, with four connections, and the

ACC with three. Post-hoc analyses identified differential longitudinal changes among rsfMRI sessions. Specifically, four connections showed significant changes after 24 hours of cohabitation: ACC-vHIP, LS-RSC and DG-dHIP showed increased functional connectivity, while the ACC-VTA showed decreased connectivity in the same period of time (Figure 4a). However, such differences were not shown between session 3 (after 2 weeks of cohabitation) and baseline (before cohabitation), suggesting acute plastic changes related to 24 hours of cohabitation. In addition, four connections showed significant differences between sessions 2 and 3: LS-ACC and LS-AON showed decreased connectivity, but mPFC-LS and mPFC-dHIP showed increased connectivity in session 3, suggesting long-term functional changes related to cohabitation (Figure 4, b-i).

Figure 4. Changes in brain functional connectivity before and after cohabitation with mating in female and male prairie voles: a) Representation of a prairie vole brain with regions (nodes) comprising the brain network that undergoes significant changes in functional connectivity after cohabitation with mating. Interregional connectivity (edges) is shown by color code. Red: increase of functional connectivity; blue: decrease of functional connectivity. ACC: anterior cingulate cortex. AON: anterior olfactory nucleus. DG: dentate gyrus. dHIP: dorsal hippocampus. LS: lateral septum. mPFC: medial prefrontal cortex. RSC: retrosplenial cortex. vHIP: ventral hippocampus. VTA: ventral tegmental area. (**b-i**) Functional Connectivity Residuals (*y axis*) show relative functional connectivity values regressed to control sex variable, displayed in boxplot graphs (25 and 75 percentile) with whiskers (min and max) data points. Connecting lines track longitudinal data of each subject between regions through specific time-points (MR sessions): Baseline, 24h and 2 weeks of cohabitation. Tukey post-hoc significant differences are shown: *<.05, **<.01 ***<.001.

Baseline functional connectivity predicts the display of affiliative behavior

Since behavioral analysis during cohabitation reflected a high level of individual variation, we tested if functional connectivity had a relationship in the display of huddling, and affiliative behavior shown to be socially relevant in the prairie vole and that may influence pair bond induction and formation (Amadei et al., 2017), we assessed the relationship between baseline functional connectivity, i.e. before cohabitation, and huddling latencies during cohabitation in voles of both sexes

(*N*=32). NBS analysis found significant negative linear relations in the following sub-networks: ACC-BLA-NAcc-DG (*p* = 0.007), RSC-VTA (*p* = 0.031), and MeA-VP (*p* = 0.032) (Figure 5). Our results show that the higher the connectivity between these regions before cohabitation, the shorter the huddling latencies during cohabitation in voles of both sexes. Additionally, *a posteriori* Spearman correlation identified the correlation strength between each connected node in the ACC-BLA-NAcc-DG sub-network: BLA-NAcc (r(30) = -0.488, p = 0.005), ACC-NAcc (r(30) = -0.500, p = 0.003), BLA-DG (r(30) = -0.382, p = 0.031) (Figure 5, a-e). These results show that the functional connectivity between these regions reflect the predisposition of the vole to display affiliative behavior, for both females and males.

Figure 5. Relationships between baseline functional connectivity and affiliative behavior (huddling) during cohabitation with mating in male and female prairie voles. Scatterplot graphs (**a-e**) showing significant interregional negative correlations (with best line fit) between baseline functional connectivity and huddling latencies (minutes) during cohabitation. The higher the connectivity between these regions before cohabitation, the shorter the huddling latencies during cohabitation in voles of both sexes. (**f**) Representation of a prairie vole brain with regions (nodes) comprising brain sub-networks that correlate with huddling latency. ACC: anterior cingulate cortex. BLA: basolateral amygdala. DG: dentate gyrus. MeA: medial amygdala. NACC: nucleus *accumbens*. RSC: retrosplenial cortex. VP: ventral pallidum. VTA: ventral tegmental area.

Partner preference measured between 48h and 72h of cohabitation predicts long-term functional connectivity between MeA and VP

Given that several network-level changes were found during cohabitation, we evaluated if functional connectivity could be related to the level of partner preference measured at 48h and 72h of cohabitation. NBS analysis found a significant positive linear relationship between the partner preference index and the MeA-VP connectivity at two weeks of cohabitation (*p*=.027) (Figure 6). This result suggests that the higher the partner preference is between 48h and 72h of cohabitation, the higher the functional connectivity will be between MeA and VP after two weeks of cohabitation.

Figure 6. The level of partner preference at 48h of cohabitation predicts long-term functional connectivity between the medial amygdala and the ventral pallidum. Scatterplot graph with best line fit (**a**) of partner preference index and functional connectivity at 2 weeks of cohabitation, suggesting that partner preference predicts levels of functional connectivity at 2 weeks of cohabitation. (**b**) Representation of a prairie vole brain with regions (nodes) comprising brain sub-networks that correlate with partner preference index. MeA: medial amygdala. VP: ventral pallidum.

Discussion

Several studies have described the relevance of different brain regions involved in pair bond induction and maintenance in prairie voles (Johnson & Young, 2017; Walum & Young, 2018; L. J. Young et al., 2005). However, longitudinal explorations of the brain before and after pair bonding are scarce (Bales et al.,

2007; Hostetler et al., 2016), especially from a network perspective. Here, by using rsfMRI, we were able to detect a brain network with significant longitudinal changes in the functional connectivity of prairie voles after pair bonding. Post-hoc analyses revealed differential short and long term connectivity changes mainly involving ACC, LS and the hippocampal formation. Furthermore, the baseline functional connectivity (before cohabitation), predicted the latency for huddling during the first hours of cohabitation, providing the potential neurofunctional substrate for the variability in affiliative behavior, further extending the recent findings that the corticostriatal electric activity modulates social bonding in prairie voles (Amadei et al., 2017). Finally, partner preference, a potential measure of pair bonding, evaluated at 48 or 72 hours of cohabitation, predicted the long term functional connectivity between the medial amygdala and the ventral pallidum. The connectivity between the latter regions at baseline also predicted affiliative behavior, further highlighting the relevance of such interaction for both initial affiliative behavior and long term pair bonding. We further discuss the potential neurophysiological basis and implications of our findings.

Longitudinal changes in functional connectivity after pair bonding

It has been proposed that pair bonding results from the convergence of the mesolimbic dopamine reward circuit and social discrimination circuits (Walum & Young, 2018). The results here presented are consistent with this model, by demonstrating both short and long-term changes in a brain network including regions largely associated with reward/motivation (ACC, VTA, NAcc) and the social decision making network, specifically related to sensory contextualization (LS,

RSC, dHIP, DG), saliency processing (LS, ACC, VTA), and memory formation and retrieval (vHIP, dHIP, DG, mPFC).

Changes in functional connectivity after 24 hours of cohabitation may result from familiarization of a new spatial context that also implies a novel social context, i.e. exposure to new housing and to a novel, opposite-sex conspecific stranger. During this process, subjects are exposed to novel stimuli and engage in their first socio-sexual interactions, forming new memories of the partner. Furthermore, functional connectivity changes detected at 2 weeks of cohabitation may be related to long-term modulation of behavior as a result of social bonding, in which the partner and its associated cues become salient/rewarding, and after-bonding behaviors such as mate and territorial guarding may appear. In general, these temporally-dynamic functional connectivity changes may be related to the modulation of socio-sexual interactions with the partner. While the nodes here identified are interconnected into a larger network and their precise contribution to complex social behavior remains elusive to our methods, their potential role will be discussed based on previous results and the connectivity patterns here identified.

The functional interactions that showed changes at 24 hours of cohabitation included the ACC and VTA, which may regulate saliency of stimuli. The ACC has been considered an important region in the decision-making process between sensory perception, motivation and final motor performance (Assadi et al., 2009). In voles, the ACC has been implicated in regulating empathy-based consoling behavior towards the partner (Burkett et al., 2016). The VTA is a region known to modulate motivational, reward, and emotionally salient stimuli (Elston & Bilkey, 2017; Gunaydin et al., 2014). There is evidence that the VTA modulates DA

release in prefrontal regions to regulate behaviorally relevant stimuli (Takahata & Moghaddam, 2002), including sexually relevant stimuli (Cortes et al., 2019). Therefore, it is likely that this functional connectivity change may be related to a regulation in sexual motivation as a result of recently-acquired sexual experience. This initial motivation may be necessary to promote further socio-sexual interaction and the likely induction of pair bonding.

Social interaction would necessarily require new memory consolidation, and possibly, formation of the neural representation of the partner. Our data showed functional connectivity changes occurred in the hippocampal formation after 24 hours of cohabitation (ACC-vHIP, DG-dHIP; Figure 4), which has demonstrated to be critical in encoding spatial and mnemonic information in rodents and humans (Kitamura et al., 2017; L. Squire, 2004). The dorsal region of CA2 has been characterized as a hub for sociocognitive memory processing in rodents, specifically for social memory encoding, consolidation, recall (Meira et al., 2018), and social recognition (Hitti & Siegelbaum, 2014; Stevenson & Caldwell, 2014). The dHIP-DG increase in connectivity may be related to partner-memory consolidation, since it has been shown that the DG plays an important role in social recognition (Kesner, 2018). Also, ACC-vHIP activity has been previously associated with the regulation of fear in novel environments and in contextual fear generalization in rodents (Bian et al., 2019). However, the ventral regions of CA1 and CA3 have also shown to contribute in social memory processing (Chiang et al., 2018; S. B. McHugh et al., 2004; Okuyama et al., 2016) and sociability modulation in rodents (Felix-Ortiz & Tye, 2014). Additionally, the ACC is thought to facilitate the integration of new information into existing internal representations, i.e. forming

learning associations (Rolls, 2015), that can motivate and modify future behavior (Bryden et al., 2011; Kolling et al., 2016). Thus, increased ACC-vHIP connectivity after 24 hours of cohabitation might be related to social memory formation, but it may also be related to potentially increased anxiety triggered by a new spatial and social environment.

At 24 hours of cohabitation, an increase of connectivity between the LS and the RSC was also identified. The LS is a structure involved in social recognition and retrieval of relevant social information (Bielsky et al., 2005; Everts & Koolhaas, 1997). In regard to the RSC, it is known to be involved in the processing of spatial memory and cognition in rodents (Mitchell et al., 2018). The increased connectivity between the LS and the RSC, at 24 hours, may be related to processing contextual information and relating social information with the spatial context. It is interesting that after 2 weeks of cohabitation, increased connectivity between these regions does not persist, suggesting that is is important at short term, potentially during memory consolidation.

After 2 weeks of cohabitation, functional connectivity between the mPFC and the dHIP was increased. The mPFC has been proposed as a critical region for remote memory retrieval and for memory consolidation, reportedly relying on the hippocampus (Euston et al., 2012). Thus, their increased interaction could support the integration of new memories into pre-existing networks (Preston & Eichenbaum, 2013). Potentially, the increased connectivity between these regions may be related to the incorporation of the partner to long-term memory representations. Indeed, OXT receptor activation in dHIP promotes the persistence of long-term social recognition memory in mice (Lin et al., 2017).

A long-term increase of connectivity was detected between the LS and the mPFC, and decreases of connectivity were observed between ACC-LS and AON-LS at 2 weeks of cohabitation. As mentioned before, the LS is involved in social recognition and retrieval of relevant social information. In rats, it has been shown that OXT release in the LS is required for the maintenance of social memory, modulated by the relevance of the social stimulus (Lukas et al., 2013). Thus, LS connectivity changes might be related to long-term social recognition and behavior modulation, mediated by afferents from prefrontal structures, i.e. the mPFC and the ACC (ACC-LS-mPFC), and by sensory-related information from the AON (AON-LS). Indeed, the aforementioned ACC may have a role in saliency regulation and in decision-making processes, while the mPFC is proposed to have a role in the initiation, maintenance and modulation of social attachment and behavior in rodents (Dalley et al., 2004; Ko, 2017; Smeltzer et al., 2006; Tanimizu et al., 2017). The latter region is also relevant for tasks in which animals must select new rules based on the available cues in order to appropriately modify ongoing behavior (Kingsbury et al., 2012). On the other hand, AON activation is essential for conspecific recognition and promotes the extraction of relevant olfactory information from the MOB in rats (Oettl et al., 2016), requiring OXT receptor activation to promote stimulus selection and facilitate memory formation (Barlow, 2012). Since AVP and OXT receptor blockade in the LS inhibits, while AVP administration enhances pair bond formation in prairie voles (Liu et al., 2001), and male voles exposed to females have increased Fos-immunoreactive cells in the LS (Wang et al., 1997); it is enticing to propose that the LS is the node that enables the integration of olfactory information, with mnemonic and motivational/saliency

information, which are necessary for conspecific recognition and subsequent modulation of behavior.

Although histological studies have evidenced sex-differentiated neurotransmitter pathways involved in pair bonding such as differential AVP expression and distribution in the BNST, MeA, LS, and VP in male voles (Bamshad et al., 1993; Lim et al., 2004; Wang, 1995) and in hippocampus OXT receptor expression (Rice et al., 2017), the functional connectivity here explored showed no significant differences between sexes, nor significant sex-session interactions. Accordingly, our results have shown modulation of consistent functional networks for both female and male prairie voles. We propose that the long-term functional connectivity changes observed in the network are related to social bonding and may lead to pair bonding induction and maintenance in both female and male prairie voles.

Correlations between sub-network functional connectivity with huddling latencies and Partner Preference Index

Even though it's been reported that in prairie voles 24 hours of cohabitation or 6 hours of *ad libitum* mating are sufficient for a pair bond to be developed (Williams et al., 1992), a considerable amount of evidence has shown that other factors influence its development and maintenance. Specifically, AVP (Okhovat et al., 2015; Ophir et al., 2008) and OXT receptor gene expression and density (King et al., 2016; Olazábal & Young, 2006; Ophir et al., 2012; Perkeybile et al., 2018), paternal nurturing (Ahern & Young, 2009; Arias del Razo y Bales, 2016; Perkeybile et al., 2018) or deprivation (Barrett et al., 2015; Tabbaa et al., 2017), have shown

to produce variability in the exhibition of prairie vole social behavior. Subjects of both sexes in this study showed a wide variability in their bonding behavior evaluated between 48 and 72h of cohabitation, even though they were under the same experimental conditions. It is likely that the sum of previously mentioned factors gives each subject a distinctive brain network configuration that ultimately relates to bonding behavior. Thence, we hypothesized there may be individual differences in functional connectivity that could explain the variability in socio-sexual behavior. Indeed, we identified three functional networks for which the functional connectivity at baseline was negatively related to huddling latencies during the first hours of cohabitation. In other words, baseline functional connectivity predicts how quickly subjects would begin affiliative huddling with an opposite sex conspecific. Huddling is a measurable affiliative behavior in prairie voles and a useful indicator of social receptiveness (Beery et al., 2018; Salo et al., 1993). The larger network included four nodes, related as follows: ACC-NAcc-BLA-DG (Figure 5), while the other two only included two nodes: MeA-VP and VTA-RSC. These subnetworks involve regions reported to have a role in social memory and recognition, spatial, and reward-seeking mechanisms. Specifically, BLA and the ACC have been found relevant in coordinating brain activity when social interaction is initiated and in the formation of social recognition memory through gene expression (Garrido Zinn et al., 2016; Tanimizu et al., 2017). In line with our results, the BLA may act as an associative site for stimulus-outcome representations (Cardinal et al., 2002), allowing an appropriate response according to previous social encounters, and may require mnemonic information encoded by the DG (BLA-DG). Part of a social interaction, once a conspecific has been

recognized, involves a reward component. The NAcc is known to translate reward-predictive information from the amygdala (BLA-NAcc) to promote cue-evoked, reward-seeking behavioral responses in rodents (Ambroggi et al., 2008; Stuber et al., 2011); additional input from the ACC (ACC-NAcc) would be necessary for a social decision-making process and motor performance (Seo & Lee, 2012; Zhong et al., 2017). Furthermore, a recent study showed that in female prairie voles, the functional connectivity of the prefrontal cortex and NAcc after the first encounter predicts affiliative huddling towards a partner, and the activation of such circuit biases later preference towards a partner (Amadei et al., 2017). Our results extend such findings, showing that a larger network including similar corticostriatal connectivity, measured even before the exposure to a potential partner, predicts affiliative behavior. Moreover, such relation is consistent for both male and females, and the circuit includes the amygdala and hippocampus, as predicted by Amadei et al., (2017).

In addition, two other circuits also predicted future affiliative behavior, the VTA-RSC and MeA-VP (Figure 5). The former may indicate a potential bias for a more efficient integration of socially-rewarding stimuli with contextual information of the partner (Hung et al., 2017; Todd and Bucci, 2015). While in the latter, it is known the VP plays a major role in reward and motivation (Smith et al., 2008), encoding reward value earlier and more robustly than the NAcc and potentially being a crucial downstream mediator of NAcc reward-related functions (Ottenheimer et al., 2018), including pair bonding (M. M. Lim & Young, 2004). Since MeA activity in rodents is necessary for social recognition (Ferguson et al., 2001) and responds to sex-specific chemosensory cues (Wang et al., 1997; Yao et

al., 2017), increased MeA-VP connectivity may improve the rewarding response of social and chemosensory cues from the partner.

Moreover, the MeA-VP connectivity after 2 weeks of cohabitation was also predicted by the Partner Preference Index at 48 and 72 hours, which captures the strength of a pair bond (Williams et al., 1992). This relation is not surprising given the recent evidence that the MeA synchronizes socio-sexual behavior by the enhancement of preference for a sexual partner (Adekunbi et al., 2018). Meanwhile, the VP is related to the hedonic or motivational impact of the partner-related stimuli (Smith et al., 2008). The identified relation suggests the strengthening of this sub-network (MeA-VP) along 2 weeks of cohabitation in voles with higher partner preference between 48 and 72 hours of cohabitation (Figure 6).

We found particularly interesting that the NAcc, MeA, VP and BLA, although deemed relevant for social bonding, are not nodes of the main network with longitudinal changes across rsfMRI sessions. However, the connectivity of these regions actually relates to the affiliative behavior of the prairie voles, highlighting the role of their neural circuits in the variability of prairie vole social behavior, i.e. "wanderer" or "resident" mating tactics; in which resident males maximize mating success through mate guarding and reduced territory space, in contrast to wanderers that overlap many home ranges to increase opportunistic mating (Getz et al., 2005; Shuster et al., 2019). This is supported by previous work demonstrating that AVP receptor expression in the RSC predicts sexual fidelity and territorial behavior in male prairie voles (Okhovat et al., 2015; Ophir et al., 2008). In addition, AVP receptor overexpression in VP induced partner preference in socially promiscuous meadow voles (Lim, Wang, et al., 2004), and variation in OXT

receptor density in the NAcc relates to partner preference in prairie voles (King et al., 2016). Consequently, sub-network functional connectivity variability may also reflect behavioral diversity influenced by genetic and environmental factors.

The prairie vole has proven to be a valuable model that has enabled the characterization of the neurobiological mechanisms behind complex socio-sexual behaviors, potentially useful to understand human social bonding and its alterations in psychological disorders. To our knowledge, this is the first study to demonstrate both short and long-term changes in the functional connectivity of multiple interacting brain regions (networks) of prairie voles in cohabitation with mating. However, there are some limitations related to the use of rsfMRI. First, the animals were scanned under anesthesia, which may potentially alter the brain functional connectivity. Yet, we have used an anesthesia protocol that practically preserves the functional interactions as in the awake state (Grandjean et al., 2014, Paasonen et al., 2018). Although functional neuroimaging of awake prairie voles is possible, the acclimation process may induce significant stress and even increase the risk of physical harm (Yee et al., 2016), limiting the longitudinal design and the interpretation of the results. Second, the precision in the definition of the ROIs is limited by the shape and the size of the voxels, being hard to assure that the defined regions specifically and uniquely include signal from the anatomical regions of interest. Nevertheless, we are certain that the defined ROIs include the actual regions of interest, and at the least, the changes here reported are related to those areas and their surrounding tissue. It is important to note that functionally connected nodes may not necessarily have direct axonal projections to each other, and the interaction between nodes could be mediated or relayed through other

structures (Friston et al., 1993). Also, contrary to other electrophysiological or neuropharmacological methods, functional MRI captures an indirect measure of neuronal activity (Kim & Ogawa, 2012; Ogawa et al., 1990). However, it poses the great advantage that it allows the longitudinal exploration of the brain, with the best spatial resolution and wider coverage that any imaging method can achieve non-invasively. The consistency of our results with recent findings using direct electrophysiology readings (Amadei et al., 2017) further support their relevance and its potential to contribute to identify the neurophysiology of complex social behaviors.

Conclusions

Our findings suggest there is a brain network that encompasses short- and long-term changes in the socio-sexual behavior of prairie voles as a consequence of social bonding and mating during cohabitation. Such functional connectivity changes may be involved in the process of pair bonding and in the formation and consolidation of the neuronal representation of the partner. Furthermore, the functional connectivity of specific subnetworks including the ACC, NAcc, BLA, DG, MeA, VP, RSC and VTA, predicts huddling latencies, suggesting there is a neurobiological predisposition to social bonding, while another network (MeA-VP) correlates with the level of early partner preference. In summary, our findings suggest: 1) baseline functional connectivity predicts the level of affiliative behavior even before sexual experience and, 2) sexual experience and long-term cohabitation induces network-wide changes in socio-sexual relevant circuits. Functional connectivity may also aid in exploring the mechanisms that underlie

individual variability in the expression of socio-sexual behavior. Overall, our findings have shown network-level changes associated with the process of social bonding and provide a novel approach to further investigate the neurophysiology of complex social behaviors displayed in the prairie vole.

Acknowledgements

We thank Deisy Gasca, Martín García, Alejandra Castilla, Leopoldo González Santos, Nuri Aranda López and Ma. De Lourdes Lara for their technical assistance. The authors thankfully acknowledge the imaging resources and support provided by the Laboratorio Nacional de Imagenología por Resonancia Magnética (LANIREM), part of the CONACYT's network of national laboratories. This research was supported by grants CONACYT 252756 to WP, 253631 to RGP, UNAM-DGAPA-PAPIIT IN202818 to WP, IN212219-3 to SA, IN203518-3 to RGP, INPER 2018-1-163 to NFD. LJY's contribution was supported by NIH grants P50MH100023 to LJY and P51OD011132 to YNPRC. M.F. López-Gutiérrez was supported by a fellwoship from CONACYT (fellwoship #626152) for her Masters of Science studies in Neurobiology at UNAM (Maestría en Ciencias [Neurobiología], UNAM).

References

Adekunbi, D. A., Li, X. F., Lass, G., Shetty, K., Adegoke, O. A., Yeo, S. H., … O'Byrne, K. T. (2018). Kisspeptin neurones in the posterodorsal medial amygdala modulate sexual partner preference and anxiety in male mice. Journal of Neuroendocrinology, 30(3), 1–9.

Ahern, T. H., & Young, L. J. (2009). The impact of early life family structure on adult social attachment, alloparental behavior, and the neuropeptide systems regulating affiliative behaviors in the monogamous prairie vole (Microtus ochrogaster). Frontiers in Behavioral Neuroscience, 3, 17.

Amadei, E. A., Johnson, Z. V., Jun Kwon, Y., Shpiner, A. C., Saravanan, V., Mays, W. D., … Liu, R. C. (2017). Dynamic corticostriatal activity biases social bonding in monogamous female prairie voles. Nature, 546(7657), 297–301.

Ambroggi, F., Ishikawa, A., Fields, H. L., & Nicola, S. M. (2008). Basolateral Amygdala Neurons Facilitate Reward-Seeking Behavior by Exciting Nucleus Accumbens Neurons. Neuron, 59(4), 648–661.

Aragona, B. J., Liu, Y., Yu, Y. J., Curtis, J. T., Detwiler, J. M., Insel, T. R., & Wang, Z. (2006). Nucleus accumbens dopamine differentially mediates the formation and maintenance of monogamous pair bonds. Nat Neurosci, 9(1), 133–139.

Arias del Razo, R., & Bales, K. L. (2016). Exploration in a dispersal task: Effects of early experience and correlation with other behaviors in prairie voles (Microtus ochrogaster). Behavioural Processes, 132, 66–75.

Assadi, S. M., Yücel, M., & Pantelis, C. (2009). Dopamine modulates neural networks involved in effort-based decision-making. Neuroscience and Biobehavioral Reviews, 33(3), 383–393.

Bales, K. L., Mason, W. A., Catana, C., Cherry, S. R., & Mendoza, S. P. (2007). Neural correlates of pair-bonding in a monogamous primate. Brain Research, 1184, 245–253.

Bamshad, M., Novak, M. A., & Vries, G. J. (1993). Sex and Species Differences in the Vasopressin Innervation of Sexually Naive and Parental Prairie Voles,

Microtus ochrogaster and Meadow Voles, Microtus pennsylvanicus. Journal of Neuroendocrinology, 5(3), 247–255.

Barlow, H. B. (2012). Possible Principles Underlying the Transformations of Sensory Messages. En Sensory Communication (pp. 216–234).

Barrett, C. E., Arambula, S. E., & Young, L. J. (2015). The oxytocin system promotes resilience to the effects of neonatal isolation on adult social attachment in female prairie voles. Translational psychiatry, 5(7), e606.

Beery, A. K., Christensen, J. D., Lee, N. S., & Blandino, K. L. (2018). Specificity in Sociality: Mice and Prairie Voles Exhibit Different Patterns of Peer Affiliation. Frontiers in behavioral neuroscience, 12, 50.

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. NeuroImage, 37(1), 90–101.

Bian, X.-L., Qin, C., Cai, C.-Y., Zhou, Y., Tao, Y., Lin, Y.-H., … Zhu, D.-Y. (2019). Anterior Cingulate Cortex to Ventral Hippocampus Circuit Mediates Contextual Fear Generalization. The Journal of neuroscience : the official journal of the Society for Neuroscience, 39(29), 5728–5739.

Bielsky, I. F., Hu, S. B., Ren, X., Terwilliger, E. F., & Young, L. J. (2005). The V1a vasopressin receptor is necessary and sufficient for normal social recognition: A gene replacement study. Neuron, 47(4), 503–513.

Biswal, B., Yetkin, F. Z., Haughton, V. M., & Hyde, J. S. (1995). Functional Connectivity in the Motor Cortex of Resting Human Brain Using Echo-Planar MRI.

Bosch, O. J., Dabrowska, J., Modi, M. E., Johnson, Z. V, Keebaugh, A. C., Barrett,

C. E., … Young, L. J. (2016). Oxytocin in the nucleus accumbens shell reverses CRFR2-evoked passive stress-coping after partner loss in monogamous male prairie voles. Psychoneuroendocrinology, 64, 66–78.

Broad, K. ., Curley, J. ., & Keverne, E. . (2006). Mother-infant bonding and the evolution of mammalian social relationships. Philosophical Transactions of the Royal Society B: Biological Sciences, 361(1476), 2199–2214.

Bryden, D. W., Johnson, E. E., Tobia, S. C., Kashtelyan, V., & Roesch, M. R. (2011). Attention for Learning Signals in Anterior Cingulate Cortex. Journal of Neuroscience, 31(50), 18266–18274.

Burkett, J. P., Andari, E., Johnson, Z. V, Curry, D. C., De Waal, F. B. M., & Young, L. J. (2016). Oxytocin-dependent consolation behavior in rodents. Science, 351(6271), 375–378.

Cardinal, R. N., Parkinson, J. A., Hall, J., & Everitt, B. J. (2002). Emotion and motivation: The role of the amygdala, ventral striatum, and prefrontal cortex. Neuroscience and Biobehavioral Reviews, 26(3), 321–352.

Carter, C. S., DeVries, A. C., & Getz, L. L. (1995). Physiological substrates of mammalian monogamy: The prairie vole model. Neuroscience and Biobehavioral Reviews, 19(2), 303–314.

Chiang, M. C., Huang, A. J. Y., Wintzer, M. E., Ohshima, T., & McHugh, T. J. (2018). A role for CA3 in social recognition memory. Behavioural Brain Research, 354, 22–30.

Cho, M. M., DeVries, A. C., Williams, J. R., & Carter, C. S. (1999). The effects of oxytocin and vasopressin on partner preferences in male and female prairie voles (Microtus ochrogaster). Behavioral Neuroscience, 113(5), 1071–1079.

Cortes, P. M., Hernández-Arteaga, E., Sotelo-Tapia, C., Guevara, M. A., Medina, A. C., & Hernández-González, M. (2019). Effects of inactivation of the ventral tegmental area on prefronto-accumbens activity and sexual motivation in male rats. Physiology & Behavior, 209(March), 112593.

Dalley, J. W., Cardinal, R. N., & Robbins, T. W. (2004). Prefrontal executive and cognitive functions in rodents: Neural and neurochemical substrates. Neuroscience and Biobehavioral Reviews, 28(7), 771–784.

Damoiseaux, J. S., Rombouts, S. A. R. B., Barkhof, F., Scheltens, P., Stam, C. J., Smith, S. M., & Beckmann, C. F. (2006). Consistent resting-state networks across healthy subjects. Proceedings of the National Academy of Sciences of the United States of America, 103(37), 13848–13853.

Elston, T. W., & Bilkey, D. K. (2017). Anterior Cingulate Cortex Modulation of the Ventral Tegmental Area in an Effort Task. Cell Reports, 19(11), 2220–2230.

Euston, D. R., Gruber, A. J., & McNaughton, B. L. (2012). The Role of Medial Prefrontal Cortex in Memory and Decision Making. Neuron, 76(6), 1057–1070.

Everts, H. G., & Koolhaas, J. M. (1997). Lateral septal vasopressin in rats: role in social and object recognition? Brain research, 760(1–2), 1–7.

Felix-Ortiz, A. C., & Tye, K. M. (2014). Amygdala Inputs to the Ventral Hippocampus Bidirectionally Modulate Social Behavior. Journal of Neuroscience, 34(2), 586–595.

Ferguson, J. N., Aldag, J. M., Insel, T. R., & Young, L. J. (2001). Oxytocin in the medial amygdala is essential for social recognition in the mouse. The Journal of neuroscience: the official journal of the Society for Neuroscience, 21(20), 8278–8285.

Fowler, C. D., Johnson, F., & Wang, Z. (2005). Estrogen Regulation of Cell Proliferation and Distribution of Estrogen Receptor-α in the Brains of Adult Female Prairie and Meadow Voles. J Comp Neurol, 489(2), 166–179.

Friston, K. J., Frith, C. D., Liddle, P. F., & Frackowiak, R. S. J. (1993). Functional connectivity: The principal-component analysis of large (PET) data sets. Journal of Cerebral Blood Flow and Metabolism, 13(1), 5–14.

Garrido Zinn, C., Clairis, N., Silva Cavalcante, L. E., Furini, C. R. G., de Carvalho Myskiw, J., & Izquierdo, I. (2016). Major neurotransmitter systems in dorsal hippocampus and basolateral amygdala control social recognition memory. Proceedings of the National Academy of Sciences, 113(33), E4914–E4919.

Getz, L. L., McGuire, B., & Carter, C. S. (2005). Social organization and mating system of free-living prairie voles Microtus ochrogaster: a review. Acta Zoologica Sinica, 51(2), 178–186.

Gorges, M., Roselli, F., Müller, H. P., Ludolph, A. C., Rasche, V., & Kassubek, J. (2017). Functional connectivity mapping in the animal model: Principles and applications of resting-state fMRI. Frontiers in Neurology, 8(MAY), 1–14.

Gozzi, A., & Schwarz, A. J. (2016). Large-scale functional connectivity networks in the rodent brain. NeuroImage, 127, 496–509.

Grandjean, J., Schroeter, A., Batata, I., & Rudin, M. (2014). Optimization of anesthesia protocol for resting-state fMRI in mice based on differential effects of anesthetics on functional connectivity patterns. NeuroImage, 102(P2), 838–847.

Gubernick, D. J. (1994). Infanticide and parental care (S. Parmigiani & F. S. Vom Saal, Eds.).

Gunaydin, L. A., Grosenick, L., Finkelstein, J. C., Kauvar, I. V., Fenno, L. E., Adhikari, A., … Deisseroth, K. (2014). Natural neural projection dynamics underlying social behavior. Cell, 157(7), 1535–1551.

Hitti, F. L., & Siegelbaum, S. A. (2014). The hippocampal CA2 region is essential for social memory. Nature, 508(1), 88–92.

Hostetler, C. M., Hinde, K., Maninger, N., Mendoza, S. P., Mason, W. A., Rowland, D. J., … Bales, K. L. (2017). Effects of pair bonding on dopamine D1 receptors in monogamous male titi monkeys (Callicebus cupreus). American journal of primatology, 79(3), 1–9.

Hung, L. W., Neuner, S., Polepalli, J. S., Beier, K. T., Wright, M., Walsh, J. J., … Malenka, R. C. (2017). Gating of social reward by oxytocin in the ventral tegmental area. Science, 357(6358), 1406–1411.

Ingberg, E., Theodorsson, A., Theodorsson, E., & Strom, J. O. (2012). Methods for long-term 17β-estradiol administration to mice. General and Comparative Endocrinology, 175(1), 188–193.

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL. NeuroImage, 62(2), 782–790.

Johnson, Z. V., Walum, H., Jamal, Y. A., Xiao, Y., Keebaugh, A. C., Inoue, K., & Young, L. J. (2016). Central oxytocin receptors mediate mating-induced partner preferences and enhance correlated activation across forebrain nuclei in male prairie voles. Hormones and Behavior, 79, 8–17.

Johnson, Z. V., Walum, H., Xiao, Y., Riefkohl, P. C., & Young, L. J. (2017). Oxytocin receptors modulate a social salience neural network in male prairie voles. Hormones and Behavior, 87, 16–24.

Johnson, Z. V., & Young, L. J. (2015). Neurobiological mechanisms of social attachment and pair bonding. Current Opinion in Behavioral Sciences, Vol. 3, pp. 38–44.

Johnson, Z. V., & Young, L. J. (2017). Oxytocin and vasopressin neural networks: Implications for social behavioral diversity and translational neuroscience. Neuroscience and Biobehavioral Reviews, 76, 87–98.

Jolles, D. D., Van Buchem, M. A., Crone, E. A., & Rombouts, S. A. R. B. (2013). Functional brain connectivity at rest changes after working memory training. Human Brain Mapping, 34(2), 396–406.

Jonckers, E., Van Audekerke, J., De Visscher, G., Van der Linden, A., & Verhoye, M. (2011). Functional Connectivity fMRI of the Rodent Brain: Comparison of Functional Connectivity Networks in Rat and Mouse. PLoS ONE, 6(4), e18876.

Kelly, A. M., & Goodson, J. L. (2014). Social functions of individual vasopressin-oxytocin cell groups in vertebrates: What do we really know? Frontiers in Neuroendocrinology, 35(4), 512–529.

Kesner, R. P. (2018, noviembre 15). An analysis of dentate gyrus function (an update). Behavioural Brain Research, Vol. 354, pp. 84–91.

Kiecolt-Glaser, J. K., Gouin, J. P., & Hantsoo, L. (2010). Close relationships, inflammation, and health. Neuroscience and Biobehavioral Reviews, 35(1), 33–38.

Kim, Seong-Gi., & Ogawa, S. (2012). Biophysical and physiological origins of blood oxygenation level-dependent fMRI signals. Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism, 32(7), 1188–1206.

King, L. B., Walum, H., Inoue, K., Eyrich, N. W., & Young, L. J. (2016). Variation in the Oxytocin Receptor Gene Predicts Brain Region–Specific Expression and Social Attachment. Biological Psychiatry, 80(2), 160–169.

Kingsbury, M. A., Gleason, E. D., Ophir, A. G., Phelps, S. M., Young, L. J., & Marler, C. A. (2012). Monogamous and promiscuous rodent species exhibit discrete variation in the size of the medial prefrontal cortex. Brain, Behavior and Evolution, 80(1), 4–14.

Kitamura, T., Ogawa, S. K., Roy, D. S., Okuyama, T., Morrissey, M. D., Smith, L. M., … Tonegawa, S. (2017). Engrams and circuits crucial for systems consolidation of a memory. Science, 356(6333), 73–78.

Ko, J. (2017). Neuroanatomical Substrates of Rodent Social Behavior: The Medial Prefrontal Cortex and Its Projection Patterns. Frontiers in Neural Circuits, 11(June), 1–16.

Kolling, N., Behrens, T., Wittmann, M., & Rushworth, M. (2016). Multiple signals in anterior cingulate cortex. Current Opinion in Neurobiology, 37, 36–43.

Lein, E. S., Hawrylycz, M. J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., … Jones, A. R. (2007). Genome-wide atlas of gene expression in the adult mouse brain. Nature, 445(7124), 168–176.

Lieberwirth, C., & Wang, Z. (2014). Social bonding: regulation by neuropeptides. Frontiers in neuroscience, 8(8 JUN), 171.

Lieberwirth, C., & Wang, Z. (2016). The neurobiology of pair bond formation, bond disruption, and social buffering. Current Opinion in Neurobiology, 40, 8–13.

Lim, Miranda M., & Young, L. J. (2004). Vasopressin-dependent neural circuits underlying pair bond formation in the monogamous prairie vole. Neuroscience,

125(1), 35–45.

Lim, Miranda M., Murphy, A. Z., & Young, L. J. (2004). Ventral Striatopallidal Oxytocin and Vasopressin V1a Receptors in the Monogamous Prairie Vole (Microtus ochrogaster). Journal of Comparative Neurology, 468(4), 555–570.

Lim, Miranda M., Wang, Z. Z., Olazábal, D. E., Ren, X., Terwilliger, E. F., & Young, L. J. (2004). Enhanced partner preference in a promiscuous species by manipulating the expression of a single gene. Nature, 429(6993), 754–757.

Lin, Yu-Ting., Hsieh, Tsan-Yu., Tsai, Tsun-Chih C., Chen, Chien-Chung C., Huang, Chiung-Chun C., & Hsu, Kuei-Sen (2017). Conditional Deletion of Hippocampal CA2/CA3a Oxytocin Receptors Impairs the Persistence of Long-Term Social Recognition Memory in Mice. The Journal of Neuroscience, 38(5), 1218–1231.

Liu, Y., Curtis, J. T., & Wang, Z. (2001). Vasopressin in the lateral septum regulates pair bond formation in male prairie voles (Microtus ochrogaster). Behavioral Neuroscience, 115(4), 910–919.

Lukas, M., Toth, I., Veenema, A. H., & Neumann, I. D. (2013). Oxytocin mediates rodent social memory within the lateral septum and the medial amygdala depending on the relevance of the social stimulus: Male juvenile versus female adult conspecifics. Psychoneuroendocrinology, 38(6), 916–926.

Mateo, C., Knutsen, P. M., Tsai, P. S., Shih, A. Y., & Kleinfeld, D. (2017). Entrainment of Arteriole Vasomotor Fluctuations by Neural Activity Is a Basis of Blood-Oxygenation-Level-Dependent "Resting-State" Connectivity. Neuron, 1–13.

McHugh, S. B., Deacon, R. M. J., Rawlins, J. N. P., & Bannerman, D. M. (2004). Amygdala and Ventral Hippocampus Contribute Differentially to Mechanisms of

Fear and Anxiety. Behavioral Neuroscience, 118(1), 63–78.

Meira, T., Leroy, F., Buss, E. W., Oliva, A., Park, J., & Siegelbaum, S. A. (2018). A hippocampal circuit linking dorsal CA2 to ventral CA1 critical for social memory dynamics. Nature Communications, 9(1), 4163.

Mitchell, A. S., Czajkowski, R., Zhang, N., Jeffery, K., & Nelson, A. J. D. (2018). Retrosplenial cortex and its role in spatial cognition. Brain and Neuroscience Advances, 2, 239821281875709.

Nasrallah, F. A., To, X. V., Chen, D. Y., Routtenberg, A., & Chuang, K. H. (2016). Functional connectivity MRI tracks memory networks after maze learning in rodents. NeuroImage, 127, 196–202.

Numan, M., & Young, L. J. (2016). Neural mechanisms of mother-infant bonding and pair bonding: Similarities, differences, and broader implications. Hormones and behavior, 77, 98–112.

O'Connell, L. A., & Hofmann, H. A. (2011). Genes, hormones, and circuits: An integrative approach to study the evolution of social behavior. Frontiers in Neuroendocrinology, 32(3), 320–335.

Oettl, L. L., Ravi, N., Schneider, M., Scheller, M. F., Schneider, P., Mitre, M., … Kelsch, W. (2016). Oxytocin Enhances Social Recognition by Modulating Cortical Control of Early Olfactory Processing. Neuron, 90(3), 609–621.

Ogawa, S., Lee, T. M., Kay, A. R., & Tank, D. W. (1990). Brain magnetic resonance imaging with contrast dependent on blood oxygenation. Proceedings of the National Academy of Sciences of the United States of America, 87(24), 9868–9872.

Okhovat, M., Berrio, A., Wallace, G., Ophir, A. G., & Phelps, S. M. (2015). Sexual

fidelity trade-offs promote regulatory variation in the prairie vole brain. Science, 350(6266), 1371–1374.

Okuyama, T., Kitamura, T., Roy, D. S., Itohara, S., & Tonegawa, S. (2016). Ventral CA1 neurons store social memory. Science, 353(6307), 1536–1541.

Olazábal, D. E., & Young, L. J. (2006). Oxytocin receptors in the nucleus accumbens facilitate "spontaneous" maternal behavior in adult female prairie voles. Neuroscience, 141(2), 559–568.

Ophir, A. G., Gessel, A., Zheng, D. J., & Phelps, S. M. (2012). Oxytocin receptor density is associated with male mating tactics and social monogamy. Hormones and Behavior, 61(3), 445–453.

Ophir, A. G., Wolff, J. O., & Phelps, S. M. (2008). Variation in neural V1aR predicts sexual fidelity and space use among male prairie voles in semi-natural settings. Proceedings of the National Academy of Sciences of the United States of America, 105(4), 1249–1254.

Ortiz, J. J., Portillo, W., Paredes, R. G., Young, L. J., & Alcauter, S. (2018). Resting state brain networks in the prairie vole. Scientific Reports, 8(1), 1231.

Ottenheimer, D., Richard, J. M., & Janak, P. H. (2018). Ventral pallidum encodes relative reward value earlier and more robustly than nucleus accumbens. Nature Communications, 9(1), 4350.

Owen, S. F., Tuncdemir, S. N., Bader, P. L., Tirko, N. N., Fishell, G., & Tsien, R. W. (2013). Oxytocin enhances hippocampal spike transmission by modulating fast-spiking interneurons. Nature, 500(7463), 458–462.

Paasonen, J., Stenroos, P., Salo, R. A., Kiviniemi, V., & Gröhn, O. (2018). Functional connectivity under six anesthesia protocols and the awake condition

in rat brain. NeuroImage, 172(October 2017), 9–20.

Perkeybile, A. M., Carter, C. S., Wroblewski, K. L., Puglia, M. H., Kenkel, W. M., Lillard, T. S., … Connelly, J. J. (2018). Early nurture epigenetically tunes the oxytocin receptor. Psychoneuroendocrinology, 99, 128–136.

Preston, A. R., & Eichenbaum, H. (2013). Interplay of Hippocampus and Prefrontal Cortex in Memory. Current Biology, 23(17), R764–R773.

Rice, M. A., Hobbs, L. E., Wallace, K. J., & Ophir, A. G. (2017). Cryptic sexual dimorphism in spatial memory and hippocampal oxytocin receptors in prairie voles (Microtus ochrogaster). Hormones and Behavior, 95(August), 94–102.

Rilling, J. K., Barks, S. K., Parr, L. A., Preuss, T. M., Faber, T. L., Pagnoni, G., … Votaw, J. R. (2007). A comparison of resting-state brain activity in humans and chimpanzees. Proceedings of the National Academy of Sciences, 104(43), 17146–17151.

Rolls, E. T. (2015). Limbic systems for emotion and for memory, but no single limbic system. Cortex, 62, 119–157.

Salo, A. L., Shapiro, L. E., & Dewsbury, D. A. (1993). Affiliative Behavior in Different Species of Voles (Microtus). Psychological Reports, 72(1), 316–318.

Seo, H., & Lee, D. (2012). Neural basis of learning and preference during social decision-making. Current opinion in neurobiology, 22(6), 990–995.

Shuster, S. M., Willen, R. M., Keane, B., & Solomon, N. G. (2019). Alternative Mating Tactics in Socially Monogamous Prairie Voles, Microtus ochrogaster. Frontiers in Ecology and Evolution, 7, 7.

Smeltzer, M. D., Curtis, J. T., Aragona, B. J., & Wang, Z. (2006). Dopamine, oxytocin, and vasopressin receptor binding in the medial prefrontal cortex of

monogamous and promiscuous voles. Neuroscience Letters, 394(2), 146–151.

Smith, K. S., Tindell, A. J., Aldridge, J. W., & Berridge, K. C. (2008). Ventral pallidum roles in reward and motivation. Behavioural brain research, 196(2), 155–167.

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., … Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during activation and rest. Proceedings of the National Academy of Sciences of the United States of America, 106(31), 13040–13045.

Squire, L. (2004). Memory systems of the brain: a brief history and current perspective. Neurobiology of learning and memory, 82(3), 171–177.

Stevenson, E. L., & Caldwell, H. K. (2014). Lesions to the CA2 region of the hippocampus impair social memory in mice. European Journal of Neuroscience, 40(9), 3294–3301.

Stuber, G. D., Sparta, D. R., Stamatakis, A. M., van Leeuwen, W. A., Hardjoprajitno, J. E., Cho, S., … Bonci, A. (2011). Excitatory transmission from the amygdala to nucleus accumbens facilitates reward seeking. Nature, 475(7356), 377–380.

Tabbaa, M., Lei, K., Liu, Y., & Wang, Z. (2017). Paternal deprivation affects social behaviors and neurochemical systems in the offspring of socially monogamous prairie voles. Neuroscience, 343, 284–297.

Takahata, R., & Moghaddam, B. (2002). Target-Specific Glutamatergic Regulation of Dopamine Neurons in the Ventral Tegmental Area. Journal of Neurochemistry, 75(4), 1775–1778.

Tanimizu, T., Kenney, J. W., Okano, E., Kadoma, K., Frankland, P. W., & Kida, S.

(2017). Functional Connectivity of Multiple Brain Regions Required for the Consolidation of Social Recognition Memory. The Journal of Neuroscience, 37(15), 4103–4116.

Todd, T. P., & Bucci, D. J. (2015). Retrosplenial Cortex and Long-Term Memory: Molecules to Behavior. Neural Plasticity, 2015, 1–9.

Ulloa, M., Portillo, W., Díaz, N. F., Young, L. J., Camacho, F. J., Rodríguez, V. M., & Paredes, R. G. (2018). Mating and social exposure induces an opioid-dependent conditioned place preference in male but not in female prairie voles (Microtus ochrogaster). Hormones and Behavior, 97(October 2017), 47–55.

Vincent, J. L., Patel, G. H., Fox, M. D., Snyder, A. Z., Baker, J. T., Van Essen, D. C., … Raichle, M. E. (2007). Intrinsic functional architecture in the anaesthetized monkey brain. Nature, 447(7140), 83–86.

Walum, H., & Young, L. J. (2018). The neural mechanisms and circuitry of the pair bond. Nature Reviews Neuroscience, 19(11), 643–654.

Wang, Z. (1995). Species differences in the vasopressin-immunoreactive pathways in the bed nucleus of the stria terminalis and medial amygdaloid nucleus in prairie voles (Microtus ochrogaster) and meadow voles (Microtus pennsylvanicus). Behavioral neuroscience, 109(2), 305–311.

Wang, Z. Hulihan, T. J., & Insel, T. R. (1997). Sexual and social experience is associated with different patterns of behavior and neural activation in male prairie voles. Brain Res, 767(2), 321–332.

Wang, Z. & Aragona, B. J. (2004). Neurochemical regulation of pair bonding in male prairie voles. Physiology and Behavior, 83(2), 319–328.

Williams, J. R., Catania, K. C., & Carter, C. S. (1992). Development of partner

preferences in female prairie voles (Microtus ochrogaster): The role of social and sexual experience. Hormones and Behavior, 26(3), 339–349.

Yamanaka, O., & Takeuchi, R. (2018). UMATracker: an intuitive image-based tracking platform. The Journal of experimental biology, 221(Pt 16), jeb182469.

Yao, S., Bergan, J., Lanjuin, A., & Dulac, C. (2017). Oxytocin signaling in the medial amygdala is required for sex discrimination of social cues. eLife, 6.

Yee, J. R., Kenkel, W. M., Kulkarni, P., Moore, K., Perkeybile, A. M., Toddes, S., … Ferris, C. F. (2016). BOLD fMRI in awake prairie voles: A platform for translational social and affective neuroscience. NeuroImage, 138, 221–232.

Young, K. A., Gobrogge, K. L., Liu, Y., & Wang, Z. (2011). The neurobiology of pair bonding: insights from a socially monogamous rodent. Frontiers in Neuroendocrinology, 32(1), 53–69.

Young, L. J., & Wang, Z. (2004). The neurobiology of pair bonding. Nature Neuroscience, Vol. 7, pp. 1048–1054.

Young, L. J., Young, A. Z. M., & Hammock, E. A. D. (2005). Anatomy and neurochemistry of the pair bond. Journal of Comparative Neurology, 493(1), 51–57.

Zalesky, A., Fornito, A., & Bullmore, E. T. (2010). Network-based statistic: Identifying differences in brain networks. NeuroImage, 53(4), 1197–1207.

Zhong, X., Deng, S., Ma, W., Yang, Y., Lu, D., Cheng, N., … Li, Z. (2017). Anterior cingulate cortex involved in social food-foraging decision-making strategies of rats. Brain and Behavior, 7(10), e00768.