1

**A putative novel role for *Eip74EF* in male reproduction in promoting sperm elongation at the**

**cost of male fecundity**

Sharif Chebbo[1], Sarah Josway[1], John M. Belote[2], and Mollie K. Manier[1*]

[1]*Department of Biological Sciences, The George Washington University, Washington, DC, USA*

[2]*Department of Biology, Syracuse University, Syracuse, NY, USA*

*Correspondence: manier@gwu.edu, +1-202-994-0126*

Running Head: *Eip74EF* promotes sperm length

Abbreviations:

*Eip74EF* - Ecdysone induced protein 74EF

GFP - green fluorescent protein

NCBI - National Center for Biotechnology Information

PAML - Phylogenetic Analysis by Maximum Likelihood

PBS - phosphate buffered saline

REML - restricted maximum likelihood

SR - seminal receptacle

RESEARCH HIGHLIGHTS

*Eip74EF* promotes oocyte maturation in *Drosophila*. We found evidence that it also promotes
sperm elongation in males, but at a cost to male fecundity. Mutant males have shorter sperm
but have higher reproductive success, while females have reduced fecundity.

ABSTRACT

Spermatozoa are the most morphologically variable cell type, yet little is known about genes

controlling natural variation in sperm shape. *Drosophila* fruit flies have the longest sperm

known, which are evolving under postcopulatory sexual selection, driven by sperm competition

and cryptic female choice. Long sperm outcompete short sperm but primarily when females

have a long seminal receptacle (SR), the primary sperm storage organ. Thus, selection on sperm

length is mediated by SR length, and the two traits are coevolving across the *Drosophila* lineage,

driven by a genetic correlation and fitness advantage of long sperm and long SR genotypes in

both males and females. *Ecdysone induced protein 74EF* (*Eip74EF*) is expressed during post-

meiotic stages of spermatogenesis, when spermatid elongation occurs, and we found that it is

rapidly evolving under positive selection in *Drosophila*. Hypomorphic knockout of the *E74A*

isoform leads to shorter sperm but does not affect SR length, suggesting that E74A may be

involved in promoting spermatid elongation but is not a genetic driver of male-female

coevolution. We also found that *E74A* knockout has opposing effects on fecundity in males and

females, with an increase in fecundity for males but a decrease in females, consistent with its

documented role in oocyte maturation. Our results suggest a novel function of *Eip74EF* in

spermatogenesis and demonstrates that this gene influences both male and female reproductive

success. We speculate on possible roles for E74A in spermatogenesis and male reproductive

success.

Keywords: ecdysone signaling, E74A, fitness, sperm length, spermatogenesis, testis

1.  INTRODUCTION

Spermatozoa are arguably the most morphologically variable cell type among metazoans,

exhibiting dramatic diversity in size and shape to encompass all possible permutations of

presence or absence of a head, midpiece, or flagellum (Pitnick et al., 2009). With this level of

morphological variation, it is not surprising that sperm can evolve rapidly on short time scales

(Pitnick et al., 2003; Manier and Palumbi, 2008; Lüpold et al., 2011; Stewart et al., 2016), with

morphological differences emerging even over the course of invasive expansion into new

environments (Kahrl and Cox, 2017). Sperm morphology can influence swimming speed

(Fitzpatrick et al., 2009; Lüpold et al., 2009; Firman and Simmons, 2010; Blengini et al., 2014) and

fertilization success (Bennison et al., 2014), and its rapid evolution has been credited to

postcopulatory sexual selection via sperm competition (Parker, 1970; Snook, 2005; Pitnick et al.,

2009; Godwin et al., 2017) and cryptic female choice (Eberhard, William G., 1986; Firman et al.,

2017).

Within the *Drosophila* lineage, sperm length exhibits among the greatest range of

phenotypic variation within a clade, spanning over two orders of magnitude from 320 μm in *D.*

*persimilis* to 58,290 μm in *D. bifurca* (Pitnick et al., 1995). Moreover, in the obscura group, males

produce heteromorphic sperm, meaning that they come in two (Snook, 1997) or three (Alpern et

al., 2019) different sizes within an ejaculate. Sperm length evolution in *Drosophila* is mediated by

both length of the female's primary sperm storage organ, the seminal receptacle or SR (Miller

and Pitnick, 2002), a genetic correlation with SR length (Lüpold et al., 2016), and moderate

fitness advantage of a long sperm genotype in both males and females (Zajitschek et al., 2019).

As a result of this tight interaction between sperm length and SR length, the two traits are

coevolving both across species (Pitnick et al., 1999) and among populations undergoing

incipient speciation (Pitnick et al., 2003).

Little is known about the genes that control natural variation in sperm length, but we

expect them to be rapidly evolving. In a screen for *Drosophila* genes showing signatures of rapid

adaptive evolution under positive selection, we identified *Ecdysone-induced protein 74EF*

(*Eip74EF*). This gene also has increased expression during post-meiotic stages of

spermatogenesis (Vibranovski et al., 2009), when spermatid elongation occurs, though it has no

known function in spermatogenesis. Neither isoform E74A nor E74B is expressed in the sperm

proteome (Dorus et al., 2006), suggesting that *Eip74EF* is more likely to be involved in

regulation of spermatogenesis rather than encoding a structural component of mature sperm.

*Eip74EF* is an E26 transformation specific (ETS) domain transcription factor, best understood in

its role as an "early response" gene in the ecdysone signaling pathway. Ecdysone regulates

many stages of development, including embryogenesis, larval molting, pupation, and

metamorphosis (Baehrecke, 1996; Yamanaka et al., 2013) and is also important in early ovary

development and virtually all stages of oogenesis (Swevers, 2019). The isoform E74A is

specifically required for  maintenance and proliferation of germline stem cells during oogenesis

(Ables and Drummond-Barbosa, 2010). In spermatogenesis, the ecdysone pathway is critical for

the maintenance and proliferation of germline and cyst stem cells (Li et al., 2014; Qian et al.,

2014), but it has no known role at later stages.

Here, we explored the role of E74A in sperm length. Because sperm length and SR length

are genetically correlated (Lüpold et al., 2016) and coevolving (Pitnick et al., 1999, 2003), we also

measured SR length in E74A hypomorphic mutant females to look for pleiotropic effects on

both sperm length and SR length. We found that E74A mutant males had shorter sperm, so we

also examined mating behavior and sperm competitive success in mutant males and fecundity

in both males and females.

2.   MATERIALS AND METHODS

**2.1 Molecular evolution**

We estimated the rate of positive selection on *Eip74EF* as a measure of adaptive molecular

divergence using DNA sequences from 22 *Drosophila* species downloaded from NCBI. The

codeml program in PAML v4.6 (Yang, 1997, 2007) compares a null model of neutral divergence

(M8a) with a model of adaptive divergence under positive selection (M8), based on the ratio

*dN/dS*, which compares the rate of non-synonymous substitutions (*dN*) with the rate of

synonymous substitutions (*dS*). *dN/dS* = $\omega$, and generally $\omega < 1$ points to purifying selection, $\omega$

= 1 indicates neutrality, and $\omega > 1$ suggests positive selection. However, $\omega$ is rarely greater than

1, since multiple sites across multiple lineages are unlikely to be under positive selection for an

extended period. Thus, $\omega$ estimates are conservative indicators of positive selection on codons.

The critical values for this test are 2.71 at 5% and 5.41 at 1%, with a null distribution of point

mass 0 and $\chi^2_1$ (Self and Liang, 1987).

**2.2 Stocks and crosses**

*Eip74EF* has only one major protein isoform expressed in adults, E74A (Liu et al., 2012). For this

study, we used a hypomorphic mutant of *Eip74EF* obtained from the Bloomington Drosophila

Stock Center (stock #12619) that has partial reduction of expression of E74A while retaining

viability (Liu et al., 2012). The mutant allele (*E74A[BG01805]*) contains a *p(GT1)* transposon insertion

in the sixth intron, within a $w^{1118}$ background. Experimental animals were reared from breeding

vials containing 10 virgin females and 20 virgin males that were transferred to new sugar-yeast-

agar food vials daily for four days, after which adults were discarded. Control animals were

reared from breeding vials containing 10 $E74A^{BG01805}$ females and 20 $w^{1118}$ males. Progeny were

collected within 8-12 hours of eclosion using light $CO_2$ anesthesia into single-sex vials at

densities of 10 females or 20 males per vial. For both experimental and control treatments,

progeny from two replicate sets of breeding vials out of four potential replicates per treatment

were phenotyped for all experiments.

For reproductive success experiments, competitor males were derived from a Canton-S

stock expressing a 3xP3-GFP construct in eye. These Canton-S Protamine-GFP lines were

generated using the piggybac germline transformation system (Handler and Harrell, 1999).  The

GFP-tagged Protamine-B construct was cut out of the *P*-element plasmid construct pW8/ProtB-

GFP1 (Manier et al., 2010) with *Kpn*I and *Bam*HI and subcloned into the pBS/2xAsc vector, a

modified pBlueScript-KS+ plasmid (Stratagene, La Jolla, CA) in which two *Asc*I sites, flanking

the multiple cloning site, had been created by site-directed mutagenesis. The insert was then cut

out with *Asc*I and ligated into the single *Asc*I site of the piggybac vector p3xP3/EGFPaf (Horn

and Wimmer, 2000) to yield p3xP3-EGFP/ProtB-GFP.  This DNA was mixed with the helper

plasmid pBΔSac (Handler and Harrell, 1999) and injected into preblastoderm embyos from a

Canton-S wild-type strain (generously provided by Mariana Wolfner, Cornell University).

Transformants were identified by green fluorescent pseudopupils and ocelli. Two tagged lines

were generated: *Canton-S pBac{3xP3-EGFP, ProtB-EGFP}27* and *Canton-S pBac{3xP3-EGFP,*

*ProtB-EGFP}16B* with the transgenes inserted on chromosome 2 and chromosome 3,

respectively. Although this stock (hereafter "Canton-S") contained a transgene, the fluorescent

eye and sperm markers were not specifically utilized in this series of experiments.

**2.3 Sperm measurements**

To obtain sperm length measurements, seminal vesicles from 5-day-old ether-anesthetized

males were dissected into a large droplet of 1X phosphate-buffered saline (PBS) on a gelatin-

coated glass slide, using fine forceps. The seminal vesicle was ruptured and dragged several

times through the droplet to release and spread motile sperm. The slide was then dried down at

55 °C in a drying oven, fixed in 3:1 methanol:acetic acid, stained with DAPI (Molecular Probes),

and mounted under a coverslip in 80% glycerol in PBS and sealed with nail polish. We then

visualized sperm using darkfield microscopy at 200X and sperm heads using fluorescence

microscopy at 400X on a Nikon Ni-U microscope and captured images with a Zyla Andor

monochrome camera. Images were measured using the segmented line tool in ImageJ software

(https://imagej.nih.gov/ij/). Up to five sperm and sperm heads were measured for each of up

to 8 males across two replicates for both the mutant and control treatments. Sperm

measurements that were abnormally short for a given male were assumed to be due to breakage

and were excluded.

**2.4 SR length**

Females were aged 5 days and then frozen in their food vials until dissection. Females were

then thawed, and thorax length was measured as a proxy for body size using a ruler reticle in a

Nikon SMZ-745 stereoscope at maximum magnification. The lower reproductive tract was

dissected with fine forceps into a droplet of 1X PBS, and the SR was carefully unraveled using a

fine insect pin. The sample was then mounted under a coverslip with clay on the corners to

allow compression of the sample into a single plane without overcompression. SRs were

immediately visualized at 100X or 200X under phase contrast microscopy, and images were

captured using a Canon EOS Rebel T3i camera. For each of two replicates for mutant and

control treatments, 10 females were measured using the segmented line tool in ImageJ.

**2.5 Fitness**

We evaluated the fitness consequences of *Eip74EF* knockdown by quantifying male and female

fecundity and male competitive fertilization success. To measure fecundity, we housed 20 males

or females per replicate individually with two Canton-S females or males, respectively, for

seven days, transferring triads (one focal individual and two mates) into new food vials every

two days, without anesthesia. We counted adult progeny that eclosed from all food vials. This

approach to counting progeny to assess fecundity does not account for effects of *Eip74EF* on

egg-to-adult viability, only on the total number of adult progeny that survived to successfully

eclose.

We assessed paternity success of mutant and control males by mating a $w^{1118}$ female first

with a Canton-S male (day 0) and giving her a four-hour opportunity to remate with a mutant

or control male over the following 4 days (days 1-4). All matings occurred in individual vials

and were observed directly, recording copulation duration and day of remating. We scored

paternity using eye color, since females, mutant males, and control males carried a recessive

white-eye mutation, while Canton-S competitor males had a dominant wild type eye color.

Paternity was expressed as $P_2$, or proportion of offspring sired by the second male (here, mutant

or control males).

**2.6 Statistical analysis**

All statistical analyses were performed in R v.3.5.3 (Team, 2018). Total sperm length and sperm

head length were analyzed using lmer within library lme4, with knockdown treatment as a

fixed effect, male identity nested within replicate as a random effect, and researcher identity as

a random effect, with restricted maximum likelihood (REML) parameter estimation. SR length

was found to be negatively correlated with body size (thorax length) (Pearson's product-

moment correlation, $t_{42}$ = -2.514; $P$ = 0.0159), so thorax length was included as a random effect in

a linear mixed model, with knockdown treatment and replicate as fixed effects, and thorax

length as a random effect. SR lengths were all measured by the same individual. Fecundity data

were analyzed using lmer with knockdown treatment as a fixed effect and replicate as a random

effect, with maximum likelihood parameter estimation, because there was only one random

effect. For all analyses implemented in lmer, P-values were estimated using a Type II Wald chi-

square test with Anova in the car library. Here, we report knockdown treatment effect sizes as

the $t$-value from the mixed model regression, along with the $\chi^2$ and p-value from the Wald's

test.

From the sperm competition experiment, we also obtained remating rate (proportion of

females that remated), copulation duration, and day of remating. Remating rate was analyzed

using a chi-square test in two different ways: with replicates as categories (N = 4) and with

knockdown treatments as categories (N = 2), both yielding the same result. Copulation duration

was analyzed using lmer with knockdown treatment as a fixed effect and replicate as a random

effect, with maximum likelihood parameter estimation. We report results as for other lmer

analyses described above. For paternity success, we used logistic regression with a logit link

function and binomial error distribution (after ensuring no overdispersion in the data),

implemented using glm.

3.   RESULTS

*Eip74EF* is rapidly evolving under positive selection, with an M8a model estimate of -20882.5,

M8 of -20875.6, and a $\chi^2$ of 13.80, well above the 1% critical threshold of 5.41. Knockdown of

*Eip74EF* resulted in shorter sperm ($t = -5.341$, $\chi^2_1 = 28.53$, $P < 0.0001$; Figure 1a) but had no effect

on sperm head length ($t = -1.48$, $\chi^2_1 = 2.191$, $P = 0.139$; Figure 1b) nor female SR length ($t = 1.132$,

$\chi^2_1 = 1.281$, $P = 0.258$; Figure 1c). Disruption of *E74A* led to an increase in male fecundity ($t =$

3.529, $\chi^2_1 = 12.45$, $P = 0.0004$; Figure 2a) but a decrease in female fecundity ($t = -10.17$, $\chi^2_1 =$

103.43, $P < 0.0001$; Figure 2b). In other words, female fecundity was adversely affected while

male fecundity was positively affected. In the sperm competition experiment, female remating

rate (replicates as groups: $\chi^2_3 = 3.143$, $P = 0.370$; treatment as groups: $\chi^2_1 = 0.762$, $P = 0.383$;

Figure 3a), day of remating ($t = 0.336$, $\chi^2_1 = 0.113$, $P = 0.737$; Figure 3b), copulation duration ($t =$

0.109, $\chi^2_1 = 0.012$, $P = 0.913$; Figure 3c), and paternity success ($P_2$; $z = 0.163$, $P = 0.871$; Figure 3d)

were all unaffected by E74A knockout.

4.   DISCUSSION

We found that *Eip74EF* is evolving rapidly under positive selection, a surprising result given

that its roles in molting and metamorphosis, which are both highly conserved processes, would

suggest that it should be under strong purifying selection. However, we found evidence that

E74A plays a novel role during sperm morphogenesis, in the positive regulation of sperm

length, a rapidly evolving trait. The decrease in sperm length in *E74A* mutants is attributable

primarily to change in sperm flagellum length, since head length was unaffected. At the same

time, mutant males had increased fecundity. It should be noted that we have only included one

genetic experiment here, and our sperm length results may be due to other genetic factors

linked to *Eip74EF* rather than this gene itself. More experimentation is needed to further

confirm the effect of E74A on sperm length including, but not limited to, examining sperm

length in a knock-in mutant with expression under control of a heat-shock promoter (Fletcher et

al., 1997).

We know little about E74A's role in testis. Within the ecdysone pathway more generally,

it is one of several genes that are activated when the steroid hormone 20-hydroxyecdysone

(ecdysone) binds to a heteromeric receptor in the nuclear envelope comprised of ecdysone

receptor (EcR) and ultraspiracle (USP) (Yamanaka et al., 2013). EcR is also a transcription factor

that directly initiates expression of a number of early response genes, including *Eip74EF*

(Uyehara and McKay, 2019). In the male germline, EcR is required for growth and proliferation

of both germline stem cells and cyst stem cells, which co-differentiate throughout

spermatogenesis (Li et al., 2014; Qian et al., 2014). Further indirect evidence for E74A function in

spermatogenesis comes from defects found in *belle* (*bel*) mutants. *Bel* is a DEAD box RNA

helicase that regulates translation of *E74A* transcripts in salivary glands during metamorphosis

(Ihry et al., 2012). Although it is unknown if *bel* similarly regulates *E74A* translation in testis,

hypomorphic mutants of *bel* have spermatogenic defects in chromosome segregation and

cytokinesis at the round spermatid stage, approximately when spermatid elongation begins

(Johnstone et al. 2005).

We can only speculate about E74A's possible function during spermatogenesis. We do

know that it is a transcription factor that regulates downstream genes in the ecdysone signaling

pathway. Ecdysone signaling generally tends to promote the maintenance of developmental

stages or to trigger a switch between stages (Gancz and Gilboa, 2013; Li et al., 2014; Yatsenko

and Shcherbata, 2018; Leiblich et al., 2019; Swevers, 2019), with parallel functions in

larval/pupal development and gametogenesis (Ables et al., 2016). Because germline

development is regulated largely by non-autonomous paracrine signaling from supporting

somatic cells (Spradling et al., 2011; Zoller and Schulz, 2012; Li et al., 2014; Qian et al., 2014;

Ables and Drummond-Barbosa, 2017), E74A is likely to be expressed in somatic cyst cells.

Indeed, ecdysone signaling is detectable in nurse cells and follicle cells throughout oogenesis

(Carney and Bender, 2000) and regulates multiple stages from development of the ovarian niche

(Gancz and Gilboa, 2013; Yatsenko and Shcherbata, 2018) to ovulation (Knapp and Sun, 2017;

Swevers, 2019). In the ovary, ecdysone signaling is orchestrated on a fine spatio-temporal scale

throughout oogenesis (Swevers 2019). In spermatogenesis, E74A is also likely to be tightly

regulated at very specific timepoints, especially since expression in the testis overall is very low

(Brown et al., 2014; Li et al., 2014). Less is known about the role of ecdysone signaling in

morphogenesis, though it is required for morphogenesis and function of Malpighian tubules

(Gautam et al., 2015). Perhaps most relevant to flagellar elongation, an early study on *Drosophila*

Kc cells in culture found that application of ecdysone induced elongation and motility in cells

mediated by the formation of filopodia through tubulin assembly (Berger et al., 1980). This

result suggests that ecdysone could also mediate flagellar assembly during spermatogenesis.

On the face of it, it is curious that knockdown of *Eip74EF* has opposing consequences in

males and females, with increased fecundity in males but decreased fecundity in females. The

result in females is more easily explained by the role that *Eip74EF* plays in oogenesis; without it,

germline stem cells arrest development and degenerate. In males, *Eip74EF* seems to promote

longer sperm but at a cost to male fecundity. There are a number of events between mating and

oviposition during which fecundity could ultimately be affected, including sperm transfer,

sperm storage, sperm retention in storage, ovulation, sperm release from storage, fertilization,

oviposition, and offspring viability to adulthood. It is possible that there is a trade-off in

*Eip74EF* mutants between producing longer sperm and producing more sperm, such that because mutants produce shorter sperm, they are now able to produce more sperm (Pitnick and Markow, 1994; Immler et al., 2011). This hypothesis is unlikely to explain the whole story, however, since fecundity in mutant males increased by approximately 50%, while sperm length only decreased by around 5%. Further experiments should examine the rate of spermatogenesis in testes of E74A mutant males.

It is also possible that fewer mutant sperm are able to enter storage. Decreased storage may be a function of sperm storage organs being able to store more shorter sperm. An alternative and non-mutually exclusive hypothesis is that E74A regulates seminal fluid composition and/or transfer. Seminal fluid proteins produced in the accessory glands can have profound effects on many aspects of female reproduction, behavior, and physiology (Avila et al. 2011), including sperm storage after mating (Neubaum & Wolfner 1999; Bloch Qazi & Wolfner 2003; Avila & Wolfner 2009). Accessory gland development and function require ecdysone receptor (EcR) (Herndon et a. 1997; Sharma et al. 2017; Leiblich et al 2019), which directly upregulates E74A in salivary glands and wing disc (Burtis et al., 1990; Thummel et al., 1990; Uyehara and McKay, 2019). It is therefore possible that E74A is also involved in accessory gland function and production of seminal fluid proteins independent of a possible role in morphogenesis during spermatogenesis.

Despite having shorter sperm, *E74A* hypomorphic mutants were not less competitive during sperm competition. Previous work found that the long-sperm advantage only occurs when mating with females with long SRs. We did not measure the SR length of females used in the paternity study, nor did we measure sperm length in the standard male competitors. It is possible that the combinations of male sperm and female SR phenotypes were not conducive to observing a fitness effect of E74A under sperm competition. More controlled experiments in

which knockout males competed against males of known sperm length within females of

known SR length would more clearly demonstrate a role for E74A in sperm competition.

What is driving the rapid evolution of *Eip74EF*? On an evolutionary level, it may reflect

selection on sperm length, mediated by length of the SR (Miller and Pitnick, 2002) and a genetic

correlation between the two traits (Lüpold et al., 2016). However, most reproductive proteins

that are rapidly evolving are ones that interact directly with female-derived molecules, such as

gamete-recognition proteins that mediate fertilization and seminal fluid proteins that modify

and are modified by molecules in the female reproductive tract (Wilburn and Swanson, 2016).

Here, we demonstrate rapid evolution of a protein that has no known direct interactions with

female-derived molecules, nor does it seem to affect length of the SR, which does directly

interact with sperm length. Dorus et al. (2006) found that many sperm proteome proteins are

well-conserved, suggesting that structural proteins that form the building blocks of

spermatozoa are evolving slowly, while the molecular builders that regulate sperm

morphogenesis evolve more rapidly. Thus, the molecular secret to rapid evolution and the vast

morphological diversity of sperm may lie not in its final structure but in its development,

harkening back to the old adage of evo-devo: we cannot understand the evolution of

morphological diversity until we understand the evolution of development.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in Dryad at

https://doi.org/10.5061/dryad.k0p2ngf4k.

ACKNOWLEDGEMENTS

Experimental assistance was provided by Michael DeNieu, Gary Hovsepian, Paul Kwon, Isvita

Marfatia, and Ponmali Photavath. This work was funded by NSF DEB-1257859 to MKM and

JMB, and the George Washington University Harlan Undergraduate Research Fellowship to SC.

The authors declare no conflicts of interest.

REFERENCES

Ables ET, Drummond-Barbosa D. 2010. The Steroid Hormone Ecdysone Functions with
351            Intrinsic Chromatin Remodeling Factors to Control Female Germline Stem Cells in
352            Drosophila. Cell Stem Cell 7:581–592.

353    Ables ET, Drummond-Barbosa D. 2017. Steroid Hormones and the Physiological Regulation of
354            Tissue-Resident Stem Cells: Lessons from the Drosophila Ovary. Curr Stem Cell Rep
355            3:9–18.

356    Ables ET, Hwang GH, Finger DS, Hinnant TD, Drummond-Barbosa D. 2016. A Genetic Mosaic
357            Screen Reveals Ecdysone-Responsive Genes Regulating Drosophila Oogenesis. G3:
358            Genes, Genomes, Genetics 6:2629–2642.

359    Alpern JHM, Asselin MM, Moehring AJ. 2019. Identification of a novel sperm class and its role
360            in fertilization in Drosophila. Journal of Evolutionary Biology 32:259–266.

361    Baehrecke EH. 1996. Ecdysone Signaling Cascade and Regulation of Drosophila
362            Metamorphosis. Archives of Insect Biochemistry and Physiology 33:231–244.

363    Bennison C, Hemmings N, Slate J, Birkhead T. 2014. Long sperm fertilize more eggs in a bird.
364            Proceedings of the Royal Society B: Biological Sciences 282:20141897–20141897.

365    Berger EM, Sloboda RD, Ireland RC. 1980. Tubulin content and synthesis in differentiating
366            drosophila cells in culture. Cell Motility 1:113–129.

367    Blengini CS, Sergio N, Gabriela C, Giojalas LC, Margarita C. 2014. Variability in sperm form and
368            function in the context of sperm competition risk in two Tupinambis lizards. Ecology
369            and Evolution 4:4080–4092.

Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, Booth BW, Wen J, Park S, Suzuki
AM, Wan KH, Yu C, Zhang D, Carlson JW, Cherbas L, Eads BD, Miller D, Mockaitis K,
Roberts J, Davis CA, Frise E, Hammonds AS, Olson S, Shenker S, Sturgill D, Samsonova
AA, Weiszmann R, Robinson G, Hernandez J, Andrews J, Bickel PJ, Carninci P, Cherbas
P, Gingeras TR, Hoskins RA, Kaufman TC, Lai EC, Oliver B, Perrimon N, Graveley BR,
Celniker SE. 2014. Diversity and dynamics of the Drosophila transcriptome. Nature
512:393–399.

Burtis KC, Thummel CS, Jones CW, Karim FD, Hogness DS. 1990. The Drosophila 74EF early
puff contains E74, a complex ecdysone-inducible gene that encodes two ets-related
proteins. Cell 61:85–99.

Carney GE, Bender M. 2000. The Drosophila ecdysone receptor (EcR) Gene Is Required
Maternally for Normal Oogenesis. Genetics 154:1203–1211.

Dorus S, Busby SA, Gerike U, Shabanowitz J, Hunt DF, Karr TL. 2006. Genomic and functional
evolution of the Drosophila melanogaster sperm proteome. Nature Genetics 38:1440–
1445.

Eberhard, William G. 1986. Female Control: Sexual Selection By Cryptic Female Choice.
Princeton: Princeton University Press.

Firman RC, Gasparini C, Manier MK, Pizzari T. 2017. Postmating Female Control: 20 Years of
Cryptic Female Choice. Trends in Ecology & Evolution 32:368–382.

Firman RC, Simmons LW. 2010. Sperm midpiece length predicts sperm swimming velocity in
house mice. Biology Letters 6:513–516.

Fitzpatrick JL, Montgomerie R, Desjardins JK, Stiver KA, Kolm N, Balshine S. 2009. Female
promiscuity promotes the evolution of faster sperm in cichlid fishes. Proceedings of the
National Academy of Sciences 106:1128–1132.

Fletcher JC, D'Avino PP, Thummel CS. 1997. A steroid-triggered switch in E74 transcription
factor isoforms regulates the timing of secondary-response gene expression. PNAS
94:4582–4586.

Gancz D, Gilboa L. 2013. Hormonal Control of Stem Cell Systems. Annual Review of Cell and
Developmental Biology 29:137–162.

Gautam NK, Verma P, Tapadia MG. 2015. Ecdysone regulates morphogenesis and function of
malpighian tubules in Drosophila melanogaster through EcR-B2 isoform.
Developmental Biology 398:163–176.

Godwin JL, Vasudeva R, Michalczyk Ł, Martin OY, Lumley AJ, Chapman T, Gage MJG. 2017.
Experimental evolution reveals that sperm competition intensity selects for longer, more
costly sperm. Evolution Letters 1:102–113.

Handler AM, Harrell RA. 1999. Germline transformation of Drosophila melanogaster with the
piggyBac transposon vector. Insect Molecular Biology 8:449–457.

Horn C, Wimmer EA. 2000. A versatile vector set for animal transgenesis. Dev Gene Evol
210:630–637.

Ihry RJ, Sapiro AL, Bashirullah A. 2012. Translational Control by the DEAD Box RNA Helicase
belle Regulates Ecdysone-Triggered Transcriptional Cascades. PLOS Genetics
8:e1003085.

Immler S, Pitnick S, Parker GA, Durrant KL, Lupold S, Calhim S, Birkhead TR. 2011. Resolving
variation in the reproductive tradeoff between sperm size and number. Proceedings of
the National Academy of Sciences 108:5325–5330.

Kahrl AF, Cox RM. 2017. Consistent Differences in Sperm Morphology and Testis Size between
Native and Introduced Populations of Three *Anolis* Lizard Species. Journal of
Herpetology 51:532–537.

Knapp E, Sun J. 2017. Steroid signaling in mature follicles is important for Drosophila ovulation.
PNAS 114:699–704.

Leiblich A, Hellberg JEEU, Sekar A, Gandy C, Mendes CC, Redhai S, Mason J, Wainwright M,
Marie P, Goberdhan DCI, Hamdy FC, Wilson C. 2019. Mating induces switch from
hormone-dependent to hormone-independent steroid receptor–mediated growth in
Drosophila secondary cells. PLOS Biology 17:e3000145.

Li Y, Ma Q, Cherry CM, Matunis EL. 2014. Steroid signaling promotes stem cell maintenance in
the Drosophila testis. Developmental Biology 394:129–141.

Liu N, Landreh M, Cao K, Abe M, Hendriks G-J, Kennerdell JR, Zhu Y, Wang L-S, Bonini NM.
2012. The microRNA miR-34 modulates ageing and neurodegeneration in Drosophila.
Nature 482:519–523.

Lüpold S, Calhim S, Immler S, Birkhead TR. 2009. Sperm morphology and sperm velocity in
passerine birds. Proceedings of the Royal Society B: Biological Sciences 276:1175–1181.

Lüpold S, Manier MK, Puniamoorthy N, Schoff C, Starmer WT, Luepold SHB, Belote JM,
Pitnick S. 2016. How sexual selection can drive the evolution of costly sperm
ornamentation. Nature 533:535–538.

Lüpold S, Westneat DF, Birkhead TR. 2011. Geographical variation in sperm morphology in the
red-winged blackbird (Agelaius phoeniceus). Evolutionary Ecology 25:373–390.

Manier MK, Belote JM, Berben KS, Novikov D, Stuart WT, Pitnick S. 2010. Resolving
Mechanisms of Competitive Fertilization Success in *Drosophila melanogaster*. Science
328:354–357.

Manier MK, Palumbi SR. 2008. Intraspecific divergence in sperm morphology of the green sea
urchin, Strongylocentrotus droebachiensis: implications for selection in broadcast
spawners. BMC Evolutionary Biology 8:283.

Miller GT, Pitnick S. 2002. Sperm-Female Coevolution in *Drosophila*. Science 298:1230–1233.

Parker GA. 1970. Sperm Competition and Its Evolutionary Consequences in the Insects.
Biological Reviews 45:525–567.

Pitnick S, Hosken DJ, Birkhead TR. 2009. 3 - Sperm morphological diversity. In: Birkhead TR,
Hosken DJ, Pitnick S, editors. Sperm Biology. London: Academic Press. p 69–149.
Available from:
http://www.sciencedirect.com/science/article/pii/B9780123725684000033

Pitnick S, Markow TA. 1994. Male Gametic Strategies: Sperm Size, Testes Size, and the
Allocation of Ejaculate Among Successive Mates by the Sperm-Limited Fly Drosophila
pachea and Its Relatives. The American Naturalist 143:785–819.

Pitnick S, Marrow T, Spicer GS. 1999. Evolution of Multiple Kinds of Female Sperm-Storage
Organs in Drosophila. Evolution 53:1804–1822.

Pitnick S, Miller GT, Schneider K, Markow TA. 2003. Ejaculate-female coevolution in *Drosophila*
mojavensis. Proceedings of the Royal Society of London Series B: Biological Sciences
270:1507–1512.

Pitnick S, Spicer GS, Markow TA. 1995. How long is a giant sperm? Nature 375:109–109.

Qian Y, Dominado N, Zoller R, Ng C, Kudyba K, Siddall NA, Hime GR, Schulz C. 2014.
Ecdysone signaling opposes epidermal growth factor signaling in regulating cyst
differentiation in the male gonad of Drosophila melanogaster. Developmental Biology
394:217–227.

Self SG, Liang K-Y. 1987. Asymptotic Properties of Maximum Likelihood Estimators and
Likelihood Ratio Tests under Nonstandard Conditions. Journal of the American
Statistical Association 82:605–610.

Snook R. 2005. Sperm in competition: not playing by the numbers. Trends in Ecology &
Evolution 20:46–53.

Snook RR. 1997. Is the Production of Multiple Sperm Types Adaptive? Evolution 51:797–808.

Spradling A, Fuller MT, Braun RE, Yoshida S. 2011. Germline Stem Cells. Cold Spring Harb
Perspect Biol 3:a002642.

Stewart KA, Wang R, Montgomerie R. 2016. Extensive variation in sperm morphology in a frog
with no sperm competition. BMC Evolutionary Biology 16:29.

Swevers L. 2019. An update on ecdysone signaling during insect oogenesis. Current Opinion in
Insect Science 31:8–13.

Team RC. 2018. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing.

Thummel CS, Burtis KC, Hogness DS. 1990. Spatial and temporal patterns of E74 transcription
during Drosophila development. Cell 61:101–111.

Uyehara CM, McKay DJ. 2019. Direct and widespread role for the nuclear receptor EcR in
mediating the response to ecdysone in Drosophila. PNAS 116:9893–9902.

Vibranovski MD, Lopes HF, Karr TL, Long M. 2009. Stage-Specific Expression Profiling of
Drosophila Spermatogenesis Suggests that Meiotic Sex Chromosome Inactivation Drives
Genomic Relocation of Testis-Expressed Genes. PLoS Genetics 5:e1000731.

Wilburn DB, Swanson WJ. 2016. From molecules to mating: Rapid evolution and biochemical
studies of reproductive proteins. Journal of Proteomics 135:12–25.

Yamanaka N, Rewitz KF, O'Connor MB. 2013. Ecdysone Control of Developmental Transitions:
Lessons from Drosophila Research. Annu Rev Entomol 58:497–516.

Yang Z. 1997. PAML: a program package for phylogenetic analysis by maximum likelihood.
Bioinformatics 13:555–556.

Yang Z. 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular Biology and
Evolution 24:1586–1591.

Yatsenko AS, Shcherbata HR. 2018. Stereotypical architecture of the stem cell niche is
spatiotemporally established by miR-125-dependent coordination of Notch and steroid
signaling. Development [Internet] 145. Available from:
https://dev.biologists.org/content/145/3/dev159178

Zajitschek S, Zajitschek F, Josway S, Shabeeb RA, Weiner H, Manier MK. 2019. Costs and
benefits of giant sperm and sperm storage organs in Drosophila melanogaster. Journal of
Evolutionary Biology 32:1300–1309.

Zoller R, Schulz C. 2012. The Drosophila cyst stem cell lineage. Spermatogenesis 2:145–157.

FIGURE CAPTIONS

**FIGURE 1** *E74A^BG01805* males had significantly shorter sperm ($P < 0.0001$) (a) with no difference

in sperm head length ($P = 0.139$) (b), and *E74A^BG01805* females had no change in SR length ($P =$

0.258) (c) relative to controls. Two replicates are shown per treatment, with the mutant in dark

gray and control in light gray.

**FIGURE 2** *E74A$^{BG01805}$* males had higher fecundity (P = 0.0004) (a), while *E74A$^{BG01805}$* females had

lower fecundity (P < 0.0001) (b). Two replicates are shown per treatment, with the mutant in

dark gray and control in light gray.

**FIGURE 3** The *E74A$^{BG01805}$* mutation had no effect on a male's ability to influence female

remating rate ($P$ = 0.323) (a), female latency to remate in days ($P$ = 0.737), illustrated as effect

sizes of the statistical model (b), copulation duration in minutes ($P$ = 0.913) (c), and P$_2$ ($P$ =

0.871) (d). Two replicates are shown per treatment, with the mutant in dark gray and control in

light gray.

Fig. 1

Fig. 2

Fig. 3