SPINT2 inhibits proteases involved in activation of both influenza viruses and metapneumoviruses

Marco R. Straus[a*], Jonathan T. Kinder[b*], Michal Segall[a], Rebecca Ellis Dutch[b] and Gary R. Whittaker[a]

[*]M.R.S and J.T.K contributed equally to this work

[a]Department of Microbiology and Immunology, College of Veterinary Medicine, Cornell University,

Ithaca, New York, United States

[b]Department of Molecular and Cellular Biochemistry, University of Kentucky, Lexington, KY, United

States.

Running Head: SPINT2 inhibits influenza virus and metapneumoviruses activation

#Address correspondence to Rebecca E. Dutch, rebecca.dutch@uky.edu or Gary R. Whittaker,

gary.whittaker@cornell.edu

Abstract

Viruses possessing class I fusion proteins require proteolytic activation by host cell proteases to mediate

fusion with the host cell membrane. The mammalian SPINT2 gene encodes a protease inhibitor that

targets trypsin-like serine proteases. Here we show the protease inhibitor, SPINT2, restricts cleavage-

activation efficiently for a range of influenza viruses and for human metapneumovirus (HMPV). SPINT2

treatment resulted in the cleavage and fusion inhibition of full-length influenza A/CA/04/09 (H1N1) HA,

A/Aichi/68 (H3N2) HA, A/Shanghai/2/2013 (H7N9) HA and HMPV F when activated by trypsin,

recombinant matriptase or KLK5. We also demonstrate that SPINT2 was able to reduce viral growth of

influenza A/CA/04/09 H1N1 and A/X31 H3N2 in cell culture by inhibiting matriptase or TMPRSS2.

Moreover, inhibition efficacy did not differ whether SPINT2 was added at the time of infection or 24 hours

post-infection. Our data suggest that the SPINT2 inhibitor has a strong potential to serve as a novel broad-

spectrum antiviral.

Keywords: Influenza, HMPV, Hemagglutinin, HMPV F, proteases, antiviral therapy, SPINT2, HAI-2

Introduction

Influenza-like illnesses (ILIs) represent a significant burden on public health and can be caused by

a range of respiratory viruses in addition to influenza virus itself (1). An ongoing goal of anti-viral drug

discovery is to develop broadly-acting therapeutics that can be used in the absence of definitive diagnosis,

such as in the case of ILIs. For such strategies to succeed, drug targets that are shared across virus families

need to be identified.

One common cause of influenza, which shaped the term ILI, are influenza A viruses, including H1N1 and

H3N2, that cross species barriers from their natural avian hosts and infect humans (2, 3). Novel emerging

viruses such as H7N9 that has caused hundreds of deaths since its appearance in China in 2013 pose an

additional concern (4). The World Health Organization (WHO) estimates that each year about 1 billion

suffer from flu infections, 3 to 5 million people worldwide are hospitalized with severe illness and

approximated 290,000 to 650,000 people die from the disease (5). Mortality rates can dramatically rise

during influenza pandemics as observed with the Spanish flu of 1918, the Asian pandemic of 1957 and the

Hong Kong pandemic of 1968 (3, 6). Vaccinations can provide an effective protection against seasonal and

pandemic outbreaks but provide limited or no protection when viruses evolve and/or acquire mutations

resulting in antigenically distinct viruses. This antigenic drift or shift requires that new vaccines be

produced quickly and in vast amounts which can be problematic especially during pandemics. In addition

to vaccines, several antiviral therapies have been applied to treat influenza A infections such as

adamantanes acting as M2 ion channel blockers (amantadine, rimantadine) and neuraminidase inhibitors

which block the cleavage of sialic acid in newly formed virions (oseltamivir, zanamivir) (7). However,

several influenza subtypes including the most common H1N1 and H3N2 have emerged globally that are

resistant against adamantanes and similar observations were made with respect to oseltamivir (7). More

recently, baloxavir marboxil (BXM) which is an inhibitor of the influenza polymerase acidic subunit (PA)

was approved as an antiviral therapy (8). Although it is fully effective against currently circulating influenza

A and B viruses clinical trial studies have already shown that treatment with BXM selects for influenza

virus with specific amino acid substitutions in PA resulting in reduced susceptibility to the drug (9, 10).

Another common cause of ILI are pneumo- and paramyxoviruses, including human metapneumovirus

(HMPV), respiratory syncytial virus, and parainfluenza viruses. Clinical presentation of these viruses

resembles many of the symptoms of influenza, where they cause significant morbidity and mortality as

well as a large economic burden (11, 12). HMPV is ubiquitous, with nearly everyone infected by the age

of 5 and reinfection is common throughout life, impacting children, the elderly and immunocompromised

individuals (13–15). While HMPV is a common cause of ILI, there are currently no approved vaccines or

antiviral therapeutics. Further research is needed to establish targets for intervention, and factors

required for infection need to be examined in more detail.

Certain influenza viruses and HMPV appear to share common activating proteases. The influenza fusion

protein hemagglutinin (HA) is synthesized as a precursor that needs to be cleaved by host cell proteases

to exert its fusogenic activity (16, 17). Cleavage separates the precursor HA_0 into HA_1 and HA_2, which

remain associated via disulfide bonds and leads to exposure of the fusion peptide at the N-terminus of

HA_2 (16, 17). Low pathogenicity avian influenza (LPAI) usually possess a monobasic cleavage site which

consists of 1 – 2 non-consecutive basic amino acids and which is generally cleaved by trypsin-like serine

proteases such trypsin, as well as members of the type II transmembrane serine protease (TTSP) family

including TMPRSS2, TMPRSS4, HAT (TMPRSS11D) and matriptase (16–18). In addition, some other

proteases such as KLK5 and KLK 12 have been implicated in influenza pathogenicity (19). In humans these

proteases are localized in the respiratory tract and therefore influenza infections are usually confined to

this tissue. In contrast, highly pathogenic avian influenza (HPAI) viruses are defined by a polybasic cleavage

site which consists of 6 – 7 basic residues allowing them to be activated by members of the proprotein

convertase (PC) family such as furin and PC6 (20). These proteases are not confined to a specific tissue

and dramatically increase the risk of a systemic infection.

Similar to influenza HA, HMPV requires the fusion protein (F) to be proteolytically processed at a single

basic residue to generate the active, metastable form. Without this cleavage process, the F protein is

unable to mediate viral entry into the target cell. However, to date, only trypsin and TMPRSS2 have been

shown to effectively cleave HMPV F and other proteases have yet to be identified (21, 22).

In addition, other respiratory viruses have been reported to utilize similar proteases for activation,

including SARS-CoV and MERS-CoV, demonstrating that targeting these proteases would inhibit multiple

respiratory pathogens (23). The fact that proteolytic activation is such a crucial step for several respiratory

viruses that predominately require a specific class of proteases makes these proteases a viable target for

the development of novel antiviral therapies (24). Earlier studies described the administration of the

serine protease inhibitor aprotinin to inhibit influenza replication and demonstrated that aprotinin

successfully inhibited IAV activation and replication (25). However, when targeting host specific factors

there are potential off target effects, and therefore the potential side effects of targeting host proteases

requires further investigation. Hamilton et al. reported that the hepatocyte growth activator inhibitor 2

(HAI-2) effectively inhibited trypsin-mediated cleavage of H1N1 and H3N2 *in vitro* and *in vivo* (26). HAI-2

is encoded by the SPINT2 gene and hereafter we will also refer to the protein as SPINT2. SPINT2 is 225

KDa plasma membrane-localized serine protease inhibitor found in epithelial cells of various tissues including the respiratory tract and all major organs (27). In most tissues, SPINT2 co-localizes with matriptase suggesting a regulatory role of SPINT2 on matriptase-mediated cleavage events. However, the finding that SPINT2 is also expressed in brain and lymph node cells indicates that it might regulate other proteases than matriptase (27). Recent reports associated the physiological role of SPINT2 with the inhibition of human serine-type proteases such as matriptase, plasmin, kallikreins (KLK) and coagulation factor XIa (28–31). SPINT2 possesses one transmembrane domain and two kunitz-type inhibitor domains that are exposed to the extracellular space and which are believed to facilitate a potent inhibition of target proteases. Wu et al. recently described that the kunitz-type domain 1 of SPINT2 is responsible for matriptase inhibition (28). A major function of SPINT2 is its role as a tumor suppressor because down-regulation diminishes the prospect of survival of several cancers such as hepatocellular carcinoma, gastric cancer, prostate cancer or melanoma (32–35). However, SPINT2 was also associated with placenta development and epithelial homeostasis (36, 37).

A previous study from our lab described the effective inhibition of trypsin by SPINT2 resulting in dramatically reduced cleavage of influenza A HA using a model protease and subsequently reduced viral growth in cell culture and mouse studies (26). Here, we report that purified SPINT2 protein inhibits several host proteases found in the human respiratory tract, such as matriptase and TMPRSS2, that are relevant for the activation of influenza viruses currently circulating and causing significant disease outbreaks. To demonstrate broad applicability, we also tested the potential of SPINT2 to inhibit the activation of the fusion protein (F) from human metapneumovirus (HMPV), a member of the pneumovirus family. We confirm the original findings that HMPV F is proteolytically processed by trypsin and TMPRSS2. In addition, we found that HAT, KLK5 and matriptase were able to cleave F, but KLK12 could not. Our results show that SPINT2 can inhibit the activation of proteases that are responsible for the activation of influenza H1N1, H3N2 and H7N9 HA as well as HMPV F. In a cell culture model, we demonstrate that viral loads are

significantly reduced in the presence of SPINT2 when infections were conducted with A/CA/04/09 and

A/X31. Moreover, the application of SPINT2 24 hours post infection inhibited the activation of influenza

A viruses with the same efficacy as when SPINT2 was added to cell culture medium at the time of infection.

Thus, SPINT2 exhibits the potential to serve as a novel and efficient antiviral therapeutic to relieve patients

from influenza A, human metapneumovirus, SARS-CoV and potentially other respiratory viruses that

require these host factors for entry.

Results

SPINT2 inhibits recombinant human respiratory tract proteases that cleave HMPV F and HA cleavage

site peptide mimics

Using a fluorogenic peptide cleavage assay that utilizes fluorogenic peptides mimicking the HA

cleavage site we previously tested the ability of SPINT2 to inhibit proteases shown to cleave HAs from

seasonal and pandemic influenza A strains that infected humans (38, 39). We found that certain HA

subtypes such as H1, H2 and H3 are cleaved by a wide variety of human respiratory proteases while others

such H5, H7 and H9 displayed more variability in cleavage by proteases and seemed less well adapted to

proteases present in the human respiratory tract (39). Here, we extended our previous study and tested

a peptide mimicking the cleavage site of the pneumovirus fusion protein of HMPV F using a variety of

proteases known for their ability to cleave the peptide mimic (Table 1) (22, 40). When we tested the

cleavage of a peptide mimicking the HMPV F cleavage site using trypsin, matriptase, KLK5, KLK12, HAT or

plasmin we found that all proteases except KLK12 were able to proteolytically cleave the peptide (Table

1). However, the Vmax values for matriptase (9.24 RFU/min), KLK5 (5.8 RFU/min) and HAT (2.99 RFU/min)

were very low compared to trypsin (135.2 RFU/min) suggesting that the three proteases have a low

affinity interaction and processivity for HMPV F.

Next, trypsin, matriptase and KLK5 were selected for the SPINT2 inhibition assays as described below.

For the SPINT2 inhibition assays, trypsin (which typically resides in the intestinal tract and expresses a very

broad activity towards different HA subtypes and HMPV F) served as a control (41). In addition, furin was

used as a negative control that is not inhibited by SPINT2. As none of the peptides used in combination

with the aforementioned proteases has a furin cleavage site we tested furin-mediated cleavage on a

peptide with a H5N1 HPAI cleavage motif in the presence of 500 nM SPINT2 (Supplementary Figure 1).

We continued by measuring the Vmax values for each protease/peptide combination in the presence of

different SPINT2 concentrations and plotted the obtained Vmax values against the SPINT2 concentrations

on a logarithmic scale (Supplementary Figure 2). Using Prism7 software, we then determined the IC_{50} that

reflects at which concentration the V_{max} of the respective reaction is inhibited by half. SPINT2 cleavage

inhibition of a representative H1N1 cleavage site by trypsin results in an IC50 value of 70.6 nM (Table 2A)

while the inhibition efficacy of SPINT2 towards matriptase, HAT, KLK5 and KLK12 ranged from 11 nM to

25 nM (Table 2A). However, inhibition was much less efficient for plasmin compared with trypsin (122

nM). We observed a similar trend when testing peptides mimicking the H3N2 and H7N9 HA cleavage sites

using trypsin, HAT, KLK5, plasmin and trypsin, matriptase, plasmin, respectively (Table 2A). With the

exception of plasmin, we found that human respiratory tract proteases are inhibited with a higher efficacy

compared to trypsin. We expanded our analysis to peptides mimicking HA cleavage sites of H2N2, H5N1

(LPAI and HPAI), H6N1 and H9N2 that all reflected the results described above (Table 2A). Only cleavage

inhibition of H6N1 HA by KLK5 did not significantly differ from the observation made with trypsin (Table

2A).

When we tested the inhibition of HMPV cleavage by trypsin, matriptase and KLK5, SPINT2 demonstrated

high inhibition efficacy for all three tested proteases with measured IC50s for trypsin, matriptase and KLK5

of 0.04 nM, 0.0003 nM and 0.95 nM, respectively (Table 2B).Compared to the IC50 values observed with

the peptides mimicking influenza HA cleavage site motifs the IC50 values for the HMPV F peptide were

very low.

Cleavage of distinct full-length HA subtypes and HMPV F is efficiently inhibited by SPINT

Cleavage of peptides mimicking cleavage sites of viral fusion proteins do not always reflect the *in vivo* situation and requires validation by expressing the full-length fusion proteins in a cell culture model to test cleavage and cleavage inhibition of the respective protease (39). However, before conducting these experiments we wanted to ensure that SPINT2 does not have a cytotoxic effect on cells. Therefore, 293T cells were incubated with various concentrations of SPINT2 over a time period of 24 hours. PBS and 500µM H_2O_2 served as cytotoxic negative and positive controls respectively. We observed a slight reduction of about 10 -15 % in cell viability when SPINT2 was added to the cells (Figure 1).

To test SPINT2-mediated cleavage inhibition of full-length HA we expressed the HAs of A/CA/04/09 (H1N1), A/x31 (H3N2) and A/Shanghai/2/2013 (H7N9) in 293T cells and added recombinant matriptase or KLK5 protease that were pre-incubated with 10nM or 500nM SPINT2. Trypsin and the respective protease without SPINT2 incubation were used as controls. Cleavage of HA_0 was analyzed via Western Blot and the signal intensities of the HA_1 bands were quantified using the control sample without SPINT2 incubation as a reference point to illustrate the relative cleavage of HA with and without inhibitor (Figure 2A and B-D). Trypsin cleaved all tested HA proteins with very high efficiency that was not observed with matriptase or KLK5 (Figure 2B-D). However, H1N1 HA was cleaved by matriptase and KLK5 to a similar extent without and with 10nM SPINT2. 500nM SPINT2 led to a relative cleavage reduction of about 70% and 50% for matriptase and KLK5, respectively (Figure 2A and B). KLK5-mediated cleavage of H3N2 HA was reduced by about 10% when KLK5 was pre-incubated with 10nM SPINT2 and by about 60% when 500nM SPINT2 was used (Figure 2A and C). When we tested the cleavage inhibition of matriptase with H7N9 HA as a substrate we found that 10nM and 500nM SPINT reduced the cleavage to 40% and 10% cleavage, respectively, compared to the control. (Figure 2A and D). In contrast, 10nM SPINT2 had no effect on KLK5-mediated cleavage of H7N9 HA while 500nM reduced relative cleavage by approximately 70% (Figure 2A and D).

In order to determine whether SPINT2 also prevented cleavage of HMPV F we first examined which

proteases, in addition to trypsin and TMPRSS2, were able to cleave HMPV F. First, we co-transfected the

full length TMPRSS2, HAT and matriptase with HMPV F in VERO cells. The F protein was then radioactively

labeled with ^{35}S methionine and cleavage was examined by quantifying the F_0 full length protein and the

F_1 cleavage product. We found that TMPRSS2 and HAT were able to efficiently cleave HMPV F while

matriptase decreased the expression of F, though it is not clear if this was due to general degradation of

protein or lower initial expression. However, matriptase demonstrated potential low-level cleavage when

co-transfected (Figure 3A and B). We then examined cleavage by the exogenous proteases KLK5, KLK12

and matriptase. Compared with the trypsin control, KLK5 and matriptase were able to cleave HMPV F,

while KLK12 was not (Figure 3C and D). In agreement with the peptide assay, cleavage of HMPV F by KLK5

and matriptase was less efficient than for trypsin and both peptide, and full-length protein assays

demonstrate that KLK12 does not cleave HMPV F. This also serves as confirmation that matriptase likely

cleaves HMPV F, but co-expression with matriptase may alter protein synthesis, stability or turnover if co-

expressed during synthesis and transport to the cell surface. Next, we tested SPINT2 inhibition of

exogenous proteases trypsin, KLK5 and matriptase. We pre-incubated SPINT2 with each protease, added

it to VERO cells expressing HMPV F and analyzed cleavage product formation. SPINT2 pre- incubation

minimally affected cleavage at a concentration of 10nM but addition of 500nM SPINT2 resulted in

inhibition of trypsin, KLK5 and matriptase-mediated cleavage of HMPV, similar to our findings for HA

(Figure 3E and F).

SPINT2 inhibits HA mediated cell-cell fusion

The above presented biochemical experiments demonstrate that SPINT2 is able to efficiently

inhibit proteolytic cleavage of HMPV F and several influenza A HA subtypes by a variety of proteases. For

a functional analysis to determine whether SPINT2 inhibition prevents cells to cell fusion and viral growth,

we examined influenza A infection in cell culture. While HMPV is an important human pathogen, Influenza

grows significantly better in cell culture compared with HMPV. First, we tested whether cleavage

inhibition by SPINT2 resulted in the inhibition of cell-cell fusion. As described above, matriptase and KLK5

were pre-incubated with 10nM and 500nM SPINT2 and subsequently added to VERO cells expressing

A/CA/04/09 (H1N1) HA or A/Shanghai/2/2013 (H7N9) HA. Cells were then briefly exposed to a low pH

buffer to induce fusion and subsequently analyzed using an immune fluorescence assay. When matriptase

and KLK5 were tested with 10nM SPINT2 and incubated with VERO cells expressing H1N1 HA, we still

observed syncytia formation (Figure 4A). However, 500mM SPINT2 resulted in the abrogation of syncytia

formation triggered by cleavage of the respective HA by matriptase and KLK5. We made the same

observation when we tested KLK5 and H7N9 HA (Figure 4B). Matriptase-mediated H7N9 HA syncytia

formation was inhibited by the addition of 10nM SPINT2 (Figure 4B). To ensure that cell-cell fusion

inhibition is a result of HA cleavage inhibition through SPINT2 but not a side effect of SPINT2 treatment

per se we expressed A/Vietnam/1204/2004 (H5N1) HA in VERO cells and treated them with the inhibitor.

H5N1 HA possesses a HPAI cleavage site and is cleaved intracellularly by furin during its maturation

process. SPINT2 does not inhibit furin and is not able to cross cell membranes. Thus, SPINT2 can not

interfere with the proteolytic processing of H5N1 HA and therefore this control allows to examine whether

SPINT2 interferes with cell-cell fusion. Figure 4C shows that H5N1 HA forms large syncytia in the absence

of SPINT2 as well as in the presence of 500 nM SPINT2. Hence, we conclude that SPINT2 does not have a

direct inhibitory effect on cell-cell fusion.

SPINT2 reduces viral growth in cell culture

To understand whether SPINT2 was able to inhibit or reduce the growth of virus in a cell culture

model over the course of 48 hours we transfected cells with human TMPRSS2 and human matriptase, two

major proteases that have been shown to be responsible for the activation of distinct influenza A subtype

viruses. TMPRSS2 is essential for H1N1 virus propagation in mice and plays a major role in the activation

of H7N9 and H9N2 viruses (42–44). Matriptase cleaves H1N1 HA in a sub-type specific manner, is involved

in the *in vivo* cleavage of H9N2 HA and our results described above suggest a role for matriptase in the

activation of H7N9 (19, 44). At 18 hours post transfection we infected MDCK cells with A/CA/04/09 (H1N1)

at a MOI of 0.1 and subsequently added SPINT2 protein at different concentrations. Non-transfected cells

served as a control and exogenous trypsin was added to facilitate viral propagation. The supernatants

were harvested 48 hours post infection and viral titers were subsequently analyzed using an immuno-

plaque assay. SPINT2 initially mitigated trypsin-mediated growth of H1N1 at a concentration of 50nM and

the extent of inhibition slightly increased with higher concentrations (Figure 5A, Table 3A). The highest

tested SPINT2 concentration of 500nM reduced viral growth by about 1 log (Figure 5A, Table 3A). We

observed a similar pattern with cells transfected with human matriptase (Figure 5B, Table 3A). Growth

inhibition started at a SPINT2 concentration of 50nM and with the application of 500nM growth was

reduced by approximately 1.5 logs (Figure 5B, Table 3A). When we infected cells expressing TMPRSS2 with

H1N1 and added SPINT2, viral growth was significantly reduced at a concentration of 150nM. Addition of

500nM SPINT2 led to a reduction of viral growth of about 1.5 logs (Figure 5C, Table 3A). We also tested

whether SPINT2 could reduce the growth of a H3N2 virus because it is major circulating seasonal influenza

subtype. However, TMPRSS2 and matriptase do not seem to activate H3N2 viruses (19, 45). Hence, trypsin

and SPINT2 were added to the growth medium of cells infected with A/X31 H3N2. Compared to control

cells without added inhibitor SPINT2 significantly inhibited trypsin-mediated H3N2 growth at a

concentration of 50nM (Figure 5D, Table 3A). At the highest SPINT2 concentration of 500nM viral growth

was reduced by about 1 log (Figure 5D).

We also examined the effect of SPINT2 inhibition of HMPV spread over time. VERO cells were

infected with rgHMPV at MOI 1 and subsequently treated with 500nM of SPINT2 and 0.3μg/mL of TPCK-

trypsin. Every 24 hours, cells were scraped and the amount of virus present was titered up to 96hpi with

SPINT2 and trypsin replenished daily. We find that un-treated cells are infected and demonstrate

significant spread through 96hpi. Conversely, cells infected and treated with SPINT2 had no detectible

virus up to 48hpi and very minimal virus detected at 72 and 96hpi, demonstrating that SPINT2 significantly

inhibition HMPV viral replication and spread (Figure 6).

Antiviral therapies are often applied when patients already show signs of disease. Therefore, we

tested if SPINT2 was able to reduce viral growth when added to cells 24 hours after the initial infection.

Cells were infected with 0.1 MOI of A/CA/04/09 (H1N1) and trypsin was added to promote viral growth.

At the time of infection, we also added 500nM SPINT2 to one sample. A second sample received 500nM

SPINT2 24 hours post infection. Growth supernatants were harvested 24 hours later, and viral growth was

analyzed. We found that viral growth was significantly reduced by regardless whether SPINT2 was added

at the time of infection or 24 hours later (Figure 7, Table 3B).

Discussion

Influenza A virus has caused four pandemics since the early 20[th] century and infects millions of

people each year as seasonal 'flu, resulting in up to 690,000 deaths annually (5). Vaccination efforts have

proven to be challenging due to the antigenic drift of the virus and emerging resistance phenotypes (46).

Moreover, the efficacy of vaccines seems to be significantly reduced in certain high-risk groups (47).

Prevalent antiviral therapies to treat influenza A virus-infected patients such as adamantanes and

neuraminidase inhibitors target viral proteins but there is increasing number of reports about circulating

influenza A subtypes that are resistant to these treatments (7). HMPV causes infections in the upper and

lower respiratory tract expressing very similar symptoms as influenza infections and resulting in significant

morbidity and mortality (11, 12). The most susceptible groups are young children, older adults and people

that are immunocompromised (13–15). Currently, there are not treatment against HMPV infections

available. In this study we focused on a novel approach that uses antiviral therapies targeting host factors

rather than viral proteins offering a more broad and potentially more effective therapeutic approach (24).

We demonstrate that SPINT2, a potent inhibitor of serine-type proteases, is able to significantly inhibit

cleavage of HMPV F and HA, impair HA-triggered fusion of cells and hence, reduce the growth of various

influenza A strains in cell culture.

We assessed cleavage of the HMPV fusion protein in vitro using a peptide cleavage assay modified from

previously work on other viral fusion proteins (38, 48). The HMPV F peptide was cleaved by trypsin,

plasmin, matriptase and KLK5 but was unable to be cleaved by KLK12. To confirm these findings in a

system in which the entire HMPV F protein was subject to cleavage, we co-expressed the fusion protein

with TMPRSS2, HAT and matriptase proteases, or treated F with exogenous proteases KLK5, KLK12 and

matriptase. These findings are the first to identify proteases besides trypsin and TMPRSS2 that are able

to cleave HMPV F. In addition, HMPV appears to utilize many of the serine proteases that influenza uses

for HA processing and therefore, offers strong potential for an antiviral target.

SPINT2 demonstrates greater advantage over other inhibitors of host proteases such as e.g. aprotinin that

was shown to be an effective antiviral but also seemed to be specific only for a subset of proteases (25).

It can be argued that a more specific protease inhibitor which inhibits only one or very few proteases

might be more advantageous because it may result in less side effects. With respect to influenza A

infections, TMPRSS2 could represent such a specific target as it was shown to a major activating proteases

for H1N1 and H7N9 in mice and human airway cells (42, 44, 49–51). However, there is no evidence that

that application of a broad-spectrum protease inhibitor results in more severe side effects than a specific

one as side effects may not be a consequence of the protease inhibition but the compound itself may act

against different targets in the body. The reports demonstrating that TMPRSS2 is crucial for H1N1 and

H7N9 virus propagation in mice and cell culture suggest that it also plays a major role in the human

respiratory tract. So far, however, it is unclear whether the obtained results translate to humans and other

studies have shown that for example human matriptase is able to process H1N1 and H7N9 (19, 52).

Our peptide assay suggests that SPINT2 has a wide variety of host protease specificity. With the exception

of plasmin, all the tested proteases in combination with peptides mimicking the cleavage site of different

HA subtypes expressed IC_{50} values in the nanomolar range. Interestingly, the IC50 values obtained for

cleavage inhibition of HMPV F were substantially lower, in the picomolar range. This suggests that the

HMPV cleavage may be more selectively inhibited by SPINT2. However, the western blot data showed

that addition of the lowest concentration (10 nM) of SPINT2 did not result in cleavage inhibition of HMPV

F by the tested proteases. Differences in sensitivity of SPINT2 between influenza HA and HMPV require

further investigation.

SPINT2 poses several potential advantages over other inhibitors that target host proteases. Cell culture

studies showed that, for example, matriptase-mediated H7N9 HA cleavage was efficiently inhibited at a

concentration of 10nM SPINT2. In contrast, the substrate range for aprotinin, a serine protease inhibitor

shown to reduce influenza A infections by targeting host proteases, seemed to be more limited (25). Other

synthetic and peptide-like molecules designed to inhibit very specific serine proteases such as TMPRSS2,

TMPRSS4 and TMPRSS11D (HAT) were only tested with those proteases and their potential to inhibit other

proteases relevant for influenza A activation remains unclear (53–55). Currently, the most promising

antiviral protein inhibitor is camostat which is already approved in Japan for the treatment of chronic

pancreatitis (56). Recently, it was demonstrated that camostat inhibited influenza replication in cell

culture and prevented the viral spread and pathogenesis of SARS-CoV in mice by inhibition of serine

proteases (55, 57). However, camostat was applied prior to the virus infection and it was administered

into the mice via oral gavage (57). A previous study showed that SPINT2 significantly attenuated influenza

A infections in mice using a concentration that was 40x lower than the described camostat concentration

and intranasal administration was sufficient (26). Our current study suggests that SPINT2 is able to

significantly inhibit viral spread during an ongoing infection and does not need to be applied prior or at

the start of an infection. The mouse study also showed that SPINT2 can be applied directly to the

respiratory tract while camostat that is currently distributed as a pill and therefore less organ specific. In

addition, camostat is synthetic whereas SPINT2 is a naturally occurring molecule which may attenuate

potential adverse effects due to non-native compounds activating the immune system. Future research

will be conducted to test if SPINT2 can be applied more efficiently via an inhaler and to explore potential

side effects in mice studies. However, when we tested the potential of SPINT2 to inhibit viral replication

in a cell culture model we were only able to achieve growth reductions of approximately 1 -1.5 logs after

48 hours with a concentration of 500nM SPINT2. One potential explanation is that 500nM SPINT2 was

unable to saturate the proteases present in the individual experiments and was not sufficient to prevent

viral growth. In addition, the continuous overexpression of matriptase and TMPRSS2 may have produced

an artificially high quantity of protein that exceeded the inhibitory capacities of SPINT2. This problem

could be solved either by using higher concentration of SPINT2 or by optimizing its inhibitory properties.

However, the data also demonstrates that SPINT2 has the ability to inhibit proteases that expressed on

the cell surface and that inhibition is not limited to proteases that were added exogenously and pre-

incubated with the inhibitor (Figure 5). SPINT2 did not express any cytotoxic effects up to a concentration

of 10mM, significantly above the therapeutic dosage required for inhibition. In comparison with other

studies, the SPINT2 concentration we used here were in the nanomolar range while other published

inhibitors require micromolar concentrations (53–55). However, we believe that future research will allow

to fully utilize the potential of SPINT2 as a broad-spectrum antiviral therapy. Wu et al., recently described

that the Kunitz domain I of SPINT2 is responsible for the inhibition of matriptase (28). In future studies we

will explore whether the inhibitory capabilities of SPINT2 can be condensed into small peptides that may

improve its efficacy. Its ability to inhibit a broad range of serine protease that are involved in the activation

of influenza A suggest that a SPINT2 based antiviral therapy could be efficient against other pathogens

too. TMPRSS2, for example, not only plays a major role in the pathogenesis of H1N1 but is also required

for the activation of SARS-CoV and MERS-CoV and HMPV (58, 59). Currently, treatment options for these

viruses are very limited and therefore SPINT2 could become a viable option if its potential as an antiviral

therapeutic can be fully exploited.

However, while SPINT2 has a therapeutic potential to treat ILIs caused by viruses that require activation

by trypsin-like serine proteases it may have its limitation to provide a treatment option for infections

caused by influenza HPAI viruses, such as H5N1 (60). These viruses are believed to be activated by furin

and pro-protein convertases that belong to the class of subtilisin-like proteases (61). Preliminary data

from our lab demonstrated that SPINT2 did not inhibit furin-mediated cleavage of HPAI cleavage site

peptide mimics as well as peptides carrying described furin cleavage sites (data not shown). In addition,

furin acts intracellularly and we have no evidence that SPINT2 is able to penetrate the cell membrane and

thus inhibiting proteases located in intracellular compartments. Therefore, it seems unlikely that SPINT2

is able to inhibit furin in cell culture-based studies or *in vivo* experiments.

In conclusion we believe SPINT2 has potential to be developed into a novel antiviral therapy. In contrast

to most similar drugs that are synthetic, SPINT2 is an endogenously expressed protein product that

confers resistance to a variety of pathogenic viruses and which can potentially be delivered directly into

the respiratory tract as an aerosol. Most importantly, SPINT2 demonstrated the ability to significantly

attenuate an ongoing viral infection in cell culture and further research will be conducted to explore the

time period during which SPINT2 demonstrates the highest efficacy.

Materials and Methods

Cells, plasmids, viruses, and proteins

293T, VERO and MDCK cells (American Type Culture Collection) used for influenza experiments

were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with 25 mM HEPES

(Cellgro) and 10% fetal bovine serum (VWR). VERO cells used for HMPV experiments were maintained in

DMEM (HyClone) supplemented with 10% FBS (Sigma). The plasmid encoding A/CA/04/09 (H1N1) HA was

generated as described (19). The plasmid encoding for HMPV F S175H434 was generated as described(62).

The plasmids encoding for A/Shanghai/2/2013 (H7N9) HA, human TMPRSS2 and human matriptase were

purchased from Sino Biological Inc. The plasmid encoding for A/Aichi/2/68 (H3N2) HA was generously

donated by David Steinhauer. A/CA/04/09 and A/X31 viruses were propagated in eggs. All recombinant

proteases were purchased as described (39).

Expression and purification of SPINT2

SPINT2 was expressed and purified as described with minor modification (26). In brief, *E. coli* RIL

(DE3) arctic express cells (Agilent) were transformed with SPINT2-pSUMO. Cells were then grown in 0.5 L

Luria Broth containing 50 µg/ml kanamycin at 37°C. At OD 0.5-0.6 cells were chilled on ice and protein

expression was induced with 0.2 mM IPTG. Cells were then grown over night at 16°C. Cells were harvested

and protein was purified as previously described (26). SPINT2 protein was eluted by a 1-hour incubation

with ULP1-6xHis. Glycerol was added to the eluted SPINT2 protein to a final concentration of 20% and

protein aliquots were stored at -80°C. Protein concentration was determined by analyzing different

dilutions of SPINT2 on an SDS-PAGE gel along with 5 defined concentrations of BSA between 100 ng and

1 µg. The gel was then stained with Coomassie, scanned with ChemiDoc Imaging system (Bio-Rad) and

bands were quantified using Image Lab software (Bio-Rad). Concentrations of the SPINT2 dilutions were

determined based on the BSA concentrations and the final SPINT2 concentration was calculated based on

the average of the SPINT2 different dilutions.

Peptide Assays

Peptide assays were carried out as described (39). The sequence of the HMPV F peptide mimicking

the HMPV F cleavage site used in this assay is ENPRQSRFVL including the same N- and C-terminal

modifications as described for the HA peptides (39). The V_{max} was calculated by graphing each replicate in

Microsoft Excel and determining the slope of the reaction for every concentration (0 nM, 1 nM, 5 nM, 10

nM, 25 nM, 50 nM, 75 nM, 150 nM, 300 nM and 500 nM). The V_{max} values were then plotted in the

GraphPad Prism 7 software against the log10 of the SPINT2 concentration to produce a negative sigmoidal

graph from which the IC_{50}, or the concentration of SPINT2 at which the V_{max} is inhibited by half, could be

extrapolated for each peptide protease mixture. Since the x-axis was the log10 of the SPINT2

concentration, the inverse log was then taken for each number to calculate the IC_{50} in nM.

Cytotoxicity assay

The cytotoxicity assay was performed with a cell counting kit-8 (Dojindo Molecular Technologies)

according to the manufacturer's instructions. In brief, approx. 2×10^3 293T cells were seeded per well of

a 96-well plate and grown over night. SPINT2 was added at the indicated concentrations. DMEM and 500

µM H_2O_2 were used as a control. 24 hours later 10 µl CKK-8 solution were added to each well and

incubated for 1 hour. Absorbance at 450 nm was measured using a SPARK microplate reader (Tecan). Per

sample and treatment three technical replicates were used and the average was counted as one biological

replicate. Experiment was conducted three times.

Metabolic protein labeling and immunoprecipitation of HMPV fusion protein

VERO cells were transfected with 2ug of pDNA using Lipofectamine and plus reagent (Invitrogen)

in opti-mem (Gibco) according to the manufacturers protocol. The following day, cells were washed with

PBS and starved in cysteine/methionine deficient media for 45 min and radiolabeled with 50uC/mL S35-

cysteine/methionine for 4 hours. Cells were lysed in RIPA lysis buffer and processed as described

previously (22) and the fusion protein of HMPV was immunoprecipitated using anti-HMPV F 54G10

monoclonal antibody (John Williams, U. Pitt). Samples were run on a 15% SDS-PAGE and visualized using

the typhoon imaging system. Band densitometry was conducted using ImageQuant software (GE).

SPINT2 cleavage inhibition by Western blot and cell-cell fusion assay

Experiments were performed as previously described with minor modifications (26). Treatment

with trypsin, recombinant matriptase and KLK5 were conducted as described (19, 63). For western blot

analysis 293T cells were transfected with Turbofect (Invitrogen) according to manufacturer's instructions.

Cell-cell fusion assays were conducted with VERO cells that were transfected with Lipofectamine

according to the manufacturer's instructions. Antibodies to detect A/CA/04/09 (H1N1) HA (NR28666),

A/Aichi/2/68 (H3N2) HA (NR3118),A/Shanghai/2/2013 (H7N9) HA (NR48765) and A/Vietnam/1204/2004

(H5N1) HA (NR2705) were obtained from the Biodefense and Emerging Infections Research Resources

Repository. Respective secondary antibodies had an Alexa488 tag (Invitrogen). Western blot membranes

were scanned using a ChemiDoc imaging system (Bio-Rad). For quantification the pixel intensity of the

individual HA_1 bands was measured using ImageJ software and cleavage efficiencies were calculated by

the following equation: HA_1 10 nM or 500 nM SPINT2/HA_1 0nM SPINT2 x 100%. Cell-cell fusion assays

were carried out as described (19).

Inhibition of influenza viral infection in cell culture

MDCK cells were seeded to a confluency of about 70% in 6-well plates. One plate each was then

transformed with a plasmid allowing for the expression of human matriptase or human TMPRSS2. One

plate was transformed with empty vector. 18 hours post transfection cells were infected with the

respective egg-grown virus at a MOI of approx. 0.1. Different SPINT2 concentrations were added as

indicated. 0.8 nM trypsin was added to the cells transformed with the empty vector. 48 hours post

infection supernatants were collected, centrifuged and stored at -80°C. Viral titers were determined using

an immune-plaque assay as described (64).

Inhibition of HMPV viral infection in cell culture

VERO (200,000) cells were plated into 24-well plates. The following day, cells were infected with

MOI 1 rgHMPV for 3 hours. Cells were washed with PBS and 500µL OPTI-mem with or without 500nM

SPINT2 and 0.3µg/mL of TPCK-trypsin was added and incubated for up to 96 hours. The SPINT2 and trypsin

was replenished in new OPTI-mem every 24 hours. For each time point, media was aspirated and 100µL

of OPTI-mem was added to cells followed by scraping and flash freezing. These samples were then titered

on confluent VERO cells up to 10^{-6} dilution to calculate viral titer. Graph shows 4 independent replicates

with internal duplicates plotted as individual points. Some data points not shown are due to sample loss,

with a minimum of 6 points per group or 3 independent replicates.

Acknowledgements

J.T.K. and R.E.D. thank Dr. John Williams from University of Pittsburgh Children's Hospital for

kindly providing the 54G10 anti-HMPV F antibody. This work was funded by National Institutes of Health

grant 5R21AI117300.

References

1.   World Health Organization. 2014. Global Epidemiological Surveillance Standards for Influenza.

https://www.who.int/influenza/resources/documents/WHO_Epidemiological_Influenza_Surveill

ance_Standards_2014.pdf.

2.   Fouchier RAM, Guan Y. 2013. Ecology and evolution of influenza viruses in wild and domestic

birds. Textb Influ 173–189.

3.   Yen H, Webster RG. 2009. Pandemic Influenza as a Current Threat, p. 3–24. *In* Current Topics in

Microbiology and Immunology.

4.   World Health Organization. 2017. WHO | Human infection with avian influenza A(H7N9) virus –

China. https://www.who.int/csr/don/26-october-2017-ah7n9-china/en/ http://www.

5.   World Health Organization. 2018. Influenza fact sheet. https://www.who.int/news-room/fact-

sheets/detail/i https://www.who.int/news-room/fact-sheets/detail/i.

6.   Kilbourne ED. 2006. Influenza pandemics of the 20th century. Emerg Infect Dis 12:9–14.

7.   Hussain M, Galvin HD, Haw TY, Nutsford AN, Husain M. 2017. Drug resistance in influenza a virus:

The epidemiology and management. Infect Drug Resist 10:121–134.

8.   Takashita E, Morita H, Ogawa R, Nakamura K, Fujisaki S, Shirakura M, Kuwahara T, Kishida N,

Watanabe S, Odagiri T. 2018. Susceptibility of influenza viruses to the novel cap-dependent

endonuclease inhibitor baloxavir marboxil. Front Microbiol 9.

9.   Hayden FG, Sugaya N, Hirotsu N, Lee N, De Jong MD, Hurt AC, Ishida T, Sekino H, Yamada K,

Portsmouth S, Kawaguchi K, Shishido T, Arai M, Tsuchiya K, Uehara T, Watanabe A. 2018.

Baloxavir marboxil for uncomplicated influenza in adults and adolescents. N Engl J Med 379:913–

923.

10.   Omoto S, Speranzini V, Hashimoto T, Noshi T, Yamaguchi H, Kawai M, Kawaguchi K, Uehara T,

Shishido T, Naito A, Cusack S. 2018. Characterization of influenza virus variants induced by

treatment with the endonuclease inhibitor baloxavir marboxil. Sci Rep 8.

11.   Taylor S, Lopez P, Weckx L, Borja-Tabora C, Ulloa-Gutierrez R, Lazcano-Ponce E, Kerdpanich A,

Angel Rodriguez Weber M, Mascareñas de Los Santos A, Tinoco J-C, Safadi MAP, Lim FS,

Hernandez-de Mezerville M, Faingezicht I, Cruz-Valdez A, Feng Y, Li P, Durviaux S, Haars G, Roy-

Ghanta S, Vaughn DW, Nolan T. 2017. Respiratory viruses and influenza-like illness: Epidemiology

and outcomes in children aged 6 months to 10 years in a multi-country population sample. J

Infect 74:29–41.

12.   Li Y, Reeves RM, Wang X, Bassat Q, Brooks WA, Cohen C, Moore DP, Nunes M, Rath B, Campbell

H, Nair H, Acacio S, Alonso WJ, Antonio M, Ayora Talavera G, Badarch D, Baillie VL, Barrera-

Badillo G, Bigogo G, Broor S, Bruden D, Buchy P, Byass P, Chipeta J, Clara W, Dang D-A, de Freitas

Lázaro Emediato CC, de Jong M, Díaz-Quiñonez JA, Do LAH, Fasce RA, Feng L, Ferson MJ, Gentile

A, Gessner BD, Goswami D, Goyet S, Grijalva CG, Halasa N, Hellferscee O, Hessong D, Homaira N,

Jara J, Kahn K, Khuri-Bulos N, Kotloff KL, Lanata CF, Lopez O, Lopez Bolaños MR, Lucero MG,

Lucion F, Lupisan SP, Madhi SA, Mekgoe O, Moraleda C, Moyes J, Mulholland K, Munywoki PK,

Naby F, Nguyen TH, Nicol MP, Nokes DJ, Noyola DE, Onozuka D, Palani N, Poovorawan Y, Rahman

M, Ramaekers K, Romero C, Schlaudecker EP, Schweiger B, Seidenberg P, Simoes EAF, Singleton

R, Sistla S, Sturm-Ramirez K, Suntronwong N, Sutanto A, Tapia MD, Thamthitiwat S, Thongpan I,

Tillekeratne G, Tinoco YO, Treurnicht FK, Turner C, Turner P, van Doorn R, Van Ranst M, Visseaux

B, Waicharoen S, Wang J, Yoshida L-M, Zar HJ. 2019. Global patterns in monthly activity of

influenza virus, respiratory syncytial virus, parainfluenza virus, and metapneumovirus: a

systematic analysis. Lancet Glob Heal 7:e1031–e1045.

13.   Kahn JS. 2006. Epidemiology of Human Metapneumovirus. Clin Microbiol Rev 19:546–557.

14.   van den Hoogen BG, de Jong JC, Groen J, Kuiken T, de Groot R, Fouchier RAM, Osterhaus ADME.

4   2001. A newly discovered human pneumovirus isolated from young children with respiratory

5   tract disease. Nat Med 7:719–724.

6   15.   Leung J, Esper F, Weibel C, Kahn JS. 2005. Seroepidemiology of Human Metapneumovirus (hMPV)

7   on the Basis of a Novel Enzyme-Linked Immunosorbent Assay Utilizing hMPV Fusion Protein

8   Expressed in Recombinant Vesicular Stomatitis Virus. J Clin Microbiol 43:1213–1219.

16.   Steinhauer D. 1999. Role of hemagglutinin cleavage for the pathogenicity of influenza virus.

Virology 258:1–20.

17.   Taubenberger JK. 1998. Influenza virus hemagglutinin cleavage into HA1, HA2: no laughing

matter. Proc Natl Acad Sci U S A 95:9713–9715.

18.   Böttcher-Friebertshäuser E, Klenk HD, Garten W. 2013. Activation of influenza viruses by

proteases from host cells and bacteria in the human airway epithelium. Pathog Dis 69:87–100.

19.   Hamilton BS, Whittaker GR. 2013. Cleavage activation of human-adapted influenza virus subtypes

by kallikrein-related peptidases 5 and 12. J Biol Chem 288:17399–17407.

20.   Horimoto T, Nakayama K, Smeekens SP, Kawaoka Y. 1994. Proprotein-processing endoproteases

PC6 and furin both activate hemagglutinin of virulent avian influenza viruses. J Virol 68:6074–

6078.

21.   Shirogane Y, Takeda M, Iwasaki M, Ishiguro N, Takeuchi H, Nakatsu Y, Tahara M, Kikuta H, Yanagi

Y. 2008. Efficient Multiplication of Human Metapneumovirus in Vero Cells Expressing the

Transmembrane Serine Protease TMPRSS2. J Virol 82:8942–8946.

22.   Schowalter RM, Smith SE, Dutch RE. 2006. Characterization of Human Metapneumovirus F

Protein-Promoted Membrane Fusion: Critical Roles for Proteolytic Processing and Low pH. J Virol

80:10931–10941.

23.   Millet JK, Whittaker GR. 2014. Host cell proteases: Critical determinants of coronavirus tropism

and pathogenesis. Virus Res.

24.   Laporte M, Naesens L. 2017. Airway proteases: an emerging drug target for influenza and other

respiratory virus infections. Curr Opin Virol 24:16–24.

25.   Zhirnov OP, Matrosovich TY, Matrosovich MN, Klenk H. 2011. Short communication Aprotinin , a

protease inhibitor , suppresses proteolytic activation of pandemic H1N1v influenza virus

174:169–174.

26.   Hamilton BS, Chung C, Cyphers SY, Rinaldi VD, Marcano VC, Whittaker GR. 2014. Inhibition of

influenza virus infection and hemagglutinin cleavage by the protease inhibitor HAI-2. Biochem

Biophys Res Commun 450:1070–1075.

27.   Szabo R, Hobson JP, List K, Molinolo A, Lin C-Y, Bugge TH. 2008. Potent inhibition and global co-

localization implicate the transmembrane Kunitz-type serine protease inhibitor hepatocyte

growth factor activator inhibitor-2 in the regulation of epithelial matriptase activity. J Biol Chem

283:29495–504.

28.   Wu SR, Teng CH, Tu YT, Ko CJ, Cheng TS, Lan SW, Lin HY, Lin HH, Tu HF, Hsiao PW, Huang HP,

Chen CH, Lee MS. 2017. The Kunitz Domain i of Hepatocyte Growth Factor Activator Inhibitor-2

Inhibits Matriptase Activity and Invasive Ability of Human Prostate Cancer Cells. Sci Rep 7:1–19.

29.   Wu SR, Lin CH, Shih HP, Ko CJ, Lin HY, Lan SW, Lin HH, Tu HF, Ho CC, Huang HP, Lee MS. 2019.

HAI-2 as a novel inhibitor of plasmin represses lung cancer cell invasion and metastasis. Br J

Cancer 120:499–511.

30.   Roversi FM, Olalla Saad ST, Machado-Neto JA. 2018. Serine peptidase inhibitor Kunitz type 2

(SPINT2) in cancer development and progression. Biomed Pharmacother 101:278–286.

31.   Delaria KA, Muller DK, Marlor CW, Brown JE, Das RC, Roczniak SO, Tamburini PP. 1997.

Characterization of Placental Bikunin, a Novel Human Serine Protease Inhibitor. J Biol Chem 272:12209–12214.

32. Fukai K, Yokosuka O, Chiba T, Hirasawa Y, Tada M, Imazeki F, Kataoka H, Saisho H. 2003. Hepatocyte growth factor activator inhibitor 2/placental bikunin (HAI-2/PB) gene is frequently hypermethylated in human hepatocellular carcinoma. Cancer Res 63:8674–9.

33. Dong W, Chen X, Xie J, Sun P, Wu Y. 2010. Epigenetic inactivation and tumor suppressor activity of HAI-2/SPINT2 in gastric cancer. Int J cancer 127:1526–34.

34. Tsai C-H, Teng C-H, Tu Y-T, Cheng T-S, Wu S-R, Ko C-J, Shyu H-Y, Lan S-W, Huang H-P, Tzeng S-F, Johnson MD, Lin C-Y, Hsiao P-W, Lee M-S. 2014. HAI-2 suppresses the invasive growth and metastasis of prostate cancer through regulation of matriptase. Oncogene 33:4643–52.

35. Hwang S, Kim H-E, Min M, Raghunathan R, Panova IP, Munshi R, Ryu B. 2015. Epigenetic Silencing of SPINT2 Promotes Cancer Cell Motility via HGF-MET Pathway Activation in Melanoma. J Invest Dermatol 135:2283–2291.

36. Szabo R, Hobson JP, Christoph K, Kosa P, List K, Bugge TH. 2009. Regulation of cell surface protease matriptase by HAI2 is essential for placental development, neural tube closure and embryonic survival in mice. Development 136:2653–2663.

37. Wu C-J, Feng X, Lu M, Morimura S, Udey MC. 2017. Matriptase-mediated cleavage of EpCAM destabilizes claudins and dysregulates intestinal epithelial homeostasis. J Clin Invest 127:623–634.

38. Jaimes JA, Millet JK, Goldstein ME, Whittaker GR, Straus MR. 2019. A Fluorogenic Peptide Cleavage Assay to Screen for Proteolytic Activity: Applications for coronavirus spike protein activation. J Vis Exp.

39. Straus MR, Whittaker GR. 2017. A peptide-based approach to evaluate the adaptability of influenza A virus to humans based on its hemagglutinin proteolytic cleavage site. PLoS One

1   12:e0174827.

2   40.   Biacchesi S, Pham QN, Skiadopoulos MH, Murphy BR, Collins PL, Buchholz UJ. 2006. Modification

3   of the trypsin-dependent cleavage activation site of the human metapneumovirus fusion protein

4   to be trypsin independent does not increase replication or spread in rodents or nonhuman

5   primates. J Virol 80:5798–806.

6   41.   Galloway SE, Reed ML, Russell CJ, Steinhauer DA. 2013. Influenza HA Subtypes Demonstrate

7   Divergent Phenotypes for Cleavage Activation and pH of Fusion: Implications for Host Range and

8   Adaptation. PLoS Pathog 9:e1003151.

42.   Hatesuer B, Bertram S, Mehnert N, Bahgat MM, Nelson PS, Pöhlman S, Schughart K. 2013.

Tmprss2 Is Essential for Influenza H1N1 Virus Pathogenesis in Mice. PLoS Pathog 9:1–10.

43.   Sakai K, Ami Y, Tahara M, Kubota T, Anraku M, Abe M, Nakajima N, Sekizuka T, Shirato K, Suzaki

Y, Ainai A, Nakatsu Y, Kanou K, Nakamura K, Suzuki T, Komase K, Nobusawa E, Maenaka K, Kuroda

M, Hasegawa H, Kawaoka Y, Tashiro M, Takeda M. 2014. The host protease TMPRSS2 plays a

major role in in vivo replication of emerging H7N9 and seasonal influenza viruses. J Virol

88:5608–16.

44.   Baron J, Tarnow C, Mayoli-Nüssle D, Schilling E, Meyer D, Hammami M, Schwalm F, Steinmetzer

T, Guan Y, Garten W, Klenk H-D, Böttcher-Friebertshäuser E. 2012. Matriptase, HAT, and

TMPRSS2 Activate the Hemagglutinin of H9N2 Influenza A Viruses. J Virol 87:1811–1820.

45.   Kühn N, Bergmann S, Kösterke N, Lambertz RLO, Keppner A, van den Brand JMA, Pöhlmann S,

Weiß S, Hummler E, Hatesuer B, Schughart K. 2016. The Proteolytic Activation of (H3N2)

Influenza A Virus Hemagglutinin Is Facilitated by Different Type II Transmembrane Serine

Proteases. J Virol 90:4298–4307.

46.   Houser K, Subbarao K. 2015. Influenza vaccines: challenges and solutions. Cell Host Microbe

17:295–300.

47.   Karlsson EA, Meliopoulos VA, van de Velde NC, van de Velde L-A, Mann B, Gao G, Rosch J,

2   Tuomanen E, McCullers J, Vogel P, Schultz-Cherry S. 2017. A Perfect Storm: Increased

3   Colonization and Failure of Vaccination Leads to Severe Secondary Bacterial Infection in Influenza

4   Virus-Infected Obese Mice. MBio 8.

5   48.   Straus MR, Whittaker GR. 2017. A peptide-based approach to evaluate the adaptability of

6   influenza A virus to humans based on its hemagglutinin proteolytic cleavage site. PLoS One

7   12:e0174827.

8   49.   Limburg H, Harbig A, Bestle D, Stein DA, Moulton HM, Jaeger J, Janga H, Hardes K, Koepke J,

9   Schulte L, Koczulla AR, Schmeck B, Klenk H-D, Böttcher-Friebertshäuser E. 2019. TMPRSS2 Is the

10   Major Activating Protease of Influenza A Virus in Primary Human Airway Cells and Influenza B

11   Virus in Human Type II Pneumocytes. J Virol 93.

12   50.   Sakai K, Ami Y, Tahara M, Kubota T, Anraku M, Abe M, Nakajima N, Sekizuka T, Shirato K, Suzaki

13   Y, Ainai A, Nakatsu Y, Kanou K, Nakamura K, Suzuki T, Komase K, Nobusawa E, Maenaka K, Kuroda

14   M, Hasegawa H, Kawaoka Y, Tashiro M, Takeda M. 2014. The Host Protease TMPRSS2 Plays a

15   Major Role in In Vivo Replication of Emerging H7N9 and Seasonal Influenza Viruses. J Virol

16   88:5608–5616.

17   51.   Tarnow C, Engels G, Arendt A, Schwalm F, Sediri H, Preuss A, Nelson PS, Garten W, Klenk H-D,

18   Gabriel G, Bottcher-Friebertshauser E. 2014. TMPRSS2 Is a Host Factor That Is Essential for

19   Pneumotropism and Pathogenicity of H7N9 Influenza A Virus in Mice. J Virol 88:4744–4751.

20   52.   Whittaker GR, Straus MR. 2019. Human matriptase/ST 14 proteolytically cleaves H7N9

21   hemagglutinin and facilitates the activation of influenza A/Shanghai/2/2013 virus in cell culture.

22   Influenza Other Respi Viruses irv.12707.

23   53.   Sielaff F, Böttcher-Friebertshäuser E, Meyer D, Saupe SM, Volk IM, Garten W, Steinmetzer T.

24   2011. Development of substrate analogue inhibitors for the human airway trypsin-like protease

HAT. Bioorganic Med Chem Lett 21:4860–4864.

54. Meyer D, Sielaff F, Hammami M, Böttcher-Friebertshäuser E, Garten W, Steinmetzer T. 2013.
    Identification of the first synthetic inhibitors of the type II transmembrane serine protease
    TMPRSS2 suitable for inhibition of influenza virus activation. Biochem J 452:331–343.

55. Yamaya M, Shimotai Y, Hatachi Y, Lusamba Kalonji N, Tando Y, Kitajima Y, Matsuo K, Kubo H,
    Nagatomi R, Hongo S, Homma M, Nishimura H. 2015. The serine protease inhibitor camostat
    inhibits influenza virus replication and cytokine production in primary cultures of human tracheal
    epithelial cells. Pulm Pharmacol Ther 33:66–74.

56. Sai JK, Suyama M, Kubokawa Y, Matsumura Y, Inami K, Watanabe S. 2010. Efficacy of camostat
    mesilate against dyspepsia associated with non-alcoholic mild pancreatic disease. J Gastroenterol
    45:335–341.

57. Zhou Y, Vedantham P, Lu K, Agudelo J, Carrion R, Nunneley JW, Barnard D, Pöhlmann S,
    McKerrow JH, Renslo AR, Simmons G. 2015. Protease inhibitors targeting coronavirus and
    filovirus entry. Antiviral Res 116:76–84.

58. Matsuyama S, Nagata N, Shirato K, Kawase M, Takeda M, Taguchi F. 2010. Efficient Activation of
    the Severe Acute Respiratory Syndrome Coronavirus Spike Protein by the Transmembrane
    Protease TMPRSS2. J Virol 84:12658–12664.

59. Gierer S, Bertram S, Kaup F, Wrensch F, Heurich A, Kramer-Kuhl A, Welsch K, Winkler M, Meyer B,
    Drosten C, Dittmer U, von Hahn T, Simmons G, Hofmann H, Pohlmann S. 2013. The Spike Protein
    of the Emerging Betacoronavirus EMC Uses a Novel Coronavirus Receptor for Entry, Can Be
    Activated by TMPRSS2, and Is Targeted by Neutralizing Antibodies. J Virol 87:5502–5511.

60. JH B, J F, AM H, FG H, R H, MD de J, S L, TK N, TH N, TH T, A N, S T, KY Y. 2005. Avian influenza
    (H5N1) infection in humans. N Engl J Med 353:1374–85.

61. Horimoto T, Nakayama K, Smeekens SP, Kawaoka Y. 1994. Proprotein-processing endoproteases

PC6 and furin both activate hemagglutinin of virulent avian influenza viruses. J Virol 68:6074–8.

62.   Kinder JT, Klimyte EM, Chang A, Williams J V, Dutch RE. 2019. Human metapneumovirus fusion

protein triggering: Increasing complexities by analysis of new HMPV fusion proteins. Virology

531:248–254.

63.   Hamilton BS, Gludish DWJ, Whittaker GR. 2012. Cleavage Activation of the Human-Adapted

Influenza Virus Subtypes by Matriptase Reveals both Subtype and Strain Specificities. J Virol

86:10579–10586.

64.   Tse L V, Marcano VC, Huang W, Pocwierz MS, Whittaker GR. 2013. Plasmin-mediated activation

of pandemic H1N1 influenza virus hemagglutinin is independent of the viral neuraminidase. J

Virol 87:5161–9.

Figure Legends

Figure 1: Cytotoxicity assay to evaluate the cytotoxic effect of SPINT2.

293T cells were incubated with indicated SPINT2 concentrations for 24 hours. DMEM and 500 μM H_2O_2

served as controls. After 24 hours cell viability was determined via a spectrophotometric assay.

Figure 2: SPINT2 inhibits cleavage of HA protein expressed in 293T cells. Cells were transfected with

plasmids encoding for the indicated HA and allowed to express the protein for ~18 hours. The recombinant

proteases were incubated for 15 minutes with the indicated SPINT2 concentrations and subsequently

added to the cells for 10 minutes (trypsin) or 90 minutes (matriptase and KLK5). Western blots were

performed and the HA_1 band was quantified using ImageJ. (A) Quantification of the HA1 band comparing

the signal intensity of the 0 nM SPINT2 samples against 10 nM and 500 nM SPINT2 of the respective

HA/protease combination. Three independent experiments were carried out and the western blots of

each experiment were analyzed. Quantifications were conducted as described in the methods section. (B

– D) Western blots showing the cleavage of (B) A/CA/04/09 H1N1 HA by matriptase and KLK5 at different

SPINT2 concentration, (C) A/Aichi/2/68 H3N2 HA by KLK5 at different SPINT2 concentration and (D)

A/Shanghai/2/2013 H7N9 HA by matriptase and KLK5 at different SPINT2 concentrations. Statistical

analysis was performed using a non-paired student's t-test comparing the samples tested with 10 nM

SPINT2 against the respective sample incubated with 500 nM SPINT2. Error bars indicate standard

deviation. * indicates p = < 0.05.

Figure 3: TMPRSS2, HAT, matriptase and KLK5 cleave HMPV F and SPINT2 is able to prevent cleavage by

exogenous proteases. HMPV F was either expressed alone or co-transfected with protease and allowed

to express for ~ 18hours. Cells were then metabolically starved of cysteine and methionine followed by

radioactive S35 labeling of protein for 4 hours in the presence of TPCK-trypsin or specified protease.

SPINT2 treated proteases were incubated at room temperature for 10 minutes and placed onto cells for

4 hours. Radioactive gels were quantified using ImageQuant software with percent cleavage equal

to $\left[\left(\frac{F_1}{F_0+F_1}\right)x100\right]$. A and B) Co-transfected proteases TMPRSS2, HAT and matriptase are able to cleave

HMPV F (n=4) while C and D) exogenous proteases KLK5 and matriptase but not KLK12 are able to cleave

HMPV F (n=5). E and F) SPINT2 prevented cleavage of HMPV F by trypsin, KLK5 and matriptase at nm

concentrations demonstrated by the loss of the F_1 cleavage product (n=3). Statistical analysis was

performed using a one-way ANOVA followed by a student's t-test with a bonferroni multiple comparisons

test correction. P<0.05 *, P<0.005 **, P<0.005 ***, P<0.001 ****. N values represent independent

replicates for each treatment group. Error bars represent SD.

Figure 4: SPINT2 inhibits HA-mediated cell-cell fusion. VERO cells were transfected with plasmids

encoding for (A) A/CA/04/09 H1N1 HA or (B) A/Shanghai/2/2013 H7N9 HA and allowed to express the

protein for ~18 hours. Recombinant matriptase and KLK5 were incubated with different SPINT2

concentrations for 15 minutes and then added to the HA-expressing cells for 3 hours. After 3 hours the

cells were briefly treated with cell fusion buffer at pH5, washed, supplemented with growth medium and

returned to the incubator for 1 hour to allow fusion. HA protein was detected using HA-specific primary

antibodies and a secondary fluorogenic Alexa488 antibody. Nuclei were stained using DAPI. Magnification

40x. (C) VERO cells expressed A/Vietnam/1204/2004 H5N1 HA that was cleaved during its maturation

process in the cell. SPINT2 was added at 0 nM or 500 nM at the time of transfection. Magnification 25x.

Figure 5: SPINT2 reduces IAV growth in cell culture. MDCK cells were transfected with plasmids encoding

for human matriptase or human TMPRSS2 and allowed to express the proteins for ~18 hours. Cells

expressing human matriptase (B) or human TMPRSS2 (C) were then infected with A/CA/04/09 H1N1 at a

MOI of 0.1 and different SPINT2 concentration were added to each well. Non-transfected cells to which

trypsin was added served as a control (A). (D) MDCK cells were infected with A/X31 H3N2 at an MOI of

0.1 and trypsin was added to assist viral propagation. Different SPINT2 concentration were added as

indicated. After 48 hours of infection the supernatants were collected and used for an immuno-plaque

assay to determine the viral loads. Experiment was repeated three times and each dot represents the viral

titer of a single experiment. Statistical analysis was performed using a non-paired student's t-test

comparing the control (0 h) against the respective sample. Error bars indicate standard deviation. *

indicates p = < 0.05. Extended horizontal line within the error bars represents mean value of the three

independent replicates.

Figure 6: SPINT2 mitigates the spread of HMPV in cell culture. VERO cells were infected with a MOI 1 of

rgHMPV. SPINT2 and TPCK-trypsin were added at 500nM and 0.3µg/mL respectively and spread was

monitored up to 96hpi. Cells not treated with SPINT2 demonstrated significant spread up through 96hpi

whereas treated samples did not show any detectable virus up to 48hpi. However, there was minimal

detected virus at 72 and 96hpi. Error bars indicate standard deviation of 4 independent samples

completed in duplicate (all data points plotted within the graph). Statistical analysis was performed using

a student's t-test. P <0.05 *, P <0.005 **, P <0.005 ***, P <0.001 ****. ND indicates that the sample was

below the limit of detection.

Figure7: SPINT2 reduces viral growth when added 24 hours post infection. MDCK cells were infected

with A/CA/04/09 H1N1 at a MOI of 0.1 and trypsin was added. 500nM SPINT2 were added at the time of

infection or 24 hours post infection. Supernatants were collected 48 hours post infection and used for an

immuno-plaque assay to determine the viral titers. Statistical analysis was performed using a non-paired

student's t-test comparing the control (0 h) against the respective sample. Error bars indicate standard

deviation. * indicates p = < 0.001. Extended horizontal line within the error bars represents mean value

of the three independent replicates.

Supplementary Figure 1: Recombinant furin was incubated with a fluorogenic peptide mimicking the

HPAI cleavage site of H5N1 with no SPINT2 present or in the presence of 500 nM SPINT2. Cleavage was

monitored by the increase of fluorescence at 390 nm and Vmax values were plotted. n = 3. Error bars

represent standard deviation.

Supplementary Figure 2: Fluorogenic peptides mimicking the cleavage sites (A) A/CA/04/09 H1N1, (B)

A/Japan/305/1957 H2N2 HA, (C) A/Aichi/2/68 H3N2 HA, (D) A/Vietnam/1203/2004 H5N1 LPAI HA, (E)

A/Vietnam/1204/2004 H5N1 HPAI HA, (F) A/Taiwan/2/2013 H6N1 HA, (G) A/Shanghai/2/2013 H7N9 HA,

(H) A/Hong Kong/2108/2003 H9N2 HA and (I) HMPV F were incubated with the indicated proteases and

different SPINT2 concentrations. Cleavage was monitored by the increase of fluorescence at 390 nm and

the resulting Vmax values were plotted against the different SPINT2 concentrations on a logarithmic scale

(x-axis). Prism 7 software was then used to calculate the IC50 values based on the graphs shown in this

figure as described in the "Material and Methods" section. In some cases, the error bars are shorter than

the height of the symbol and therefore are not displayed.

Table 1: Cleavage profile of the HMPV F: A fluorogenic peptide mimicking the cleavage site of HMPV F

was incubated with the indicated proteases and cleavage was monitored by the increase of fluorescence

at 390 nm. RFU = relative fluorescence units. StDev = Standard Deviation.

Table 2: IC50 values of all protease/peptide combinations. : Fluorogenic peptides mimicking the cleavage

sites of A/CA/04/09 H1N1, A/Japan/305/1957 H2N2 HA, A/Aichi/2/68 H3N2 HA, A/Vietnam/1203/2004

H5N1 LPAI HA, A/Vietnam/1204/2004 H5N1 HPAI HA, A/Taiwan/2/2013 H6N1 HA, A/Shanghai/2/2013

H7N9 HA, A/Hong Kong/2108/2003 H9N2 HA and HMPV F were incubated with the indicated proteases

and different SPINT2 concentrations. Cleavage was monitored by the increase of fluorescence at 390 nm

and the resulting Vmax values were used to calculate the IC50 values as described in the "Material and

Methods" section. (A) IC50 values of influenza A fluorogenic cleavage site peptide mimics. Concentrations

are in nanomolar. (B) IC50 values of the HMPV F cleavage site peptide mimic. Concentrations are in

picomolar. NT = Not tested.

Table 3: Viral titers measured in the infection studies: Table shows the average viral titers and standard

deviation calculated from the 3 independent biological replicates depicted in Figures 6 and 7. (A) Titers

and standard deviation from infection studies shown in Figure 5. (B) Titers and standard deviation shown

in Figure 7.

1 **Table 1**

	Vmax (RFU/min)	StDev
Trypsin	135.25	17.55
Matriptase	9.24	0.39
KLK5	5.80	0.12
HAT	2.99	0.41
KLK12	0.00	0.00
Plasmin	37.86	2.07

3 **Table 2**

4 **A**

	IC50 values (nM)					
	Trypsin	Matriptase	HAT	KLK5	KLK12	Plasmin
H1N1 HA	70.57	25.00	24.08	28.34	11.70	122.10
H2N2 HA	155.40	NT	34.74	8.27	6.88	NT
H3N2 HA	207.90	NT	15.33	9.54	NT	106.00
H5N1 HPAI HA	1135.00	194.10	NT	NT	NT	1166.00
H5N1 LPAI HA	145.80	5.32	23.56	6.38	3.66	160.10
H6N1 HA	30.52	NT	NT	29.93	14.04	NT
H7N9 HA	20.97	8.00	NT	NT	NT	77.59
H9N2 HA	99.16	9.02	12.74	11.79	NT	134.00

6 **B**

	IC50 values (nM)					
	Trypsin	Matriptase	HAT	KLK5	KLK12	Plasmin
HMPV F	0.04	0.0003	NT	0.95	NT	NT

Table 3

A

SPINT2	Trypsin - H1N1		Matriptase - H1N1		TMPRSS2 - H1N1		Trypsin - X31	
(nM)	Average pfu/ml	StDev	Average pfu/ml	StDev	Average pfu/ml	StDev	Average pfu/ml	StDev
0	1.99E+08	3.91E+07	1.95E+09	4.39E+08	4.90E+08	1.16E+08	9.47E+06	9.02E+05
10	1.14E+08	3.70E+07	1.73E+09	2.95E+08	2.93E+08	7.57E+06	7.80E+06	1.04E+06
50	5.93E+07	3.06E+06	9.24E+08	4.16E+08	2.08E+08	6.16E+07	4.27E+06	1.40E+06
150	4.60E+07	8.72E+06	5.09E+08	2.44E+08	7.80E+07	4.65E+07	3.27E+06	1.62E+06
500	1.40E+07	5.29E+06	9.53E+07	3.43E+07	1.40E+07	5.29E+06	8.00E+05	5.29E+05

5 B

SPINT2	Trypsin - H1N1	
(nM)	Average pfu/ml	StDev
0	1.88E+08	1.06E+07
500	2.80E+07	1.00E+07
500 24hpi	4.07E+07	2.00E+07

1   **Figure 1**

     Figure 2

1   **Figure 3**

1   **Figure 4**

1   **Figure 5**

A

B

C

D

1   **Figure 6**

1   **Figure 7**

1   **Supplementary Figure 1**

1   **Supplementary Figure 2**

C

D

E

F

G

H

I