Population-level sensitivity to landscape variables reflects individual-based habitat selection in a woodland bat species

Pierre-Loup Jan[1,2,*,†], Diane Zarzoso-Lacoste[3,4,*], Damien Fourcy[1], Alice Baudouin[3], Olivier Farcy[5], Josselin Boireau[6], Pascaline Le Gouar[3], Sébastien J. Puechmaille[7,8], Eric J. Petit[1]

[1] UMR 985 ESE, INRA, Agrocampus Ouest, 35042 Rennes cedex, France

[2] UMR 7372 CEBC, CNRS, La Rochelle Université, 79360 Villiers-en-Bois, France

[3] UMR 6553 ECOBIO, Université Rennes 1, Campus de Beaulieu, 35042 Rennes cedex, France

[4] UMR 8079 ESE, Université Paris-Sud/CNRS/AgroParisTech, Université Paris-Saclay, 91405 Orsay Cedex, France

[5] Bretagne Vivante, 29221 Brest cedex 2, France

[6] Groupe Mammalogique Breton, 29450 Sizun, France

[7] Zoological Institute and Museum, Greifswald University, 17489 Greifswald, Germany

[8] ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France.

* Co-first-authors

† Corresponding author : pierreloup.jan@gmail.com

Abstract

Understanding the relationship between habitat quality and population dynamics is fundamental for long-term management and range predictions in ecology. However, habitat suitability is generally only investigated at the individual scale, as it is the case for the lesser horseshoe bat (*Rhinolophus hipposideros*), a species of conservation concern. Using a statistical modelling approach and census data of 94 lesser horseshoe bat colonies located in Brittany (France), we analysed the effect of landscape composition and configuration on the demography of surveyed maternity colonies (i.e. colony size, fecundity and growth rate), and compared our result to those provided by individual-based studies. Our results validated that the landscape in a 500-meter buffer around colonies (core foraging area) is crucial for population size and dynamics, and confirmed the positive influence of broadleaved woodland proportion on bat colony size. We revealed a positive effect of lakeshores and riverbanks on colony size and growth rate, underlying the importance of these habitats for the long-term conservation of this non-migratory forest species. Importantly, our results refine previous knowledge concerning the threat posed by the intensification of human activities (e.g. urbanization, agriculture, habitat fragmentation), and highlight the negative effect of large and regular patches of artificial and crop lands and of open land patches shape complexity on all demographic variables investigated. Finally, by confirming and specifying the conclusions of previous individual level studies about lesser horseshoe bats' habitat selection and quality requirements, our results support the dependence of population dynamics and associated conservation management to individual behaviour and sensitivity.

Keywords: *Rhinolophus hipposideros*, population dynamics, habitat requirements, landscape composition and configuration.

1. Introduction

Better understanding the relationship between population dynamics and habitat quality is fundamental both for predicting range shifts under global change and for establishing sound conservation strategies (Elith and Leathwick, 2009; Hodgson et al., 2009). Recently, Species Distribution Modelling (SDMs) has become the standard to inform scenarios of biodiversity under global change (e.g. climate and land use changes) by correlating species occurrence with eco-geographical information related to species climate and/or habitat niche (Elith and Leathwick, 2009). These models, however, do not explicitly consider the demographic processes (e.g. fecundity or survival) relating population-level metrics (e.g. presence, density, growth rate, carrying capacity) to habitat quality (Thuiller et al., 2013), while they are essential for species to follow environmental conditions favourable to their survival (species range dynamics- Aitken et al., 2008; Auffret et al., 2017; Gaillard et al., 2010; Kokko and López-Sepulcre, 2006). Therefore, research questions that target population-level processes should use metrics that relate to population dynamic parameters (Gaillard et al., 2010).

Yet, studies that directly evaluate relationships between population-level metrics and habitat quality are seldom both because demographic data are not often gathered over many populations (Gurevitch et al., 2016) and because habitat quality is too rarely assessed where population data are collected (Pulliam, 2000). If population-level species-habitat relationships are so rarely documented, where does our basic ecological knowledge come from? Historically, from the study of resource selection by individuals (Manly et al., 2007).

The fact that population-level studies of habitat suitability are not frequent has triggered evaluations of the necessary conditions for using one metric (i.e. population-level metrics) as a proxy for another (i.e. habitat suitability - Thuiller et al., 2014). For instance, though neither species occurrence neither resource selection by individual directly relates population-level dynamics to environmental conditions, both have been used to predict abundance (Acevedo et al., 2017; Boyce et al., 2016; Street et al., 2016). Studies relating population-level metrics and processes to habitat quality are needed to test their validity as surrogates for inferring this relationship. More precisely, scaling up resource selection by individuals to abundances is informative, at least when environmental variables that are important for variations in population sizes were included in the study of resource use (Boyce et al., 2016; Street et al., 2016). In general, establishing in which conditions environmental variables that come out of individual-based studies are also relevant at the population-scale would give more weight to conservation actions based on data gathered at the individual level, and enlarge the range of studies that can feed more realistic predictions of species range shifts.

The lesser horseshoe bat, *Rhinolophus hipposideros,* has undergone a massive decline in the northern part of its distribution range during the last century for reasons that are still not fully understood, although pesticides use, decline in insect prey, and habitat destruction were suspected causes (Bontadina et al., 2000). This stressed the need for a better understanding of its habitat requirement, leading to numerous telemetry studies (e.g Bontadina et al., 2002; Downs et al., 2016; Reiter et al., 2013; Schofield, 1996) and, more recently, to larger scale presence/absence studies (Le Roux et al., 2017; Tournant et al., 2013). These studies mainly focused on habitat selection by females during the crucial period of parturition, during which they gather in maternity colonies. Individual-level studies revealed that the lesser horseshoe bat

forages in woodland, especially broadleaved ones, and spend most of their foraging time in close vicinity to their roost (within a radius of 500-600 meters - Bontadina et al., 2002; Reiter et al., 2013). The importance of woodland proportion on the presence and abundance of *R. hipposideros* was confirmed by several studies (e.g. Reiter, 2004; Tournant et al., 2013), and presence data revealed a negative effect of artificial land cover on the presence of maternity colonies (Tournant et al., 2013). Habitat requirements of this species may dramatically constrain the opportunity offered by a changing climate to its predicted northward expansion (Rebelo et al., 2010).

In this study, we tested the hypothesis that important environmental variables known from individual radio-tracking studies or presence surveys also influence the demography of the species. We used count data obtained from a fifteen years monitoring scheme at 94 colonies situated in Brittany (France), to investigate the population-habitat relationship in *R. hipposideros*. We used these data to infer variables that are related to the population dynamics of the colonies before testing their relationship with environmental variables.

2. Material and Methods

2.1. Demographic data for *R. hipposideros* colonies

From 2000 to 2014, 94 *R. hipposideros* maternity colonies were monitored in Brittany, France (Fig. 1, see details in Jan et al., 2017). The following demographic variables were compiled for each colony: mean colony size (the number of adults averaged over the years), fecundity (the juveniles/adults ratio averaged over the years), and growth rate. The last variable was estimated with the density-independent Ricker model using the package "PVAclone" (Nadeem and Solymos, 2016). Five clones in the data-cloning algorithm and 20 000 iterations were performed. The density independent Ricker model is a parsimonious model that considers that the observed colony size variation through time is only the result of growth rate and observation error, and allows to estimate the growth with a known precision. The reasons behind the choice of the model are detailed in Supp. Inf. A.

2.2. Landscape attributes

We built a geographic information system (GIS) of land cover associated with other descriptors of the landscape surrounding the colonies. To obtain information precise enough and relevant for *R. hipposideros,* the land cover dataset was built from different databases: CORINE Land Cover - European Commission 2006, BD Topo 2.1 - IGN 2013, BD Forest v1 - IGN 2012 and Graphic Parcel Register 1.0 (RPG) - IGN 2012. The land cover map construction included procedures for correcting topological errors, homogenizing information resolution and harmonizing attribute data. Finally, we obtained a map featuring 38 classes of land use (Supp. Inf. B) with a minimum mapping unit (MMU) of 100 m².

Figure 1: Map of the 94 colonies monitored in Brittany, France.

2.3. Assessment of the effect of landscape variables on bat population dynamics

We considered the landscape within a circular buffer surrounding each colony. Two buffer radius were tested: 2500 m, reflecting the radius commonly used in recent studies on the lesser horseshoe bat (Afonso et al., 2016), and 500 m, reflecting the core area of the foraging zone (Bontadina et al., 2002; Reiter et al., 2013). Before clipping the land cover layer with each buffer, the 38 cover types were aggregated in six land-cover classes (Supp. Inf. B) according to previous knowledge gathered at the individual level on lesser horseshoe bat's ecology (Bontadina et al., 2002; Tournant et al., 2013), namely: broadleaved woodland, coniferous woodland, artificial area, water bodies, cropland, and open land (other than crops and artificial). For each buffer we calculated six FRAGSTATS metrics (McGarigal and Marks, 1995) describing the composition and the configuration of the landscape pattern: the proportion of the landscape occupied by each class (%LAND), the mean SHAPE index (MSI) of each class, the Simpson's diversity index (SIDI), the Shannon's diversity index (SHDI), the Shannon's evenness index (SHEI) and the mean patch size (MPS). Since a MSI value of 1 corresponds to a circular patch, and because this value increases with patch shape irregularity, a high MSI value reflects an increase in edge density of a particular land cover, for an equivalent area.

Statistical model averaging was then performed to estimate the effect of land cover variables on population dynamic parameters. Full models considered either the colony size (Generalized linear model with negative-binomial distribution - Jan et al. 2017), fecundity (linear model) or the growth rate (weighted least squares regressions) as the response variable, and the proportion and shape of the six land cover, the mean patch size and land cover diversity as explanatory variables (without interaction). Explanatory variables were centred and scaled before model computation. Weighted least square regressions were performed to take into account the uncertainty around growth rate estimates by weighting each data point by the inverse of the standard deviation given by the estimation (Ryan, 2008).

These models were then used in the glmulti R package to obtain every possible combination of explanatory variables (without interaction) and order them by AICc criterion (Calcagno and de Mazancourt, 2010). We then performed model averaging by calculating the Akaike weight of each model within 2ΔAICc of the best model (Burnham and Anderson, 2003). The model-averaged regression coefficients of the predictors and their 85% confidence intervals (CI) were calculated based on the cumulative weights of the models including the variable (Calcagno and de Mazancourt, 2010). 85% confidence intervals are more relevant than 95% interval confidence for AIC-based model averaging (Arnold, 2010), and explanatory variables were then considered as having a meaningful positive or negative impact on the response variable if their 85% confidence interval did not include zero. Because growth rates and their associated standard deviations obtained through PVAClone can slightly vary from one run to another, we decided to run ten times the growth rates estimation and the model averaging associated with this response variable. Only the explanatory variables that were significant ten times out of ten were considered as significant in our results.

Land cover areas were considered through their proportions, inevitably leading to correlation between variables. We prevented this by discarding models including more than four proportion variables for model averaging. Correlations between coefficients were thus lower than 0.8 for every remaining models. The three diversity indexes (SIDI, SHDI and SHEI) were also highly correlated, and we selected the index which explained most variability for each demographic variable by building simple linear models with every response variable prior to model averaging.

Since we previously showed that some climatic variables impact colony size and fecundity (Jan et al., 2017), we averaged these variables (June and October precipitation and May and November mean temperature for Colony size; April and October precipitation and April and July minimum temperature for fecundity) along the sampling period for each colony and included them in all models using colony size or fecundity as a response variable to control for the variability explained by the weather. The description of the complete models which were the basis of model averaging are presented in Supp. Inf. C. Our data set provides population dynamic parameters on a large spatio-temporal scale, and a sample size large enough to investigate numerous environmental variables. Complete models included the whole 94 colony dataset to test the significance of no more than 14 environmental variables (climatic variables being present in the model but not tested), ensuring than the ratio sample size (N) / tested predictor (k) was high enough to avoid overparametrisation (N/k = 6.71, Forstmeier and Schielzeth, 2011).

To determine which radius (500 or 2500m) best explained variations in our response variables, we compared the AICc and R² of the models explaining the population dynamics parameters with the significant predictors detected in the previous model averaging.

3. Results

Shannon's evenness index (SHEI) was the diversity index explaining most variability for the three demographic variables, and was thus the only diversity index included in model averaging. Significant landscape variables were detected for each response variable when considering a 500 m radius buffer around the colony: four for colony size, one for fecundity and six for the growth rate (Tab. 1). The proportion of artificial land negatively impacted the three investigated response variables. The growth rate was also negatively impacted by the proportion of crop areas, while colony size was positively influenced by the proportion of broadleaved woodland. Mean Shape Indices (MSI) associated with water bodies were positively related to colony size and growth rates, which were also positively influenced by the MSI of crop and artificial lands. Finally, MSI of open land had a negative impact on colony size and growth rates.

Only six significant predictors were detected within the 2500 m radius buffer, two of them being concordant with the results of the 500 m radius buffer (Tab. 1). When comparing the AICc and R² of the model given by the two buffer sizes, fixed effect detected with the 500m buffer always performed better (Tab. 2). Models constructed with significant land cover predictors explained 42% of the variance for colony size, and 8% for fecundity. These values reached 53% and 13% when significant climatic predictors previously detected for colony size and fecundity were added to the corresponding models (Jan et al., 2017). The six significant predictors found in growth rate models explained 80% ($\pm$3%) of this variable, with crop land proportion being the predictor with the largest effect size and explaining 53% ($\pm$5%) of the variation.

Table 1: Results of a model averaging of population dynamics parameters as a function of the proportion of land (%LAND) and Mean Shape Index (MSI) of different cover type, the mean patch size and SHEI (Shannon's evenness index). NS: non-significant predictor; +: positive significant predictor; - : negative significant predictor. When results differ between the two buffer sizes, results given by the 2500m buffer are indicated between brackets. Cells with significant results with the 500m buffer are shaded.

	Broadleaved woodland		Coniferous woodland		Artificial		Crop land		Open land		Water bodies		SHEI	Mean patch size
	%LAND	MSI	%LAND	MSI	%LAND	MSI	%LAND	MSI	%LAND	MSI	%LAND	MSI		
Colony size	+	NS	NS	NS	- (NS)	NS	NS	NS	NS	- (NS)	NS	+ (NS)	NS	NS
Fecundity	NS	NS	NS	NS	- (NS)	NS	NS	NS	NS	NS	NS	NS	NS	NS
Growth rate	NS	NS(+)	NS	NS	- (NS)	+ (NS)	- (NS)	+	NS	- (NS)	NS (+)	+ (-)	NS	NS (+)

Table 2: Residual degree of freedom (df), AICc and R² of models that included land cover significant predictors for the two buffer scales. For colony size and fecundity, R² of models that include significant climatic predictor detected in a previous study (Jan et al. 2017) are also presented. AICc and R² of growth rate model are averaged over the ten runs we performed for growth rate calculation, and resented with the corresponding standard deviation

Response variable	Radius	Fixed effects	df	AICc	R²	R² with climatic variables
Colony size	500 m.	Broadleaved woodland and artificial proportion Open land and water land MSI	89	775.9	0.42	0.53
	2500 m.	Broadleaved woodland proportion	92	796.5	0.07	0.23
Fecundity	500 m.	Artificial proportion	92	-116.7	0.08	0.13
	2500 m.	-	-	-	-	-
Growth rate	500 m.	Artificial and crop land proportion Artificial, crop land, water bodies and open land MSI	87	272.0 (±3.8)	0.80 (±0.03)	-
	2500 m.	Water bodies proportion Broadleaved woodland, crop land water bodies MSI Mean patch size	88	344.2 (±3.8)	0.55 (±0.05)	-

4. Discussion

The current accelerating rate of biodiversity erosion combined with the threat posed by already ongoing global changes have prompt the development of anticipatory predictive models to forecast the future of biodiversity (e.g. species distribution).

Contrarily to explanatory predictions, that aim to corroborate hypotheses on the mechanisms underpinning observed changes using an hypothetico-deductive reasoning, anticipatory predictions are decoupled from the explanatory process: they usually assume that the theory or the putative link between considered causes and effects is true and do not aim at testing models and theory (Mouquet et al., 2015).

Considering the increasing availability in the literature of information on species occurrence and habitat selection or use (e.g. from individual telemetry or observation studies) that can be used to perform anticipatory predictions, it is important to empirically test whether individual- and population-level responses to habitat quality are convergent to evaluate the predictive power of the assumed link between habitat selection and population dynamics.

In this study, we assessed the population dynamic consequences of habitat selection in the lesser horseshoe bat based on environmental variables selected from previous individual level studies. Because we used demographic data censused over multiple years and sites, we were able to correlate both land cover area and shape on all the investigated bat demography parameters (i.e. colony size, fecundity, growth rate). Following paragraphs describe how these results fit and refine previous conclusions of individual level studies.

4.1. Scale of habitat dependency and management implication

We showed that investigating the effect of landscape composition and configuration on the dynamics of *R. hipposideros* maternity colonies is more relevant at the smallest of the two investigated scales. Indeed, the AICc and R² values of models including landscape significant predictors always supported the importance of the landscape surrounding colonies within a 500 m radius (lower AICc and larger part of explained variance R² - Tab. 2). This result is consistent with those of individual level telemetry studies demonstrating that individuals forage for at least half of their time within a 500-600 m radius around maternity roosts, although few individuals can eventually forage at distances up to 5 km around colonies (Bontadina et al., 2002; Reiter et al., 2013). Nevertheless, further investigations are required to precise the most relevant spatial scale at which habitat (affecting colony sizes and dynamics) should be protected or managed for an optimal and cost effective conservation of this species, while taking into account connectivity between populations.

4.2. Identification of key habitats for *R. hipposideros* population dynamics and conservation

Colony size was positively influenced by broadleaved woodland proportion at both spatial scales. This validates and refines previous studies, in which woodland areas was one of the best landscape predictors of maternity roost presence (Bontadina et al., 2002; Reiter, 2004; Schofield, 1996; Tournant et al., 2013). Indeed, broadleaf woodlands, often offering a higher richness and abundance of insects than coniferous plantations (Benton et al., 2002; Goiti et al.,

2004), are more intensively used as foraging areas by *R. hipposideros* than any other habitat (Bontadina et al., 2002; Schofield, 1996).

We also found a significant positive effect of water bodies shape complexity on colony size and growth rate. This result reveals that long lakeshores or riverbanks near maternity colonies favour colony size and the long-term establishment of this species. Shores with sufficient riparian vegetation to provide food and shelter to a great density of insect is known to drive the distribution and foraging behaviour of insectivorous bats (Warren et al., 2000), as observed in *R. hipposideros* (Bontadina et al., 2002; Holzhaider et al., 2002; Reiter, 2004). These habitats must represent high quality foraging areas for this bat since a non-negligible part of its main prey (e.g. Trichoptera, Nematocera, Lepidoptera – Arlettaz et al., 2000; Lino et al., 2014) have an aquatic larval development.

The preservation of large broadleaved woodland and high density of lakeshores and riverbanks in the direct vicinity of maternity colonies appears to be the priority for conserving viable colonies of this non-migratory forest species. These high quality cover types, in terms of resources and protection, are suspected to positively influence bat population dynamics by enhancing individual survival. This knowledge would be highly beneficial to objectively guide bat conservation, for instance by identifying where to favour the settlement of new colonies through roost construction (e.g. Mering and Chambers, 2014) and/or by prioritising the management or renovation of roosting sites (buildings, abandoned mines, caves - e.g. Lacki, 2000).

4.3. Landscape variables threatening *R. hipposideros* population dynamics and conservation

Our study highlights that fragmentation and degradation of habitat are a major concern for *R. hipposideros* dynamics.

Indeed, our results revealed a significant and negative effect of open land MSI on population size and growth rate. This means that the complexity of the spatial configuration of open land patches (e.g. patch number, shape) surrounding maternity colonies, as a measurement of habitat fragmentation (Wang et al., 2014), affects the dynamics and persistence of this bat. This may reveal a barrier effect of this generally avoided cover type (Bontadina et al., 2002; Holzhaider et al., 2002; Reiter, 2004; Tournant et al., 2013), possibly by hindering individual bat foraging movements, especially if forest corridors and hedge rows are scarce (Downs and Racey, 2006; Fischer et al., 2010; Lumsden and Bennett, 2005). Indeed, most insectivorous bats are known to use scattered trees or hedgerows as transit and foraging corridors because these natural vertical structures offer shelter from predators and generally present higher insect densities than open areas (Downs and Racey, 2006; Fischer et al., 2010; Lumsden and Bennett, 2005). This seems to be supported by the apparent inability of this species to adapt its spatial foraging behaviour in degraded landscapes, and by the high energetic cost of flight that probably constrains its travelling distance, especially for pregnant or lactating females that mostly compose maternity colonies (Rainho and Palmeirim, 2011; Reiter et al., 2013).

Impact of habitat quality degradation could be assessed with the effects of artificial and crop lands. Proportions of both cover types have negative impact on *R. hipposideros* dynamics. In addition, their complexity indexes have positive effect which suggests a negative impact of

large and regular urban or crop patches around the colonies probably by decreasing the edge density and the interactions with other more favourable types of landscape cover. The negative effect of urbanization on colony sizes, fecundity and growth rates, is in accordance with a study based on occupancy data only (Tournant et al., 2013). It may be due to the low quality of this cover type (scarcity of favourable foraging areas, low quality or quantity of food resource, higher exposition to light pollution, accidental mortality, predation, disturbance, stress or pollution – see Fensome and Mathews, 2016; Gaston et al., 2013; O'shea et al., 2016; Stone et al., 2009; Ziembicki et al., 2015).

The major and negative effect of crop land proportion on maternity colony growth rates that we detected is probably due to the unfavourable nature of this habitat for *R. hipposideros* (Bontadina et al., 2002). Direct and indirect effects of agriculture intensification have been proposed as one of the main cause implicated in the decline of various organisms (Bayat et al., 2014; Gibbons et al., 2014; Hallmann et al., 2014) including *R. hipposideros* (Bontadina et al., 2000; Wickramasinghe et al., 2004). Pesticides can have various additive or synergistic deleterious effects on survival and health of long-lived insectivorous bats (Hsiao et al., 2016). Their use in conventional farmlands reduces insect abundance and diversity (especially three key families for many insectivorous bats: lepidopteran, dipteran, coleopteran) compared to organic farmlands, the latter being more frequently used as foraging sites by *R. hipposideros* (Wickramasinghe et al., 2004). Agrochemicals also impact insectivorous bats by direct poisoning through dermal contact, inhalation or ingestion in their diet, and because of the accumulation and biomagnification of these pollutants in bat organism, and their remobilization from fat tissues during hibernation or migration (Bayat et al., 2014; Mineau and Callaghan, 2018). During lactation, juveniles strongly accumulate organochlorine pesticides (Lüftl et al., 2005; Streit et al., 1995) through the feeding from their mother, and their survival is more affected by pesticide contamination than adult survival (Frick et al., 2007). Moreover, when foraging, inexperienced juveniles fly at much greater distances and are less selective on habitat quality than reproductive females (Bontadina et al., 2002). They are therefore potentially more exposed to factors affecting their survival (e.g. low prey availability, agrochemicals, predation) in such unfavourable habitats which can in turn negatively affect population growth rates.

4.4. Consistency between individual and population level effect of habitat quality

Overall, our results support the dependence of population dynamics to patterns of habitat selection by individuals with a positive relationship between habitat quality and bat abundance/density : habitats predicted as high quality by individual-level studies (i.e. broadleaved woodlands and long lakeshores or riverbanks) were associated to large colony sizes, contrarily to those predicted as low quality habitats (i.e. artificial and open lands) associated with low colony sizes, but also with a low fecundity and/or growth rate. Such relationships between elements of landscapes that favour or limit the dynamics of maternity colonies are key information both for the long-term conservation of this non-migratory species and to refine range shift predictions under climate change.

Nevertheless, other studies showed that habitat selection can vary depending on the spatial and temporal scales, both between and within species (e.g. according to the sex and/or life stage), and affect individual fitness and population dynamics through different demographic

processes (e.g. sex or stage specific survival rates, fecundity - Bjørneraas et al., 2012; Dyson et al., 2018; Jacques et al., 2015; King et al., 2006; Low et al., 2010; Street et al., 2016). Considering the variety of the possible mechanisms, mediated by individual fitness, driving population-level response to habitat quality, we strongly encourage further cross-validation studies aiming at testing these relationships, for example using a habitats-to-populations (HTP) framework (Matthiopoulos et al., 2019).

4.5. Conclusion

We underlined the importance of broadleaved woodland proportion and lakeshores and riverbanks density in *R. hipposideros* core foraging area (500 m radius around maternity colonies), probably because their associated vegetation provides particularly favourable shelter, movement and foraging corridors. We also pinpointed the threat posed by the intensification of human activities (i.e. urbanization and agriculture intensification inducing habitat fragmentation), resulting in the spread of unfavourable and degraded habitats (i.e. with low availability of their prey, high exposure to predation and pollution), on the long-term persistence of large *R. hipposideros* populations.

The links we found between landscape variables and the demography of this non-migratory forest species will help refining range shift predictions based on climatic variables only (Rebelo et al., 2010). Concomitantly, considering the connectivity of favorable habitats and the threats posed by the intensification of human activities (urbanization, agriculture, pollution), for example under different land use change scenarii, will allow to more accurately predict the fate of the species towards the end of the century.

<u>Acknowledgements</u>

We are very grateful to all volunteers from Bretagne Vivante and Le Groupe Mammalogique Breton for providing visual count data of maternity colonies in Brittany that allowed us to perform this study as well as the companion study published in Oecologia (Jan et al. 2017). We thank Benoît Persyn and Météo-France for providing us with meteorological data through the research convention Météo-France/INRA. This work benefited from funding by the French National Forest Office (ONF).

<u>References</u>

Acevedo, P., Ferreres, J., Escudero, M.A., Jimenez, J., Boadella, M., Marco, J., 2017. Population dynamics affect the capacity of species distribution models to predict species abundance on a local scale. Divers. Distrib. 23, 1008–1017.

Afonso, E., Tournant, P., Foltête, J.-C., Giraudoux, P., Baurand, P.-E., Roué, S., Canella, V., Vey, D., Scheifler, R., 2016. Is the lesser horseshoe bat (*Rhinolophus hipposideros*) exposed to causes that may have contributed to its decline? A non-invasive approach. Glob. Ecol. Conserv. 8, 123–137.

Aitken, S.N., Yeaman, S., Holliday, J.A., Wang, T., Curtis-McLane, S., 2008. Adaptation, migration or extirpation: climate change outcomes for tree populations. Evol. Appl. 1, 95–111.

Arlettaz, R., Godat, S., Meyer, H., 2000. Competition for food by expanding pipistrelle bat populations (*Pipistrellus pipistrellus*) might contribute to the decline of lesser horseshoe bats (*Rhinolophus hipposideros*). Biol. Conserv. 93, 55–60.

Arnold, T.W., 2010. Uninformative parameters and model selection using Akaike's Information Criterion. J. Wildl. Manag. 74, 1175–1178.

Auffret, A.G., Rico, Y., Bullock, J.M., Hooftman, D.A., Pakeman, R.J., Soons, M.B., Suárez-Esteban, A., Traveset, A., Wagner, H.H., Cousins, S.A., 2017. Plant functional connectivity–integrating landscape structure and effective dispersal. J. Ecol. 105, 1648–1656.

Bayat, S., Geiser, F., Kristiansen, P., Wilson, S.C., 2014. Organic contaminants in bats: trends and new issues. Environ. Int. 63, 40-52.

Benton, T.G., Bryant, D.M., Cole, L., Crick, H.Q., 2002. Linking agricultural practice to insect and bird populations: a historical study over three decades. J. Appl. Ecol. 39, 673–687.

Bjørneraas, K., Herfindal, I., Solberg, E.J., Sæther, B.E., van Moorter, B., Rolandsen, C.M., 2012. Habitat quality influences population distribution, individual space use and functional responses in habitat selection by a large herbivore. Oecologia 168, 231-243.

Bontadina, F., Arlettaz, R., Fankhauser, T., Lutz, M., Mühlethaler, E., Theiler, A., Zingg, P., 2000. The lesser horseshoe bat *Rhinolophus hipposideros* in Switzerland: present status and research recommendations. Le Rhinolophe 14, 69–83.

Bontadina, F., Schofield, H., Naef-Daenzer, B., 2002. Radio-tracking reveals that lesser horseshoe bats (*Rhinolophus hipposideros*) forage in woodland. J. Zool. 258, 281–290.

Boyce, M.S., Johnson, C.J., Merrill, E.H., Nielsen, S.E., Solberg, E.J., Van Moorter, B., 2016. Can habitat selection predict abundance? J. Anim. Ecol. 85, 11–20.

Burnham, K.P., Anderson, D.R., 2003. Model selection and multimodel inference: a practical information-theoretic approach. Springer Science & Business Media.

Calcagno, V., de Mazancourt, C., 2010. glmulti: an R package for easy automated model selection with (generalized) linear models. J. Stat. Softw. 34, 1–29.

Downs, N.C., Cresswell, W.J., Reason, P., Sutton, G., Wells, D., Wray, S., 2016. Sex-specific habitat preferences of foraging and commuting lesser horseshoe bats *Rhinolophus hipposideros* (Borkhausen, 1797) in lowland England. Acta Chiropt. 18, 451–465.

Downs, N.C., Racey, P.A., 2006. The use by bats of habitat features in mixed farmland in Scotland. Acta Chiropterologica 8, 169–185.

Dyson, M.E., Schummer, M.L., Barney, T.S., Fedy, B.C., Henry, H.A., Petrie, S.A., 2018. Survival and habitat selection of wood duck ducklings. J. Wildl. Manag. 82, 1725-1735.

Elith, J., Leathwick, J.R., 2009. Species distribution models: ecological explanation and prediction across space and time. Annu. Rev. Ecol. Evol. Syst. 40, 677–697.

Fensome, A.G., Mathews, F., 2016. Roads and bats: a meta-analysis and review of the evidence on vehicle collisions and barrier effects. Mammal Rev. 46, 311–323.

Fischer, J., Stott, J., Law, B.S., 2010. The disproportionate value of scattered trees. Biol. Conserv. 143, 1564–1567.

Forstmeier, W., Schielzeth, H., 2011. Cryptic multiple hypotheses testing in linear models: overestimated effect sizes and the winner's curse. Behav. Ecol. Sociobiol. 65, 47–55.

Frick, W.F., Rainey, W.E., Pierson, E.D., 2007. Potential effects of environmental contamination on *Yuma myotis* demography and population growth. Ecol. Appl. 17, 1213–1222.

Gaillard, J.-M., Hebblewhite, M., Loison, A., Fuller, M., Powell, R., Basille, M., Van Moorter, B., 2010. Habitat–performance relationships: finding the right metric at a given spatial scale. Philos. Trans. R. Soc. Lond. B Biol. Sci. 365, 2255–2265.

Gaston, K.J., Bennie, J., Davies, T.W., Hopkins, J., 2013. The ecological impacts of nighttime light pollution: a mechanistic appraisal. Biol. Rev. 88, 912–927.

Gibbons, D., Morrissey, C., Mineau, P., 2014. A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate wildlife. Environ. Sci. Pollut. Res. 22, 103-118.

Goiti, U., Aihartza, J.R., Garin, I., 2004. Diet and prey selection in the Mediterranean horseshoe bat *Rhinolophus euryale* (*Chiroptera,*

Rhinolophidae) during the pre-breeding season. Mammalia 68, 397–402.

Gurevitch, J., Fox, G.A., Fowler, N.L., Graham, C.H., 2016. Landscape Demography: Population Change and its Drivers Across Spatial Scales. Q. Rev. Biol. 91, 459–485.

Hallmann, C.A., Foppen, R.P.B., van Turnhout, C.A.M., de Kroon, H., Jongejans, E., 2014. Declines in insectivorous birds are associated with high neonicotinoid concentrations. Nature 511, 341-343.

Hodgson, J.A., Thomas, C.D., Wintle, B.A., Moilanen, A., 2009. Climate change, connectivity and conservation decision making: back to basics. J. Appl. Ecol. 46, 964–969.

Holzhaider, J., Kriner, E., Bernd-Ulric, R., Zahn, A., 2002. Radio-tracking a Lesser horseshoe bat (*Rhinolophus hipposideros*) in Bavaria: an experiment to locate roosts and foraging sites. Myotis 40, 47–54.

Hsiao, C.-J., Lin, C.-L., Lin, T.-Y., Wang, S.-E., Wu, C.-H., 2016. Imidacloprid toxicity impairs spatial memory of echolocation bats through neural apoptosis in hippocampal CA1 and medial entorhinal cortex areas. Neuroreport 27, 462–468.

Jacques, C.N., Jenks, J.A., Grovenburg, T.W., Klaver, R.W., 2015. Influence of habitat and intrinsic characteristics on survival of neonatal pronghorn. PloS One 10, e0144026.

Jan, P.-L., Farcy, O., Boireau, J., Le Texier, E., Baudoin, A., Le Gouar, P., Puechmaille, S.J., Petit, E.J., 2017. Which temporal resolution to consider when investigating the impact of climatic data on population dynamics? The case of the lesser horseshoe bat (*Rhinolophus hipposideros*). Oecologia 184, 749–761.

King, D.I., Degraaf, R.M., Smith, M.L., Buonaccorsi, J.P., 2006. Habitat selection and habitat-specific survival of fledgling ovenbirds (*Seiurus aurocapilla*). J. Zool. 269, 414-421.

Kokko, H., López-Sepulcre, A., 2006. From individual dispersal to species ranges: perspectives for a changing world. Science 313, 789–791.

Lacki, M.J., 2000. Effect of trail users at a maternity roost of Rafinesque's big-eared bats. J. Cave Karst Stud. 62, 163–168.

Le Roux, M., Redon, M., Archaux, F., Long, J., Vincent, S., Luque, S., 2017. Conservation planning with spatially explicit models: a case for horseshoe bats in complex mountain landscapes. Landsc. Ecol. 32, 1005–1021.

Lino, A., Fonseca, C., Goiti, U., Pereira, M.J.R., 2014. Prey selection by *Rhinolophus hipposideros* (*Chiroptera, Rhinolophidae*) in a modified forest in Southwest Europe. Acta Chiropt. 16, 75–83.

Low, M., Arlt, D., Eggers, S., Pärt, T., 2010. Habitat-specific differences in adult survival rates and its links to parental workload and on-nest predation. J. Anim. Ecol. 79, 214-224.

Lüftl, S., Freitag, B., Deutz, A., Steineck, T., Tataruch, F., 2005. Concentrations of organochlorine pesticides and PCBs in the liver of European bats (microchiroptera). Fresenius Environ. Bull. 14, 167–172.

Lumsden, L.F., Bennett, A.F., 2005. Scattered trees in rural landscapes: foraging habitat for insectivorous bats in south-eastern Australia. Biol. Conserv. 122, 205–222.

Manly, B.F.L., McDonald, L., Thomas, D.L., McDonald, T.L., Erickson, W.P., 2007. Resource selection by animals: statistical design and analysis for field studies. Springer Science & Business Media.

Matthiopoulos, J., Field, C., MacLeod, R., 2019. Predicting population change from models based on habitat availability and utilization. Proc. R. Soc. B 286, 20182911.

McGarigal, K., Marks, B.J., 1995. FRAGSTATS: spatial pattern analysis program for quantifying landscape structure. Gen Tech Rep PNW-GTR-351 Portland US Dep. Agric. For. Serv. Pac. Northwest Res. Stn. 122 P 351.

Mering, E.D., Chambers, C.L., 2014. Thinking outside the box: a review of artificial roosts for bats. Wildl. Soc. Bull. 38, 741–751.

Mineau, P., Callaghan, C., 2018. Neonicotinoid insecticides and bats: an assessment of the direct and indirect risks. Canadian Wildlife Federation.

Mouquet, N., Lagadeuc, Y., Devictor, V., Doyen, L., Duputié, A., Eveillard, D., ... , Jabot, F., 2015.

Predictive ecology in a changing world. J. Appl. Ecol. 52, 1293-1310.

Nadeem, K., Solymos, P., 2016. PVAClone: population viability analysis with data cloning. R package version 0.1-6. https://CRAN.R-project.org/package=PVAClone

O'shea, T.J., Cryan, P.M., Hayman, D.T., Plowright, R.K., Streicker, D.G., 2016. Multiple mortality events in bats: a global review. Mammal Rev. 46, 175–190.

Pulliam, H.R., 2000. On the relationship between niche and distribution. Ecol. Lett. 3, 349–361.

Rainho, A., Palmeirim, J.M., 2011. The importance of distance to resources in the spatial modelling of bat foraging habitat. PLoS One 6, e19227.

Rebelo, H., Tarroso, P., Jones, G., 2010. Predicted impact of climate change on European bats in relation to their biogeographic patterns. Glob. Change Biol. 16, 561–576.

Reiter, G., 2004. The importance of woodland for *Rhinolophus hipposideros* (*Chiroptera, Rhinolophidae*) in Austria. Mammalia 68, 403–410.

Reiter, G., Pölzer, E., Mixanig, H., Bontadina, F., Hüttmeir, U., 2013. Impact of landscape fragmentation on a specialised woodland bat, *Rhinolophus hipposideros*. Mamm. Biol.-Z. Für Säugetierkd. 78, 283–289.

Ryan, T.P., 2008. Modern regression methods. John Wiley & Sons.

Schofield, H.W., 1996. The ecology and conservation biology of *Rhinolophus hipposideros*, the lesser horseshoe bat (PhD Thesis). University of Aberdeen.

Stone, E.L., Jones, G., Harris, S., 2009. Street lighting disturbs commuting bats. Curr. Biol. 19, 1123–1127.

Street, G.M., Fieberg, J., Rodgers, A.R., Carstensen, M., Moen, R., Moore, S.A., Windels, S.K., Forester, J.D., 2016. Habitat functional response mitigates reduced foraging opportunity: implications for animal fitness and space use. Landsc. Ecol. 31, 1939–1953.

Streit, B., Winter, S., Nagel, A., 1995. Bioaccumulation of selected organochlorines in bats and tits: Influence of chemistry and biology. Environ. Sci. Pollut. Res. 2, 194–199.

Thuiller, W., Münkemüller, T., Lavergne, S., Mouillot, D., Mouquet, N., Schiffers, K., Gravel, D., 2013. A road map for integrating eco-evolutionary processes into biodiversity models. Ecol. Lett. 16, 94–105.

Thuiller, W., Münkemüller, T., Schiffers, K.H., Georges, D., Dullinger, S., Eckhart, V.M., Edwards, T.C., Gravel, D., Kunstler, G., Merow, C., Moore, K., Piedallu, C., Vissault, S., Zimmermann, N.E., Zurell, D., Schurr, F.M., 2014. Does probability of occurrence relate to population dynamics? Ecography 37, 1155–1166.

Tournant, P., Afonso, E., Roué, S., Giraudoux, P., Foltête, J.-C., 2013. Evaluating the effect of habitat connectivity on the distribution of lesser horseshoe bat maternity roosts using landscape graphs. Biol. Conserv. 164, 39–49.

Wang, X., Blanchet, F.G., Koper, N., 2014. Measuring habitat fragmentation: an evaluation of landscape pattern metrics. Methods Ecol. Evol. 5, 634–646.

Warren, R.D., Waters, D.A., Altringham, J.D., Bullock, D.J., 2000. The distribution of Daubenton's bats (*Myotis daubentonii*) and pipistrelle bats (*Pipistrellus pipistrellus*)(Vespertilionidae) in relation to small-scale variation in riverine habitat. Biol. Conserv. 92, 85–91.

Wickramasinghe, L.P., Harris, S., Jones, G., Vaughan Jennings, N., 2004. Abundance and species richness of nocturnal insects on organic and conventional farms: effects of agricultural intensification on bat foraging. Conserv. Biol. 18, 1283–1292.

Ziembicki, M.R., Woinarski, J.C., Webb, J.K., Vanderduys, E., Tuft, K., Smith, J., Ritchie, E.G., Reardon, T.B., Radford, I.J., Preece, N., 2015. Stemming the tide: progress towards resolving the causes of decline and implementing management responses for the disappearing mammal fauna of northern Australia. Therya 6, 169–225.

Data accessibility

R scripts used in this study were deposited at Figshare through the DOI 10.6084/m9.figshare.8337437

Supplementary Information:

Supplementary Information A: Selection of population growth model.
Supplementary Information B: Aggregation of 38 cover types in six-landscape class.
Supplementary Information C: Description of the complete models used for model averaging.