Electrophysiological network alterations in adults with copy number variants associated with high neurodevelopmental risk

Diana C. Dima[1,2*], Rachael Adams[1,3], Stefanie C. Linden[2,3,4], Alister Baird[3], Jacqueline Smith[3,4], Sonya Foley[1], Gavin Perry[1], Bethany C. Routley[1], Lorenzo Magazzini[1], Mark Drakesmith[1,2], Nigel Williams[2,3,4], Joanne Doherty[1,3,4], Marianne B.M. van den Bree[2,3,4], Michael J. Owen[2,3,4], Jeremy Hall[2,3,4], David E. J. Linden[1,2,3,4,5], and Krish D. Singh[1]

Author affiliations:

[1] Cardiff University Brain Research Imaging Centre (CUBRIC), School of Psychology, Cardiff University, Maindy Road, Cardiff, CF24 4HQ, United Kingdom

[2] Neuroscience and Mental Health Research Institute (NMHRI), Cardiff University, Maindy Road, Cardiff, CF24 4HQ, United Kingdom

[3] Division of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, CF24 4HQ, UK

[4] MRC Centre for Neuropsychiatric Genetics and Genomics, School of Medicine, Cardiff University, Maindy Road, Cardiff, CF24 4HQ, United Kingdom

[5] School of Mental Health and Neuroscience, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands

*Corresponding author: DimaDC@cardiff.ac.uk

Abstract

Rare copy number variants associated with increased risk for neurodevelopmental and psychiatric disorders (referred to as ND-CNVs) are characterized by heterogeneous phenotypes thought to share a considerable degree of overlap. Altered neural integration has often been linked to psychopathology and is a candidate marker for potential convergent mechanisms through which ND-CNVs modify risk; however, the rarity of ND-CNVs means that few studies have assessed their neural correlates. Here, we used magnetoencephalography (MEG) to investigate resting-state oscillatory connectivity in a cohort of 42 adults with ND-CNVs, including deletions or duplications at 22q11.2, 15q11.2, 15q13.3, 16p11.2, 17q12, 1q21.1, 3q29, and 2p16.3, and 42 controls. We observed decreased connectivity between occipital, temporal and parietal areas in participants with ND-CNVs. This pattern was common across genotypes and not exclusively characteristic of 22q11.2 deletions, which were present in a third of our cohort. Furthermore, a data-driven graph theory framework enabled us to successfully distinguish participants with ND-CNVs from unaffected controls using differences in node centrality and network segregation. Together, our results point to alterations in electrophysiological connectivity as a putative common mechanism through which genetic factors confer increased risk for neurodevelopmental and psychiatric disorders.

1 Introduction

A number of rare genetic variants occurring through the deletion or duplication of
chromosomal segments are associated with significantly increased risk for a range of
neurodevelopmental disorders (ND), including schizophrenia, autism spectrum disorder
(ASD), and developmental delay (Kirov et al., 2014). Although the underlying mechanisms
remain poorly understood, these copy number variants (referred to hereafter as ND-CNVs) are
thought to increase the risk for psychopathology through alterations in neural structure and
function. Thus, neuroimaging studies in participants with ND-CNVs provide a unique
opportunity to study intermediate phenotypes of mental disorders.

Recent work suggests that CNV-specific phenotypic outcomes are limited, pointing instead to
a large degree of similarity across phenotypes associated with different ND-CNVs (Chawner
et al., 2019; Niarchou et al., 2019). Focusing on convergent neural alterations across different
genotypes can thus help elucidate the mechanisms linking ND-CNVs at different loci to a
shared psychopathology and increase in neurodevelopmental risk.

Failures of functional neural integration are considered a hallmark of schizophrenia (Friston et
al., 2016; Dong et al., 2018) and have also been found in ASD (Hull et al., 2017; O'Reilly et
al., 2017) and across diagnostic boundaries (Sha et al., 2019). ND-CNVs are thought to increase
disorder risk by acting on large-scale neural integration through molecular and cellular
mechanisms (Karayiorgou et al., 2010). Synchronous oscillatory activity thought to support
communication between brain areas is thus of particular interest as a potential biomarker of
neurodevelopmental risk, and can be measured at rest using electro- and magneto-
encephalography (EEG/MEG).

However, the rarity of ND-CNVs means that evidence of their functional connectivity
correlates is scarce. Of the genetic imaging studies conducted so far, most have focused on the

22q11.2 deletion syndrome. This deletion is associated with a number of physical phenotype manifestations as well as high risk for psychopathology (Jonas et al., 2014; Niarchou et al., 2014; Schneider et al., 2014; Owen and Doherty, 2016). The presence of a 22q11.2 deletion has been linked to alterations in brain structure and function (Zinkstok and Amelsvoort, 2005; Boot and van Amelsvoort, 2013; Reddaway et al., 2018; Sun et al., 2018), including disrupted structural connectivity (Ottet et al., 2013; Villalón-Reina et al., 2019). Although fewer studies have investigated functional connectivity, they report similarly disrupted networks using functional MRI (Debbané et al., 2012; Padula et al., 2015; Scariati et al., 2016b) and EEG (Tomescu et al., 2014).

Despite emerging evidence of white matter alterations associated with other ND-CNVs (Berman et al., 2016; Drakesmith et al., 2019; Silva et al., 2019), very few studies have investigated their functional correlates. Recent electrophysiological research reported increased beta-band activity in participants with 15q11.2-q13.1 duplications (Frohlich et al., 2016, 2019) and 16p11.2 deletions (Hinkley et al., 2019), as well as delayed evoked responses in the latter (Berman et al., 2016; Jenkins et al., 2016). Based on current evidence it is difficult to assess the extent of functional connectivity alterations, especially for rarer ND-CNVs.

To address this, we investigated oscillatory connectivity measured with MEG in participants with ND-CNVs at nine different loci. Given the common phenotypic outcomes associated with ND-CNVs (Chawner et al., 2019), this approach can identify convergent endophenotypes of potentially higher clinical relevance. Because a third of our cohort had 22q11.2 deletions, we also investigated alterations in connectivity separately in this subgroup and in the group of participants with other ND-CNVs. This allowed us to assess the specificity of the effects, especially considering previous findings of widespread neural alterations associated with 22q11.2 deletions.

In both subgroups, we found evidence of disrupted alpha and beta-band oscillatory connectivity

in posterior brain regions. Furthermore, using graph theory measures of network topology and

information transfer, we were able to identify participants with ND-CNVs based on their

individual connectivity maps. The two approaches highlighted common patterns of

dysconnectivity in participants with ND-CNVs, as well as specific network features that might

be linked to CNV pathogenicity.

Methods

Participants

MEG data were acquired in 42 adults with ND-CNVs targeted for their high penetrance for

neurodevelopmental disorders (22 female; mean age 38.5±12.5 years). ND-CNVs at nine

different loci were represented in the cohort, with 14 (33%) participants carrying 22q11.2

deletions (Table 1).

Recruitment was performed through NHS genetics clinics and relevant support groups within

the UK. Written consent was obtained in accordance with The Code of Ethics of the World

Medical Association (Declaration of Helsinki), and all procedures were approved by the South

East Wales Research Ethics Committee.

All participants with ND-CNVs were assessed using the Psychiatric Assessment Schedule for

Adults with Developmental Disability (PAS-ADD; Moss et al., 1997). 24 were also assessed

using the Structured Interview for Prodromal Symptoms (SIPS; Miller et al., 2003), and 26

using the Structured Clinical Interview for DSM (SCID II; First and Gibbon, 2004). Diagnoses

were assigned by research psychologists and verified by an adult psychiatrist using the

Diagnostic and Statistical Manual for Mental Disorders, 4th and 5th Editions. Of the 42

participants, 34 had at least one diagnosis; 22 had anxiety disorders, 15 had mood disorders,

and 15 had neurodevelopmental disorders (including 6 with intellectual disability and 6 with

autism spectrum disorders). Five participants exhibited full or attenuated psychotic symptoms, of whom four met the criteria for a psychotic disorder, with two schizophrenia diagnoses in the 22q11.2 deletion group. There was no significant difference in number of diagnoses between the 22q11.2 deletion group and the other ND-CNV group (t(33.7) = 0.14, P = 0.89).

IQ tests were administered using the Wechsler Adult Intelligence Scale (WAIS-III). We report Verbal IQ (VIQ), Performance IQ (PIQ) and Full Scale IQ scores (FSIQ). There was no significant difference in IQ between the 22q11.2 deletion group and the other ND-CNV group (t(22.7-25.1) < 0.57, P > 0.58).

In the ND-CNV group, 62% of participants were taking medication for physical, neurological or mood disorders (e.g. high blood pressure, asthma, pain/migraine, depression/anxiety), with the most common medications including gabapentin, co-codamol (combination of codeine and paracetamol), and fluoxetine (N = 3). Given the high variability of medications, their effects could not be systematically investigated; however, their impact was alleviated in MEG analysis by tests of generalizibility (e.g. resampling).

Controls were selected among resting-state datasets acquired at CUBRIC as part of the "100 Brains"' and "UK MEG Partnership" projects. These cohorts included healthy participants aged 18-65 with no history of neurological or neuropsychiatric disorders, and 42 controls were chosen to match the gender and age of the ND-CNV carriers as closely as possible (22 female; mean age 33.3±9.6 years). These measurements were acquired under protocols approved by the Cardiff University School of Psychology Ethics Committee.

Since a third of the ND-CNV cohort consisted of participants with 22q11.2 deletions, we assessed the impact of this subgroup by repeating all analyses on (1) participants with other ND-CNVs (except 22q11.2 deletions) and their matched controls (N = 56), and (2) participants with 22q11.2 deletions and their matched controls (N = 28).

CNV and critical region (hg19)	Total N	N female	Age (mean±SD)	N diag (mean±SD)	VIQ (mean±SD)	PIQ (mean±SD)	FSIQ (mean±SD)
All CNV	*42*	*22*	*38.53±12.55*	*2.93±2.54*	*88.72±18.84*	*96.63±14.63*	*91.24±17.15*
22q11.2del chr22:19,037,332-21,466,726	14	9	38.97±16.43	3±2.07	86.28±21.23	95.28±15.6	89.93±17.88
22q11.2dup chr22:19,037,332-21,466,726	4	13	38.32±10.44	2.89±2.78	90±17.77	97.33±14.36	91.89±17.08
17q12dup chr17:34,815,904-36,217,432	2						
16p11.2del chr16:29,650,840-30,200,773	1						
16p11.2dup (distal) chr16:28,823,196-29,046,783	1						
15q13.1-13.3del (BP4-5) chr15:31,080,645-32,462,776	2						
15q13.1-13.3dup (BP4-5) chr15:31,080,645-32,462,776	1						
15q11.2del (BP1-2) chr15:22,805,313-23,094,530	7						
15q11.2dup(BP1-2) chr15:22,805,313-23,094,530	1						
15q11.2q12dup(BP2-3; PWS/AS) chr15:22,805,313-28390339	1						
3q29del chr3:195,720,167-197,354,826	1						
2p16.3del (NRXN1) chr2:50145643-51259674	1						
1q21.1del chr1:146,527,987-147,394,444	5						
1q21.1dup chr1:146,527,987-147,394,444	1						
Controls	*42*	*22*	*33.35±9.58*	-	-	-	-

Table 1. Participant information (ND-CNV status, demographic information, number of psychiatric diagnoses, and IQ). Note that the 22q11.2 deletion group included two atypical deletions and one adjacent deletion.

Genotyping

Participants with ND-CNVs were genotyped using the Illumina HumanCoreExome whole genome SNP array, which contained an additional 27 000 genetic variants at loci previously linked to neurodevelopmental disorders, including CNVs. The raw intensity data was

processed using Illumina Genome Studio software (version 2011.1). PennCNV (version 1.0.3)

was used to perform CNV calling in order to confirm the presence of the ND-CNV in each

case sample, with each CNV being required to span a minimum of 10 informative SNPs and to

be at least 10 kb in length. CNV coordinates were specified according to genome version hg19,

and the boundaries of each CNV were confirmed by manually inspecting the Log R ratio and

B allele frequency plots at each of the genomic regions of interest (Table 1). Genetic

information was not available for control participants; given the rarity of these genotypes in

the general population, they were assumed to carry no ND-CNVs.

Data collection

Five-minute resting-state MEG recordings were made using a 275-channel CTF radial

gradiometer system (CTF, Vancouver, Canada) at a sampling rate of 1200 Hz. Three of the

sensors were turned off due to excessive noise, and 29 reference channels were recorded to

improve noise cancellation(Vrba, 2002). During the recordings, participants were seated

upright and fixated their eyes on a red fixation point presented centrally on either a CRT

monitor or LCD projector. Three electromagnetic coils were placed at fiducial locations (nasion

and pre-auricular) for head localization.

To aid in source localization, structural T1-weighted MRI scans were also acquired using a 3T

General Electric or Siemens MRI scanner with a 1 mm isotropic FSPGR/MPRAGE pulse

sequence.

Data analysis

Pre-processing

To remove muscle artifacts, a semi-automatic procedure was implemented using the FieldTrip

toolbox (Oostenveld et al., 2011) and MATLAB R2015a. Sensor time-series were bandpass-

filtered between 110 and 140 Hz and z-transformed; segments exceeding a participant-specific

z-score threshold were removed. Next, eye movement and cardiac artifacts were projected out

of the data using Independent Component Analysis (ICA). Noisy channels exhibiting high

variance were also removed from the data where necessary. There was no significant difference

in recording duration after artifact rejection between the ND-CNV and control groups (t(81.8)

= 1.61, P = 0.11, mean duration 255.88±29.31 s and 245.33±30.86 s respectively).

Head motion was monitored continuously in 18/42 ND-CNV datasets and 40/42 control

datasets, and head localization was performed at the start and end of the recording in the

remaining datasets. There was no significant difference between the ND-CNV and control

groups in maximum head coil displacement between the beginning and end of the recording

(t(79.8) = 0.85, P = 0.39, mean displacement 2.07±3.62 mm and 2.7±3.06 mm respectively).

In datasets with continuous head localization, the maximum distance of the head coils from

their average position across the entire recording did not significantly differ between groups

(t(39.4) = 1.44, P = 0.16, mean distance 4.74±3.5 mm and 3.21±4.2 mm respectively).

Prior to source localization, coregistration was performed by manually marking head coil

locations on each participant's MRI using FieldTrip. The data were downsampled to 600 Hz

and bandpass-filtered in six different frequency bands: delta (2-4 Hz), theta (4-8 Hz), alpha (8-

13 Hz), beta (13-30 Hz), low gamma (40-60 Hz), and high gamma (60-90 Hz).

Figure 1. Overview of the analysis pipeline. Resting-state MEG data were preprocessed, filtered into six frequency bands, and projected into source space. Hilbert envelopes were calculated at 90 AAL-atlas-based virtual sensor locations, and correlated to obtain functional connectivity matrices. These were z-scored and rank-thresholded at the group level for between-group analyses, and at the subject level for data-driven prediction of ND-CNV status using graph theory.

Estimating functional connectivity

To assess group differences in resting-state connectivity (Figure 1), we focused on amplitude-amplitude coupling of source-localized oscillatory signals (Colclough et al., 2016). Continuous data in each of the six frequency bands were projected into source space using a Linearly Constrained Minimum Variance (LCMV) beamformer. Sources were reconstructed on a 6 mm template grid warped to each participant's MRI, using a multiple local-spheres forward model (Huang et al., 1999). To alleviate the depth bias, beamformer weights were normalized by their vector norm (Hillebrand et al., 2012).

Next, 90 nodes corresponding to cortical regions of interest (ROI) in the Automated

Anatomical Labelling (AAL) atlas (Tzourio-Mazoyer et al., 2002) were identified by

performing a frequency analysis on all sources within each ROI and selecting the source with

the largest temporal standard deviation. Continuous virtual sensor timecourses corresponding

to the 90 nodes were then reconstructed and bandpass-filtered into the frequency bands of

interest.

To avoid spurious correlations, the node time-series were orthogonalized using a multivariate

symmetric orthogonalization approach (Colclough et al., 2015). A Hilbert transform was used

to calculate oscillatory amplitude envelopes, which were then despiked using a median filter,

downsampled to 1 Hz, and trimmed to avoid filter and edge effects. To obtain connectivity

matrices, pairwise correlations were calculated between the 90 Hilbert envelopes. Next, a

Fisher transform was applied to obtain z-scores with zero mean and unit variance across

connections in each participant's map. This procedure corrected for possible systematic

differences across participants, for example due to differences in data quality (Siems et al.,

2016).

Intracranial volume (ICV), quantified as the number of 1-mm isotropic voxels inside the brain,

was smaller in the ND-CNV group than the control group ($t(65.5) = -2.19$, $P = 0.03$), in line

with some previous reports (Hu et al., 2017; Sønderby et al., 2018; Warland et al., 2019). The

potential impact of this difference on the MEG results was alleviated through the source

localization procedure and the z-scoring of the connectivity matrices.

In addition to the six frequency bands listed above, a combined measure of connectivity was

obtained by calculating the vector-sum of connectivity matrices across all frequency bands

(Koelewijn et al., 2019).

Group differences in resting-state connectivity

To reduce the impact of noise, a conservative ranking procedure(Koelewijn et al., 2019) was used to threshold the connectivity maps for the purposes of between-group comparisons. This consisted of calculating the rank of each connection in participant-level connectivity matrices and averaging the resulting rank map across participants in each group. Only the top 20% edges in the average rank map were considered "valid" and selected for further analysis. To ensure that large differences in signal across cohorts were not discarded by this procedure, the rank-thresholding procedure was performed separately in each cohort, and connections determined as "valid" in either cohort were included in further analysis.

To assess differences between groups, Welch's t-tests were conducted at each valid edge. Significant edges were identified using an uncorrected $\alpha = 0.05$. Correction for multiple comparisons was applied using a randomization procedure with 10000 sign-shuffling iterations and maximal statistic thresholding (omnibus $\alpha = 0.05$)(Nichols and Holmes, 2001).

The robustness of cohort differences was evaluated through a resampling procedure. Increases and decreases in connectivity between groups were tabulated using random samples of half of each group. This was repeated across 10000 iterations, and edges showing a consistent effect direction across at least 95% of iterations were considered robust.

To control for potential confounds (for example, resulting from imperfect age matching between the ND-CNV and control groups), an additional multiple regression analysis was performed. Combined-frequency connectivity matrices were entered as response variables with a categorical predictor (ND-CNV presence) and three covariates (age, gender, and ICV). A resampling procedure as described above was performed to assess the robustness of between-group differences. The sign of the regression slope associated with the main predictor was tabulated across 1000 split-half cohort randomizations. Edges showing consistent effects across 95% of iterations were considered robust.

Individual networks: identifying participants with ND-CNVs using graph theory

Next, a data-driven graph theory approach was used to assess whether participants with ND-CNVs could be distinguished from unaffected controls using functional connectivity features. To this aim, the cohort was divided into training and test sets using an iterated cross-validation procedure.

This analysis focused on individual networks by selecting the top 20% of connections in each participant's normalized connectivity map as the basis for undirected graphs. Networks were then characterized using six nodal graph theory metrics. These included three measures of node connectedness: *degree* (the number of connections linking each node to other nodes); *betweenness centrality* (the fraction of shortest paths between nodes containing a given node); and *eccentricity* (the maximal shortest path from a node to any other node). *Global efficiency* (the average inverse shortest path between a node and all others) was evaluated as a measure of network integration. Finally, two metrics captured network modularity: *local efficiency* (the average inverse shortest path between a node and its neighbours) and *clustering coefficient* (the fraction of connected node triplets around a node). All metrics with the exception of node degrees were weighted by the inverse of the normalized connectivity matrices; in other words, stronger amplitude-amplitude coupling was taken to reflect shorter paths between nodes. Graph theory analyses were performed using MATLAB R2019a and the Brain Connectivity Toolbox (Rubinov and Sporns, 2010).

To discriminate between groups, a linear support vector machine (SVM) classifier (Boser et al., 1992) was trained on each of the node metrics. Additionally, a pooled feature vector was created by combining the six metrics to maximize the amount of complementary information input to the classifier. This approach makes use of information across all nodes while avoiding the need for multiple testing.

Classification was performed between the ND-CNV and control groups, as well as between the

two ND-CNV subgroups (22q11.2 deletions and other ND-CNVs) and their matched controls.

To avoid overfitting, model performance was evaluated using 100 iterations of stratified five-

fold cross-validation. This entailed iteratively leaving out a fifth of the data for testing and

training the model on the remaining data, whilst ensuring balanced group representation in each

fold. Performance was quantified using accuracy (proportion correctly classified observations),

sensitivity (true positive rate) and specificity (true negative rate) in order to highlight any

asymmetries in ND-CNV and control identification. Furthermore, significance was assessed

by shuffling the true labels 5000 times and recomputing classifier accuracy to estimate the

empirical chance level and calculate a one-tailed p-value (Nichols and Holmes, 2001).

Figure 2. A. Differences in resting-state connectivity (amplitude correlations) between participants with ND-CNVs and controls. Connectivity increases and decreases in the ND-CNV group are shown in red and blue respectively. The rows show (top to bottom): valid connections after mean-rank thresholding in each frequency band; uncorrected (*P*<0.05) differences between groups; multiple comparison-corrected (omnibus *P*<0.05) differences between groups; and connections showing consistent increases/decreases in 95% of cohort resampling iterations. **B.** As in A, for subgroups excluding participants with 22q11.2 deletions and their matched controls (left) or including only participants with 22q11.2 deletions and their matched controls (right). To facilitate comparison, "valid" connections were the same as in A. Only frequency bands with surviving "valid" connections are shown. **C.** Nodes displayed in the circular plots, labelled and colour-coded by region. Connections shown are

251 the result of the conjunction analysis between the 22q11.2 deletion group and the other ND-CNV group (also see

252 Figure 3: blue connections are decreased in both groups; red connections have opposing signs).

Results

Connectivity alterations associated with ND-CNVs

The analysis of group differences in oscillatory connectivity revealed the largest number of

valid connections in the alpha and beta bands (Figure 2). Most of the significantly different

connections showed a decrease in oscillatory connectivity between posterior, parietal and

temporal nodes in the ND-CNV group, with the exception of a few right-hemisphere edges.

More extensive cohort effects were detected using the combined frequency maps (61 edges

exceeded the uncorrected threshold, compared to 1, 28 and 42 in the theta, alpha and beta

bands). These patterns were robust to random sub-sampling of the cohort, suggesting that they

were not driven by individual subjects. A small number of left-hemisphere connections,

including the precuneus, early visual cortex, and parietal regions, survived omnibus correction

for multiple comparisons.

Importantly, a similar pattern of hypoconnectivity was observed even after excluding

participants with 22q11.2 deletions and their matched controls (Figure 2B-C). Both ND-CNV

subgroups showed decreased posterior connectivity (Figure 3; 18 connections decreased in

both groups), indicating that the overall pattern was not driven by the 22q11.2 deletion group.

This pattern occurred despite higher heterogeneity in the mixed ND-CNV group (Figure 4) and

might thus reflect convergent alterations across genotypes. On the other hand, participants with

22q11.2 deletions exhibited more right-hemisphere hyperconnectivity compared to controls.

These effects spanned the precuneus and parietal cortex, as well as frontal regions, suggesting

some overlap with the default mode network (DMN).

To ensure that these differences were not affected by potential confounds, the cohort

resampling tests on combined frequency maps were repeated as a multiple linear regression

with age, gender and ICV as covariates. This analysis revealed fewer connections (65 compared

to the original 92 in the whole cohort analysis), but largely similar patterns of dysconnectivity

(Figure 3B).

Furthermore, although IQ scores could not be included in this analysis because they were not

available for the control group, IQ scores in the ND-CNV group significantly correlated with

connectivity strength at only four edges (max Pearson's r = 0.5, P<0.05).

Figure 3. Differences in connectivity are not driven by age, gender, and intracranial volume. Connections meeting

the 95% confidence criterion in the cohort resampling test are displayed for all group tests (first three columns).

The last column shows supra-threshold connections in both the 22q11.2 deletion group and the other ND-CNV

group; they are shown in blue where they are decreased in both groups, and in red where they have opposite signs.

Line width increases with effect robustness. **A**. Connections exhibiting robust differences based on the cohort

resampling test of combined frequency matrices, plotted on the template brain. **B.** As in A, after including age,

gender and intracranial volume as covariates in a multiple linear regression with "ND-CNV presence" as a main

categorical predictor.

Figure 4. Variability in connection z-scores across participant groups. The distribution of standard deviations

(SD) across valid connections in the combined frequency maps are shown for: **A.** the ND-CNV and control groups;

B. the two ND-CNV subgroups. The mixed ND-CNV group displays larger variability. Plots were generated using

the RainCloudPlots toolbox (Allen et al., 2019).

Network features as predictors of ND-CNV status

A graph theory framework was employed to identify participants with ND-CNVs from their

functional networks based on combined frequency maps. This approach has the advantage of

reducing dimensionality and complements the edge-focused group testing approach described

above.

Graph theory metrics were successful predictors of ND-CNV participants relative to unaffected

controls. The best prediction accuracy was achieved by combining the 6 node features (Table

2; Figure 5A; maximum accuracy 71%±3.44, *P*=0.0002). However, participants with 22q11.2

deletions were more consistently correctly classified than those with other ND-CNVs (Figure

5B).

Figure 5. Classifying participants with ND-CNVs and unaffected controls from individual MEG functional networks using graph theory. **A:** Classification performance on the three groups, using different metrics to characterize the networks (eccentricity, degree, betweenness centrality, global and local efficiency, clustering coefficient, and the pooled model). Above-chance accuracies (permutation testing) are marked with 1, 2, and 3 dots respectively for *p*<0.05, *p*<0.01, and *p*<0.001. **B.** How well are different ND-CNVs classified? The plot shows the mean predicted label for each of the 42 participants with ND-CNVs across 100 cross-validation iterations using pooled node features. Participants with 22q11.2 deletions are most consistently correctly identified. Two participants with schizophrenia diagnoses are marked with "S".

This was confirmed by subgroup classification analyses, which also pointed to subgroup differences. When excluding participants with 22q11.2 deletions, the best decoding accuracies were achieved using node eccentricities (65.62%±3.91, *P*=0.0016), node degrees, and the joint feature model. On the other hand, all node features were successful in discriminating participants with 22q11.2 deletions from their matched controls, with the best performance obtained using the clustering coefficient (87.61%±4.95, *P*=0.0002).

These results point to commonalities in network features (such as decreased centrality) that allow for the successful classification of participants with ND-CNVs across distinct genotypes. On the other hand, the features are specific enough to allow successful discrimination between participants with 22q11.2 deletions and other ND-CNVs (Table 2). Given the higher overall burden of 22q11.2 deletions in neurodevelopmental disorders (Kirov et al., 2014), this suggests that increased neurodevelopmental risk may be associated with more salient alterations in network function and may underpin specific genotype effects.

| | ND-CNV vs Controls (whole cohort) | | | | | | | 22q11.2del vs other ND-CNV |
	Eccentricity	Degree	Betweenness centrality	Global efficiency	Local efficiency	Clustering coefficient	Pooled node features	Pooled node features
Accuracy (%)	69.41	62.74	59.25	62.1	64.44	58.9	71	72.43
SD	3.14	3.25	4.3	3.91	3.7	3.72	3.44	1.63
P-value	0.0002	0.0012	0.0108	0.001	0.0006	0.022	0.0002	0.0036
Sensitivity	0.6	0.59	0.56	0.59	0.6	0.54	0.67	0.67
Specificity	0.79	0.67	0.63	0.65	0.68	0.64	0.75	0.75
	ND-CNV vs Controls (excluding 22q11.2del and matched controls)							Train on 22q11.2del, test on other ND-CNV
Accuracy (%)	65.62	60.27	52.45	56.69	56.06	57.82	63.43	55.36
SD	3.91	4.12	4.65	4.14	4.7	4.72	4.16	-
P-value	0.0016	0.0242	0.2967	0.1038	0.1032	0.053	0.0068	0.225
Sensitivity	0.58	0.54	0.51	0.56	0.55	0.53	0.58	0.21
Specificity	0.74	0.66	0.54	0.57	0.58	0.62	0.69	0.89
	ND-CNV vs Controls (22q11.2del vs matched controls only)							Train on other ND-CNV, test on 22q11.2del
Accuracy (%)	75.31	64.56	66.91	72.75	85.53	87.61	87.07	75
SD	4.7	5.69	6.52	4.89	3.94	4.95	3.69	-
P-value	0.0002	0.0296	0.0118	0.003	0.0002	0.0002	0.0002	0.0048
Sensitivity	0.58	0.54	0.61	0.62	0.78	0.83	0.8	0.57
Specificity	0.93	0.75	0.73	0.84	0.93	0.92	0.94	0.93

Table 2. Network-based classification results. Note that cross-decoding across ND-CNV subgroups achieves high

specificity but not sensitivity (middle-right and lower-right cells). This suggests that classifier performance is

driven mainly by network features in the control group.

For classification between participants with 22q11.2 deletions and other ND-CNVs (top right cell), 100 subsamples

of 14 participants were drawn from the 28 carriers of other ND-CNVs, gender-matched to the 22q11.2 deletion

group. The subgroups were similar in age, ensuring balanced classes in terms of sample size, gender, and age.

Predicted labels were then averaged across cohort resamplings (mean 50±5.57 repetitions per participant).

Discussion

To our knowledge, the present study provides the first insight into oscillatory connectivity

alterations in people with rare ND-CNVs. Using both an established group analysis pipeline

and a data-driven graph theory framework, we uncovered a pattern of functional

dysconnectivity affecting posterior regions in participants with ND-CNVs. These patterns were

robust to effects of age, gender, and intracranial volume, and emerged despite a conservative

thresholding approach restricted to the most reproducible connections.

Effects originated in the alpha and beta frequency bands, which are thought to underpin long-

range communication between brain areas (Schnitzler and Gross, 2005). Connections linking

parietal, temporal, and occipital areas were most consistently affected in both the 22q11.2

deletion group and the other ND-CNV group. Similar patterns have been previously reported

in schizophrenia patients (Brookes et al., 2016), including alpha-band parietal

hypoconnectivity in first-episode schizophrenia (Phalen et al., 2019). Furthermore, posterior

structural network alterations have been identified as an early marker of ASD (Lewis et al.,

2014), suggesting a link between such alterations and increased neurodevelopmental risk.

Similar connectivity changes in the visual processing system and the default mode network

have been shown in people with 22q11.2 deletions using structural and functional MRI

(Schreiner et al., 2014; Scariati et al., 2016a; Larsen et al., 2019). Here, we found that these

effects are not restricted to the 22q11.2 deletion group, suggesting that long-range connectivity

could act as a common marker across genetic variants. Although non-invasive measurements

cannot provide direct mechanistic insight, this is consistent with potential alterations in

excitatory-inhibitory balance (Deco et al., 2014; Gu et al., 2019) as a mechanism for pleiotropic

genetic effects underlying neurodevelopmental disorders (Gao and Penzes, 2015; Foss-Feig et

al., 2017; Lee et al., 2017). This is thought to occur through increased excitation or disinhibition

caused by gene haploinsufficiency and mediated by impaired GABA and NMDA receptor

function (Kehrer et al., 2008; Ramamoorthi and Lin, 2011; Pocklington et al., 2015).

Despite sample size limitations, differences between the two subgroups also point to effects

specific to the highly penetrant 22q11.2 deletions. Hypoconnectivity was more extensive in

people with other ND-CNVs, while the 22q11.2 deletion group exhibited more focused

patterns; these were robust to cohort resampling, suggesting that they are unlikely to be driven

by individual cases. These differences were reflected in the graph theory analysis. Although all

network features were altered in the 22q11.2 deletion group, their increased modularity was

particularly discriminative, in line with previous reports of increased structural network

segregation in people with 22q11.2 deletions (Ottet et al., 2013; Scariati et al., 2016b; Sandini

et al., 2018). For other ND-CNVs, the only predictive features were centrality measures

(specifically the node eccentricity and degree), reflecting hypoconnectivity in the ND-CNV

cohort. Between and within-group classification results (Table 2) highlight the ability of graph

theory metrics to capture both convergent and specific network alterations, which could help

elucidate the link between CNV pathogenicity and neural system function.

We alleviated concerns of potential systematic between-group differences unrelated to

genotype by using conservative approaches (e.g. rejecting weaker connections that may

introduce noise) and resampling procedures. Although head motion did not appear to differ

between groups, this was not continuously measured in all participants and so we could only

partially rule out its effects.

Although the present study was able to evaluate ND-CNV effects independently of the

contribution of highly penetrant 22q11.2 deletions, the limited sample size remains a concern

common in CNV imaging research. The high genotype variability within the cohort makes the

specificity of these effects difficult to assess, particularly with regard to differences between

the 22q11.2 deletion group and other ND-CNVs. Variability within the heterogeneous ND-

CNV group was higher than within the 22q11.2 deletion group (Figure 4), suggesting that our

focus on convergent alterations may obscure specific effects. However, the fact that we see

differences robust to resampling in this group, despite its heterogeneity, points to common

connectivity alterations across distinct genotypes. Studies recruiting larger samples of

participants with ND-CNVs, for example through large multi-site collaborations, are necessary

to evaluate the generalizability of connectivity patterns and graph theory metrics as

"fingerprints" associated with ND-CNV status.

In summary, the present study assessed oscillatory long-range connectivity as a potential

marker of pathogenic genetic effects across a range of rare ND-CNVs. Occipital, parietal and

temporal brain areas were characterized by consistent hypoconnectivity across genotypes,

which was not exclusively driven by the presence of a large number of participants with highly-

penetrant 22q11.2 deletions. Functional networks in the ND-CNV group exhibited decreased

node centrality and alterations in network efficiency and structure. Furthermore, features

specific to highly penetrant variants were present alongside convergent network alterations and

enabled successful ND-CNV classification. These results are consistent with a common

mechanism for genetic risk, based on an altered balance between excitatory and inhibitory

synaptic processes and leading to network dysfunction. We propose that these functional

connectivity alterations are an intermediate phenotype on the pathway from synaptic molecular

changes to disruption of cognitive function and psychotic illness.

Conflict of interest

The authors declare no competing interests.

Acknowledgements

The authors wish to thank: Ffion Evans and Kali Barawi for assistance with psychometric and

clinical data collection; Eirini Messaritaki for helpful comments on the graph theory analysis;

George Kirov for help with genetic data; and Alexander Shaw for the SourceMesh visualization

toolbox. This work was supported by a Wellcome Trust Strategic Award (100202/Z/12/Z), the

MRC UK MEG Partnership Grant (MR/K005464/1), and the MRC Doctoral Training Grant

(MR/K501086/1).

References

Allen M, Poggiali D, Whitaker K, Marshall TR, Kievit RA (2019) Raincloud plots: A multi-platform
tool for robust data visualization [version 1; peer review: 2 approved]. Wellcome Open Res 4.

Berman JI, Chudnovskaya D, Blaskey L, Kuschner E, Mukherjee P, Buckner R, Nagarajan S, Chung
WK, Sherr EH, Roberts TPL (2016) Relationship between M100 Auditory Evoked Response
and Auditory Radiation Microstructure in 16p11.2 Deletion and Duplication Carriers. AJNR Am
J Neuroradiol 37:1178–1184.

Boot E, van Amelsvoort TAMJ (2013) Neuroimaging Correlates of 22q11.2 Deletion Syndrome:
Implications for Schizophrenia Research. Curr Top Med Chem 12:2303–2313.

Boser BE, Guyon IM, Vapnik VN (1992) A training algorithm for optimal margin classifiers. In:
Proceedings of the fifth annual workshop on Computational learning theory - COLT '92, pp

144–152. New York, New York, USA: ACM Press.

Brookes MJ, Tewarie PK, Hunt BAE, Robson SE, Gascoyne LE, Liddle EB, Liddle PF, Morris PG

(2016) A multi-layer network approach to MEG connectivity analysis. Neuroimage 132:425–

438.

Chawner SJRA, Owen MJ, Holmans P, Raymond FL, Skuse D, Hall J, van den Bree MBM (2019)

Genotype-phenotype associations in children with copy number variants associated with high

neuropsychiatric risk in the UK (IMAGINE-ID): a case-control cohort study. The Lancet

Psychiatry 6:493–505.

Colclough GL, Brookes MJ, Smith SM, Woolrich MW (2015) A symmetric multivariate leakage

correction for MEG connectomes. Neuroimage 117:439–448.

Colclough GL, Woolrich MW, Tewarie PK, Brookes MJ, Quinn AJ, Smith SM (2016) How reliable

are MEG resting-state connectivity metrics ? Neuroimage 138:284–293.

Debbané M, Lazouret M, Lagioia A, Schneider M, Van De Ville D, Eliez S (2012) Resting-state

networks in adolescents with 22q11.2 deletion syndrome: Associations with prodromal

symptoms and executive functions. Schizophr Res 139:33–39.

Deco G, Ponce-Alvarez A, Hagmann P, Romani GL, Mantini D, Corbetta M (2014) How local

excitation-inhibition ratio impacts the whole brain dynamics. J Neurosci 34:7886–7898.

Dong D, Wang Y, Chang X, Luo C, Yao D (2018) Dysfunction of Large-Scale Brain Networks in

Schizophrenia: A Meta-analysis of Resting-State Functional Connectivity. Schizophr Bull

44:168–181.

Drakesmith M, Parker GD, Smith J, Linden SC, Rees E, Williams N, Owen MJ, van den Bree M, Hall

J, Jones DK, Linden DEJ (2019) Genetic risk for schizophrenia and developmental delay is

associated with shape and microstructure of midline white-matter structures. Transl Psychiatry

9:102.

First MB, Gibbon M (2004) The Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I)

and the Structured Clinical Interview for DSM-IV Axis II Disorders (SCID-II). In:

Comprehensive handbook of psychological assessment, Vol. 2. Personality assessment

(Hilsenroth MJ, Segal DL, eds), pp 134–143. Hoboken, NJ, US: John Wiley & Sons Inc.

Foss-Feig JH, Adkinson BD, Ji JL, Yang G, Srihari VH, McPartland JC, Krystal JH, Murray JD,

Anticevic A (2017) Searching for Cross-Diagnostic Convergence: Neural Mechanisms

Governing Excitation and Inhibition Balance in Schizophrenia and Autism Spectrum Disorders.

Biol Psychiatry 81:848–861.

Friston K, Brown HR, Siemerkus J, Stephan KE (2016) The dysconnection hypothesis (2016).

Schizophr Res 176:83–94.

Frohlich J, Reiter LT, Saravanapandian V, DiStefano C, Huberty S, Hyde C, Chamberlain S, Bearden

CE, Golshani P, Irimia A, Olsen RW, Hipp JF, Jeste SS (2019) Mechanisms underlying the EEG

biomarker in Dup15q syndrome. Mol Autism 10:29.

Frohlich J, Senturk D, Saravanapandian V, Golshani P, Reiter LT, Sankar R, Thibert RL, DiStefano

C, Huberty S, Cook EH, Jeste SS (2016) A Quantitative Electrophysiological Biomarker of

Duplication 15q11.2-q13.1 Syndrome Ravel N, ed. PLoS One 11:e0167179.

Gao R, Penzes P (2015) Common mechanisms of excitatory and inhibitory imbalance in

schizophrenia and autism spectrum disorders. Curr Mol Med 15:146–167.

Gu H, Hu Y, Chen X, He Y, Yang Y (2019) Regional excitation-inhibition balance predicts default-

mode network deactivation via functional connectivity. Neuroimage 185:388–397.

Hillebrand A, Barnes GR, Bosboom JL, Berendse HW, Stam CJ (2012) Frequency-dependent

functional connectivity within resting-state networks : An atlas-based MEG beamformer

solution. Neuroimage 59:3909–3921.

Hinkley LBN, Dale C, Luks TL, Findlay AM, Bukshpun P, Pojman N, Thieu T, Chung WK, Berman

J, Roberts TPL, Mukherjee P, Sherr EH, Nagarajan SS (2019) Sensorimotor cortical oscillations

during movement preparation in 16p11.2 deletion carriers. J Neurosci:3001–3017.

Hu M-L, Zong X-F, Mann JJ, Zheng J-J, Liao Y-H, Li Z-C, He Y, Chen X-G, Tang J-S (2017) A

Review of the Functional and Anatomical Default Mode Network in Schizophrenia. Neurosci

Bull 33:73–84.

Huang MX, Mosher JC, Leahy RM (1999) A sensor-weighted overlapping-sphere head model and

exhaustive head model comparison for MEG. Phys Med Biol 44:423–440.

Hull J V., Dokovna LB, Jacokes ZJ, Torgerson CM, Irimia A, Van Horn JD (2017) Resting-State

Functional Connectivity in Autism Spectrum Disorders: A Review. Front Psychiatry 7:205.

Jenkins J, Chow V, Blaskey L, Kuschner E, Qasmieh S, Gaetz L, Edgar JC, Mukherjee P, Buckner R,

Nagarajan SS, Chung WK, Spiro JE, Sherr EH, Berman JI, Roberts TPL (2016) Auditory

Evoked M100 Response Latency is Delayed in Children with 16p11.2 Deletion but not 16p11.2

Duplication. Cereb Cortex 26:1957–1964.

Jonas RK, Montojo CA, Bearden CE (2014) The 22q11.2 deletion syndrome as a window into

complex neuropsychiatric disorders over the lifespan. Biol Psychiatry 75:351–360.

Karayiorgou M, Simon TJ, Gogos JA (2010) 22q11.2 microdeletions: linking DNA structural

variation to brain dysfunction and schizophrenia. Nat Rev Neurosci 11:402–416.

Kehrer C, Maziashvili N, Dugladze T, Gloveli T (2008) Altered excitatory-inhibitory balance in the

NMDA-hypofunction model of schizophrenia. Front Mol Neurosci 1:6.

Kirov G, Rees E, Walters JTR, Escott-Price V, Georgieva L, Richards AL, Chambert KD, Davies G,

Legge SE, Moran JL, McCarroll SA, Donovan MCO, Owen MJ (2014) The penetrance of copy

number variations for schizophrenia and developmental delay. Biol Psychiatry 75:378–385.

Koelewijn L, Lancaster TM, Linden D, Dima DC, Routley BC, Magazzini L, Barawi K, Brindley L,

Adams R, Tansey KE, Bompas A, Tales A, Bayer A, Singh K (2019) Oscillatory hyperactivity

and hyperconnectivity in young APOE-ε4 carriers and hypoconnectivity in Alzheimer's disease.

Elife 8:1–25.

Larsen KM, Dzafic I, Siebner HR, Garrido MI (2019) Alteration of functional brain architecture in

22q11.2 deletion syndrome – Insights into susceptibility for psychosis. Neuroimage 190:154–

171.

Lee E, Lee J, Kim E (2017) Excitation/Inhibition Imbalance in Animal Models of Autism Spectrum

Disorders. Biol Psychiatry 81:838–847.

Lewis JD, Evans AC, Pruett JR, Botteron K, Zwaigenbaum L, Estes A, Gerig G, Collins L,

Kostopoulos P, McKinstry R, Dager S, Paterson S, Schultz RT, Styner M, Hazlett H, Piven J

(2014) Network inefficiencies in autism spectrum disorder at 24 months. Transl Psychiatry

4:e388–e388.

Miller TJ, McGlashan TH, Rosen JL, Cadenhead K, Ventura J, McFarlane W, Perkins DO, Pearlson

QD, Woods SW (2003) Prodromal Assessment With the Structured Interview for Prodromal

Syndromes and the Scale of Prodromal Symptoms: Predictive Validity, Interrater Reliability,

and Training to Reliability. Schizophr Bull 29:703–715.

Moss S, Ibbotson B, Prosser H, Goldberg D, Patel P, Simpson N (1997) Validity of the PAS-ADD for

detecting psychiatric symptoms in adults with learning disability (mental retardation). Soc

Psychiatry Psychiatr Epidemiol 32:344–354.

Niarchou M, Chawner SJRA, Doherty JL, Maillard AM, Jacquemont S, Chung WK, Green-Snyder L,

Bernier RA, Goin-Kochel RP, Hanson E, Linden DEJ, Linden SC, Raymond FL, Skuse D, Hall

J, Owen MJ, Bree MBM van den (2019) Psychiatric disorders in children with 16p11.2 deletion

and duplication. Transl Psychiatry 9:8.

Niarchou M, Zammit S, van Goozen SHM, Thapar A, Tierling HM, Owen MJ, van den Bree MBM

(2014) Psychopathology and cognition in children with 22q11.2 deletion syndrome. Br J

Psychiatry 204:46–54.

Nichols TE, Holmes AP (2001) Nonparametric Permutation Tests For Functional Neuroimaging : A

Primer with Examples. Hum Brain Mapp 25:1–25.

O'Reilly C, Lewis JD, Elsabbagh M (2017) Is functional brain connectivity atypical in autism? A

systematic review of EEG and MEG studies Gozzi A, ed. PLoS One 12:e0175870.

Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for advanced

analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci

2011:156869.

Ottet M-C, Schaer M, Debbané M, Cammoun L, Thiran J-P, Eliez S (2013) Graph theory reveals

dysconnected hubs in 22q11DS and altered nodal efficiency in patients with hallucinations.

Front Hum Neurosci 7:402.

Owen MJ, Doherty JL (2016) What can we learn from the high rates of schizophrenia in people with

22q11.2 deletion syndrome? World Psychiatry 15:23–25.

Padula MC, Schaer M, Scariati E, Schneider M, Van De Ville D, Debbané M, Eliez S (2015)

Structural and functional connectivity in the default mode network in 22q11.2 deletion

syndrome. J Neurodev Disord 7:23.

Phalen H, Coffman BA, Ghuman A, Sejdić E, Salisbury DF (2019) Non-Negative Matrix

Factorization Reveals Resting-State Cortical Alpha Network Abnormalities in the First Episode

Schizophrenia-Spectrum. Biol Psychiatry Cogn Neurosci Neuroimaging.

Pocklington AJ, Rees E, Walters JTR, Han J, Kavanagh DH, Chambert KD, Holmans P, Moran JL,

McCarroll SA, Kirov G, O'Donovan MC, Owen MJ (2015) Novel Findings from CNVs

Implicate Inhibitory and Excitatory Signaling Complexes in Schizophrenia. Neuron 86:1203–

1214.

Ramamoorthi K, Lin Y (2011) The contribution of GABAergic dysfunction to neurodevelopmental

disorders. Trends Mol Med 17:452–462.

Reddaway JT, Doherty JL, Lancaster T, Linden D, Walters JT, Hall J (2018) Genomic and Imaging

Biomarkers in Schizophrenia. Curr Top Behav Neurosci 40:325–352.

Rubinov M, Sporns O (2010) Complex network measures of brain connectivity: Uses and

interpretations. Neuroimage 52:1059–1069.

Sandini C, Scariati E, Padula MC, Schneider M, Schaer M, Van De Ville D, Eliez S (2018) Cortical

Dysconnectivity Measured by Structural Covariance Is Associated With the Presence of

Psychotic Symptoms in 22q11.2 Deletion Syndrome. Biol Psychiatry Cogn Neurosci

Neuroimaging 3:433–442.

Scariati E, Padula MC, Schaer M, Eliez S (2016a) Long-range dysconnectivity in frontal and midline

structures is associated to psychosis in 22q11.2 deletion syndrome. J Neural Transm 123:823–

839.

Scariati E, Schaer M, Karahanoglu I, Schneider M, Richiardi J, Debbané M, Van De Ville D, Eliez S

(2016b) Large-scale functional network reorganization in 22q11.2 deletion syndrome revealed

by modularity analysis. Cortex 82:86–99.

Schneider M et al. (2014) Psychiatric Disorders From Childhood to Adulthood in 22q11.2 Deletion

Syndrome: Results From the International Consortium on Brain and Behavior in 22q11.2

Deletion Syndrome. Am J Psychiatry 171:627–639.

Schnitzler A, Gross J (2005) Normal and pathological oscillatory communication in the brain. Nat

Rev Neurosci 6:285–296.

Schreiner MJ, Karlsgodt KH, Uddin LQ, Chow C, Congdon E, Jalbrzikowski M, Bearden CE (2014)

Default mode network connectivity and reciprocal social behavior in 22q11.2 deletion

syndrome. Soc Cogn Affect Neurosci 9:1261–1267.

Sha Z, Wager TD, Mechelli A, He Y (2019) Common Dysfunction of Large-Scale Neurocognitive

Networks Across Psychiatric Disorders. Biol Psychiatry 85:379–388.

Siems M, Pape A-A, Hipp JF, Siegel M (2016) Measuring the cortical correlation structure of

spontaneous oscillatory activity with EEG and MEG. Neuroimage 129:345–355.

Silva AI, Ulfarsson MO, Stefansson H, Gustafsson O, Walters GB, Linden DEJ, Wilkinson LS,

Drakesmith M, Owen MJ, Hall J, Stefansson K (2019) Reciprocal White Matter Changes

Associated With Copy Number Variation at 15q11.2 BP1-BP2: A Diffusion Tensor Imaging

574 Study. Biol Psychiatry 85:563–572.

575 Sønderby IE et al. (2018) Dose response of the 16p11.2 distal copy number variant on intracranial

576 volume and basal ganglia. Mol Psychiatry.

577 Sun D et al. (2018) Large-scale mapping of cortical alterations in 22q11.2 deletion syndrome:

578 Convergence with idiopathic psychosis and effects of deletion size. Mol Psychiatry:1–13.

579 Tomescu MI, Rihs TA, Becker R, Britz J, Custo A, Grouiller F, Schneider M, Debbané M, Eliez S,

580 Michel CM (2014) Deviant dynamics of EEG resting state pattern in 22q11.2 deletion syndrome

581 adolescents: A vulnerability marker of schizophrenia? Schizophr Res 157:175–181.

582 Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B,

583 Joliot M (2002) Automated Anatomical Labeling of Activations in SPM Using a Macroscopic

584 Anatomical Parcellation of the MNI MRI Single-Subject Brain. Neuroimage 15:273–289.

585 Villalón-Reina JE et al. (2019) Altered white matter microstructure in 22q11 . 2 deletion syndrome : a

586 multisite diffusion tensor imaging study. Mol Psychiatry.

587 Vrba J (2002) Magnetoencephalography: The art of finding a needle in a haystack. Phys C Supercond

588 its Appl 368:1–9.

589 Warland A, Kendall KM, Rees E, Kirov G, Caseras X (2019) Schizophrenia-associated genomic copy

590 number variants and subcortical brain volumes in the UK Biobank. Mol Psychiatry:1.

591 Zinkstok J, Amelsvoort T van (2005) Neuropsychological Profile and Neuroimaging in Patients with

592 22Q11.2 Deletion Syndrome: A Review. Child Neuropsychol 11:21–37.