Sawfish, Read in Tooth and Saw: rostral teeth as endogenous
chemical records of movement and life-history in a critically
endangered species

Jens C. Hegg[1], Breanna Graves[1], Chris M. Fisher[2]*

[*]jhegg@uidaho.edu

[1]*Department of Fish & Wildlife Sciences, University of Idaho, Moscow, ID 83844-1136, USA*
[2]*Centre for Exploration Targeting, School of the Earth Sciences, The University of Western*
Australia, Crawley, WA 6009, Australia

Abstract

1) The ecology of endangered and rare species can be difficult to study due to their low
abundances and legal limits on scientist's ability to catch, sample, and track them. This is
particularly true of sawfish (family Pristidae) whose numbers have declined
precipitously, placing all five species on the IUCN Red List of Threatened Species
worldwide. Best known for their distinctive, toothed rostrum, the ecology, movement,
and life-history of sawfish is poorly understood.
2) Sawfish rostral teeth are modified placoid scales, which grow continuously throughout
the life of the fish. This continuous growth, combined with their stable calcified makeup,
makes sawfish rostral teeth a potential source of temporal records of chemical and
isotopic changes through the life of the fish.
3) Rostral teeth are often preserved in museums and as curios, potentially providing a
source of life-history data to inform conservation actions without the need for field
study, or as an important compliment to it. This is the first study to recover temporally
explicit chemical data from sawfish rostral teeth.
4) Using archived samples of largetooth sawfish (*Pristis pristis*) we show that multiple
chemical tracers can be recovered from sawfish rostral teeth, and that these tracers can
be used to understand movement across salinity gradients. We further show that
sawfish rostral teeth contain repeated structures and indistinct banding which could
potentially be used for aging or growth analysis of fish.

Keywords: Pristis pristis, largetooth sawfish, elemental chemistry, strontium isotopes,
geochemistry

1. Introduction

Understanding the ecology of endangered and cryptic species can be challenging. Low abundance, and the ethical limitations on human interaction for highly endangered species, limit our ability to apply traditional techniques to understand the ecology of these species at appropriate temporal and spatial scales (Cuthill, 1991; Mackenzie et al., 2005; Chadès et al., 2008; Parris et al., 2010). However, effective conservation of these species often requires detailed understanding of their life-history, migration patterns, and ontogeny. This leaves ecologists to find creative ways to uncover this important information.

The largetooth sawfish (*Pristis pristis*), a large ray with an iconic toothed rostrum, is a critically endangered species with extremely low abundances whose life-history is poorly understood (Burgess, Fernandez-Carvalho & Imhoff, 2009; Kyne, Carlson & Smith, 2013; Fernandez-Carvalho et al., 2014). Along with habitat loss and black market trade in fins and rostra, a major threat to sawfish is entanglement in nets as bycatch for other species (Seitz & Poulakis, 2006; Dulvy et al., 2016). All species of sawfish move across a range of salinities from salty to brackish, and largetooth sawfish, are found in salty and fresh water (Thorson, 1982; Poulakis et al., 2011; Scharer et al., 2012; Faria et al., 2013).

Behavior of Australian largetooth sawfish populations in relation to salinity are relatively better understood (Peverell, 2005; Thorburn et al., 2007; Whitty et al., 2009; Gleiss et al., 2017). However, regional variation in freshwater use in largetooth sawfish is as evidenced by historical populations in Lake Nicaragua and reports as far upstream as Manaus in the Amazon river indicating extensive freshwater use in some populations (Thorson, 1982; Burgess, Fernandez-Carvalho & Imhoff, 2009). Despite this, the ecology and distribution of largetooth sawfish in Central and South America is not well studied and many published accounts are limited to opportunistic sightings (Nunes et al., 2016; Feitosa, Martins & Nunes, 2017; Schmid & Giarrizzo, 2017). However, understanding the distribution and ecology of sawfish, especially in their remaining stronghold in the Amazon region, is a necessary precursor to conservation actions beyond the legal limits on catch and trade in sawfish parts (Kyne, Carlson & Smith, 2013; Dulvy et al., 2016). Uncovering patterns of habitat use and migration, and their relationship to net fishing which is a large source of mortality, could help target conservation and management actions to life-stages, locations, or seasons where the risk to sawfish is greatest. Further, understanding the locations of nursery or rearing areas in fresh and brackish water could promote conservation of these locations. However, low population densities and large expanses of potential habitat make collecting movement data using capture or tracking technologies particularly difficult.

Sawfish conservation across a large portion of their range is limited by a lack of knowledge related to their presence or absence, and the location of critical habitats (Dulvy et al., 2016). This lack of specific spatial knowledge, and the reality of extremely low catch rates, is reflected in recent innovative use of indirect abundance-occupancy models to assess the likelihood of sawfish abundance and occupancy worldwide (Yan et al., 2021). The presence of sawfish in much of Latin America, Africa, and Asia remains listed as "uncertain" or "possibly extinct" for multiple species (Dulvy et al., 2016). The range of the most widely distributed species, the largetooth sawfish, has primarily been defined by historical accounts across much of their range, with the vast majority of our detailed knowledge coming from Lake Nicaragua

and Northern Australia, and even stongholds in the Amazon region have not been
systematically surveyed (Kyne, Carlson & Smith, 2013; Kyne et al., 2021). Further, the
opportunity to collect new information in many areas is considered low, due to this decrease in
population abundance (Kyne et al., 2021).
Where detailed presence or absence data is missing, conservation plans are often built
based upon community knowledge of sawfish catches, with opportunistic collection of size and
catch data where samples are available (Leeney & Poncelet, 2015; Jabado et al., 2017; Mendoza
et al., 2017; Haque, Leeney & Biswas, 2020; López-Angarita et al., 2021). While historical data
may not reflect the current state of the population, it increasingly plays a role in planning and
management within marine science, as conservation often requires an understanding of both
historical range and current information to define conservation goals (Máñez et al., 2014).
However, community-based approaches and opportunistic catch data is often of low-
resolution. Further, designation of important habitat is often done without the opportunity for
thorough assessment of local behavior and ecology, as evidenced by habitat designation of the
smalltooth sawfish in the United States which is only now being supplemented by direct studies
of movement and location using telemetry (Wiley & Simpfendorfer, 2010; Norton et al., 2012;
Graham et al., 2020), and presence/absence studies using environmental DNA (eDNA)
techniques (Simpfendorfer et al., 2016; Lehman et al., 2020).
In areas where funding is available for systematic and large-scale population research,
telemetry and eDNA data is extremely useful. However, both methods require significant
funding and investment, funding which is often unavailable in much of the sawfishes range.
Further, these methods can only reveal the movements and location of a severely depleted
species which has lost significant range, and presumably similarly significant levels of life-
history diversity.
The importance of habitat and life-history diversity on the resilience of species, and the
importance of identifying and prioritizing life-history diversity within conservation plans is well
understood (Beechie et al., 2006; Schindler et al., 2010; Schroeder et al., 2016). New tools
which can increase the data richness of historical samples are needed, and attempts are
currently underway to use archived rostral samples to understand historical genetic diversity
and population structure of sawfish worldwide (Phillips, Nicole., Wueringer, 2015; Fearing et al.,
2018). However, no tools currently exist which can elucidate life history in these archived
rostral samples, or tie historical genetic information gleaned from them to the movement and
behavioral aspects of sawfish ecology. Further, few opportunities exist to determine adult life-
history of sawfish, one of the least understood areas of sawfish ecology (Kyne et al., 2021), yet
many archived rostrum represent samples of large adult fish.
One tool that has been revolutionary in understanding movement and ecology of
teleosts (and increasingly chondrichthyans) is the use of chemistry stored in fish hard parts to
determine provenance, migration, and other aspects of life history (Kennedy et al., 1997;
Hobson, Barnett-Johnson & Cerling, 2010; Tillett et al., 2011; Smith, Miller & Heppell, 2013;
Walther, 2019; Wang, Walther & Gillanders, 2019). These isotopic and trace elemental
techniques are particularly powerful when applied to calcareous structures which grow
throughout the life of the fish, as other Group 2 elements readily substitute for calcium in the
matrix, and other elements are incorporated in the crystalline structure (Elsdon et al., 2008).

Strontium isotope composition, $^{87}Sr/^{86}Sr$, has been shown to be recorded without
fractionation in fish otoliths (Kennedy et al., 1997, 2000). This isotope ratio can vary greatly in
nature and has been used to trace fish movement with kilometer precision in fresh water, and
the presence of a global marine signature worldwide can be used as a marker of entry into salt
water (Hamann & Kennedy, 2012; Courter et al., 2013; Hegg et al., 2013).
The ratio of elemental Sr and Ba to calcium (Sr/Ca and Ba/Ca) in hard parts have been
shown to be useful tracers of salinity changes due to generally higher bulk strontium in the
ocean than fresh water, and the flocculation of barium in the presence of salinity (Kalish, 1990;
Zimmerman, 2005; Shippentower, Schreck & Heppell, 2011; Hamer et al., 2015). This has been
used to understand entry of anadromous fish into salt water, as well as habitat use in estuarian
and euryhaline species, and to distinguish provenance of fully marine coastal species as well
(Kraus & Secor, 2004; Warner et al., 2005; Mohan et al., 2012; Schaffler, Miller & Jones, 2014;
Rohtla & Vetemaa, 2016; Seeley & Walther, 2017). As well, trace element ratios of Mg (Mg/Ca)
have been strongly related to growth and metabolism with potentially higher influx into otolith
endolymph through ion channels during higher metabolic activity (Limburg et al., 2018).
Manganese is rarely found in the water column in an ionic form suitable for incorporation, and
is released only in or near reducing, anoxic substrates (Altenritter, Cohuo & Walther, 2018;
Limburg & Casini, 2018). Therefore, Mg/Ca has been linked to exposure to reducing
environments such as mangroves, and exposure to hypoxic stress in several species has been
well described (Mohan et al., 2014; Paillon et al., 2014; Turner, Limburg & Palkovacs, 2015;
Smith et al., 2016; Altenritter, Cohuo & Walther, 2018; Limburg & Casini, 2018). Further,
numerous studies in anadromous, catadromous, coastal and estuarine species utilize a suite of
these elements and isotopes to determine provenance. These include tracing provenance and
of reef fish to natal seagrass beds, natal origin of American shad across the Atlantic seaboard,
nursery areas of menhaden in Chesapeake Bay, entry of juvenile salmon into the ocean, and
facultative oligohaline habitat use in Atlantic tarpon (Chittaro et al., 2006; Walther & Thorrold,
2008; Hegg, Kennedy & Fremier, 2013; Schaffler, Miller & Jones, 2014; Woodcock & Walther,
2014).
Until recently the use of chemical signatures in sharks and rays has lagged behind their
use in teleost species and has been confined mainly to elemental ratios due to the utility of
elemental ratios in fully marine species. However, recent studies have shown that
elasmobranch vertebra function as temporal records of elemental chemistry and location
(Tillett et al., 2011; Smith, Miller & Heppell, 2013; Raoult et al., 2016). Numerous studies have
now been published using suites of elemental markers to determine location and stock
structure in elasmobranchs (Izzo et al., 2016; McMillan et al., 2017; Mohan et al., 2018;
Coiraton, Amezcua & Ketchum, 2020). Further work utilized strontium and barium elemental
ratios to recover movements across salinity gradients in several species of shark, including
repeated spawning movements into fresh water for bull shark, and likely proximity to mangrove
areas based on manganese concentrations (Tillett et al., 2011; Lewis et al., 2016; Feitosa,
Dressler & Lessa, 2020). Recent work has shown a similar relationship of salinity to Sr/Ca in
smalltooth sawfish vertebra as that shown in other elasmobranchs and teleosts (Scharer et al.,
2012).
The incorporation of these elemental signatures into elasmobranch hard parts appears
to function in similar ways to teleosts, though fewer controlled studies have been done. Two

controlled tank and mesocosm experiments have shown Sr/Ca, Ba/Ca, Mg/Ca, and Mn/Ca to be
similarly useful as natural tags in shark vertebra along with other elemental ratios (Tillett et al.,
2011; Smith, Miller & Heppell, 2013). These studies indicate that strontium and barium are
taken up in proportion to the element's concentration in the environment, while elements such
as Mg and Mn are more highly regulated at the cellular level and therefore reflect growth and
physiological conditions to a greater degree. The effect of temperature on Sr and Ba
incorporation in vertebra is conflicting, as it is in the teleost literature, however the magnitude
of the temperature effect is small (Tillett et al., 2011). Turnover rate of these signatures within
the elasmobranch body is unknown and is likely mediated by the blood supply and speed of
growth in the tissue being examined, as opposed to teleost otoliths which lack blood supply but
show changes at a scale of weeks (Campana, 1999; Hegg, Kennedy & Chittaro, 2018).
We propose that sawfish rostral teeth may provide a useful record of age, growth and
environmental chemistry which could hold information on life-history and movement in
individual fish. Rostral teeth are not true teeth but instead highly modified dermal denticles
which grow through the life of the fish (Welten et al., 2015). Made of bio-apatite (Miller, 1974;
Shellis & Berkovitz, 1980), rostral teeth grow continuously from their base and are not replaced
if they are lost (Slaughter & Springer, 1968; Wueringer, Squire & Collin, 2009). Rostral teeth
presumably incorporate similar trace-elements and isotopes in their chemical matrix similar to
other calcium carbonate and calcium phosphate hard parts such as shells, true teeth, and
otoliths (Campana & Thorrold, 2001; Schöne et al., 2004; Begg et al., 2005; Thornton, 2011).
While the morphology of sawfish teeth varies across species, their basic formation and growth
is similar. These characteristics indicate that sawfish rostral teeth may store useful ecological
data as an endogenous record.
Importantly, sawfish rostra are often dried and kept as curios and many are kept in
museums, academic and private collections worldwide (Seitz & Poulakis, 2006; Melo Palmeira
et al., 2013). These collections provide a reservoir of samples by which to uncover details of
sawfish life-history without harming living individuals (Phillips et al., 2009; Whitty et al., 2014).
While work from archived collections are necessarily reconstructions of the past, archived
samples also potentially reflect records of behavior and life history before and during the
ongoing declines in abundance of these species. Because of this, they may hold important
information that is no longer evident in extant populations. Further, there is additional
potential to utilize natural mortalities and samples seized by law enforcement to infer limited
information about current populations.
Only one unpublished study has investigated the use of sawfish rostral teeth as an
endogenous record of growth and provenance (Field, Meekan & Bradshaw, 2009). This work
indicated that the trace element chemistry of whole sawfish rostral teeth could be used to
distinguish sawfish populations around the coast of Australia. The authors also demonstrated
that growth banding could be observed on the outside of whole rostral teeth which could be
used to estimate age. Despite this, no work has been published investigating whether rostral
teeth record higher-resolution chemical or structural data throughout the life of the fish.
However, Sawfish vertebra, much like those of sharks and other rays, have been shown to
contain yearly growth bands as well as a chemical record of salinity (Scharer et al., 2012).
In this study we present data from the analysis of rostral teeth taken from two
largetooth sawfish rostra housed in a university collection in the United States. Our first

objective was to demonstrate methods of sectioning, treating and imaging sawfish rostral teeth
to quantify age and growth information they contain. Our second objective was to document
the potential of sawfish rostral teeth as a chemical record of location and movement using laser
ablation-inductively coupled plasma mass spectrometry (LA-ICPMS) of $^{87}Sr/^{86}Sr$ isotope ratio, as
well as trace element ratios of Sr/Ca, Ba/Ca, Mn/Ca, Mg/Ca. Finally, we analyzed these chemical
and growth data to investigate their utility in understanding lifetime sawfish movement across
salinity and the existence of other environmental or ontogenetic information recorded during
the life of the fish.
222
223 ## 2. Methods
224
225 *Sample Information*
226
All data in this study were taken from two sawfish rostra included in the ichthyology
teaching collection at University of Idaho in Moscow, ID USA (Figure 1). The first was the
rostrum of a largetooth sawfish embryo. Information attached to the rostrum in the collection
read, "Mother caught off Belize (British Honduras) coast, 1969. 8ft long with 5 embryos ≈ 1.5 ft
long. Blade covered with protective sheath when embryo was removed from mother." Thus,
the total length (TL) of the embryo was approximately 45 cm. This rostrum has a total rostral
length (TRL, length from tip of the rostrum to the attachment to the head where the rostrum
begins to flare *sensu* (Whitty et al., 2014)) of 14.2 cm assuming the rostrum was removed at
this attachment point, and a standard rostral length (SRL, length from tip of rostrum to the
most distal tooth) of 13.6 cm (Figure 1A)
The second sample is a larger rostrum from a largetooth sawfish with a TRL of 57 cm
and a standard rostral length of 55 cm (Figure 1B). No information was available for the
provenance of this rostrum. Using the mean SRL to total length (SRL/TL) relationship from
Whitty et al. (2014) results in a TL of 238.7 cm for the fish from which the rostrum was taken.
This length would indicate the fish was a sub-adult as 240 cm is the smallest published length of
maturity for males, with 280 cm for males and 300 cm for females being more widely accepted
(Wueringer, Squire & Collin, 2009; Dulvy et al., 2016).
244
245 *Structural Analysis*
246
The most distal tooth on the left-hand side of each rostrum was removed using a utility
knife and a Dremel tool. The embryo tooth showed no signs of tooth wear. The sub-adult tooth
was highly worn on the anterior edge of the tooth extending beyond the rostrum, and
significantly pitted and eroded on the posterior edge. Rostral teeth were cleaned of tissue by
soaking in water followed by scraping with a scalpel blade. Prior to sectioning, the surface of
each tooth was stained using silver nitrate to highlight growth ridges as described by Field et al.
(2009). The structure of sawfish teeth is extremely dense, and as noted by Bradford (1957) is
difficult even to dissolve using nitric acid. Therefore, this surface treatement is unlikely to affect
further chemical analysis within the sectioned tooth.

256	Rostral teeth were then embedded in epoxy resin and sectioned using an Isomet low
257	speed saw and a diamond wafering blade (Beuhler, beuhler.com) in a manner described for
258	large otoliths and fin rays (Secor, Dean & Laban, 1992; Koch & Quist, 2007). The sub-adult tooth
259	was curved both distally and along the dorsal-ventral axis. To allow sectioning through the
260	center of the entire tooth, given the complex curvature, it was cut in half near the point at
261	which it protruded from the rostrum. The embryo tooth and the two pieces of the sub-adult
262	tooth were embedded in epoxy and an approximately 1mm section in the dorsal-ventral plane
263	was taken that included the center core of each tooth (Figure 2). A further two sections were
264	taken in the same plane from the sub-adult tooth, and a second section was taken from the
265	embryo tooth to facilitate staining while maintaining one unstained section for chemical
266	analysis. Each section was polished on 3600 grit and 6000 grit Micro Mesh wet-dry sandpaper
267	(Micro Mesh, micromesh.com).
268	The sub-adult tooth sections were stained using both silver nitrate and Mutvei's solution
269	(Dunca, Fiebig & Pfeiffer, 2005) to enhance any growth banding present in the rostral teeth.
270	Sections were submerged in silver nitrate for approximately 5 minutes prior to light exposure.
271	The second section was submerged in Mutvei's solution (dilute acetic acid, gluteraldahyde, and
272	alcian blue) for 6 hours. The remaining embedded sample was burned using a handheld butane
273	micro-torch (BernzOmatic, bernzomatic.com) until the sample surface was golden brown. The
274	sample was then polished using the same sequence of wet-dry sandpaper to enhance the rings.
275	The embryo tooth was not stained for growth analysis as the small size of the tooth caused the
276	second thin section to be non-representative of the core of the tooth and very little internal
277	structure was present to indicate growth analysis would be fruitful. Therefore, we opted to
278	preserve the best thin section for chemical analysis.
279	Each section was imaged in sections at 10X magnification using a Leica DFC450 digital
280	camera (leicacamerausa.com) attached to a Zeiss AxioImager A.1 microscope (zeiss.com). These
281	images were subsequently stitched into a larger mosaic using the Photomerge tool in Adobe
282	Photoshop (adobe.com) with the "collage" layout option and "blend images together" option
283	checked. These color images were then transformed into grayscale and the color palette was
284	inverted to determine an imaging method which would best highlight any banding or sequential
285	growth structures that were present. Each image was subsequently optimized for contrast
286	using a levels adjustment.
287	The number of bands and/or sequential structures inside the sub-adult tooth were
288	counted by two readers using ImageJ analysis software (https://imagej.nih.gov). Points and
289	distances between them were measured along a line adjacent to the proximal edge of the
290	tooth, on the outside of the dentin layer and inside the enamel layer where it was still present.
291	Distances were standardized to the length of the laser scan for comparison with the chemical
292	data.
293	
294	*Trace-element and Isotopic Analysis*
295	
296	The unstained sections of the tip and base of the sub-adult rostral tooth, as well as the
297	embryo, were analyzed for strontium composition ($^{87}Sr/^{86}Sr$) as well as concentration of a suite
298	of trace elements (^{86}Sr, ^{138}Ba, ^{55}Mn, ^{25}Mg). The structure of sawfish teeth is dentin, primarily
299	composed of calcium phosphate (Miller, 1974; Shellis & Berkovitz, 1980). Thus, calcium was

used as the internal standard to correct for variation in laser ablation efficiency across the
sample and all trace element data is expressed as a ratio to calcium (Sr/Ca, Ba/Ca, Mn/Ca,
Mg/Ca).
Analysis was conducted at the Radiogenic Isotope and Geochronology Laboratory (RIGL)
at Washington State University. Strontium ratio and trace element data were collected
simultaneously using a laser ablation split stream (LASS) methodology (Hegg & Fisher, 2020).
Laser ablation was conducted using a Teledyne Analyte Excite 193nm ArF laser using a 35μm
spot diameter, 10μm/sec rastering speed, operating at 20Hz, and with laser fluence of ~4J/cm^2.
Ablated material was then split between a NeptunePlus multicollector inductively coupled mass
spectrometer (MC-ICPMS) for ^{87}Sr/^{86}Sr measurement and Element2 ICPMS for trace element
measurement. The laser was used to sample along the center of the sampled tooth,
perpendicular to the growth axis, from the tip to the base of the tooth.
Trace element concentrations were normalized to the NIST-610 glass standard, while
^{87}Sr/^{86}Sr was corrected for K and Rb interferences and normalized to a modern marine shell and
a ^{87}Sr/^{86}Sr of 0.70918 for modern ocean water, as described by (Hegg & Fisher, 2020). Trace
element analysis was calibrated using the NIST 610 standard as described in (Hegg & Fisher,
2020).
All data reduction was completed using a custom built data reduction reduction scheme
(DRS) operating within the Iolite software package (Fisher et al., 2017; Hegg & Fisher, 2020). All
plotting and analyses were completed in R statistical software (https://www.r-project.org).
320
321

3. Results

323
324   *Structural Analysis*
325
Whole rostral teeth, both unstained and after silver nitrate staining, did not exhibit the
external growth banding described by Field et al. (2009). The interior of both rostral teeth
contained an inner, clear layer and an outer opaque layer in areas which had not been
subjected to wear (Figure 2A). The rostral teeth contain a mesh-like network of interconnected
tubules as described by Miller (Miller, 1974). These tubules followed a general pattern of
intertwining tubes near the central core, with downward "legs" curving distally toward the edge
of the tooth (Figure 2A&D). These descending legs appear to be spaced somewhat regularly,
suggesting that they could mark growth or age.
Burning of the sample showed some evidence of layering within the tooth but it was too
indistinct to be easily quantified. Silver nitrate staining of the tooth sections did not reveal
additional layering or banding structure. Mutvei's solution was moderately successful in
bringing out faint alternating banding in the middle portion of the sub-adult tooth (Figure 2B,
D&E). The alcian blue stain in Mutvei's solution infiltrated the tubules which broke the surface
of each tooth section, making clear which tubules were on the surface and which were below.
Both readers agreed that the inverted image of the section stained with Mutvei's
solution provided the best image for measurement (Figure 2C). This image rendered the surface
tubules in a distinct orange which was easier to discern than the blue of the non-inverted

image. Quantifying of the apparent banding created by the tubule legs and indistinct light-dark
bands was done by both readers. Both readers picked 26 growth bands. The first 15 of these
bands, starting from the youngest growth at the tooth tip, matched between the readers
(Figure 2D). In the basal section of the tooth, where the tubules were most indistinct, 6 of the
remaining 11 points matched between the readers (Figure 2E). A consensus measurement was
then created which contained 29 bands. When plotted the distance between measurements
displayed a linear relationship.
350
351 *Trace-element and Isotopic Analysis*
352
Laser ablation and ICP-MS analysis was successful, with 35µm spot size allowing useful
resolution across the sample (Figure 3). The marine shell standard measured 0.70915 ($\pm$0.00038
2SE, global marine signature 0.70918). Trace element measurements exceeded the limits of
detection (LOD) by at least three orders of magnitude in all cases (Average LOD across
standards and samples: Sr=0.24 ppm, Ba=0.07 ppm, Mn=0.36, ppm, Mg=0.54 ppm). All analyses
exhibited repeated, low level, spikes in signal. While tubules at the base of the rostral tooth
created voids which unavoidably resulted in large changes to the signal, the smaller, repeated
spikes appeared not to correlate with voids in the surface of the tooth or with changes in the
signal and were therefore assumed to be environmental.
Strontium isotope ratio (Figure 3A) exhibited a convex pattern in the sub-adult tooth,
starting below 0.708 in the earliest growth periods, before increasing in the middle of the fish's
life toward the global marine signature (0.70918), then gradually decreasing to near 0.707 at
the base of the tooth. In the embryo, ^{87}Sr/^{86}Sr was very steady and near the global marine
signature of .70918 (Figure 3A).
The Sr/Ca was relatively steady in the sub-adult fish, with slightly higher values in the
earliest period of growth and at mid-life (Figure 3B). In comparison the Sr/Ca was significantly
elevated in the embryo, exhibiting lower values at the start of growth, increasing to their
highest values between 5 and 10 mm from the tip, and decreasing slightly at the tooth base
(Figure 3B).
In the sub-adult tooth, Ba/Ca exhibited the highest values at the earliest growth stages,
decreasing at mid-life, and increasing again toward the end of life at the tooth base (Figure 3C.
The embryo tooth contained very low levels of barium throughout, contrasting with Sr/Ca
which showed the opposite pattern as would be expected in the case of movement across
salinity (Figure 3C).
Magnesium to calcium ratios showed pronounced, repeated peaks throughout the life
of both fish, with large decreases near the tooth base (Figure 3D). Manganese to calcium ratios
showed similar repeated peaks throughout the life of the fish (Figure 3E). The highest Mn/Ca
values were recorded at the tip and base of each tooth. Extreme spikes in Mn/Ca near the base
of the tooth were likely caused by voids in the base of the tooth, and the extreme spike at
~12mm is likely related to contamination rather than a biological process. Repeated peaks,
however, are assumed to be related to fish movement or metabolism.
384
385

4. Discussion

The highly endangered status of sawfishes presents challenges to researchers and conservationists (Simpfendorfer, 2000; Kyne, Carlson & Smith, 2013; Dulvy et al., 2016). Modern, long-term, studies of living sawfish in the wild are rare and are often skewed toward smaller, juvenile fish because they are most likely to be captured (Thorson, 1982; Peverell, 2008; Poulakis et al., 2011). Further, conservation status and abundance preclude most lethal sampling of sawfish, largely limiting the promise of vertebra as an aging and microchemical tool for understanding sawfish life history and provenance, as sawfish vertebra are rarely preserved (Whitty et al., 2014). Developing methods which can address questions of life-history and location with larger sample sizes, especially of adult fish, are required to understand sawfish ecology and inform conservation efforts of sawfish. Because sawfish rostra, as opposed to vertebra, have value and are often retained, there is a larger store of historical samples available for analysis (Whitty et al., 2014; Fearing et al., 2018).

If sawfish rostral teeth do indeed contain chemical records of movement and location, as shown in this study, then archived samples may allow larger sample sizes and increased investigation of life-history questions. In particular, very little is known about the ecology and life-history of large adult sawfish due to their rarity. This has been identified as a problem in understanding basic questions of sawfish growth and maturation, and also affects our understanding of their adult movement and migration patterns (Kyne et al., 2021). Current knowledge of the locations of sawfish populations is limited in many areas, with particularly little detailed understanding of nursery areas outside of a few heavily studied nursery hotspots in Australia (largetooth sawfish, green sawfish) and Florida (smalltooth sawfish) (Poulakis et al., 2011; Feldheim et al., 2017; Huston et al., 2017; Scharer et al., 2017). Feeding areas, or movement corridors for adult sawfish of all species are less well studied (Graham et al., 2020).

These questions of habitat use and movement are critical to sawfish conservation. While sawfish are legally protected in many locations, more granular understanding of their distribution and ecology is required to provide the site and area protection and resource and habitat protection noted as a needed conservation action by the IUCN (Kyne, Carlson & Smith, 2013). If provenance and life history diversity can be investigated through the use of rostral tooth chemistry, as in teleosts and other chondricthyans, opportunities for identifying provenance and habitat use with more specificity could provide more spatially explicit ecological information to guide conservation. While the catch location of a preserved rostrum provides a single data point on species presence, analysis of rostral tooth structure and chemistry could provide temporally explicit chemical information useful for identifying nearby nursery, rearing, and feeding areas, as well as lifetime movement patterns. One of the most clear-cut examples of movement data available from calcified hard parts is movement across salinity, due to the unique dynamics of strontium and barium in relation to salinity (Zimmerman, 2005; Scharer et al., 2012; Hamer et al., 2015).

Movements across salinity can be determined using a combination of ^{87}Sr/^{86}Sr, Sr/Ca, and Ba/Ca, whose dynamics in teleosts and elasmobranchs appear to be similar (Tillett et al., 2011; Smith, Miller & Heppell, 2013). Strontium isotope composition is well mixed in ocean water, such that all ocean water exhibits a signature of 0.70918 (Faure & Mensing, 2004). Deviations from this ratio indicate movement into fresh or brackish water. Further, barium

concentrations quickly decrease as salinity increases in estuaries, resulting in commensurate
decreases in Ba/Ca in hard structures, signaling the transition from fresh to brackish water
(Kalish, 1990; Walther & Limburg, 2012). Strontium concentrations, and Sr/Ca in hard parts,
exhibit the opposite dynamic, increasing quickly as brackish water transitions to the higher
salinity of the ocean (Zimmerman, 2005).
The sub-adult sample in this study shows likely movement across fresh and brackish
water throughout its life. High Ba/Ca, low Sr/Ca, and ^{87}Sr/^{86}Sr far below 0.70918 indicate the
earliest recorded period of the sub-adult fish was spent in low salinity, likely fresh water (Figure
3A, B&C). During the middle portion of its life (15-35mm) Ba/Ca decreases and ^{87}Sr/^{86}Sr
increases toward the global marine signature which indicates movement toward more saline
water. The lack of marked increase in Sr/Ca and the fact that ^{87}Sr/^{86}Sr stays slightly below the
global marine signature indicates the fish was in saline water, but not in fully saline water. By
the end of its life the fish appears to have moved back into fresh or mildly saline water. The
smaller spikes and troughs in these signals may indicate smaller-scale forays into different
salinities, either through movement or through seasonal changes in salinity as reported by
Sharer et al. (2012).
This result is consistent with the known life-history of largetooth sawfish, which inhabit
shallow, brackish, nearshore water as well as fresh water. Further, largetooth sawfish juveniles
are known to utilize freshwater habitats as nursery areas (Wueringer, Squire & Collin, 2009).
These chemical results indicate an expected pattern of freshwater utilization in early life,
followed by use of multiple salinities later in life.
In contrast, the embryo chemistry indicates provenance in a much higher salinity
environment. Throughout its length the embryo rostral tooth exhibits high but somewhat
variable Sr/Ca, extremely low Ba/Ca, and a stable ^{87}Sr/^{86}Sr signature very near the global
marine signature. Together, these indicate that the embryo spent the entirety of the gestation
in a saline environment. The variation in Sr/Ca indicates some change of salinity, but the lack of
variation in Ba/Ca and ^{87}Sr/^{86}Sr indicates that this change was likely not into significantly lower
salinity.
Of course, the environment recorded by the embryo tooth is mediated by the internal
chemistry of the mother, which presents some potential complexity in interpretation. However,
it has been shown in multiple species that Sr/Ca, Ba/Ca are deposited in relation to the
surrounding concentration (Bath et al., 2000; Campana & Thorrold, 2001; Smith, Miller &
Heppell, 2013). Similarly, ^{87}Sr/^{86}Sr has been shown to deposit in hard parts without biological
fractionation in the body across species (Kennedy et al., 1997, 2000; Flockhart et al., 2015).
Thus, it is likely that the mother inhabited saline water during some amount of the early natal
period of the embryo.
To pass these signatures on to the embryo the mother's body must have been
equilibrated to a saline environment at least until formation of the egg. As seen in salmon,
provisioning by the yolk sac maintains the progeny chemistry at that of the maternal signature
with little change, due to relatively low ion exchange with the outside environment (Kalish,
1990; Hegg, Kennedy & Chittaro, 2018). The same may be true of the developing sawfish whose
provisioning is largely from the yolk. The rate at which the mother's chemistry would be
reflected in the embryo as it equilibrates to entry into fresh water would likely depend on the
degree of ion exchange with the embryo, the speed of change within the blood, and the

turnover time of the body in its entirety. None of these parameters are known in detail, though
plots in Smith et al. (2013) show equilibration in Ba/Ca occurring at rates consistent with a
weekly equilibration period in *Urobatis halleri*.
The Mg/Ca ratio has been called the 'chemical clock' in some fish species due to its
association with seasonal growth rates, and one controlled elasmobranch study has found
limited evidence that Mg incorporation increases with condition (Limburg et al., 2018; Pistevos
et al., 2019). Without additional information about sawfish in particular it is impossible to
determine, but the repeated sharp peaks in this tracer are suggestive of periods of higher and
lower growth (figure 3D), a relationship which merits further study. The marked decrease in
Mg/Ca near the base of the tooth (30-45mm) coincides with the onset of cloudiness in the
tooth matrix. This pattern also occurs in the embryo tooth, with a markedly similar shape but
scaled to the size of the smaller embryo tooth. This indicates that the change in Mg chemistry
may be related to some unknown process of tooth formation occurring near the root, rather
than a tracer of changes in the environment.
In teleosts, otolith Mn/Ca has been shown to loosely follow concentrations in the water,
however endogenous processes related to growth further mediate it's deposition (Mohan et
al., 2012; Turner, Limburg & Palkovacs, 2015). Manganese in the water column is most closely
associated with reduction of Mn oxides by microbes on the substrate under anerobic conditions
and is otherwise rarely found in solution, however anthropogenic inputs from sewers have
been documented as well as proximity to mangroves, which is a known habitat of largetooth
sawfish (Thorrold & Shuttleworth, 2000; Mohan et al., 2012; Paillon et al., 2014; Smith et al.,
2016). However the kinetics of dissolved Mn are such that it is unlikely to remain dissolved for
long periods before being oxidized, and therefore increased levels in rostral teeth likely indicate
close proximity to reducing environments (Laslett, 1995).
Sawfish are known to inhabit shallow, nearshore habitats and mangrove swamps where
anerobic decomposition occurs, and increases in Mn/Ca could indicate forays into these
environments, while faster growth during these periods would result in further apparent
increases. These increases do not necessarily indicate the fish are making forays through
hypoxic water, but that they may be close enough to reducing environments to incorporate
dissolved Mn. It is interesting to note, however, that the peaks in Mn/Ca exhibited in both the
sub-adult and embryo rostral teeth (Figure 3D) rise above the threshold set by Limburg and
Casini (2018) for hypoxia exposure in teleosts, and multiple studies have shown the effects of
hypoxic conditions recorded in otoliths (Altenritter, Cohuo & Walther, 2018; Limburg & Casini,
2018). However, it is unlikely that the embryo, whose gestation likely occurred in ocean water
according to its Sr/Ca and Ba/Ca signatures, would have been exposed to hypoxia, indicating
that incorporation of Mn into rostral teeth may differ from that of otoliths.
Analysis of banding structures in the sub-adult tooth showed that the distance between
the putative growth bands formed a linear relationship rather than characteristic decrease in
distance with age expected from a von Bertalanffy growth function. There are two possible
reasons for this. The first is that these structures form at regular intervals along the tooth and
are not influenced by growth. The second is related to tooth wear. Sawfish rostral teeth are
often heavily worn, as the sub-adult sample in our study was. The amount of tooth that has
worn away is difficult to know, but it is possible that the only remaining growth record in our
sample is the relatively linear, sub-adult portion of the growth curve. The furthest distal rostral

teeth in our rostrum were also some of the most worn, indicating that future studies should
target rostral teeth with less wear and curvature, elsewhere on the rostrum.
While the length-at-age relationships of sawfish are poorly understood for larger adults,
comparisons of our data to existing growth curves suggest that the putative growth bands are
sub-annual. Based on the SRL/TL relationship reported by Whitty et al. (2014), the sub-adult
sample was taken from a 238 cm fish. Simpfendorfer (2000) predicts the sub-adult would have
been ~5 years old, or ~12 years old based on an alternative growth curve assuming a halved
growth rate. More recent growth curves from Peverell (2008) all predict ages less than 6 years
old. Thus, if the structures observed are related to age or growth, the relationship would need
to be established through a relationship to other age information such as counts of annual
vertebral bands.
Taken together this study presents compelling evidence that largetooth sawfish rostral
teeth are a recorder of chemical information throughout the life of individual fish. Further, this
chemical data records long-term, high-resolution, information about sawfish life-history and
movement that is otherwise difficult to obtain from a live specimen. Our data further suggest
that the structures in sectioned rostral teeth are periodic in nature and may be useful in
determining growth, though they do not appear to be annual in nature.
This record of chemistry and movement may provide a useful tool in the conservation of
largetooth sawfish, and likely other sawfish species. Hard-part chemistry in other fish species
demonstrates the power of these chemical techniques to uncover important information about
provenance, movement, and life-history which has been instrumental in conservation planning
(Walther, 2019). When combined with isoscape techniques and other abiotic spatial data, hard-
part chemistry is capable of uncovering important habitat and behavior with great specificity
(Hobson, Barnett-Johnson & Cerling, 2010; Hamann et al., 2014; Smith et al., 2016; Brennan &
Schindler, 2017). A reservoir of historical samples on which to apply these techniques exist
because rostra are often saved (Whitty et al., 2014). These archived samples potentially contain
information such as population-specific abiotic affinities, movement patterns, nursery hotspots,
and adult movement and life-history. Combining chemical information with ongoing genetic
efforts using archived rostra could also be synergistic, providing even greater detail in
population connectivity and movement (Fearing et al., 2018). While these samples are
necessarily historical, the information they contain is likely to be applicable to current
populations, and in many cases where populations have declined precipitously may be the only
information available. Further, currently a saved rostrum is used as often only a single data
point recording presence of a species, often with poor spatial resolution. Rostral tooth
chemistry could expand the data available from a single archived rostrum to include temporally
explicit movement data, as well as population specific habitat and life history information that
could support more specific conservation actions.
However, this study points out significant questions which must be answered to validate
the use of sawfish rostral teeth as records of movement, life history, or growth. First, more
must be known about the biology of chemical incorporation into rostral teeth, and the rates at
which the chemistry of a new environment is reflected in rostral teeth. This is an ongoing
question in the use of elasmobranch vertebral chemistry, and understandings from vertebral
studies may provide answers applicable to sawfish rostral teeth as well.

Second, the rate of rostral tooth wear must be understood in order to understand how
far back into the life of a fish the information in a rostral tooth extends. As teeth are worn,
information from the early life of the fish is lost at the tip, while the most current data is
available nearest the base of the tooth. Even without the entire life of the fish intact it is clear
that important movement information is retained. Thus, it may still be possible to identify
important nursery, rearing, or foraging locations based on chemical patterns in the existing
tooth. However, understanding the rate of wear in relation to age could be used to put bounds
on the time period and ages represented in sampled rostral teeth. Comparison of teeth in
captive and wild sawfish, whose rate of tooth wear differs, could be used to examine this
phenomenon. Sawfish rostral teeth that are entirely lost do not appear to regrow (Miller,
1974), but teeth have been reported to fracture to the pulp and return to a normal shape in at
least one non-peer reviewed report (Clippinger, 1993). Methods for identifying these fractured
and regrown teeth, in the same manner that regrown scales are excluded from teleost growth
studies, may be important. The degree to which habitat specific factors increase or decrease
wear may also be an important consideration.
The results reported here are necessarily preliminary given the limitations of our sample
size. While our results are limited to largetooth sawfish, the similarities in morphology and
formation of rostral teeth in other species indicate that our results may be useful for other
species of sawfish as well. Larger samples sizes, including fish of known location and origin
which can be compared to known water chemistries, are necessary to confirm our findings.
Despite these limitations, our data supports further investigation of archived sawfish rostral
teeth as a method of reconstructing movement, life-history, age and growth without the need
to disturb extant fish. The large numbers of rostra housed at museums, academic institutions,
and in private collections worldwide could potentially be employed to study elements of
sawfish ecology which are currently difficult to study in extant populations.
586

Acknowledgements

588
589 This study would not have been possible without the help of C. Caudill who provided access to
590 the samples. Thanks to J. Vervoort and the RIGL lab at Washington State Univeristy for access to
591 analytical equipment. Thanks to P. Charvet and V. Faria for independent species identification.
592 In memory of P. Anders for his editing and unwavering support for this work. The authors have
593 no conflicts of interest to report.

594
595

References Cited

Altenritter, M.E., Cohuo, A. & Walther, B.D. (2018). Proportions of demersal fish exposed to
sublethal hypoxia revealed by otolith chemistry. *Marine Ecology Progress Series*, 589, 193–
208. https://doi.org/10.3354/meps12469
Bath, G.E., Thorrold, S.R., Jones, C.M., Campana, S.E., McLaren, J.W. & Lam, J.W.H. (2000).
Strontium and barium uptake in aragonitic otoliths of marine fish. *Geochimica et*
Cosmochimica Acta, 64(10), 1705–1714. https://doi.org/10.1016/S0016-7037(99)00419-6
Beechie, T., Buhle, E., Ruckelshaus, M., Fullerton, A. & Holsinger, L. (2006). Hydrologic regime

and the conservation of salmon life history diversity. *Biological Conservation*, 130(4), 560–
572. https://doi.org/10.1016/j.biocon.2006.01.019

Begg, G.A., Campana, S.E., Fowler, A.J. & Suthers, I.M. (2005). Otolith research and application:
Current directions in innovation and implementation. *Marine and Freshwater Research*,
56(5), 477–483. https://doi.org/10.1071/MF05111

Bradford, E.W. (1957). The structure of rostral teeth and the rostrum of *pristis microdon*.
Journal of Dental Research, 36(5), 663–668.
https://doi.org/10.1177/00220345570360050401

Brennan, S.R. & Schindler, D.E. (2017). Linking otolith microchemistry and dendritic isoscapes to
map heterogeneous production of fish across river basins: *Ecological Applications*, 27(2),
363–377. https://doi.org/10.1002/eap.1474

Burgess, G.H., Fernandez-Carvalho, J. & Imhoff, J.L. (2009). An Evaluation of the Status of the
Largetooth Sawfish, Pristis perotteti, Based on Historic and Recent Distribution and
Qualitative Observations of Abundance. *Florida Museum of Natural History Report 2009*,
(2009), 1–93.

Campana, S. (1999). Chemistry and composition of fish otoliths: pathways, mechanisms and
applications. *Marine Ecology Progress Series*, 188, 263–297.
https://doi.org/10.3354/meps188263

Campana, S.E. & Thorrold, S.R. (2001). Otoliths, increments, and elements: keys to a
comprehensive understanding of fish populations? *Canadian Journal of Fisheries and
Aquatic Sciences*, 58(1), 30–38. https://doi.org/10.1139/f00-177

Chadès, I., McDonald-Madden, E., McCarthy, M.A., Wintle, B., Linkie, M. & Possingham, H.P.
(2008). When to stop managing or surveying cryptic threatened species. *Proceedings of the
National Academy of Sciences of the United States of America*, 105(37), 13936–13940.
https://doi.org/10.1073/pnas.0805265105

Chittaro, P.M., Hogan, J.D., Gagnon, J., Fryer, B.J. & Sale, P.F. (2006). In situ experiment of
ontogenetic variability in the otolith chemistry of *Stegastes partitus*. *Marine Biology*,
149(5), 1227–1235. https://doi.org/10.1007/s00227-006-0280-x

Clippinger, D.H. (1993). Oservations of Rostral Tooth Regrowth in Smalltooth Sawfish, *Pristis
pectinata*. *Drum and Croaker*, 24(1), 31–33.

Coiraton, C., Amezcua, F. & Ketchum, J.T. (2020). New insights into the migration patterns of
the scalloped hammerhead shark *Sphyrna lewini* based on vertebral microchemistry.
Marine Biology, 167(5), 58. https://doi.org/10.1007/s00227-020-3668-0

Courter, I.I., Child, D.B., Hobbs, J.A., Garrison, T.M., Glessner, J.J.G., Duery, S. et al. (2013).
Resident rainbow trout produce anadromous offspring in a large interior watershed.
Canadian Journal of Fisheries and Aquatic Sciences, 70(5), 701–710.
https://doi.org/10.1139/cjfas-2012-0457

Cuthill, I. (1991). Field experiments in animal behaviour: methods and ethics. *Animal Behaviour*,
42(6), 1007–1014. https://doi.org/10.1016/S0003-3472(05)80153-8

Dulvy, N.K., Davidson, L.N.K., Kyne, P.M., Simpfendorfer, C.A., Harrison, L.R., Carlson, J.K. et al.
(2016). Ghosts of the coast: Global extinction risk and conservation of sawfishes. *Aquatic
Conservation: Marine and Freshwater Ecosystems*, 26(1), 134–153.

https://doi.org/10.1002/aqc.2525

Dunca, E., Fiebig, J. & Pfeiffer, M. (2005). Mutvei's solution: An ideal agent for resolving
microgrowth structures of biogenic carbonates. *Palaeogeography, Palaeoclimatology,*
Palaeoecology, 228(1–2), 149–166. https://doi.org/10.1016/J.PALAEO.2005.03.054

Elsdon, T.S., Wells, B.K., Campana, S.E., Gillanders, B.M., Jones, C.M., Limburg, K.E. et al. (2008).
Otolith chemistry to describe movements and life-history parameters of fishes:
hypotheses, assumptions, limitations and inferences. *Oceanography and Marine Biology:*
an annual review, 46(1), 297–330. https://doi.org/10.1201/9781420065756.ch7

Faria, V. V, McDavitt, M.T., Charvet, P., Wiley, T.R., Simpfendorfer, C.A. & Naylor, G.J.P. (2013).
Species delineation and global population structure of Critically Endangered sawfishes
(Pristidae). *Zoological Journal of the Linnean Society*, 167(1), 136–164.
https://doi.org/10.1111/j.1096-3642.2012.00872.x

Faure, G. & Mensing, T.M. (2004). *Isotopes: principles and applications*. John Wiley & Sons Inc.

Fearing, A., Smith, K., Wiley, T., Whitty, J., Feldheim, K., Kyne, P. et al. (2018). Looking Back for
the Future: Utilizing Sawfish Saws from Natural History Collections to Conserve the
Critically Endangered Largetooth Sawfish (*Pristis pristis*). *Biodiversity Information Science*
and Standards. p. e25806.

Feitosa, L.M., Dressler, V. & Lessa, R.P. (2020). Habitat Use Patterns and Identification of
Essential Habitat for an Endangered Coastal Shark With Vertebrae Microchemistry: The
Case Study of *Carcharhinus porosus*. *Frontiers in Marine Science*, 7(125), 1–12.
https://doi.org/10.3389/fmars.2020.00125

Feitosa, L.M., Martins, A.P.B. & Nunes, J.L.S. (2017). Sawfish (Pristidae) records along the
Eastern Amazon coast. *Endangered Species Research*, 34, 229–234.

Feldheim, K., Fields, A., Chapman, D., Scharer, R. & Poulakis, G. (2017). Insights into
reproduction and behavior of the smalltooth sawfish *Pristis pectinata*. *Endangered Species*
Research, 34, 463–471. https://doi.org/10.3354/esr00868

Fernandez-Carvalho, J., Imhoff, J.L., Faria, V. V, Carlson, J.K. & Burgess, G.H. (2014). Status and
the potential for extinction of the largetooth sawfish *Pristis pristis* in the Atlantic Ocean.
Aquatic Conservation: Marine and Freshwater Ecosystems, 24(4), 478–497.
https://doi.org/10.1002/aqc.2394

Field, I., Meekan, M. & Bradshaw, C. (2009). *Development of non-lethal methods for*
determining age and habitat use of sawfishes from northern Australia. Australian
Government.

Fisher, C.M., Paton, C., Pearson, D.G., Sarkar, C., Luo, Y., Tersmette, D.B. et al. (2017). Data
Reduction of Laser Ablation Split-Stream (LASS) Analyses Using Newly Developed Features
Within Iolite: With Applications to Lu-Hf + U-Pb in Detrital Zircon and Sm-Nd +U-Pb in
Igneous Monazite. *Geochemistry, Geophysics, Geosystems*, 18(12), 4604–4622.
https://doi.org/10.1002/2017GC007187

Flockhart, D.T., Keyser, T.K., Don, C., Miller, N.G. & Norris, R.D. (2015). Experimental evidence
shows no fractionation of strontium isotopes (87Sr/86Sr) among soil , plants , and
herbivores : implications for tracking wildlife and forensic science. *Isotopes in*
Environmental and Health Studies, 202, 123–140.

https://doi.org/http://dx.doi.org/10.1080/10256016.2015.1021345

Gleiss, A.C., Morgan, D.L., Whitty, J.M., Keleher, J.J., Fossette, S. & Hays, G.C. (2017). Are
vertical migrations driven by circadian behaviour? Decoupling of activity and depth use in a
large riverine elasmobranch, the freshwater sawfish (*Pristis pristis*). *Hydrobiologia*, 787(1),
181–191. https://doi.org/10.1007/s10750-016-2957-6

Graham, J., Kroetz, A., Poulakis, G., Scharer, R., Carlson, J., Lowerre-Barbieri, S. et al. (2020).
Large-scale space use of large juvenile and adult smalltooth sawfish *Pristis pectinata*:
implications for management. *Endangered Species Research*, 44, 45–59.
https://doi.org/10.3354/esr01088

Hamann, E.J. & Kennedy, B.P. (2012). Juvenile dispersal affects straying behaviors of adults in a
migratory population. *Ecology*, 93(4), 733–740. https://doi.org/10.1890/11-1009.1

Hamann, E.J., Kennedy, B.P., Whited, D.C. & Stanford, J.A. (2014). Spatial Variability in Spawning
Habitat Selection by Chinook Salmon (*Onchorhynchus Tshawytscha*) in a Wilderness River.
River Research and Applications, 30(9), 1099–1109. https://doi.org/10.1002/rra.2704

Hamer, P., Henderson, A., Hutchison, M., Kemp, J., Green, C. & Feutry, P. (2015). Atypical
correlation of otolith strontium:calcium and barium:calcium across a marine-freshwater
life history transition of a diadromous fish. *Marine and Freshwater Research*, 66(5), 411–
419. https://doi.org/10.1071/MF14001

Haque, A.B., Leeney, R.H. & Biswas, A.R. (2020). Publish, then perish? Five years on, sawfishes
are still at risk in Bangladesh. *Aquatic Conservation: Marine and Freshwater Ecosystems*,
30(12), 2370–2383. https://doi.org/10.1002/aqc.3403

Hegg, J.C. & Fisher, C.M. (2020). Simultaneous collection of ^{87}Sr/^{86}Sr and trace-element data in
otoliths and other sclerochronological hard structures. *bioRxiv*, 1–17.
https://doi.org/10.1101/2020.04.24.060640

Hegg, J.C., Kennedy, B.P. & Chittaro, P.M. (2018). What did you say about my mother? The
complexities of maternally derived chemical signatures in otoliths. *Canadian Journal of
Fisheries and Aquatic Sciences*, 14(April), 1–14. https://doi.org/10.1139/cjfas-2017-0341

Hegg, J.C., Kennedy, B.P., Chittaro, P.M. & Zabel, R.W. (2013). Spatial structuring of an evolving
life-history strategy under altered environmental conditions. *Oecologia*, 172(4), 1017–
1029. https://doi.org/10.1007/s00442-012-2564-9

Hegg, J.C., Kennedy, B.P. & Fremier, A.K. (2013). Predicting strontium isotope variation and fish
location with bedrock geology: Understanding the effects of geologic heterogeneity.
Chemical Geology, 360–361, 89–98. https://doi.org/10.1016/j.chemgeo.2013.10.010

Hobson, K.A., Barnett-Johnson, R. & Cerling, T. (2010). Using isoscapes to track animal
migration. *Isoscapes: understanding movement, pattern, and process on Earth through
isotope mapping*. Springer Verlag, pp. 273–298.

Huston, C., Stevens, P., Blaxton, R., Tolley, S., Scharer, R., Tornwall, B. et al. (2017). Diel
movements of juvenile smalltooth sawfish: implications for defining the size of a nursery
hotspot. *Endangered Species Research*, 34, 311–322. https://doi.org/10.3354/esr00851

Izzo, C., Huveneers, C., Drew, M. & A Bradshaw, C.J. (2016). Vertebral chemistry demonstrates
movement and population structure of bronze whaler. *Marine Ecology Progress Series*,
556, 195–207. https://doi.org/10.3354/meps11840

Jabado, R., Al Baharna, R., Al Ali, S., Al Suwaidi, K., Al Blooshi, A. & Al Dhaheri, S. (2017). Is this
the last stand of the Critically Endangered green sawfish *Pristis zijsron* in the Arabian Gulf?
Endangered Species Research, 32(1), 265–275. https://doi.org/10.3354/esr00805

Kalish, J.M. (1990). Use of Otolith Microchemistry to Distinguish the Progeny of Sympatric
Anadromous and Non-anadromous Salmonids. *Fisheries Bulletin*, 88(4), 657–666.

Kennedy, B.P., Blum, J.D., Folt, C.L. & Nislow, K.H. (2000). Using natural strontium isotopic
signatures as fish markers: methodology and application. *Canadian Journal of Fisheries and*
Aquatic Sciences, 57(11), 2280–2292. https://doi.org/10.1139/f00-206

Kennedy, B.P., Folt, C.L., Blum, J.D. & Chamberlain, C.P. (1997). Natural isotope markers in
salmon. *Nature*, 387(6635), 766–767. https://doi.org/0.1038/42835

Koch, J.D. & Quist, M.C. (2007). A Technique for Preparing Fin Rays and Spines for Age and
Growth Analysis. *North American Journal of Fisheries Management*, 27(3), 782–784.
https://doi.org/10.1577/m06-224.1

Kraus, R.T. & Secor, D.H. (2004). Incorporation of strontium into otoliths of an estuarine fish.
Journal of Experimental Marine Biology and Ecology, 302(1), 85–106.
https://doi.org/10.1016/j.jembe.2003.10.004

Kyne, P., Oetinger, M., Grant, M. & Feutry, P. (2021). Life history of the Critically Endangered
largetooth sawfish: a compilation of data for population assessment and demographic
modelling. *Endangered Species Research*, 44, 79–88. https://doi.org/10.3354/esr01090

Kyne, P.M., Carlson, J. & Smith, K. (2013). *Pristis pristis (errata version published 2019)*.

Laslett, R.E. (1995). Concentrations of dissolved and suspended particulate Cd, Cu, Mn, Ni, Pb
and Zn in surface waters around the coasts of England and Wales and in adjacent seas.
Estuarine, Coastal and Shelf Science, 40(1), 67–85. https://doi.org/10.1016/0272-
7714(95)90014-4

Leeney, R.H. & Poncelet, P. (2015). Using fishers' ecological knowledge to assess the status and
cultural importance of sawfish in Guinea-Bissau. *Aquatic Conservation: Marine and*
Freshwater Ecosystems, 25(3), 411–430. https://doi.org/10.1002/aqc.2419

Lehman, R.N., Poulakis, G.R., Scharer, R.M., Schweiss, K.E., Hendon, J.M. & Phillips, N.M. (2020).
An environmental DNA tool for monitoring the status of the Critically Endangered
Smalltooth Sawfish, *Pristis pectinata*, in the western Atlantic. *Conservation Genetics*
Resources, 12(4), 621–629. https://doi.org/10.1007/s12686-020-01149-5

Lewis, J.P., Patterson, W.F., Carlson, J.K. & McLachlin, K. (2016). Do vertebral chemical
signatures distinguish juvenile blacktip shark (*Carcharhinus limbatus*) nursery regions in
the northern Gulf of Mexico? *Marine and Freshwater Research*, 67(7), 1014–1022.
https://doi.org/10.1071/MF15088

Limburg, K.E. & Casini, M. (2018). Effect of marine hypoxia on Baltic Sea cod *Gadus morhua*:
Evidence from otolith chemical proxies. *Frontiers in Marine Science*, 5(DEC), 482.
https://doi.org/10.3389/fmars.2018.00482

Limburg, K.E., Wuenschel, M.J., Hüssy, K., Heimbrand, Y. & Samson, M. (2018). Making the
Otolith Magnesium Chemical Calendar-Clock Tick: Plausible Mechanism and Empirical
Evidence. *Reviews in Fisheries Science and Aquaculture*, 26(4), 479–493.
https://doi.org/10.1080/23308249.2018.1458817

López-Angarita, J., Camilo Cubillos-M, J., Villate-Moreno, M., Del Cid, A., Díaz, J.M., Cooke, R. et al. (2021). Identifying priority areas for research and conservation of the largetooth sawfish *Pristis pristis* in Colombia and Panama. *BioRxiv*, 1–16. https://doi.org/10.1101/2021.01.14.426658

Mackenzie, D.I., Nichols, J.D., Sutton, N., Kawanishi, K. & Bailey, L.L. (2005). Improving Inferences in Population Studies of Rare Species That are Detected Imperfectly. *Special Feature Ecology*, 86(5), 1101–1113. https://doi.org/10.1890/04-1060

Máñez, K.S., Holm, P., Blight, L., Coll, M., MacDiarmid, A., Ojaveer, H. et al. (2014). The future of the oceans past: Towards a global marine historical research initiative. *PLoS ONE*, 9(7). https://doi.org/10.1371/journal.pone.0101466

McMillan, M.N., Izzo, C., Junge, C., Albert, O.T., Jung, A. & Gillanders, B.M. (2017). Analysis of vertebral chemistry to assess stock structure in a deep-sea shark, *Etmopterus spinax. ICES Journal of Marine Science*, 74(3), 793–803. https://doi.org/10.1093/icesjms/fsw176

Melo Palmeira, C.A., da Silva Rodrigues-Filho, L.F., de Luna Sales, J.B., Vallinoto, M., Schneider, H. & Sampaio, I. (2013). Commercialization of a critically endangered species (largetooth sawfish, *Pristis perotteti*)in fish markets of northern Brazil: Authenticity by DNA analysis. *Food Control*, 34(1), 249–252. https://doi.org/10.1016/j.foodcont.2013.04.017

Mendoza, A., Kelez, S., Cherres, W.G. & Maguiño, R. (2017). The largetooth sawfish, Pristis pristis (Linnaeus, 1758), is not extirpated from Peru: New records from Tumbes. *Check List*, 13(4), 261–265. https://doi.org/10.15560/13.4.261

Miller, W.A. (1974). Observations on the Developing Rostrum and Rostral Teeth of Sawfish: *Pristis perotteti* and *P. cuspidatus. Copeia*, 1974(2), 311–318. https://doi.org/10.2307/1442525

Mohan, J., Rahman, M.S., Thomas, P. & Walther, B. (2014). Influence of constant and periodic experimental hypoxic stress on Atlantic croaker otolith chemistry. *Aquatic Biology*, 20(1), 1–11. https://doi.org/10.3354/ab00542

Mohan, J.A., Miller, N.R., Herzka, S.Z., Sosa-Nishizaki, O., Kohin, S., Dewar, H. et al. (2018). Elements of time and place: manganese and barium in shark vertebrae reflect age and upwelling histories. *Proceedings of the Royal Society B: Biological Sciences*, 285(1890), 20181760. https://doi.org/10.1098/rspb.2018.1760

Mohan, J.A., Rulifson, R.A., Corbett, D.R. & Halden, N.M. (2012). Validation of Oligohaline Elemental Otolith Signatures of Striped Bass by Use of In Situ Caging Experiments and Water Chemistry. *Marine and Coastal Fisheries*, 4(1), 57–70. https://doi.org/10.1080/19425120.2012.656533

Norton, S.L., Wiley, T.R., Carlson, J.K., Frick, A.L., Poulakis, G.R. & Simpfendorfer, C.A. (2012). Designating Critical Habitat for Juvenile Endangered Smalltooth Sawfish in the United States. *Marine and Coastal Fisheries*, 4(1), 473–480. https://doi.org/10.1080/19425120.2012.676606

Nunes, J.L.S., Rincon, G., Piorski, N.M. & Martins, A.P.B. (2016). Near-term embryos in a *Pristis pristis* (Elasmobranchii: Pristidae) from Brazil. *Journal of Fish Biology*. pp. 1112–1120.

Paillon, C., Wantiez, L., Kulbicki, M., Labonne, M. & Vigliola, L. (2014). Extent of mangrove nursery habitats determines the geographic distribution of a coral reef fish in a South-

Pacific Archipelago. C Savage, Ed. *PLoS ONE*, 9(8), e105158.
https://doi.org/10.1371/journal.pone.0105158

Parris, K.M., McCall, S.C., McCarthy, M.A., Minteer, B.A., Steele, K., Bekessy, S. et al. (2010).
Assessing ethical trade-offs in ecological field studies. *Journal of Applied Ecology*, 47(1),
227–234. https://doi.org/10.1111/j.1365-2664.2009.01755.x

Peverell, S.C. (2005). Distribution of sawfishes (Pristidae) in the Queensland Gulf of Carpentaria,
Australia, with notes on sawfish ecology. *Environmental Biology of Fishes*, 73(4), 391–402.
https://doi.org/10.1007/s10641-005-1599-8

Peverell, S.C. (2008). Sawfish (Pristidae) of the Gulf of Carpentaria, Queensland Australia. .
James Cook University.

Phillips, Nicole., Wueringer, B. (2015). Sawfish: Ancient predators in need of modern
conservation tools. *Wildlife Australia*, 52(1), 14–17.

Phillips, N., Chaplin, J., Morgan, D. & Peverell, S. (2009). Extraction and amplification of DNA
from the dried rostra of sawfishes (Pristidae) for applications in conservation genetics.
Pacific Conservation Biology, 15(2), 128. https://doi.org/10.1071/PC090128

Pistevos, J.C.A., Reis-Santos, P., Izzo, C. & Gillanders, B.M. (2019). Element composition of shark
vertebrae shows promise as a natural tag. *Marine and Freshwater Research*. pp. 1722–
1733.

Poulakis, G.R., Stevens, P.W., Timmers, A.A., Wiley, T.R. & Simpfendorfer, C.A. (2011). Abiotic
affinities and spatiotemporal distribution of the endangered smalltooth sawfish, *Pristis*
pectinata, in a south-western Florida nursery. *Marine and Freshwater Research*, 62(10),
1165–1177. https://doi.org/10.1071/MF11008

Raoult, V., Peddemors, V.M., Zahra, D., Howell, N., Howard, D.L., De Jonge, M.D. et al. (2016).
Strontium mineralization of shark vertebrae. *Scientific Reports*, 6(1), 1–10.
https://doi.org/10.1038/srep29698

Rohtla, M. & Vetemaa, M. (2016). Otolith chemistry chimes in: migratory environmental
histories of Atlantic tarpon (*Megalops atlanticus*) caught from offshore waters of French
Guiana. *Environmental Biology of Fishes*, 99(8–9), 593–602.
https://doi.org/10.1007/s10641-016-0501-1

Schaffler, J.J., Miller, T.J. & Jones, C.M. (2014). Spatial and temporal variation in otolith
chemistry of juvenile atlantic menhaden in the Chesapeake Bay. *Transactions of the*
American Fisheries Society, 143(4), 1061–1071.
https://doi.org/doi:10.1080/00028487.2014.889748

Scharer, R.M., Patterson, W.F., Carlson, J.K. & Poulakis, G.R. (2012). Age and Growth of
Endangered Smalltooth Sawfish (*Pristis pectinata*) Verified with LA-ICP-MS Analysis of
Vertebrae. *PLoS ONE*, 7(10), 1–8. https://doi.org/10.1371/journal.pone.0047850

Scharer, R.M., Stevens, P.W., Shea, C.P. & Poulakis, G.R. (2017). All nurseries are not created
equal: Large-scale habitat use patterns in two smalltooth sawfish nurseries. *Endangered*
Species Research, 34, 473–492. https://doi.org/10.3354/ESR00871

Schindler, D.E., Hilborn, R., Chasco, B., Boatright, C.P., Quinn, T.P., Rogers, L.A. et al. (2010).
Population diversity and the portfolio effect in an exploited species. *Nature*, 465(7298),
609–612. https://doi.org/10.1038/nature09060

Schmid, K. & Giarrizzo, T. (2017). New Record of a Largetooth Sawfish Specimen from the
Amazon River Estuary in Northern Brazil. *Fisheries*, 42(5), 254–255.
https://doi.org/10.1080/03632415.2017.1276357

Schöne, B.R., Dunca, E., Mutvei, H. & Norlund, U. (2004). A 217-year record of summer air
temperature reconstructed from freshwater pearl mussels (*M. margarifitera*, Sweden).
Quaternary Science Reviews, 23(16–17), 1803–1816.
https://doi.org/10.1016/J.QUASCIREV.2004.02.017

Schroeder, R.K., Whitman, L.D., Cannon, B., Olmsted, P. & Rennie, M. (2016). Juvenile life-
history diversity and population stability of spring Chinook salmon in the Willamette River
basin, Oregon. *Canadian Journal of Fisheries and Aquatic Sciences*, 73(6), 921–934.
https://doi.org/10.1139/cjfas-2015-0314

Secor, D.H., Dean, J.M. & Laban, E.H. (1992). Otolith removal and preparation for
microstructural analysis. In: , Campana SE, & , Stevenson DK (eds). *Otolith Microstructure*
Examination and Analysis. Ottawa, pp. 19–57.

Seeley, M. & Walther, B. (2017). Facultative oligohaline habitat use in a mobile fish inferred
from scale chemistry. *Marine Ecology Progress Series*, 598, 1–13.
https://doi.org/10.3354/meps12223

Seitz, J.C. & Poulakis, G.R. (2006). Anthropogenic effects on the smalltooth sawfish (*Pristis*
pectinata) in the United States. *Marine Pollution Bulletin*, 52(11), 1533–1540.
https://doi.org/10.1016/J.MARPOLBUL.2006.07.016

Shellis, R.P. & Berkovitz, B.K.B. (1980). Dentine structure in the rostral teeth of the sawfish
Pristis (Elasmobranchii). *Archives of Oral Biology*, 25(5), 339–343.
https://doi.org/10.1016/0003-9969(80)90043-6

Shippentower, G.E., Schreck, C.B. & Heppell, S.A. (2011). Who's your momma? Recognizing
maternal origin of juvenile steelhead using injections of strontium chloride to create
transgenerational marks. *Transactions of the American Fisheries Society*, 140(5), 1330–
1339. https://doi.org/10.1080/00028487.2011.620488

Simpfendorfer, C., Kyne, P., Noble, T., Goldsbury, J., Basiita, R., Lindsay, R. et al. (2016).
Environmental DNA detects Critically Endangered largetooth sawfish in the wild.
Endangered Species Research, 30(1), 109–116. https://doi.org/10.3354/esr00731

Simpfendorfer, C.A. (2000). Predicting population recovery rates for endangered western
Atlantic sawfishes using demographic analysis. *Environmental Biology of Fishes*, 58, 371–
377. https://doi.org/10.1023/A:1007675111597

Slaughter, B.H. & Springer, S. (1968). Replacement of Rostral Teeth in Sawfishes and Sawsharks.
Copeia, 1968(3), 499. https://doi.org/10.2307/1442018

Smith, W.D., Miller, J.A. & Heppell, S.S. (2013). Elemental Markers in Elasmobranchs: Effects of
Environmental History and Growth on Vertebral Chemistry. D Steinke, Ed. *PLoS ONE*, 8(10),
e62423. https://doi.org/10.1371/journal.pone.0062423

Smith, W.D., Miller, J.A., Márquez-Farías, F. & Heppell, S.S. (2016). Elemental signatures reveal
the geographic origins of a highly migratory shark: Prospects for measuring population
connectivity. D Steinke, Ed. *Marine Ecology Progress Series*, 556(10), 173–193.
https://doi.org/10.3354/meps11844

Thorburn, D.C., Morgan, D.L., Rowland, A.J. & Gill, H.S. (2007). Freshwater Sawfish *Pristis*
microdon Latham, 1794 (Chondrichthyes: Pristidae) in the Kimberley region of Western
Australia. *Zootaxa*, 1471(1471), 27–41. https://doi.org/10.11646/zootaxa.1471.1.3

Thornton, E.K. (2011). Reconstructing ancient Maya animal trade through strontium isotope
(^{87}Sr/^{86}Sr) analysis. *Journal of Archaeological Science*, 38(12), 3254–3263.
https://doi.org/10.1016/j.jas.2011.06.035

Thorrold, S.R. & Shuttleworth, S. (2000). In situ analysis of trace elements and isotope ratios in
fish otoliths using laser ablation sector field inductively coupled plasma mass
spectrometry. *Canadian Journal of Fisheries and Aquatic Sciences*, 57(6), 1232–1242.
https://doi.org/10.1139/f00-054

Thorson, T.B. (1982). Life history implications of a tagging study of the largetooth sawfish,
Pristis perotteti, in the Lake Nicaragua-Río San Juan system. *Environmental Biology of*
Fishes, 7(3), 207–228. https://doi.org/10.1007/BF00002497

Tillett, B., Meekan, M., Parry, D., Munksgaard, N., Field, I., Thorburn, D. et al. (2011). Decoding
fingerprints: elemental composition of vertebrae correlates to age-related habitat use in
two morphologically similar sharks. *Marine Ecology Progress Series*, 434, 133–142.
https://doi.org/10.3354/meps09222

Turner, S.M., Limburg, K.E. & Palkovacs, E.P. (2015). Can different combinations of natural tags
identify river herring natal origin at different levels of stock structure? *Canadian Journal of*
Fisheries and Aquatic Sciences, 72(February), 845–854. https://doi.org/10.1139/cjfas-
2014-0403

Walther, B.D. (2019). The art of otolith chemistry: Interpreting patterns by integrating
perspectives. *Marine and Freshwater Research*, 70(12), 1643–1658.
https://doi.org/10.1071/MF18270

Walther, B.D. & Limburg, K.E. (2012). The use of otolith chemistry to characterize diadromous
migrations. *Journal of Fish Biology*, 81(2), 796–825. https://doi.org/10.1111/j.1095-
8649.2012.03371.x

Walther, B.D. & Thorrold, S.R. (2008). Continental-scale variation in otolith geochemistry of
juvenile American shad (*Alosa sapidissima*). *Canadian Journal of Fisheries and Aquatic*
Sciences, 65(12), 2623–2635. https://doi.org/10.1139/F08-164

Wang, C.-H., Walther, B.D. & Gillanders, B.M. (2019). Introduction to the 6th International
Otolith Symposium. *Marine and Freshwater Research*, 70(12), i–iii.
https://doi.org/10.1071/mfv70n12_ed

Warner, R.R., Swearer, S.E., Caselle, J.E., Sheehy, M. & Paradis, G. (2005). Natal Trace-Elemental
Signatures in the Otoliths of an Open-Coast Fish. *Limnology and Oceanography*, 50(5),
1529–1542. https://doi.org/10.4319/lo.2005.50.5.1529

Welten, M., Smith, M.M., Underwood, C. & Johanson, Z. (2015). Evolutionary origins and
development of saw-teeth on the sawfish and sawshark rostrum (Elasmobranchii;
Chondrichthyes). *Royal Society Open Science*, 2(9), 150189.
https://doi.org/10.1098/rsos.150189

Whitty, J.M., Morgan, D.L., Peverell, S.C., Thorburn, D.C. & Beatty, S.J. (2009). Ontogenetic
depth partitioning by juvenile freshwater sawfish (*Pristis microdon*: Pristidae) in a riverine

environment. *Marine and Freshwater Research*, 60(4), 306–316.
https://doi.org/10.1071/MF08169

Whitty, J.M., Phillips, N.M., Thorburn, D.C., Simpfendorfer, C.A., Field, I., Peverell, S.C. et al.
(2014). Utility of rostra in the identification of Australian sawfishes *(Chondrichthyes:*
Pristidae). Aquatic Conservation: Marine and Freshwater Ecosystems, 24(6), 791–804.
https://doi.org/10.1002/aqc.2398

Wiley, T.R. & Simpfendorfer, C.A. (2010). Using public encounter data to direct recovery efforts
for the endangered smalltooth sawfish *Pristis pectinata. Endangered Species Research*,
12(3), 179–191. https://doi.org/10.3354/esr00303

Woodcock, S.H. & Walther, B.D. (2014). Trace elements and stable isotopes in Atlantic tarpon
scales reveal movements across estuarine gradients. *Fisheries Research*, 153, 9–17.
https://doi.org/10.1016/j.fishres.2014.01.003

Wueringer, B.E., Squire, L. & Collin, S.P. (2009). The biology of extinct and extant sawfish
(Batoidea: Sclerorhynchidae and Pristidae). *Reviews in Fish Biology and Fisheries*, 19(4),
445–464. https://doi.org/10.1007/s11160-009-9112-7

Yan, H.F.H.F., Kyne, P.M.P.M., Jabado, R.W.R.W., Leeney, R.H.R.H., Davidson, L.N.K., Derrick,
D.H.D.H. et al. (2021). Overfishing and habitat loss drives range contraction of iconic
marine fishes to near extinction. *Science Advances*, 7(7), 1–11.
https://doi.org/10.1126/sciadv.abb6026

Zimmerman, C.E. (2005). Relationship of otolith strontium-to-calcium ratios and salinity:
experimental validation for juvenile salmonids. *Canadian Journal of Fisheries and Aquatic*
Sciences, 62(1), 88–97. https://doi.org/10.1139/f04-182

Figures

964
965   *Figure 1* – Rostra from two largetooth sawfish were used in this study. A sub-adult rostrum (A)
966   of unknown origin had a TRL of 57cm and a SRL of 55cm. The embryo rostrum collected in
967   Belize in 1969 had a TRL of 14.2cm and a SRL of 13.6cm. The most distal tooth on the left-hand
968   side was taken for analysis from each rostrum.

969
970
Figure 2 – Sub-adult largetooth sawfish rostral tooth sectioned on the dorsal-ventral plane (A)
shows internal tubule structures and faint banding. Staining with Mutvei's solution (acetic acid,
glutaraldehyde, and alcian blue) highlighted the tubules which broke the surface in darker blue
(B). Inverting the color spectrum of the stained section (C) further improved visualization of the
surface tubules. Tubules overlapped in curved bands (D), with "legs" descending distally from
the core of the tooth to the edge. These tubules were less distinct near the base of the tooth,
however faint banding was also visible. Putative growth was quantified by two readers (reader
1 – red dots, reader 2 – purple dots). Measurements agreed despite being offset where
descending tubule "legs" were prominent (D). Agreement was lower in sections with less
prominent tubules (E).
981
982
983
984

Figure 3 – Results of split stream LA-ICPMS analysis of sectioned rostral teeth from to
largetooth sawfish specimens, sub-adult (left-hand column) and an embryo (right-hand column)
showed significant variation across the tooth. Relative age of the fish is indicated above the
plots, with the youngest age represented by zero microns, and older ages with increasing
distance along the tooth. Strontium ratio ($^{87}Sr/^{86}Sr$, A) indicated movement within fresh or
brackish water, with movement toward the global marine signature (blue horizontal line) in
mid-life indicating movement into higher salinity water. Meanwhile the embryo $^{87}Sr/^{86}Sr$
appears to match the global marine signature throughout life. Strontium to calcium ratio
(Sr/Ca, B) did not rise significantly during mid-life in the sub-adult, confirming residence in
brackish rather than fully saline water. The embryo Sr/Ca, in contrast, is elevated, confirming a
likely ocean residence during provisioning or development. Barium to calcium ratios (Ba/Ca, C)
increase in fresh water and decrease quickly in the presence of salinity, confirming sub-adult
and embryo movements inferred from the prior tracers. Magnesium to calcium ratios (Mg/Ca,
D) are linked to growth rate in other species. Peaks in Mg/Ca could indicate growth rate
changes as well as movement into new locations. Manganese to calcium ratios (Mn/Ca, E) are
linked to reducing conditions such as mangrove swamps, as well as hypoxic stress in some
teleosts, as ionic Mn is only available in significant quantities in the water column due to a
redox reactions in low oxygen environments. Changes in Mn/Ca has also been shown to be
related to fish condition in some studies.
1005