The versican-hyaluronan complex provides an essential extracellular matrix niche for

Flk1+ hematoendothelial progenitors

[1] Sumeda Nandadasa, PhD, [1] Anna O'Donnell BS, MBA, [2]Ayako Murao BS, [2]Yu Yamaguchi,

MD, PhD, [1] Ronald J. Midura PhD, [3]Lorin Olson, PhD and [1*]Suneel S. Apte, MBBS, DPhil

[1] Department of Biomedical Engineering (ND20), Cleveland Clinic Lerner Research Institute,

9500 Euclid Avenue, Cleveland, OH 44195

[2] Human Genetics Program, Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA

92037

[3]Cardiovascular Biology Research Program, Oklahoma Medical Research Foundation,

Oklahoma City, OK 73104

Human Genetics Program, Sanford Prebys Medical Discovery Institute

[*]Corresponding author: Suneel S. Apte, MBBS, D.Phil., Department of Biomedical Engineering-

ND20, Lerner Research Institute, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195,

USA. Phone: 216.445.3278; E-mail: aptes@ccf.org

Short title: Versican and hyaluronan in vasculogenesis

Total word count

22
23
24

Abstract

Little is known about extracellular matrix (ECM) contributions to formation of the earliest cell

lineages in the embryo. Here, we show that the proteoglycan versican and glycosaminoglycan

hyaluronan are associated with emerging Flk1$^+$ hematoendothelial progenitors at gastrulation.

The mouse versican mutant *Vcan*^{hdf} lacks yolk sac vasculature, with attenuated yolk sac

hematopoiesis. CRISPR/Cas9-mediated *Vcan* inactivation in mouse embryonic stem cells

reduced vascular endothelial and hematopoietic differentiation in embryoid bodies, which

generated fewer blood colonies, and had an impaired angiogenic response to VEGF$_{165}$. HA was

severely depleted in *Vcan*^{hdf} embryos, with corresponding increase in the HA-depolymerase

TMEM2. Conversely, HA-deficient mouse embryos also had vasculogenic suppression but with

increased versican proteolysis. VEGF$_{165}$ and Indian hedgehog, crucial vasculogenic factors,

utilized the versican-HA matrix, specifically versican chondroitin sulfate chains, for binding.

Versican-HA ECM is an obligate requirement for vasculogenesis and primitive hematopoiesis,

providing a vasculogenic factor-enriching microniche for Flk1$^+$ progenitors from their origin at

gastrulation.

KEYWORDS: Proteoglycan; extracellular matrix; hyaluronan; Tie-2; versican;

glycosaminoglycan; vasculogenesis; angiogenesis; vascular endothelium; embryonic stem cell;

TMEM2;ADAMTS;

Introduction

The first blood vessels in amniotes form *de novo* in the yolk sac prior to initiation of hemodynamic forces, indicating an in situ differentiation driven by the local environment, which includes extracellular matrix (ECM)[1]. This vasculogenic process is inextricably linked to subsequent primitive hematopoiesis, i.e., formation of the first erythroid and myeloid cells [2], implying a close lineage relationship between the first vascular endothelial and blood cells. The relevant Flk1$^+$ progenitor cells arise from mesodermal precursors at gastrulation [3], migrate from the primitive streak, and aggregate in blood islands in the proximal extra-embryonic yolk sac on the seventh day of gestation (E7) in mouse embryos [4, 5]. Several key transcription factors and soluble effectors of vasculogenesis/angiogenesis and hematopoiesis are known [1]. Although ECM and ECM-derived proteolytic fragments are recognized as angiogenesis regulators [6] and sulfated glycosaminoglycans can promote endothelial differentiation of mesenchymal stem cells [7], specific ECM proteoglycans that influence vasculogenesis are unknown. Another early cell lineage, primordial germ cells, are accompanied by a protective "traveling niche" of steel factor/stem cell factor-producing cells during migration to the gonads [8]. Whether a cell-associated ECM supports other early lineages such as hematoendothelial progenitors is unknown.

ECM surrounds all mammalian cells, forming a distinct pericellular matrix in some cell types, and the interstitial matrix of tissues. It influences cell behavior by modulating cell adhesion, migration and tissue mechanics, sequesters growth factors and cytokines, and its proteolysis can generate bioactive fragments. Proteoglycans, which are ECM and cell-surface molecules with one or more glycosaminoglycan chains covalently attached to a core protein, participate in all these mechanisms. For example, chondroitin sulfate proteoglycans (CSPGs) such as versican, which are bulky and inherently anti-adhesive, regulate focal adhesion formation and cell migration [9-11]. Heparan sulfate proteoglycans (HSPGs) sequester growth factors, including angiogenic growth factors such as VEGF$_{165}$ and FGF2 through their HS chains and act as co-receptors in

angiogenic signaling [12]. CSPGs can also bind $VEGF_{165}$ and may overlap functionally with

HSPGs in VEGF-induced sprouting angiogenesis [13], but physiological in vivo contexts for this

regulatory activity remain unidentified for CSPGs.

Versican is a widely distributed CSPG that aggregates with the glycosaminoglycan

hyaluronan (HA), through its N-terminal G1 domain [14-17]. Its C-terminal G3 domain binds to

ECM components fibronectin, fibrillins and tenascins [18] and was shown to interact with $VEGF_{165}$

in biochemical assays [19]. HA is anchored to cell-surface receptors such as CD44 and RHAMM

as well as the membrane-localized hyaluronan synthases [20], providing a means by which

versican can localize to the pericellular matrix. Pericellular versican was previously shown to

modulate smooth muscle cell differentiation via regulation of cell adhesion [9, 21, 22]. Four

versican isoforms (V0, V1-V3) arise from alternative splicing of large exons, numbered 7 and 8,

encoding CS-bearing domains GAGα and GAGβ, respectively [16, 23]. A versican insertional

mutant mouse allele ($Vcan^{hdf}$), which lacks all isoforms, demonstrated an essential role for

versican in early cardiac development [24, 25]. *Has2* null embryos, lacking the major HA

synthase, die by 10.5 days of gestation with similar cardiac defects as $Vcan^{hdf/hdf}$ [26], consistent

with the versican-HA complex being a major constituent of cardiac jelly. Here, upon identification

of abnormal vasculature in $Vcan^{hdf/hdf}$ yolk sac, we investigated its role in vasculogenesis and early

hematopoiesis, with investigation of the key findings in embryos with inactivation of hyaluronan

synthase (*Has*) genes. An intimate association of versican and HA with $Flk1^+$ hematoendothelial

progenitor cells from their origin at gastrulation is demonstrated to have profound significance for

vasculogenesis and primitive hematopoiesis.

Results

$Vcan^{hdf/hdf}$ yolk sacs lack a vascular plexus:

$Vcan^{hdf/hdf}$ embryos (Figure 1-Supplement 1) did not survive past E10.5, and were recognizable

by consistently smaller size and dilated pericardial sac at E9.5 (Fig. 1A). Although major

developmental milestones, including axial rotation of the embryo and initiation of cardiac

contraction were completed by E9.5, *Vcan*^{hdf/hdf} yolk sacs and embryos were less vascular (Fig.

1B,C). Whole mount CD31 immunostaining revealed the lack of a vascular network in E9.5

Vcan^{hdf/hdf} yolk sacs, instead of which scattered vascular CD31+ cells were present (Fig. 1C).

In E9.5 wild-type yolk sac versican localized to the mesoderm, but no staining was present

in *Vcan*^{hdf/hdf} yolk sacs (Fig. 1D). HA-binding protein (HABP) staining overlapped with versican in

wild-type yolk sac mesoderm but was absent in *Vcan*^{hdf/hdf} yolk sacs (Fig. 1D). Fibronectin

immunostaining in contrast, showed increased staining in *Vcan*^{hdf/hdf} yolk sacs, whereas collagen

IV staining was unaffected (Fig. 1D), suggesting that loss of HA did not reflect global ECM

reduction in *Vcan*^{hdf/hdf} yolk sacs. Similar to *Has2*, *Itga5* and *Fn1*-null alleles, which have failed

yolk sac vasculogenesis and hematopoiesis [26, 27], *Vcan*^{hdf/hdf} yolk sacs consistently showed

mesoderm detachment from visceral endoderm with few evident blood islands (Fig 1D, Fig.1

Supplement 2 A-C). Intense F-actin staining was consistently observed in *Vcan*^{hdf/hdf} visceral

endoderm (Fig. 1 D, Fig.1 Supplement 2A), and rounded, rather than spindle-shaped nuclei were

observed in the yolk sac mesothelial layer (Fig.1 Supplement 2B). Transmission electron

microscopy (TEM), undertaken with fixation conditions that preserved cell-matrix interactions [28]

showed complete coverage of blood islands by vascular endothelium in wild-type yolk sac but

discontinuous vascular endothelial cells with numerous membrane protrusions in *Vcan*^{hdf/hdf} yolk

sacs (Fig.1 Supplement 2C). Thus, versican is essential for proper yolk sac morphogenesis.

E8.5 *Vcan*^{hdf/hdf} embryos had a normal shape, but appeared pale and their yolk sacs lacked

a visible vascular network (Fig. 2A). Whole-mount CD31 immunostaining of E8.5 embryos

identified severe, widespread attenuation of vasculature (Fig. 2B). *En face* imaging of whole-

mount wild-type yolk sac stained with anti-CD31 demonstrated a well-formed vascular plexus

lacking in *Vcan*^{hdf/hdf} yolk sacs (Fig. 2C). Immunostaining of versican and CD31, together with HA-

staining in E8.5 wild-type yolk sacs revealed versican and HA co-localization in patches

corresponding to individual cells on the mesothelial aspect of blood islands (Fig. 2D). mRNA *in*

situ hybridization with *Vcan* exon 7 (GAGα), or exon 8 (GAGβ)-specific probes revealed that only

Figure-1

Figure 1. *Vcan^{hdf/hdf}* yolk sacs are avascular and lack hyaluronan.
(A) E9.5 wild type and *Vcan^{hdf/hdf}* embryos. The red arrowhead shows the dilated pericardial sac in the latter.
(B) E9.5 yolk sacs imaged in situ demonstrates the absence of vasculature in the E9.5 *Vcan^{hdf/hdf}* yolk sac.
(C) Three-dimensional (3D) maximum-intensity projections of whole-mount yolk sacs stained with anti-CD31 (red) showing absence of the vascular network in E9.5 *Vcan^{hdf/hdf}* yolk sacs (n=3 yolk sacs of each genotype).
(D) Versican GAGβ staining (green) is present throughout the mesoderm (Me) of E9.5 wild type yolk sacs and absent in the *Vcan^{hdf/hdf}* yolk sac. F-actin (red) staining highlights visceral endoderm (VE), detaches from mesoderm in *Vcan^{hdf/hdf}* yolk sacs. HA and fibronectin (green) were similarly distributed as versican in wild type yolk sac, but HA staining was absent in *Vcan^{hdf/hdf}* yolk sac and fibronectin staining was more intense. Collagen IV staining intensity was similar in wild type and *Vcan^{hdf/hdf}* yolk sacs (n=3 yolk sacs of each genotype). Scale bar in A, B=1mm, C= 100μm, D= 50μm.

the exon 8 probe hybridized to E8.5 yolk sac mesoderm (Fig 2E), suggesting exclusively versican

V1 isoform expression at E8.5. Both *Vcan* probes hybridized strongly to E8.5 wild-type hearts

(Fig. 2F) but not $Vcan^{hdf/hdf}$ hearts (e.g., exon 8 probe, Fig. 2G), indicative of specificity.

Versican and HA are associated with yolk sac Flk1^{+} cells

Flk1, a well characterized hematoendothelial progenitor marker indispensable for both vascular

and blood lineage development [29], showed complete overlap with versican and HA on the

mesothelial aspect of blood islands in wild-type E8.5 yolk sac (Fig. 3A). $Vcan^{hdf/hdf}$ yolk sacs, in

contrast, showed small blood islands with dramatically attenuated HA and Flk1 staining (Fig.

3A). *En face* confocal microscopy of whole-mount wild-type E8.5 yolk sacs revealed discrete

patches of versican throughout the mesoderm (Fig. 3B). CD41, which is expressed upon

specific commitment to the blood lineage [30], was strongly expressed in blood islands in the

mid-mesoderm plane in wild-type yolk sacs (Fig. 3B). Versican did not precisely overlap with

CD41+ cells and was restricted to intense patches corresponding to individual endothelial cells

in wild-type yolk sacs, evident both in the surface (mesothelial) plane and cross-sections,

whereas $Vcan^{hdf/hdf}$ yolk sacs lacked CD41-stained blood islands entirely (Fig. 3B). High

magnification images showed a few cells with weak CD41 staining in $Vcan^{hdf/hdf}$ yolk sac

mesoderm (arrowheads in Fig. 3B), contrasting with well-demarcated CD41+ wild-type blood

islands. Combined, these results show that versican and HA distribution at E8.5 is restricted and

specifically associated with uncommitted Flk1^{+} hemogenic endothelial cells, which are crucial for

both blood and vascular development in the mouse embryo. Versican and HA are not

specifically associated with subsequently differentiated blood cells (Fig. 3B,C).

Versican is required for blood formation

qRT-PCR revealed significantly lower *Flk1* expression in $Vcan^{hdf/hdf}$ yolk sac and

embryos (Fig. 3D), in agreement with reduced Flk1 staining observed in the mutants (Fig. 3A).

Neither expression of *Brachyury*, a mesoderm marker, nor *Has2*, encoding the major HA

synthase in the embryo, were altered in $Vcan^{hdf/hdf}$ yolk sacs or embryos (Fig. 3D).   Expression

of genes encoding blood lineage commitment markers *Itga2b* (CD41) and *Runx1* were similarly

reduced in the $Vcan^{hdf/hdf}$ yolk sacs and embryos, as well as erythroid and myeloid transcripts

Figure-2

Figure 2. Impaired vasculogenesis in the *Vcan^hdf/hdf* embryo and yolk sac.
(A) E8.5 *Vcan^hdf/hdf* yolk sacs are avascular yet dissected embryos appear morphologically similar to wild type.
(B) Maximum intensity projections of whole mount E8.5 wild type and *Vcan^hdf/hdf* embryos stained with anti-CD31 (red) and DAPI (blue). Boxed areas are shown at higher magnification with red channel only in the lower panels. Residual wild-type yolk sac is marked by a white dotted line (n=3 embryos from each genotype).
(C) Three-dimensional (3D) maximum intensity projections of yolk sacs stained *en face* with anti-CD31 (red) show a well-formed vascular plexus in wild type yolk sac and random CD31+ cells in *Vcan^hdf/hdf* yolk sac (n=3 yolk sacs from each genotype).
(D) Cross section of E8.5 wild type yolk sac blood islands co-stained with versican (green), HAbp (red) and CD31 (white). Versican and HA co-localize with CD31+ cells on the mesothelial aspect of blood islands (arrowheads). The blood island imaged on the left is enlarged in the bottom right-hand panel (n=3 wild type yolk sacs). VE, visceral endoderm, Bi, blood island,
(E) RNAscope in situ hybridization of E8.5 wild-type yolk sac shows expression of *Vcan* isoforms containing exon 8 (V1), but not exon 7 (V0, V2) in mesoderm adjacent to blood islands (Bi), VE, visceral endoderm.
(F) *Vcan* exon 7 and exon 8 probes both hybridize to myocardium (Mc) of E8.5 wild-type embryos. En, Endocardium.
(G) *Vcan* probes (exon 8 shown) do not hybridize to *Vcan^hdf/hdf* heart. Scale bar in D=10μm, E= 25μm, 50μm in F-G.

Hbb (β-globin), *Gata1* and *Ptprc* (CD45) (Fig. 3D). CD31 (*Pecam1*) mRNA expression was

greatly reduced in the *Vcan^hdf/hdf* yolk sacs and embryos (Fig. 3D), consistent with the observed

lack of vasculature (Fig. 2B). Methylcellulose colony formation assays demonstrated fewer

colony-forming units (CFUs) in E8.5 *Vcan^hdf/hdf* embryos and yolk sacs (Fig. 3E), in agreement

with fewer CD41-stained blood islands observed in the mutant yolk sacs.

Versican and HA are associated with the earliest Flk1$^+$ cells at gastrulation

Since Flk1$^+$ cells arise at gastrulation, we analyzed versican and HA distribution in wild-type

E7.5 mouse embryos. Versican and HA co-localized to the extraembryonic mesoderm in the

blood island ring (Fig. 4 A,B), and colocalized with Flk1$^+$ cells (Fig. 4 C,D). Correspondingly,

RNA *in situ* hybridization of serial sections from several embryos showed consistent overlap

between *Vcan*, *Has2* and *Flk1* expressing cells in the extraembryonic mesoderm (Fig. 4E, Fig.4

Supplement 1). *Vcan* exon 7-containing probes gave a weaker signal than exon 8 probes. In

contrast, *Runx1*, which identifies blood lineage-committed cells, was expressed by cells in the

extraembryonic mesoderm distinct from those expressing *Flk1*, *Vcan* and *Has2* (Fig. 4E, Fig.4

Figure-3

Figure 3. Versican and HA localize to Flk1+ cells and are essential for yolk sac blood island formation.

(**A**) E8.5 wild type and *Vcan*^{hdf/hdf} yolk sac cross-sections co-stained with versican (magenta), HAbp (red) and Flk1 (green). In wild type yolk sac versican-HA staining colocalizes with Flk1. In *Vcan*^{hdf/hdf} yolk sac, blood islands (Bi) are smaller and both HAbp staining and Flk1 staining are weak. Arrowhead: versican-HA-Flk1 co-stained patches, asterisks: cell shown in high magnification in the right-hand panels (n=4 yolk sacs from each genotype). VE, visceral endoderm.

(**B**) *En face* confocal imaging of E8.5 yolk sac versican (red) and CD41 (green) staining with the mesothelial aspect facing the objective (bottom). Surface optical sections show versican-rich loci throughout wild type yolk sac but absent in *Vcan*^{hdf/hdf} yolk sac. The mid-optical image shows versican and CD41 co-stained cells in blood islands. Versican is associated with wild-type blood islands but does not overlap with CD41. No CD41+ cells were observed in *Vcan*^{hdf/hdf} yolk sac. The high magnification image shows the distinct cell populations marked by versican and CD41. Arrowheads mark weak CD41 staining in *Vcan*^{hdf/hdf} images (n=3 yolk sacs from each genotype).

(**C**) Cross-section of an E8.5 wild type blood showing no overlap of HA (red) and CD41 (green) (n=3 yolk sacs).

 (**D**) qRT-PCR analysis of wild type and *Vcan*^{hdf/hdf} yolk sacs and embryos shows significantly lower *Flk1* expression but not *Has2* or *Brachyury* expression in *Vcan*^{hdf/hdf} mutants. CD41 *(Itga2b)* and *Runx1* expression were significantly lower in *Vcan*^{hdf/hdf} yolk sac and embryos. Blood markers *β-globin*, *Gata1* and CD45 *(Ptprc)* and the vascular endothelial marker CD31 *(Pecam1)*, were reduced in *Vcan*^{hdf/hdf} yolk sacs and embryos (n=3 yolk sacs and embryos from each genotype, error bars= S.E.M;*, $p<0.05$; **, $p<0.01$; ***, $p<0.001$).

E. Methylcellulose assay shows significantly fewer blood colony forming units (CFUs) in *Vcan*^{hdf/hdf} yolk sacs and embryos (n=3 yolk sacs and three embryos from each genotype, error bars= S.D.,*, $p<0.05$; **, $p<0.001$).

170  Supplement 1). Thus, versican-HA ECM is intimately associated with the hematoendothelial

171  lineage emerging at gastrulation.

172    Data mining of a single cell RNA sequencing (scRNA-seq) atlas of early (6.5 to 8.5 day-

173  old) mouse embryos [31] supported specific *Vcan* expression by hematoendothelial progenitors.

174  High *Vcan* expression was first detected in hematoendothelial progenitors emerging at E6.75,

175  as well as in nascent and uncommitted mesoderm (Fig. 4 Supplement 2A-C). *Vcan* was

176  expressed in *Kdr* (encoding *Flk1*)-expressing cells at E6.75, E7.0 and E7.5, spanning mouse

177  gastrulation (Fig. 4 Supplement 2A-C). *Kdr*⁺ cells also expressed *Has2* and the gene encoding

178  another HA receptor, CD44 (Fig. 4 Supplement 2B). These observations suggest a potential cell

179  autonomous role for versican in Flk1⁺ cells, bound to HA and thus to the cell surface via CD44

180  and HAS2. After E7.5 (e.g., at E 8.5)*, Vcan* was broadly expressed in mesoderm, the

181  developing brain, allantois and cardcardiomyocytes (Fig. 4 Supplement 2C) and *Kdr* expression

Figure-4

Figure 4. Versican and HA co-localize with Flk1+ cells from their origin at gastrulation.
(A) Maximum intensity projection image of an E7.5 embryo shows strong versican staining (green) in the putative blood island (BI) ring (white brackets) in the proximal extraembryonic (Exe) region. Emb, embryo
(B-E) Serial sections of E7.5 wild type embryos analyzed by immunostaining **(B-D)**, or by RNA in situ hybridization **(E)**. **(B-C)** Versican (green) and HAbp (red) colocalize in extraembryonic mesoderm corresponding to the blood island ring (white brackets and arrowheads). The blood islands on the left are shown at high magnification in **(C-D)**. Arrowheads show colocalization of versican, HAbp and Flk1 in extraembryonic mesoderm (N=4 embryos). Epc, ectoplacental cavity; Ch, chorion; Exc, exocelomic cavity; Am, amnion; Ac, amnion cavity; Ne, neural ectoderm; Me, mesoderm; Ve, visceral endoderm.
(E) *In situ* hybridization for *Vcan* exon 7, *Vcan* exon 8, *Has2*, *Flk1* and *Runx1*. *Vcan* exon 8, *Has2* and *Flk1* mRNAs have near-identical expression patterns (red) corresponding to primitive streak cells migrating toward extra-embryonic mesoderm. *Vcan* exon 7 (GAGα) shows weaker overlapping expression. *Runx1* marks committed blood cells (n=4 embryos). Scale bars in **B,E**= 100μm.

extended into endothelial cells, although hematoendothelial progenitors subsequently continued

to express *Vcan* and *Kdr* (Fig. 4 Supplement 2C). scRNA-seq data showed that HA link proteins

(*Hapln1-4*) were not expressed in hematoendothelial progenitors (Fig. 4 Supplement 3A), and

the major versican-binding link protein, HAPLN1, was absent in E7.5 embryo sections (Fig. 4

Supplement 3B). Staining for ADAMTS-cleaved versican (using anti DPEAAE) revealed no

staining in wild-type blood islands, suggesting that versican does not undergo proteolysis by

ADAMTS proteases at this time (Fig.4 Supplement 3C).

Loss of versican dramatically reduces yolk sac and embryo HA levels

In addition to reduced HA staining in *Vcan*hdf/hdf yolk sac blood islands, we observed dramatic

loss of HAbp staining throughout *Vcan*hdf/hdf embryos (Fig. 5A). Since *Has2* mRNA expression

was unaltered in *Vcan*hdf/hdf embryos and yolk sacs (Fig. 3D), reduced HABP staining may have

resulted from increased breakdown in situ, or alternatively, extraction of unliganded HA from

tissue sections during staining. Therefore, fluorophore-assisted carbohydrate electrophoresis

(FACE) [32-34] was undertaken on snap-frozen whole embryos (Fig. 5B). FACE analysis

demonstrated that versican is a major CSPG in E8.5 embryos, contributing nearly all unsulfated

(0S) CS, since both the overall level of CS and its 0S form were dramatically reduced in

*Vcan*hdf/hdf embryos (Fig. 5B). In addition, HA-FACE showed reduction of HA compared to wild-

type littermates, supporting HA loss intrinsically within the embryo rather than during sample

Figure-5

Figure 5. *Vcan* inactivation results in loss of hyaluronan.
(A) Versican (green) and HAbp (red) staining of E8.5 wild type and *Vcan*^{hdf/hdf} embryos showing severely reduced HAbp staining in *Vcan*^{hdf/hdf} embryos. Arrowheads indicate versican and HAbp-stained yolk sac blood islands.
(B) *(Top)* Fluorophore-assisted carbohydrate electrophoresis (FACE) analysis for chondroitin sulfate (CS-FACE) and hyaluronan (HA-FACE) in E8.5 littermate embryos. Versican is the major CS-proteoglycan in E8.5 embryos. HA is significantly reduced in *Vcan*^{hdf/hdf} embryos. *(Bottom)* Quantification of CS and HA FACE, normalized to DNA (n=4 embryos from each genotype, error bars= S.E.M.,**, p<0.01; ***, p<0.001; ****, p<0.0001).
(C) 1 μm thick, toluidine blue-stained Eponate 12 sections from E8.5 wild type and *Vcan*^{hdf/hdf} embryos show compaction of craniofacial mesenchymal cells (Me) and loss of cardiac jelly (CJ) between the myocardium (Mc) and endocardium (En). The red line indicates the boundary of neural epithelium (NE) with mesenchyme.
(D) E8.5 wild *Vcan*^{hdf/hdf} embryo sections stained with F-actin (red) and fibronectin (green) show stronger fibronectin staining.
(E) qRT-PCR shows increased *Tmem2* transcription in *Vcan*^{hdf/hdf} embryos and yolk sacs (N=3 embryos and yolk sacs from each genotype, error bars= S.E.M.;***, p<0.001; ****, p<0.0001). Scale bar in **A**= 200μm, 50μm and 20μm in **C**.

preparation (Fig. 5B). Consistent with the role of versican and HA as an expansile complex,

craniofacial mesenchyme and myocardium were severely compacted in *Vcan*^{hdf/hdf} embryos (Fig.

5C). We also asked whether the absence of versican affected fibronectin, a major embryonic

ECM component that is essential for angiogenesis and observed a dramatic increase both in

fibronectin staining and mRNA in *Vcan*^{hdf/hdf} embryos (Fig. 5D, Fig. 5 Supplement 1A,B), which

is presently unexplained. In combination with unaltered *Has2* expression, these findings suggest

increased HA turnover in *Vcan*^{hdf/hdf} embryos. TMEM2 was recently identified as a major

extracellular hyaluronidase [35-37] and qRT-PCR revealed significantly raised*Tmem2*

expression in *Vcan*^{hdf/hdf} yolk sac and embryos (Fig. 5E). indicating that hyaluronan catabolism

was activated in the absence of versican.

HA deficiency impairs vasculogenesis

Since HA was lost in the absence of versican, we inquired if the converse were true, and

whether HA-deficiency indeed led to defective yolk sac vasculogenesis, as previously

suggested by morphology of *Has2* germline mutants [26]. *Has2*^{−/−} embryos die by E 9.5 due to

heart defects [26] whereas *Has1*^{−/−} and *Has3*^{−/−} mice are both viable and do not exhibit

noticeable embryonic phenotypes [38, 39]. To completely deplete HA from developing embryos,

we generated triple compound embryos lacking all three *Has* genes (*Has1*[−/−]*;Has2*[−/−]*;Has3*[−/−];

referred to as *Has1-3*[TKO]). We generated HA-deficient mouse embryos by intercrossing *Has1+3*[-

[/-]*; Has2*[+/-] mice. At E9.5, *Has1-3*[TKO] embryos, like *Vcan*[hdf/hdf] and *Has2*-epiblast deleted embryos

[26] showed loss of yolk sac vasculature (Fig. 6A). HA staining in the *Has1-3*[TKO] yolk sac and

embryos was absent, as expected and the sections also showed a near-complete lack of

versican staining (Fig. 6B,C). In contrast, immunostaining for cleaved versican (anti-DPEAAE)

showed robust and widespread versican processing in the absence of HA (Fig. 6 B,D). Epiblast-

specific *Has2* conditional deletion (*Sox2*Cre;*Has2*[Fl/Fl]) showed a similar loss of yolk sac

vasculature and cardiac defects (Fig.6 Supplement 1A). RNA in situ hybridization showed a low

level *Has2* expression in these embryos and slightly reduced *Vcan* transcript levels (Fig. 6

Supplement 1B). Phenocopying *Has1-3*[TKO] embryos, the *Sox2*Cre; *Has2*[Fl/Fl] embryos and yolk

sacs also showed loss of both HA and versican staining and increased ADAMTS-mediated

versican cleavage (Fig. 6 Supplement 1C). Taken together, these data suggest that HA may

protect versican from ADAMTS-mediated degradation.

Embryoid bodies formed by *Vcan*-null ES cells have impaired angiogenesis and

hematopoiesis.

Since the dramatic loss of vasculature could result from perturbed hemodynamics due to cardiac

defects in *Vcan*[hdf/hdf] embryos, we undertook an orthogonal (in vitro) approach for evaluation of

versican in vasculogenesis and hematopoiesis. We used gene editing by CRISPR/Cas9 [40, 41]

to introduce *Vcan*-null mutations in R1 mouse embryonic stem cells (mESC) [42]. *Vcan* exon 2

(containing the start codon) and exon 3 (start of the G1 domain) were targeted independently to

obtain mESC clones D8 and F9 respectively, each with defined frameshift mutations that

generated null alleles (Fig. 7A). Since exons 2 and 3 are included in all *Vcan* splice isoforms, no

versican was produced, as shown by western blot with anti-GAGβ antibody (Fig. 7B). The mutated

clones retained normal expression of *Oct4, Sox2, Nanog* and *C-myc* (Fig. 7C) indicating unaltered

pluripotency, validating their suitability for *in vitro* differentiation. *Vcan*-null and wild-type ES cells

Figure-6

Figure 6. Versican is severely reduced in *Has1+2+3* null embryos.
(A) E9.5 *Has1+3* null embryos (control) and *Has1+2+3* null embryos showing the dilated pericardial sac in the latter (arrowhead) similar to *Vcan$^{hdf/hdf}$*.
(B) *Has1+3* null yolk sac (control) and *Has1+2+3* null yolk sac co-stained with HAbp (green) and versican or cleaved versican (DPEAAE) (red) show loss of both HA and versican although cleaved versican (DPEAAE, red, bottom) is present in *Has1+2+3* null yolk sac.
(C-D) *Has1+3* null yolk sac (control) and *Has1+2+3* null embryos co-stained with HAbp and versican **(C)** or cleaved versican (DPEAAE) **(D)**, show loss of HA and versican staining and weaker DPEAAE staining in the *Has1+2+3* null embryos. Scale bars = 200μm in in **C**,**D**.

were allowed to form embryoid bodies (EBs), in which random differentiation into various lineages

occurs. RT-qPCR analysis of *Has2* showed no change in *Vcan*-null EBs (Fig. 7 Supplement 1A)

while the mesoderm differentiation marker *Brachyury/T* showed a significant decrease in the F9

but not the D8 *Vcan*-null EBs (Fig. 7 Supplement 1B). Similar to *Vcan$^{hdf/hdf}$* embryos, *Vcan* null

embryoid bodies showed reduced *Kdr* expression (Fig. 7 Supplement 1C), and the blood lineage

commitment markers *Itga2b* (CD41) and *Runx1* (Fig. 7 Supplement 1D), as well as of

differentiated-blood and vascular endothelial markers (Fig. 7 Supplement 1E, F). Since EB

differentiation is random and independent of a closed pulsatile circulation, the data from embryoid

bodies and the mutant mice together suggests that versican acts directly on Flk1^{+}

hematoendothelial progenitors rather than indirectly *via* abnormal cardiac development.

Additionally, we separately evaluated the endothelial and hematopoietic potential of *Vcan*-

null ESC. Sprouting angiogenesis was induced by culturing 4-day differentiated EBs in 3-

dimensional collagen I gels and treatment with VEGF-A$_{165}$ for 12 days [43]. Robust angiogenic

sprouting identified by CD31 and smooth muscle α-actin staining was seen in wild-type EBs, but

not in the majority of *Vcan*-null EBs (Fig. 7 D,E). The few *Vcan*-null EBs with sprouting had fewer

sprouts per EB, and these were significantly shorter than wild-type sprouts (Fig. 7 F,G). When 10-

day old EBs were disaggregated and the cells were plated in a 3-dimensional methylcellulose

matrix containing a complete set of blood differentiation cytokines, significantly fewer blood

colonies were formed by *Vcan* null EBs (Fig. 7 H).

Figure-7

Figure 7. Embryoid bodies (EBs) from *Vcan*-null mouse embryonic stem cells (mESCs) are unresponsive to VEGF and form few blood colonies.

(A) *Vcan* locus showing targeting by independent guide RNAs (gRNAs) targeting exons 2 and 3. Scissors indicate the Cas9 cleavage site 3 bp from the protospacer adjacent motif (PAM, orange lettering). An exon 2 mutant mESC clone (D8) had a homozygous 1 bp insertion (bold red letters) and an exon 3 mutant clone (F9) had heterozygous targeting: deletion of T/A in one allele and a 19-bp deletion in the other resulting in frame-shifts.

(B) Western blot of 10-day differentiated EBs shows no versican in EBs derived from clones D8 and F9.

(C) qRT-PCR analysis showing unaffected expression of pluripotency markers *Oct4*, *Sox2*, *Nanog* and *C-myc* in D8 and F9 (n=3 independent batches of EBs from each genotype, error bars= S.E.).

(D) 4-day-old *Vcan*-null embryoid bodies embedded in collagen I and treated with VEGF$_{165}$ for 12 days lack vascular sprouting identified by CD31 (green) or α-SMA (red) immunostaining (n=3 independent batches of EBs from each genotype).

(E) D8 and F9 EBs show significantly fewer vascular sprouts (n=3 independent batches of EBs from each genotype, error bars= S.E. **, $p<0.01$; ***, $p<0.001$).

(F) Fewer sprouts/EB were seen in D8 and F9 EBs. (n=3 independent batches of EBs from each genotype, error bars= S.D. ***, $p<0.001$; ****, $p<0.0001$).

(G) Significantly shorter sprouts in *Vcan* null lines (n=3 independent batches of EBs from each genotype, error bars= S.D. ****, $p<0.0001$).

(H) Methylcellulose colony assay using dissociated 10-day old EBs shows significantly fewer colonies in *Vcan* null EBs. (N=3 independent batches of EBs from each genotype, error bars= S.D. **, $p<0.01$). Scale bar in D is 200 μm.

The versican-HA complex sequesters growth factors essential for vasculogenesis and

hematopoiesis.

These observations raised questions about the underlying mechanisms by which versican-HA

matrix would support survival and expansion of Flk1[+] cells. An *ADAMTS9* null RPE1 cell line

(D12) lacks ADAMTS9, a key versican-degrading protease [22] and thus demonstrates

constitutively strong versican staining in ECM (Figure 8A). HABP staining of these cells

demonstrated stronger HA staining than in wild type RPE1 cells, with formation of long HA

cables decorated with versican (Fig. 8A, Fig. 8 Supplement 1A,B); such cables are well-known

to occur in 25 mM glucose-containing medium [44] [45]. qRT-PCR revealed *HAS2* & HAS3

expression were reduced in D12 vs. wt (Fig. 8B) with no change in *HAS2* expression but drastic

reduction of *TMEM2* expression was observed in the RPE1-D12 cells (Fig. 8B). Intriguingly,

upon siRNA-mediated versican depletion [46, 47], RPE1-D12 cells showed dramatically

Figure-8

Figure 8. Versican regulates HA abundance and cable formation and sequesters VEGf and Ihh *via* chondroitin sulfate (CS) chains in a cell culture model.
(A) Increased versican (red) and HAbp staining (green) in *ADAMTS9*-deficient (D12) RPE-1 cells. Versican and HA co-stained cables are formed in D12 cells.
(B) qRT-PCR showing no change in *HAS1-3* or *CD44* but dramatically reduced *TMEM2* expression in D12 cells. (n=3 independent RNA extractions, error bars= S.E.M., *, $p<0.05$; ****, $p<0.0001$).
(C) *VCAN* knockdown reduces both versican (red) and HA staining in D12 cells.
(D) Recombinant $VEGF_{165}$ or Ihh (green) co-staining with versican (red) in control siRNA-transfected D12 cells is lost in *VCAN* siRNA-transfected D12 cells.
(E) The schematic illustrates chondroitinase ABC removal of versican GAG chains and the experimental timeline used.
(F) Fluorescence microscope of VEGF and Ihh with versican, with or without chondroitinase ABC treatment prior to addition of recombinant $VEGF_{165}$ or Ihh (green) shows reduced binding of both growth factors after CS-chain removal, but does not affect versican core protein staining (red).

reduced versican and HA staining and absence of HA/versican cables (Fig. 8C, Fig. 8

Supplement 2A). Conversely, qRT-PCR showed increased *TMEM2* expression after *VCAN*

knockdown (Fig. 8 Supplement 2B), suggesting regulation of the HA-depolymerase *TMEM2* by

the level of versican as a possible mechanism of concurrent HA loss.

Next, to test if versican can sequester growth factors known to regulate vasculogenesis,

we treated wild-type and RPE1-D12 cells with increasing concentrations of recombinant

$VEGF_{165}$ and Ihh. Each showed dose-dependent punctate staining in RPE1-D12 monolayers

which co-localized with the versican-HA cables, imaged by super-resolution fluorescent

microscopy. (Fig. 8D, Fig. 8 Supplement 3A-B, Fig. 8 Supplement 4A-B). Upon *VCAN* depletion

in RPE1-D12 cells both $VEGF_{165}$ and Ihh staining were significantly decreased (Fig. 8D).

Enzymatic removal of proteoglycan chondroitin sulfate chains prior to addition of recombinant

$VEGF_{165}$ and Ihh to D12 cultures also dramatically reduced $VEGF_{165}$ and Ihh staining intensity

without affecting versican core protein staining (Fig. 8E,F), suggesting that versican CS chains

mediated VEGF and Ihh sequestration.

Discussion

Although versican is widely expressed during organogenesis and in adult tissues [14], the

chronology of its expression in early vascular development undertaken here and independently

derived from recently published single cell RNA-seq data, shows specific association with Flk1$^+$

cells from their earliest origin in the gastrulating embryo until establishment of yolk sac

vasculature and primitive hematopoiesis. Analysis of *Vcan* mutant mice *in vivo* and *Vcan*-null

embryoid bodies *in vitro* suggest that versican has an indispensable role in vasculogenesis and

primitive hematopoiesis. Taken together, RNA *in situ* hybridization data, scRNA-seq analysis of

Flk1$^+$ cells and immunostaining of embryos suggests that versican is a product of Flk1$^+$ cells that

associates with them via HA and CD44 to form a crucial, possibly cell-autonomously acting

ECM. Furthermore, in vitro analysis undertaken in REP1-D12 cells for their high versican levels,

showed dose-dependent binding of VEGF and Ihh to the CS-chains of versican.

Two novel, unexpected findings of this study were the substantial loss of embryonic and

extraembryonic HA in the absence of versican and reciprocally, of versican in the absence of

HA. Also unexpected was the transcriptional effect of versican expression on TMEM2. As in

Vcan$^{hdf/hdf}$ yolk sac, avascularity was previously noted in *Has2* null yolk sac [26], although it was

not studied further. *Has2* and *Vcan* null cardiac defects are similar, which suggested that an

obligate versican-HA complex in cardiac jelly and endocardial cushions was required for cardiac

morphogenesis [24-26]. The present work similarly demonstrates that a versican-HA complex,

rather than versican alone, is necessary for vasculogenesis and primitive hematopoiesis (Fig.

S12). HA-versican aggregates have a net negative charge, and exert swelling pressure *via*

absorption of water (the Gibbs-Donnan effect) [48]. As shown here and previously, embryonic

tissues compact in their absence [24, 49]. Although versican-deficient yolk sac showed

extensive structural disorganization at E9.5, it is clear that the association and effect of versican

vis-à-vis Flk1$^+$ cells occurs much earlier, i.e., shortly after gastrulation. Reduced

vasculogenesis, angiogenesis and hematopoiesis in *Vcan*-null embryoid bodies supports a

direct, local role of versican on hematoendothelial progenitors than a secondary effect of cardiac

anomalies on vasculogenesis, or of yolk sac disorganization on blood island formation. We

therefore conclude that versican-HA aggregates act directly and as early as E6.75 within the

microenvironment of Flk1$^+$ cells.

HA, which is extruded directly from HA synthases on the plasma membrane, is decorated

with versican, and is retained at the cell surface via HA synthases or CD44 and other HA receptors

[50]. *Cd44* is coordinately expressed with *Vcan* and *Has2* by Flk1$^+$ cells during early

embryogenesis. Nevertheless, *Cd44* null mice survive and are not known to have vasculogenesis

or primitive erythropoiesis defects [51], suggesting that HAS2 acting as a *de facto* receptor, may

participate in forming a cell-associated versican-HA complex. The versican-HA pericellular matrix

has a well-established impact on different cells. We previously found that the amount of versican

in the pericellular matrix in fibroblasts and vascular and myometrial smooth muscle cells [9, 21,

52] was a determinant of phenotype modulation. Versican is anti-adhesive [9, 52], which may

allow Flk1$^+$ cells to migrate and proliferate efficiently after their emergence at gastrulation. Another

possible role of versican-HA is sequestration of vasculogenic factors such as VEGF-A. Versican

binding to VEGF-A and Ihh *via* its CS-chains (this study) or G3 domain [19], may generate high

concentrations around Flk1$^+$ cells. Versican is associated with Flk1$^+$CD41$^-$ cells at their origin, but

not with the Flk1$^-$CD41$^+$ yolk sac cells subsequently, suggesting that the versican-HA pericellular

ECM specifically regulates the fate of Flk1$^+$ CD41$^-$ cells. We conclude that as the dominant CSPG

in the embryo, specifically localized to the vicinity of Flk1$^+$ cells, versican may sequester essential

factors such as VEGF-A and Ihh to provide a high local concentration that sustains the Flk1$^+$

population.

Although lack of versican led to higher *Fn1* expression, *Vcan*$^{hdf/hdf}$ defects are unlikely to

result from excess fibronectin, because previous work has shown that deficiency, not excess of

fibronectin or the fibronectin receptor subunit α5 integrin results in reduced embryo and yolk sac

vascularity [27, 53]. Since versican binds fibronectin through the G3 domain [19], we conclude

that the three major components of the provisional embryonic ECM (versican, HA and fibronectin)

are each crucial for establishment of the first blood vessels in the embryo. In the absence of

versican, the HA catabolism rate *via* TMEM2 likely exceeds the HA synthesis rate since *Has2*

mRNA levels were relatively unchanged. It was previously noted that HA deposition was reduced

to 85% of normal in fibroblasts taken from a mouse hypomorphic *Vcan* mutant (*Vcan*$^{\Delta3/\Delta3}$) having

75% reduction of versican [54]. In association with reduced HA deposition, *Vcan*$^{\Delta3/\Delta3}$ fibroblasts

had accelerated senescence, suggesting a possible mechanism for the lack of Flk1$^+$ cells in

Vcan$^{hdf/hdf}$ yolk sac, although this study did not specifically address the fate of Flk1+ cells.

We observed stronger association of exon 8-containing *Vcan* transcripts than exon 7-

containing transcripts with blood islands at gastrulation and exon 7 transcripts were absent at

E8.5. Autosomal dominant splice site mutations affecting exon 8 (leading to its exclusion) and

exon 8 deletions in humans cause Wagner syndrome [55-57], which is characterized by impaired

vision and defects of the ocular vitreous and retina, but lacks consistent extra-ocular

manifestations. Exon 7 inactivation in mice led to specific neural anomalies and subtle cardiac

anomalies [58, 59]. Thus, neither individual exon mutation in mice or humans is associated with

embryonic lethality, defective vasculogenesis or impaired hematopoiesis. We conclude that both

exon 7 and exon 8-containing transcripts, possibly included in V0, the transcript containing both

exons, are required for vasculogenesis and hematopoiesis.

The GAGβ domain encoded by exon 8 has a unique N-terminal sequence, which is

cleaved by ADAMTS proteases at the E^{441}-A^{442} peptide bond in several contexts, notably cardiac

valve development, interdigital web regression, umbilical cord development, neural tube and

palate closure and myometrial activation [9, 22, 60]. The resulting N-terminal versican V1

fragment, G1-DPEAAE^{441}, named versikine, is bioactive in interdigital web regression and

myeloma growth [61, 62]. *Vcan* knock-in mouse mutants in which E^{441}-A^{442} was mutated to render

it uncleavable, complete gestation and havenormal yolk sac avascularity, suggesting that E^{441}-

A^{442} cleavage is not involved in hematoendothelial development ([63] and Nandadasa et al.,

manuscript in preparation).

With the added role in vasculogenesis and hematopoiesis elucidated here, it is not an

exaggeration to state that versican is crucial for development of the entire circulatory system.

Indeed, versican-HA matrix may also have a broad role in formation of vasculature by

angiogenesis in a variety of physiological and disease settings. Recent work found that syngeneic

B16F10 tumors in adult $Vcan^{hdf/+}$ mice had significantly impaired angiogenesis and reduced

growth [64]. Relevant to the overlap of versican-HA with Flk1 expression, a recent study utilizing

the same cancer model in $Flk1+/-$ mice also found reduced angiogenesis during tumor growth

[65]. Furthermore, VEGF binding of the versican-HA ECM which is quantitatively increased in the

presence of high glucose, may be relevant to diabetic retinopathy, a common complication of

diabetes where the increased activity of VEGF is well-established and indeed, a current target of

treatment [66].

Methods

Mice

$Vcan^{Tg(Hoxa1)1Chm}$ ($Vcan^{hdf}$) mice [25] (Figure-1 Supplement 1) were obtained under a material

transfer agreement from Roche. Generation of $Has1^{-/-};Has3^{-/-}$ double knockout mice was

described previously [67]. A $Has2$-null allele ($Has2^-$) was created from a $Has2^{flox}$ allele [68] by

crossing $Has2^{flox/flox}$ mice with the germline deleter $Meox2$-Cre mice [69]. All of these mouse

lines were backcrossed to C57BL/6J for more than 10 generations. $Has1^{-/-};Has2^{+/-};Has3^{-/-}$ mice

were bred from these mutant mice. Triple Has knockout embryos were generated by crossing

$Has1^{-/-};Has2^{+/-};Has3^{-/-}$ female and male mice. In a second approach to generate HA-deficient

mouse embryos, $Has2^{Fl/Fl}; Sox2Cre^{tg}$ mice were generated by crossing $Has2^{Fl/Fl}$ males with

$Has2^{+/Fl}; Sox2Cre^{tg}$ females. Mouse experiments were conducted with IACUC approval

(Cleveland clinic protocols 2015:1530 and 2018:2045). Mice were maintained in a fixed light-

dark cycle with food and water *ad libitum*. For genotyping E8.5 embryos with intact yolk sacs,

the allantois was dissected out and lysed in 10 μl DirectPCR (Tail) digest reagent (Qiagen,

catalog no. 102-T) supplemented with 1 μl of proteinase K overnight at 55°C. Tails were used to

genotype E9.5 embryos. *Vcan^hdf* and wild-types were identified with a specific genotyping

strategy based on the genetic interruption (Figure-1 S1). Details of the *Has1-3* mutant mouse

genotyping is provided in the Supplemental File. *Vcan^{hdf/hdf}* embryos were compared to wild-type

littermates in all experiments.

Mouse embryonic stem cell (mESC) culture.

R1 mESC [42] were cultured on 0.3% type B gelatin (Sigma-Aldrich, catalog no. G9382) coated

60 mm cell culture plates in Iscove's Modified Dulbecco's Medium (IMDM) containing 4mM L-

glutamine and 1mM sodium pyruvate, supplemented with 20% fetal bovine serum (Hyclone,

catalog no. SH30071), 0.1 mM cell culture grade 2-mercaptoethanol (Gibco, Life Technologies,

catalog no. 21985), 0.1mM nonessential amino acids (Gibco, Life Technologies, catalog no.

11140-050), 50 μg/mL penicillin/streptomycin and 1×10^6 units/mL leukemia inhibitory factor (LIF;

ESGRO, EMD Millipore, catalog no. ESG1106) in a humidified 5% CO_2, 37°C environment. mESC

were maintained at 60-80% confluence, with daily medium change and passaged every other day

in a 1:5 split.

CRISPR/Cas9 targeting of mESC *Vcan*

2.5 μg of CRISPR/Cas9 plasmids in the U6gRNA-Cas9-2A-GFP vector, targeting *Vcan* exon 2 or

exon 3 (Sigma-Aldrich, target IDs MM0000080027 and MM0000080028) were transfected into

R1 mESCs at 60-80% confluence in 6-well plates coated with 0.3% gelatin, using FuGene 6

(Promega, catalog no. E2691). 24 hrs post-transfection, individual GFP+ mESCs were sorted

into 96-well plates coated with 0.3% gelatin using a FACSAria-II cell sorter (BD Biosciences).

Fast-growing wells (containing >1000 cells/well) were trypsinized and expanded to 24-well cell

culture plates after 7-10 days. Culture medium was replaced daily. Genomic DNA from clones

was isolated using DirectPCR (Tail) reagent (Viagen, catalog no. 102-T) and *Vcan* exon 2 and

exon 3 were amplified using Phusion *Taq* (NEB, catalog no. F530L) (see SI). Amplicons were

excised from 2% agarose gels, purified using the QIAquick Gel Extraction kit (QIAGEN, catalog

no. 28704) and cloned into pCR-Blunt II-TOPO vector using the Zero blunt PCR cloning kit (Life

Technologies, Invitrogen, catalog no. K2800-40). Plasmid DNA was harvested from bacterial

colonies for Sanger-sequencing to determine the precise mutations. Two independent *Vcan*-null

mESC clones, D8 and F9, were established. Western blotting and immunostaining of 10-day old

embryoid bodies from these lines (see below) confirmed *Vcan* inactivation. Embryoid body

formation and induction of vascular sprouts is described in the **Online Supplement**.

Immunostaining and fluorescence microscopy

Immunostaining of E 9.5 and E 8.5 yolk sac was carried out on 30 μm thick vibratome sections

[22] or paraffin-embedded 7 μm sections. Immunostaining of collagen-embedded embryoid

bodies were carried out in 4-chamber cell culture slides (Fisher Scientific, catalog no. 354114).

Confocal microscopy images of whole-mount mouse embryos and sections were acquired using

a Leica TCS SP5 II multiphoton confocal microscope equipped with a 25X water immersion

objective (Leica Microsystems, Wetzlar, Germany). For 3D-projection of whole-mount Z-stacks,

the Volocity 3D imaging software was used (version 6.3, PerkinElmer, Inc., Waltham, MA) in

maximum intensity projection method.

Methylcellulose colony formation assay

Single cell suspensions of E8.5 embryos, yolk sacs or day-10 embryoid bodies were generated

by incubation with trypsin for 10 minutes followed by disaggregation by pipetting with a 200 μL

pipette tip until complete. 50,000 cells from each experimental group were transferred to a single

35 mm culture dish containing 1 mL of MethoCult GF M3534 culture medium (Methylcellulose

medium with recombinant cytokines for mouse cells, Stem Cell Technologies, Vancouver, CA,

catalog no. 03534) using a 3 mL syringe and 16-gauge needle, following the manufacturer's

protocol. Triplicate cultures from each genotype were incubated for 14 days in a humidified, 5%

CO_2, 37°C cell culture incubator. Blood colonies were counted using an inverted microscope.

Aggregates with >50 cells were considered a colony-forming unit (CFU).

Additional details of reagents and procedures including primary antibodies used,

RNAscope in situ hybridization, western blotting, transmission electron microscopy, and

fluorophore-assisted carbohydrate electrophoresis (FACE) are described in the **Online**

Supplement.

Acknowledgments
453

This work was supported by the NIH-NHLBI Program of Excellence in Glycosciences award

HL107147 (to S.S.A., R.J.M.,), NIH RF1 AG057579 (to Y.Y.) and by the Allen Distinguished

Investigator Program, through support made by The Paul G. Allen Frontiers Group and the

American Heart Association (to S.S.A), the David and Lindsay Morgenthaler Postdoctoral

Fellowship and the Mark Lauer pediatric research grant (to S.N.). Purchase of the Leica

SP8 confocal microscope was supported by NIH SIG grant 1S10RR026820-01. We thank Dr.

David LePage and Dr. Ron Conlon at the CWRU transgenic core for R1 mES cells, Eric Schultz

and Joseph Gerow of the LRI Flow Cytometry Core for mES cell sorting, Valbona Cali for FACE

assays, Dr. Judy Drazba and Mei Yin of the LRI Imaging Core for guidance with confocal and

electron microscopy and the Apte laboratory members for valuable discussions.

Author contributions

S.N. and A.O conducted experiments. S.N., R.J.M., and S.S.A. conceived experiments and

interpreted the data. S.N. and S.S.A. wrote the paper. All authors read and edited the paper.

Declaration of interests

The authors declare no competing interests
469

References:

1. Qiu, J. and K.K. Hirschi, *Endothelial Cell Development and Its Application to Regenerative Medicine.* Circ Res, 2019. **125**(4): p. 489-501.
2. Risau, W. and I. Flamme, *Vasculogenesis.* Annu Rev Cell Dev Biol, 1995. **11**: p. 73-91.
3. Lugus, J.J., et al., *Both primitive and definitive blood cells are derived from Flk-1+ mesoderm.* Blood, 2009. **113**(3): p. 563-6.
4. Ferkowicz, M.J. and M.C. Yoder, *Blood island formation: longstanding observations and modern interpretations.* Exp Hematol, 2005. **33**(9): p. 1041-7.
5. Dzierzak, E. and N.A. Speck, *Of lineage and legacy: the development of mammalian hematopoietic stem cells.* Nat Immunol, 2008. **9**(2): p. 129-36.
6. Red-Horse, K., et al., *Endothelium-microenvironment interactions in the developing embryo and in the adult.* Dev Cell, 2007. **12**(2): p. 181-94.
7. Lozito, T.P., et al., *Human mesenchymal stem cells express vascular cell phenotypes upon interaction with endothelial cell matrix.* J Cell Biochem, 2009. **107**(4): p. 714-22.
8. Gu, Y., et al., *Steel factor controls primordial germ cell survival and motility from the time of their specification in the allantois, and provides a continuous niche throughout their migration.* Development, 2009. **136**(8): p. 1295-303.
9. Mead, T.J., et al., *ADAMTS9-Regulated Pericellular Matrix Dynamics Governs Focal Adhesion-Dependent Smooth Muscle Differentiation.* Cell Rep, 2018. **23**(2): p. 485-498.
10. Perris, R., et al., *Inhibitory effects of PG-H/aggrecan and PG-M/versican on avian neural crest cell migration.* Faseb J, 1996. **10**(2): p. 293-301.
11. Yamagata, M., et al., *Regulation of cell-substrate adhesion by proteoglycans immobilized on extracellular substrates.* J Biol Chem, 1989. **264**(14): p. 8012-8.
12. van Wijk, X.M. and T.H. van Kuppevelt, *Heparan sulfate in angiogenesis: a target for therapy.* Angiogenesis, 2014. **17**(3): p. 443-62.
13. Le Jan, S., et al., *Functional overlap between chondroitin and heparan sulfate proteoglycans during VEGF-induced sprouting angiogenesis.* Arterioscler Thromb Vasc Biol, 2012. **32**(5): p. 1255-63.
14. Bode-Lesniewska, B., et al., *Distribution of the large aggregating proteoglycan versican in adult human tissues.* J Histochem Cytochem, 1996. **44**(4): p. 303-12.
15. Kimata, K., et al., *A large chondroitin sulfate proteoglycan (PG-M) synthesized before chondrogenesis in the limb bud of chick embryo.* J Biol Chem, 1986. **261**(29): p. 13517-25.
16. Shinomura, T., et al., *cDNA cloning of PG-M, a large chondroitin sulfate proteoglycan expressed during chondrogenesis in chick limb buds. Alternative spliced multiforms of PG-M and their relationships to versican.* J Biol Chem, 1993. **268**(19): p. 14461-9.
17. Zimmermann, D.R. and E. Ruoslahti, *Multiple domains of the large fibroblast proteoglycan, versican.* Embo J, 1989. **8**(10): p. 2975-81.
18. Wu, Y.J., et al., *The interaction of versican with its binding partners.* Cell Res, 2005. **15**(7): p. 483-94.
19. Zheng, P.S., et al., *Versican/PG-M G3 domain promotes tumor growth and angiogenesis.* FASEB J, 2004. **18**(6): p. 754-6.
20. Slevin, M., et al., *Hyaluronan-mediated angiogenesis in vascular disease: uncovering RHAMM and CD44 receptor signaling pathways.* Matrix Biol, 2007. **26**(1): p. 58-68.
21. Hattori, N., et al., *Pericellular versican regulates the fibroblast-myofibroblast transition: a role for ADAMTS5 protease-mediated proteolysis.* J Biol Chem, 2011. **286**(39): p. 34298-310.
22. Nandadasa, S., C.M. Nelson, and S.S. Apte, *ADAMTS9-Mediated Extracellular Matrix Dynamics Regulates Umbilical Cord Vascular Smooth Muscle Differentiation and Rotation.* Cell Rep, 2015. **11**(10): p. 1519-28.

23. Dours-Zimmermann, M.T. and D.R. Zimmermann, *A novel glycosaminoglycan*
attachment domain identified in two alternative splice variants of human versican. J Biol
Chem, 1994. **269**(52): p. 32992-8.
24. Mjaatvedt, C.H., et al., *The Cspg2 gene, disrupted in the hdf mutant, is required for right*
cardiac chamber and endocardial cushion formation. Dev Biol, 1998. **202**(1): p. 56-66.
25. Yamamura, H., et al., *A heart segmental defect in the anterior-posterior axis of a*
transgenic mutant mouse. Dev Biol, 1997. **186**(1): p. 58-72.
26. Camenisch, T.D., et al., *Disruption of hyaluronan synthase-2 abrogates normal cardiac*
morphogenesis and hyaluronan-mediated transformation of epithelium to mesenchyme.
J Clin Invest, 2000. **106**(3): p. 349-60.
27. Francis, S.E., et al., *Central roles of alpha5beta1 integrin and fibronectin in vascular*
development in mouse embryos and embryoid bodies. Arterioscler Thromb Vasc Biol,
2002. **22**(6): p. 927-33.
28. Hunziker, E.B., W. Herrmann, and R.K. Schenk, *Ruthenium hexammine trichloride*
(RHT)-mediated interaction between plasmalemmal components and pericellular matrix
proteoglycans is responsible for the preservation of chondrocytic plasma membranes in
situ during cartilage fixation. J Histochem Cytochem, 1983. **31**(6): p. 717-27.
29. Shalaby, F., et al., *A requirement for Flk1 in primitive and definitive hematopoiesis and*
vasculogenesis. Cell, 1997. **89**(6): p. 981-90.
30. Ferkowicz, M.J., et al., *CD41 expression defines the onset of primitive and definitive*
hematopoiesis in the murine embryo. Development, 2003. **130**(18): p. 4393-403.
31. Pijuan-Sala, B., et al., *A single-cell molecular map of mouse gastrulation and early*
organogenesis. Nature, 2019. **566**(7745): p. 490-495.
32. Calabro, A., et al., *Microanalysis of enzyme digests of hyaluronan and*
chondroitin/dermatan sulfate by fluorophore-assisted carbohydrate electrophoresis
(FACE). Glycobiology, 2000. **10**(3): p. 273-81.
33. Calabro, A., V.C. Hascall, and R.J. Midura, *Adaptation of FACE methodology for*
microanalysis of total hyaluronan and chondroitin sulfate composition from cartilage.
Glycobiology, 2000. **10**(3): p. 283-93.
34. Midura, R.J., et al., *Quantification of hyaluronan (HA) using a simplified fluorophore-*
assisted carbohydrate electrophoresis (FACE) procedure. Methods Cell Biol, 2018. **143**:
p. 297-316.
35. De Angelis, J.E., et al., *Tmem2 Regulates Embryonic Vegf Signaling by Controlling*
Hyaluronic Acid Turnover. Dev Cell, 2017. **40**(2): p. 123-136.
36. Schinzel, R.T., et al., *The Hyaluronidase, TMEM2, Promotes ER Homeostasis and*
Longevity Independent of the UPR(ER). Cell, 2019. **179**(6): p. 1306-1318.e18.
37. Yamamoto, H., et al., *A mammalian homolog of the zebrafish transmembrane protein 2*
(TMEM2) is the long-sought-after cell-surface hyaluronidase. J Biol Chem, 2017.
292(18): p. 7304-7313.
38. Bai, K.J., et al., *The role of hyaluronan synthase 3 in ventilator-induced lung injury.* Am J
Respir Crit Care Med, 2005. **172**(1): p. 92-8.
39. Mack, J.A., et al., *Enhanced inflammation and accelerated wound closure following*
tetraphorbol ester application or full-thickness wounding in mice lacking hyaluronan
synthases Has1 and Has3. J Invest Dermatol, 2012. **132**(1): p. 198-207.
40. Doudna, J.A. and E. Charpentier, *Genome editing. The new frontier of genome*
engineering with CRISPR-Cas9. Science, 2014. **346**(6213): p. 1258096.
41. Jinek, M., et al., *A programmable dual-RNA-guided DNA endonuclease in adaptive*
bacterial immunity. Science, 2012. **337**(6096): p. 816-21.
42. Nagy, A., et al., *Derivation of completely cell culture-derived mice from early-passage*
embryonic stem cells. Proc Natl Acad Sci U S A, 1993. **90**(18): p. 8424-8.

43.   Baker, M., et al., *Use of the mouse aortic ring assay to study angiogenesis.* Nat Protoc,
572   2012. **7**(1): p. 89-104.
573   44.   Wang, A. and V.C. Hascall, *Hyaluronan structures synthesized by rat mesangial cells in
574   response to hyperglycemia induce monocyte adhesion.* J Biol Chem, 2004. **279**(11): p.
575   10279-85.
576   45.   Selbi, W., et al., *Characterization of hyaluronan cable structure and function in renal
577   proximal tubular epithelial cells.* Kidney Int, 2006. **70**(7): p. 1287-95.
578   46.   Gueye, N.A., et al., *Versican Proteolysis by ADAMTS Proteases and Its Influence on
579   Sex Steroid Receptor Expression in Uterine Leiomyoma.* J Clin Endocrinol Metab, 2017.
580   **102**(5): p. 1631-1641.
581   47.   Keire, P.A., et al., *Inhibition of versican expression by siRNA facilitates tropoelastin
582   synthesis and elastic fiber formation by human SK-LMS-1 leiomyosarcoma smooth
583   muscle cells in vitro and in vivo.* Matrix Biol, 2016. **50**: p. 67-81.
584   48.   Buschmann, M.D. and A.J. Grodzinsky, *A molecular model of proteoglycan-associated
585   electrostatic forces in cartilage mechanics.* J Biomech Eng, 1995. **117**(2): p. 179-92.
586   49.   Szabo, A., et al., *In vivo confinement promotes collective migration of neural crest cells.*
587   J Cell Biol, 2016. **213**(5): p. 543-55.
588   50.   Evanko, S.P., et al., *Hyaluronan-dependent pericellular matrix.* Adv Drug Deliv Rev,
589   2007. **59**(13): p. 1351-65.
590   51.   Protin, U., et al., *CD44-deficient mice develop normally with changes in subpopulations
591   and recirculation of lymphocyte subsets.* J Immunol, 1999. **163**(9): p. 4917-23.
592   52.   Evanko, S.P., J.C. Angello, and T.N. Wight, *Formation of hyaluronan- and versican-rich
593   pericellular matrix is required for proliferation and migration of vascular smooth muscle
594   cells.* Arterioscler Thromb Vasc Biol, 1999. **19**(4): p. 1004-13.
595   53.   George, E.L., et al., *Defects in mesoderm, neural tube and vascular development in
596   mouse embryos lacking fibronectin.* Development, 1993. **119**(4): p. 1079-91.
597   54.   Suwan, K., et al., *Versican/PG-M Assembles Hyaluronan into Extracellular Matrix and
598   Inhibits CD44-mediated Signaling toward Premature Senescence in Embryonic
599   Fibroblasts.* J Biol Chem, 2009. **284**(13): p. 8596-604.
600   55.   Kloeckener-Gruissem, B., et al., *Identification of the genetic defect in the original
601   Wagner syndrome family.* Mol Vis, 2006. **12**: p. 350-5.
602   56.   Kloeckener-Gruissem, B., et al., *Novel VCAN mutations and evidence for unbalanced
603   alternative splicing in the pathogenesis of Wagner syndrome.* Eur J Hum Genet, 2013.
604   **21**(3): p. 352-6.
605   57.   Rothschild, P.R., et al., *De novo splice mutation in the versican gene in a family with
606   Wagner syndrome.* JAMA Ophthalmol, 2013. **131**(6): p. 805-7.
607   58.   Burns, T.A., et al., *Imbalanced expression of Vcan mRNA splice form proteins alters
608   heart morphology and cellular protein profiles.* PLoS One, 2014. **9**(2): p. e89133.
609   59.   Dours-Zimmermann, M.T., et al., *Versican V2 assembles the extracellular matrix
610   surrounding the nodes of ranvier in the CNS.* J Neurosci, 2009. **29**(24): p. 7731-42.
611   60.   Nandadasa, S., S. Foulcer, and S.S. Apte, *The multiple, complex roles of versican and
612   its proteolytic turnover by ADAMTS proteases during embryogenesis.* Matrix Biol, 2014.
613   **35**: p. 34-41.
614   61.   Hope, C., et al., *Immunoregulatory roles of versican proteolysis in the myeloma
615   microenvironment.* Blood, 2016. **128**(5): p. 680-5.
616   62.   McCulloch, D.R., et al., *ADAMTS metalloproteases generate active versican fragments
617   that regulate interdigital web regression.* Dev Cell, 2009. **17**(5): p. 687-98.
618   63.   Islam, S., et al., *Accumulation of versican facilitates wound healing: implication of its
619   initial ADAMTS-cleavage site.* Matrix Biol, 2019.
620   64.   Asano, K., et al., *Stromal Versican Regulates Tumor Growth by Promoting
621   Angiogenesis.* Sci Rep, 2017. **7**(1): p. 17225.

65. Oladipupo, S.S., et al., *Impaired tumor growth and angiogenesis in mice heterozygous*
for Vegfr2 (Flk1). Sci Rep, 2018. **8**(1): p. 14724.
66. Zhao, Y. and R.P. Singh, *The role of anti-vascular endothelial growth factor (anti-VEGF)*
in the management of proliferative diabetic retinopathy. Drugs Context, 2018. **7**: p.
212532.
67. Arranz, A.M., et al., *Hyaluronan deficiency due to Has3 knock-out causes altered*
neuronal activity and seizures via reduction in brain extracellular space. J Neurosci,
2014. **34**(18): p. 6164-76.
68. Matsumoto, K., et al., *Conditional inactivation of Has2 reveals a crucial role for*
hyaluronan in skeletal growth, patterning, chondrocyte maturation and joint formation in
the developing limb. Development, 2009. **136**(16): p. 2825-35.
69. Tallquist, M.D. and P. Soriano, *Epiblast-restricted Cre expression in MORE mice: a tool*
to distinguish embryonic vs. extra-embryonic gene function. Genesis, 2000. **26**(2): p.
113-5.
636

The versican-hyaluronan complex is an extracellular matrix microniche for Flk1+

precursors in vasculogenesis and primitive hematopoiesis

Sumeda Nandadasa, PhD, Anna O'Donnell BS, MBA, Ayako Murao BS, Yu Yamaguchi, MD,

PhD, Ronald J. Midura PhD, Lorin Olson, PhD and Suneel S. Apte, MBBS, DPhil[*]

SUPPLEMNTAL FIGURES AND TABLE

Figure-1, Supplement-1

Figure 1, Supplement 1: Characterization of the *Vcan^hdf* mutation and genotyping strategy.
(A) *Vcan* locus, showing the retroviral insertion site in intron 7. *Vcan* exons are indicated in blue and the insertion in red. Primer sequences and amplicons used for genotyping are shown at the bottom. **(B)** Sequence of the 401bp *Vcan^hdf* amplicon and the primer sequences used (underlined). The *Vcan* intron 7 sequence is in blue text and sequence corresponding to the retroviral insertion is in red text.

Figure-1, Supplement-2

Figure 1, Supplement 2: Severe disorganization and cellular changes in *Vcan^{hdf/hdf}* yolk sacs

(A) E9.5 *Vcan^{hdf/hdf}* yolk sacs show dramatically increased F-actin staining.

(B) DAPI staining shows altered nuclear morphology and disrupted cellular organization of the *Vcan^{hdf/hdf}* yolk sac mesoderm (Me) and mesothelium and (with **A**) separation from visceral endoderm (VE). High magnification of the boxed areas (red line) in the upper panels of **B** are shown below. Arrowheads identify the nuclei of yolk sac mesothelium, which are rounded in *Vcan^{hdf/hdf}* yolk sac.

(C) Transmission electron microscopy (TEM) images of wild type and *Vcan^{hdf/hdf}* yolk sacs showing endothelium-lined blood islands in the wild type and detached vascular endothelial cells in the *Vcan^{hdf/hdf}* yolk sac. The red line marks the basement membrane of the visceral endoderm, En=vascular endothelium. Scale bars in **A-B**= 50μm, 5μm in **C**.

Figure 4, Supplement 1:RNAScope *in situ* hybridization of *Vcan* exon 7, *Vcan* exon 8, *Has2*, *Flk1* and *Runx1* probes to consecutive sections from four individual E7.5 wild type embryos. Hybridization signal is red, sections were counterstained with hematoxylin (blue). Scale bar = 100μm.

Figure-4, Supplement-2

Figure 4, Supplement 2: Data extracted from a published single cell transcriptome atlas of mouse gastrulation illustrates specific *Vcan* and *Has2* expression by *Kdr*-expressing hematoendothelial cells.

(A) Force-directed graph layout of blood-related lineages showing the lineage-specific and time-resolved expression of *Kdr, Vcan* and *Has2*. The stippled red line encircles hematoendothelial progenitors.

(B) Violin plots of *Kdr, Vcan*, *Has2* and *Cd44* expression by cells of various lineages from E6.75 embryos. Note strong co-expression of these genes by single hematoendothelial progenitors.

(C) Uniform manifold approximation and projection (UMAP) plots showing overlap between single cell *Kdr* and *Vcan* expression at different gestational ages. High magnification inserts of hematoendothelial progenitors are indicated within red boxes.

Figure-4, Supplement-3

Figure 4, Supplement 3: Data extracted from a published single cell transcriptome atlas of mouse gastrulation illustrates lack of link protein gene expression by hematoendothelial progenitors and versican cleavage in blood islands.

(A) Violin plots for *Hapln1-4* expression by cells of various lineages from E6.75 embryos. No expression of *Hapln1-4* is detected in hematoendothelial progenitors (red arrowhead) at this stage.

(B) HAPLN1 staining was not detected in E7.5 blood islands marked by versican staining (white arrowheads).

(C) ADAMTS-cleaved versican staining (anti-DPEAAE) is undetectable in E7.5 blood islands (white arrowheads). Scale bars = 100μm in **B** and **C**.

Figure-5, Supplement-1

A *Fn1*

Wt

Vcan^{hdf/hdf}

B

Embryos

Yolk sacs

**

***

Fold change

Wt HDF Wt HDF

Figure 5, Supplement 1: Increased *Fn1* transcription in *Vcan^{hdf/hdf}* embryos
(A) RNAscope *in-situ* hybridization of E9.5 wild type and *Vcan^{hdf/hdf}* embryos showing stronger *Fn1* RNA expression in the mutant embryo.
(B) qRT-PCR analysis of *Fn1* transcript in wild type and *Vcan^{hdf/hdf}* embryos and yolk sacs showing significantly increased expression in the mutant. (n=3 embryos and yolk sacs each genotype, error bars= S.E.M., **, p<0.01; ***, p<0.0001). Scale bars = 100µm in **A**.

Figure-6, Supplement-1

Figure 6, Supplement 1: Loss of yolk sac vasculature, HA and versican staining in *Has2* conditionally deleted embryos

(A) Whole mount images of *Has2^{Fl/Fl}; Sox2*Cre yolk sacs and embryos showing loss of yolk sac vasculature and heart defect (red arrowhead) compared to control littermates (N=8 each genotype).
(B) RNAScope in situ hybridization for *Has2, Vcan* exon 7 (GAG α) and exon 8 (GAG β) shows residual *Has2* transcript and slightly reduced *Vcan* transcript labeling in *Has2^{Fl/Fl}; Sox2* Cre embryos (n=4 embryos each probe).
(C) *Has2^{Fl/Fl}; Sox2* Cre yolk sacs show loss of HA and versican staining (upper panels) and increased versican catabolism is revealed by DPEAAE staining (lower panels) (N=4 each group).
(D) *Has2^{Fl/Fl}; Sox2* Cre embryos show loss of HA and versican staining (N=3 each group).
(E) *Has2^{Fl/Fl}; Sox2* Cre embryos show increased DPEAAE staining (N=3 each group).

Figure-7, Supplement-1

Figure 7, Supplement 1: Formation of blood and vascular lineages is impaired in *Vcan*-null embryoid bodies.

(A-F) 10-day differentiated embryoid bodies from three independent experiments were analyzed by qRT-PCR. *Has2* transcription in embryoid bodies was unaffected by the loss of versican. mRNA for the mesoderm marker *Brachyury* was significantly lower in F9 but not in D8 embryoid bodies contrasting with *Vcan*^{hdf/hdf} embryos (main Fig. 3E). Expression of the hematovascular progenitor marker *Flk1,* blood lineage commitment progenitor markers *Itga2b* (CD41) and *Runx1*, differentiated blood markers *Hbb* (β-globin), *Gata1* and *Ptprc* (CD45) and endothelial marker *Pecam1* (CD31) was reduced in *Vcan*-null embryoid bodies (N=3 independent batches of 10 day embryoid bodies from each genotype, error bars= S.E.M.,*, p<0.05; **, p<0.01; ***, p<0.0001).

Figure-8, Supplement-1

Figure 8, Supplement 1: Formation of hyaluronan and HA cables in *ADAMTS9* deficient RPE-1 cells in 25mM glucose

(A) Four independent representative areas of wild type RPE-1 and *ADAMTS9* deficient RPE-1 (D12) cell cultures stained with HAbp showing HA accumulation and numerous long HA-cables present in D12 cultures, but not parental RPE1 cells (white arrowheads).

(B) High-magnification confocal images showing two examples of long HA cables (green) decorated with versican (red). Scale bars = 50μm in **A** and 5μm in **B**.

A D12+ siControl   D12+ *VCAN* siRNA

B

VCAN · HAS1 · HAS2 · HAS3 · TMEM2 · CD44

Figure 8, Supplement 2: Loss of HA and HA cables in *ADAMTS9*-deficient RPE-1 cells upon *VCAN* siRNA treatment
(A) Four independent representative areas of *ADAMTS9*-deficient RPE-1 (clone D12) cell cultures transfected with control siRNA (siControl) or *VCAN* siRNA, stained with HAbp showing loss of HA after *VCAN* knockdown, whereas HA-cables persist in siControl D12 cells (white arrowheads).
(B) qRT-PCR of siControl and *VCAN* siRNA treated D12 cells showing decreased *VCAN* transcript levels upon siRNA treatment and upregulation of *TMEM2* expression (n=3 independent siRNA treatment experiments, error bars= S.E.M., *, p<0.05; **, p<0.01; ***, p<0.001;   ****, p<0.0001). Scale bar = 50μm in **A**.

Figure-8, Supplement-3

Figure 8, Supplement 3: VEGF$_{165}$ co-localizes with versican in extracellular matrix of RPE1 D12 cells in a dose-dependent manner.
(A) Wild type RPE-1 and *ADAMTS9*-deficient RPE-1 (D12) cells treated with PBS or increasing concentrations of recombinant biotinylated VEGF$_{165}$ (green) showing dose-dependent VEGF staining intensity and colocalization with versican (red) in D12 cells.
(B) Super-resolution confocal images of a long versican/HA cable showing patches of VEGF$_{165}$ (green) deposited along the versican stained cable (red). Nuclei are stained blue with DAPI. The upper merged panel shows some strongly co-staining regions (arrowheads), whereas the center panel shows co-staining in the X-Z plane. The lower panels show the single-color images of the areas marked as Box 1 and Box 2 in the upper panel. Scale bars = 50μm in **A** and 10μm in **B**.

Figure-8, Supplement-4

Figure 8, Supplement 4: Ihh co-localizes with versican in extracellular matrix in a dose-dependent manner.
(A) Wild type RPE-1 and *ADAMTS9* deficient (D12) cells treated with PBS or increasing concentrations of recombinant Ihh N-terminal domain (green) showing dose-dependent deposition and increased co-staining (yellow) with versican (red) in D12 cells.   Nuclei are stained blue with DAPI.
(B) Super-resolution confocal image of a versican-HA stained cable showing Ihh staining (green) along a versican stained cable (red). Strongly co-stained regions are indicated by arrowheads. The center panel shows the Y-Z plane. Bottom panels show single-color images of the area imaged in the upper panel. Scale bars = 20μm in **A** and 10μm in **B**.

106   **Supplemental Table 1. Oligonucleotide primers used for qRT-PCR.**

18S-F	5'-TTGACGGAAGGGCACCACCAG-3'
18S-R	5'-GCACCACCACCCACGGAATCG-3'
Actb-F	5'-GTGGGCCGCTCTAAGGCACCA-3'
Actb-R	5'-TGGCCTTAGGGTGCAGGGGG-3'
Brachyury-F	5'-GCTCTAAGGAACCACCGGTCATC-3'
Brachyury-R	5'-ATGGGACTGCAGCATGGACAG-3'
Myc-F	5'-TGACCTAACTCGAGGAGGAGCTGGAATC-3'
Myc-R	5'-AAGTTTGAGGCAGTTAAAATTATGGCTGAAGC-3'
Fn1-F	5'- GTCTAGGCGAAGGCAATGG-3'
Fn1-R	5'- CCTATAGGATGTCCGGGTGT-3'
Flk1-F	5'-ACTGCAGTGATTGCCATGTTCT-3'
Flk1-R	5'-TCATTGGCCCGCTTAACG-3'
Gata1-F	5'-AGCCTATTCTTCCCCCAAGTTTC-3'
Gata1-R	5'-CTCCACAGTTCACACACTCTCTGG-3'
Has2-F	5'-GGTCCAAGTGCCTTACTGAAAC-3'
Has2-R	5'-TGTAGAGCCACTCTCGGAAGTA-3'
Hbb-F	5'-GCTCTTGCCTGTGAACAATG-3'
Hbb-R	5'-GTCAGAAGACAGATTTTCAAATG-3'
Itga2b-F	5'-GAACCTGGAAGAAGCTGGCG-3'
Itga2b-R	5'-CTCCACTGTGGCTTCAGCTTG-3'
Nanog-F	5'-GGTTGAAGACTAGCAATGGTCTGA-3'
Nanog-R	5'-TGCAATGGATGCTGGGATACTC-3'
Oct4-F	5'-TCAGGTTGGACTGGGCCTAGT-3'
Oct4-R	5'-GGAGGTTCCCTCTGAGTTGCTT-3'
Pecam1-F	5'-CTGCCAGTCCGAAAATGGAAC-3'
Pecam1-R	5'-CTTCATCCACCGGGGCTATC-3'
Ptprc-F	5'-GAACATGCTGCCAATGGTTCT-3'
Ptprc-R	5'-TGTCCCACATGACTCCTTTCC-3'
Runx1-F	5'-GACCATCACCGTCTTTACAAATC-3'
Runx1-R	5'-TTGGTCTGATCATCTAGTTTCTGC-3'
Sox2-F	5'-GCACATGAACGGCTGGAGCAACG-3'
Sox2-R	5'-TGCTGCGAGTAGGACATGCTGTAGG-3'
Tmem2-F	5'-TCCTGGAGAAGCCAGAACTCAC-3'
Tmem2-R	5'-AGATGCACTGGCGTTGGTCGTT-3'
GAPDH-F	5'-AGCCTCAAGATCATCAGCAATG-3'
GAPDH-R	5'-CTTCCACGATACCAAAGTTGTCAT-3'
CD44-F	5'-CCAGAAGGAACAGTGGTTTGGC-3'
CD44-R	5'-ACTGTCCTCTGGGCTTGGTGTT-3'

HAS1-F	5'-CTGCGATACTGGGTAGCCTTCA-3'
HAS1-R	5'-CCAGGAACTTCTGGTTGTACCAG-3'
HAS2-F	5'-CCGATAACAGAAAATCTCTTTTTC-3'
HAS2-R	5'-AGCAGCCCATTGAACCAGAGAC-3'
HAS3-F	5'-AGCACCTTCTCGTGCATCATGC-3'
HAS3-R	5'-TCCTCCAGGACTCGAAGCATCT-3'
TMEM2-F	5'-ACCGAGCACATTCCAACTACCG-3'
TMEM2-R	5'-GGCAGAGATGATTGAGAGGAACG-3'

SUPPLEMENTAL METHODS

Mouse genotyping

Vcan^Tg(Hoxa1)1Chm (*Vcan*^hdf) genotyping is described in Figure-1, Supplement-1. *Has2*^flox genotyping

was carried out using the following primers; forward primer: 5'

-TGCAGAATTTAGGGGCGAATTGGGAGCTAA-3', reverse primer: 5'-

ATGAGGTTAGAGATTAGCAAGACTGAGTTC-3' which results in a 441bp band for wild type

and a 550bp band for the floxed allele. Sox2-Cre mice were genotyped using the following

three-primer combination, primer-1: 5'-CTTGTGTAGAGTGATGGCTTGA-3', primer-2: 5'-

TAGTGCCCCATTTTTGAAGG-3', primer-3: 5'-CCAGTGCAGTGAAGCAAATC-3' which results

in a 207bp band for wild type and a 165bp band for Sox2-Cre. The *Has2* null allele was

genotyped using forward primer: 5'-CTTGAACCTTGAGTGTGCCATTTTGTAGTC-3', reverse

primer: 5'-CATTCTTGTTTTGAAGTTTGTTTCCTTGAC-3' which results in a 346bp band.

Immunostaining and fluorescence microscopy

Immunostaining of E9.5 and E8.5 yolk sac was carried out on 30μm thick vibratome sections [1]

or paraffin-embedded 7 μm sections. Immunostaining of cultured RPE-1 cells were carried out in

8-chamber cell culture slides (Fisher Scientific, catalog no. 354118). Immunostaining of collagen-

embedded embryoid bodies were carried out in 4-chamber cell culture slides (Fisher Scientific,

catalog no. 354114). Confocal microscopy images of whole mount mouse embryos and sections

were acquired using a Leica TCS SP5 II multiphoton confocal microscope equipped with a 25X

water immersion objective (Leica Microsystems, Wetzlar, Germany). For 3D-projection of whole

mount Z-stacks, the Volocity 3D imaging software was used (version 6.3, PerkinElmer, Inc.,

Waltham, MA) in maximum intensity projection method. Confocal and super resolution

microscopy of RPE-1 cells were carried out using a Leica TCS SP8 confocal microscopy equipped

with Huygens deconvolution (HyVolution) capability as previously described [2] .

Methylcellulose colony formation assay

Single cell suspensions of E8.5 embryos, yolk sacs or day-10 embryoid bodies (EBs) were

generated by incubation with trypsin for 10 minutes followed by disaggregation by pipetting with

a 200 μL pipette tip until complete. 50,000 cells from each experimental group were transferred

to a single 35 mm culture dish containing 1 mL of MethoCult GF M3534 culture medium

(Methylcellulose medium with recombinant cytokines for mouse cells, Stem Cell Technologies,

Vancouver, CA, catalog no. 03534) using a 3 mL syringe and 16-gauge needle, following the

manufacturer's protocol. Triplicate cultures from each genotype were incubated for 14 days in a

humidified, 5% CO_2, 37°C cell culture incubator. Blood colonies were counted using an inverted

microscope. Aggregates with >50 cells were considered a colony-forming unit (CFU).

Primary antibodies and dilutions

For staining whole mount embryos, yolk sacs, vibratome sections and cell culture chamber slides

the following antibodies, staining reagents and dilutions were used: Rabbit polyclonal anti-mouse

versican GAGβ domain (Millipore-Sigma, catalog no. AB1033) 1:200; Rabbit polyclonal anti-

versican cleavage site (anti-Vc) [3] 1:400; Rabbit polyclonal anti- versican V_0/V_1 neo cleavage

antibody (DPEAAE) (Thermo Fisher, catalog no. PA1-1748A) 1:400; biotinylated hyaluronan

binding protein (HAbp) (Millipore-Sigma, Calbiochem, catalog no. 385911) 1:100; rabbit

polyclonal anti-fibronectin (Abcam, catalog no. Ab2413) 1:200; rabbit polyclonal anti-collagen-IV

(Rockland antibodies and assays, catalog no. 600-401-106) 1:400; rat monoclonal anti-mouse

Flk1 (clone Avas12, Thermo Fisher, catalog no. 17-5821-81) 1:200; hamster anti-mouse CD31

(Milipore-Sigma, catalog no. MAB1398Z) 1:400; FITC-conjugated anti-mouse CD41 (Biolegend,

catalog no.133904) 1:100; mouse monoclonal Cy3 conjugated smooth muscle α-actin (Sigma-

Aldrich, catalog no. C6198) 1:600;Alexa Fluor-568 phalloidin (Life Technologies, catalog no.

A12380) 1:500; Streptavidin-FITC (Invitrogen, catalog no. SA1001) 1:400; Goat anti-mouse Ihh

N-terminus (R&D Systems, catalog no. AF1705) 1:400; Mouse anti-Hapln1 (Link protein)

(DSHB, 9/30/8-A-4-C) 1:100. All primary antibodies were diluted in 5% normal goat serum in

PBST (PBS+ 0.1% Tween 20) and incubated overnight at 4°C. Alexa488, 568 or 647-conjugated

secondary antibodies against the corresponding species, or streptavidin conjugated Alexa488 or

568-labeled antibodies for HAbp detection were purchased from Invitrogen and were used at

1:400 dilution at room temperature for 2-3hrs. Slides were photographed using an Olympus BX51

upright microscope (Olympus, Center Valley, PA) connected to a Leica DFC7000T camera and

Leica Application Suite v4.6 imaging software (both from Leica, Wetzlar, Germany).

Quantitative real time RT-PCR (qRT-PCR) analysis

RNA from whole embryos, dissected yolk sacs, cultured cells or EBs was extracted using TRIzol

reagent (ThermoFisher Scientific, catalog no. 15596026) according to manufacturer

recommendations. 2 μg of RNA from each sample (embryo, yolk sac or EB pool) was used for

cDNA synthesis using the high capacity cDNA synthesis kit (Applied Biosystems, Thermo Fisher

Scientific, catalog no. 4368814). qRT-PCR was carried out using a CFX96 Touch Real-Time PCR

detection system (Bio-Rad Laboratories) with the Bullseye EvaGreen qPCR mix (MIDSCI, catalog

no. BEQPCR-S), using primer pairs listed in the Supplemental Table-1. 18s ribosomal RNA or

Actb RNA were used for normalization of mouse genes while *GAPDH* was used for normalizing

human gene expression and relative expression of genes was calculated by the ΔΔCt

quantification method. An unpaired, two-tailed Student t-test was used to determine statistical

significance.

RNAscope *In-situ* hybridization and RNAseq database mining

4% PFA fixed mouse embryos and yolk sacs were paraffin embedded and 7 μm thick sections

were collected immediately prior to in-situ hybridization. All RNAscope in situ probes and reagents

were from Advanced Cell Diagnostics. The probes were: *Vcan* exon 7 (GAG-α) (catalog no.

428311), *Vcan* exon 8 (GAG-β) (catalog no. 428321), *Has2* (catalog no. 465171), *Flk1 (Kdr)*

(catalog no. 414811) and *Runx1* (catalog no. 406671). In situ hybridization was carried out

following the user manual for RNAScope 2.5 HD Red detection kit using the RNAscope HybEZ

oven. Slides were photographed as described above.

A database containing recent analysis of 116,312 single cells (Pijuan-Sala B, et al.,

2019) obtained from gastrulating mouse embryos spanning E6.5 to E8.5 was searched at

specific gestational ages for expression of *Vcan, Kdr, Has2* and *Cd44*. Individual data panels

were generated by the web portal for the deposited scRNAseq data accessed via:

https://marionilab.cruk.cam.ac.uk/MouseGastrulation2018/

Western blotting

7.5% reducing SDS-PAGE was used for western blotting. Embryoid bodies were lysed in PBS

containing 1% Tween 20 supplemented with a protease inhibitor cocktail (Roche, catalog no.

11873580001). Supernatants obtained after centrifugation at 1200 rpm for 10 minutes were

treated with BSA free *Proteus vulgaris* chondroitinase ABC (Sigma-Aldrich, catalog no. 3667)

for 2 h at 37°C. Samples were boiled in Laemmli sample buffer (0.375% mM Tris.HCL, 9%

SDS, 50% glycerol, 0.03% bromophenol blue, with 9% (v/v) β-mercaptoethanol) for 10 minutes

followed by SDS-PAGE. Nitrocellulose membranes with transferred proteins were blocked in LI-

COR Odyssey PBS blocking buffer (LI-COR Biosciences, Lincoln, NE catalog no. 927-40000)

for 30 minutes in RT. Primary antibodies rabbit anti-mouse versican (GAGβ) antibody (1:1000)

(Millipore-Sigma, catalog no. AB1033) or mouse monoclonal GAPDH antibody (1:5000)

(Millipore-Sigma, catalog no. MAB374) were diluted in blocking buffer and incubated overnight

at 4°C with gentle rocking. LI-COR IR dye secondary antibodies against mouse and rabbit

antibodies (LI-COR Biosciences, Lincoln, NE) were used at 1:10000 dilutions in PBS to detect

primary antibodies. A LI-COR Odyssey CLx scanner and the LI-COR Image Studio (ver. 4.0)

was used to image western blots.

Fluorophore-assisted carbohydrate electrophoresis

FACE procedures used were essentially as previously published [4-6]. In brief, tissue was

extensively digested with proteinase K and released glycosaminoglycans (GAGs) were isolated

by ice-cold ethanol precipitation and centrifugation. GAGs were resuspended in 0.1 M ammonium

acetate, pH 7.0 and then digested overnight with *Streptomyces* hyaluronidase (200 mU) at 37° C

for HA-FACE. HA digestion products were separated from other intact GAGs using ice-cold

ethanol precipitation and centrifugation. The supernatant was then dried under vacuum

centrifugation and reacted with 6.25 mM 2-aminoacridone (AMAC) and 625 mM sodium

cyanoborohydride (NaBH$_3$CN) overnight at 37° C. The pellet was resuspended in 0.1 M

ammonium acetate, pH 7.0 and then digested overnight with chondroitinase ABC (25 mU) at 37°

C for CS-FACE. CS digestion products were separated from other intact GAGs using ice-cold

ethanol precipitation and centrifugation. The supernatant was then dried under vacuum

centrifugation and reacted with 6.25 mM AMAC and 625 mM NaBH$_3$CN overnight at 37° C.

AMAC-labeled HA and CS digestion products were loaded onto 20% (w/v) acrylamide gels and

these products were resolved by electrophoresis at 500 V constant current over a 50-60 minute

run time. FACE gels were imaged while still housed in the gel plates on a UVP ChemiDoc-It^2 515

system. Digital gel images were analyzed for AMAC-labeled HA and CS digestion product band

intensities using ImageJ (NIH, Bethesda MD).

Transmission electron microscopy

E9.5 yolk sacs were harvested from timed pregnancies and fixed in 4% PFA + 2.5%

glutaraldehyde in PBS overnight. Samples were dehydrated in an ethanol:PBS gradient and fully

dehydrated samples were embedded in pure Eponate12 resin and allowed to polymerize

overnight. Ultrathin sections (85 nm) were stained with uranyl acetate and lead citrate and viewed

and imaged on a FEI Tecnai G2 Spirit BioTWIN Transmission Electron Microscope (FEI company,

Hillsboro, OR) equipped with an Orius 832 CCD camera (Gatan, Inc., Pleasanton, CA).

Some E8.5 embryos were fixed in 4% PFA + 2.5% glutaraldehyde + 0.7% w/v ruthenium

hexamine trichloride (Sigma-Aldrich, catalog no. 262005) in PBS overnight to optimize

proteoglycan and cell membrane preservation [7] and embedded in Eponate 12 resin as

described above. 1 micron thick sections were stained with 1% toluidine blue containing 2%

sodium borate for 30 seconds on a hot plate and washed with distilled water. Slides were mounted

with Cytoseal XYL mounting media (ThermoFisher Scientific, catalog no. 22-050-262) and imaged

as described for routine microscopy.

VEGF$_{165}$ and Ihh treatment of RPE-1 cells

Wild type hTERT RPE-1 (ATCC, CRL-4000) cells and ADAMTS9 deficient RPE-1 cells [2] were

cultured in 8-chamber cell culture slides (Fisher Scientific, catalog no. 354118) with a seeding

density of 50,000 cells/chamber in DMEM F-12 culture medium containing 10% FBS for 48 hrs

and washed with PBS and cultured in DMEM F-12 culture medium without FBS for an additional

24hrs in a cell culture incubator at 37C with 5% CO_2. Cells were cultured in DMEM F-12 medium

containing 25, 50, 100 ng/ml recombinant human biotinylated VEGF$_{165}$ (R&D Systems, Cat. No.

BT293-010) or recombinant human/mouse Ihh N-terminus (C2811) (R&D Systems, Cat. no. 1705-

HH-025) for 6hrs. Cells were washed three times with PBS and fixed in 4% PFA. For versican

depletion experiments, a previously validated highly potent *VCAN* siRNA (Ambion, Cat. no.

S229335) [8-10], or control siRNA (Ambion, cat. no. 4390843) were added to the cell cultures

24hrs after seeding using the Lipofectamine RNAiMAX protocol (Invitrogen, Cat. no. 13778) and

cultured for 24hrs prior to serum starvation and growth factor treatment. For treatment with

chondroitinase ABC (0.2 U/ml in PBS, Sigma-Aldrich, Cat. no. C3667), cells were incubated

overnight prior to the addition of recombinant VEGF$_{165}$ or IHH. For co staining, recombinant

VEGF$_{165}$ was detected using Streptavidin FITC (1:400, Invitrogen, SA1001), recombinant IHH

using anti mIhh-N antibody (1:400, R&D Systems, Cat. no. AF1705) and versican using the pVC

antibody at a 1:400 dilution as previously described [2, 3]. Fixed cell culture chambers were

blocked and incubated with primary antibodies in 5% normal goat serum for VEGF$_{165}$ experiments

and 5% normal horse serum (Sigma-Aldrich, Cat. no. H0146-10ml) for Ihh experiments. Goat anti

rabbit Alexa-568 secondary antibody (Invitrogen, Cat. no. A11011) and donkey anti goat Alexa-

488 (Invitrogen, Cat. no. A11055) secondary antibodies were used at 1:600 dilution for detecting

versican and Ihh primary antibodies. Super resolution confocal microscopy was conducted as

previously described [2].

Embryoid body formation and VEGF$_{165}$ induced angiogenic-sprouting

mESCs were cultured for two days till 100% confluent, gently washed with PBS and incubated

with IMDM without LIF for an additional 48 h. IMDM without LIF was used in subsequent

manipulations to form EBs. The medium was removed and 250μl of trypsin was added to each

plate and aspirated out after a 30 second incubation leaving a thin film of trypsin. Upon

detachment and rounding of cells, 1 mL of IMDM was added to each plate, and a small pipette tip

was used to generate small aggregates mechanically from the mESC sheet. The aggregates were

transferred to 60 mm (non-adhesive) bacteriological dishes (Corning, catalog no. 351007)

containing 5 mL of IMDM, where most curled and formed free-floating spheroids after 24 hrs.

These 1-day EBs were transferred to a fresh bacteriological dish with 5 mL of IMDM and cultured

for an additional 3 days. 10-20 4-day old EBs from each genotype were transferred to a 4-well

chamber (Fisher Scientific, catalog no. 354114) and embedded in bovine type 1 collagen

(Corning, catalog no. 354231) containing 30 ng/mL recombinant VEGF$_{165}$ (PreproTech, catalog

no. 100-20). 500 μL IMDM containing 30 ng/mL recombinant VEGF$_{165}$, was added atop each

collagen gel and replaced very 4 days for a further 12 days. The collagen-embedded EBs were

washed with PBS and fixed in 4% PFA for 15 min, permeabilized by three washes in PBS

containing 0.3% Tween 20 for 30 minutes prior to immunostaining with anti-CD31 or anti-smooth

muscle α-actin (SMA). For qRT-PCR, EBs in floating culture were incubated for 10 days (day-10

EBs). 300 μL of TRIzol reagent (ThermoFisher scientific, catalog no. 15596026) was used to

harvest total RNA from each 60 mm plate.

Embryoid body vascular sprout quantifications

For quantifying EB vascular sprouts, collagen-embedded, fixed embryoid bodies in 4-chamber

slides were imaged using an inverted bright field microscope. Images were analyzed by Image

J-FIJI (NIH, Bethesda, MD), and the freehand drawing tool was used to trace the length of

individual vascular sprouts. A total N= 72 wild type, 68 D8 and 68 F9 EBs were analyzed in 3

independent experiments. Lengths of 402 sprouts from wild type, 91 from D8 and 97 from F9

EBs were measured.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Primary antibodies		
Hamster monoclonal anti-mouse CD31	Millipore-Sigma	MAB1398Z
Rabbit polyclonal anti-versican	Millipore-Sigma	AB1033
Rabbit polyclonal anti-versican cleavage site (pVC)	Lab generated	N/A
Rabbit polyclonal anti-versican V_0/V_1 Neo cleavage (DPEAAE) antibody	Thermo Fisher	PA1-1748A
Rabbit polyclonal anti-fibronectin	Abcam	Ab2413
Rabbit polyclonal anti-collagen-IV	Rockland antibodies and assays	600-401-106
Rat monoclonal anti-mouse Flk1 (clone Avas12)	Thermo Fisher	17-5821-81
Rat monoclonal anti-mouse CD41	Biolegend	133904
Mouse monoclonal anti-smooth muscle α-actin	Sigma-Aldrich	C6198
Mouse monoclonal anti-GAPDH	Millipore-Sigma	MAB374
Goat anti-moue IHH N-terminus antibody	R&D Systems	AF1705
Mouse monoclonal anti-HAPLN1 (Link protein)	DSHB	9/30/8-A-4-C
Bacterial and Virus Strains		
One shot TOP10 chemically competent *E. coli* cells	Thermo Fisher	K280020
Biological Samples		
N/A	N/A	N/A
Chemicals, Peptides, and Recombinant Proteins		
Leukemia inhibitory factor (LIF)	EMD-Millipore	ESG1106
Recombinant $VEGF_{165}$	PreproTech	100-20
Recombinant biotinylated $VEGF_{165}$	R&D Systems	BT293-010
Recombinant human/mouse IHH N-terminus	R&D Systems	1705-HH-025
Bovine type 1 collagen	Corning	354231
Proteus vulgaris chondroitinase ABC	Sigma-Aldrich	3667
Biotinylated hyaluronan binding protein (HAbp)	Millipore-Sigma, Calbiochem	385911

Alexa Fluor-568 phalloidin	Invitrogen	A12380
Streptavidin-FITC	Invitrogen	SA1001
Ruthenium hexamine trichloride	Sigma-Aldrich	262005
Protease inhibitor cocktail	Roche	11873580001
Critical Commercial Assays		
RNAscope mouse *Vcan* exon 7 (GAG-α)	Advanced Cell Diagnostics	428311
RNAscope mouse *Vcan* exon 8 (GAG-β)	Advanced Cell Diagnostics	428321
RNAscope mouse *Has2*	Advanced Cell Diagnostics	465171
RNAscope mouse *Kdr* (Flk1)	Advanced Cell Diagnostics	414811
RNAscope mouse *Runx1*	Advanced Cell Diagnostics	406671
MethoCult GF M3534	Stem Cell Technologies	03534
Bullseye EvaGreen qPCR mix	MIDSCI	BEQPCR-S
Deposited Data		
N/A	N/A	N/A
Experimental Models: Cell Lines		
R1 mouse embryonic stem cells (mESC)	CWRU Transgenic & Targeting Facility	[11] ATCC, SCRC-1011
Experimental Models: Organisms/Strains		
Mouse: *Vcan*$^{Hdf/+}$: Vcan$^{Tg(Hoxa1)1Chm}$/C57BL/6J	N/A	MGI:1857689 [12]
Mouse: *Has1*$^{-/-}$;*Has3*$^{-/-}$	N/A	[13]
Mouse: *Has2* $^{tm1.1Yama}$	N/A	MGI:4360705 [14]
Mouse: *Sox2Cre*^{tg}	N/A	JAX:008454 [15]
Oligonucleotides		
See Supplemental Table 1 for qRT-PCR primers	This paper	N/A
VCAN siRNA	Ambion	S229335
Control siRNA	Ambion	4390843
Recombinant DNA		
mVcan-E2-U6gRNA-Cas9-2A-GFP	Sigma-Aldrich	target ID MM0000080027
mVcan-E3-U6gRNA-Cas9-2A-GFP	Sigma-Aldrich	target ID MM0000080028
pCR-Blunt II-TOPO	Life Technologies	K2800-40
Software and Algorithms		
LI-COR Image Studio (ver. 4.0)	LI-COR	N/A
Volocity 3D imaging software (ver. 6.3)	PerkinElmer	N/A
Image J	NIH	https://imagej.nih.gov/ij/
Other		
N/A		

1. Nandadasa, S., C.M. Nelson, and S.S. Apte, *ADAMTS9-Mediated Extracellular Matrix*
Dynamics Regulates Umbilical Cord Vascular Smooth Muscle Differentiation and
Rotation. Cell Rep, 2015. **11**(10): p. 1519-28.
2. Nandadasa, S., et al., *Secreted metalloproteases ADAMTS9 and ADAMTS20 have a*
non-canonical role in ciliary vesicle growth during ciliogenesis. Nat Commun, 2019.
10(1): p. 953.
3. Foulcer, S.J., et al., *Determinants of versican-V1 proteoglycan processing by the*
metalloproteinase ADAMTS5. J Biol Chem, 2014. **289**(40): p. 27859-73.
4. Calabro, A., et al., *Microanalysis of enzyme digests of hyaluronan and*
chondroitin/dermatan sulfate by fluorophore-assisted carbohydrate electrophoresis
(FACE). Glycobiology, 2000. **10**(3): p. 273-81.
5. Calabro, A., V.C. Hascall, and R.J. Midura, *Adaptation of FACE methodology for*
microanalysis of total hyaluronan and chondroitin sulfate composition from cartilage.
Glycobiology, 2000. **10**(3): p. 283-93.
6. Midura, R.J., et al., *Quantification of hyaluronan (HA) using a simplified fluorophore-*
assisted carbohydrate electrophoresis (FACE) procedure. Methods Cell Biol, 2018. **143**:
p. 297-316.
7. Hunziker, E.B., W. Herrmann, and R.K. Schenk, *Ruthenium hexammine trichloride*
(RHT)-mediated interaction between plasmalemmal components and pericellular matrix
proteoglycans is responsible for the preservation of chondrocytic plasma membranes in
situ during cartilage fixation. J Histochem Cytochem, 1983. **31**(6): p. 717-27.
8. Francis-West, P., et al., *Signalling interactions during facial development.* Mech Dev,
1998. **75**(1-2): p. 3-28.
9. Gueye, N.A., et al., *Versican Proteolysis by ADAMTS Proteases and Its Influence on*
Sex Steroid Receptor Expression in Uterine Leiomyoma. J Clin Endocrinol Metab, 2017.
102(5): p. 1631-1641.
10. Keire, P.A., et al., *Inhibition of versican expression by siRNA facilitates tropoelastin*
synthesis and elastic fiber formation by human SK-LMS-1 leiomyosarcoma smooth
muscle cells in vitro and in vivo. Matrix Biol, 2016. **50**: p. 67-81.
11. Nagy, A., et al., *Derivation of completely cell culture-derived mice from early-passage*
embryonic stem cells. Proc Natl Acad Sci U S A, 1993. **90**(18): p. 8424-8.
12. Yamamura, H., et al., *A heart segmental defect in the anterior-posterior axis of a*
transgenic mutant mouse. Dev Biol, 1997. **186**(1): p. 58-72.
13. Arranz, A.M., et al., *Hyaluronan deficiency due to Has3 knock-out causes altered*
neuronal activity and seizures via reduction in brain extracellular space. J Neurosci,
2014. **34**(18): p. 6164-76.
14. Matsumoto, K., et al., *Conditional inactivation of Has2 reveals a crucial role for*
hyaluronan in skeletal growth, patterning, chondrocyte maturation and joint formation in
the developing limb. Development, 2009. **136**(16): p. 2825-35.
15. Hayashi, S., et al., *Efficient gene modulation in mouse epiblast using a Sox2Cre*
transgenic mouse strain. Mech Dev, 2002. **119 Suppl 1**: p. S97-S101.
334