Genetic analysis of hybrid incompatibility suggests transposable elements increase reproductive isolation in the *D. virilis* clade

Dean M. Castillo[*,1] and Leonie C. Moyle[*]

*Department of Biology, Indiana University, Bloomington IN 47405

1. Current address:
School of Biological Sciences, University of Utah, Salt Lake City UT 84112

47
48   Running title: TEs contribute to postzygotic isolation
49
50
51
52   Keywords: Speciation, transposable elements, dysgenesis, Drosophila
53

54   Corresponding Author
55   Dean Castillo
56   School of Biological Sciences
57   257 South 1400 E, Room 201
58   Salt Lake City UT 84112
59
60   dean.castillo@utah.edu   801-581-7919
61

Abstract

Although observed in many interspecific crosses, the genetic basis of most hybrid

incompatibilities is still unknown. Mismatches between parental genomes in selfish elements and

the genes that regulate these elements are frequently hypothesized to underlie hybrid

dysfunction. We evaluated the potential role of transposable elements (TEs) in hybrid

incompatibilities by examining hybrids between *Drosophila virilis* strains polymorphic for TEs

that cause dysgenesis and a closely related species that appears to lack these elements. Using

genomic data, we confirmed copy number differences in potentially causal TEs between the

dysgenic-causing *D. virilis* (TE+) strain and a sensitive *D. virilis* (TE-) strain and *D. lummei*

genotype. We then contrasted isolation phenotypes in a cross where dysgenic TEs are absent

from both *D. virilis* (TE-) and *D. lummei* parental genotypes, to a cross where dysgenic TEs are

present in the *D. virilis* (TE+) parent and absent in the *D. lummei* parent, predicting increased

reproductive isolation in the latter cross. Using F1 and backcross experiments that account for

alternative hypotheses, we demonstrated amplified reproductive isolation specifically in the

interspecific cross involving TE+ *D. virilis*, consistent with the action of dysgenesis-inducing

TEs. These experiments demonstrate that TEs can contribute to hybrid incompatibilities via

presence/absence polymorphisms.

Introduction

Post-zygotic reproductive isolation in hybrid individuals is generally proposed to be a consequence of negative genetic interactions among two or more loci (Dobzhansky 1937; Muller 1942). Genic differences are obvious candidates to underlie these hybrid incompatibilities. However because the exact loci involved are known in only a few cases (Johnson 2010; Presgraves 2010; Castillo and Barbash 2017), the importance of other genetic mechanisms in generating hybrid incompatibilities is still undetermined. One alternative mechanism involves the action of selfish genetic elements, including repetitive elements (Michalak 2009; Johnson 2010; Werren 2011; Crespi and Nosil 2013). The proposal that transposable elements (TEs) contribute to reproductive isolation was met with early skepticism, largely due to a failure to detect their proposed mutational effects—including increases in the mutation rate—in crosses between different species with marked postzygotic isolation (Coyne 1986; Hey 1988; Coyne 1989). However, we now understand that phenotypes related to 'dysgenesis'—a specific incompatibility phenotype caused by TEs—instead reflect more complex effects of de-repression of TE transcription (Martienssen 2010; Khurana *et al.* 2011) and genome wide DNA damage (Khurana *et al.* 2011), rather than strictly mutational effects on essential genes (Petrov *et al.* 1995; Blumenstiel and Hartl 2005). This potential to cause negative genetic interactions via TE de-repression in hybrids, places TEs within the classical Dobzhansky-Muller framework for the evolution of incompatibilities (Johnson 2010; Castillo and Moyle 2012; Crespi and Nosil 2013).

A role for selfish elements in postzygotic reproductive isolation has been strongly suggested by empirical studies across plants and animals that correlate differences in repetitive elements between species with increased repeat transcription and dysfunction in hybrids (Labrador *et al.* 1999; Martienssen 2010; Brown *et al.* 2012; Dion-Cote *et al.* 2014).

Nonetheless there have been few attempts to directly assess the role of TEs in the expression of

reproductive isolation. Clades with differences in TE content within and between species can be

leveraged to infer the roles of TEs in reproductive isolation by conducting specific crosses that

are predicted a priori to display elevated hybrid dysfunction based on the presence or absence of

lineage specific TEs. Dysgenic systems in *Drosophila* are among the best characterized in terms

of phenotypic effects of TEs that could be relevant to reproductive isolation. Most notable

among these, the dysgenic syndrome within *Drosophila virilis* and the *P-element* system within

D. melanogaster are typically defined by gonad atrophy when a strain carrying transposable

elements is crossed with a strain lacking these elements (Kidwell 1985; Lozovskaya *et al.* 1990).

The dysgenic phenomenon depends on the direction of the cross—dysgenesis occurs when the

female parent lacks the TE elements—and the copy number in the paternal parent (Srivastav and

Kelleher 2017; Serrato-Capuchina *et al.* 2020b). These intraspecific dysgenic systems can be

used to evaluate the contribution of TEs to reproductive isolation with other species, by

contrasting the magnitude of interspecific hybrid dysfunction generated from lines that do and do

not carry active dysgenic elements. This contrast requires two genotypes/lines within a species

that differ in the presence of specific TEs known to cause dysgenesis in intraspecies crosses, and

a second closely related species that lacks these TEs. If the TEs responsible for hybrid

dysfunction within a species also influence reproductive isolation, interspecific crosses involving

the TE-carrying line should exhibit greater hybrid incompatibility than crosses involving the

lines in which TEs are absent.

In this study, we evaluate whether TEs with known effects on intraspecific hybrid

dysgenesis also affect interspecific reproductive isolation, using the *D. virilis* dysgenic system

and the closely related species *D. lummei*. Species in the *D. virilis* clade are closely related,

easily crossed with one another, and vary in transposable element abundance with *D. lummei*

lacking full length active TEs such as *Penelope* (Zelentsova *et al.* 1999; Evgen'ev *et al.* 2000)

and other elements (see below). Dysgenesis was first described within *D. virilis* by Lozovskaya

and colleagues (Lozovskaya *et al.* 1990) when they observed gonadal atrophy and sterility in

males and females in crosses between strains that vary in transposable elements. Initially one

major element, *Penelope,* was implicated in the dysgenic phenotype (Evgenev *et al.* 1997), but

more recent studies have inferred that *Penelope* cannot be the causal agent and have proposed

other specific elements as the primary cause of dysgenesis (Rozhkov *et al.* 2013; Funikov *et al.*

2018; Hemmer *et al.* 2019). Therefore dysgenesis in *D. virilis* may be more complex than

described in *D. melanogaster* (Petrov et al. 1995) because several elements might contribute to

the phenotype.

Using these features and the established role of hybrid dysgenesis among *D. virilis* strains

we contrast a cross where dysgenic TEs are absent from both *D. virilis* (TE-) and *D. lummei*

parental genotypes to a cross where dysgenic TEs is present in the *D. virilis* (TE+) parent and

absent in the *D. lummei* parent, to evaluate the connection between postzygotic reproductive

isolation and differences in TE composition. In intraspecific crosses between female *D. virilis*

that lack relevant TEs (TE-) and male *D. virilis* that carry these TEs (TE+)*,* strong dysgenesis is

observed. Therefore, in interspecific crosses we expect the strongest reproductive isolation

specifically between *D. lummei* females that lack dysgenic inducing TEs and *D. virilis* males that

are TE+—an asymmetrical pattern that is important to distinguish a model of TEs from

alternative explanations. With these expectations, using a series of directed crosses and

backcrosses, we infer that patterns of reproductive isolation observed in our study are consistent

with a causal role for TEs in the expression of postzygotic species barriers.

Materials and Methods

Fly stocks

Two *D. virilis* stocks were used in the experiments. Strain 9 is a wild type strain that was

collected in Georgia, USSR in 1970 and is consistently used as a parent in dysgenic crosses

because it lacks inducing repetitive elements (Lozovskaya *et al.* 1990; Blumenstiel and Hartl

2005). For the purposes of this study we refer to this strain as TE-. The reference genome strain

was used as our inducer strain (UCSD stock number 15010-1051.87) because it is known to

induce dysgenesis (Blumenstiel 2014), and we refer to this strain as TE+. The genome strain was

used instead of Strain 160, which is commonly used in dysgenic studies, because we experienced

significant premating isolation between Strain 160 and *D. lummei* in single pair crosses, that

precluded further genetic analysis and examination of postzygotic isolation. The reference

genome strain is closely related to Strain 160 as both were created by combining several

phenotypic markers to make multiply marked lines (genome strain genotype *b; tb, gp-L2; cd; pe*)

(Lozovskaya *et al.* 1990). The strain of *D. lummei* was acquired from the UCSD stock center

(15010-1011.07).

Estimating copy number of dysgenic causing elements

We used publicly available genomes from several *D. virilis* strains and species in the virilis clade

to examine the copy number of potentially dysgenic causing elements. The SRA accession

numbers and strain information for all samples are listed in Supplemental Table 1. We focused

on candidate TEs that are associated with dysgenesis (Funikov *et al.* 2018) but mapped reads to

all inserts from a previously compiled TE library (Erwin *et al.* 2015) to prevent bias that might

arise from mapping reads from the other species to a TE library generated from *D. virilis*. All

copy number estimates were made using deviaTE (Weilguny and Kofler 2019), which estimates

haploid copy number based on single-copy "control" genes. For our control genes we chose the

D. virilis homologs of Beta-tubulin60D and alpha-tubulin84B as essential single copy genes in

the *Drosophila virilis* genome.

We first compared copy number differences between the genomes of the inducing strain

D. virilis Strain 160 and *D. virilis* Strain 9 (TE-), the two strains typically used in studies of

dysgenesis. We determined the reliability of our estimates for identifying potentially causal TEs,

by comparing our estimate for the enrichment of TE copy in Strain 160 to other studies that have

used read mapping abundance from genomic data and read abundance for piRNA from ovaries

of each strain (Erwin *et al.* 2015; Funikov *et al.* 2018). Although the majority of work on

dysgenesis in *D. virilis* has focused on *Penelope*, new evidence suggests that *Penelope* does not

play a role in dysgenesis (Rozhkov *et al.* 2013; Funikov *et al.* 2018). Therefore, we looked at

relationships between candidate TEs and *Penelope* across *D. virilis* genomes, to determine which

of these TEs had similar copy number distribution as *Penelope*. We specifically focused on

Penelope, Polyphemus, Paris, Helena, Skippy, and *Slicemaster,* given their greater mapping

abundance in inducer vs non-inducer strains (Funikov *et al.* 2018). We used hierarchical

clustering to determine the correspondence of copy number of each candidate TE and *Penelope*

across strains. A strong correspondence suggests a causal role for this element in the dysgenic

studies that have previously focused on *Penelope*.

We next determined which candidates had a higher average copy number in the two

inducing strains (the genome strain TE+ and 160) compared to the non- inducing Strain 9 (TE-)

as well as the copy number average of the two *D. lummei*. We retained the threshold of two-fold

increase in the average TE copy number, based on the analysis above. We also examined the

differences in allele frequencies and polymorphic sites in the *D. lummei* strains compared to *D.*

virilis strains. We filtered SNPs using default settings in deviaTE to calculate the average major

allele frequency at each site as well as the number of sites that were polymorphic. TEs that are

homogenous (have an average major allele frequency close to one) reflect either new invasions

or active TEs, whereas evidence for the accumulation of variable sites indicates older inactive

elements (Beall *et al.* 2002; Erwin *et al.* 2015; Weilguny and Kofler 2019).

Intra- and interspecies crosses to determine the nature of reproductive isolation

Virgin males and females from each stock were collected as they eclosed and aged for seven

days prior to matings. All crosses involved single pairs of the focal female and male genotype

combination (a minimum of 20, range 20-23, replicates were completed for a given female ×

male genotype combination). In total there were three <u>intra-strain</u> crosses that we completed to

account for fecundity differences in subsequent analysis (TE- × TE-, TE+ × TE+, and *D. lummei*

× *D. lummei*). As expected, in the three intra-strain crosses we did not see any evidence of male

or female gonad atrophy (classical dysgenic phenotypes) or skewed sex ratios. There were

differences in the total number of progeny produced, hatchability, and proportion of embryos

that had reached the pre-blastoderm stage (hereafter 'proportion fertilized'). We used these

values as baseline estimates of maternal effects which might influence these phenotypes in

subsequent inter-strain analyses (Table 1; Table 2).

For <u>inter-strain</u> comparisons, we performed all reciprocal crosses between the three

strains to analyze reproductive isolation on F1 progeny. These crosses fell into three categories.

I-Intraspecific Crosses Polymorphic for TEs: Crosses between TE- and TE+ were used to

confirm the expression of intraspecific male dysgenesis when the inducing (TE+) strain was used

as a father. **II-Interspecies crosses with no TEs present:** Crosses between TE- and *D. lummei*

were used to determine the nature of reproductive isolation between the species in the absence of

dysgenic inducing TEs. **III-Interspecies crosses with TEs present:** Crosses between TE+ and

D. lummei were used to determine the additional contribution, if any, of TEs to interspecific

post-mating reproductive isolation (Supplemental Fig 1). This contrast assumes that the two *D.*

virilis strains, TE- and TE+, do not differ from each other in their genic Dobzhansky-Muller

incompatibilities with *D. lummei*, an assumption that we could also evaluate with data from

crosses performed using hybrid genotypes (described further below).

Backcross design to test models of hybrid incompatibility

We used a set of backcross experiments to compare the fit of several models of hybrid

incompatibility to our data for intraspecies dysgenesis and interspecific reproductive isolation.

F1 males from TE+ and *D. lummei* crosses were generated in both directions. These specific F1

males were created because this genotype was used previously to infer the genetic basis of

dysgenesis (Lovoskaya et al. 1990; see below). These F1 males were then backcrossed

individually to TE- and *D. lummei* females. We evaluated two main classes of models with

fitness data from the resulting BC hybrids. Both classes broadly describe Dobzhansky-Muller

incompatibilities, because they involve negative epistatic interactions between loci in the two

parental strains, however they differ in whether they assume that the interactions occur between

genes or between TEs and TE suppressors.

Comparing observed patterns to theoretical expectations

The phenotypes that we observed for postzygotic reproductive isolation could be the

product of several non-mutually exclusive genetic mechanisms. Here we describe the unique

predictions that are made by different incompatibility models, focusing on two major classes:

models that assume interactions between genes from each parental genome (Genic models) and

models that assume an asymmetric copy number of TEs between the parental genomes (TE

models).

Genic models of hybrid incompatibilities

The first set of models we compared to our empirical observations assume that alternative

alleles at incompatible loci have diverged in each lineage and have a dominant negative

interaction in F1 and backcross progeny that carry these alleles (Coyne and Orr 2004). When two

unlinked loci are involved we expect to see discrete classes of backcross (BC1) genotypes that

exhibit hybrid incompatibilities, specifically half of the progeny would carry the incompatible

genotype and half would not (Supplemental Fig. 2). If the hybrid incompatibility is not fully

penetrant then, on average, any specific replicate BC1 cross would appear to have half the

number of progeny exhibiting the hybrid incompatibility compared to F1s from the cross

between the parental species. This provides an expectation for us to compare observed levels of

incompatibility between F1 and BC1 crosses (see Statistical Analysis below).

We also expanded this general model to include interactions between specific

chromosomes (e.g. X-autosome interactions), as well more than two interacting loci. More

complex models are described in the Supplemental Information. For our purposes we considered

an X-autosome model where there is a negative interaction between the X chromosome of the

TE- or *D. lummei* parent and an autosome of the TE+ parent (Supplemental Fig. 3). We

considered this specific X-autosome interaction because this generates

dysgenesis/incompatibility in both sexes for the intraspecific dysgenic cross (TE- × TE+), but no

male gonad dysgenesis/incompatibility in the reciprocal cross, consistent with previous reports

(Lozovskaya *et al.* 1990). For models in which higher order (greater than 2 loci) interactions are

responsible for the incompatibility observed in the F1s, we would expect to observe a smaller

proportion of affected individuals in the backcross compared to the F1 cross (Supplemental

Table 2); the specific expected degree of hybrid incompatibility in the backcross depends on the

total number of loci required for hybrid incompatibility and the number of loci carried by each

parental strain.

TE models of hybrid incompatibility

To contrast with genic models of hybrid incompatibility we assessed the fit of our

observations to a model is based on transposable element copy number. Under this model we

expect asymmetric reproductive isolation in crosses between parental species in F1 crosses, and a

reduction of hybrid incompatibilities in the backcross. In particular, if the F1 parent used in the

backcross has an intermediate copy number of dysgenic causing TEs that is below a minimum

copy number necessary to elicit dysgenesis, then we would expect very low or no hybrid

incompatibilities in the backcross. This dosage/copy number effect for the *D. virilis* system was

first suggested by Vieira et al. (Vieira *et al.* 1998) and has been supported in the *P*-element

system in *D. melanogaster* and *D. simulans* (Serrato-Capuchina *et al.* 2020b). The specific

prediction of low or no hybrid incompatibility in backcrosses comes from previous observations

in crosses between TE- females and males heterozygous for dysgenic factors (F1 males from

TE+ × *D. lummei* crosses), where the progeny are not dysgenic (Lozovskaya et al. 1990). We

interpret this observation as indicating that the copy number of dysgenic factors (TEs) in the

male parent influences the amount of dysgenesis, and is insufficiently high in F1 males of this

cross to cause a dysgenic phenotypic in their offspring. This expectation is not predicted from the

genic hybrid incompatibility models.

Phenotypes measured to infer dysgenesis and post-zygotic reproductive isolation

The phenotypes that we measured to infer levels of post-zygotic isolation were: egg to adult

viability and the total number of progeny produced (both forms of inviability), the sex ratio of

resulting progeny, and male gonad atrophy (a form of sterility common in dysgenic systems).

These phenotypes were measured in both F1 and backcross experiments and could measure both

germline and somatic effects of TEs. After the initial pairing of individuals, the parents were

kept for seven days. On day 4 the parental individuals were transferred to a new fresh vial. On

day 7 the parents were transferred to a vial that had been previously dyed with blue food coloring

to assist counting of eggs. All vials were cleared and parental individuals discarded at day 8. To

ensure that we compared progeny of the same age among different crosses, we did not use any

progeny from the first three days of the cross because both intra-strain and inter-strain crosses

involving *D. lummei* females would often produce few or no progeny within this time. To be

used in further analyses, an individual cross must have produced at least 10 progeny over the 4

day period (days 4-7).

To estimate the total number of progeny produced and their sex ratio we counted the

number of male and female progeny from days 4-7. We determined the viability of individuals

for each cross by counting the number of eggs in the day 7 vial. We then allowed these eggs to

hatch and develop and counted the number of pupae and adults. The number of progeny

produced and egg to adult viability gave the same results; since the total number of progeny

produced was collected from more replicates and could be summarized by normal distributions

we only present those data.

To estimate male gonad atrophy we collected all flies that eclosed from days 4-6 of the

cross and saved males into new vials that we supplemented with extra yeast paste, with <10

males per vial. We aged these flies on yeast for five days before scoring gonad atrophy on a

dissecting scope by looking for the presence of atrophied testes (Blumenstiel and Hartl 2005). In

crosses between Strain 9 and Strain 160 it has been noted that both, one, or zero testes can be

atrophied (Blumenstiel and Hartl 2005). Our estimate for analysis was the proportion of testes

atrophied for the total number of testes scored. We also examined female ovary atrophy for a

small sample from each interspecies cross but did not observe any atrophy, so we focus

exclusively on male gonad atrophy.

To determine if differences in viability and total proportion of progeny produced for F1

crosses could be explained by differences in embryo lethality, we directly assayed hatchability

and also assessed evidence for fertilization versus early embryo defects via embryo staining.

First, to assay hatchability we set up population cages consisting of 25 males and 25 females

using the same genotype combinations that produced F1s described above (3 replicate

populations cages were set up per cross type). Males and females were left in the cages for one

week and were given grape agar plates supplemented with yeast culture daily. After the

acclimation period, a fresh agar plate was given to each population cage. After 24 hours the

plates were replaced (Day 1). All the eggs on the plate were counted. Seventy-two hours later

(Day 4) all eggs and larvae on the plate were counted. We calculated hatchability as the

proportion of eggs that failed to hatch/total number of eggs. This was confirmed by counting

larvae on the plates. The day after plates were collected for the hatchability assay (Day 2)

another agar plate was collected (eggs had been oviposited for 24 hours) and embryos were

collected, fixed, and stained with DAPI following basic protocols (Rothwell and Sullivan 2000;

Rothwell and Sullivan 2007; Ferree and Barbash 2009). Briefly, embryos were collected in a

nitex basket using 0.1% Triton X-100 Embryo Wash (7g NaCl, 3mL Triton X, 997 mL H20).

Embryos were dechorionated using 50% bleach for 3 minutes and rinsed well with water.

Embryos were fixed with 37% formaldehyde and n-Heptane and then vitelline membrane was

removed by serially shaking and washing in methanol. To stain embryos they were first

rehydrated by rinsing with PBS and then blocked by incubating with PBTB, followed by rinsing

with PBS before staining with PBTB+ 1:1000 DAPI for 5 minutes. Embryos were visualized

using a fluorescent scope immediately after staining to determine if development had progressed.

Embryos were considered to have begun development if we could see nuclei patterns typical of

preblastoderm, blastoderm, and gastrula stages. Since we did not score fertilization by co-

staining and looking for presence of sperm or whether the second phase of meiosis (formation of

polar bodies) had been initiated, we may have missed any very early stage embryo lethality and

considered these eggs unfertilized. Thus our estimates of embryo lethality are conservative and

they potentially underestimate embryo lethality. All females were dissected after this last embryo

collection to confirm they had mated by looking for the presence of sperm stored in the

reproductive tract.

Statistical analysis

<u>F1 reproductive isolation</u>

All analyses were performed in R v3.3.2. For traits measured in the F1 crosses we performed

three separate analyses, based on the identity of the female used in the cross.

To determine if there were differences in trait values based on the male parent in the cross we

used linear and generalized linear models. All models had the form

362
$$y = \mu + \beta_i x_i + \beta_j x_j$$

μ represents the trait value for the intra-strain cross. β_i and β_j represent the effect sizes for males

that were from different strains/species than the female in the specific analysis. For example,

when we analyzed data where TE- was the female parent, β_i could represent the deviation

caused by having the TE+ strain as the father and β_j could represent the deviation caused by

having *D. lummei* as the father. For a given phenotype there was significant reproductive

isolation when β_i or β_j was significantly less than zero, indicating lower trait value than the intra-

strain cross. In the case of sex-ratio any deviation of the correlation coefficient from zero could

be interpreted as sex specific effects. The dysgenesis and total progeny produced variables were

analyzed with simple linear regressions. Since the sex ratio data are more accurately represented

by binomial sampling we analyzed these data using binomial regression (a generalized linear

model).

To examine whether there was asymmetry in reproductive isolation for the *D. lummei* and

TE+ strain cross, it was necessary to convert these phenotypes into relative values, based on the

intra-strain cross. For example, TE+ female × *D. lummei* male measurements can be converted to

relative values by finding the difference from the TE+ intra-strain cross average value. These

relative measurements were compared in pairwise t-tests.

<u>Backcross reproductive isolation</u>

The results for the backcross experiment were analyzed in the same way as the F1 crosses. We

compared phenotypes for crosses based on the female parent. Specifically, we compared the

phenotype values for the control TE- (*D. virilis*) intra-strain cross with the crosses between TE-

females and TE+ or F1 male genotypes. We then compared the phenotype values for the control

D. lummei intra-strain cross with the crosses involving *D. lummei* females mated to TE+ or F1

male genotypes.

For a two locus genic model of incompatibility with a negative interaction in F1 progeny

between two unlinked autosomal loci, we predict there would be no sex specific effects

(Supplemental Fig. 2). When F1 males are used as parents only ½ of the resulting progeny,

regardless of sex, should exhibit the hybrid incompatibility. If the epistatic interaction does not

have complete penetrance then the effect size in this BC should be ½ compared to the parental

cross. For example if 30% of F1 progeny are dysgenic then only 15% of the backcross progeny

are expected to be dysgenic. Under a genic model with an X-autosome interaction between the X

chromosome of the TE- parent and an autosome of the TE+ parent, sex specific effects will only

be observed in the F1 progeny and not in the backcross (Supplemental Fig. 3). Lastly, for the TE

model there are no expected sex-specific effects and the level of incompatibility in the backcross

will be the same as the intra-strain control, and significantly less than what is predicted from the

genic models of incompatibility. For the gonad atrophy data, we analyzed only the data from the

F1 male that carried the X chromosome from TE+ since the other genotype was already

determined to not be significantly different from zero (see results). For the data on numbers of

progeny produced, we pooled all backcrosses for this analyses since they produced the same

number of progeny on average (see results).

Comparison of hatchability and embryo development measurements

Any decrease in progeny production could be the result of different mechanisms (prezygotic or

postzygotic isolation or both) depending on the specific cross. To determine if lower hatchability

was a product of unfertilized eggs (prezygotic isolation) or embryo lethality (postzygotic

isolation), we compared these independent measures using contingency tables. We used the

proportion of eggs that had begun development past the pre-blastoderm stage as our measure of

fertilization rate. Since there were few eggs for a given embryo collection replicate, we pooled

across all three replicates. We then compared the proportion of eggs hatching to the proportion of

eggs fertilized using a Chi-square test of independence. We could also compare differences in

the proportion of eggs fertilized between crosses that shared a common female genotype, to infer

premating isolation.

Data availability

All data and R code used is available through dryad (doi to be determined).

Results

Estimates of copy number and divergence in dysgenic causing TEs

We confirmed that candidate dysgenic TE copy numbers were consistently different

between Strain 160 (TE+) and Strain 9 (TE-) as expected (Erwin *et al.* 2015; Funikov *et al.*

2018). Specifically *Penelope, Polyphemus, Paris, Helena, Skippy,* and *Slicemaster* were

enriched in Strain 160 compared to Strain 9 (Fig 1A). Using hierarchical clustering across the *D.*

virilis strains we determined *Polyphermus* had a similar copy number distribution as *Penelope*

and formed a cluster that did not include the other candidate TEs (Fig. 1B). The correlation in

copy number between *Penelope* and *Polyphemus* across strains was r=0.68 (Fig. 1C, P=0.0073).

When one outlier (Dvir87) was removed the correlation increased to r=0.866 (P=0.0001). Either

correlation indicates a strong relationship between copy number of these two elements.

Using our copy number estimates, we identified two elements, *Polyphemus* and

Slicemaster, but not *Penelope*, that had higher average copy number in the inducing *D. virilis*

strains , Genome Strain (TE+) and Strain 160, compared to the non-inducing *D. virilis* Strain 9

(TE-) and two *D. lummei* strains. Interestingly, we were able to find copies of *Penelope* in the *D.*

lummei genome despite previous reports of the absence of this element (Zelentsova *et al.* 1999).

We used the patterns of allele frequency to assess the likelihood that *Polyphemus* and

Slicemaster were active in inducing strains but not in non-inducing strains. We expected active

elements to have homogenous insertions with very few polymorphic sites (Erwin et al. 2015).

The pattern for *Polyphemus* was consistent with our expectation for the inducing *D. virilis* strains

and non-inducing *D. virilis* strain, although there was a difference in the estimates between the

D. lummei samples (Supplemental Fig. 4)—one *D. lummei* sample was nearly identical to the

non-inducing Strain 9 (TE-), while the other strain had a higher CN and average major allele

frequency. Nonetheless, both factors were still lower in both *D. lummei* samples than in the

inducing strains (Supplemental Fig. 4). In contrast, the major allele frequency data for

Slicemaster was consistent with recent broad invasion the *D. virilis* clade because all strains,

with the exception of one *D. lummei* strain, had high average major allele frequency. As a result,

for *Slicemaster*, there was no consistent pattern differentiating inducing *D. virilis* and non-

inducing D. virilis, making it an unlikely candidate contributing to dysgenesis.

Combined, our observations from within *D. virilis* and comparison between *D. virilis* and

D. lummei suggest that copy number differences in *Polyphemus* have the potential to play

important roles in hybrid dysgenesis. This is consistent with analysis that suggest *Polyphemus*

induced DNA breaks in dysgenic progeny (Hemmer *et al.* 2019).

Dysgenesis confirmed in intraspecies crosses polymorphic for TEs

In the classic dysgenic cross (TE- female × TE+ male) we observed 38% testes atrophy (Fig. 2).

This frequency of gonad atrophy was significantly greater than the TE- intra-strain cross

(β=0.3807; *P*<0.0001). In the reciprocal non-dysgenic cross (TE+ female × TE- male) we

observed rare gonad atrophy events at a frequency not significantly different from zero. The non-

dysgenic cross produced significantly more offspring than the TE- female × TE- male intra-strain

comparison (β=26.732; *P*=0.0136); this increased productivity may reflect some outbreeding

vigor between these long-established lab lines. The number of fertilized eggs differed between

crosses so comparing the number that were actually fertilized was critical. There was no

deviation in the proportion of eggs hatching compared to the proportion fertilized in either cross,

indicating there was no embryo inviability (Table 2).

Prezygotic isolation occurs in interspecies crosses when TEs are absent

The TE- × *D. lummei* cross produced, on average, 22 fewer adult progeny than the TE- intra-

strain control cross (μ=123.318; *P*<0.0001; β=-22.509; *P*=0.0341). This reduction in progeny

production likely represents reduced fertilization and prezygotic isolation rather than reduced F1

embryo viability in this cross, based on data from two comparisons. First a comparison of the

proportion of fertilized embryos to the proportion of embryos hatching within this cross shows

that these did not differ, indicating no evidence for embryo lethality. Second, we compared the

proportion of fertilized embryos between this cross and the intra-strain control, and found that

the proportion of fertilized embryos was significantly less for the TE- × *D. lummei* cross

compared to the TE- intra-strain control cross (Table 2), consistent with a prezygotic fertilization

barrier.

In comparison, the reciprocal cross (*D. lummei* female × TE-) had no evidence of

prezygotic isolation or F1 embryo lethality. We actually observed a higher proportion of hatched

embryos compared to our estimate for proportion fertilized; this likely reflects an underestimate

of fertilization rate because this measure relies on handling embryos with multiple washing and

fixing steps which can lead to loss of embryos in the final sample (Rothwell and Sullivan

2007b). In contrast to prezygotic isolation, gonad atrophy was not observed for males or females

in either reciprocal cross between TE- and *D. lummei*.

Strong F1 postzygotic isolation occurs in interspecies crosses when TEs are present

The cross between TE+ females and *D. lummei* males produced rare male gonad atrophy,

at frequencies similar to the intraspecies non-dysgenic cross. We also saw no atrophy in the

reciprocal *D. lummei* × TE+ cross where gonad atrophy might be expected (Fig 2). Instead,

reduced viability was observed in both the reciprocal crosses, resulting in significantly fewer

progeny produced compared to control crosses (Fig 3). The TE+ female × *D. lummei* male cross

produced 19 fewer progeny compared to the TE+ intra-strain control cross (mu=50.52;

$P<0.0001$; $\beta=-31.37$; $P<0.0001$). In the reciprocal cross, *D. lummei* female × TE+ male, 17

fewer progeny were produced compared to the *D. lummei* control cross (mu=34.50; $P<0.0001$;

$\beta=-16.95$; $P=0.0027$). To test for asymmetrical effects, we compared the relative reduction in

viability in these reciprocal crosses, taking into account the differences in intra-strain and

maternal effects. For the TE+ female × *D. lummei* male cross, progeny production was reduced

by 62%; for the reciprocal cross (*D. lummei* × TE+) progeny production was reduced by 50%.

These reductions were not significantly different from one another (*t*=1.547; df=41;*P*=0.1527).

In both crosses involving *D. lummei* and TE+, the proportion of embryos hatched was

significantly lower than the proportion of embryos fertilized, indicating that the reduced number

of progeny was due to early embryo lethality in both cases (Table 2). In contrast, in the TE+

female × *D. lummei* male cross only, inviability was also specifically increased for males,

resulting in a significant excess of females and a sex ratio that was 63% female (*β*=0.1171;

P<0.0001; Fig 3).

Together our data for F1s suggested that there are possibly two mechanisms contributing

to hybrid incompatibilities in this pair. The TE model alone predicts asymmetric reproductive

isolation—specifically reduced F1 fitness in the TE+ female × *D. lummei* male cross—so our

observation of F1 inviability in both directions suggests an additional mechanism contributing in

reciprocal direction, that we also further dissected with crosses (below). This deficit of males is

not consistent with an X-autosome incompatibility between the *D. lummei* X and the TE+

autosome, because the males that are inviable carry the TE+ X chromosome.

Backcross experiments support a role for TEs in postzygotic isolation

Using data on gonad atrophy and hybrid inviability in our backcrosses, we aimed to

determine whether the observed patterns of hybrid incompatibility were more consistent with a

TE model or genic incompatibility models (Supplemental Fig. 2 and 3). First we assessed the

relationship between TE dosage and gonad atrophy with the specific strains used in this

experiment by testing whether F1 males from crosses between *D. lummei* and TE+ produced

dysgenic sons when crossed to TE- females (similar to Lovoskaya et al. 1990). For the TE model

we would expect to observe rescue of gonad atrophy in a backcross using F1 males because these

males will have fewer TEs than required to cause dysgenesis. For the genic incompatibility

models we would expect this backcross to have half the amount of gonad atrophy observed in the

F1s, because only 1/2 of their progeny would carry both incompatible alleles. The 38% F1 gonad

atrophy we observed the classic intraspecific dysgenic cross (TE- × TE+ cross) served as our

baseline expectation. We found that both backcrosses had significantly less than 19% gonad

atrophy, results that are inconsistent with the two locus genic models. When F1 males carried

the X chromosome from *D. lummei* the level of dysgenesis was not significantly different than

zero (β=0.0264; *P*=0.1366). When F1 males carried the X chromosome from TE+ the level of

dysgenesis was ~6%, significantly less than the TE+ parent (β=0.0638; *P*=0.0011; CI= 0.0261,

0.1016) and less than 19% atrophy. Therefore data for gonad atrophy in our backcrosses are

consistent with strong rescue of fertility predicted from the TE incompatibility model.

Given our results for gonad atrophy we proceeded to examine support for the different

incompatibility models in explaining the other hybrid phenotype—inviability, including sex-

specific inviability—we had observed in the interspecies crosses, using similar logic as for the

gonad atrophy phenotype. Specifically, we examined progeny production and sex ratio in

backcrosses between *D. lummei* females and F1 males. For the TE copy number model we

expect rescue of progeny viability when either F1 male is used as the male parent, and no sex

specific lethality. For an autosome-autosome model or X-autosome model, we expect to observe

~26 progeny produced, which is ½ of the reduction in progeny observed when comparing the *D.*

lummei female × TE+ male control cross (where ~17.5 progeny were produced) to the intrastrain

control cross (where ~34.5 progeny were produced). In both backcrosses the number of progeny

produced was not significantly different than the *D. lummei* × *D. lummei* control cross and

represents complete rescue of viability. The strength of this rescue is consistent with the TE copy

number model but not the two-locus incompatibility model (Table 3; Fig 3). Using a one-sample

t-test we determined that both the F1 male carrying the *D. lummei* X chromosome (mean=39.76,

se=5.26) and the F1 male carrying the X chromosome from the TE+ (mean=38.41,se=5.56)

produced significantly more progeny when crossed with the *D. lummei* female than the expected

26 progeny under the two-locus incompatibility model (Table 3).

Two-locus genic incompatibility models might be an oversimplification when

considering hybrid incompatibilities, so we also considered models with higher order (greater

than two independent/freely-recombining loci). For a three-locus autosomal interaction the

greatest expected rescue would be 1/4 of the original effect (therefore 9.5% dysgenesis with TE-

females and ~30.26 progeny produced with D. *lummei* females), while for a four-locus

autosomal interaction the largest rescue would be 1/8 the original effect (4.75% dysgenesis with

TE- females and 32.38 progeny produced with *D. lummei* females) (Supplemental Table 2).

Using a one sample t-test we rejected the hypothesis that our data can be explained by a three-

locus incompatibility for gonadal dysgenesis (H0: $\mu < 9.5$; $t = -627.2$, df = 10, $P < 0.0001$) and for

number progeny produced (H0: $\mu > 30.26$; $t=2.36$, df=28, $P=0.0126$). We were also able to reject

the hypothesis for the four-locus incompatibility for gonadal dysgenesis (H0: $\mu < 4.75$; $t = -311.48$,

df = 10, $P < 0.0001$) and number progeny produced (H0: $\mu > 32.38$; $t=1.802$, df=28, $P=0.041$). We

did not explore interactions beyond four loci because *D. virilis* only has four major autosomes

and, since we used F1 males that lack recombination, each chromosome is transmitted as a single

locus in our crosses.

In addition to progeny production, we also had data on sex ratio in our backcrosses.

Interestingly, in the cross involving *D. lummei* females and F1 males that carried the X

chromosome from TE+ there was a significant deviation from 50% female sex ratio (β =0.25;

p=0.037). This deviation could be caused by a hybrid incompatibility, however neither the TE

incompatibility model or the specific X chromosome incompatibility model we considered

predicted sex-specific effects past the F1 generation (see methods). Since we had already

observed a pattern of sex-specific inviability from the cross between TE+ females and *D. lummei*

males, we also used the backcross data to determine the genetic factors responsible for this

additional male-specific hybrid inviability. The only two other crosses where we observed a

skewed sex ratio were also female-biased and involved F1 males that carried the X chromosome

from TE+ and *D. lummei* Y crossed with *D. lummei* or TE- females (Fig. 4). In each of these

three crosses, males had different X chromosome genotypes but consistently had the *D. lummei*

Y chromosome. From this, we conclude that this additional incompatibility is caused by an

autosome-Y interaction.

Discussion

In this study, we used a powerful set of crosses to test whether the presence of TEs with

known effects on intraspecies hybrid dysgenesis also influences reproductive isolation. Using the

D. virilis dysgenic system we predicted that the *D. virilis* strain carrying dysgenic-inducing TEs

should show increased reproductive isolation when crossed with *D. lummei* lacking these TEs,

compared to the cross involving the *D. virilis* strain that lacks dysgenic-inducing TEs. Taken

together, our genetic data support the inference that TEs contribute to elevated reproductive

isolation observed in the interspecific cross that involves dysgenic TEs, and are inconsistent with

alternative hybrid incompatibility models to explain this elevated isolation. In addition, with

genomic data we also assessed which candidate TE might be responsible for observed dysgenesis

patterns among inducing *D. virilis* strains compared to non-inducing *D. virilis* and *D. lummei,*

and confirmed *Polyphemus* as a strong candidate (Funikov *et al.* 2018; Hemmer *et al.* 2019).

A role for accumulated selfish genetic elements in the expression of postzygotic

reproductive isolation is consistent with the Dobzhansky-Muller model of hybrid incompatibility

(Johnson 2010; Castillo and Moyle 2012; Crespi and Nosil 2013), but little emphasis has been

placed on the difficulty of disentangling the effects of these selfish elements from the effects of

other hybrid incompatibilities. For example, one way TEs are thought to contribute to

reproductive isolation is through transcriptional misregulation that causes sterility or inviability

(Martienssen 2010; Michalak 2010; Dion-Cote *et al.* 2014). However, this is not a unique feature

of TE-based hybrid incompatibility; divergence in trans and cis-regulatory elements are common

and can also cause misregulation and hybrid incompatibilities (reviewed in (Mack and Nachman

2017)). To disentangle TE-specific from other effects it would therefore also be necessary to

differentiate whether TE misregulation is symptomatic of a common misregulation phenomena

in hybrids or if TE divergence drives global misregulation. Instead of focusing on analyzing

patterns of misregulation, here we took the approach of isolating phenotypic effects due to TEs

by using strains with defined differences in TE abundance, and potential activity, making explicit

predictions about which hybrid classes should exhibit increased reproductive isolation, and

evaluating these using classical genetic crosses.

The strength of this approach relies on our ability to make and test predictions that are

exclusive to a TE incompatibility model, compared to alternative genic models, and thereby

differentiate effects due to TE- and non-TE causes. For example, we observed that postzygotic

incompatibility occurred only in crosses involving the TE-carrying *D. virilis* strain, and not in the

cross between TE- *D. virilis* and *D. lummei,* clearly implicating the presence of TEs in this

postzygotic isolation. For all of these incompatibilities to instead be genic would require that

they all happened to accumulate in the TE-carrying *D. virilis* strain, a scenario that is possible in

principle but biologically implausible. Nonetheless, the observation of equally strong

postzygotic isolation in both directions of the cross does suggest the additional contribution of

other non-TE factors between our *D. virilis* TE+ line and *D. lummei*.

Our ability to disentangle the effects of TEs vs non-TE incompatibilities in backcross

offspring rested on the observation that F1 males failed to produce dysgenic sons when crossed

to TE- females (Lovoskaya et al. 1990). The mechanism we propose based on previous studies

(Vieira *et al.* 1998; Srivastav and Kelleher 2017; Serrato-Capuchina *et al.* 2020b), is that F1s

have sufficiently low copy number of TEs such that dysgenesis does not occur when F1 males

are themselves used as fathers. This pattern allowed us to compare the TE incompatibility model

with genic incompatibility models because they make contrasting predictions about the level of

incompatibility expected. We found that both the rescue of non-atrophied testes in crosses

between TE- females and F1 males, and the rescue of viability in crosses between *D. lummei*

females and F1 males, is consistent with the TE incompatibility model; significantly greater

rescue was observed in these crosses than predicted by the genic incompatibility models. Using

these backcrosses we were also able to determine that additional non-TE loci contributed to the

incompatibility observed in the TE+ × *D. lummei* cross. This inviability was male specific,

suggested by a female biased sex ratio, and resulted from an autosome-Y incompatibility.

Together these observations support the operation of two distinct mechanisms of hybrid

inviability in our crosses, one of which directly implicates TEs in the expression of this

postzygotic isolating barrier.

A second inference from our results is that TEs can cause different hybrid incompatibility

phenotypes—that is, sterility or inviability—depending on context. TEs misregulation can cause

substantial cellular or DNA damage which has the potential to produce multiple different

incompatibility phenotypes (Tasnim and Kelleher ; Phadnis N. *et al.* 2015). Although the

mechanistic link between damage and hybrid incompatibility needs further exploration, these

observations suggest that the occurrence of hybrid sterility vs hybrid inviability could result from

misregulation of the same TEs at different developmental stages. Moreover, while transposable

elements have typically been thought to restrict their damage to the germline, this is not always

the case, including in the dysgenic *D. virilis* system. For example, direct evidence that TEs that

can cause inviability phenotypes when active in somatic tissue comes from a *P-element* mutant

that was engineered to have expression in somatic cells (Engels *et al.* 1987); in this case,

inviability that is dependent on *P*-element copy number was observed in crosses involving this

somatically-expressed *P-element* mutant line. Overall, it is mechanistically plausible that this

element is responsible for both of our observed incompatible phenotypes. The involvement of

TEs in both germline and somatic effects in hybrids would provide a potentially direct

mechanistic connection between the effects of a molecular change (the accumulation of TEs) and

both intraspecies dysgenesis and interspecies reproductive isolation.

Finally, our observations also identify a mechanism that could contribute to variation

among intraspecific strains in the strength of their isolation from other species. Evidence for

polymorphic incompatibilities is typically inferred from the observation that there is among-

population variation in isolation phenotypes when crossed with a second species, because the

specific alleles underlying this variation is usually unknown (Reed and Markow 2004;

Kozlowska *et al.* 2012). These polymorphic incompatibility loci are typically assumed to be

based on genic Dobzhansky-Muller incompatibilities (Cutter 2012). In this study, we show that

the presence/absence of TEs segregating in populations and species can also contribute to

intraspecific variation in the strength of isolation between species. Moreover, the rapid pace with

which TE abundance and identity can change within a lineage also indicates that such differences

could rapidly accumulate between species; this suggests that this mechanism is most likely to

influence species pairs in which there is dynamic turnover in the identity of their active TEs.

Overall, here we have provided evidence for a role of TEs in increased reproductive

isolation between lineages that differ in the presence of specific active TE families, by

leveraging known TE biology to differentiate TE vs non-TE effects in a set of targeted crosses.

The generality of when and how TEs contribute to reproductive isolation could be further

addressed by similarly focusing on other *a priori* cases where species are known to vary in the

number or identity of TEs, rather than relying on interpreting patterns of TE misregulation that

arise in hybrid crosses. For instance, increased reproductive isolation cause by transposable

elements was recently reported in crosses between *D. simulans*, that are polymorphic for *P-*

elements, and a sister species, *D. sechellia*, that completely lack *P-elements* (Serrato-Capuchina

et al. 2020a). Moreover, studies that examine lineages at a range of stages of TE differentiation

would be particularly valuable for addressing the contribution of TEs to the accumulation of

hybrid incompatibilities over evolutionary time. Our findings suggest that the rapid evolution of

TEs in different lineages could potentially explain the presence of polymorphic incompatibilities

in many systems that experience dynamic changes in these selfish genetic elements.

676
677
678   **Literature Cited**

680   Beall, E. L., M. B. Mahoney and D. C. Rio, 2002 Identification and analysis of a hyperactive
681   mutant form of Drosophila P‑element transposase. Genetics 162**:** 217-227.

Blumenstiel, J. P., 2014 Whole genome sequencing in *Drosophila virilis* identifies *Polyphemus*, a recently activated Tc1-like transposon with a possible role in hybrid dysgenesis. Mobile DNA 5: 6.

Blumenstiel, J. P., and D. L. Hartl, 2005 Evidence for maternally transmitted small interfering RNA in the repression of transposition in *Drosophila virilis*. Proceedings of the National Academy of Sciences of the United States of America 102: 15965-15970.

Brown, J. D., V. Piccuillo and R. J. O'Neill, 2012 Retroelement Demethylation Associated with Abnormal Placentation in Mus musculus x Mus caroli Hybrids. Biology of Reproduction 86: NIL_0272-NIL_0281.

Castillo, D., and L.C. Moyle, 2012 Evolutionary implications of mechanistic models of TE-mediated hybrid incompatability. International Journal of Evolutionary Biology 2012.

Castillo, D. M., and D. A. Barbash, 2017 Moving speciation genetics forward: modern techniques build on foundational studies in *Drosophila*. Genetics 207: 825-842.

Coyne, J. A., 1986 Meiotic segregation and male recombination in interspecific hybrids of *Drosophila*. Genetics 114: 485-494.

Coyne, J. A., 1989 Mutation rates in hybrids between sibling species of *Drosophila*. Heredity 63: 155-162.

Coyne, J. A., and H. A. Orr, 2004 *Speciation*. Sinauer Associates.

Crespi, B., and P. Nosil, 2013 Conflictual speciation: species formation via genomic conflict. Trends in Ecology & Evolution 28: 48-57.

Cutter, A. D., 2012 The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities. Trends in Ecology & Evolution 27.

Dion-Cote, A. M., S. Renaut, E. Normandeau and L. Bernatchez, 2014 RNA-seq Reveals Transcriptomic Shock Involving Transposable Elements Reactivation in Hybrids of Young Lake Whitefish Species. Molecular Biology and Evolution 31: 1188-1199.

Dobzhansky, T., 1937 *Genetics and the Origin of Species*. Columbia University Press.

Engels, W. R., W. K. Benz, C. R. Preston, P. L. Graham, R. W. Phillis *et al.*, Somatic effects of *P* element activity in *Drosophila melanogaster:* pupal lethality. Genetics 117: 745-757.

Erwin, A. A., M. A. Galdos, M. L. Wickersheim, C. C. Harrison, K. D. Marr *et al.*, 2015 piRNAs are associated with diverse transgenerational effects on gene and transposon expression in a hybrid dysgenic syndrome of *D. virilis*. Plos Genetics 11: e1005332.

Evgen'ev, M. B., H. Zelentsova, H. Poluectova, G. T. Lyozin, V. Veleikodvorskaja *et al.*, 2000 Mobile elements and chromosomal evolution in the virilis group of Drosophila. Proceedings of the National Academy of Sciences of the United States of America 97: 11337-11342.

Evgenev, M. B., H. Zelentsova, N. Shostak, M. Kozitsina, V. Barskyi *et al.*, 1997 Penelope, a new family of transposable elements and its possible role in hybrid dysgenesis in Drosophila virilis. Proceedings of the National Academy of Sciences of the United States of America 94: 196-201.

Ferree, P. M., and D. A. Barbash, 2009 Species-specific heterochromatin prevents mitotic chromosome segregation to cause hybrid lethality in *Drosophila*. Plos Genetics 7: e10000234.

Funikov, S. Y., D. A. Kulikova, G. S. G.S. Krasnov, A. P. A.P. Rezvykh, L. N. L.N. Chuvakova *et al.*, 2018 Spontaneous gain of susceptibility suggests a novel mechanism of resistance to hybrid dysgenesis in *Drosophila virilis*. Plos Genetics 14: e1007400.

Hemmer, L. W., G. Dias, B. Smith, K. Van Vaerenberghe, A. Howard *et al.*, 2019 The meiotic
recombination landscape of Drosophila virilis is robust to mitotic damage during hybrid
dysgenesis. bioRxiv https://doi.org/10.1101/342824.
Hey, J., 1988 Speciation via hybrid dysgenesis: Negative evidence from the *Dorosophila affinis*
subgroup. Genetica 78: 97-104.
Johnson, N. A., 2010 Hybrid incompatibility genes: remnants of a genomic battlefield? Trends in
Genetics 26: 317-325.
Khurana, J. S., J. Wang, J. Xu, B. S. Koppetsch, T. C. Thomson *et al.*, 2011 Adaptation to *P*
element transposon invasion in *Drosophila melanogaster*. Cell 147: 1551-1563.
Kidwell, M. G., 1985 Hybrid dysgenesis in *Drosophila melanogaster*: nature and inheritance of
P-element regulation. Genetics 111: 337-350.
Kozlowska, J. L., A. R. Ahmad, E. Jahesh and A. D. Cutter, 2012 Genetic variation for post-
zygotic reproductive isolation between *Caenorhabditis briggsae* and *Caenorhabditis sp.*
9. Evolution 66: 1180-1195.
Labrador, M., M. Farre, F. Utzet and A. Fontdevila, 1999 Interspecific hybridization increases
transposition rates of Osvaldo. Molecular Biology and Evolution 16: 931-937.
Lozovskaya, E. R., V. S. Scheinker and M. B. Evgenev, 1990 A hybrid dysgenesis syndrome in
Drosophila virilis. Genetics 126.
Mack, K. L., and M. W. Nachman, 2017 Gene regulation and speciation. Trends in Genetics 33:
68-80.
Martienssen, R., 2010 Heterochromatin, small RNA and post-fertilization dysgenesis in
allopolyploid and interploid hybrids of *Arabidopsis*. New Phytologist 186: 46-53.
Michalak, P., 2009 Epigenetic, transposon and small RNA determinants of hybrid dysfunctions.
Heredity 102: 45-50.
Michalak, P., 2010 An eruption of mobile elements in genomes of hybrid sunflowers. Heredity
104: 329-330.
Muller, H. J., 1942 Isolationg mechanisms, evolution and temperature. Biological Symposia 6:
71-125.
Petrov, D. A., J. L. Schutzman, D. L. Hartl and E. R. Lozovskaya, 1995 Diverse transposable
elements are mobilized in hybrid dysgenesis in *Drosophila virilis*. Proceedings of the
National Academy of Sciences of the United States of America 92: 8050-8054.
Phadnis N., E. P. E. P. Baker, J. C. J. C. Cooper, K. A. K. A. Frizzell, E. E. Hsieh *et al.*, 2015 An
essential cell cycle regulation gene causes hybrid inviability in *Drosophila*. Science 350:
1552-1555.
Presgraves, D. C., 2010 The molecular evolutionary basis of species formation. Nature Reviews
Genetics 11: 175-180.
Reed, L. K., and T. A. Markow, 2004 Early events in speciation: polymorphism for hybrid male
sterility in Drosophila. Proceedings of the National Academy of Sciences of the United
States of America 101: 9009-9012.
Rothwell, W. F., and W. Sullivan, 2000 Flourescent analysis of *Drosophila* embryos in
Drosophila Protocols, edited by S. e. al. Cold Springs Harbor Laboratory Press, Cold
Springs Harbor, NY.
Rothwell, W. F., and W. Sullivan, 2007 Drosophila embryo collection. Cold Spring Harbor
Protocols Sep 1.

Rozhkov, N. V., N. G. Schostak, E. S. Zelentsova, I. A. Yushenova, O. G. Zatsepina *et al.*, 2013
Evolution and dynamics of small RNA response to a retroelement invasion in
Drosophila. Molecular Biology and Evolution 30**:** 397-408.
Serrato-Capuchina, A., E.R.R. D'Agostino, D. Peede, B. Roy, K. Isbell *et al*., 2020a *P*-elements
strengthen reproductive isolation within the *Drosophila simulans* species complex.
bioRxiv https://doi.org/10.1101/2020.08.17.254169.
Serrato-Capuchina, A., J. Wang, E. Earley, D. Peede, K. Isbell *et al.*, 2020b Paternally inherited
P-element copy number affects the magnitude of hybrid dysgenesis in *Drosophila*
simulans and *D. melanogaster*. Genome Biology and Evolution 12**:** 808-826.
Srivastav, S. P., and E. S. Kelleher, 2017 Paternal induction of hybrid dysgenesis in *Drosophila*
melanogaster is weakly correlated with both *P-element* and *hobo* element dosage. G3 7**:**
1487-1497.
Tasnim, S., and E. S. Kelleher, *p53* is required for female germline stem cell maintenance in *P*-
element hybrid dysgenesis. Developmental Biology 434**:** 215-220.
Vieira, J., C. P. Vieira, D. L. Hartl and E. R. Lozovskaya, 1998 Factors contributing to the hybrid
dysgenesis syndrome in Drosophila virilis. Genetical Research 71**:** 109-117.
Weilguny, L., and R. Kofler, 2019 DeviaTE: assembly-free analysis and visualization of mobile
genetic element composition. Molecular Ecology Resources 19**:** 1346-1354.
Werren, J. H., 2011 Selfish genenetic elements, genetic conflict, and evolutionary innovation.
Proceedings of the National Academy of Sciences of the United States of America 108**:**
10863-10870.
Zelentsova, H., H. Poluectova, L. Mnjoian, G. Lyozin, V. Veleikodvorskaja *et al.*, 1999
Distribution and evolution of mobile elements in the *virilis* species group of *Drosophila*.
Chromosoma 108**:** 443-456.
795
796

798 Table 1. Average trait means for control intrastrain crosses. These values are used as baseline in
799 statistical models testing for increased reproductive isolation in interspecis crosses. For male
800 dysgensis (n) refers to the number of male progeny scored. se=standard error. The number of
801 replicates used to calculate the standard error are n=22 (TE-), n=23 (TE+), and n=20 (*D. lummei*)
802

Female	Male	Dysgenesis (n)	Sex Ratio (se)	Viability (se)
TE-	TE-	0.0 (1398)	0.488 (0.016)	0.844 (0.055)
TE+	TE+	0.0 (575)	0.516 (0.014)	0.571 (0.057)
D. lummei	*D. lummei*	0.0 (335)	0.521 (0.015)	0.411 (0.067)

803
Table 2. F1 postzygotic reproductive isolation, as observed in crosses between TE+ and *D.*
lummei, is a product of embryo inviability. Inviability is inferred when fewer eggs hatch than
expected based on the proprtion of eggs that initiated development (proportion fertilized). The
samples sizes (n) are the pooled number of embryos examined across replicate crosses. *
represents significant differences in the proportion fertilized from proportion hatched. †
represents a significant difference in the proportion fertilized for an interspecific cross compared
to the control cross of the maternal strain.
811

Female	Male	Proportion Fertilized (n)	Proportion Hatched (n)	χ^2_1
TE-	TE-	0.85 (179)	0.79 (344)	2.76
TE+	TE+	0.58 (155)	0.55 (224)	0.2
D. lummei	*D. lummei*	0.61 (218)	0.67 (260)	1.75
TE-	TE+	0.85 (180)	0.80 (283)	1.00
TE+	TE-	0.57 (153)	0.57 (206)	0.00
TE-	*D. lummei*	0.33† (389)	0.39 (79)	0.59
D. lummei	TE-	0.60 (161)	0.75 (337)	12.18*
TE+	*D. lummei*	0.55 (126)	0.42 (235)	5.41*
D. lummei	TE+	0.43† (212)	0.31 (208)	6.06*

Table 3. Progeny production is rescued when F1 males are crossed to *D. lummei* females. For
the backcrosses we tested whether progeny rescue was greater than the expectation for two locus
genic incompatibility models. F1-1 are males from the *D. lummei* × TE+ cross and F1-2 males
are from the reciprocal cross. For the progeny produced we report the number of replicate
crosses (n) and the standard error (se). *indicates a significant difference in progeny production
from the intrastrain control.
819

Female	Male	Mean Progeny Produced (n, se)	*t* statistic H0: mu=26 (df)	*P*-value
D. lummei	*D. lummei*	34.50 (20, 4.28)	NA	NA
D. lummei	TE+	17.54* (22, 2.78)	NA	NA
D. lummei	F1-1	39.76 (17,5.26)	2.61 (16)	0.009
D. lummei	F1-2	38.41 (12, 5.56)	2.22 (11)	0.023

820
821

822   **Figure 1.** Identification of dysgenic causing transposable elements in *D. virilis*. A) Candidate
823   TEs have higher copy number in the inducing Strain 160 compared to the noninducing Strain 9
824   (TE-). Red dashed line represents 2-fold increase in copy number. B) Clustering of candidate
825   TEs across strains of *D. virilis*. C) Positive relationship between the copy number of *Penelope*
826   and *Polyphemus* across *D. virilis* strains.
827
828
Figure 2. Gonad atrophy characteristic of the classic dysgenic phenotype. A) Intrastrain control
crosses, intraspecific crosses, and interspecific crosses demonstrating the lack of dysgenesis
except in the intraspecific cross. B) Dysgenesis in backcross progeny compared to TE- × TE+
cross showing that the F1 males do not induce dysgenesis to the same extent as the pure TE+
males. F1-1 are males from the *D. lummei* × TE+ cross and F1-2 males are from the reciprocal
cross. The thick red line is the median, and thin red lines the inter-quartile range for each
distribution   * indicates significantly different from zero (*P*<0.05). ** indicates significant
difference between GS and F1 males (*P*<0.05). The dashed line represents the expected effect
size based on the two-locus incompatibility models.
838
Figure 3. The number of progeny produced for each cross demonstrating significantly fewer
progeny produced in crosses between *D. lummei* and TE+ in both directions. A) Intrastrain
control crosses, intraspecific crosses, and interspecific crosses. B). Backcross progeny compared
to the *D. lummei* intrastrain cross showing that the F1 males do not induce progeny lethality, and
progeny production is recused. F1-1 are males from the *D. lummei* × TE+ cross and F1-2 males
are from the reciprocal cross. The thick red line is the median, and thin red lines the inter-quartile
range for each distribution * indicates significantly different from the intrastrain cross of the
maternal line (*P*<0.05).
847
Figure 4. The sex ratio of each cross used to examine deviations from a 50:50 sex ratio caused
by sex specific lethality. A) Intrastrain control crosses, intraspecific crosses, and interspecific
crosses. B). The sex ratio of backcross progeny showing sex biased progeny ratios occur when
D. lummei Y is in combination with some complement of TE+ autosomes. F1-1 are males from
the *D. lummei* × TE+ cross and F1-2 males are from the reciprocal cross. The thick red line is the
median, and thin red lines the inter-quartile range for each distribution * indicates significant
deviation from 50:50 sex ratio (*P*<0.05).
855

857 **Figure 1**

903 **Figure 2**

906   **Figure 3**

914 **Figure 4**