Main Manuscript

NODAL/TGFβ signalling mediates the self-sustained stemness induced by *PIK3CA*[H1047R] homozygosity in pluripotent stem cells

Ralitsa R. Madsen[1,2,3,8,*], James Longden[4,5], Rachel G. Knox[2,3], Xavier Robin[4], Franziska Völlmy[4], Kenneth G. Macleod[6], Larissa S. Moniz[7], Neil O. Carragher[6], Rune Linding[4,5], Bart Vanhaesebroeck[7], Robert K. Semple[1*]

[1]Centre for Cardiovascular Science, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK.
[2]Metabolic Research Laboratories, Wellcome Trust-MRC Institute of Metabolic Science, University of Cambridge, Cambridge, UK.
[3]The National Institute for Health Research Cambridge Biomedical Research Centre, Cambridge, UK.
[4]Biotech Research and Innovation Centre, University of Copenhagen, Copenhagen, Denmark.
[5]Humboldt-Universität zu Berlin, Berlin, Germany.
[6]Edinburgh Cancer Research UK Centre, Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Crewe Road South, Edinburgh, UK.
[7]University College London Cancer Institute, Paul O'Gorman Building, University College London, London, UK.
[8]Current Address: Cell Signalling, University College London Cancer Institute, Paul O'Gorman Building, University College London, London, UK.

*Corresponding authors: Ralitsa R. Madsen (R.R.M.), Robert K. Semple (R.K.S.)

Email: r.madsen@ucl.ac.uk (R.R.M.); rsemple@ed.ac.uk (R.K.S.)

0000-0001-8844-5167 (R.R.M.)

0000-0001-6539-3069 (R.K.S.)

Keywords

PI3K, *PIK3CA*, stemness, pluripotent stem cells

Author Contributions

Overall conceptualisation and study design by R.R.M. and R.K.S., with important contributions from B.V. (PI3K signalling biology, MEF data), R.L. and J.L. (total proteomics and WGCNA). R.R.M. and R.G.K. performed all hPSC experiments. F.V. performed the mass spectrometry experiments, and X.R. performed MCMC computational analysis. L.M. performed the MEF experiment. K.M. and N.C. were responsible for RPPA sample processing, and R.R.M carried out statistical analysis. R.R.M. performed all RNA sequencing quantitation and IPA analyses on transcriptomic and proteomic datasets. R.R.M., X.R., J.L., F.V. and R.G.K. were responsible for data curation. R.R.M., B.V. and R.K.S. wrote the manuscript. R.L., J.L., N.C., X.R., L.M., and F.V. reviewed and edited the final version.

This PDF file includes:

       Main Text
       Figures 1 to 5

Abstract

Activating *PIK3CA* mutations are known "drivers" of human cancer and developmental overgrowth syndromes.
We recently demonstrated that the "hotspot" *PIK3CA^{H1047R}* variant exerts unexpected allele dose-dependent
effects on stemness in human pluripotent stem cells (hPSCs). In the present study, we combine high-depth
transcriptomics, total proteomics and reverse-phase protein arrays to reveal potentially disease-related
alterations in heterozygous cells, and to assess the contribution of activated TGFβ signalling to the stemness
phenotype of *PIK3CA^{H1047R}* homozygous cells. We demonstrate signalling rewiring as a function of oncogenic
PI3K signalling dose, and provide experimental evidence that self-sustained stemness is causally related to
enhanced autocrine NODAL/TGFβ signalling. A significant transcriptomic signature of TGFβ pathway activation
in *PIK3CA^{H1047R}* heterozygous was observed but was modest and was not associated with the stemness
phenotype seen in homozygous mutants. Notably, the stemness gene expression in *PIK3CA^{H1047R}*
homozygous iPSCs was reversed by pharmacological inhibition of TGFβ signalling, but not by pharmacological
PI3Kα pathway inhibition. Altogether, this provides the first in-depth analysis of PI3K signalling in human
pluripotent stem cells and directly links dose-dependent PI3K activation to developmental NODAL/TGFβ
signalling.
57

Introduction

Class IA phosphoinositide 3-kinases (PI3Ks) are evolutionarily conserved enzymes that catalyse formation of the membrane-bound second messenger phosphatidylinositol-3,4,5-trisphosphate (PIP_3). PI3Ks are activated downstream of receptor tyrosine kinases, with the ensuing increase in PIP_3 and its derivative $PI(3,4)P_2$ triggering a widespread signalling network, best known for the activation of the serine/threonine kinases AKT and mTORC1. PI3K activation promotes cell survival, glucose uptake, anabolic metabolism, cell proliferation and cell migration (1). Among the class IA PI3K isoforms (PI3Kα, PI3Kβ, PI3Kδ), the ubiquitously-expressed PI3Kα (encoded by the *PIK3CA* gene in humans), is the main regulator of organismal growth, development and survival (2).

Activating mutations in *PIK3CA* are among the most common somatic point mutations in cancer, together with inactivation or loss of the tumour suppressor *PTEN* (a negative regulator of PI3K) (3–5). The same mutations in *PIK3CA*, when acquired postzygotically during development, also cause a range of largely benign overgrowth disorders, for which the term *PIK3CA*-related overgrowth spectrum (PROS) has been coined (6). Motivated by the need to understand the role of PI3K signalling in a human developmental context, we previously generated an allelic series of human induced pluripotent stem cells (iPSCs) with heterozygous or homozygous expression of the activating mutation *PIK3CA^H1047R^*, the most commonly observed *PIK3CA* mutation in both cancer and PROS (7). Despite the severe developmental disorders caused by heterozygosity for *PIK3CA^H1047R^* in humans *in vivo*, we found little discernible effect on germ layer specification from heterozygous iPSCs. In sharp contrast, homozygosity for *PIK3CA^H1047R^* led to self-sustained stemness and resistance to spontaneous differentiation *in vitro* and *in vivo* (7). This suggested a previously unappreciated quantitative relationship between the strength of PI3K signalling and the gene regulatory network (GRN) in pluripotent stem cells.

The core pluripotency GRN features a feedforward, autoregulatory circuit comprising three transcription factors, namely SRY box 2 (SOX2), Octamer-binding transcription factor 3/4 (OCT3/4; encoded by POU5F1), and the homeobox transcription factor NANOG (8–10). SOX2 helps sustain OCT3/4 expression, which is required for establishment and maintenance of the pluripotent state (11). However, even modest overexpression of OCT3/4 destabilises the pluripotency network and triggers differentiation (12, 13). In contrast, NANOG, while dispensable for maintenance of pluripotency (14), stabilises the pluripotency gene regulatory network. Overexpression of NANOG by as little as 1.5-fold leads to sustained self-renewal (or "stemness") of murine and human PSCs (15–18). In hPSCs, NANOG expression is activated by the transcription factors SMAD2/3 (19), which in turn are activated by receptors binding TGFβ, Activin or NODAL (20). Overexpression of NODAL thus results in self-sustained stemness of hPSCs even in differentiation-promoting conditions (21, 22).

Given the unexpected and surprisingly mild phenotype caused by heterozygous *PIK3CA^H1047R^* expression in iPSCs, we reasoned that more sensitive assays would allow us to discern small but disease-relevant alterations in these cells. Thus, in this study, we first applied high depth transcriptomics, and proteomics to seek evidence of disease-related phenotypes in heterozygous cells, and to investigate how high-dose PI3K signalling leads to self-sustained stemness in homozygous *PIK3CA^H1047R^* iPSCs. We demonstrate that heterozygous cells do exhibit significant transcriptomic changes, although these are a weak echo of the widespread changes seen in homozygous cells. The mild transcriptional consequences of heterozygous expression of disease-relevant *PIK3CA* mutations were also validated in additional model systems and contrast with previous findings of major transcriptional rewiring in immortalised, non-transformed breast epithelial cells (23, 24). We demonstrate that the stemness phenotype of *PIK3CA^H1047R/H1047R^* iPSCs is maintained by self-sustained NODAL/TGFβ signalling, in line with increased *PIK3CA*-mediated *NODAL* expression, and that it is not reversible by PI3Kα-specific inhibition. This work provides the first in-depth characterisation of dose-dependent PI3K signalling effects in hPSCs and establishes dose-dependent PI3Kα-induced NODAL/TGFβ signalling as the main mechanism for self-sustained stemness in homozygous *PIK3CA^H1047R^* iPSCs. We discuss the implications of our findings for understanding developmental disorders and cancers driven by genetic PI3K activation.

Results

A sharp PI3K activity threshold determines gene expression changes in *PIK3CA^H1047R^* iPSCs

109   We previously generated isogenic human iPSCs with heterozygous or homozygous knock-in of the
110   "hotspot" *PIK3CA^{H1047R}* mutation. Surprisingly, heterozygous cells showed few phenotypic changes and
111   differentially expressed protein-coding transcripts. In contrast, homozygous *PIK3CA^{H1047R/H1047R}* cells exhibited
112   marked morphological changes and altered gene expression, with strong enrichment for cancer-associated
113   pathways (7).

114   To substantiate the apparent PI3K activity threshold manifest in *PIK3CA^{H1047R}*-driven gene
expression changes, and to look for further disease-related changes in heterozygous cells, we undertook RNA
sequencing at substantially greater depth, also increasing the sample size to four independently-derived,
previously unstudied iPSC cultures for each *PIK3CA* genotype. As before, homozygous mutant cells clearly
separated from heterozygous and wild-type cells, which overlapped on multidimensional scaling (**Fig. 1A**), but
we now detected a reduction in the levels of 451 transcripts and an increase in the levels of 710 transcripts in
PIK3CA^{WT/H1047R} iPSCs (**Fig. 1B**). This dropped to 149 and 343 transcripts, respectively, after applying a fold-
change cut-off of 1.3 (**Fig. 1B and Dataset S1**), indicative of the small magnitude of many expression changes
in heterozygous mutants (**Fig. S1A**). Use of the same cut-off of 1.3, in sharp distinction, yielded 2873 and 2771
transcripts of decreased or increased abundance, respectively, in homozygous iPSC mutants (**Fig. 1B and**
Dataset S2). Not only was the number of gene expression changes higher by an order of magnitude in
homozygous cells, but many expression changes were large compared to wild-type controls (**Fig. S1A**). The
magnitudes of gene expression changes in *PIK3CA^{H1047R/H1047R}* cells correlated strongly with our previous
findings (Spearman's rho = 0.74, p < 2e-16) (**Fig. S1B**), whereas correlation was low (Spearman's rho = 0.1, p
< 2e-16) for *PIK3CA^{WT/H1047R}* iPSCs (**Fig. S1C**).

129   Given prior reports that *PIK3CA^{H1047R}* heterozygosity in breast epithelial cells extensively remodels
gene expression (23, 24), we undertook further transcriptional profiling in two unrelated cellular models of
genetic *PIK3CA* activation. First, we examined iPSCs derived from a woman with clinically obvious but mild
PROS due to mosaicism for *PIK3CA^{E418K}* (**Fig. 1C**) (25). Heterozygous iPSCs were compared to wild-type lines
established simultaneously from dermal fibroblasts from the same skin biopsy, which is possible due to genetic
mosaicism of the sampled skin. Like *PIK3CA^{WT/H1047R}* iPSCs, *PIK3CA^{WT/E418K}* iPSCs closely clustered with
isogenic wild-type controls on multidimensional scaling (MDS) plotting (**Fig. 1D**), with only 30 differentially
expressed genes (**Dataset S3**). We also studied previously reported *Pik3ca^{WT/H1047R}* mouse embryonic
fibroblasts (MEFs) 48 h after *Cre*-mediated *Pik3ca^{H1047R}* induction (26). Wild-type and *Pik3ca^{WT/H1047R}* MEFs
were superimposable on an MDS plot (**Fig. 1E**), with only 192 downregulated and 77 upregulated genes
(**Dataset S4**). Our findings suggest that there are *bona fide* transcriptional changes induced by heterozygosity
for *PIK3CA^{H1047R}*, but these are dramatically smaller in number and magnitude than changes induced by
homozygosity for *PIK3CA^{H1047R}*.

142   To assess whether transcriptional changes observed in iPSCs were mirrored in the proteome, we
applied label-free proteomics to the iPSC lines used in our previous study (7). Around 4,600 protein ratios were
obtained for both heterozygous *versus* wild-type and homozygous *versus* wild-type iPSC comparisons, as
estimated using a novel Bayesian approach based on the Markov Chain Monte Carlo (MCMC) method (27).
In contrast to other algorithms, the MCMC method generates an error estimate alongside each protein
concentration which permits more confident determination of proteins with the most robust differential
expression. The number of differentially-expressed proteins correlated with *PIK3CA^{H1047R}* allele dosage, with 54
and 258 differentially expressed proteins in *PIK3CA^{WT/H1047R}* and *PIK3CA^{H1047R/H1047R}* cells, respectively (**Fig. 1F,**
Datasets S5 and S6). Of these, 27 proteins were differentially expressed in both heterozygous and
homozygous *PIK3CA^{H1047R}* iPSCs (**Dataset S7**), with 16 changing in opposite directions (**Fig. 1F**). There was
a strong correlation between differentially-expressed proteins and corresponding transcripts in
PIK3CA^{H1047R/H1047R} iPSCs (**Fig. S2A, S2B**), but not in heterozygous mutants (**Fig. S2C, S2D**). As for the
relatively weak correlation seen between transcriptomic experiments for heterozygous cells, this likely reflects
the small magnitude of gene expression changes induced by heterozygous *PIK3CA^{H1047R}* (**Fig. 1B, S1C**).

156   Collectively, these findings corroborate the existence of a threshold of PI3K pathway activity which
determines the large majority of gene expression changes in *PIK3CA^{H1047R/H1047R}* iPSCs in a near-binary
manner. While deeper sequencing did reveal statistically significant gene expression changes in heterozygous
iPSCs, and while these changes may contribute to growth-related phenotypes in PROS when sustained across
development, effect sizes were modest and more variable. Similar findings in heterozygous MEFs suggest that
this may be generalisable to differentiated cell types, irrespective of species. This consolidates the view that only
homozygosity for *PIK3CA^{H1047R}* results in robust and widespread transcriptional changes in otherwise normal,

diploid cells, arguing against a universal "butterfly" effect of heterozygosity suggested based on studies of a
genetically abnormal breast epithelial cell line (23, 24).

***PIK3CA^H1047R/H1047R* iPSCs show evidence of signalling "rewiring"**

We previously demonstrated a graded increase in AKT (S473) phosphorylation across heterozygous
and homozygous *PIK3CA^H1047R* iPSCs (7). To assess in more detail whether the near-binary gene expression
difference between heterozygous and homozygous *PIK3CA^H1047R* cells is underpinned by corresponding
differences in indices of PI3K pathway activation, we profiled phosphorylation of a wider repertoire of pathway
components using reverse phase phosphoprotein array (RPPA) technology.
Changes in protein phosphorylation were surprisingly modest, with the largest change a two-fold
increase in AKT phosphorylation (on S473 and T308) in *PIK3CA^H1047R/H1047R* cells. Contrasting with the near-
binary response seen at the transcriptional level, heterozygous and homozygous *PIK3CA^H1047R* expression
generally produced graded phosphorylation of PI3K pathway components, with slightly higher levels in
homozygous iPSCs (**Fig. 2A**). None of the mutant genotypes showed consistently increased phosphorylation
of the mTORC1 target P70S6K or its downstream substrate S6 (**Fig. S3A**), perhaps reflecting saturation at this
level of the pathway due to other stimuli for mTORC1 in the complete culture medium (e.g. amino acids) (28).
When deprived of growth factors for 1 h prior to RPPA profiling, both heterozygous and homozygous mutant
did exhibit increased P70S6K phosphorylation, whereas S6 phosphorylation remained similar to wild-type cells
(**Fig. 2B**).
Inhibition of PI3Kα activity with the PI3Kα-selective inhibitor BYL719 for 24 h fully reversed canonical
PI3K signalling-related changes in phosphorylation of downstream proteins including AKT, GSK3, FOXO1,
TSC2 and P70S6K (**Fig. 2B**). Consistent with these signalling changes, we previously showed that the same
dose of BYL719 (100 nM) abolishes the increased tolerance to growth factor deprivation-induced death
conferred by heterozygous or homozygous *PIK3CA^H1047R* in iPSCs (7). Despite its effects on the primary PI3K
signalling cascade, PI3Kα inhibition failed to reverse other changes observed in *PIK3CA^H1047R/H1047R* iPSCs,
including increased phosphorylation of SMAD2 and ERK1/2 and increased expression of c-MYC and IGF1R
(**Fig. 2B, Fig. S3B**). This suggests signalling rewiring in *PIK3CA^H1047R/H1047R* iPSCs that is partially resistant to
relatively short-term inhibition of the inducing stimulus.

Pathway and network analyses implicate TGFβ signalling in *PIK3CA^H1047R* dose-dependent stemness

Pathway and network analyses were next applied to proteomic and transcriptomic data to identify
candidate mechanism(s) mediating *PIK3CA^H1047R* dose-dependent stemness. Consistent with our previous
study (7), TGFβ1 was again the most significant predicted upstream activator according to Ingenuity® Pathway
Analysis (IPA) of the top 2000 upregulated and top 2000 downregulated transcripts in *PIK3CA^H1047R/H1047R*
iPSCs (**Fig. 3A**). TGFβ1 was also the most significant upstream activator predicted by analysis of
PIK3CA^H1047R/H1047R proteomic data (**Fig. 3B**). This strongly suggests activation of the TGFβ pathway in
homozygous *PIK3CA^H1047R* iPSCs.
Although *PIK3CA^WT/H1047R* iPSCs showed around 10-fold fewer differentially expressed genes than
homozygous iPSC cells, IPA in heterozygous iPSCs also revealed multiple TGFβ pathway-related stimuli
among predicted upstream activators (**Fig. 3C**). Moreover, TGFβ1 was predicted as one of only two significant
upstream activators when analysis was performed on genes concordantly differentially expressed (N = 180) in
PIK3CA^H1047R mutant iPSCs *versus* wild-type controls (**Fig. 3C and Dataset S8**).
The other significant upstream regulator common to heterozygous and homozygous *PIK3CA^H1047R* was
MAPK1 (also known as ERK2), consistent with RPPA findings and immunoblot evidence of increased ERK
kinase phosphorylation in *PIK3CA^H1047R* mutant iPSCs (Ref. (7), **Fig. 2A and Fig. S3A**). The significance of
predicted TGFβ activation in heterozygous *PIK3CA^H1047R* iPSCs (overlap p-value = 1.7e-05) was much lower
than in homozygous (overlap p-value = 4.3e-21) mutants. This points towards a critical role for the TGFβ
pathway in mediating the allele dose-dependent effect of *PIK3CA^H1047R* in human iPSCs.
To complement IPA analysis, which is based on highly curated, proprietary datasets, we undertook non-
hypothesis-based Weighted Gene Correlation Network Analysis (WGCNA) – a network-based data reduction
method that seeks to determine gene correlation patterns across multiple samples, irrespective of the function
of individual genes (29). Using all transcripts expressed in wild-type, heterozygous and homozygous
PIK3CA^H1047R iPSCs (**Fig. 4A**), this analysis returned 43 modules (or clusters) of highly interconnected genes

(**Fig. 4B**). Of the two modules with the highest correlation with the homozygous trait, one showed enrichment
for several KEGG pathway terms relevant to stemness of *PIK3CA^{H1047R/H1047R}* iPSCs, notably including
"Signalling pathways regulating pluripotency in stem cells" (**Fig. 4C**).

Given prior evidence of strong activation of TGFβ signalling in homozygous mutant cells, we next
constructed the minimal network of differentially expressed genes in *PIK3CA^{H1047R/H1047R}* iPSCs that linked
pluripotency, PI3K and TGFβ signalling pathways (**Fig. 4D**). This approach allowed us to navigate the signalling
rewiring and to link strong PI3K pathway activation, stemness and TGFβ signalling in an unbiased manner.
Indeed, the resulting network exhibited high interconnectivity, with multiple shared nodes across all three
pathways, suggesting close crosstalk between PI3K and TGFβ signalling in stemness regulation. That most
nodes represented genes with increased expression in homozygous mutants strengthens the notion that strong
oncogenic PI3Kα activation stabilises the pluripotency network in human iPSCs. The MYC oncogene stood out
as the only network node intersecting with all three signalling pathways, suggesting it may comprise a key
mechanistic link in the observed phenotype.

Inhibition of TGFβ signalling destabilises the pluripotency gene network in *PIK3CA^{H1047R/H1047R}* iPSCs

TGFβ signalling plays a critical role in pluripotency regulation (19, 22, 30), and a differentiation-resistant
phenotype has been reported in *NODAL*-overexpressing iPSCs (21). Together with increased *NODAL*
expression in homozygous *PIK3CA^{H1047R}* iPSCs and computational identification of enhanced TGFβ pathway
activity in PI3K-driven "constitutive" stemness (Ref. (7) and current study), this led us to hypothesise that strong
PI3Kα-dependent induction of *NODAL* underlies establishment of the differentiation-resistant phenotype of
homozygous *PIK3CA^{H1047R}* iPSCs. Specifically, we hypothesised that autocrine NODAL enhances TGFβ
signalling in *PIK3CA^{H1047R/H1047R}* iPSCs, with resulting increased *NANOG* expression "locking" the cells in
perpetual stemness (19).

Testing this hypothesis in iPSCs is challenging for biological and technical reasons, including lack of
specific pharmacological inhibitors of NODAL, and difficulty in detecting subtle early phenotypic consequences
of partial destabilisation of the iPSC pluripotency gene regulatory network. Moreover, the widely adopted
maintenance medium and coating substrate we used for cell culture both contain TGFβ ligands (31, 32), which
may mask effects of *NODAL* repression by PI3Kα-specific inhibition. We previously found that treatment of
PIK3CA^{H1047R/H1047R} iPSCs in this 'complete' maintenance medium with 500 nM BYL719 reduces *NODAL*
mRNA expression within 24 h, but has no discernible effect on increased *NANOG* mRNA levels (7).

To minimise confounding effects of exogenous TGFβ ligands, we prepared medium with and without
recombinant NODAL supplementation, and assessed expression of *NODAL* and *NANOG* as surrogate
markers of stemness over 72 h of culture. We also reduced the BYL719 concentration to 250 nM given
increased iPSC toxicity observed with 500 nM BYL719 (7); and pilot experiments (not shown) in which 24 h
treatment with 250 nM but not 100 nM BYL719 in complete medium reduced *NODAL* mRNA expression in
PIK3CA^{H1047R/H1047R} iPSC clones. Within 48 h, exclusion of NODAL from the medium resulted in the expected
downregulation of *NODAL* and *NANOG* expression in wild-type iPSCs, and this was greater still at 72 h (**Fig. 5**
and Fig. S4A). In *PIK3CA^{H1047R/H1047R}* iPSCs, however, *NODAL* removal had no effect on the increased *NODAL*
and *NANOG* expression (**Fig. 5 and Fig. S4A**), in line with a self-sustained stemness phenotype. Exposure of
NODAL-free *PIK3CA^{H1047R/H1047R}* cultures to 250 nM BYL719 had a visible colony growth-inhibitory effect (**Fig.**
S5) and decreased *NODAL* expression within 24 h, and this continued to decrease subsequently (**Fig. 5**). This
is consistent with NODAL's known ability to control its own expression through a feed-forward loop (33). Despite
a 55% reduction in *NODAL* mRNA after 72 h, however, little effect on *NANOG* expression was seen (**Fig. 5**).
This may reflect the short time course studied (to avoid confounding effect of passaging), or the exquisite
sensitivity of iPSCs to residual upregulation of *NODAL* in homozygous *PIK3CA^{H1047R}* iPSCs. This may be
compounded by residual low levels of TGFβ-like ligands in the coating substrate, or possibly by increased
expression of two other TGFβ superfamily ligands, *GDF3* and *TGFB2*, observed in homozygous mutant cells
(**Dataset S2**).

To confirm that TGFβ signalling is required for maintenance of stemness in *PIK3CA^{H1047R/H1047R}* iPSCs,
the cells were treated with SB431542 – a specific inhibitor of TGFβ and NODAL type I receptors (34). This
completely repressed *NODAL* expression within 24 h, accompanied by downregulation of *NANOG* expression
(**Fig. 5**). A similar effect was observed on *POU5F1* expression, consistent with destabilisation of the pluripotency

gene regulatory network in *PIK3CA^H1047R/H1047R* iPSCs (**Fig. 5**). Confirming this, we used a lineage-specific gene
expression array to demonstrate similar reduction in expression of several other well-established stemness
markers (*MYC*, *FGF4*, *GDF3*) with increased expression in *PIK3CA^H1047R/H1047R* iPSCs, performing the analysis
after 48 h of TGFβ pathway inhibition (**Fig. S4B**). Despite the short treatment, we found evidence for the
expected neuroectoderm induction upon inhibition of the TGFβ pathway (21, 35), reflected by increased
expression of *CDH9*, *MAP2*, *OLFM3* and *PAPLN* (**Fig. S4B**).
Collectively, these data suggest that the stemness phenotype of *PIK3CA^H1047R/H1047R* iPSCs is
mediated by self-sustained TGFβ signalling, most likely through PI3K dose-dependent increase in *NODAL*
expression, and that this is amenable to reversal through inhibition of the TGFβ pathway but not of PI3Kα itself.
277

Discussion

PIK3CA^H1047R is the most common activating *PIK3CA* mutation in human cancers and in PROS (6).
We recently found that *PIK3CA*-associated cancers often harbour multiple mutated *PIK3CA* copies, and
demonstrated that homozygosity but not heterozygosity for *PIK3CA^H1047R* leads to self-sustained stemness in
human pluripotent stem cells (hPSCs) (7). High-depth transcriptomics in this study confirmed that
heterozygosity for *PIK3CA^H1047R* induces significant but very modest transcriptional changes; observed both in
CRISPR-edited hPSCs with long-term *PIK3CA^H1047R* expression and in mouse embryonic fibroblasts (MEFs)
upon acute *PIK3CA^H1047R* induction by *Cre*, with canonical PI3K pathway activation seen in both cases (current
study and Ref. (7, 26, 36)). Similarly, hPSCs with heterozygous expression of *PIK3CA^E418K*, a "non-hotspot"
mutation, were transcriptionally indistinguishable from their isogenic wild-type controls. In contrast to the mild
transcriptional consequences of these heterozygous variants, however, homozygosity for *PIK3CA^H1047R* was
associated with differential expression of nearly a third of the hPSC transcriptome, suggesting widespread
epigenetic reprogramming. This near-binary response is not a consequence of a similar quantitative difference
in PI3K pathway activation, as assessed by phosphoprotein profiling, which instead showed a relatively modest
and graded increase in homozygous *versus* heterozygous *PIK3CA^H1047R* hPSCs. This implies that the apparent
sharp PI3K signalling threshold that determines the cellular response in hPSCs is "decoded" distal to the
canonical pathway activation.
Using a combination of computational analyses and targeted experiments, the current study further
provides evidence for self-sustained TGFβ pathway activation as the main mechanism through which
PIK3CA^H1047R homozygosity "locks" hPSCs in a differentiation-resistant state that has become independent of
the driver mutation and the associated PI3K pathway activation. We suggest that homozygosity but not
heterozygosity for *PIK3CA^H1047R* promotes sufficient TGFβ pathway activity to induce increased *NODAL* and
downstream *NANOG* expression to levels that stabilise the stem cell state, yet are not high enough to tip the
balance towards mesendoderm differentiation (7). Exactly how PI3K activation regulates *NODAL* expression
remains unknown. A potential mechanism involves increased expression of the stem cell reprogramming factor
MYC, which was observed at both mRNA and protein level in homozygous but not heterozygous *PIK3CA^H1047R*
iPSCs. Furthermore, MYC was the only node in the WGCNA-based network of pluripotency, PI3K and TGFβ
pathway components that was shared by all three pathways (**Fig. 4D**). MYC has previously been shown to
exert oncogenic effects that depend on a sharp threshold of MYC expression, reminiscent of the effects we
observe for dose-dependent *PIK3CA* activation (37). Elevated MYC has also been shown to allow
PIK3CA^H1047R-induced murine breast cancers to become independent of continuous *PIK3CA^H1047R* expression
(38).
Stabilisation of the stemness phenotype in hPSCs by strong genetic PI3K pathway activation may
be generalisable beyond the iPSC model system. BYL719 (alpelisib; Novartis), the PI3Kα-selective inhibitor
used in our cellular studies, was recently approved for use in combination with anti-estrogen therapy in ER-
positive breast cancers (39). In a separate study focusing on human breast cancer (Madsen *et al.*, manuscript
submitted), we have described use of computational analyses to demonstrate a strong, positive relationship
between a transcriptomically-derived PI3K activity score, stemness gene expression and tumour grade in
breast cancer. Prior reports have suggested a role for NODAL in driving breast cancer stemness and
aggressive disease (40, 41), with potential links to mTORC1 activation (42, 43). Our findings that BYL719 fails
to fully reverse the increased *NODAL* and stemness gene expression in homozygous *PIK3CA^H1047R* iPSCs
suggests that inhibition of TGFβ signalling as a pro-differentiation therapy warrants investigation as co-therapy

with PI3K inhibitors in breast tumours with strong PI3K pathway activation. Finally, the lack of widespread
transcriptional changes upon heterozygous expression of mutant *PIK3CA* in otherwise genetically normal cell
models may explain the low oncogenicity of this genotype in isolation *in vivo*.
323

Materials and Methods

326   All cell lines used in this study are listed in **Table S1**. Unless stated otherwise, standard chemicals
327   were acquired from Sigma Aldrich, with details for the remaining reagents included in **Table S2**.

iPSC culture and treatments

Maintenance

The derivation of the iPSC lines, including associated ethics statements, has been
described previously (7). All lines were grown at 37°C and 5% CO_2 in Essential 8 Flex (E8/F)
medium on Geltrex-coated plates, in the absence of antibiotics. For maintenance, cells at 70-90%
confluency were passaged as aggregates with ReLeSR, using E8 supplemented with RevitaCell
(E8/F+R) during the first 24 h to promote survival. A detailed version of this protocol is available via
protocols.io (doi: dx.doi.org/10.17504/protocols.io.4rtgv6n).
All cell lines were tested negative for mycoplasma and genotyped routinely to rule out
cross-contamination during prolonged culture. Short tandem repeat profiling was not performed. All
experiments were performed on cells within 10 passages since thawing.
339
Collection for RNA sequencing and total proteomics

For RNA sequencing and total proteomics, subconfluent cells were fed fresh E8/F 3 h prior to snap-
freezing on dry ice and subsequent RNA or protein extraction. Relative to the results in Ref. (7),
the current transcriptomic data of *PIK3CA^H1047R* were obtained more than 6 months following the
first study, with cells at different passages, and were thus independent from one another. Moreover,
sample collection for the second transcriptomics experiment was conducted over three days
according to a block design, thus allowing us to determine transcriptional differences that are robust
to biological variability.

Cell lysate collection for RPPA

For RPPA in growth factor-replete conditions, cells were fed fresh E8/F 3h before
collection. To assess variability due to differences in collection timing, clones from each iPSC
genotype were collected on each one of three days according to a block design, giving rise to a
total of 22 cultures. To test the effect of the PI3Kα-specific inhibitor BYL719, cells were treated with
100 nM drug (or DMSO only as control treatment) for 24 h and exposed to growth factor removal
within the last hour before collection. All cells were washed in DPBS prior to collection to rinse off
residual proteins and cell debris.

TGFβ/NODAL signalling studies

Wild-type or homozygous *PIK3CA^H1047R* iPSCs were seeded in 12-well plates all coated
with Geltrex from the same lot (#2052962; diluted in DMEM/F12 lot #RNBH0692). Cells were
processed for seeding at a ratio of 1:15 according to the standard maintenance protocol. One day
after seeding, individual treatments were applied to triplicate wells. Briefly, cells were first washed
twice with 2 and 1 ml of Dulbecco's PBS (DPBS) to remove residual growth factors. The base
medium for individual treatments was Essential 6 supplemented with 10 ng/ml heat-stable FGF2.

This was combined with one of the following reagents or their diluent equivalents: 100 ng/ml
NODAL (diluent: 4 mM HCl), 250 nM BYL719 (diluent: DMSO), 5 μM SB431542 (diluent: DMSO).
Cells were snap-frozen on dry ice after 24, 48 and 72 h following a single DPBS wash. Individual
treatments were replenished daily at the same time of day to limit temporal confounders.

Mouse embryonic fibroblast (MEF) culture

368 The derivation and culture of the wild-type and $PIK3CA^{WT/H1047R}$ MEFs used in this study
369 have been reported previously (26). Cell pellets were collected on dry ice 48 h after induction of
370 heterozygous $PIK3CA^{H1047R}$ expression, without prior starvation.

RNA sequencing

Induced pluripotent stem cell lysates were collected in QIAzol and processed for RNA
extraction with the DirectZol Kit as per the manufacturer's instructions. The final RNA was subjected
to quantification and quality assessment on an Agilent Bioanalyzer using the RNA 6000 Nano Kit,
confirming that all samples had a RIN score of 10. For $PIK3CA^{H1047R}$ iPSCs and corresponding
wild-types, an Illumina TruSeq Stranded mRNA Library Prep Kit was used to synthesise 150-bp-
long paired-end mRNA libraries, followed by sequencing on an Illumina HiSeq 4000, with average
depth of 70 million reads per sample. $PIK3CA^{WT/E418K}$ and isogenic control iPSCs were subjected
to 50-bp-long single-end RNA sequencing (RNAseq) at an average depth of 20 million reads per
sample.
MEF RNA was extracted using Qiagen's RNAeasy miniprep (with QIAshredder). All
samples had a confirmed Agilent Bioanalyzer RIN score of 10. An Illumina TruSeq Unstranded
mRNA kit was used to prepare 100-bp-long paired-end libraries, followed by Illumina HiSeq 2000
sequencing.
Details of the subsequent data analyses (raw read mapping, counting, statistical testing,
pathway and network analyses) are provided in SI Appendix.

Label-free total proteomics

Sample preparation

Cells were cultured to subconfluence in Geltrex-coated T175 flasks, and protein was
harvested by lysis in 3 ml modified RIPA buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% NP-40,
0.5% Na-deoxycholate, 1 mM EDTA) supplemented with phosphatase inhibitors (5 mM ß-
glycerophosphate, 5 mM NaF, 1 mM Na$_3$VO$_4$) and protease inhibitors (Roche cOmplete ULTRA
Tablets, EDTA-free). The lysates were sonicated on ice (4x 10s bursts, amplitude = 60%; Bandelin
Sonopuls HD2070 sonicator) and spun down for 20 min at 4300g. Ice-cold acetone was added to
the supernatant to achieve a final concentration of 80% acetone, and protein was left to precipitate
overnight at -20°C. Precipitated protein was pelleted by centrifugation at 2000g for 5 min and
solubilised in 6 M urea, 2 M thiourea, 10 mM HEPES pH 8.0. Protein was quantified using the
Bradford assay and 8 mg of each sample were reduced with 1 mM dithiothritol, alkylated with 5 mM
chloroacetamide and digested with endopeptidase Lys-C (1:200 v/v) for 3 h. Samples were diluted
to 1 mg/ml protein using 50 mM ammonium bicarbonate and incubated overnight with trypsin (1:200
v/v). Digested samples were acidified and urea removed using SepPak C18 cartridges. Peptides
were eluted, and an aliquot of 100 µg set aside for total proteome analysis. The peptides were
quantified using the Pierce quantitative colorimetric peptide assay. The equalised peptide amounts
were lyophilised and resolubilised in 2% acetonitrile and 1% trifluoroacetic acid in order to achieve
a final 2 µg on-column peptide load.

Mass spectrometry (MS) data acquisition

All spectra were acquired on an Orbitrap Fusion Tribrid mass spectrometer (Thermo Fisher
Scientific) operated in data-dependent mode coupled to an EASY-nLC 1200 liquid chromatography
pump (Thermo Fisher Scientific) and separated on a 50 cm reversed phase column (Thermo Fisher
Scientific, PepMap RSLC C18, 2 μM, 100A, 75 μm x 50 cm). Proteome samples (non-enriched)
were eluted over a linear gradient ranging from 0-11% acetonitrile over 70 min, 11-20% acetonitrile
for 80 min, 21-30% acetonitrile for 50 min, 31-48% acetonitrile for 30 min, followed by 76%
acetonitrile for the final 10 min with a flow rate of 250 nl/min.
Survey-full scan MS spectra were acquired in the Orbitrap at a resolution of 120,000 from
m/z 350-2000, automated gain control (AGC) target of $4x10^5$ ions, and maximum injection time of
20 ms. Precursors were filtered based on charge state ($\geq$2) and monoisotopic peak assignment,
and dynamic exclusion was applied for 45s. A decision tree method allowed fragmentation for ion
trap MS2 via electron transfer dissociation (ETD) or higher-energy collision dissociation (HCD),
depending on charge state and m/z. Precursor ions were isolated with the quadrupole set to an
isolation width of 1.6 m/z. MS2 spectra fragmented by ETD and HCD (35% collision energy) were
acquired in the ion trap with an AGC target of 1e4. Maximum injection time for HCD and ETD was
80 ms for proteome samples.
Details of the subsequent data analyses (FASTA file generation, mass spectrometry
searches) are provided in SI Appendix.
425
Reverse phase protein array (RPPA)

For RPPA, snap-frozen cells were lysed in ice-cold protein lysis buffer containing: 50 mM
HEPES, 150 mM NaCl, 1.5 mM $MgCl_2$, 10% (v/v) glycerol, 1% (v/v) TritonX-100, 1 mM EGTA, 100
mM NaF, 10 mM $Na_4P_2O_7$, 2 mM Na_3VO_4 (added fresh), 1X EDTA-free protease inhibitor tablet,
1X PhosStop tablet. Protein concentrations were measured using BioRad's DC protein assay, and
all concentrations were adjusted to 1 mg/ml with lysis buffer and 1X SDS sample buffer (10%
glycerol, 2% SDS, 62.5 mM Tris-HCl pH 6.8) supplemented with 2.5% β-mercaptoethanol.
The protein lysates were processed for slide spotting and antibody incubations as
described previously (44). Briefly, a four-point dilution series was prepared for each sample and
printed in triplicate on single pad Avid nitrocellulose slides (Grace Biolabs) consisting of 8 arrays
with 36x12 spots each. Next, slides were blocked and incubated in primary and secondary
antibodies. The processed arrays were imaged using an Innopsys 710 slide scanner. Non-specific
signals were determined for each slide by omitting primary antibody incubation step. For
normalisation, sample loading on each array was determined by staining with Fast Green dye and
recording the corresponding signal at 800 nm. Details for all primary and secondary RPPA
antibodies are included in **Table S3**.
Details of all subsequent data analyses, including statistical testing, are provided in SI
Appendix.
444
Reverse transcription-quantitative PCR (RT-qPCR)

Cellular RNA was extracted as described above for RNA Sequencing, and 200 ng used for
complementary DNA (cDNA) synthesis with Thermo Fisher's High-Capacity cDNA Reverse
Transcription Kit. Subsequent SYBR Green-based qPCRs were performed on 2.5 ng total cDNA.
TaqMan hPSC Scorecards (384-well) were used according to the manufacturer's instructions with
minor modifications. Further details on protocol modifications and all data analysis steps are
provided in SI Appendix.

Statistical analyses

Bespoke statistical analyses are specified in the relevant sections above and in SI
Appendix.

Data and materials availability

Acknowledgements

We thank Dominique McCormick and Ineke Luijten for help with RNA extraction and cDNA synthesis, Cornelia Gewert for technical support, and Marcella Ma, Brian Lam and Michelle Dietzen for technical support with RNA sequencing and genomics analyses, respectively. We thank Evelyn K. Lau for help with GEO upload of the MEF RNAseq data. We are grateful to Prof Siddhartan Chandran and his group for iPSC culturing facilities and to Pamela Brown (SURF Biomolecular Core, University of Edinburgh) for access to qPCR facilities.

Funding:

R.R.M. and R.K.S. are supported by the Wellcome Trust (105371/Z/14/Z, 210752/Z/18/Z) and United Kingdom (UK) NIHR Cambridge Biomedical Research Centre, and R.R.M. by a Boak Student Award from Clare Hall. Work in the laboratory of B.V. is supported by Cancer Research UK (C23338/ A25722) and the UK NIHR University College London Hospitals Biomedical Research Centre. Metabolic Research Laboratories Core facilities are supported by the Medical Research Council Metabolic Diseases Unit (MC_UU_12012/5) and a Wellcome Major Award (208363/Z/17/Z). R.L. is funded by a Lundbeck Foundation Fellowship. K.G.M., N.O.C. and the University of Edinburgh RPPA facility is supported by a Cancer Research UK Centre award. O.M.R. and C.C. are supported by Cancer Research UK.

Competing interests

R.K.S. is a consultant for HotSpot Therapeutics (Boston, MA, USA). B.V. is a consultant for Karus Therapeutics (Oxford, UK), iOnctura (Geneva, Switzerland) and Venthera (Palo Alto, CA, USA) and has received speaker fees from Gilead Sciences (Foster City, US). N.O.C. is a director of Ther-IP Ltd (Edinburgh, UK) and founder, shareholder and advisor for PhenoTherapeutics Ltd (Edinburgh, UK) and a member of the advisory board and shareholder of Amplia Therapeutics Ltd (Melbourne, Australia).

References

1. D. A. Fruman, *et al.*, The PI3K Pathway in Human Disease. *Cell* **170**, 605–635 (2017).
2. B. Bilanges, Y. Posor, B. Vanhaesebroeck, PI3K isoforms in cell signalling and vesicle trafficking. *Nat. Rev. Mol. Cell Biol.* **20**, 515–534 (2019).
3. M. T. Chang, *et al.*, Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity. *Nat. Biotechnol.* **34**, 155–163 (2015).
4. P. J. Campbell, *et al.*, Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2020).
5. F. Sanchez-Vega, *et al.*, Oncogenic Signaling Pathways in The Cancer Genome Atlas. *Cell* **173**, 321-337.e10 (2018).
6. R. R. Madsen, B. Vanhaesebroeck, R. K. Semple, Cancer-Associated PIK3CA Mutations in Overgrowth Disorders. *Trends Mol. Med.* **24**, 856–870 (2018).
7. R. R. Madsen, *et al.*, Oncogenic PIK3CA promotes cellular stemness in an allele dose-dependent manner. *Proc. Natl. Acad. Sci.* **116**, 8380–8389 (2019).
8. M. Li, J. C. I. Belmonte, Ground rules of the pluripotency gene regulatory network. *Nat. Rev. Genet.* **18**, 180–191 (2017).
9. L. Boyer, *et al.*, Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* **122**, 947–956 (2005).
10. Y. H. Loh, *et al.*, The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nat. Genet.* **38**, 431–440 (2006).
11. J. Nichols, *et al.*, Formation of pluripotent stem cells in the mammalian embryo dependes on the POU transcription factor Oct4. *Cell* **95**, 379–391 (1998).
12. H. Niwa, J. Miyazaki, A. G. Smith, Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat. Genet.* **24**, 372–376 (2000).
13. A. Radzisheuskaya, *et al.*, A defined Oct4 level governs cell state transitions of pluripotency entry and differentiation into all embryonic lineages. *Nat. Cell Biol.* **15**, 579–590 (2013).
14. I. Chambers, *et al.*, Nanog safeguards pluripotency and mediates germline development. *Nature* **450**, 1230–1234 (2007).
15. N. Ivanova, *et al.*, Dissecting self-renewal in stem cells with RNA interference. *Nature* **442**, 533–538 (2006).
16. K. Mitsui, *et al.*, The Homeoprotein Nanog Is Required for Maintenance of Pluripotency in Mouse Epiblast and ES Cells. *Cell* **113**, 631–642 (2003).
17. I. Chambers, *et al.*, Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell* **113**, 643–655 (2003).
18. H. Darr, Overexpression of NANOG in human ES cells enables feeder-free growth while inducing primitive ectoderm features. *Development* **133**, 1193–1201 (2006).
19. R. H. Xu, *et al.*, NANOG is a Direct Target of TGFβ/Activin-Mediated SMAD Signaling in Human ESCs. *Cell Stem Cell* **3**, 196–206 (2008).
20. S. Pauklin, L. Vallier, Activin/Nodal signalling in stem cells. *Development* **142**, 607–619 (2015).
21. L. Vallier, D. Reynolds, R. A. Pedersen, Nodal inhibits differentiation of human embryonic stem cells along the neuroectodermal default pathway. *Dev. Biol.* **275**, 403–421 (2004).
22. L. Vallier, M. Alexander, R. A. Pedersen, Activin/Nodal and FGF pathways cooperate to maintain pluripotency of human embryonic stem cells. *J. Cell Sci.* **118**, 4495–4509 (2005).
23. V. Y. Kiselev, *et al.*, Perturbations of PIP3 signalling trigger a global remodelling of mRNA landscape and reveal a transcriptional feedback loop. *Nucleic Acids Res.* **43**, 9663–79 (2015).
24. J. R. Hart, *et al.*, The butterfly effect in cancer: A single base mutation can remodel the cell. *Proc. Natl. Acad. Sci.* **112**, 1131–1136 (2015).
25. V. E. R. Parker, *et al.*, Safety and efficacy of low-dose sirolimus in the PIK3CA-related overgrowth spectrum. *Genet. Med.* **21**, 1189–1198 (2019).
26. L. S. Moniz, *et al.*, Phosphoproteomic comparison of Pik3ca and Pten signalling identifies

the nucleotidase NT5C as a novel AKT substrate. *Sci. Rep.* **7**, 39985 (2017).

27. X. Robin, *et al.*, Probability-based detection of phosphoproteomic uncertainty reveals rare
signaling events driven by oncogenic kinase gene fusion. *bioRxiv*, 621961 (2019).

28. A. J. Valvezan, B. D. Manning, Molecular logic of mTORC1 signalling as a metabolic
rheostat. *Nat. Metab.* (2019) https:/doi.org/10.1038/s42255-019-0038-7.

29. P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).

30. D. Mesnard, M. Guzman-Ayala, D. B. Constam, Nodal specifies embryonic visceral
endoderm and sustains pluripotent cells in the epiblast before overt axial patterning.
Development **133**, 2497–505 (2006).

31. G. Chen, *et al.*, Chemically defined conditions for human iPSC derivation and culture. *Nat.*
Methods **8**, 424–429 (2011).

32. S. Vukicevic, *et al.*, Identification of multiple active growth factors in basement membrane
Matrigel suggests caution in interpretation of cellular activity related to extracellular matrix
components. *Exp. Cell Res.* **202**, 1–8 (1992).

33. C. S. Hill, Spatial and temporal control of NODAL signaling. *Curr. Opin. Cell Biol.* **51**, 50–
57 (2018).

34. G. J. Inman, *et al.*, SB-431542 is a potent and specific inhibitor of transforming growth
factor-beta superfamily type I activin receptor-like kinase (ALK) receptors ALK4, ALK5,
and ALK7. *Mol. Pharmacol.* **62**, 65–74 (2002).

35. L. Vallier, *et al.*, Activin/Nodal signalling maintains pluripotency by controlling Nanog
expression. *Development* **136**, 1339–49 (2009).

36. I. M. Berenjeno, *et al.*, Oncogenic PIK3CA induces centrosome amplification and
tolerance to genome doubling. *Nat. Commun.* **8**, 1773 (2017).

37. D. J. Murphy, *et al.*, Distinct Thresholds Govern Myc's Biological Output In Vivo. *Cancer*
Cell **14**, 447–457 (2008).

38. P. Liu, *et al.*, Oncogenic PIK3CA-driven mammary tumors frequently recur via PI3K
pathway-dependent and PI3K pathway-independent mechanisms. *Nat. Med.* **17**, 1116–
1120 (2011).

39. F. André, *et al.*, Alpelisib for PIK3CA-mutated, hormone receptor-positive advanced breast
cancer. *N. Engl. J. Med.* **380**, 1929–1940 (2019).

40. M. Bar-Eli, Back to the embryonic stage: Nodal as a biomarker for breast cancer
progression. *Breast Cancer Res.* **14**, 105 (2012).

41. N. V. Margaryan, *et al.*, The stem cell phenotype of aggressive breast cancer cells.
Cancers (Basel). **11**, 1–11 (2019).

42. M. Jewer, *et al.*, Translational control of breast cancer plasticity. *Nat. Commun.* **11**, 2498
(2020).

43. Y. Katsuno, *et al.*, Chronic TGF-b exposure drives stabilized EMT, tumor stemness, and
cancer drug resistance with vulnerability to bitopic mTOR inhibition. *Sci. Signal.* **12**,
eaau8544 (2019).

44. K. G. Macleod, B. Serrels, N. O. Carragher, *Proteomics for Drug Discovery*, I. M. Lazar,
M. Kontoyianni, A. C. Lazar, Eds. (Springer New York, 2017).

45. Y. Perez-Riverol, *et al.*, The PRIDE database and related tools and resources in 2019:
Improving support for quantification data. *Nucleic Acids Res.* (2019)
https:/doi.org/10.1093/nar/gky1106.

592
593
594

595
596 **Figures and Tables**
597
598

Figure 1. Transcriptomic and proteomic analyses of human and mouse cell lines with endogenous
expression of oncogenic *PIK3CA*. (A) Multidimensional scaling (MDS) plot of the transcriptomes of wild-type
(WT), *PIK3CA^{WT/H1047R}* (HET) and *PIK3CA^{H1047R/H1047R}* (HOM) human iPSCs. The numbers in brackets indicate

the presence of two closely overlapping samples. **(B)** The number of differentially expressed genes in iPSCs
heterozygous or homozygous for $PIK3CA^{H1047R}$ before and after application of an absolute fold-change cut-off
$\geq$ 1.3 (FDR $\leq$ 0.05, Benjamini-Hochberg). The data are based on four iPSCs cultures from minimum two clones
per genotype. See also Fig. S1. **(C)** Woman with asymmetric overgrowth caused by mosaicism for cells with
heterozygous expression of $PIK3CA^{E418K}$. Skin biopsies obtained from unaffected and affected tissues were
used to obtain otherwise isogenic dermal fibroblasts for subsequent reprogramming into iPSCs. This image
was reproduced from Ref. (25) **(D)** MDS plot of the transcriptomes of wild-type (WT) and $PIK3CA^{WT/E418K}$ iPSCs
(based on 3 independent mutant clones and 3 wild-type cultures from 2 independent clones). **(E)** MDS plot of
the transcriptomes of wild-type (WT) and $PIK3CA^{WT/H1047R}$ (HET) mouse embryonic fibroblasts (MEFs) following
48 h of mutant induction (N = 4 independent clones per genotype). **(F)** Venn diagram showing the number of
differentially expressed proteins in $PIK3CA^{H1047R/H1047R}$ (HOM) and $PIK3CA^{WT/H1047R}$ (HET) iPSCs relative to
wild-type controls, profiled by label-free total proteomics on three clones per genotype. An absolute fold-change
and z-score $\geq$ 1.2 were used to classify proteins as differentially expressed. The number of discordant and
concordant changes in the expression of total proteins detected in both comparisons are indicated. See also
Fig. S2.

Figure 2. Reverse Phase Protein Array (RPPA) of *PIK3CA^WT/H1047R* **(HET) and** *PIK3CA^H1047R/H1047R* **(HOM)**
human iPSCs. (A) Left: Diagram of PI3K pathway-related phosphorylated proteins, with colour code used to
signify differentially expressed targets in *PIK3CA^H1047R* mutant iPSCs *versus* isogenic wild-type controls. Color-
coded targets were significant at FDR $\leq$ 0.05 (Benjamini-Hochberg). Right: Barplots show representative
examples of differentially expressed protein targets, revealing relatively modest quantitative changes.
Phosphorylated proteins were normalised to the corresponding total protein when available. The data are based

on 10 wild-type cultures (3 clones), 5 *PIK3CA^WT/H1047R* cultures (3 clones) and 7 *PIK3CA^H1047R/H1047R* cultures (2
clones) as indicated. See also Fig. S3A. **(B)** Unsupervised hierarchical clustering based on target-wise
correlations of RPPA data from wild-type (WT), *PIK3CA^WT/H1047R* (HET) and *PIK3CA^H1047R/H1047R* (HOM) iPSCs
following short-term growth factor removal (1 h), +/- 100 nM BYL719 (PI3Kα inhibitor) for 24 h. The data are
from two independent experiments, each performed using independent clones. For each row, the colours
correspond to Fast Green-normalised expression values in units of standard deviation (z-score) from the mean
(centred at 0) across all samples (columns). Groups of phosphorylated proteins exhibiting a consistent
expression pattern in BYL719-treated *PIK3CA^H1047R/H1047R* iPSCs are specified. See also Fig. S3B.

Figure 3. Ingenuity® pathway analyses (IPA) predict activation of TGFβ signalling in heterozygous
and homozygous *PIK3CA^H1047R* iPSCs. (A) IPA of upstream regulators using the list of top 2000
upregulated and top 2000 downregulated mRNA transcripts in *PIK3CA^H1047R/H1047R* iPSCs (for RNAseq
details, see Fig. 1B). Red points signify transcripts with absolute predicted activation z-score > 2 and overlap
P-value < 0.001 (Fisher's Exact Test). The red rectangle highlights the most significant upstream regulator,
TGFβ1. **(B)** As in (A) but using the list of differentially-expressed proteins identified by total proteomics and
red-colouring targets with predicted activation z-score > 2 and overlap P-value < 0.05 (Fisher's Exact Test).
(C) As in (A) but using the list of differentially expressed total proteins in *PIK3CA^WT/H1047R* iPSCs and red-
colouring upstream regulators with absolute predicted bias-corrected z score > 2 and overlap P-value < 0.05
(Fisher's Exact Test). Red rectangles highlight the two upstream regulators (TGFβ1 and MAPK1) with
absolute predicted bias-corrected z score > 2 that remained significant (overlap P-value < 0.05) when the
analysis was repeated using the list of shared and concordant differentially expressed genes (N = 180) in
heterozygous and homozygous *PIK3CA^H1047R* iPSCs *vs* wild-type controls.

Figure 4. Weighted gene correlation network analysis (WGCNA) identifies links among pluripotency components, TGFβ and PI3K signalling. (A) Schematic of the WGCNA workflow and subsequent data selection for visualisation. **(B)** Unsigned WGCNA modules identified using the list of transcripts expressed in wild-type, *PIK3CA*^{WT/H1047R} and *PIK3CA*^{H1047R/H1047R} iPSCs (for RNAseq details, see Fig. 1B). **(C)** The two network modules with genes whose module membership correlated strongest with differential expression in homozygous *PIK3CA*^{H1047R} iPSCs. The colour of each module corresponds to its colour in the module

dendrogram in (B). Representative KEGG pathways with significant enrichment in each gene network module
are listed (hypergeometric test with two-sided uncorrected P < 0.05). **(D)** The minimal network connecting
KEGG pluripotency, PI3K/AKT and TGFβ pathway components within the turquoise gene network module. Fill
colour and shape are used to specify direction of differential mRNA expression in $PIK3CA^{H1047R/H1047R}$ iPSCs
and pathway membership, respectively. Fill colour saturation represents gene expression fold-change (FC;
log2) in $PIK3CA^{H1047R/H1047R}$ (HOM) *vs* wild-type (WT) iPSCs. MYC is highlighted as the only network component
intersecting all three KEGG pathways, suggesting it may comprise a key mechanistic link in the observed
phenotype.

Figure 5. TGFβ signalling-dependent regulation of stemness in *PIK3CA^H1047R/H1047R* iPSCs. Gene expression time course of *NODAL*, *NANOG* and *POU5F1* in wild-type (WT) or *PIK3CA^H1047R/H1047R* iPSCs following the indicated treatments for 24 h, 48 h or 72 h. B250: 250 nM BYL719 (PI3Kα-selective inhibitor); E6/FGF2: Essential 6 medium supplemented with 10 ng/ml basic fibroblast growth factor 2 (FGF2). SB431542 is a specific inhibitor of the NODAL type I receptors ALK4/7 and the TGFβ type I receptor ALK5; used at 5 μM. When indicated, cultures were supplemented with 100 ng/ml NODAL. The data are from two independent experiments, with each treatment applied to triplicate cultures of three wild-type and two homozygous iPSC clones. To aid interpretation, gene expression values are normalised to the E6/FGF2 condition within each genotype and time-point. An alternative visualisation that illustrates the differential expression of *NODAL* and *NANOG* between mutant and wild-type cells is shown in Fig. S4A. For analysis of additional lineage markers, see Fig. S4B. For representative micrographs of *PIK3CA^H1047R/H1047R* iPSCs exposed to the different treatments, see Fig. S5. A.U., arbitrary units.