Leveraging host metabolism for bisdemethoxycurcumin production in *Pseudomonas putida*

Matthew R. Incha[1,2,3], Mitchell G. Thompson[1,2,3], Jacquelyn M. Blake-Hedges[1,2,4], Allison N.

Pearson[1,2], Matthias Schmidt[1,2], Adam M. Deutschbauer[3,5], Jay D. Keasling[1,2,6,7,8]

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.

[2]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720, USA.

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720,

USA

[4]Department of Chemistry, University of California, Berkeley, CA 94720, USA

[5]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National

Laboratory, Berkeley, California, USA

[6]Department of Chemical and Biomolecular Engineering, University of California, Berkeley, CA

94720, USA

[7]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark,

Denmark

[8]Center for Synthetic Biochemistry, Institute for Synthetic Biology, Shenzhen Institutes for
Advanced Technologies, Shenzhen, China

ABSTRACT

Pseudomonas putida is a saprophytic bacterium with robust carbon metabolisms and

strong solvent tolerance making it an attractive host for metabolic engineering and

bioremediation. Due to its diverse carbon metabolisms, its genome encodes an array of proteins

and enzymes that can be readily applied to produce valuable products. In this work we sought to

identify design principles and bottlenecks in the production of type III polyketide synthase

(T3PKS)-derived compounds in *P. putida*. T3PKS products are widely used as nutraceuticals and

medicines and often require aromatic starter units, such as coumaroyl-CoA, which is also an

intermediate in the native coumarate catabolic pathway of *P. putida*. Using a randomly barcoded

transposon mutant (RB-TnSeq) library, we assayed gene functions for a large portion of aromatic

catabolism, confirmed known pathways, and proposed new annotations for two aromatic

transporters. The tetrahydroxynapthalene synthase of *Streptomyces coelicolor* (RppA), a

microbial T3PKS, was then used to rapidly assay growth conditions for increased T3PKS

product accumulation. The feruloyl/coumaroyl CoA synthetase (Fcs) of *P. putida* was used to

supply coumaroyl-CoA for the curcuminoid synthase (CUS) of *Oryza sativa*, a plant T3PKS. We

identified that accumulation of coumaroyl-CoA in this pathway results in extended growth lag

times in *P. putida*. Deletion of the second step in coumarate catabolism, the enoyl-CoA

hydratase lyase (Ech), resulted in 'drop-in' production of the type III polyketide

bisdemethoxycurcumin.

1 INTRODUCTION

Secondary metabolites of fungi, plants, and bacteria have been used as medicines and

supplements for millennia [1]. Compounds such as naringenin, raspberry ketone, resveratrol, and

curcumin are widely used as nutraceuticals and are biosynthesized through similar pathways

[2,3]. Commercially, these chemicals are either extracted directly from plants or produced

synthetically, as in the case of raspberry ketone [4]. Renewable microbial production of these

compounds will decrease reliance on agriculture and fossil fuel-derived chemical synthesis. The

biosynthesis of these compounds (naringenin, raspberry ketone, resveratrol, and curcumin) relies

on a class of enzymes called type III polyketide synthases (T3PKSs). T3PKSs carry out iterative

Claisen condensation reactions typically with coenzyme A (CoA)-based starter and extender

units, both of which vary widely among these enzymes [5]. In the case of the

tetrahydroxynapthalene synthase of *Streptomyces coelicolor* (RppA) the starter and extender

units are simply malonyl-CoA, while in many plant T3PKSs the starter unit is a

phenylpropenoyl-CoA thioester, usually derived from ferulate, coumarate, or cinnamate [6,7].

64   Coumarate and ferulate are components of lignin found in lignocellulosic hydrolysate

(LH), which has been proposed for use as a renewable feedstock for biocatalysis [8,9]. However,

the limited capabilities of commonly used model organisms, such as *Escherichia coli*, have

hindered progress toward making LH a viable feedstock [10]. Characteristics such as high

solvent tolerance and diverse carbon metabolisms are lacking in many *E. coli* strains, but are

essential for a microbe to be used in LH valorization. With this in mind, it has become apparent

that progress needs to be made in developing microbes with these robust traits. Among these

emergent organisms is the saprophytic bacterium *Pseudomonas putida. P. putida* has been

studied for its ability to catabolize phenolics and withstand solvents, making it an attractive host

for LH valorization [11–13].

74   In this work we sought to leverage the native catabolism of *P. putida* KT2440 for use in

the biosynthesis of a plant T3PKS product from coumarate. RppA of *S. coelicolor* was first

expressed to determine optimal production conditions for T3PKS product accumulation. Next we

sought to identify alternative catabolic pathways that could result in reduced titers. Using a

randomly barcoded transposon mutant (RB-TnSeq) library of *P. putida*, we assayed for genes

involved in the catabolism of coumarate and seven other related aromatic compounds. The native

feruloyl/coumaroyl-CoA synthetase (Fcs) of *P. putida* was then used to produce the coumaroyl-

CoA starter unit to the curcuminoid synthase (CUS) of *Oryza sativa*. In the process, we found

that accumulation of coumaroyl-CoA was toxic to *P. putida*. Finally, relying on the native

expression levels of *fcs* from the chromosome and plasmid-based expression of *CUS* resulted in

drop-in production of bisdemethoxycurcumin (BDC) from coumarate (Figure 1).

Figure 1: Diagram of the engineered *P. putida* strain used to produce bisdemethoxycurcumin.
Fcs is the feruloyl/coumaroyl-CoA synthetase native to *P. putida* KT2440. Ech is the native
enoyl-CoA hydratase-lyase which carries out the second step in coumarate catabolism and was
knocked out in our production host. CUS is the curcuminoid synthase from *O. sativa*. BDC is the
final curcuminoid product, bisdemethoxycurcumin.
91
2 RESULTS

2.1 RppA as a screen for optimal T3PKS production conditions

The tetrahydroxynaphthalene synthase of *Streptoyces coelicolor* was recently applied as a

biosensor for malonyl-CoA concentrations in several hosts including *P. putida* [14]. We sought

to utilize this system to rapidly assay growth conditions for increased T3PKS product

accumulation. RppA was initially codon optimized for expression in *P. putida* and cloned into

the construct pBADT-*rppA*-OW. We then constructed two more vectors, one with the complete

rppA cloned from the *S. coelicolor* genome (pBADT-*rppA*-NW) and another with a 5' truncation

of 75 base pairs (pBADT-*rppA*-NT) as this had been described previously to increase enzymatic

activity [6].

Figure 2: Flaviolin production as a reference for T3PKS activity. (A) Diagram of the pathway to
the red pigment, flaviolin, from RppA and subsequent spontaneous oxidation. (B) Absorbance
measurements at 340 nm of supernatants from cells expressing RppA variants in LB containing
different concentrations of glucose. Error bars indicate the standard deviation of three trials.
RppA-NW contains natural *S. coelicolor* codons, RppA-NT is the same as RppA-NW but with a
C-terminal truncation of 25 amino acids, and RppA-OW is translated from a transcript that was
codon optimized for *P. putida*.
109
Since the product of RppA, 1,3,6,8-tetrahydroxynaphthalene, spontaneously oxidizes to

the red pigment flaviolin, we used a colorimetric assay to determine how glucose concentrations

affect the production of T3PKS products (Figure 2a). Concentrations of glucose ranging from 0 -

100 mM supplemented into LB medium were tested, and the production of flaviolin was

measured by absorbance at 340 nm, as previously reported [14]. Unexpectedly, the *rppA*

expression vectors using the native *S. coelicolor* codons produced more flaviolin than the *P.*

putida codon optimized variant (Figure 2b). In all three constructs tested, there was an increase

in the accumulation of flaviolin in cultures containing greater than 25 mM glucose (Figure 2b).

Given this result, production experiments for bisdemethoxycurcumin were conducted in media

supplemented with 100 mM glucose.

Coumarate

Ferulate

-3.9 PP_3356 (*fcs*) -3.7

-3.5 PP_3358 (*ech*) -2.7

-4.7 PP_3357 (*vdh*) -3.1

4-Hydroxybenzoate

Vanillate

PP_3621-2 Vanillin
-2.1

Phenylacetate

-4.7 PP_3537 (*pobA*)
-3.4

-5.0
-5.5 PP_3736-7(*vanAB*)
-5.7

PP_3271 (*phaK*) -3.5
PP_3272 (*actP-III*)

Benzoate

PP_3161-3 (*benABC*) -3.3

PP_1376 (*pcaK*) Protocatechuate
-3.5

PP_3279 (*paaK*) -6.0

-5.5 -5.3
PP_4655-6 (*pcaGH*) -4.5 -6.2
-6.1 -5.0

PP_3164 (*benD*) -7.9

PP_3274-8 (*paaECBA*)
-5.3

PP_3713 (*catA-1*) N.D.**
PP_3166 (*catA-2*)

-6.0 -6.1
PP_1379 (*pcaB*) -6.2 -6.5
-6.8 -6.8

PP_3283 (*paaG*) -5.6

PP_3715 (*catB*) -6.3

PP_1381 (*pcaC*) N.D.*

PP_3270 (*paaZ*) -5.5

PP_3714 (*catC*)
-6.2

-6.0 -5.1 -6.0
PP_1380 (*pcaD*) -5.8 -6.0
-5.8 -5.1

PP_3280 (*paaJ*) -5.1

-7.7 -7.0 -6.8
PP_3951-2 (*pcaIJ*) -6.0 -5.7
-7.6 -6.4

PP_3284 (*paaF*) -6.3

PP_1377 (*pcaF-I*)
PP_3280 (*paaJ*)***

PP_3282 (*paaH*)
-6.1

-7.5 -7.1 -6.5
-5.9 -6.6 -5.1***
-6.1 -6.1

Figure 3: Overview of aromatic catabolism in *P. putida* KT2440. The colored and labelled
compounds depicted were fed as sole carbon sources to the barcoded transposon library. Fitness
values for each gene are the average of 2 replicate RB-TnSeq assays, and are colored
corresponding to the carbon source (gray: benzoate, red: coumarate, blue: 4-hydroxybenzoate,
light green: ferulate, dark green: vanillate, purple: vanillin, yellow: protocatechuate, pink:
phenylacetate). For reactions where multiple genes are necessary, i.e. enzyme complexes, the
fitness values for each gene involved in the reaction were averaged. *The t-score for *pcaC* was
insignificant ($|t_{score}| < 4.0$) and was excluded from our analysis. **Fitness values for these genes
were mild (fitness > -2.0) and excluded. ***Fitness value corresponds to *paaJ*.
132
2.2 Functional genomics to validate aromatic catabolisms of *P. putida*

While aromatic catabolism has been extensively studied in *P. putida*, genes implicated in

these pathways are still being discovered [15]. The genes involved in the first steps of coumarate

catabolism reside in an operon with a putative acyl-CoA dehydrogenase (*PP_3354*) and a

putative beta-ketothiolase (*PP_3355*), which have been proposed to be involved in an alternative

catabolic pathway [16]. As functional redundancy in coumarate catabolism could result in loss of

the type III polyketide precursor, coumaroyl-CoA, we sought to identify any pathways that could

potentially impact product titers. To assay for genes involved in coumarate and related

metabolisms, we grew a randomly barcoded transposon mutant (RB-TnSeq) library of *P. putida*

KT2440 in minimal medium with a variety of different aromatic compounds often found in LH

(p-coumarate, ferulate, benzoate, p-hydroxybenzoate, protocatechuate, vanillin, vanillate,

phenylacetate) and glucose as sole carbon sources. The fitness of each gene was calculated by

comparing the abundance of barcodes before versus after growth selection, using barcode

sequencing (BarSeq) [17,18]. Negative values indicate that the gene was important for growth in

that condition.

The results of the RB-TnSeq assay indicate that the primary route for ferulate and

coumarate catabolism is through the feruloyl/coumaroyl-CoA synthetase (Fcs) and the enoyl-

CoA hydratase lyase (Ech) (Figure 3 and Figure S1). The genes in the proposed secondary

pathway of coumarate and ferulate catabolism, *PP_3354* and *PP_3355*, had no significant fitness

phenotype, indicating that these genes are not necessary for coumarate or ferulate catabolism

[19].

Of all the known reactions depicted in the map of aromatic catabolism (Figure 3), we

observed significant negative fitness values for all but three of their corresponding genes: *catA-1*,

catA-2, and *pcaC*. In the case of *catA-1* and *catA-2* we believe the cause for no fitness phenotype

is the functional redundancy of these two genes, i.e. a cell carrying a transposon insertion in one

will still have the other gene present. In the case of *pcaC*, we noticed that there was a strong

phenotype ($-\log_2 < -2.0$), but the significance fell below our cutoff in all conditions tested ($|t_{score}|$

< 4.0). This is likely due to the low frequency of transposon insertions into this gene in the

library (n=4). While our results heavily support the current models of aromatic metabolism in *P.*

putida, our data also indicated that some gene annotations should be revised. *PP_3272* is

currently annotated as encoding an acetate permease but given this data and its homology to

other systems [20], it should be reannotated as the phenylacetate transporter (phaJ). The

PP_1376 gene is annotated as encoding a 4-hydroxybenzoate transporter; however, we only

observed a fitness detriment for this gene with protocatechuate as the sole carbon source.

Because of this, *PP_1376* should be reannotated as a protocatechuate transporter.

2.3 Accumulation of coumaroyl-CoA is toxic to *P. putida*

The first step in the biosynthetic pathway for bisdemethoxycurcumin is the activation of

coumarate with coenzyme A (CoASH) (Figure 1). Because *Pseudomonas putida* KT2440

natively produces coumaroyl-CoA during coumarate catabolism, we knocked out the subsequent

gene in the native catabolic pathway, *ech*, to prevent *P. putida* from consuming this necessary

precursor (Figure 1) [20]. Initial production experiments in Δ*ech* strains overexpressing *fcs* and

CUS from a synthetic operon resulted in an extended lag phase (data not shown). To determine

the cause, we overexpressed *fcs* alone under control of the arabinose-inducible *araBAD* promoter

(P$_{BAD}$) in the presence and absence of coumarate (Figure 4A). Increasing the inducer

concentration resulted in increased lag times only when coumarate was present in the medium

(Figure 4B). This suggested that the coumaroyl-CoA intermediate is toxic to *P. putida*. We

therefore sought to minimize the burden of this intermediate by relying on the native

chromosomal expression of *fcs* in future experiments.

Figure 4: Production of coumaroyl-CoA inhibits growth of *P. putida.* (A) Diagram of the
induction system used to interrogate coumaroyl-CoA toxicity. Fcs was expressed alone from the
P$_{BAD}$ promoter and coumarate was added to the media. (B) 24 hour growth curves of KT2440
Δ*ech* carrying pBADT-*fcs* in LB induced with varying concentrations of arabinose and no added
coumarate (left), or 5 mM coumarate (right). Error bars are +/- the standard deviation of three
replicates.
187
2.4 Production of bisdemethoxycurcumin

In order to produce bisdemethoxycurcumin in *P. putida*, we employed the biochemical

pathway outlined in Figure 1. Exogenously added coumarate is activated by the native

feruloyl/coumaroyl-CoA synthetase of *P. putida* (Fcs), then two resultant coumaroyl-CoA

molecules are condensed with malonyl-CoA by the curcuminoid synthase of *O. sativa* (CUS) to

yield bisdemethoxycurcumin [21]. The CoA synthetase, *fcs*, was expressed from its native

chromosomal locus, while *CUS* was expressed from the pBADT plasmid and induced with L-

arabinose [22].

Our initial production strategy was to induce *CUS* until the cultures reached stationary

phase. Then the cultures were pelleted and resuspended in fresh LB medium supplemented with

5 mM coumarate. These samples were then incubated for another 72 hours. A similar approach

had been used successfully in *E. coli* [23]; however, our titers were less than 0.5 mg/L (Figure

S2). Given that bisdemethoxycurcumin is insoluble in water [24], a 10 % v/v oleyl alcohol

overlay was used to extract the product as the fermentation progressed. Production levels were

low (approximately 0.1 mg/L) in *P. putida* Δ*ech* when the medium was supplemented with 10

mM coumarate, likely due to the toxicity of coumaroyl-CoA (Figure 5). We therefore lowered

the concentration of externally-supplied coumarate in an attempt to decrease precursor toxicity.

Supplementation with 5 mM coumarate resulted in a ~5-fold increase in bisdemethoxycurcumin

titers in the *P. putida* Δ*ech* strain relative to wild-type (Figure 4A).

Figure 5: Bisdemethoxycurcumin titers in *P. putida* KT2440 Δ*ech* and in wildtype KT2440
harboring pBADT-CUS. *CUS* was induced with 2 % w/v L-arabinose. Error bars are +/- the
standard deviation of three biological replicates.
210

3. DISCUSSION

Pseudomonas putida is among the most well studied saprophytic bacteria. Its diverse

metabolisms enable it to catabolize a wide variety of complex carbon sources, including

lignocellulosic hydrolysate [25]. The robust catabolic pathways of *P. putida*, while useful for

producing valuable molecules from diverse carbon sources, can also serve as an obstacle to

achieving high product titers as it can often metabolize the desired products [26]. Given recent

advances in gene editing techniques [27–31] and our ability to rapidly assay for gene function

with transposon site sequencing [17,18,32], engineering non-model hosts like *P. putida* for

industrial applications has become less challenging.

Using RB-TnSeq mutant fitness assays we were able to rapidly confirm entire pathways

of aromatic catabolism (Figure 3 and S1). The genes downstream of *fcs*, previously described as

a possible alternative route for coumarate/ferulate catabolism [16,20], showed no significant

fitness detriment on any of the carbon sources tested. These genes may be structural remnants of

a β-oxidation pathway that eventually evolved into the coumarate/ferulate pathway requiring *fcs*,

ech, and *vdh* [33]. We then revised the annotations of two genes required for transport of the

aromatic compounds, phenylacetate and protocatechuate, an important plant hormone and lignin

metabolite respectively [34]. There is a large amount of information in these data about

regulatory and structural genetic elements that could be useful to engineers and biologists.

Heterologous expression of bacterial T3PKSs, including the tetrahydroxynapthalene

synthase of *S. coelicolor* (RppA), has previously been demonstrated in *P. putida* KT2440

[14,35,36]. Using variants of *rppA*, we were able to rapidly screen for optimal T3 polyketide

production conditions. Expressing the codon optimized variant we created in this study, *rppA*-

OW, resulted in less flaviolin production than the native codon variants *rppA*-NW and *rppA*-NT.

It is possible that there are some factors affecting heterologous protein expression that are not

sufficiently accounted for in current codon optimization algorithms [37]. However, we

demonstrated in all our constructs that increasing the glucose concentration had a considerable

effect on the production of flaviolin (Figure 2). These T3PKS "sensors" have broad utility in

both rapidly assaying culture conditions, as described here, and as high-throughput screens of

genetic libraries for increased malonyl-CoA accumulation [14]. Future work could employ these

sensors to screen for increased intracellular malonyl-CoA concentrations from a complex

substrate like LH.

To provide the coumaroyl-CoA substrate for the bisdemethoxycurucumin T3PKS, CUS,

we sought to use the native CoA synthetase (Fcs) of *P. putida*. Plasmid-based induction of *fcs*

expression in the presence of coumarate, however, resulted in an increase in lag times due to the

build up of the toxic coumaroyl-CoA intermediate. This toxicity has been observed in

Acinetobacter baylyi following disruptions in the gene encoding its enoyl-CoA hydratase lyase

and in *E. coli* expressing the *A. baylyi fcs* homolog in the presence of coumarate [38]. This

toxicity could also have been limiting other systems using bacterial coumaroyl-CoA synthetases,

but the defective growth phenotype may not have been observed due to differences in

experimental design [39,40]. The exact cause for coumaroyl-CoA toxicity is unclear and will be

the subject of future investigations, as it is likely to interfere with other biosynthetic pathways

requiring this intermediate.

To engineer *P. putida* for bisdemethoxycurcumin production, we deleted the native

enoyl-CoA hydratase lyase (*ech*) responsible for the conversion of coumaroyl-CoA to acetyl-

CoA and p-hydroxybenzaldehyde. In order to relieve coumaroyl-CoA toxicity, we relied on the

native genomic copy of *fcs* instead of a plasmid-based system. We demonstrated that native

expression of *fcs* generates a sufficient coumaroyl-CoA pool for the curcuminoid synthase

(CUS). Extraction of the product during growth using an oleyl alcohol overlay also significantly

enhanced titers (Figure S2 and 5). In the final *P. putida* production strain, we achieved 'drop-in'

production of bisdemethoxycurcumin at titers of 2.15 mg/L (Figure 5).

This work is a significant first step towards the production of plant T3PKS-derived

compounds in *P. putida*, but several issues remain to be addressed. The use of an overlay and the

exogenous addition of coumarate are not feasible for an industrial process. Recent *in vitro* and *in*

vivo experiments suggest that malonyl-CoA is a limiting factor in plant T3PKS pathways

[41,42]. Titers could be improved by modulating this substrate pool. This approach has been

successfully used to optimize titers of the T3PKS product naringenin, resulting in titers of 191.9

mg/L [35]. Future work could employ a similar strategy in *P. putida* to achieve higher titer

production of plant T3PKS products. As we continue to understand more about its metabolism

and engineer it more effectively, *P. putida* will become a more attractive host for renewable

chemical biosynthesis.

4. METHODS

<u>4.1 Media, chemicals, and culture conditions</u>

General *E. coli* cultures were grown in Luria-Bertani (LB) Miller medium (BD

Biosciences, USA) at 37 °C, while *P. putida* was grown at 30 °C. MOPS minimal media was

used where indicated and comprised of the following: 32.5 µM $CaCl_2$, 0.29 mM K_2SO_4, 1.32 mM

K_2HPO_4, 8 µM $FeCl_2$, 40 mM MOPS, 4 mM tricine, 0.01 mM $FeSO_4$, 9.52 mM NH_4Cl, 0.52 mM

$MgCl_2$, 50 mM NaCl, 0.03 µM $(NH_4)_6Mo_7O_{24}$, 4 µM H_3BO_3, 0.3 µM $CoCl_2$, 0.1 µM $CuSO_4$,

0.8 µM $MnCl_2$, and 0.1 µM $ZnSO_4$ [43]. Cultures were supplemented with kanamycin (50 mg/L,

Sigma Aldrich, USA) when indicated. Technical grade oleyl alcohol was acquired from Alfa

Aesar (Alfa Aesar, Thermo Fisher Scientific). All other compounds were purchased through

Sigma Aldrich (Sigma Aldrich, USA). Construction of *P. putida* deletion mutants was performed

as described previously [44].

4.2 Strains and plasmids

Bacterial strains and plasmids used in this work are listed in Table 1. All strains and

plasmids created in this work are available through the public instance of the JBEI registry

https://public-registry.jbei.org. All plasmids were designed using Device Editor and Vector

Editor software, while all primers used for the construction of plasmids were designed using j5

software [45–47]. Plasmids were assembled via Gibson Assembly using standard protocols [48],

or Golden Gate Assembly using standard protocols [49]. Plasmids were routinely isolated using

the Qiaprep Spin Miniprep kit (Qiagen, USA), and all primers were purchased from Integrated

DNA Technologies (IDT, Coralville, IA).

Table 1: Strains and plasmids used in this study

Strain	JBEI Part ID	Reference
E. coli DH10B		[50]
E. coli BL21(DE3)		Novagen
P. putida KT2440		ATCC 47054
P. putida KT2440 Δ*ech*		Park et al (Unpublished)
Plasmids		

pET28		Novagen
pBADT		[22]
pBADT-*fcs-CUS*	TBD	This work
pBADT-*CUS*	TBD	This work
pBADT-*fcs*	TBD	This work
pBADT-*rppA*-NW	TBD	This work
pBADT-*rppA*-NT	TBD	This work
pBADT-*rppA*-OW	TBD	This work
pMQ30		[51]
pMQ30-*PP_3358*		Park et al. (unpublished)

4.3 Plate based growth assays

Growth studies of bacterial strains were conducted using microplate reader kinetic assays.

Overnight cultures were inoculated into 10 mL of LB medium from single colonies, and grown

at 30 °C. These cultures were then diluted 1:100 into 500 μL of LB medium with appropriate

concentrations of arabinose and p-coumarate in 48-well plates (Falcon, 353072). Plates were

sealed with a gas-permeable microplate adhesive film (VWR, USA), and then optical density and

fluorescence were monitored for 48 hours in an Biotek Synergy 4 plate reader (BioTek, USA) at

30 °C with fast continuous shaking. Optical density was measured at 600 nm.

4.4 HPLC detection of bisdemethoxycurcumin

HPLC analysis was performed on an Agilent Technologies 1200 series liquid

chromatography instrument coupled to a Diode Array Detector (Agilent Technologies, USA).

Compounds were separated at a constant flow rate of 0.4 mL/min over a Kinetex C18 column

(2.6 μm diameter, 100 Å particle size, dimensions 100 x 3.00 mm, Phenomenex, USA) held at 50

°C. The mobile phase consisted of H_2O + 0.1% trifluoroacetic acid (A) and acetonitrile + 0.1%

trifluoroacetic acid (B). Separation was performed using the following gradient method: 0-3

minutes 95% A, 3-15 minutes 95-5% A, 15-17 minutes 5% A, 17-17.5 minutes 5-95% A, 17.5-

20 minutes 95% A. The presence of bisdemethoxycurcumin was monitored and quantified at 440

nm.

4.5 RB-TnSeq experiments and analysis

BarSeq-based experiments utilized the *P. putida* RB-TnSeq library, JBEI-1, which has

been described previously [44]. An aliquot of JBEI-1 was thawed on ice, diluted into 25 mL of

LB medium supplemented with kanamycin and grown to an OD_{600} of 0.5 at 30 °C. Three 1 mL

aliquots of the library were pelleted and stored at -80 °C to later serve as the t_0 of gene

abundance. Libraries were then washed in MOPS minimal medium, and diluted 1:50 in MOPS

minimal medium with 10 mM p-coumarate, ferulate, benzoate, p-hydroxybenzoate,

protocatechuate, vanillin, vanillate, phenylacetate, or D-glucose. Cells were grown in 600 μL of

medium in 96-well deep well plates (VWR). Plates were sealed with a gas-permeable microplate

adhesive film (VWR, USA), and then grown at 30 °C in an INFORS HT Multitron (Infors USA

Inc.), with shaking at 700 rpm. Two 600-μL samples were combined, pelleted, and stored at -

80 °C until analysis by BarSeq , which was performed as previously described [17,18]. All

fitness data is publically available at http://fit.genomics.lbl.gov.

4.5 Curcuminoid production

For production of bisdemthoxycurcumin without an overlay, an overnight culture of *P.*

putida KT2440 Δ*ech* + pBADT-CUS was diluted 1:100 into 5 mL of LB supplemented with 50

mg/L kanamycin, 1 % w/v L-arabinose, and 100 mM glucose. The culture was grown to

stationary phase over 12 hours then pelleted in a centrifuge at 5000 xg for 5 minutes. The cell

pellets were resuspended in 2 mL of fresh LB with 50 mg/L kanamycin, 0.5 % L-arabinose, 100

mM glucose, and 5 mM coumarate. The culture was allowed to proceed for 72 hours.

Overnights of *P. putida* harboring pBADT-CUS were diluted 1:100 into 25 mL of fresh

LB supplemented with 50 mg/L kanamycin, and 100 mM glucose. Arabinose and coumarate

were added at the beginning of the fermentation at concentrations indicated. A 2.5 mL overlay of

oleyl alcohol was added to extract the bisdemethoxycurcumin during growth. The fermentation

was allowed to proceed for 72 hours.

4.6 Curcuminoid extraction

For cultures lacking an overlay, 0.5 mL of culture was acidified to pH 3 with 3 N HCl.

Bisdemethoxycurcumin was then extracted with an equal volume of ethyl acetate. 250 μL of the

ethyl acetate layer was removed and the solvent was allowed to evaporate overnight. The dried

samples were then resuspended in 50 μL of acetonitrile for analysis with HPLC-DAD.

For cultures with an overlay, the cultures were acidified to pH 3 with 3 N HCl. Acidified

cultures were then pelleted in a centrifuge and the oleyl alcohol overlays were extracted. To

quantify bisdemethoxycurcumin, 100 μL of the extracted overlays were added to a black, clear

bottom 96-well plate and absorbance was measured at 425 nm in a Biotek Synergy 4 plate reader

(BioTek, USA). A standard curve was made with bisdemethoxycurcumin standards dissolved in

oleyl alcohol (Figure S3)

4.7 Flaviolin production

Colonies of *P. putida* KT2440 strains carrying pBADT-*rppA*-NW, pBADT-*rppA*-NT,

and pBADT-*rppA*-OW were used to inoculate LB medium with 50 mg/L kanamycin and

cultured overnight. The overnight culture was then diluted 1/100 into fresh LB with 0.2 % w/v L-

arabinose, 50 mg/L kanamycin, and 100, 50, 25, 12.5, 6.25, 3.125, 1.5625 or 0 mM glucose.

Cultures were conducted in 24-well deep-well plates, and allowed to proceed for 48 hours.

Cultures were pelleted in a centrifuge at 5000 xg for 5 minutes. Supernatants were removed,

aliquoted into a 96-well black clear bottom plate, and the absorbance was measured at 340 nm in

a Biotek Synergy 4 plate reader (BioTek, USA). For cultures expressing *rppA*-NW and *rppA*-NT,

the supernatants were too opaque with red product and needed to be diluted 1:2 in fresh LB for

accurate absorbance measurements.

Acknowledgements

We would like to thank Professor Mattheos Koffas for providing us with the plasmid

pEMT6-4CL-CUS. We thank Jesus Barajas for his careful reading of this manuscript and helpful

suggestions during preparation. We also thank Morgan Price for assistance in analyzing RB-

TnSeq data.

This work was part of the DOE Joint BioEnergy Institute (https://www.jbei.org)

supported by the U. S. Department of Energy, Office of Science, Office of Biological and

Environmental Research, supported by the U.S. Department of Energy, Energy Efficiency and

Renewable Energy, Bioenergy Technologies Office, through contract DE-AC02-05CH11231

between Lawrence Berkeley National Laboratory and the U.S. Department of Energy. The views

and opinions of the authors expressed herein do not necessarily state or reflect those of the

United States Government or any agency thereof. Neither the United States Government nor any

agency thereof, nor any of their employees, makes any warranty, expressed or implied, or

assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any

Contributions

Conceptualization, M.R.I., M.G.T., J.M.B; Methodology, M.R.I., M.G.T., J.M.B.; Investigation,

M.R.I., M.G.T., J.M.B., M.S., A.N.P.; Writing – Original Draft, M.R.I.; Writing – Review and

Editing, All authors.; Resources and supervision, A.M.D., J.D.K.

Competing Interests

J.D.K. has financial interests in Amyris, Lygos, Demetrix, Napigen and Maple Bio

Supplementary Figures

Figure S1: Heatmap of RB-TnSeq mutant fitness data. Genes shown have fitness defects ($-\log_2$

< -2) and with statistical significance ($|t_{score}| > 4$) in at least one tested condition which show no

significant fitness defects when grown on glucose.

Fitness (-log$_2$)

Scale: −6 to 6

Gene	Vanillate_1	Vanillate_2	Vanillin_1	Vanillin_2	Ferulate_1	Ferulate_2	Coumarate_1	Coumarate_2	p-Hydroxybenzoate_1	p-Hydroxybenzoate_2	Benzoate_1	Benzoate_2	Protocatechuate_1	Protocatechuate_2	Phenylacetate_1	Phenylacetate_2	Glucose_1	Glucose_2	Annotation
PP_0120	-1.4	-1.7	-2.2	-1.7	-1.8	-1.6	-0.68	-0.66	-0.33	-0.16	-0.25	-0.5	-0.69	-0.57	-0.44	-0.41	-0.45	-0.16	putative zinc ABC transporter
PP_0368	-1.8	-1.9	-3.4	-3.5	-1.1	-1	-1.8	-1.7	-1	-0.84	-1.4	-1.3	-1.7	-1.7	-1.6	-1.7	-0.31	0.06	putative Acyl-CoA dehydrogenase
PP_1292	-0.87	-0.8	-3.2	-3.8	-1.1	-1.1	-0.69	-0.65	-0.11	-0.64	-0.22	-0.33	-0.65	-0.44	-0.46	-0.88	-0.26	-0.35	Cyclic pyranopterin monophosphate synthase accessory protein
PP_1294	-0.6	-0.39	-2.4	-3.4	-0.78	-0.95	-0.47	-0.47	-0.53	-0.31	-0.72	-0.32	-0.12	-0.88	-0.59	-0.53	-0.23	-0.11	molybdopterin synthase large subunit
PP_1375	-5.1	-7	-4.7	-6.2	-5.4	-5.1	-5.4	-5.2	-5.8	-4.8	-6.8	-6.7	-6.5	-6	-0.09	-0.12	-0.02	0.1	transcription regulatory protein (pca regulon)
PP_1376	-0.51	-0.58	-0.37	-0.41	-0.07	-0.29	-0.75	-0.48	-0.43	-0.31	0.12	-0.26	-3.5	-3.5	-0.06	-0.19	-0.23	0.02	4-hydroxybenzoate transporter
PP_1377	-5.7	-6.6	-5.6	-7.5	-6.5	-7.7	-7.3	-7.6	-5.3	-6.5	-6.3	-6.6	-5.7	-6.4	-0.19	-0.31	-0.15	0.23	beta-ketoadipyl-CoA thiolase
PP_1379	-6	-7.6	-5.4	-7.5	-6	-6.1	-6.9	-6.6	-6.8	-5.7	-0.91	-0.85	-7	-6.7	-0.19	-0.31	-0.15	0.03	3-carboxy-cis,cis-muconate cycloisomerase
PP_1380	-4.9	-5.3	-4.9	-7.1	-5.7	-4.5	-5.9	-6	-6.2	-5.3	-6.4	-5.5	-5.5	-5.9	-0.04	-0.22	-0.23	0.18	3-oxoadipate enol-lactonase 2
PP_1615	-2.2	-2	-2.1	-2.2	-2.8	-2.3	-0.28	0.01	0.14	-0.23	-0.03	-0.13	0.06	0.04	-0.16	-0.15	-0.1	-0.18	Transcriptional regulator, LysR family
PP_1616	-3.2	-4.2	-4.9	-4.6	-4.2	-4.5	-0.37	-0.65	-0.48	-0.46	-0.88	-0.57	-0.52	-0.51	-0.49	-0.6	-0.28	-0.07	formaldehyde dehydrogenase, glutathione-dependent
PP_2123	-0.58	-0.76	-2.9	-3.8	-1.1	-0.99	-0.85	-0.51	-0.18	-0.31	-0.47	-0.61	-0.55	-0.71	-0.72	-0.78	-0.18	-0.38	molybdenum::molybdopterin ligase
PP_2136	-1.4	-1.2	-1.6	-2.2	-1.2	-1.4	-2	-1.5	-0.63	-0.82	-0.78	-1.6	-0.47	-0.47	-2.5	-1.1	-0.08	-0.48	enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase
PP_2354	-1.3	-1.4	-1.9	-1.7	-1.5	-1.5	-1.5	-1.4	-1.4	-1.5	-1.6	-1.6	-1.6	-1.2	-1.5	-2	-0.05	-0.09	Sensory box histidine kinase/response regulator
PP_2356	-1.5	-1.7	-2.2	-2.1	-2.1	-2.2	-1.9	-1.9	-1.5	-1.7	-2	-1.9	-1.5	-2	-2.6	-2.9	-0.31	-0.09	putative Phytochrome family protein
PP_3159	-0.11	-0.04	-0.06	-0.25	-0.09	0.02	0.07	-0.01	0.06	-0.15	-3.2	-3	-0.01	0.08	0.06	-0.09	-0.04	-0.1	BenABC operon transcriptional activator
PP_3161	-0.24	0	-0.16	-0.16	-0	-0.01	0.02	-0.06	-0.11		-2.9	-2.9	0.05	0.02	0.09	0.02	-0.11	0.1	benzoate 1,2-dioxygenase subunit alpha
PP_3162	0.21	-0.11	0.02	-0.02	-0.09	-0.16	-0.28	-0.13	0.03	-0.29	-2.7	-4	0.07	0.21	-0.21	0.16	-0.01	-0.08	benzoate 1,2-dioxygenase subunit beta
PP_3163	0.09	0.02	-0.07	-0.16	-0.32	-0.11	0.02	-0.21	0.07	-0.02	-3.7	-3.5	-0.45	-0.32	0.02	0.16	-0.16	-0.18	benzoate 1,2-dioxygenase electron transfer component
PP_3164	-0.49	-0.48	-0.29	0	-0.4	-0.28	-0.49	0.34	0.21	-0.1	-7.9	-7.8	0.05	0.08	-0.41	0.17	0.02	0.17	1,6-dihydroxycyclohexa-2,4-diene-1-carboxylate dehydrogenase
PP_3270	-0.04	0.02	0.08	-0.07	0.1	0.18	0.03	-0.04	-0.06	-0.04	0.09	-0.15	-0.02	0	-4.8	-6.2	0	0.09	oxepin-CoA hydrolase/3-oxo-5,6-dehydrosuberyl-CoA semialdehyde dehydrogenase
PP_3271	-0.14	0.06	0.01	0.07	-0.11	0.02	0	-0.01	-0.14	-0.1	-0.13	-0.22	-0.07	0.07	-1.8	-2.1	-0.13	0.05	phenylacetic acid-specific porin
PP_3272	-0	-0.07	0.06	0.07	-0.07	-0.02	0.12	0.11	0.1	0.01	0.04	0.03	-0.01	-0.01	-3.5	-6.5	0.04	-0	acetate permease
PP_3273	-0.15	-0.12	0.14	-0.17	0.14	0.01	0.31	0.11	0.07	0.29	-0.03	0.18	-0.01	0.08	-3.5	-3.8	0.22	0.22	conserved protein of unknown function
PP_3274	-0.13	-0.15	-0.14	-0.2	-0.23	-0.07	0.17	0.03	-0.05	0.01	-0.11	-0.26	-0.05	-0.05	-4.2	-6	-0.16	-0.12	ring 1,2-phenylacetyl-CoA epoxidase, reductase subunit
PP_3276	0.05	0.17	-0.09	0.18	0.18	0.1	-0.19	-0.45	-0.34	-0.01	-0.23	-0.15	0.04	0.09	-4.5	-7.3	0.1	-0.03	ring 1,2-phenylacetyl-CoA epoxidase beta subunit
PP_3277	0.17	0.11	-0.03	0.02	0.04	-0.34	-0.25	-0.13	0.24	-0.26	-0.32	-0.32	-0.22	-0.12	-5.2	-6.1	-0.28	-0.36	ring 1,2-phenylacetyl-CoA epoxidase regulatory subunit
PP_3278	0.04	-0.1	-0.07	0.27	0.06	0.2	-0	0.01	-0.01	-0.3	-0.15	-0.05	0.16	-0.1	-3.5	-5.8	0.02	0.03	ring 1,2-phenylacetyl-CoA epoxidase alpha subunit
PP_3279	-0.05	0.13	0.17	0.05	0.29	0.03	0.05	0.04	-0.08	-0.02	-0.03	-0.05	-0.24	0.06	-4.8	-7.2	0.08	0.02	phenylacetate-CoA ligase
PP_3280	0.14	0.23	0.15	0.48	0.13	-0.36	0.41	0.12	-0.05	-0.29	0.35	-0.32	0.18	-0.53	-3.7	-6.4	-0.22	-0.19	3-oxoadipyl-CoA/3-oxo-5,6-dehydrosuberyl-CoA thiolase
PP_3281	-0.16	0.31	-0.18	-0.19	0.14	-0.01	0.09	-0.03	-0.25	-0.08	-0	0.15	0.17	-0.11	-2.5	-3.2	-0.04	0.08	hydroxyphenylacetyl-CoA thioesterase
PP_3282	-0.05	-0.01	-0.04	-0.09	-0.03	-0.04	0.09	0.09	-0.06	-0.05	0.14	-0.15	-0.04	-0.14	-4.5	-7.6	0.07	0.11	3-hydroxyadipyl-CoA dehydrogenase
PP_3283	-0.03	-0.07	-0.21	0.24	0.15	-0.19	-0.17	0.08	-0.02	0.09	0.16	-0.11	-0.25	0.22	-4.8	-6.4	-0.09	-0.27	1,2-epoxyphenylacetyl-CoA isomerase
PP_3284	-0.02	0.34	0.26	0.1	0.22	0.27	-0.32	-0.04	-0.18	0.03	-0.01	-0.24	0.31	-0.01	-4.7	-7.8	-0.14	-0.15	enoyl-CoA hydratase-isomerase
PP_3285	0.19	0.09	0.1	0.09	-0.43	0.21	0.01	0.42	-0.04	0.07	-0.11	0.41	-0.06	-0.28	-4.6	-7.4	0.18	-0.03	detoxifying thioesterase of the phenylacetate degradation pathway
PP_3356	0.02	0.15	-0.05	-0.11	-3.3	-4	-4.2	-3.6	-0.05	-0.06	-0	-0.05	0.1	-0.03	0.08	-0.03	0.15	-0.06	feruloyl-CoA-synthetase
PP_3357	0.12	-0.1	0.18	0.06	-3.6	-2.5	-5.2	-4.2	0.24	-0.1	-0.04	-0.06	-0.03	0.23	0.09	0.17	0.14	0.13	vanillin dehydrogenase
PP_3358	0.09	0.12	0.05	0.2	-2.9	-2.5	-3.1	-3.8	0.17	0.07	-0	0.23	0.03	-0.02	-0.01	0.08	0.04	0.02	hydroxycinnamoyl-CoA hydratase-lyase
PP_3493	-1.4	-1	-1.5	-1.5	0.11	0.14	0.06	-0.05	0.16	-0.23	-1.7	-1.8	0.02	0.04	-2.7	-3.8	0.01	0.07	Transcriptional regulator, LysR family
PP_3494	-1.5	-0.93	-1.4	-1.1	0.01	-0.01	0	-0.15	-0.18	-0.16	-0.77	-0.95	-0.22	0.04	-3.5	-3.2	0.01	-0.19	conserved exported protein of unknown function
PP_3537	0.09	0.21	-0.05	-0.04	0.11	-0.11	-4.6	-4.8	-3.2	-3.6	0.02	-0.02	0.1	-0.02	0.15	-0.03	-0.05	-0.02	p-hydroxybenzoate hydroxylase
PP_3538	0.2	0.14	0.14	0.12	0.04	0.06	-4.6	-4.7	-3	-3.8	-0	0.05	0.02	0.16	0.09	0.06	0.04		transcriptional regulator
PP_3590	-3	-3.2	-1.8	-2	-3.6	-2.9	-2.1	-1.8	-2.7	-3.5	-3.4	-3.3	-1.2	-1.6	-1.6	-2	-0.19	-0.08	D-lysine aminotransferase
PP_3603	-1.4	-1.3	-1.7	-2.2	-0.79	-1.2	-0.47	-0.76	-0.49	-1.2	-0.62	-0.55	-0.79	-0.44	-1.1	-1	-0.4	-0.32	Transcriptional regulator, GntR family
PP_3621	-0.36	-0.23	-1.7	-2.4	-0.15	-0.25	-0.35	-0.15	-0.26	0.12	-0.07	-0.53	-0.12	-0.28	-0.02	0.36	0.18	-0.07	isoquinoline 1-oxidoreductase subunit alpha
PP_3622	-0.11	0.08	-1.9	-2.3	0.09	-0.04	-0.19	-0.24	0.03	0.01	0.08	0.03	-0.15	0.07	0.07	-0.03	0.02		putative Isoquinoline 1-oxidoreductase, beta subunit
PP_3662	-1.5	-1.4	-1.8	-1.6	-1.1	-1.3	-0.5	-0.6	-1.1	-0.89	-1.9	-1.4	-0.72	-0.73	-1.7	-2.7	-0.41	-0.11	conserved protein of unknown function
PP_3714	0.19	0.34	-0.38	0.23	-0.38	-0.08	-0.03	-0.11	-0.09	0	-6.7	-5.6	-0.26	-0.51	-0.31	0.02	-0.47	-0.14	Muconolactone Delta-isomerase
PP_3715	0.18	0.75	-0.08	0.12	-0.18	0.11	-0.24	-0.13	0.06	0.28	-5.7	-6.9	-0.27	-0.28	0.34	-0.01	0.14	0.31	Muconate cycloisomerase 1
PP_3716	-0.15	-0.13	-0.1	-0.05	-0.01	-0.02	-0.09	-0.04	-0.04	-0.04	-5.6	-6	-0.2	-0.07	-0.06	-0.31	0	0.07	HTH-type transcriptional regulator
PP_3736	-5.8	-6.4	-4.3	-7.2	-5.2	-6.2	-0.09	0.01	-0.03	0.08	0.03	-0.32	0.14	0.25	0.12	-0.02	0.08	-0.02	vanillate O-demethylase oxygenase subunit
PP_3737	-4.8	-6	-4.5	-6	-4.4	-4.2	-0.04	0.14	0.04	-0.07	-0.04	-0.04	-0.01	0.18	0.11	-0.05	-0.24	0.03	vanillate O-demethylase oxidoreductase
PP_3828	-0.45	-0.54	-2.5	-3.9	-0.71	-0.35	-0.48	-0.64	-0.38	-0.24	-0.24	-0.33	-0.36	-0.28	-0.28	-0.52	-0.37	-0.07	Molybdate-binding periplasmic protein
PP_3829	-0.5	-0.45	-3.1	-3.3	-0.76	-0.75	-0.47	-0.69	-0.23	-0.19	-0.42	-0.01	-0.59	-0.27	-0.23	-0.43	-0.12		Molybdenum transport system permease protein ModB
PP_3830	-0.09	-0.3	-1.7	-2.2	-0.29	-0.66	-0.32	-0.28	-0.17	-0.2	-0.2	-0.16	-0.29	-0.29	-0.23	-0.51	-0.1	-0	Molybdenum import ATP-binding protein ModC
PP_3951	-5.2	-7.3	-3.7	-6.4	-7.2	-7.1	-7.8	-8.1	-5.5	-5.6	-5.1	-8.1	-8	-7.9	-0.23	0.03	-0.25	0.21	3-oxoadipate CoA-transferase subunit A
PP_3952	-6.4	-6.6	-4.6	-8.2	-6.8	-7	-7.3	-7.6	-6.8	-6	-6	-8	-7.1	-7.4	0.12	-0.38	-0.22	-0.08	3-oxoadipate CoA-transferase subunit A
PP_4008	-0.74	-0.11	-0.78	-1.1	-1.1	-0.99	-0.81	-0.68	-0.43	-1	-0.46	-0.44	-0.48	-0.28	-1.2	-2.2	-0.25	0.1	ATP-dependent serine protease
PP_4058	-1.9	-2.2	-2.3	-1.9	-1.5	-1.6	-0.5	-0.3	-0.7	-0.6	-0.15	-0.06	-0.5	-0.51	-3.1	-3.6	-0.19	-0.03	1,4-alpha-glucan branching enzyme
PP_4098	-1.2	-1.7	-0.98	-0.7	-1.3	-0.43	-3.4	-0.57	-1.3	-1	-1.1	-0.79	-2.5	-0.48	-0.48	-0.97	0.01	-0.29	UvrABC system protein C
PP_4231	0.02	0.08	-1.9	-2.5	-0.01	-0.03	-0.3	0.17	-0.19	0.09	-0.08	-0.08	0.04	-0.13	-0.09	0.1	0.01	-0.12	putative Xanthine dehydrogenase accessory factor
PP_4597	-0.4	-0.46	-1.9	-2.9	-0.16	-0.73	-0.31	-0.03	-0.41	-0.06	0	-0.5	0.07	-0.31	-0.51	-0.21	-0.33	-0.08	Cyclic pyranopterin monophosphate synthase
PP_4622	-0.99	-0.99	-2.1	-1.1	-0.18	-0.17	-0.13	-0.12	-0.23	0.17	-0.06	-0.19	-0.49	-0.17	-0.81	-0.91	-0.04	-0.07	Hmg transcriptional repressor
PP_4655	-4	-5.1	-4.9	-6.5	-4.9	-5	-4.2	-4.9	-5.2	-3.9	0.1	0.19	-6.3	-5.1	0.23	0.02	0.03	0.05	protocatechuate 3,4-dioxygenase alpha chain
PP_4656	-3.9	-6.8	-5.4	-8	-5.1	-6.1	-6.8	-6	-4.5	-4.6	0.11	0.27	-6.5	-6.4	-0.32	-0.14	-0.31	0.04	protocatechuate 3,4-dioxygenase beta chain
PP_5085	-1.3	-1.3	-1.6	-0.85	-0.81	-0.63	-2.3	-2.5	-4.5	-4.4	-0.83	-0.57	-0.75	-0.54	-1.3	-1.3	0.34	0.27	malic enzyme B
PP_5331	0.9	0.62	0.34	0.91	0.47	0.51	-0.44	-0.69	-2	-2	-0.95	-1.6	0.22	0.26	-1.2	-0.52	-0.21	-0.41	Long-chain acyl-CoA thioester hydrolase

395 **Figure S2:** HPLC trace of curcumin extracted from *P. putida* KT2440 Δ*ech* + pBADT-CUS
396 overlaid with a 3.125 mg/L standard
397

398
399 **Figure S3:** Standard curve of bisdemethoxycurcumin. Absorbance measurements at 425 nm
400 from 50 to 0 mg/L of bisdemethoxycurcumin dissolved in oleyl alcohol.

401
402
Bibliography

[1]   S.J. Hewlings, D.S. Kalman, Curcumin: A review of its' effects on human health.,
Foods. 6 (2017). doi:10.3390/foods6100092.
[2]   S. Schanz, G. Schröder, J. Schröder, Stilbene synthase from Scots pine (*Pinus*
***sylvestris*), FEBS Lett. 313 (1992) 71–74. doi:10.1016/0014-5793(92)81187-Q.**
[3]   W. Heller, K. Hahlbrock, Highly purified "flavanone synthase" from parsley
catalyzes the formation of naringenin chalcone, Arch. Biochem. Biophys. 200 (1980)
617–619. doi:10.1016/0003-9861(80)90395-1.
[4]   L.R. Smith, Rheosmin ("Raspberry Ketone") and Zingerone, and Their Preparation

by Crossed Aldol-Catalytic Hydrogenation Sequences, Chem. Educator. 1 (1996) 1–18. doi:10.1007/s00897960034a.

[5]   D. Yu, F. Xu, J. Zeng, J. Zhan, Type III polyketide synthases in natural product biosynthesis., IUBMB Life. 64 (2012) 285–295. doi:10.1002/iub.1005.

[6]   M. Izumikawa, P.R. Shipley, J.N. Hopke, T. O'Hare, L. Xiang, J.P. Noel, et al., Expression and characterization of the type III polyketide synthase 1,3,6,8-tetrahydroxynaphthalene synthase from Streptomyces coelicolor A3(2)., J. Ind. Microbiol. Biotechnol. 30 (2003) 510–515. doi:10.1007/s10295-003-0075-8.

[7]   T. Wakimoto, H. Morita, I. Abe, Engineering of plant type III polyketide synthases., Meth. Enzymol. 515 (2012) 337–358. doi:10.1016/B978-0-12-394290-6.00016-1.

[8]   J.M. Lawther, R. Sun, The fractional characterisation of polysaccharides and lignin components in alkaline treated and atmospheric refined wheat straw, Industrial Crops and Products. 5 (1996) 87–95. doi:10.1016/0926-6690(96)00002-7.

[9]   L.J. Jönsson, B. Alriksson, N.-O. Nilvebrant, Bioconversion of lignocellulose: inhibitors and detoxification., Biotechnol. Biofuels. 6 (2013) 16. doi:10.1186/1754-6834-6-16.

[10]   T.Y. Mills, N.R. Sandoval, R.T. Gill, Cellulosic hydrolysate toxicity and tolerance mechanisms in Escherichia coli., Biotechnol. Biofuels. 2 (2009) 26. doi:10.1186/1754-6834-2-26.

[11]   T. Nakazawa, T. Yokota, Benzoate metabolism in Pseudomonas putida(arvilla) mt-2: demonstration of two benzoate pathways., J. Bacteriol. 115 (1973) 262–267.

[12]   M.T. Guarnieri, M. Ann Franden, C.W. Johnson, G.T. Beckham, Conversion and assimilation of furfural and 5-(hydroxymethyl)furfural by Pseudomonas putida KT2440., Metab. Eng. Commun. 4 (2017) 22–28. doi:10.1016/j.meteno.2017.02.001.

[13]   J.G. Linger, D.R. Vardon, M.T. Guarnieri, E.M. Karp, G.B. Hunsinger, M.A. Franden, et al., Lignin valorization through integrated biological funneling and chemical catalysis., Proc Natl Acad Sci USA. 111 (2014) 12013–12018. doi:10.1073/pnas.1410657111.

[14]   D. Yang, W.J. Kim, S.M. Yoo, J.H. Choi, S.H. Ha, M.H. Lee, et al., Repurposing type III polyketide synthase as a malonyl-CoA biosensor for metabolic engineering in bacteria., Proc Natl Acad Sci USA. 115 (2018) 9835–9844. doi:10.1073/pnas.1808567115.

[15]   M.N. Price, J. Ray, A.T. Iavarone, H.K. Carlson, E.M. Ryan, R.R. Malmstrom, et al., Oxidative pathways of deoxyribose and deoxyribonate catabolism., MSystems. 4 (2019). doi:10.1128/mSystems.00297-18.

[16]   J. Overhage, H. Priefert, J. Rabenhorst, A. Steinbüchel, Biotransformation of eugenol to vanillin by a mutant of Pseudomonas sp. strain HR199 constructed by disruption of the vanillin dehydrogenase (vdh) gene, Appl. Microbiol. Biotechnol. 52 (1999) 820–828. doi:10.1007/s002530051598.

[17]   K.M. Wetmore, M.N. Price, R.J. Waters, J.S. Lamson, J. He, C.A. Hoover, et al.,

Rapid quantification of mutant fitness in diverse bacteria by sequencing randomly
bar-coded transposons., MBio. 6 (2015) e00306-15. doi:10.1128/mBio.00306-15.

[18]   **J.M. Rand, T. Pisithkul, R.L. Clark, J.M. Thiede, C.R. Mehrer, D.E. Agnew, et al.,**
A metabolic pathway for catabolizing levulinic acid in bacteria., Nat. Microbiol. 2
(2017) 1624–1634. doi:10.1038/s41564-017-0028-z.

[19]   **R. Plaggenborg, J. Overhage, A. Steinbüchel, H. Priefert, Functional analyses of**
genes involved in the metabolism of ferulic acid in Pseudomonas putida KT2440.,
Appl. Microbiol. Biotechnol. 61 (2003) 528–535. doi:10.1007/s00253-003-1260-4.

[20]   **J.I. Jiménez, B. Miñambres, J.L. García, E. Díaz, Genomic analysis of the aromatic**
catabolic pathways from Pseudomonas putida KT2440., Environ. Microbiol. 4 (2002)
824–841. doi:10.1046/j.1462-2920.2002.00370.x.

[21]   **Y. Katsuyama, M. Matsuzawa, N. Funa, S. Horinouchi, In vitro synthesis of**
curcuminoids by type III polyketide synthase from Oryza sativa., J. Biol. Chem. 282
(2007) 37702–37709. doi:10.1074/jbc.M707569200.

[22]   **C. Bi, P. Su, J. Müller, Y.-C. Yeh, S.R. Chhabra, H.R. Beller, et al., Development of**
a broad-host synthetic biology toolbox for Ralstonia eutropha and its application to
engineering hydrocarbon biofuel production., Microb. Cell Fact. 12 (2013) 107.
doi:10.1186/1475-2859-12-107.

[23]   **Y. Katsuyama, N. Funa, I. Miyahisa, S. Horinouchi, Synthesis of unnatural**
flavonoids and stilbenes by exploiting the plant biosynthetic pathway in Escherichia
coli., Chem. Biol. 14 (2007) 613–621. doi:10.1016/j.chembiol.2007.05.004.

[24]   **I. Javeri, N. Chand, Curcumin, in: Nutraceuticals, Elsevier, 2016: pp. 435–445.**
doi:10.1016/B978-0-12-802147-7.00031-0.

[25]   **Y. Wang, F. Horlamus, M. Henkel, F. Kovacic, S. Schläfle, R. Hausmann, et al.,**
Growth of engineered*Pseudomonas putida* KT2440 on glucose, xylose and arabinose:
hemicellulose hydrolysates and their major sugars as sustainable carbon sources,
Glob. Change Biol. Bioenergy. (2019). doi:10.1111/gcbb.12590.

[26]   **M.G. Thompson, L.E. Valencia, J.M. Blake-Hedges, P. Cruz-Morales, A.E.**
Velasquez, A.N. Pearson, et al., Omics-driven identification and elimination of
valerolactam catabolism in Pseudomonas putida KT2440 for increased product titer,
Metab. Eng. Commun. (2019) e00098. doi:10.1016/j.mec.2019.e00098.

[27]   **T.B. Cook, J.M. Rand, W. Nurani, D.K. Courtney, S.A. Liu, B.F. Pfleger, Genetic**
tools for reliable gene expression and recombineering in Pseudomonas putida., J.
Ind. Microbiol. Biotechnol. 45 (2018) 517–527. doi:10.1007/s10295-017-2001-5.

[28]   **T. Aparicio, V. de Lorenzo, E. Martínez-García, CRISPR/Cas9-Based**
Counterselection Boosts Recombineering Efficiency in Pseudomonas putida.,
Biotechnol. J. 13 (2018) e1700161. doi:10.1002/biot.201700161.

[29]   **T. Aparicio, V. de Lorenzo, E. Martínez-García, CRISPR/Cas9-enhanced ssDNA**
recombineering for Pseudomonas putida., Microb. Biotechnol. 12 (2019) 1076–1089.
doi:10.1111/1751-7915.13453.

[30]   E. Martínez-García, V. de Lorenzo, Engineering multiple genomic deletions in
493   Gram-negative bacteria: analysis of the multi-resistant antibiotic profile of
494   Pseudomonas putida KT2440., Environ. Microbiol. 13 (2011) 2702–2716.
495   doi:10.1111/j.1462-2920.2011.02538.x.

496   **[31]**   N.T. Wirth, E. Kozaeva, P.I. Nikel, Accelerated genome engineering of
497   Pseudomonas putida by I-SceI-mediated recombination and CRISPR-Cas9
498   counterselection., Microb. Biotechnol. (2019). doi:10.1111/1751-7915.13396.

499   **[32]**   M.N. Price, K.M. Wetmore, R.J. Waters, M. Callaghan, J. Ray, H. Liu, et al.,
500   Mutant phenotypes for thousands of bacterial genes of unknown function., Nature.
501   557 (2018) 503–509. doi:10.1038/s41586-018-0124-0.

502   **[33]**   H. Priefert, J. Rabenhorst, A. Steinbüchel, Biotechnological production of vanillin.,
503   Appl. Microbiol. Biotechnol. 56 (2001) 296–314. doi:10.1007/s002530100687.

504   **[34]**   F. Wightman, D.L. Lighty, Identification of phenylacetic acid as a natural auxin in
505   the shoots of higher plants, Physiol. Plant. 55 (1982) 17–24. doi:10.1111/j.1399-
506   3054.1982.tb00278.x.

507   **[35]**   S. Zhou, Y. Lyu, H. Li, M.A.G. Koffas, J. Zhou, Fine-tuning the (2S)-naringenin
508   synthetic pathway using an iterative high-throughput balancing strategy.,
509   Biotechnol. Bioeng. 116 (2019) 1392–1404. doi:10.1002/bit.26941.

510   **[36]**   A. Martinez, S.J. Kolvek, C.L.T. Yip, J. Hopke, K.A. Brown, I.A. MacNeil, et al.,
511   Genetically modified bacterial strains and novel bacterial artificial chromosome
512   shuttle vectors for constructing environmental libraries and detecting heterologous
513   natural products in multiple expression hosts., Appl. Environ. Microbiol. 70 (2004)
514   2452–2463. doi:10.1128/aem.70.4.2452-2463.2004.

515   **[37]**   G. Cambray, J.C. Guimaraes, A.P. Arkin, Evaluation of 244,000 synthetic sequences
516   reveals design principles to optimize translation in Escherichia coli., Nat. Biotechnol.
517   36 (2018) 1005–1015. doi:10.1038/nbt.4238.

518   **[38]**   D. Parke, L.N. Ornston, Toxicity caused by hydroxycinnamoyl-coenzyme A
519   thioester accumulation in mutants of Acinetobacter sp. strain ADP1., Appl. Environ.
520   Microbiol. 70 (2004) 2974–2983.

521   **[39]**   C.N.S. Santos, M. Koffas, G. Stephanopoulos, Optimization of a heterologous
522   pathway for the production of flavonoids from glucose., Metab. Eng. 13 (2011) 392–
523   400. doi:10.1016/j.ymben.2011.02.002.

524   **[40]**   S.R. Park, J.A. Yoon, J.H. Paik, J.W. Park, W.S. Jung, Y.-H. Ban, et al.,
525   Engineering of plant-specific phenylpropanoids biosynthesis in Streptomyces
526   venezuelae., J. Biotechnol. 141 (2009) 181–188. doi:10.1016/j.jbiotec.2009.03.013.

527   **[41]**   Y. Zang, J. Zha, X. Wu, Z. Zheng, J. Ouyang, M.A.G. Koffas, In Vitro Naringenin
528   Biosynthesis from p-Coumaric Acid Using Recombinant Enzymes., J. Agric. Food
529   Chem. (2019). doi:10.1021/acs.jafc.9b00413.

530   **[42]**   Z. Fang, J.A. Jones, J. Zhou, M.A.G. Koffas, Engineering Escherichia coli Co-
531   Cultures for Production of Curcuminoids From Glucose., Biotechnol. J. 13 (2018)

532   e1700576. doi:10.1002/biot.201700576.

[43]   A.E. LaBauve, M.J. Wargo, Growth and laboratory maintenance of Pseudomonas
534   aeruginosa., Curr. Protoc. Microbiol. Chapter 6 (2012) Unit 6E.1.
535   doi:10.1002/9780471729259.mc06e01s25.

[44]   M.G. Thompson, J.M. Blake-Hedges, P. Cruz-Morales, J.F. Barajas, S.C. Curran,
537   C.B. Eiben, et al., Massively Parallel Fitness Profiling Reveals Multiple Novel
538   Enzymes in Pseudomonas putida Lysine Metabolism., MBio. 10 (2019).
539   doi:10.1128/mBio.02577-18.

[45]   T.S. Ham, Z. Dmytriv, H. Plahar, J. Chen, N.J. Hillson, J.D. Keasling, Design,
541   implementation and practice of JBEI-ICE: an open source biological part registry
542   platform and tools., Nucleic Acids Res. 40 (2012) e141. doi:10.1093/nar/gks531.

[46]   J. Chen, D. Densmore, T.S. Ham, J.D. Keasling, N.J. Hillson, DeviceEditor visual
544   biological CAD canvas., J. Biol. Eng. 6 (2012) 1. doi:10.1186/1754-1611-6-1.

[47]   N.J. Hillson, R.D. Rosengarten, J.D. Keasling, j5 DNA assembly design automation
546   software., ACS Synth. Biol. 1 (2012) 14–21. doi:10.1021/sb2000116.

[48]   D.G. Gibson, L. Young, R.-Y. Chuang, J.C. Venter, C.A. Hutchison, H.O. Smith,
548   Enzymatic assembly of DNA molecules up to several hundred kilobases., Nat.
549   Methods. 6 (2009) 343–345. doi:10.1038/nmeth.1318.

[49]   C. Engler, R. Kandzia, S. Marillonnet, A one pot, one step, precision cloning method
551   with high throughput capability., PLoS ONE. 3 (2008) e3647.
552   doi:10.1371/journal.pone.0003647.

[50]   S.G. Grant, J. Jessee, F.R. Bloom, D. Hanahan, Differential plasmid rescue from
554   transgenic mouse DNAs into Escherichia coli methylation-restriction mutants., Proc
555   Natl Acad Sci USA. 87 (1990) 4645–4649.

[51]   R.M.Q. Shanks, D.E. Kadouri, D.P. MacEachran, G.A. O'Toole, New yeast
557   recombineering tools for bacteria., Plasmid. 62 (2009) 88–97.
558   doi:10.1016/j.plasmid.2009.05.002.