***Staphylococcus aureus* can survive in the absence of c-di-AMP upon inactivation of**

the main glutamine transporter

Merve S. Zeden[a^], Igor Kviatkovski[a^], Vinai Thomas Chittezham[b], Paul D. Fey[b] and Angelika

Gründling[a#]

[a] Section of Microbiology and Medical Research Council Centre for Molecular Bacteriology

and Infection, Imperial College London, London SW7 2AZ.

[b] University of Nebraska Medical Center, Department of Pathology and Microbiology,

Omaha, Nebraska, USA.

^ These authors contributed equally

[#] To whom correspondence should be addressed: Angelika Gründling –

a.grundling@imperial.ac.uk

Keywords: *Staphylococcus aureus*, c-di-AMP, glutamine transporter, AlsT, glutamate

transporter, GltS, amino acid transport, OpuD, osmolyte transport

Running title: Regulation of c-di-AMP by glutamine transport

Summary (250 words)

The nucleotide second messenger c-di-AMP negatively regulates potassium and osmolyte uptake in *Staphylococcus aureus* and many other bacteria. c-di-AMP is also important for growth and an *S. aureus* strain deleted for the c-di-AMP cyclase gene *dacA* is unable to survive in rich medium unless it acquires compensatory mutations. Previously, we have shown that an *S. aureus dacA* mutant can grow after the acquisition of inactivating mutants in *opuD*, encoding the main glycine-betaine osmolyte transporter, or mutations in *alsT*, encoding a predicted amino acid transporter. Using the size of bacterial cells as a proxy for their osmotic balance, we show that inactivation of OpuD helps bacteria to re-establish their osmotic balance, while inactivation of AlsT does not and bacteria remain enlarged, a characteristic of *S. aureus* cells unable to produce c-di-AMP. We show that AlsT is the main glutamine transporter in *S. aureus*, thus revealing that *S. aureus* can survive without c-di-AMP when glutamine uptake is prevented. Using a bioinformatics approach combined with uptake assays, we identified GltS as the main glutamate transporter in *S. aureus*. Using WT and mutant strain, we show that glutamine is preferred over glutamate for bacterial growth and that its uptake represses c-di-AMP production. Glutamine and glutamate are important players in osmotic regulation, but their cellular levels also serve as a key indicator of nitrogen availability in bacterial cells. Therefore, we not only provide a further connection between the c-di-AMP signalling network and osmotic regulation in *S. aureus* but also to central nitrogen metabolism.

Introduction

In the human host, S*taphylococcus aureus* can infect various niches, including, but not limited to, blood, kidneys, bones, heart, soft tissue and lungs (Kluytmans *et al.*, 1997, Fridkin *et al.*, 2005). *S. aureus* can colonize these different tissues thanks to its refined regulatory mechanisms that allow it to rapidly respond to external stimuli. Amongst others, this allows the organism to adapt its metabolism and utilize different carbon and nitrogen sources available in each specific niche (Fridkin *et al.*, 2005, Spahich *et al.*, 2016, Vitko *et al.*, 2015, Crooke *et al.*, 2013, Fuller *et al.*, 2011, Richardson *et al.*, 2008, Halsey *et al.*, 2017, Lehman *et al.*, 2019).

Glucose is the preferred carbon source for *S. aureus,* but it can be limiting during infection due to the host immune response (Kelly & O'Neill, 2015, Spahich *et al.*, 2016, Halsey *et al.*, 2017, Lehman *et al.*, 2019). In glucose-limiting conditions, *S. aureus* instead catabolizes secondary carbon sources; amino acids, particularly glutamate and proline, serve as major carbon sources during growth in the absence of glucose (Halsey *et al.*, 2017). However, not much is known about amino acid uptake and catabolism in *S. aureus* and how the availability of certain nutrients can affect virulence factor expression and invasion of the host. While a large number of amino acid transporters and oligopeptide permeases can be identified bioinformatically, their actual substrate specificities and functions in *S. aureus* have not yet been studied in detail. Also, predicting the substrates for transporters bioinformatically remains difficult and hence such questions need to be addressed experimentally.

Secondary messenger molecules are crucial in allowing bacteria to rapidly adapt to different environmental and host cell niches (Römling, 2008, Hengge, 2009). There is now considerable evidence that one such messenger, cyclic di-adenosine monophosphate (c-di-AMP) plays a significant role in osmoregulation in bacteria (Pham *et al.*, 2018, Pham & Turner, 2019, Quintana *et al.*, 2019, Zarrella *et al.*, 2018, Teh *et al.*, 2019, Fahmi *et al.*, 2019, Devaux *et al.*, 2018, Bai *et al.*, 2014, Zeden *et al.*, 2018, Corrigan *et al.*, 2011, Rocha *et al.*, 2019, Gundlach *et al.*, 2017b, Gundlach *et al.*, 2017a, Witte *et al.*, 2013, Whiteley *et*

al., 2015, Whiteley et al., 2017). c-di-AMP binds to and negatively regulates a number of different potassium and osmolyte importers (Rocha et al., 2019, Quintana et al., 2019, Kim et al., 2015, Corrigan et al., 2013, Moscoso et al., 2015, Chin et al., 2015, Huynh et al., 2016, Schuster et al., 2016, Pham & Turner, 2019, Pham et al., 2018, Devaux et al., 2018, Zarrella et al., 2018, Gundlach et al., 2017b, Gundlach et al., 2017a, Gundlach et al., 2017c). c-di-AMP is essential for bacterial growth under standard growth conditions but it is also toxic at high levels in many Firmicutes, hence its cellular levels must be tightly regulated (Gundlach et al., 2015b, Mehne et al., 2013, Corrigan et al., 2011, Corrigan et al., 2015, Woodward et al., 2010, Witte et al., 2013). In S. aureus and Listeria monocytogenes, deletion of the diadenylate cyclase gene dacA, the enzyme responsible for the synthesis of c-di-AMP, was only possible in chemically defined medium (Whiteley et al., 2015, Zeden et al., 2018, Devaux et al., 2018), whereas in Bacillus subtilis all three c-di-AMP cyclases could only be inactivated in minimal medium also containing low amounts of potassium (Gundlach et al., 2017b).

Previously, we found that inactivation of the main glycine betaine transporter OpuD as well as the predicted amino acid transporter AlsT (SAUSA300_1252) allows an S. aureus dacA mutant to grow in rich medium in the absence of c-di-AMP (Zeden et al., 2018). In several other Firmicutes, including B. subtilis, Lactococcus lactis, Streptococcus pneumoniae, Streptococcus pyogenes and L. monocytogenes, inactivating mutations have also been identified in osmolyte and potassium transport systems that allow these bacteria to grow in the absence of c-di-AMP (Pham et al., 2018, Pham & Turner, 2019, Quintana et al., 2019, Zarrella et al., 2018, Teh et al., 2019, Fahmi et al., 2019, Devaux et al., 2018, Bai et al., 2014, Zeden et al., 2018, Corrigan et al., 2011, Rocha et al., 2019, Gundlach et al., 2017b, Gundlach et al., 2017a, Witte et al., 2013, Whiteley et al., 2015, Whiteley et al., 2017). This is consistent with the idea that in the absence of c-di-AMP, potassium and osmolyte transporters are more active, resulting in the accumulation of toxic levels of potassium and osmolytes in the cell. Consistent with a key function of c-di-AMP in regulating the osmotic balance in the cell, we found that S. aureus cells show significant differences in cell size

depending on their intracellular c-di-AMP levels (Zeden *et al.*, 2018, Corrigan *et al.*, 2011).

Bacteria of the high c-di-AMP level *S. aureus* mutant strain LAC*gdpP* show a decrease in

cell size, while cells of the low level c-di-AMP strain LAC*dacA*$_{G206S}$ show an increase in cell

size (Zeden *et al.*, 2018, Corrigan *et al.*, 2011). As c-di-AMP negatively regulates potassium

and osmolyte uptake (Corrigan *et al.*, 2013, Moscoso *et al.*, 2015, Schuster *et al.*, 2016), the

increase in cell size is consistent with the hypothesis that an increase in potassium and

osmolyte uptake and retention of water at reduced c-di-AMP levels leads to the observed

increase in cell size. As part of this study, we further investigated the mechanisms by which

inactivation of the main glycine betaine transporter OpuD and the predicted amino acid

transporter AlsT allow *S. aureus* to survive in the absence of c-di-AMP.

The counterion of potassium in the cell is glutamate, which at the same time also

serves as main nitrogen donor for cellular metabolites and macromolecules. A correlation

between cellular levels of c-di-AMP, glutamate and glutamine has been reported for several

Firmicutes, including *B. subtilis* and *L. monocytogenes* (Whiteley *et al.*, 2017, Gundlach *et*

al., 2015b, Gundlach *et al.*, 2017a, Sureka *et al.*, 2014). A two-fold increase in cellular c-di-

AMP levels was observed in *B. subtilis* when bacteria where grown in Spizizen minimal

medium with glutamate (Glu) as compared to growth in the same medium with glutamine

(Gln) as nitrogen source (Gundlach *et al.*, 2015b). In *L. monocytogenes,* c-di-AMP is a

negative regulator of the key TCA cycle enzyme pyruvate carboxylase (Sureka *et al.*, 2014).

The depletion of c-di-AMP in the cell resulted in an increased flux of glucose into the

production of glutamine and glutamate, likely due to increased pyruvate carboxylase activity

and increased flux into the TCA cycle at reduced c-di-AMP levels (Sureka *et al.*, 2014).

Interestingly, an *L. monocytogenes dacA* mutant that also lacks *citZ*, which codes for the key

TCA cycle enzyme citrate synthase, is viable in rich medium (Sureka *et al.*, 2014, Whiteley

et al., 2017). In a *citZ* mutant the TCA cycle is blocked and it was shown that depletion of c-

di-AMP in this mutant no longer results in the accumulation of glutamate and glutamine in

the cell (Sureka *et al.*, 2014). Therefore, the amino acids glutamine and glutamate not only

play an important function in the osmotic regulation in bacterial cells, but also have a

particularly critical role in TCA cycle function and their cellular ratio has an important role in

signalling nitrogen-limiting (high glutamate levels and low glutamine levels) or nitrogen

excess (high glutamine and low glutamate levels) conditions (Halsey *et al.*, 2017, Gundlach

et al., 2017a).

As part of this study, we further investigated why inactivation of the main glycine-

betaine transporter OpuD and the predicted amino acid transporter AlsT allows *S. aureus* to

grow in the absence of c-di-AMP. Our results indicate that inactivation of OpuD helps

bacteria to re-establish their osmotic balance, while inactivation of AlsT, which we show here

is the main glutamine transporter in *S. aureus,* functions differently to bypass the essentiality

of c-di-AMP and possible mechanisms are discussed. We also identified the *S. aureus* GltS

protein as the main glutamate transporter in *S. aureus* and show that glutamine but not

glutamate uptake represses the production of c-di-AMP. With this study, we not only provide

a further link between the c-di-AMP signalling network and osmotic regulation in bacterial

cells but also to the central nitrogen metabolism in *S. aureus.*

Results

Inactivation of OpuD but not AlsT reduces the cell size of an *S. aureus dacA* mutant

In previous work, we reported a correlation between the cell size and c-di-AMP levels in *S.*

aureus: bacteria with high c-di-AMP level are smaller, whereas bacteria with low c-di-AMP

levels (strain LAC*$dacA_{G206S}$) are larger as compared to wild-type bacteria (Zeden *et al.*,

2018, Corrigan *et al.*, 2011). We also reported that inactivating mutations in *opuD*

(*SAUSA300_1245*) coding for the main glycine betaine osmolyte transporter and *alsT*

(*SAUSA300_1252*) coding for a predicted amino acid sodium symporter, rescue the growth

defect observed for the c-di-AMP negative *S. aureus* strain LAC*$dacA::kan$ in rich medium

(Zeden *et al.*, 2018). To investigate the mechanism by which the growth defect of the *dacA*

mutant strain is rescued in the LAC*$dacA/opuD$ and LAC*$dacA/alsT$ suppressor strains and

to assess if restoring the osmotic imbalance could be a contributing factor, the cell size of

WT   and   different   *S. aureus*   mutants   was   determined.   Initially,   the   cell   size   of

LAC*dacA/opuD and LAC*dacA/alsT suppressor strain bacteria was compared to that of WT

LAC* and the low-level c-di-AMP LAC*dacA$_{G206S}$ strain after growth in TSB medium. As

expected, the low-level c-di-AMP bacteria showed an increase in cell size as compared to

WT bacteria (Fig. 1A and B). While a similar increase in cell size was still observed for

bacteria of strain LAC*dacA/alsT, the cell size of LAC*dacA/opuD bacteria, while still

increased as compared WT, was significantly smaller as compared to the low-level

LAC*dacA$_{G206S}$ strain (Fig. 1A and B). As TSB medium is not suitable for the growth of the c-

di-AMP null strain LAC*dacA::kan, bacterial cell sizes were also determined for the WT and

mutant S. aureus strains following growth in TSB supplemented with 0.4 M NaCl, which is

permissive for the growth of the dacA mutant (Fig. 1C-F). Bacteria from all strains had a

reduced cell size when grown in TSB supplemented with 0.4 M NaCl compared to bacteria

grown in TSB (Fig. 1). Similar to what was observed for the low level c-di-AMP dacA$_{G206S}$

mutant strain, the size of bacteria from the c-di-AMP null strain LAC*dacA::kan was

significantly increased compared to WT bacteria. As observed before, the cell size was not

rescued for bacteria of the LAC*dacA/alsT suppressor strain (Fig. 1C-F). On the other hand,

the size of LAC*dacA/opuD bacteria was similar to that of WT bacteria (Fig. 1C-F). Taken

together, these data indicate that inactivation of OpuD, and hence reduced glycine betaine

transport, likely helps bacteria to survive in the absence of c-di-AMP by re-establishing the

osmotic balance in the mutant while a different mechanism is at play for the alsT mutant.

Bacteria lacking alsT have an altered amino acid uptake profile

AlsT (SAUSA300_1252) is a predicted amino acid transporter protein and annotated as an

alanine/sodium symporter. However, no difference in the uptake of radiolabelled alanine was

detected between a WT and the LAC*dacA/alsT mutant strain in our previous study (Zeden

et al., 2018), indicating that AlsT is not an alanine transporter. To identify potential

substrates for the S. aureus AlsT transporter, we followed the depletion of different amino

acids from the culture supernatant during the growth of the WT and alsT mutant strains in

TSB medium. To this end, strain LAC*alsT::tn containing a transposon insertion in alsT was

constructed by phage transducing the *alsT::tn* region from the Nebraska Transposon Mutant

Library (NMTL) strain NE142 (Fey *et al.*, 2013) into the *S. aureus* LAC* strain background.

WT LAC* and the *alsT* mutant strain LAC**alsT::tn* showed similar growth rates when grown

in TSB medium (Fig. 2A). Next, their ability to take up different amino acids was assessed by

determining the levels of the individual amino acids in the culture supernatant at the start of

the experiment (T = 0 h) as compared to 6, 10 and 12 h following their growth in TSB

medium. While no significant differences were observed for most amino acids (Fig. S1), a

slight increase in the utilization of aspartate and a slight decrease in the uptake of serine

was observed (Fig. 2B and 2C), suggesting that AlsT could potentially be a serine

transporter. To test this, uptake assays were performed with radiolabelled serine using the

WT LAC* strain, the *alsT* mutant strain LAC**alsT::tn* piTET as well as the complementation

strain LAC**alsT::tn* piTET-*alsT*. However, no significant differences in the uptake rate of

serine were observed between the strains (Fig. 3A), indicating that AlsT is not the main

serine transporter in *S. aureus*. While slight differences in the amino acid uptake profile were

observed between the WT and *alsT* mutant strain, this analysis did not allow us to identify

the main substrate for AlsT. However, it is of note that using this method one cannot

distinguish between glutamine and glutamate or asparagine and aspartate utilization.

Additionally, tryptophan was not measured due to the limitations of the method used.

AlsT is the main glutamine transporter in *S. aureus*

Next, a more detailed bioinformatics analysis was performed to identify potential AlsT

substrates. A BlastP search against the *B. subtilis* 168 genome led to the identification of

four close homologs of AlsT (SAUSA300_1252), namely AlsT (e-value: e-166), GlnT (e-

value: e-149), YrbD (e-value: e-117) and YflA (e-value: 2e-72). Of note, an AlsT homologue

SAUSA300_0914 is also present in the *S. aureus* FPR3757 genome (Fig. S2). While *S.*

aureus AlsT (SAUSA300_1252) shows the highest similarity to the *B. subtilis* AlsT and GlnT

proteins, SAUSA300_0914 has the highest similarity with the *B. subtilis* YrbD protein. AlsT is

annotated in *B. subtilis* as a potential glutamine sodium symporter, but to the best of our

knowledge, this has not yet been experimentally verified. Expression of *alsT* is controlled in

B. subtilis by GlnR and TnrA, the two main transcriptional regulators adjusting gene

expression in response to nitrogen availability (Randazzo *et al.*, 2017, Yoshida *et al.*, 2003,

Mirouze *et al.*, 2015). GlnT is a confirmed glutamine transporter in *B. subtilis* (Satomura *et*

al., 2005), whose production is induced in the presence of glutamine as nitrogen source by

the two-component system GlnKL. To test if *S. aureus* AlsT is a potential glutamine or

glutamate transporter, uptake assays were performed with radiolabelled glutamine and

glutamate using the WT *S. aureus* strain LAC*, the *alsT* mutant LAC**alsT::tn* piTET and the

complementation strain LAC**alsT::tn* piTET-*alsT*. Uptake of glutamine, but not of glutamate,

was severely reduced in the *alsT* mutant when compared to the WT (Fig. 3B). This defect

was restored upon expression of *alsT* in the complementation strain (Fig. 3C). To confirm

that *alsT* also functions as main glutamine transporter in the LAC**dacA/alsT* suppressor

strain, uptake assays were also performed with strain LAC**dacA/alsT* and compared to that

of the WT LAC* and LAC**dacA::kan* control strains (Fig. 3D-F). Similar to what was

observed for the *alsT* mutant, glutamine uptake was severely attenuated in strain

LAC**dacA/alsT* when compared to the control strains (Fig. 3D-F). These data highlight that

under the growth conditions tested, AlsT functions as the main glutamine transporter in *S.*

aureus. A slight reduction in glutamine uptake was seen in the absence of c-di-AMP,

suggesting that c-di-AMP levels can impact glutamine uptake in *S. aureus.* But perhaps

most importantly, our data suggest that bacteria lacking c-di-AMP can survive in rich

medium when glutamine uptake is blocked.

Investigating the contribution of SAUSA300_0914 and GlnPQ to glutamine and

glutamate transport in *S. aureus*

S. aureus SAUSA300_0914 codes for a predicted amino acid symporter, which shows 41%

identity with the *S. aureus* AlsT protein. After assigning AlsT a function as glutamine

transporter, we wanted to test if SAUSA300_0914 might also play a role in glutamine or

glutamate transport. To this end, strain LAC**0914::tn* was constructed by transducing the

genomic region from the NMTL strain NE1463 (Fey *et al.*, 2013) containing a transposon

insertion in *SAUSA300_0914* into the *S. aureus* LAC* background. Subsequently uptake of

radiolabelled glutamine and glutamate was assessed (Fig. 4A-B). No significant differences

in the uptake of these amino acids was observed between WT LAC* and strain

LAC**0914::tn*, showing that SAUSA300_0914 does not function as glutamine or glutamate

transporter under our assay conditions.

AlsT and SAUSA300_0914 are members of the amino acid-sodium symporter family of

transporters, which are composed of a single multimembrane spanning protein. Besides this

type of transporter, GlnPQ-type ABC transporters play a major role in glutamine and

glutamate transport in other bacteria (Schuurman-Wolters & Poolman, 2005). *S. aureus*

contains a *glnPQ (SAUSA300_1808 - SAUSA300_1807)* operon with *glnP* coding for a

substrate binding domain-permease fusion protein and *glnQ* coding for the cytoplasmic

nucleotide-binding ATPase domain. The results from a previous study suggested that this

transporter functions as glutamine transporter in *S. aureus*, as a *glnP* mutant was more

resistant to the toxic glutamine analogue gamma-L-glutamyl hydrazide (Zhu *et al.*, 2009). To

assess the contribution of the GlnPQ transporter to glutamine and glutamate transport in *S.*

aureus under our assay conditions*,* strain LAC**glnQ::tn* was generated by transducing the

glnQ::tn region from the NMTL strain NE153 (Fey *et al.*, 2013) into the LAC* background.

The resulting LAC**glnQ::tn* mutant strain displayed no difference in glutamine or glutamate

uptake compared to WT LAC* (Fig. 4C-D), indicating that the ABC transporter GlnPQ does

not function as a main glutamate or glutamine transporter under our assay conditions.

GltS (SAUSA300_2291) is the main glutamate transporter in *S. aureus*

Cellular glutamine levels are key in signalling nitrogen availability in bacterial cells (Fisher,

1999, Gunka & Commichau, 2012). In addition, differences in cellular c-di-AMP levels were

reported in *B. subtilis* depending on the presence of glutamine or glutamate as available

nitrogen source. More specifically, an increase in cellular c-di-AMP levels was observed in

the presence of glutamate as compared to glutamine (Gundlach *et al.*, 2015b). *S. aureus*

does not only take up glutamine but also shows robust glutamate uptake (Figs 3 and 4).

However, none of the transporters (AlsT, SAUSA300_0914 and GlnPQ) investigated so far

plays a major role in glutamate uptake under our growth conditions. In *B. subtilis* GltT,

belonging to the dicarboxylate/amino acid cation symporter (DAACS) family of proteins, is a

major high-affinity Na^+-coupled glutamate/aspartate symporter and can also mediate the

uptake of glyphosate (Wicke *et al.*, 2019). An additional two paralogs, DctP and GltP are

found in *B. subtilis* of which GltP has also been shown to be a glutamate transporter (Tolner

et al., 1995). The *S. aureus* protein SAUSA300_2329 (from here on referred to as GltT)

shows a high degree of similarity (52% identity) to the *B. subtilis* GltT protein. In addition, we

identified SAUSA300_2330 (from here on referred to as GltS) as a potential glutamate

transporter in *S. aureus* due to its similarity to the *E. coli* glutamate permease GltS (e-value:

6e-77; 38% identity) (Deguchi *et al.*, 1990). To test if *S. aureus* GltT or GltS are glutamate

transporters or impact glutamine uptake in *S. aureus*, strains LAC**gltT::tn* and LAC**gltS::tn*

were constructed by moving the respective *gltT and gltS* transposon insertion regions from

the NMTL strains NE566 and NE560 (Fey *et al.*, 2013) into the LAC* strain background.

Next, the uptake of radiolabelled glutamine and glutamate was assessed for the WT LAC*

strain and the LAC* *gltT::tn* and LAC* *gltS::tn* mutants. No difference in the uptake of

glutamine was observed between the strains (Fig. 5A) and in the case of LAC**gltT::tn*, also

no difference in the uptake of glutamate was observed. However, a significant reduction in

glutamate uptake was observed for strain LAC**gltS::tn* when compared to the WT (Fig. 5B).

The glutamine uptake defect could be restored in a complementation strain harbouring

plasmid piTET-*gltS* allowing for inducible *gltS* expression (Fig. 5C). Indeed, increased

glutamate uptake was observed in the complementation strain, indicating increased *gltS*

expression in the complementation strain as compared to a WT strain. Taken together,

these data reveal that under the growth conditions tested, GltS is the main glutamate

transporter in *S. aureus*.

Glutamine but not glutamate stimulates the growth of *S. aureus* in CDM lacking ammonium as nitrogen source.

Glutamine and glutamate are important amino acids that can serve as nitrogen sources for the synthesis of many other cellular metabolites. To examine the effect of these amino acids on the growth of *S. aureus* as well as to evaluate the contribution of the glutamine (AlsT) and glutamate (GltS) transporters for growth, growth curves were performed with WT, *alsT::tn* and *gltS::tn* mutant strains in chemically defined medium (CDM) lacking ammonium and containing either glutamine (CDM+Gln) or glutamate (CDM+Glu). Reduced growth was seen when the *S. aureus* strains were grown in glutamate as compared to glutamine containing medium, suggesting that glutamine but not glutamate can stimulate the growth of *S. aureus* in CDM lacking ammonium as nitrogen source (Fig. 6A). Consistent with this, the *alsT::tn* mutant strain, which is deficient in glutamine uptake, showed a similar growth reduction even if grown in the glutamine-containing medium (CDM+Gln) (Fig. 6A). The observed growth defect for the *alsT* mutant in CDM+Gln could be restored in the complementation strain harbouring the plasmid piTET-*alsT* (Fig. 6B). On the other hand, the *gltS::tn* mutant, which is defective in glutamate uptake, grew similar to the WT strain under all conditions tested, suggesting that glutamate uptake does not impact the growth of *S. aureus* under the test conditions. Taken together, these data indicate that glutamine is preferred over glutamate for the growth of *S. aureus* in CDM lacking ammonium as nitrogen source.

Glutamine uptake leads to a reduction in the cellular c-di-AMP levels in *S. aureus*

In a previous study, it has been reported that the presence of glutamine or glutamate in the growth medium can affect cellular c-di-AMP levels in *B. subtilis* and it was proposed that glutamate uptake leads to an activation of c-di-AMP synthesis in this organism (Gundlach *et al.*, 2015b). To assess if the presence of glutamine or glutamate would also affect c-di-AMP

levels in *S. aureus*, the intracellular c-di-AMP concentrations were determined for the WT *S.*

aureus strain LAC* following growth in CDM+Gln or CDM+Glu medium. c-di-AMP levels

were significantly higher in the presence of glutamate as compared to glutamine (Fig. 7A).

As previously reported (Corrigan *et al.*, 2011), in the absence of the c-di-AMP

phosphodiesterase GdpP (strain LAC*gdpP::kan) c-di-AMP levels were increased as

compared to a WT strain (Fig. 7A). Interestingly and similar as observed for the WT strain, c-

di-AMP levels were also higher in the *gdpP* mutant strain in the presence of glutamate as

compared to glutamine. This indicates that the observed regulation of c-di-AMP synthesis

depending on the presence of glutamine or glutamate is at the level of synthesis and not

degradation. To test if glutamine uptake inhibits or glutamate uptake activates c-di-AMP

production, c-di-AMP levels were also determined for the *alsT::tn* and *gltS::tn* mutants, which

are defective in glutamine or glutamate uptake, respectively. Following growth in glutamate

containing medium, all strains produced high and comparable levels of c-di-AMP (Fig. 7B

white columns). However, some variation in the relative c-di-AMP levels produced by the

gltS mutant compared to the WT strain was observed between experiments (Fig. 7B white

columns and Fig. S3). The reason for this is currently not known. But taken together, our

data suggest that the ability of *S. aureus* to take up glutamate does not drastically affect c-di-

AMP production. On the other hand, clear differences in c-di-AMP levels were observed for

strain *alsT::tn*, which is unable to take up glutamine. The WT and the *gltS::tn* mutant strains

produced low c-di-AMP amounts following growth in CDM+Gln, while the c-di-AMP levels

remained high in strain *alsT::tn.* (Fig. 7B grey columns and Fig. S3), suggesting that

glutamine uptake inhibits c-di-AMP production. The c-di-AMP levels in the *alsT* mutant strain

could be restored back to WT levels in the complementation strain harbouring plasmid

piTET-*alsT* (Fig. 7C). Taken together, these results highlight that glutamine uptake blocks c-

di-AMP production in *S. aureus* and that eliminating glutamine from the medium or

preventing its uptake stimulates c-di-AMP production. Such an activation is likely achieved

through stimulating the activity of the c-di-AMP cyclase DacA, rather than preventing its

degradation by GdpP and possible mechanisms for this will be discussed.

Discussion

Under standard laboratory growth conditions, c-di-AMP is essential for growth in many bacteria (Mehne *et al.*, 2013, Corrigan *et al.*, 2015, Woodward *et al.*, 2010, Gundlach *et al.*, 2015a), but the exact molecular mechanisms behind this have remained unclear. Here we have shown that inactivation of OpuD likely helps a *S. aureus* c-di-AMP null strain survive by allowing bacteria to re-establish their osmotic balance. Bacteria unable to produce c-di-AMP are larger than WT cells but bacteria that are unable to produce c-di-AMP and also lack OpuD, the main transporter for the osmolyte glycine betaine in *S. aureus,* are similar in size to WT bacteria. This indicates that in these cells the osmotic balance has been restored. Furthermore, we show that mutations in *alsT*, which we identify as part of this study to encode for the main glutamine transporter in *S. aureus*, suppress the essentiality of c-di-AMP in a different way and potential mechanisms for this are discussed here.

Over the last decade, considerable evidence has emerged that c-di-AMP plays a major role in osmotic regulation, primarily by positively regulating potassium export or negatively regulating potassium and osmolyte uptake (Rocha *et al.*, 2019, Quintana *et al.*, 2019, Kim *et al.*, 2015, Corrigan *et al.*, 2013, Moscoso *et al.*, 2015, Chin *et al.*, 2015, Huynh *et al.*, 2016, Schuster *et al.*, 2016, Pham & Turner, 2019, Pham *et al.*, 2018, Devaux *et al.*, 2018, Zarrella *et al.*, 2018, Gundlach *et al.*, 2017b, Gundlach *et al.*, 2017a, Gundlach *et al.*, 2017c, Gundlach *et al.*, 2019). However, individual c-di-AMP target proteins identified thus far are themselves not essential. Therefore, the essentiality of c-di-AMP is likely due to its ability to regulate multiple target proteins simultaneously. Furthermore, in the absence of this molecule, many transporters are activated rather than inactivated, likely leading to accumulation of toxic levels of metabolites, such as potassium and osmolytes. Consistent with this idea, inactivating mutations in potassium uptake systems, oligopeptide and osmolyte transporters have been reported to rescue the growth defect of bacteria unable to produce c-di-AMP (Whiteley *et al.*, 2015, Whiteley *et al.*, 2017, Gundlach *et al.*, 2017b, Gundlach *et al.*, 2017c, Pham *et al.*, 2018, Devaux *et al.*, 2018, Zeden *et al.*, 2018). We have previously shown that in *S. aureus* inactivation of the main glycine betaine transporter

OpuD bypasses the requirement of c-di-AMP for the growth of *S. aureus* in rich medium

(Zeden *et al.*, 2018). Using bacterial cell size as a proxy for the osmotic balance of cells, we

show here that inactivation of OpuD likely helps an c-di-AMP null strain survive by allowing

bacteria to re-establish their osmotic balance, as *dacA/opuD* mutant bacteria, which cannot

produce c-di-AMP but are also lacking the main glycine betaine transport, are similar in size

to WT bacteria (Fig. 1). Here it is interesting to note that while the carnitine osmolyte

transporter OpuCA has been shown to be a direct target of c-di-AMP in both *S. aureus* and

L. monocytogenes (Huynh *et al.*, 2016, Schuster *et al.*, 2016), no direct interaction between

OpuD or any other glycine-betaine transporter and c-di-AMP has been reported for *S.*

aureus. Hence, it remains unclear if glycine betaine osmolyte transport is directedly

regulated by c-di-AMP in *S. aureus* and hence the absence of c-di-AMP leads to an excess

in glycine-betaine uptake and therefore inactivation of OpuD prevents such excess in

uptake. Or alternatively, only potassium and carnitine uptake might be increased in the

absence of c-di-AMP since transporters of these molecules are direct targets of c-di-AMP,

and glycine betaine uptake remains unaffected but becomes toxic upon accumulation of

potassium and carnitine. Therefore, it remains to be determined if the observed increase in

cell size in the absence of c-di-AMP is solely caused by an increase in potassium transport

by the Ktr, Kpd and potentially KimA systems and carnitine osmolyte transport by OpuCA or

if also glycine betaine transport is directly affected by cellular c-di-AMP levels in *S. aureus*. A

direct role for c-di-AMP in the control of glycine betaine or betaine transporters has been

proposed for other bacteria where c-di-AMP binds to the transcriptional regulator BusR,

which controls the expression of the predicted glycine betaine or betaine transporter BusAB

(Devaux *et al.*, 2018, Pham *et al.*, 2018). However, such a system does not appear to be

present in *S. aureus*.

Bacteria of the *dacA/alsT* suppressor strain, which survive in the absence of c-di-

AMP, remained enlarged, indicating that the essentiality of c-di-AMP is bypassed in this

strain through a different mechanism. Here, we show that AlsT is the main glutamine

transporter in *S. aureus* (Fig. 3). Glutamine as well as proline accumulate under NaCl stress

conditions in *S. aureus,* indicating that glutamine also plays an important role in osmotic

regulation (Anderson & Witter, 1982). However, under the osmotic stress conditions tested

in this previous study, glutamine accumulation was proposed to be due to synthesis rather

than uptake (Anderson & Witter, 1982). In terms of other functions of the glutamine

transporter AlsT in *S. aureus;* it is interesting to note that in a recent study investigating

genetic determinants required for eDNA during biofilm formation, it was found that

inactivation of GdpP as well as AlsT results in a significant decrease in eDNA release

(DeFrancesco *et al.*, 2017). Since we show here that in an *alsT* mutant, which is unable to

import glutamine, cellular c-di-AMP levels can be significantly higher as compared to a WT

strain (Fig. 7), similar to a *gdpP* mutant, this could mean that the underlying mechanistic

bases for the decrease in eDNA release observed for the *gdpP* and *alsT* mutant strains

might be related.

There are several (not mutually exclusive) possibilities how preventing glutamine

uptake could rescue the growth of a c-di-AMP null strain (Fig. 8). The cellular

glutamine/glutamate ratio serves as a key indicator of nitrogen availability in bacterial cells

with a high glutamine/glutamate ratio indicating nitrogen availability (Forchhammer, 2007).

The cellular glutamate concentration is usually higher than the glutamine concentration and

excess glutamine can be readily converted to glutamate via the GOGAT pathway by the

glutamine oxoglutarate aminotransferase composed in *S. aureus* of the GltB and GltD

proteins subunits (Gunka & Commichau, 2012). Hence, glutamine uptake and its availability

in the cell will provide a flux towards glutamate synthesis and glutamate is the counterion of

potassium in the cell. Therefore, glutamine uptake and its conversion to glutamate could

indirectly facilitate further potassium uptake, which becomes toxic in a c-di-AMP null strain,

in which potassium influx is already increased. As a result, reduction in glutamine uptake, as

observed in the *alsT* mutant, and preventing its flux to glutamate, could also prevent the

intoxication of cells with potassium.

As stated above, a high glutamine/glutamate ratio indicates nitrogen availability and

in the absence of other limitations, this will boost the general metabolism of bacterial cells

(see Model Fig. 8). In a study on *L. monocytogenes,* an increased flux of pyruvate into the

TCA cycle has been described for bacteria unable to produce c-di-AMP (Sureka *et al.*,

2014). As a consequence of this increased TCA cycle activity, an accumulation of citrate and

increased carbon flux into glutamine and glutamate was observed (Sureka *et al.*, 2014). Of

note, the authors did not distinguish between glutamine and glutamate in this study (Sureka

et al., 2014). This provided experimental evidence that a decrease in c-di-AMP levels leads

to increased TCA cycle activity and accumulation of cellular metabolites such as citrate and

glutamine/glutamate. The essentiality of c-di-AMP in *L. monocytogenes* could be reversed

by mutating *citZ,* coding for the citrate synthase, which prevented the accumulation of high

levels of citrate as well as glutamine and glutamate pool (again analyzed as a combined

pool) in bacterial cells (Sureka *et al.*, 2014). Perhaps similar to the observations in *L.*

monocytogenes, the absence of c-di-AMP could also boost the metabolism and potentially

TCA cycle activity in *S. aureus*. As part of this study, we provide evidence that in *S. aureus*

glutamine is preferred over glutamate for growth in CDM lacking ammonium as nitrogen

source. The growth of an *S. aureus* strain in this glucose-containing but ammonium free

medium was improved by the addition of glutamine but not glutamate and the uptake of

glutamine mediated by AlsT was required for this growth improvement (Fig. 6). Hence, the

lack of c-di-AMP combined with glutamine uptake could fuel the bacterial metabolism and

the resulting metabolic imbalance might become toxic to the cell, similar as observed for *L.*

monocytogenes (Sureka *et al.*, 2014, Whiteley *et al.*, 2017). This futile cycle might be

blocked by preventing glutamine uptake and reducing the metabolic activity of cells.

The actual stimuli and underlying molecular mechanisms that regulate c-di-AMP

production in bacterial cells are at the moment poorly understood. As part of this study, we

show that glutamine uptake negatively impacts c-di-AMP production in *S. aureus* and

bacteria grown in medium lacking glutamine or inactivated for the main glutamine transporter

AlsT have significantly increased cellular c-di-AMP levels (Fig. 7B and 7C). The increase in

c-di-AMP production in the absence of glutamine uptake is likely achieved by activation of

the c-di-AMP cyclase DacA and not inhibition of the c-di-AMP phosphodiesterase GdpP, as

an increase in cellular c-di-AMP levels was also detected in a *gdpP* mutant strain when

grown in medium lacking glutamine (Fig. 7A). Current evidence suggests that the activity of

DacA can be regulated through the interaction with two proteins: the membrane anchored

regulator protein YbbR (also name CdaR in other bacteria) and the phosphoglucomutase

enzyme GlmM (Tosi *et al.*, 2019, Zhu *et al.*, 2016, Gundlach *et al.*, 2015b, Pham *et al.*,

2016) (see Model Fig. 8). YbbR and GlmM are encoded in the same operon with DacA.

GlmM converts glucosamine-6-P to glucosamine-1-P, an essential precursor for UDP-

GlcNAc production and hence peptidoglycan synthesis. GlmM can block the c-di-AMP

cyclase activity of DacA through a direct interaction (Tosi *et al.*, 2019, Zhu *et al.*, 2016).

Glutamine is a key precursor for the production of the GlmM substrate glucosamine-6-P

since it and fructose-6-P are converted by GlmS to glutamate and glucosamine-6-P.

Therefore, the cellular glutamine levels will impact GlmS activity and hence also the

availability of the GlmM substrate, which could in turn impact the ability of GlmM to interact

with DacA. Based on our findings that c-di-AMP levels are increased in the absence of

glutamine in the medium or in the *alsT* mutant would suggest that low glutamine levels

would prevent an interaction between GlmM and DacA, resulting in an increase in c-di-AMP

production. *Vice versa*, high glutamine level could stimulate the interaction between GlmM

and DacA and in this manner reduce c-di-AMP production and perhaps stimulate the

bacterial metabolism (Fig. 8). In other bacteria it has been reported that YbbR can,

depending on the growth conditions, either act as an activator or repressor of DacA (Mehne

et al., 2013, Rismondo *et al.*, 2016). It is thought that this regulation is achieved through a

direct interaction between the membrane spanning helix of YbbR and the transmembrane

domain in DacA (Gundlach *et al.*, 2015b). YbbR has an extracellular sensor domain, which

could respond to changes in the peptidoglycan structure (e.g. caused by changes in

glutamine levels and flux of precursors towards peptidoglycan synthesis) or alternatively

biophysical changes in the membrane bilayer depending on osmotic status and/or nitrogen

availability. In this manner the availability of glutamine and resulting cellular changes could

also be sensed by YbbR and transduced to DacA (Fig. 8).

Taken together, with this work, we provide a further connection between the c-di-

AMP signalling network and osmotic regulation in *S. aureus* but also to central nitrogen

metabolism. It will be interesting to determine in future studies the mechanistic bases for the

observed changes in cellular c-di-AMP levels depending on glutamine uptake and the

involvement of GlmM and YbbR in this process.

Experimental Procedures

Bacterial strains and culture conditions

Bacterial strains used in this study are listed in Table 1. *S. aureus* strains were grown in

Tryptic Soy Broth (TSB), Tryptic Soy Agar (TSA) or Chemically Defined Medium (CDM).

CDM was prepared as described previously (Zeden *et al.*, 2018). Where indicated, certain

amino acids were removed from the CDM recipe during uptake assays and when needed

the TSB was supplemented with 0.4 M NaCl. An ammonium free chemically defined

medium, CDM+Glu, was prepared the same way as CDM containing the standard glutamine

concentration of 100 mg/L but lacking ammonium sulphate and CDM+Gln was prepared the

same way as CDM+Glu but replacing the glutamate with 100 mg/ L glutamine. *Escherichia*

coli strains were grown in Lysogeny Broth (LB). Where appropriate, antibiotics and/or

inducers were added to the media at the following concentration: 200 ng/ml

anhydrotetracyline (Atet), 90 µg/ml Kanamycin (Kan), 10 µg/ml Erythromycin (Erm), 7.5 or

10 µg/ml Chloramphenicol (Cam), Ampicillin (Amp) 100 µg/ml.

Bacterial strain construction

All strains used in this study are listed in Table 1 and primers used in this study are listed in

Table 2. The transposon insertion sites in the Nebraska transposon mutant library (NTML)

strains (Fey *et al.*, 2013) used as part of this study were confirmed by PCR and sequencing.

The transposon and surrounding region were also moved by phage transduction using

phage 85 into the *S. aureus* LAC* strain background. This resulted in the generation of *S.*

aureus strains LAC*alsT::tn (*SAUSA300_1252::tn*; ANG4803), LAC*0914::tn

(*SAUSA300_0914::tn*; ANG5141), LAC**glnQ::tn* (*SAUSA300_0914::tn;* ANG5070),

LAC**gltT::tn* (*SAUSA300_1252::tn*; ANG5366) and LAC**gltS::tn* (*SAUSA300_2291::tn*;

ANG5367). The transposon insertion in the respective gene was again confirmed by PCR

and sequencing. For complementation analysis, the Atet inducible single copy integration

plasmids piTET-*alsT* and piTET-*gltS* were constructed. To this end, *alsT* (*SAUSA300_1252*)

and *gltS* (*SAUSA300_2291*) were amplified using LAC* chromosomal DNA and primers

ANG2250/ANG2251 and ANG3209/ANG3210, respectively. The products as well as piTET

were digested with AvrII and SacII and then ligated. Plasmid piTET-*alsT* was recovered in *E.*

coli strain XL1-Blue (yielding strain ANG3937), shuttled through *E. coli* strain IM08B (strain

ANG3955) and then introduced into LAC**alsT::tn* (ANG4803), yielding strain LAC**alsT::tn*

piTET-*alsT* (ANG4854). As a control, plasmid piTET was also introduced into LAC**alsT::tn*

(ANG4803) yielding strain LAC**alsT::tn* piTET (ANG4853). Plasmid piTET-*gltS* was

transformed into *E. coli* XL1-Blue (yielding strain ANG5494), shuttled through *E. coli* IM08B

(yielding strain ANG5495) and transformed into *LAC* gltS::tn*, yielding the complement strain

LAC* *gltS::tn* piTET-*gltS* (ANG5493). As a control, the piTET plasmid was transformed into

LAC* *gltS*:tn strain, yielding the strain LAC* *gltS*:tn piTET (ANG5492). Correct plasmid

integration into the *geh* locus was confirmed by PCR and the sequences of all plasmid

inserts were confirmed by fluorescent automated sequencing.

Bacterial growth curves and amino acid analysis in culture supernatants

S. aureus strains LAC* and LAC**alsT::tn* were grown overnight in TSB supplemented with

10 µg/ml erythromycin where appropriate. Overnight cultures were then diluted to an OD_{600}

of 0.01 into 50 ml of fresh TSB. Cultures were incubated at 37°C with aeration, and OD_{600}

values determined every hour. The experiment was performed with three biological

replicates and the average OD_{600} values and standard deviations were plotted. Using the

same cultures, supernatant samples were prepared at time 0, 6, 10 and 12 h after growth

and amino acid levels determined as previously described using an amino acid analyser

(Halsey *et al.*, 2017). For measuring the growth of *S. aureus* strains LAC*, LAC* piTET,

LAC*_alsT::tn_, LAC*_alsT::tn_ piTET, LAC*_alsT::tn_ piTET-_alsT,_ LAC*_gltS::tn_ in CDM+Gln and

CDM+Glu, the bacteria were grown overnight in TSB medium supplemented with

chloramphenicol and erythromycin where appropriate. Next day, bacteria from a 1 ml aliquot

were washed twice in PBS and diluted to OD_{600} of 0.005 in either CDM-Glu or CDM+Gln

supplemented with 200 ng/ml Atet were indicated. One hundred µl of the diluted cultures (6

technical replicates) were transferred into well of a 96-well plate and the plate was then

incubated with shaking (500 rpm) in a plate reader and OD_{600} measured every 30 min. The

experiment was performed with three biological replicates and one representative graph is

shown.

Microscopic analysis and cell size measurements

Microscopic analysis to determine bacterial cell sizes was performed essentially as

previously described (Zeden _et al._, 2018). Briefly, _S. aureus_ strains LAC*, LAC*_dacA::kan_,

LAC*_dacA_$_{G206S}$, LAC*_dacA/opuD_ (ANG3835) and LAC*_dacA/alsT_ (ANG3838) were grown

overnight at 37°C in TSB or TSB supplemented with 0.4 M NaCl where stated. Next day, the

cultures were diluted to an OD_{600} of 0.01 and grown for 3 hours at 37°C. 100 µl of these

cultures were then stained for 20 min at 37°C with Vancomycin-BODIPY FL at a final

concentration of 2 µg/ml. 1.5 µl of each sample was spotted onto a thin 1.5% agarose gel

patch prepared in H_2O or in 0.4 M NaCl and the bacteria subsequently imaged at 1000 x

magnification using an Axio Imager A2 Zeiss microscope equipped with a GFP filter set.

Images were acquired using the ZEN 2012 (blue edition) software. The bacterial cell

diameters were determined using the Fiji software. Only non-dividing cells (cells without any

obvious fluorescent dots or lines at the mid-cell), were used for cell diameter measurements.

The cell diameters of 50 cells were measured and the average cell diameter determined.

The experiment was conducted with three or four biological replicates (as indicated in the

figure legend) and the averages and standard deviations of the average cell diameters

plotted.

Uptake assays using [14]C-labelled amino acids

Uptake assays were conducted as previously described with some minor modifications (Zeden *et al.*, 2018). Briefly, *S. aureus* strains were streaked on TSA or TSA 0. 4M NaCl plates with appropriate antibiotics and the plates incubated overnight at 37°C. Bacteria were subsequently scraped off from the plates and suspended in 1 ml PBS pH 7.4 buffer and the OD_{600} determined. Fifty ml of CDM (where indicated with 200 ng/ml of the inducer Atet added) was inoculated with the appropriate bacterial suspensions to an OD_{600} of 0.05. The cultures were grown at 37°C to an OD_{600} between 0.4 and 0.9 and then bacteria from an OD_{600} equivalent of 8 were harvested by centrifugation for 10 min at 19,000 x g at RT. Supernatants were discarded and the bacterial pellets were suspended in 2 ml of CDM without glutamate (for glutamine and glutamate uptake assays), CDM without serine (for serine uptake assays). The OD_{600} of the cell suspensions were measured and the cells diluted to an OD_{600} of approximately 1. The OD_{600} was re-measured and this measurement used for normalization purposes. Five hundred and fifty µl of these cell suspensions were aliquoted into 50 ml conical tubes and 100 µl used to measure the background radiation, by filtering the cells onto a nitrocellulose membrane filter, followed by a wash step with 16 ml PBS. Then, 6.2 µl of Glutamine, L-[14C(U)] (Hartmann Analytic, MC1124), Glutamic acid, L-[14C(U)] (Hartmann Analytic, MC156), or serine L-[14C(U)] (Hartmann Analytic, MC265) was added to the remaining 450 µl sample. Hundred µl aliquots were filtered at 0, 3, 6 and 9 minutes and the filters were then washed with 2 x 16 ml of PBS pH 7.4. The filters were subsequently dissolved in 9 ml of scintillation cocktail Filter Count (Perkin Elmer) and the radioactivity measured in counts per minute (CPM) using a Wallac 1409 DSA liquid scintillation counter. CPM for each sample were then normalized by the OD_{600} of the final cell suspension and the means and standard deviations of the CPM/ml OD_{600} = 1 of three or four (as indicated in the figure legends) independent experiments were plotted.

Determination of cellular c-di-AMP levels by competitive ELISA.

Intracellular c-di-AMP levels in WT LAC* and the indicated *S. aureus* mutant strains were

determined using a previously described competitive ELISA method (Underwood *et al.*,

2014) and a slightly modified method for the preparation of *S. aureus* samples (Bowman *et*

al., 2016). Briefly, a single colony of the WT and different *S. aureus* mutant strains were

picked from TSA plates and used to inoculate 5 ml of either CDM+Gln or CDM+Glu and the

cultures were incubated for 18 h at 37°C with shaking. Next, bacteria from 4.5 ml culture

were collected by centrifugation, washed three times with PBS and subsequently suspended

in 0.75 to 1 ml 50 mM Tris pH 8 buffer supplemented with 20 ng/ml lysostaphin and the cells

were lysed by bead beating. The lysates were cleared by centrifugation for 5 min at 17,000 x

g and the supernatant transferred to a new tube. A small sample aliquot was removed, and

the protein concentration determined using a Pierce BCA protein assay kit for normalization

purposes (Thermo Sientific, Waltham, MA, USA). The remainder of the sample was heated

to 95°C for 10 min. For the ELISA assay, the samples were diluted to a protein concentration

of 100, 200, 400 or 500 µg/ml as, appropriate. ELISA plates were prepared by adding 100 µl

of coating buffer (50 mM Na_2CO_3, 50 mM $NaHCO_3$, pH 9.6) containing 10 µg/ml of the c-di-

AMP binding $CpaA_{SP}$ protein to each well of a NUNC MaxiSorp 96 well plate (Thermo

Sientific, Waltham, MA, USA) and the plate was incubated for approximately 18 h at 4°C.

Next, the plate was washed three times with 200 µl PBST pH 7.4 (10 mM Na_2HPO_4, 1.8 mM

KH_2PO_4 137 mM NaCl , 2.7 mM KCl , 0.05% (v/v) Tween 20), blocked for 1 h at 18°C with

150 µl blocking solution (1% BSA in PBS pH 7.4) and washed three times with 200 µl PBST.

Fifty µl of the samples (three biological replicates and three technical replicates) or

standards (two technical replicates) were mixed with 50 µl of a 50 nM biotinylated c-di-AMP

solution prepared in 50 mM Tris pH 8 buffer. For the standard curve, c-di-AMP standards

were prepared in 50 mM Tris pH 8 buffer at concentrations of 0, 12.5, 25, 37.5, 50, 75, 100,

200 nM. Following the addition of the samples and the standards, the plate was incubated

for 2 h at 18°C and then washed three times with PBST. Next, 100 µl of a high sensitivity

streptavidin-HRP solution (Thermo Scientific, Waltham, MA, USA) diluted 1:500 in PBS was

added to each well and the plate was incubated for 1 h at 18°C. The plate was washed

again 3 x with 200 µl PBST and 100 µl of a developing solution (0.103 M NaHPO$_4$, 0.0485 M

citric acid, 500 mg/l o-phenylenediamine dihydrochloride, 0.03% H$_2$O$_2$) was added to each

well and the plate incubated for 15 min at 18°C. The reaction was then stopped by adding

100 µl of 2 M H$_2$SO$_4$ solution. The absorbance was measured in a plate reader at a

wavelength of 490 nm and c-di-AMP concentrations were calculated as ng c-di-AMP / mg

protein.

Acknowledgments: This work was funded by the Wellcome Trust grants 100289/Z/12/Z

and 210671/Z/18/Z to AG and the NIH/NIAID grants P01AI083211 and R01AI125588 to

P.D.F. MSZ was supported by a Medical Research Council Centre for Molecular

Bacteriology and Infection (MRC CMBI) studentship. The funders had no role in the study

design, data collection and interpretation, or the decision to submit the work for publication.

Author contribution: MSZ, IK and AG design the study, MSZ and IK acquired the data,

MSZ, IK, VTC, PF and AG, designed experiments, analyzed and interpreted the data, MSZ,

IK and AG prepared the figures and writing of the manuscript. All authors approved the final

version of the manuscript.

Graphical abstract:
652

653
654
655

Abbreviated Summary:

A large number of amino acid transporters and oligopeptide permeases are encoded in

bacterial genomes. However, their actual substrate specificity and functions are hard to

predict bioinformatically. In this study, we report that GltS and AlsT are main glutamate and

glutamine transporters in *Staphylococcus aureus*, respectively and show that glutamine

uptake inhibits the production of the nucleotide signalling molecule c-di-AMP.

Table 1: Bacterial strains used in this study

Unique ID	Strain name and resistance	Source
***Escherichia coli* strains**		
ANG284	XL1-Blue piTET; AmpR	(Gründling & Schneewind, 2007)
ANG2154	DH10B pIMAY; CamR	(Monk *et al.*, 2012)
ANG3724	IM08B	(Monk *et al.*, 2015)
ANG3928	IM08B piTET; AmpR	(Zeden *et al.*, 2018)
ANG3937	XL1-Blue piTET-*alsT*; AmpR	This study
ANG3955	IM08B piTET-*alsT*; AmpR	This study
ANG5494	XL1-Blue piTET-*gltS*; AmpR	This study
ANG5495	IM08B piTET-*gltS;* AmpR	This study
***Staphylococcus aureus* strains**		
AH1263	LAC* Erm sensitive CA-MRSA USA300 strain (ANG1575)	(Boles *et al.*, 2010)
ANG1961	LAC**gdpP::kan;* KanR	(Corrigan *et al.*, 2011)
ANG3664	LAC**dacA*~G206S~; KanR	(Bowman *et al.*, 2016)
ANG3666	LAC**dacA::kan* (*dacA*) KanR	(Zeden *et al.*, 2018)
ANG3835	LAC**dacA::kan*-S7 (LAC**dacA/opuD*); KanR	(Zeden *et al.*, 2018)
ANG3838	LAC**dacA::kan*-S10 (LAC**dacA/alsT*)]; KanR	(Zeden *et al.*, 2018)
ANG3940	NE142 (*alsT::tn*) – NMTN strain	(Fey *et al.*, 2013)
ANG4054	LAC* piTET; CamR	(Zeden *et al.*, 2018)
ANG4803	LAC**alsT::tn*; ErmR	This study
ANG4854	LAC**alsT::tn* piTET-*alsT* ; ErmR CamR	This study
ANG4853	LAC**alsT::tn* piTE ; ErmR CamR	This study
ANG4968	NE1463 (JE2 SAUSA300_*0914::tn*) – NMTN strain	(Fey *et al.*, 2013)
ANG5070	NE153 (JE2 *glnQ::tn*) – NMTN strain	(Fey *et al.*, 2013)
ANG5141	LAC**0914::tn*; ErmR	This study
ANG5242	LAC**glnQ::tn;* ErmR	This study
ANG5309	NE566 (JE2 *gltT::tn*) – NMTN strain	(Fey *et al.*, 2013)
ANG5310	NE560 (JE2 *gltS::tn*) – NMTN strain	(Fey *et al.*, 2013)
ANG5366	LAC**gltT::tn;* ErmR	This study
ANG5367	LAC**gltS::tn;* ErmR	This study
ANG5492	LAC**gltS::tn* piTET; ErmR CamR	This study
ANG5493	LAC**gltS::tn* piTET-*gltS*; ErmR CamR	This study

666   **Table 2:** Cloning primers used in this study

Primer ID	Name	Sequence
ANG2250	5-AvrII-*alsT*	AGTCCCTAGGCGGTCTAATTTTATAGAAGG
ANG2251	3-SacII-*alsT*	TCCCCGCGGGGTTTATTTGATTTTTATATAATGAATCG
ANG3209	5-AvrII-*gltS*	ATACCTAGGAGGGAGAGGGATATTCAACAAGGGGGATTTG
ANG3210	3-SacII-*gltS*	GCCCGCGGTTTAACTAAACCATTGTATGAATCCCATAATG

Figures and Legends

FIG 1: Inactivation of the glycine betaine transporter OpuD rescues the cell size of *S. aureus dacA* mutant bacteria. (A, C, E) Microscopy images of *S. aureus* cells stained with fluorescently labelled vancomycin. Cultures of *S. aureus* LAC* (WT), LAC*dacA$_{G206S}$ (*dacA*$_{G206S}$) (panels A and C only), LAC**dacA::kan* (*dacA*) (panel E only) and the suppressor strains LAC**dacA/alsT* (*dacA/alsT*) and LAC**dacA/opuD* (*dacA/opuD*) were grown in (A) TSB or (C and E) TSB 0.4 M NaCl medium and subsequently stained with fluorescently labelled vancomycin. The bacteria were then viewed using a fluorescent microscope and representative images are shown. Scale bars are 1 µm. (B, D, F) Bacterial cell diameter measurements. The diameter of non-dividing bacterial cells was measured as described in the Materials and Method section for *S. aureus* strains grown in (B) TSB or grown in (D and F) TSB 0.4 M NaCl medium. The diameters of 50 cells were determined and the average diameter calculated. The experiment was performed in triplicate (B and F) or quadruplicate (D) and the averages and SD of the average cell diameters plotted. For statistical analysis, one-way ANOVA followed by Dunnett's multiple comparison tests were performed (ns = not significant, * = $p<0.01$, ** = $p<0.001$, *** = $p<0.0001$).

FIG 2: Amino acid uptake analysis to determine the function of AlsT. (A) Bacterial

growth curves. *S. aureus* strains LAC* (WT) and LAC*alsT::tn (*alsT::tn*) were grown in TSB

medium and OD_{600} readings determined at hourly intervals and the average and standard

deviations from three biological replicates plotted. (B and C) Quantification of amino acid

levels in culture supernatants. Spent medium samples from the cultures shown in panel A

were prepared at the 0, 6, 10 and 12 h time points and (B) aspartate and (C) serine levels

determined as previously described using an amino acid analyzer (Halsey *et al.*, 2017). The

average values and standard deviations from 3 biological replicates were plotted. The plots

for all other amino acids measured are shown in Figure S1.

FIG 3: AlsT is a main glutamine transporter in *S. aureus*. Amino acid uptake assays. (A-

C) *S. aureus* strain LAC* (WT), the *alsT* mutant LAC**alsT::tn* piTET (*alsT::tn* piTET) and the

complementation strain LAC**alsT::tn* piTET-*alsT* (*alsT::tn* piTET-*alsT*) were grown to mid-log

phase in CDM supplemented with 200 ng/ml Atet for the strains containing the piTET

plasmids. Subsequently radiolabelled (A) serine, (B) glutamate or (C) glutamine was added

to culture aliquots, samples removed and filtered at the indicated time points and the

radioactivity accumulated in the cells measured. The average values and standard

deviations from three (A,C,D,F) or four (B,E) experiments were plotted. (D-F) The same

uptake assay experiment was performed as described in (A-C) but using *S. aureus* strains

LAC**dacA::kan* (*dacA*) and LAC**dacA/alsT* (*dacA/alsT*). The amino acid uptake curve for the

LAC* (WT) strain is the same as shown in panels A-C, as all strains were grown and

processed at the same time.

711
712

FIG 4: LAC*_0914::tn_ and LAC*_glnQ::tn_ strains do not show a defect in glutamine or

glutamate uptake. (A and B) Amino acid uptake assays. _S. aureus_ strains LAC* (WT) and

LAC*_0914::tn_ were grown to mid-log phase in CDM. Subsequently radiolabelled (A)

glutamine and (B) glutamate was added to culture aliquots, samples removed and filtered at

the indicated time points and the radioactivity accumulated in the cells measured. The

average values and standard deviations from three experiments were plotted. (C and D)

Amino acid uptake assays. Amino acid uptake assays were performed and the data plotted

as described in panels A and B, using _S. aureus_ LAC* (WT) and LAC*_glnQ::tn_ (_glnQ::tn_).

FIG 5: GltS is a main glutamate transporter in *S. aureus*. Amino acid uptake assays. (A

and B) *S. aureus* strains LAC* (WT), LAC*gltT::tn and LAC*gltS::tn were grown to mid-log

phase in CDM. Subsequently radiolabelled (A) glutamine or (B) glutamate was added to

culture aliquots, samples removed and filtered at the indicated time points and the

radioactivity accumulated in the cells measured. (C) Same as (B) but using *S. aureus* strains

LAC* (WT), LAC*gltS::tn piTET and the complementation strain LAC*gltS::tn piTET-gltS and

supplementing the CDM medium with 200 ng/μl Atet. The average values and standard

deviations from three experiments were plotted.

Figure 6. Addition of glutamine but not glutamate can stimulate the growth of *S.*

***aureus* in CDM lacking ammonium as nitrogen source. (A)** Growth curves of *S. aureus*

strains LAC* (WT), LAC*gltS::tn (*gltS::tn*) and LAC*alsT::tn (*alsT::tn*). The strains were

grown in CDM medium lacking ammonium and containing either glutamine (CDM+Gln) or

glutamate (CDM+Glu). OD_{600} readings were measured every 30 min. Average and standard

deviations of six technical replicates were plotted. This experiment is a representative result

of three independent experiments. **(B)** Same as (A) but using the *S. aureus* strains LAC*

(WT), LAC*alsT::tn piTET and the complementation strain LAC*alsT::tn piTET-alsT grown in

CDM+Gln medium supplemented with 200 ng/µl Atet.

Figure 7. Glutamine uptake inhibits c-di-AMP production in *S. aureus*. (A) *S. aureus* strains LAC* (WT) and LAC* *gdpP::kan* (*gdpP::kan*) were grown in CDM+Gln or CDM+Glu. Cell extracts were prepared, and cellular c-di-AMP levels determined using a competitive ELISA assay. The average values and standard deviation from three biological replicates were determined and c-di-AMP levels plotted as ng c-di-AMP/ mg protein. For statistical analysis f-tests and Student's t-tests were performed and c-di-AMP levels were found to be statistically significantly different for both WT and the *gdpP* mutant, when grown in the different media (* = p<0.01 and ** p<0.001). **(B)** *S. aureus* strains LAC* (WT), LAC**gltS::tn* (*gltS::tn*) and LAC**alsT::tn* (*alsT::tn*) were grown in CDM+Glu (white columns) or CDM+Gln (grey columns) and c-di-AMP concentrations determined and plotted as described in (A). **(C)** *S. aureus* strains LAC* piTET, LAC**alsT::tn* piTET and the complementation strain LAC**alsT::tn* piTET-*alsT* were grown in CDM+Gln supplemented with 200 ng/µl Atet and c-di-AMP levels determined as described in (A). For panels B and C, one-way ANOVA followed by Dunnett's multiple comparison tests were performed to identify statistically significant differences in c-di-AMP levels between WT and the mutant strains (ns = not significant, *** = p<0.0001).

Figure 8. Model of how AlsT-mediated glutamine uptake impacts bacterial physiology and c-di-AMP production. As shown as part of this study, the proteins AlsT (SAUSA300_1252) and GltS (SAUSA300_2291) are the main glutamine (Gln) and glutamate (Glu) transporters in *S. aureus*, respectively. A high glutamine/glutamate ratio signals nitrogen availability and hence glutamine uptake is thought to stimulate the bacterial metabolism. Glutamine is further converted by the GOGAT enzymes GltB and GltD to glutamate, the cellular counterion of potassium. Glutamine is also an important precursor for the synthesis of the essential peptidoglycan precursor UDP-GlcNAc. Glutamine and fructose-6-P (F6P) are converted by GlmS to glucosamine-6-P (GlcN6P), which is converted by GlmM to glycosamine-1-P (GlcN1P) and then utilized by GlmU for the production of UDP-GlcNAc. GlmM has been shown to directly interact and inhibit the activity of the c-di-AMP cyclase DacA. Based on the data presented in this study, glutamine uptake could potentially stimulate the GlmM/DacA interaction resulting in the observed reduction in c-di-AMP production. Alternatively, a potential increased flow of glutamine into peptidoglycan precursor and actual peptidoglycan synthesis could be sensed by the proposed regulator protein YbbR. Once such a signal is perceived by YbbR, it will be transduced to the c-di-AMP cyclase DacA leading to reduced c-di-AMP production. Reactions predicted to increase upon glutamine update are indicated by green arrows and the reduced c-di-AMP synthesis (potentially mediated by interaction with GlmM or YbbR) is indicated by a red arrow.

FIG S1: Amino acid uptake analysis to determine the function of AlsT. (A-O) Quantification of amino acid levels in culture supernatants. Spent medium samples from the cultures shown in Fig. 2 panel A were prepared at the 0, 6, 10 and 12 h time points and amino acid levels as indicated above each panel determined as previously described using an amino acid analyzer (Halsey *et al.*, 2017). The average values and standard deviations from 3 independent biological replicates were plotted.

FIG S2: CLUSTAL Omega alignment of *B. subtilis* and *S. aureus* AlsT/GlnT homologs.

The *S. aureus* AlsT (SAUSA300_1252) proteins was used as query sequence in a BLAST

search to identity homologous proteins encoded in *S. aureus* FPR3757 and *B. subtilis* 168.

One protein SAUSA300_0914 was identified in *S. aureus* and four close homologues AlsT$_{BS}$,

GlnT$_{BS}$, YrbD$_{BS}$ (e-value: 353 e-117) and YflA$_{BS}$ (e-value: 237 2e-72) were found in *B.*

subtilis. A Clustal-Omaga alignment was performed with the six proteins and is shown.

Using BOXSHADE, identical residues were boxed dark blue and similar residues light blue.

Figure S3. Glutamine uptake inhibits c-di-AMP production in *S. aureus,* but no clear

differences are seen upon glutamate uptake. (A and B) *S. aureus* strains LAC* (WT),

LAC**gltS::tn* (*gltS::tn*) and LAC**alsT::tn* (*alsT::tn*) were grown in CDM+Glu (white columns)

or CDM+Gln (grey columns) and c-di-AMP concentrations determined and plotted as

described in Figure 7. The data shown in panels A and B are two additional experimental

replicates of the data shown in Fig. 7B. One-way ANOVA tests followed by a Dunnett's

multiple comparison tests were performed to identify statistically significant differences in c-

di-AMP levels between WT and the mutant strains (ns = not significant; * = p<0.01; ** =

p<0.001; *** = p<0.0001).

References

References

Anderson, C.B., and Witter, L.D. (1982) Glutamine and proline accumulation by
Staphylococcus aureus with reduction in water activity. *Appl Environ Microbiol* **43**:
1501-1503.
Bai, Y., Yang, J., Zarrella, T.M., Zhang, Y., Metzger, D.W., and Bai, G. (2014) Cyclic di-AMP
impairs potassium uptake mediated by a cyclic di-AMP binding protein in
Streptococcus pneumoniae. *J Bacteriol* **196**: 614-623.
Boles, B.R., Thoendel, M., Roth, A.J., and Horswill, A.R. (2010) Identification of genes
involved in polysaccharide-independent *Staphylococcus aureus* biofilm formation.
PLoS One **5**: e10146.
Bowman, L., Zeden, M.S., Schuster, C.F., Kaever, V., and Gründling, A. (2016) New Insights
into the Cyclic Di-adenosine Monophosphate (c-di-AMP) Degradation Pathway and
the Requirement of the Cyclic Dinucleotide for Acid Stress Resistance in
Staphylococcus aureus. *J Biol Chem* **291**: 26970-26986.
Chin, K.H., Liang, J.M., Yang, J.G., Shih, M.S., Tu, Z.L., Wang, Y.C., Sun, X.H., Hu, N.J.,
Liang, Z.X., Dow, J.M., Ryan, R.P., and Chou, S.H. (2015) Structural Insights into the
Distinct Binding Mode of Cyclic Di-AMP with SaCpaA_RCK. *Biochemistry* **54**: 4936-
4951.
Corrigan, R.M., Abbott, J.C., Burhenne, H., Kaever, V., and Gründling, A. (2011) c-di-AMP is
a new second messenger in *Staphylococcus aureus* with a role in controlling cell size
and envelope stress. *PLoS Pathog* **7**: e1002217.
Corrigan, R.M., Bowman, L., Willis, A.R., Kaever, V., and Gründling, A. (2015) Cross-talk
between two nucleotide-signaling pathways in *Staphylococcus aureus*. *J Biol Chem*
290: 5826-5839.
Corrigan, R.M., Campeotto, I., Jeganathan, T., Roelofs, K.G., Lee, V.T., and Gründling, A.
(2013) Systematic identification of conserved bacterial c-di-AMP receptor proteins.
Proc Natl Acad Sci U S A **110**: 9084-9089.
Crooke, A.K., Fuller, J.R., Obrist, M.W., Tomkovich, S.E., Vitko, N.P., and Richardson, A.R.
(2013) CcpA-independent glucose regulation of lactate dehydrogenase 1 in
Staphylococcus aureus. *PLoS One* **8**: e54293.
DeFrancesco, A.S., Masloboeva, N., Syed, A.K., DeLoughery, A., Bradshaw, N., Li, G.W.,
Gilmore, M.S., Walker, S., and Losick, R. (2017) Genome-wide screen for genes
involved in eDNA release during biofilm formation by *Staphylococcus aureus*. *Proc*
Natl Acad Sci U S A **114**: E5969-E5978.
Deguchi, Y., Yamato, I., and Anraku, Y. (1990) Nucleotide sequence of *gltS*, the
Na+/glutamate symport carrier gene of *Escherichia coli B*. *J Biol Chem* **265**: 21704-
21708.
Devaux, L., Sleiman, D., Mazzuoli, M.V., Gominet, M., Lanotte, P., Trieu-Cuot, P., Kaminski,
P.A., and Firon, A. (2018) Cyclic di-AMP regulation of osmotic homeostasis is
essential in Group B *Streptococcus*. *PLoS Genet* **14**: e1007342.
Fahmi, T., Faozia, S., Port, G., and Cho, K.H. (2019) The Second Messenger c-di-AMP
Regulates Diverse Cellular Pathways Involved in Stress Response, Biofilm
Formation, Cell Wall Homeostasis, SpeB Expression and Virulence in *Streptococcus*
pyogenes. *Infect Immun*.
Fey, P.D., Endres, J.L., Yajjala, V.K., Widhelm, T.J., Boissy, R.J., Bose, J.L., and Bayles,
K.W. (2013) A genetic resource for rapid and comprehensive phenotype screening of
nonessential *Staphylococcus aureus* genes. *MBio* **4**: e00537-00512.
Fisher, S.H. (1999) Regulation of nitrogen metabolism in *Bacillus subtilis*: vive la difference!
Mol Microbiol **32**: 223-232.
Forchhammer, K. (2007) Glutamine signalling in bacteria. *Front Biosci-Landmrk* **12**: 358-
370.
Fridkin, S.K., Hageman, J.C., Morrison, M., Sanza, L.T., Como-Sabetti, K., Jernigan, J.A.,
Harriman, K., Harrison, L.H., Lynfield, R., Farley, M.M., and Active Bacterial Core
Surveillance Program of the Emerging Infections Program, N. (2005) Methicillin-

resistant *Staphylococcus aureus* disease in three communities. *N Engl J Med* **352**:
1436-1444.

Fuller, J.R., Vitko, N.P., Perkowski, E.F., Scott, E., Khatri, D., Spontak, J.S., Thurlow, L.R.,
and Richardson, A.R. (2011) Identification of a lactate-quinone oxidoreductase in
Staphylococcus aureus that is essential for virulence. *Front Cell Infect Microbiol* **1**:
19.

Gründling, A., and Schneewind, O. (2007) Genes required for glycolipid synthesis and
lipoteichoic acid anchoring in *Staphylococcus aureus*. *J Bacteriol* **189**: 2521-2530.

Gundlach, J., Commichau, F.M., and Stülke, J. (2017a) Perspective of ions and
messengers: an intricate link between potassium, glutamate, and cyclic di-AMP. *Curr*
Genet.

Gundlach, J., Dickmanns, A., Schröder-Tittmann, K., Neumann, P., Kaesler, J., Kampf, J.,
Herzberg, C., Hammer, E., Schwede, F., Kaever, V., Tittmann, K., Stülke, J., and
Ficner, R. (2015a) Identification, characterization, and structure analysis of the cyclic
di-AMP-binding PII-like signal transduction protein DarA. *J Biol Chem* **290**: 3069-
3080.

Gundlach, J., Herzberg, C., Hertel, D., Thurmer, A., Daniel, R., Link, H., and Stülke, J.
(2017b) Adaptation of *Bacillus subtilis* to Life at Extreme Potassium Limitation. *MBio*
8.

Gundlach, J., Herzberg, C., Kaever, V., Gunka, K., Hoffmann, T., Weiss, M., Gibhardt, J.,
Thurmer, A., Hertel, D., Daniel, R., Bremer, E., Commichau, F.M., and Stülke, J.
(2017c) Control of potassium homeostasis is an essential function of the second
messenger cyclic di-AMP in *Bacillus subtilis*. *Sci Signal* **10**.

Gundlach, J., Krüger, L., Herzberg, C., Turdiev, A., Poehlein, A., Tascon, I., Weiß, M.,
Hertel, D., Daniel, R., Hänelt, I., Lee, V.T., and Stülke, J. (2019) Sustained sensing in
potassium homeostasis: Cyclic di-AMP controls potassium uptake by KimA at the
levels of expression and activity. *J Biol Chem* **294**: 9605-9614.

Gundlach, J., Mehne, F.M., Herzberg, C., Kampf, J., Valerius, O., Kaever, V., and Stülke, J.
(2015b) An Essential Poison: Synthesis and Degradation of Cyclic Di-AMP in
Bacillus subtilis. *J Bacteriol* **197**: 3265-3274.

Gunka, K., and Commichau, F.M. (2012) Control of glutamate homeostasis in *Bacillus*
subtilis: a complex interplay between ammonium assimilation, glutamate
biosynthesis and degradation. *Mol Microbiol* **85**: 213-224.

Halsey, C.R., Lei, S., Wax, J.K., Lehman, M.K., Nuxoll, A.S., Steinke, L., Sadykov, M.,
Powers, R., and Fey, P.D. (2017) Amino Acid Catabolism in *Staphylococcus aureus*
and the Function of Carbon Catabolite Repression. *MBio* **8**.

Hengge, R. (2009) Principles of c-di-GMP signalling in bacteria. *Nat Rev Microbiol* **7**: 263-
273.

Huynh, T.N., Choi, P.H., Sureka, K., Ledvina, H.E., Campillo, J., Tong, L., and Woodward,
J.J. (2016) Cyclic di-AMP targets the cystathionine beta-synthase domain of the
osmolyte transporter OpuC. *Mol Microbiol*.

Kelly, B., and O'Neill, L.A. (2015) Metabolic reprogramming in macrophages and dendritic
cells in innate immunity. *Cell Res* **25**: 771-784.

Kim, H., Youn, S.J., Kim, S.O., Ko, J., Lee, J.O., and Choi, B.S. (2015) Structural Studies of
Potassium Transport Protein KtrA Regulator of Conductance of K+ (RCK) C Domain
in Complex with Cyclic Diadenosine Monophosphate (c-di-AMP). *J Biol Chem* **290**:
16393-16402.

Kluytmans, J., van Belkum, A., and Verbrugh, H. (1997) Nasal carriage of *Staphylococcus*
aureus: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol*
Rev **10**: 505-520.

Lehman, M.K., Nuxoll, A.S., Yamada, K.J., Kielian, T., Carson, S.D., and Fey, P.D. (2019)
Protease-Mediated Growth of *Staphylococcus aureus* on Host Proteins Is *opp3*
Dependent. *MBio* **10**.

Mehne, F.M., Gunka, K., Eilers, H., Herzberg, C., Kaever, V., and Stülke, J. (2013) Cyclic di-
AMP homeostasis in *Bacillus subtilis*: both lack and high level accumulation of the
nucleotide are detrimental for cell growth. *J Biol Chem* **288**: 2004-2017.
Mirouze, N., Bidnenko, E., Noirot, P., and Auger, S. (2015) Genome-wide mapping of TnrA-
binding sites provides new insights into the TnrA regulon in *Bacillus subtilis*.
Microbiologyopen **4**: 423-435.
Monk, I.R., Shah, I.M., Xu, M., Tan, M.W., and Foster, T.J. (2012) Transforming the
untransformable: application of direct transformation to manipulate genetically
Staphylococcus aureus and *Staphylococcus epidermidis*. *MBio* **3**.
Monk, I.R., Tree, J.J., Howden, B.P., Stinear, T.P., and Foster, T.J. (2015) Complete Bypass
of Restriction Systems for Major *Staphylococcus aureus* Lineages. *MBio* **6**: e00308-
00315.
Moscoso, J.A., Schramke, H., Zhang, Y., Tosi, T., Dehbi, A., Jung, K., and Gründling, A.
(2015) Binding of Cyclic Di-AMP to the *Staphylococcus aureus* Sensor Kinase KdpD
Occurs via the Universal Stress Protein Domain and Downregulates the Expression
of the Kdp Potassium Transporter. *J Bacteriol* **198**: 98-110.
Pham, H.T., Nhiep, N.T.H., Vu, T.N.M., Huynh, T.N., Zhu, Y., Huynh, A.L.D., Chakrabortti,
A., Marcellin, E., Lo, R., Howard, C.B., Bansal, N., Woodward, J.J., Liang, Z.X., and
Turner, M.S. (2018) Enhanced uptake of potassium or glycine betaine or export of
cyclic-di-AMP restores osmoresistance in a high cyclic-di-AMP *Lactococcus lactis*
mutant. *PLoS Genet* **14**: e1007574.
Pham, H.T., and Turner, M.S. (2019) Onwards and [K(+)]upwards: a new potassium
importer under the spell of cyclic-di-AMP. *J Bacteriol*.
Pham, T.H., Liang, Z.X., Marcellin, E., and Turner, M.S. (2016) Replenishing the cyclic-di-
AMP pool: regulation of diadenylate cyclase activity in bacteria. *Curr Genet* **62**: 731-
738.
Quintana, I.M., Gibhardt, J., Turdiev, A., Hammer, E., Commichau, F.M., Lee, V.T., Magni,
C., and Stülke, J. (2019) The KupA and KupB proteins of *Lactococcus lactis* IL1403
are novel c-di-AMP receptor proteins responsible for potassium uptake. *J Bacteriol*.
Randazzo, P., Aucouturier, A., Delumeau, O., and Auger, S. (2017) Revisiting the in vivo
GlnR-binding sites at the genome scale in *Bacillus subtilis*. *BMC Res Notes* **10**: 422.
Richardson, A.R., Libby, S.J., and Fang, F.C. (2008) A nitric oxide-inducible lactate
dehydrogenase enables *Staphylococcus aureus* to resist innate immunity. *Science*
319: 1672-1676.
Rismondo, J., Gibhardt, J., Rosenberg, J., Kaever, V., Halbedel, S., and Commichau, F.M.
(2016) Phenotypes Associated with the Essential Diadenylate Cyclase CdaA and Its
Potential Regulator CdaR in the Human Pathogen Listeria monocytogenes. *J*
Bacteriol **198**: 416-426.
Rocha, R., Teixeira-Duarte, C.M., Jorge, J.M.P., and Morais-Cabral, J.H. (2019)
Characterization of the molecular properties of KtrC, a second RCK domain that
regulates a Ktr channel in *Bacillus subtilis*. *J Struct Biol* **205**: 34-43.
Römling, U. (2008) Great times for small molecules: c-di-AMP, a second messenger
candidate in Bacteria and Archaea. *Sci Signal* **1**: pe39.
Satomura, T., Shimura, D., Asai, K., Sadaie, Y., Hirooka, K., and Fujita, Y. (2005)
Enhancement of glutamine utilization in *Bacillus subtilis* through the GlnK-GlnL two-
component regulatory system. *J Bacteriol* **187**: 4813-4821.
Schuster, C.F., Bellows, L.E., Tosi, T., Campeotto, I., Corrigan, R.M., Freemont, P., and
Gründling, A. (2016) The second messenger c-di-AMP inhibits the osmolyte uptake
system OpuC in *Staphylococcus aureus*. *Sci Signal* **9**: ra81.
Schuurman-Wolters, G.K., and Poolman, B. (2005) Substrate specificity and ionic regulation
of GlnPQ from *Lactococcus lactis*. An ATP-binding cassette transporter with four
extracytoplasmic substrate-binding domains. *J Biol Chem* **280**: 23785-23790.
Spahich, N.A., Vitko, N.P., Thurlow, L.R., Temple, B., and Richardson, A.R. (2016)
Staphylococcus aureus lactate- and malate-quinone oxidoreductases contribute to
nitric oxide resistance and virulence. *Mol Microbiol* **100**: 759-773.

Sureka, K., Choi, P.H., Precit, M., Delince, M., Pensinger, D., Huynh, T.N., Jurado, A.R.,
Goo, Y.A., Sadilek, M., Iavarone, A.T., Sauer, J.D., Tong, L., and Woodward, J.J.
(2014) The cyclic dinucleotide c-di-AMP is an allosteric regulator of metabolic
enzyme function. *Cell* **158**: 1389-1401.
Teh, W.K., Dramsi, S., Tolker-Nielsen, T., Yang, L., and Givskov, M. (2019) Increased
Intracellular Cyclic di-AMP Levels Sensitize *Streptococcus gallolyticus* subsp.
gallolyticus to Osmotic Stress and Reduce Biofilm Formation and Adherence on
Intestinal Cells. *J Bacteriol* **201**.
Tolner, B., Ubbink-Kok, T., Poolman, B., and Konings, W.N. (1995) Characterization of the
proton/glutamate symport protein of *Bacillus subtilis* and its functional expression in
Escherichia coli. *J Bacteriol* **177**: 2863-2869.
Tosi, T., Hoshiga, F., Millership, C., Singh, R., Eldrid, C., Patin, D., Mengin-Lecreulx, D.,
Thalassinos, K., Freemont, P., and Gründling, A. (2019) Inhibition of the
Staphylococcus aureus c-di-AMP cyclase DacA by direct interaction with the
phosphoglucosamine mutase GlmM. *PLoS Pathog* **15**: e1007537.
Underwood, A.J., Zhang, Y., Metzger, D.W., and Bai, G. (2014) Detection of cyclic di-AMP
using a competitive ELISA with a unique pneumococcal cyclic di-AMP binding
protein. *J Microbiol Methods* **107**: 58-62.
Vitko, N.P., Spahich, N.A., and Richardson, A.R. (2015) Glycolytic dependency of high-level
nitric oxide resistance and virulence in *Staphylococcus aureus*. *MBio* **6**.
Whiteley, A.T., Garelis, N.E., Peterson, B.N., Choi, P.H., Tong, L., Woodward, J.J., and
Portnoy, D.A. (2017) c-di-AMP modulates *Listeria monocytogenes* central
metabolism to regulate growth, antibiotic resistance and osmoregulation. *Mol*
Microbiol **104**: 212-233.
Whiteley, A.T., Pollock, A.J., and Portnoy, D.A. (2015) The PAMP c-di-AMP Is Essential for
Listeria monocytogenes Growth in Rich but Not Minimal Media due to a Toxic
Increase in (p)ppGpp. *Cell Host Microbe* **17**: 788-798.
Wicke, D., Schulz, L.M., Lentes, S., Scholz, P., Poehlein, A., Gibhardt, J., Daniel, R.,
Ischebeck, T., and Commichau, F.M. (2019) Identification of the first glyphosate
transporter by genomic adaptation. *Environ Microbiol* **21**: 1287-1305.
Witte, C.E., Whiteley, A.T., Burke, T.P., Sauer, J.D., Portnoy, D.A., and Woodward, J.J.
(2013) Cyclic di-AMP is critical for *Listeria monocytogenes* growth, cell wall
homeostasis, and establishment of infection. *MBio* **4**: e00282-00213.
Woodward, J.J., Iavarone, A.T., and Portnoy, D.A. (2010) c-di-AMP secreted by intracellular
Listeria monocytogenes activates a host type I interferon response. *Science* **328**:
1703-1705.
Yoshida, K., Yamaguchi, H., Kinehara, M., Ohki, Y.H., Nakaura, Y., and Fujita, Y. (2003)
Identification of additional TnrA-regulated genes of *Bacillus subtilis* associated with a
TnrA box. *Mol Microbiol* **49**: 157-165.
Zarrella, T.M., Metzger, D.W., and Bai, G. (2018) Stress Suppressor Screening Leads to
Detection of Regulation of Cyclic di-AMP Homeostasis by a Trk Family Effector
Protein in *Streptococcus pneumoniae*. *J Bacteriol* **200**.
Zeden, M.S., Schuster, C.F., Bowman, L., Zhong, Q., Williams, H.D., and Gründling, A.
(2018) Cyclic di-adenosine monophosphate (c-di-AMP) is required for osmotic
regulation in *Staphylococcus aureus* but dispensable for viability in anaerobic
conditions. *J Biol Chem* **293**: 3180-3200.
Zhu, Y., Pham, T.H., Nhiep, T.H., Vu, N.M., Marcellin, E., Chakrabortti, A., Wang, Y.,
Waanders, J., Lo, R., Huston, W.M., Bansal, N., Nielsen, L.K., Liang, Z.X., and
Turner, M.S. (2016) Cyclic-di-AMP synthesis by the diadenylate cyclase CdaA is
modulated by the peptidoglycan biosynthesis enzyme GlmM in *Lactococcus lactis*.
Mol Microbiol **99**: 1015-1027.
Zhu, Y., Xiong, Y.Q., Sadykov, M.R., Fey, P.D., Lei, M.G., Lee, C.Y., Bayer, A.S., and
Somerville, G.A. (2009) Tricarboxylic acid cycle-dependent attenuation of
Staphylococcus aureus in vivo virulence by selective inhibition of amino acid
transport. *Infect Immun* **77**: 4256-4264.