Identification of the main glutamine and glutamate transporters in *Staphylococcus*

***aureus* and their impact on c-di-AMP production**

Merve S. Zeden[a,^], Igor Kviatkovski[a,^], Christopher F. Schuster[a], Vinai C. Thomas[b], Paul D.

Fey[b] and Angelika Gründling[a,#]

[a] Section of Molecular Microbiology and Medical Research Council Centre for Molecular

Bacteriology and Infection, Imperial College London, London SW7 2AZ.

[b] University of Nebraska Medical Center, Department of Pathology and Microbiology,

Omaha, Nebraska, USA.

^ These authors contributed equally

[#] To whom correspondence should be addressed: Angelika Gründling –

a.grundling@imperial.ac.uk

Keywords: *Staphylococcus aureus*, c-di-AMP, glutamine transporter, AlsT, glutamate

transporter, GltS, amino acid transport, OpuD, osmolyte transport

Running title: Regulation of c-di-AMP production by glutamine transport

Summary (200 words)

A *Staphylococcus aureus* strain deleted for the c-di-AMP cyclase gene *dacA* is unable to survive in rich medium unless it acquires compensatory mutations. Previously identified mutations were in *opuD*, encoding the main glycine-betaine transporter, and *alsT*, encoding a predicted amino acid transporter. Here, we show that inactivation of OpuD restores the cell size of a *dacA* mutant to near wild-type size, while inactivation of AlsT does not, suggesting two different mechanisms for the growth rescue. AlsT was identified as an efficient glutamine transporter, indicating that preventing glutamine uptake in rich medium rescues the growth of the *S. aureus dacA* mutant. In addition, GltS was identified as a glutamine transporter. By performing growth curves with WT, *alsT* and *gltS* mutant strains in defined medium supplemented with ammonium, glutamine or glutamate, we revealed that ammonium and glutamine, but not glutamate promote the growth of *S. aureus*. This suggests that besides ammonium also glutamine can serve as a nitrogen source under these conditions. Ammonium and uptake of glutamine via AlsT inhibited c-di-AMP production, while glutamate uptake had no effect. These findings provide, besides the previously reported link between potassium and osmolyte uptake, a connection between nitrogen metabolism and c-di-AMP signalling in *S. aureus*.

Introduction

In the human host, S*taphylococcus aureus* can grow in virous tissues such as kidneys, bones, heart, soft tissues and lungs (Kluytmans *et al.*, 1997, Fridkin *et al.*, 2005). Sensitive regulatory mechanisms enable this organism to rapidly respond to external stimuli and environmental changes. Amongst others, this allows bacteria to adapt their metabolism and utilize different carbon and nitrogen sources available in each specific niche (Fridkin *et al.*, 2005, Spahich *et al.*, 2016, Vitko *et al.*, 2015, Crooke *et al.*, 2013, Fuller *et al.*, 2011, Richardson *et al.*, 2008, Halsey *et al.*, 2017, Lehman *et al.*, 2019).

Glucose is the preferred carbon source for *S. aureus,* but it can be limiting during infection due to the host immune response (Kelly & O'Neill, 2015, Spahich *et al.*, 2016, Halsey *et al.*, 2017, Lehman *et al.*, 2019). In glucose-limiting conditions, *S. aureus* instead catabolizes secondary carbon sources and amino acids, particularly glutamate and proline, which serve as major carbon sources during growth in the absence of glucose (Halsey *et al.*, 2017). However, not much is known about amino acid uptake and catabolism in *S. aureus* and how the availability of certain nutrients can affect virulence factor expression and invasion of the host. While a large number of amino acid transporters and oligopeptide permeases can be identified bioinformatically, their actual substrate specificities and functions in *S. aureus* have not yet been studied in detail. Predicting the substrates for transporters bioinformatically remains challenging and hence such questions need to be addressed experimentally.

Secondary messenger molecules are crucial in allowing bacteria to rapidly adapt to different environmental and host cell niches (Römling, 2008, Hengge, 2009). There is now considerable evidence that one of these messengers, cyclic di-adenosine monophosphate (c-di-AMP) plays a key role in osmoregulation in bacteria (Pham *et al.*, 2018, Pham & Turner, 2019, Quintana *et al.*, 2019, Zarrella *et al.*, 2018, Teh *et al.*, 2019, Fahmi *et al.*, 2019, Devaux *et al.*, 2018, Bai *et al.*, 2014, Zeden *et al.*, 2018, Corrigan *et al.*, 2011, Rocha *et al.*, 2019, Gundlach *et al.*, 2017a, Gundlach *et al.*, 2018, Witte *et al.*, 2013, Whiteley *et al.*, 2015, Whiteley *et al.*, 2017). c-di-AMP binds to and negatively regulates a number of

different potassium and osmolyte importers (Rocha *et al.*, 2019, Quintana *et al.*, 2019, Kim

et al., 2015, Corrigan *et al.*, 2013, Moscoso *et al.*, 2015, Chin *et al.*, 2015, Huynh *et al.*,

2016, Schuster *et al.*, 2016, Pham & Turner, 2019, Pham *et al.*, 2018, Devaux *et al.*, 2018,

Zarrella *et al.*, 2018, Gundlach *et al.*, 2017a, Gundlach *et al.*, 2018, Gundlach *et al.*, 2017b).

In many Firmicutes, c-di-AMP is essential for bacterial growth under standard rich medium

growth conditions, but it is also toxic at high levels. Hence, the cellular levels must be tightly

regulated (Gundlach *et al.*, 2015, Mehne *et al.*, 2013, Corrigan *et al.*, 2011, Corrigan *et al.*,

2015, Woodward *et al.*, 2010, Witte *et al.*, 2013). In *S. aureus, Streptococcus agalactiae*

and *Listeria monocytogenes,* deletion of *dacA* (also referred to as *cdaA* in many bacteria),

coding for the diadenylate cyclase enzyme and responsible for the synthesis of c-di-AMP,

was only possible in chemically defined medium (Whiteley *et al.*, 2015, Zeden *et al.*, 2018,

Devaux *et al.*, 2018), whereas in *Bacillus subtilis* all three c-di-AMP cyclases could only be

inactivated in minimal medium also containing low amounts of potassium (Gundlach *et al.*,

2017a).

Previously, we found that inactivation of the main glycine betaine transporter OpuD

(SAUSA300_1245) as well as the predicted amino acid transporter AlsT (SAUSA300_1252)

allows an *S. aureus dacA* mutant to grow in rich medium in the absence of c-di-AMP (Zeden

et al., 2018). In several other Firmicutes, including *B. subtilis, Lactococcus lactis,*

Streptococcus pneumoniae, S. agalactiae, Streptococcus pyogenes and *L. monocytogenes*,

inactivating mutations have also been identified in osmolyte and potassium transport

systems that allow these bacteria to grow in the absence of c-di-AMP (Pham *et al.*, 2018,

Pham & Turner, 2019, Quintana *et al.*, 2019, Zarrella *et al.*, 2018, Teh *et al.*, 2019, Fahmi *et*

al., 2019, Devaux *et al.*, 2018, Bai *et al.*, 2014, Zeden *et al.*, 2018, Corrigan *et al.*, 2011,

Rocha *et al.*, 2019, Gundlach *et al.*, 2017a, Gundlach *et al.*, 2018, Witte *et al.*, 2013,

Whiteley *et al.*, 2015, Whiteley *et al.*, 2017). This suggests that potassium and osmolyte

transporters are more active in the absence of c-di-AMP, resulting in the accumulation of

toxic levels of potassium and osmolytes in the cell. Consistent with a key function of c-di-

AMP in regulating the osmotic balance in the cell, we found that *S. aureus* cells show

significant differences in cell size depending on their intracellular c-di-AMP levels (Zeden *et*

al., 2018, Corrigan *et al.*, 2011). Cells of strain LAC*gdpP, which have high c-di-AMP levels,

show a decrease in cell size, while cells of strain LAC*dacA$_{G206S}$, containing low levels of c-

di-AMP, show an increase in cell size (Zeden *et al.*, 2018, Corrigan *et al.*, 2011). As c-di-

AMP negatively regulates potassium and osmolyte uptake (Corrigan *et al.*, 2013, Moscoso

et al., 2015, Schuster *et al.*, 2016), the increase in cell size is consistent with the hypothesis

that an increase in potassium and osmolyte uptake and retention of water at reduced c-di-

AMP levels leads to the observed increase in cell size. c-di-AMP levels affecting bacterial

cell size has also been observed for other bacteria such as *S. pneumonia* and *L.*

monocytogenes (Commichau *et al.*, 2019, Bai *et al.*, 2013).

AlsT is a predicted amino acid transporter and a correlation between cellular levels of

c-di-AMP and the amino acids glutamate and glutamine has been reported for *B. subtilis* and

L. monocytogenes (Whiteley *et al.*, 2017, Gundlach *et al.*, 2015, Gundlach *et al.*, 2018,

Sureka *et al.*, 2014)*. A two-fold increase in cellular c-di-AMP levels was observed in *B.*

subtilis when bacteria where grown in Spizizen minimal medium with glutamate (Glu) as

compared to growth in the same medium with glutamine (Gln) as a nitrogen source

(Gundlach *et al.*, 2015). In *L. monocytogenes*, c-di-AMP was identified as a negative

regulator of the key TCA cycle enzyme pyruvate carboxylase (Sureka *et al.*, 2014)*.

Depletion of c-di-AMP resulted in an increased flux into the TCA cycle and as a

consequence an increase in the cellular glutamine/glutamate pool (Sureka *et al.*, 2014). In a

citZ mutant, which lacks the TCA cycle enzyme citrate synthase and thus has an early block

in the TCA cycle, the depletion of c-di-AMP no longer resulted in the accumulation of

glutamate/glutamine in the cell (Sureka *et al.*, 2014). Interestingly, an *L. monocytogenes*

dacA/*citZ* double mutant was again viable in rich medium (Sureka *et al.*, 2014, Whiteley *et*

al., 2017).

As part of this study, we further investigated why inactivation of the main glycine-

betaine transporter OpuD and the predicted amino acid transporter AlsT allows *S. aureus* to

grow in the absence of c-di-AMP in rich medium. We show that AlsT is a main glutamine

transporter in *S. aureus* and that AlsT-mediated glutamine uptake represses c-di-AMP

production. Similarly, growth in ammonium-containing defined medium but not in glutamate-

containing medium, repressed c-di-AMP production. The repression of c-di-AMP production

was independent of the activity of the c-di-AMP phosphodiesterase GdpP and the predicted

cyclase regulator YbbR. With this study, we not only provide a further link between the c-di-

AMP signalling network and osmotic regulation in bacterial cells but also with the uptake of

specific nitrogen sources and amino acids in *S. aureus.*

Results

Inactivation of OpuD but not AlsT reduces the cell size of an *S. aureus dacA* mutant

In previous work, we reported a correlation between the cell size and c-di-AMP levels in *S.*

aureus: bacteria with high c-di-AMP level are smaller, whereas bacteria with low c-di-AMP

levels (strain LAC*dacA_{G206S}) are larger as compared to wild-type bacteria (Zeden *et al.*,

2018, Corrigan *et al.*, 2011). We also reported that inactivating mutations in *opuD*

(*SAUSA300_1245*) coding for the main glycine betaine osmolyte transporter and *alsT*

(*SAUSA300_1252*) coding for a predicted amino acid sodium symporter, rescue the growth

defect observed for the c-di-AMP negative *S. aureus* strain LAC*dacA::kan* in rich medium

TSB (Zeden *et al.*, 2018). Here, we investigated further the mechanism by which the growth

defect of the *dacA* mutant strain is rescued in the LAC*dacA/opuD* and LAC*dacA/alsT*

suppressor strains. Initially, we compared the cell size of bacteria from the suppressor

strains LAC*dacA/opuD* and LAC*dacA/alsT* to that of WT LAC* and the low c-di-AMP level

strain LAC*dacA_{G206S} after growth in the rich medium TSB. As expected, the bacteria with

low-levels of c-di-AMP showed an increase in cell size as compared to WT bacteria (Fig. 1A

and B). While a similar increase in cell size was still observed for bacteria of strain

LAC*dacA/alsT*, the cell size of LAC*dacA/opuD* bacteria, while still increased as compared

to the wild type, was significantly smaller as compared to the low-level LAC*dacA_{G206S} strain

(Fig. 1A and B). Because regular TSB medium is not suitable for the growth of the c-di-AMP

null strain LAC*dacA::kan,* bacterial cell sizes were also determined following growth in TSB

medium supplemented with 0.4 M NaCl, which is permissive for the growth of the *dacA*

mutant (Fig. 1C-F). Similar to what was observed for the low c-di-AMP level *dacA*$_{G206S}$

mutant strain, the size of bacteria from the c-di-AMP null strain LAC**dacA::kan* was

significantly increased compared to WT bacteria. As observed before, the cell size was not

rescued for bacteria of the LAC**dacA/alsT* suppressor strain (Fig. 1C-F). On the other hand,

the size of LAC**dacA/opuD* bacteria was similar to that of WT bacteria (Fig. 1C-F). Taken

together, the observed differences in cell size indicate that the underlying molecular

mechanisms enabling the *opuD* and *alsT* mutant strains to survive in the absence of c-di-

AMP in rich medium might be different.

AlsT is a glutamine transporter in *S. aureus*

AlsT (SAUSA300_1252) is a predicted amino acid transporter and annotated in the InterPro

database (www.ebi.ac.uk/interpro) as alanine/sodium symporter. However, in a previous

study no difference in the uptake of radiolabelled alanine was detected between a WT and

the LAC**dacA/alsT* mutant strain (Zeden *et al.*, 2018), indicating that AlsT is not an alanine

transporter. To identify potential substrates for the *S. aureus* AlsT transporter, we initially

followed the depletion of different amino acids from the culture supernatant during the

growth of the WT LAC* strain and an isogenic *alsT* transposon mutant strain in TSB

medium, where both strains exhibit similar growth rates (Fig. S1A). Of note, using this

method, tryptophan uptake cannot be measured and it is also not possible to distinguish

between glutamine/glutamate or asparagine/aspartate utilization. While no significant

differences were observed for most amino acids (Fig. S1), a slight increase in the utilization

of aspartate/asparagine and a slight decrease in the uptake of serine was observed (Fig.

S1D and Fig. S1O), suggesting that AlsT could potentially be a serine transporter. To test

this, uptake assays were performed with radiolabelled serine using WT LAC*, the *alsT*

mutant strain LAC**alsT::tn* piTET as well as the complementation strain LAC**alsT::tn* piTET-

alsT. However, no significant differences in the uptake rate of serine were observed between

the strains (Fig. 2A), indicating that AlsT is not a main serine transporter in *S. aureus*. Next,

a more detailed bioinformatics analysis was performed to identify potential AlsT substrates.

A BlastP search against the *B. subtilis* 168 genome led to the identification of four proteins

showing significant homology to the *S. aureus* AlsT (SAUSA300_1252) protein, namely AlsT

(e-value: e-166), GlnT (e-value: e-149), YrbD (e-value: e-117) and YflA (e-value: 2e-72).

Also in *S. aureus* a second AlsT homologue, SAUSA300_0914 (e-value 9e-108), could be

identified (Fig. S2), which is encoded at a different chromosomal region. AlsT is annotated in

B. subtilis as a potential glutamine sodium symporter (Zhu & Stülke, 2018), but to the best of

our knowledge, this has not yet been experimentally verified. To test if *S. aureus* AlsT is a

potential glutamine or glutamate transporter, uptake assays were performed with

radiolabelled glutamine and glutamate using the WT *S. aureus* strain LAC*, the *alsT* mutant

LAC*alsT::tn* piTET and the complementation strain LAC*alsT::tn* piTET-*alsT*. Uptake of

glutamine, but not of glutamate, was severely reduced in the *alsT* mutant when compared to

the WT strain (Fig. 2B-C). This defect was restored upon expression of *alsT* in the

complementation strain (Fig. 2C). To confirm that *alsT* functions as main glutamine

transporter also in the LAC*dacA/alsT* suppressor strain, uptake assays were also

performed with strain LAC*dacA/alsT* along with the WT LAC* and LAC*dacA::kan* control

strains (Fig. 2D-F). Similar as observed for the *alsT* single mutant, serine and glutamate

uptake were only marginally affected in strain LAC*dacA/alsT* (Fig. 2D-E), whereas

glutamine uptake was severely reduced in strain LAC*dacA/alsT* when compared to the

control strains (Fig. 2F). These data suggest that under the uptake assay conditions tested,

AlsT functions as the main glutamine transporter in *S. aureus.* A slight reduction in glutamine

and serine uptake was seen in the absence of c-di-AMP (Fig. 2D and Fig. 2F), suggesting

that c-di-AMP levels can impact glutamine and serine uptake in *S. aureus.* Taken together,

our data suggest that *S. aureus* cells that are unable to produce c-di-AMP can survive in rich

medium such as TSB, when glutamine uptake is reduced or blocked. However, we cannot

formally exclude that AlsT is able to transport other amino acid or substrates present in rich

medium.

Investigating the contribution of SAUSA300_0914 and GlnQ to glutamine and glutamate transport in *S. aureus*

S. aureus SAUSA300_0914 codes for a predicted amino acid symporter, which shows 41% identity to the *S. aureus* AlsT protein. After assigning AlsT a function as glutamine transporter, we wanted to test if SAUSA300_0914 might also play a role in glutamine or glutamate transport. To this end, strain LAC**0914::tn* was constructed by transducing the genomic region from the NMTL strain NE1463 (Fey *et al.*, 2013) containing a transposon insertion in *SAUSA300_0914* into the *S. aureus* LAC* background. Subsequently the uptake of radiolabelled glutamine and glutamate was assessed (Fig. 3A-B). No significant differences in the uptake of these amino acids was observed between WT LAC* and strain LAC**0914::tn*, showing that SAUSA300_0914 does not function as a major glutamine or glutamate transporter under our assay conditions (Fig. 3A-B). AlsT and SAUSA300_0914 are members of the amino acid/sodium symporter family of transporters, which are composed of a single multimembrane spanning protein. Besides this type of transporter, GlnPQ-type ABC transporters play a major role in glutamine and glutamate transport in other bacteria (Schuurman-Wolters & Poolman, 2005). *S. aureus* contains a *glnPQ* *(SAUSA300_1808 - SAUSA300_1807)* operon with *glnP* coding for a substrate binding domain-permease fusion protein and *glnQ* coding for the cytoplasmic nucleotide-binding ATPase domain. The results from a previous study suggested that this transporter functions as glutamine transporter in *S. aureus*, as a *glnP* mutant was more resistant to the toxic glutamine analogue γ-L-glutamyl hydrazide (Zhu *et al.*, 2009). To assess the contribution of the GlnPQ transporter to glutamine and glutamate transport in *S. aureus* LAC*, the strain LAC**glnQ::tn* was generated by transducing the *glnQ::tn* region from the NMTL strain NE153 (Fey *et al.*, 2013) into the LAC* background. The resulting *glnQ* mutant strain LAC**glnQ::tn* displayed no difference in glutamine or glutamate uptake compared to WT LAC* (Fig. 3C-D). This indicates that the ABC transporter GlnPQ does not function under our assay conditions and in the *S. aureus* LAC* strain background as a main glutamate or glutamine transporter.

Inactivation of AlsT but not SAUSA300_0914 or GlnQ reduces the susceptibility of *S. aureus* to the toxic glutamine analogue γ-L-glutamyl hydrazide

To further validate the findings from the uptake assays and verify that AlsT is the main glutamine transporter, we performed growth curves in the presence of increasing concentrations of the toxic glutamine analogue γ-L-glutamyl hydrazide with the WT and LAC**alsT::tn* mutant strains. Strains LAC**0914::tn* and LAC**glnQ::tn* were also included in these assays, to uncover a potential low level glutamine uptake activity for the SAUSA300_0914 and GlnPQ transporters. Strains defective in taking up this glutamine analogue are expected to show reduced susceptibility to this toxic compound. In the absence of the compound, all strains grew similarly in the chemically defined medium used for this assay (Fig. 4A). As expected, addition of γ-L-glutamyl hydrazide reduced the growth of the WT LAC* strain, in a dose-deponent manner (Fig. 4B and Fig. S3). Similar growth inhibition curves to that of the WT strain were obtained for strains LAC**0914::tn* and LAC**glnQ::tn*, while strain LAC**alsT::tn* showed increased resistance to the compound (Fig. 4B and Fig. S3). These findings support our earlier conclusion that AlsT is the main glutamine transporter in *S. aureus*, while GlnPQ and SAUSA300_0914 are either unable to take up glutamine or play only a minor role in its uptake under our growth conditions.

GltS (SAUSA300_2291) is a glutamate transporter in *S. aureus*

S. aureus does not only take up glutamine but also shows robust glutamate uptake (Fig. 2 and Fig. 3). However, none of the transporters (AlsT, SAUSA300_0914 and GlnPQ) investigated so far plays a major role in glutamate uptake under our growth conditions. In *B. subtilis* GltT, belonging to the dicarboxylate/amino acid cation symporter (DAACS) family of proteins, is a major high-affinity Na^+-coupled glutamate/aspartate symporter and can also mediate the uptake of glyphosate (Wicke *et al.*, 2019). Two paralogs, DctP and GltP, are found in *B. subtilis* of which GltP has also been shown to be a glutamate transporter (Tolner *et al.*, 1995). The *S. aureus* protein SAUSA300_2329 (from here on referred to as GltT)

shows a high degree of similarity (52% identity) to the *B. subtilis* GltT protein. In addition,

SAUSA300_2291 (from here on referred to as GltS) is annotated in UniProt

(www.uniprot.org) as a potential glutamate transporter in *S. aureus*. To experimentally test if

GltT or GltS impact glutamate transport in *S. aureus*, strains LAC**gltT::tn* and LAC**gltS::tn*

were constructed by moving the respective *gltT and gltS* transposon insertion regions from

the NMTL strains NE566 and NE560 (Fey *et al.*, 2013) into the LAC* strain background.

Next, the uptake of radiolabelled glutamine and glutamate was assessed for WT LAC* and

strains LAC* *gltT::tn* and LAC* *gltS::tn*. No difference in the uptake of glutamine was

observed between the strains (Fig. 5A) and in the case of LAC**gltT::tn*, also no difference in

the uptake of glutamate was observed (Fig. 5B). However, a significant reduction in

glutamate uptake was observed for strain LAC**gltS::tn* when compared to the WT strain

(Fig. 5B). The glutamate uptake defect could be restored in a complementation strain

harbouring plasmid piTET-*gltS* allowing for inducible *gltS* expression (Fig. 5C). Indeed,

increased glutamate uptake was observed in the complementation strain, indicating

increased *gltS* expression in the complementation strain as compared to the WT strain.

Taken together, these data indicate that under the growth conditions tested, GltS is the main

glutamate transporter in *S. aureus*.

Ammonium and glutamine but not glutamate stimulate the growth of *S. aureus* in

defined medium containing glucose as carbon source

Glutamine and glutamate are important amino acids that can serve, together with

ammonium, as nitrogen sources for the synthesis of many other cellular metabolites. Since

S. aureus is phenotypically auxotroph for many amino acids and at the same time can use

several amino acids as carbon and nitrogen sources, it is not possible to grow this organism

in any of the typical minimal media that are used to assess the ability of bacteria to

specifically use ammonium, glutamine or glutamate as nitrogen sources. However, to begin

to examine the effect of these compounds on the growth of *S. aureus*, growth curves were

performed with the WT LAC* strain in glucose containing defined medium (GDM), containing

essential vitamins, metals and 17 amino acids but lacking ammonium, glutamine and

glutamate as potential nitrogen/amino acid sources (see Table S1 for medium composition).

In addition, the WT LAC* was also grown in GDM containing glutamine (GDM+Gln),

glutamate (GDM+Glu), ammonium (GDM+NH$_3$), glutamine and ammonium (GDM+Gln+NH$_3$)

or glutamate and ammonium (GDM+Glu+NH$_3$). The addition of glutamine or ammonium

alone or in combinations stimulated the growth of the WT LAC* strain as compared to its

growth in GDM (Fig. 6A). On the other hand, no growth improvement was seen in the

presence of glutamate (GDM+Glu) (Fig. 6A). To examine the contribution of the glutamine

and glutamate transporters AlsT and GltS, additional growth curves were performed in the

different media with WT LAC* as well as strains LAC*alsT::tn and LAC*gltS::tn (Fig. S4).

Similar growth profiles were observed for all strains in the different media (Fig. S4), except in

GDM+Gln, in which the alsT mutant strain exhibited reduced growth compared to the WT

and gltS mutant strains (Fig. S4B). The growth defect could be restored in the alsT

complementation strain harbouring plasmid piTET-alsT (Fig. 6B). Taken together, these data

indicate that ammonium and glutamine are preferred over glutamate for the growth of S.

aureus. The observation that the addition of ammonium improves the growth of S. aureus

indicates that our base medium is likely nitrogen limiting and suggests that glutamine but not

glutamate can likely also serve as nitrogen source under these growth conditions. Finally,

these data further confirm the importance of AlsT for glutamine uptake in S. aureus.

Ammonium and glutamine uptake lead to a reduction in c-di-AMP levels in *S. aureus*

For *B. subtilis,* it has been reported that the addition of glutamine, glutamate or ammonium

to a defined growth medium can affect cellular c-di-AMP levels (Gundlach et al., 2015). It

was further proposed that glutamate uptake and to some extent also ammonium uptake

leads to an activation of c-di-AMP synthesis in this organism (Gundlach et al., 2015). To

assess if the presence of glutamine, glutamate or ammonium would also affect c-di-AMP

levels in *S. aureus*, the intracellular c-di-AMP concentrations were determined for the WT *S.*

aureus strain LAC* following growth in GDM, GDM+Gln, GDM+Glu, GDM+NH$_3$,

GDM+Gln+NH$_3$ and GDM+Glu+NH$_3$. Using a competitive ELISA assay, c-di-AMP could be

readily detected in bacteria grown in GDM, our base medium (Fig. 7A). Similar amounts of c-

di-AMP were detected in bacteria grown in the glutamate-containing medium (GDM+Glu),

however the c-di-AMP levels were significantly lower in bacteria grown in medium containing

either ammonium or glutamine (GDM+Gln, GDM+NH$_3$, GDM+Gln+NH$_3$, GDM+Glu+NH$_3$)

(Fig. 7A). To verify that the addition of glutamine reduces c-di-AMP production and to

investigate the contribution of the glutamine transporter AlsT to this inhibition, c-di-AMP

levels were determined for WT piTET, the *alsT* mutant LAC**alsT::tn* piTET and the

complementation strain LAC**alsT::tn* piTET-*alsT* following growth in the glutamine

containing medium GDM+Gln. While again low c-di-AMP levels were detected for the WT

strain, the c-di-AMP levels increased significantly in the *alsT* mutant and were restored back

to wild-type levels in the complementation strain (Fig. 7B). A similar experiment was

performed with the *gltS* mutant and complementation strain in the glutamate containing

medium GDM+Glu. High and similar c-di-AMP levels were detected for all strains (Fig. 7C),

indicating that neither the addition of glutamate to the medium nor its uptake impacts c-di-

AMP production in *S. aureus* under our test conditions. Taken together, these data highlight

that ammonium as well as AlsT-mediated glutamine uptake represses c-di-AMP production

in *S. aureus*.

The inhibition of the c-di-AMP production by glutamine and ammonium is not

mediated by GdpP or YbbR

The observed reduction of c-di-AMP levels in the presence of glutamine or ammonium could

potentially be achieved through an increase in the activity of the c-di-AMP specific

phosphodiesterase GdpP. To investigate this, cellular c-di-AMP levels were compared

between the WT LAC* and the isogenic *gdpP* mutant strain LAC**gdpP::kan*. As previously

reported for strain LAC*gdpP::kan following growth in TSB medium (Corrigan et al., 2011),

the c-di-AMP levels were also increased in the gdpP mutant compared to the WT strain

following growth in GDM, the glucose containing defined medium used as part of this study

(Fig. 8A). However, a significant reduction in the cellular c-di-AMP levels was also seen for

the gdpP mutant following the addition of glutamine or ammonium to the medium (Fig. 8A).

This indicates that the reduction in c-di-AMP levels upon addition of glutamine or ammonium

is likely due to decreased synthesis by DacA and not increased degradation by GdpP. We

next tested the involvement of YbbR, a proposed c-di-AMP cyclase regulator, by comparing

the cellular c-di-AMP levels produced by WT LAC* and strain LAC*ΔybbR. Similar c-di-AMP

levels were detected in the WT and ybbR mutant in GDM medium (Fig. 8B). The addition of

glutamine or ammonium to the medium led also to a large reduction in the cellular c-di-AMP

in the ybbR mutant strain (Fig. 8B). These data suggest that the observed reduction of c-di-

AMP production in the presence of glutamine and ammonium is neither mediated by GdpP

nor YbbR, and hence involves a different regulator protein, or that the cellular glutamine and

nitrogen levels are directly sensed by the cyclase DacA.

Discussion

Over the last decade, considerable evidence has emerged that c-di-AMP plays a

major role in osmotic regulation in bacteria, primarily by positively regulating potassium

export or negatively regulating potassium and osmolyte uptake (Rocha et al., 2019,

Quintana et al., 2019, Kim et al., 2015, Corrigan et al., 2013, Moscoso et al., 2015, Chin et

al., 2015, Huynh et al., 2016, Schuster et al., 2016, Pham & Turner, 2019, Pham et al.,

2018, Devaux et al., 2018, Zarrella et al., 2018, Gundlach et al., 2017a, Gundlach et al.,

2018, Gundlach et al., 2017b, Gundlach et al., 2019). However, individual c-di-AMP target

proteins identified thus far are themselves not essential. Therefore, the essentiality of c-di-

AMP is likely due to its ability to regulate multiple target proteins simultaneously.

Furthermore, in the absence of this molecule, many transporters are activated rather than

inactivated, likely leading to accumulation of toxic levels of metabolites, such as potassium

and osmolytes. Consistent with this idea, inactivating mutations in potassium uptake

systems, oligopeptide and osmolyte transporters have been reported to rescue the growth

defect of bacteria unable to produce c-di-AMP (Whiteley *et al.*, 2015, Whiteley *et al.*, 2017,

Gundlach *et al.*, 2017a, Gundlach *et al.*, 2017b, Pham *et al.*, 2018, Devaux *et al.*, 2018,

Zeden *et al.*, 2018). We have previously shown that inactivation of the main glycine betaine

transporter OpuD bypasses the requirement of c-di-AMP for the growth of *S. aureus* in rich

medium (Zeden *et al.*, 2018). We hypothesize that inactivation of OpuD might help a c-di-

AMP null strain survive by allowing bacteria to re-establish their osmotic balance. Bacteria of

the *dacA/opuD* mutant strain, which cannot produce c-di-AMP but are also defective in

glycine betaine transport, are similar in size to WT bacteria (Fig. 1). At this point it is not

known if c-di-AMP can directly bind to and regulate the activity of the *S. aureus* OpuD

protein. We attempted to address this question; however, despite using multiple different

approaches, we were unable to produce sufficient amounts of the full-length OpuD

membrane protein to perform protein/nucleotide interaction studies. On the other hand, a

direct role for c-di-AMP in the control of glycine betaine or betaine transporters has been

reported in *S. agalactiae* and *L. lactis*. In these organisms, c-di-AMP binds to the

transcriptional regulator BusR, which controls the expression of the predicted glycine betaine

or betaine transporter BusAB (Devaux *et al.*, 2018, Pham *et al.*, 2018).

Bacteria of the *dacA/alsT* suppressor strain, which survive in the absence of c-di-

AMP, remained enlarged, indicating that the essentiality of c-di-AMP is bypassed in this

strain potentially through a different mechanism. Here, we show that AlsT is an efficient

glutamine transporter in *S. aureus* (Fig. 2). These findings indicate that eliminating or

reducing the ability of *S. aureus* to take up glutamine from rich growth medium rescues the

growth of an *S. aureus* unable to produce c-di-AMP. There are several (not mutually

exclusive) possibilities how preventing glutamine uptake could rescue the growth of a c-di-

AMP null strain in rich medium. Glutamine as well as proline have been shown to

accumulate in *S. aureus* under NaCl stress conditions (Anderson & Witter, 1982). While it

has been suggested that the glutamine accumulation is due to synthesis rather than uptake

(Anderson & Witter, 1982), these data highlight that glutamine likely plays an important role

in osmotic regulation in *S. aureus.* Despite the cell size not being restored in the *dacA/alsT*

suppressor strain, blocking glutamine uptake could potentially still help bacteria to better

balance their cellular osmolality during growth in rich medium in the absence of c-di-AMP.

Another possible explanation how eliminating glutamine uptake could allow *S. aureus* to

grow in the absence of c-di-AMP could be related to changes in metabolism and TCA cycle

activity. In *L. monocytogenes,* an increased flux of pyruvate into the TCA cycle has been

described for a strain unable to produce c-di-AMP (Sureka *et al.*, 2014). As a consequence,

an accumulation of citrate and increased carbon flux into glutamine/glutamate was

observed, which resulted in a metabolic imbalance and growth defect (Sureka *et al.*, 2014).

Perhaps similar to the observations in *L. monocytogenes*, the absence of c-di-AMP could

also boost TCA cycle activity in *S. aureus*, thus leading to glutamine accumulation and a

metabolic imbalance. Hence, the lack of c-di-AMP combined with active glutamine uptake

could fuel the bacterial metabolism and the resulting metabolic imbalance might become

toxic to the cell, similar as observed for *L. monocytogenes* (Sureka *et al.*, 2014, Whiteley *et*

al., 2017).

In a recent study investigating genetic determinants required for eDNA release

during biofilm formation, it was found that inactivation of GdpP as well as of AlsT, resulted in

a significant decrease in eDNA release and in an increase in resistance to Congo red

(DeFrancesco *et al.*, 2017). Therefore, inactivation of AlsT and preventing/reducing

glutamine uptake might lead to alterations in the bacterial cell wall that make bacteria more

resistant to cell lysis. Such changes could also be an advantage during osmotic stress or c-

di-AMP deficiency. Indeed, we have recently shown a correlation between specific changes

in the peptidoglycan structure and the NaCl stress resistance in *S. aureus* (Schuster *et al.*,

2019). In addition, since the cellular c-di-AMP levels are significantly higher in the *gdpP* as

well as the *alsT* mutant strains compared to WT (Fig. 7B and Fig. 8A), the underlying

mechanistic bases for the decrease in eDNA release observed for the *gdpP* and *alsT* mutant

strains might be related.

The actual stimuli and underlying molecular mechanisms that regulate c-di-AMP

production in bacterial cells are at the moment poorly understood. As part of this study, we

show that ammonium and AlsT-mediated glutamine uptake but not GltS-mediated glutamate

uptake negatively impacts c-di-AMP production (Fig. 7). Changes in cellular c-di-AMP levels

depending on the presence of ammonium, glutamine or glutamate have already been

reported for *B. subtilis* (Gundlach *et al.*, 2015). For *B. subtilis* it has been suggested that

glutamine and to some extent ammonium uptake stimulates c-di-AMP production (Gundlach

et al., 2015). Here we show that in *S. aureus* ammonium and glutamine uptake leads to an

inhibition of c-di-AMP production rather than glutamate promoting its synthesis (Fig. 7). The

decrease in c-di-AMP production in the presence of ammonium or glutamine is likely

achieved by reducing the activity of the c-di-AMP cyclase DacA and not by activation of the

c-di-AMP specific phosphodiesterase GdpP. This conclusion is based on our observation

that the cellular c-di-AMP levels also decreased in a *gdpP* mutant strain upon addition of

glutamine or ammonium (Fig. 8A). Current evidence suggests that the activity of DacA can

be regulated through the interaction with two proteins: the membrane anchored and

proposed DacA regulator protein YbbR (also name CdaR in other bacteria) and the

phosphoglucomutase enzyme GlmM (Tosi *et al.*, 2019, Zhu *et al.*, 2016, Gundlach *et al.*,

2015, Pham *et al.*, 2016). We could exclude that the observed reduction in cellular c-di-AMP

levels in the presence of ammonium or glutamine is mediated by YbbR, as <u>a</u> *ybbR* mutant

showed a similar decrease in the c-di-AMP levels as observed for the WT strain (Fig. 8B).

GlmM has been shown to be a negative regulator of DacA activity both *in vivo* and *in vitro*

(Tosi *et al.*, 2019, Pham *et al.*, 2018). However, since GlmM is likely an essential enzyme in

S. aureus, we were unable to construct a *glmM* mutant and test its involvement in the

observed repression of c-di-AMP synthesis in the presence of ammonium or glutamine as

we did for the *gdpP* and *ybbR* mutant strains. Nevertheless, with this work, we not only

identified main glutamine and glutamate transporters in *S. aureus,* but we also linked the c-

di-AMP signalling network to central nitrogen metabolism in *S. aureus.* It will be interesting to

determine in future studies the mechanistic bases for the observed changes in cellular c-di-

AMP levels depending on ammonium and glutamine uptake and the involvement of GlmM or

other factors in this process.

Experimental Procedures

Bacterial strains and culture conditions

Bacterial strains used in this study are listed in Table 1. *S. aureus* strains were grown in

Tryptic Soy Broth (TSB), Tryptic Soy Agar (TSA) or Glucose Defined Medium (GDM). GDM

was prepared similar to the chemically defined medium (CDM) reported in an earlier study

(Zeden *et al.*, 2018), with some modifications. The detailed content of the GDM (which

contains essential vitamins, trace metals, amino acids and glucose as a carbon source but

lacks ammonium, glutamine or glutamate as a potential nitrogen or amino acid source) is

shown in Table S1. In addition to the GDM, GDM containing glutamine (GDM+Gln),

glutamate (GDM+Glu), ammonium (GDM+NH_3), glutamine and ammonium (GDM+Gln+NH_3)

or glutamate and ammonium (GDM+Glu+NH_3) were used as part of this study (for exact

composition see Table S1). Where indicated, certain amino acids were removed from the

GDM recipe during uptake assays and when needed the TSB was supplemented with 0.4 M

NaCl. *Escherichia coli* strains were grown in Lysogeny Broth (LB). Where appropriate,

antibiotics and/or inducers were added to the media at the following concentration: 200

ng/ml anhydrotetracycline (Atet), 90 µg/ml Kanamycin (Kan), 10 µg/ml Erythromycin (Erm),

7.5 or 10 µg/ml Chloramphenicol (Cam), Ampicillin (Amp) 100 µg/ml.

Bacterial strain construction

All strains used in this study are listed in Table 1 and primers used in this study are listed in

Table 2. The transposon insertion sites in the Nebraska transposon mutant library (NTML)

strains (Fey *et al.*, 2013) used as part of this study were confirmed by PCR and sequencing.

The transposon and surrounding regions were moved by phage transduction using phage 85

into the *S. aureus* LAC* strain background. This resulted in the generation of *S. aureus*

strains LAC**alsT::tn* (*SAUSA300_1252::tn*; ANG4803), LAC**0914::tn* (*SAUSA300_0914::tn*;

ANG5141), LAC*glnQ::tn (SAUSA300_1807::tn; ANG5070), LAC*gltT::tn

(SAUSA300_2329::tn; ANG5366) and LAC*gltS::tn (SAUSA300_2291::tn; ANG5367). The

transposon insertion in the respective gene was again confirmed by PCR and sequencing.

For complementation analysis, the Atet inducible single copy integration plasmids piTET-

alsT and piTET-gltS were constructed. To this end, alsT (SAUSA300_1252) and gltS

(SAUSA300_2291) were amplified using LAC* chromosomal DNA and primers

ANG2250/ANG2251 and ANG3209/ANG3210, respectively. The products as well as piTET

were digested with AvrII and SacII and then ligated. Plasmid piTET-alsT was recovered in E.

coli strain XL1-Blue (yielding strain ANG3937), shuttled through E. coli strain IM08B (strain

ANG3955) and then introduced into LAC*alsT::tn (ANG4803), yielding strain LAC*alsT::tn

piTET-alsT (ANG4854). As a control, plasmid piTET was also introduced into LAC*alsT::tn

(ANG4803) yielding strain LAC*alsT::tn piTET (ANG4853). Plasmid piTET-gltS was

transformed into E. coli XL1-Blue (yielding strain ANG5494), shuttled through E. coli IM08B

(yielding strain ANG5495) and transformed into LAC*gltS::tn, yielding the complement strain

LAC*gltS::tn piTET-gltS (ANG5493). As a control, the piTET plasmid was transformed into

LAC*gltS::tn strain, yielding the strain LAC*gltS:tn piTET (ANG5492). Correct plasmid

integration into the geh locus was confirmed by PCR and the sequences of all plasmid

inserts were confirmed by fluorescent automated sequencing.

Bacterial growth curves and amino acid analysis in culture supernatants

S. aureus strains LAC* and LAC*alsT::tn were grown overnight in TSB supplemented with

10 µg/ml erythromycin where appropriate. Overnight cultures were then diluted to an OD_{600}

of 0.01 into 50 ml of fresh TSB. Cultures were incubated at 37°C with aeration, and OD_{600}

values determined every hour. The experiment was performed with three biological

replicates and the average OD_{600} values and standard deviations (SDs) were plotted. Using

the same cultures, supernatant samples were prepared at the 0, 6, 10 and 12 h time points

and amino acid levels determined as previously described using an amino acid analyser

(Halsey et al., 2017). For measuring the growth of S. aureus strains LAC*, LAC* piTET,

LAC*alsT::tn, LAC*alsT::tn piTET, LAC*alsT::tn piTET-alsT and LAC*gltS::tn in GDM,

GDM+Gln, GDM+Glu, GDM+NH_3, GDM+Gln+NH_3 or GDM+Glu+NH_3, the bacteria were

grown overnight in TSB medium supplemented with chloramphenicol and erythromycin

where appropriate. Next day, bacteria from a 1 ml aliquot were washed twice with PBS and

diluted to OD_{600} of 0.005 in the indicated GDM. LAC* WT, LAC*alsT::tn and LAC*gltS::tn

were grown in GDM, GDM+Gln, GDM+Glu, GDM+NH_3, GDM+Gln+NH_3 and GDM+Glu+NH_3

while LAC* piTET, LAC*alsT::tn piTET and LAC*alsT::tn piTET-alsT were grown in

GDM+Gln supplemented with 200 ng/ml Atet. One hundred µl of the diluted cultures (six

technical replicates) were transferred into wells of a 96-well plate and the plate was then

incubated with shaking (500 rpm) in a plate reader and OD_{600} readings determined every 30

min. The average values of the technical replicates were determined for each strain. The

experiment was performed three times and the average readings and standard deviations

were plotted.

γ-L-glutamyl hydrazide susceptibility assay

The susceptibility of *S. aureus* LAC*, LAC*alsT::tn (ANG4803), LAC*0914::tn (ANG5141)

and LAC*glnQ::tn (ANG5242) to the toxic glutamine analogue γ-L-glutamyl hydrazide (Alfa

Aesar, MA, USA) was determined using a similar method as previously reported (Zhu *et al.*,

2009). Briefly, the different strains were grown overnight at 37°C in 5 ml TSB medium,

supplemented with 10 µg/ml erythromycin where appropriate. Next day, the bacteria were

washed twice with PBS, diluted to an OD_{600} of 0.005 in GDM+NH_3. Next, 10 µl of water (0

mM control) or 10 µl of a γ-L-glutamyl hydrazide solution dissolved in water was added to

0.99 ml aliquots of these bacterial suspension to give a final concentration of 20, 40, 60 or

80 µg/ml, respectively. One hundred µl were subsequently transferred in four replicates into

wells of a 96 well plates and the plate incubated at 37°C with shaking (500 rpm) in a plate

reader and OD_{600} readings determined every 10 min for 12 h. The experiment was

performed three times and the average OD_{600} values of the three experiments presented as

growth curves. The average values and SDs of the OD_{600} values from the 7 h time point

were also plotted against the different γ-L-glutamyl hydrazide concentrations.

Microscopic analysis and cell size measurements

The microscopic analysis to determine bacterial cell sizes was performed essentially as

previously described (Zeden *et al.*, 2018). Briefly, *S. aureus* strains LAC*, LAC**dacA::kan*,

LAC**dacA*$_{G206S}$, LAC**dacA/opuD* (ANG3835) and LAC**dacA/alsT* (ANG3838) were grown

overnight at 37°C in TSB or TSB supplemented with 0.4 M NaCl where stated. Next day, the

cultures were diluted to an OD_{600} of 0.01 and grown for 3 h at 37°C to mid-exponential phase

(OD_{600} of 0.5-0.9). One hundred μl of these cultures were then stained for 20 min at 37°C

with Vancomycin-BODIPY FL at a final concentration of 2 μg/ml. One and a half μl of each

sample was spotted onto a thin 1.5% agarose gel patch prepared in H_2O or in 0.4 M NaCl

and the bacteria subsequently imaged at 1000 x magnification using an Axio Imager A2

Zeiss microscope equipped with a GFP filter set. Images were acquired using the ZEN 2012

(blue edition) software. The bacterial cell diameters were determined using the Fiji software.

Only non-dividing cells (cells without any obvious fluorescent dots or lines at the mid-cell),

were used for cell diameter measurements. The cell diameters of 50 cells were measured

and the average cell diameter determined. The experiment was conducted three or four

times (as indicated in the figure legend) and the averages and standard deviations of the

average cell diameters plotted.

Uptake assays using ^{14}**C-labelled amino acids**

Uptake assays were conducted as previously described with some minor modifications

(Zeden *et al.*, 2018). Briefly, *S. aureus* strains were streaked on TSA or TSA 0. 4M NaCl

plates with appropriate antibiotics and the plates incubated overnight at 37°C. Bacteria were

subsequently scraped off from the plates and suspended in 1 ml PBS pH 7.4 buffer and the

OD_{600} determined. Fifty ml of GDM+Glu+NH_3 (where indicated with 200 ng/ml of the inducer

Atet added) was inoculated with the appropriate bacterial suspensions to an OD_{600} of 0.05.

The cultures were grown at 37°C to an OD_{600} between 0.4 and 0.9 and bacteria from an

OD_{600} equivalent of 8 were harvested by centrifugation for 10 min at 19,000 x g at RT.

Supernatants were discarded and the bacterial pellets were suspended in 2 ml of $GDM+NH_3$

(for glutamine and glutamate uptake assays) or $GDM+Glu+NH_3$ without serine (for serine

uptake assays). The OD_{600} of the cell suspensions were measured and the cells diluted to

an OD_{600} of approximately 1. The OD_{600} was re-measured and this measurement used for

normalization purposes. Five hundred and fifty µl of these cell suspensions were aliquoted

into 50 ml conical tubes and 100 µl used to measure the background radiation, by filtering

the cells onto a nitrocellulose membrane filter, followed by a wash step with 16 ml PBS.

Then, 6.2 µl of glutamine, L-[14C(U)] (Hartmann Analytic, MC1124), glutamic acid, L-

[14C(U)] (Hartmann Analytic, MC156), or serine L-[14C(U)] (Hartmann Analytic, MC265)

was added to the remaining 450 µl sample. One hundred µl aliquots were filtered 0, 3, 6 and

9 min after addition of the radiolabelled amino acid and the filters were then washed two

times with 16 ml of PBS pH 7.4. The filters were subsequently dissolved in 9 ml of

scintillation cocktail Filter Count (Perkin Elmer) and the radioactivity measured in counts per

minute (CPM) using a Wallac 1409 DSA liquid scintillation counter. The CPMs were then

normalized to the OD_{600} reading of the final cell suspension and the means and standard

deviations of the CPM/ml OD_{600} = 1 of three or four (as indicated in the figure legends)

independent experiments were plotted.

Determination of cellular c-di-AMP levels by competitive ELISA

Intracellular c-di-AMP levels in WT LAC* and the indicated *S. aureus* mutant strains were

determined using a previously described competitive ELISA method (Underwood *et al.*,

2014) and a slightly modified method for the preparation of *S. aureus* samples (Bowman *et*

al., 2016). Briefly, single colonies of the WT LAC* strain were picked from TSA plates and

used to inoculate 5 ml of GDM, GDM+Gln, GDM+Glu, $GDM+NH_3$, $GDM+Gln+NH_3$ and

$GDM+Glu+NH_3$. Colonies of the strains LAC*gdpP::kan and LAC*ΔybbR were inoculated

into 5 ml of GDM, GDM+Glu and GDM+Gln. Colonies of strains LAC* piTET, LAC*alsT::tn

piTET and LAC*alsT::tn piTET-alsT were inoculated into GDM+Gln containing 200 ng/ml

Atet and colonies of strains LAC* piTET, LAC*gltS::tn piTET and LAC*gltS::tn piTET-gltS

were inoculated into GDM+Glu supplemented with 200 ng/ml Atet. All cultures were

incubated for 18 h at 37°C with shaking. Next, bacteria from 4.5 ml culture were collected by

centrifugation, washed three times with PBS and subsequently suspended in 0.75 to 1 ml 50

mM Tris pH 8 buffer supplemented with 20 ng/ml lysostaphin and the cells were lysed by

bead beating. The lysates were cleared by centrifugation for 5 min at 17,000 x g and the

supernatant transferred to a new tube. A small sample aliquot was removed, and the protein

concentration determined for normalization purposes using a Pierce BCA protein assay kit

(Thermo Scientific, Waltham, MA, USA). The remainder of the sample was heated to 95°C

for 10 min. For the competitive ELISA assay, the samples were diluted to a protein

concentration of 100, 200, 400 or 500 µg/ml as, appropriate. ELISA plates were prepared by

adding 100 µl of coating buffer (50 mM Na_2C0_3, 50 mM $NaHCO_3$, pH 9.6) containing 10

µg/ml of the c-di-AMP binding protein $CpaA_{SP}$ to each well of a 96 well NUNC MaxiSorp

plate (Thermo Scientific, Waltham, MA, USA) and the plate was incubated for approximately

18 h at 4°C. Next, the plate was washed three times with 200 µl PBST pH 7.4 (10 mM

Na_2HPO_4 , 1.8 mM KH_2PO_4 137 mM NaCl , 2.7 mM KCl , 0.05% (v/v) Tween 20), blocked for

1 h at 18°C with 150 µl blocking solution (1% BSA in PBS pH 7.4) and washed three times

with 200 µl PBST. Fifty µl of the samples (three biological replicates and three technical

replicates) or standards (two technical replicates) were mixed with 50 µl of a 50 nM

biotinylated c-di-AMP solution prepared in 50 mM Tris pH 8 buffer. For the standard curve,

c-di-AMP standards were prepared in 50 mM Tris pH 8 buffer at concentrations of 0, 12.5,

25, 37.5, 50, 75, 100, 200 nM. Following the addition of the samples and the standards, the

plate was incubated for 2 h at 18°C and then washed three times with PBST. Next, 100 µl of

a high sensitivity streptavidin-HRP solution (Thermo Scientific, Waltham, MA, USA) diluted

1:500 in PBS was added to each well and the plate was incubated for 1 h at 18°C. The plate

was washed again 3 x with 200 µl PBST and 100 µl of a developing solution (0.103 M

$NaHPO_4$, 0.0485 M citric acid, 500 mg/l o-phenylenediamine dihydrochloride, 0.03% H_2O_2)

was added to each well and the plate incubated for 15 min at 18°C. The reaction was then

stopped by adding 100 µl of 2 M H_2SO_4 solution. The absorbance was measured in a plate

reader at a wavelength of 490 nm and c-di-AMP concentrations were calculated as ng c-di-

AMP / mg protein.

Acknowledgments: This work was funded by the Wellcome Trust grants 100289/Z/12/Z

and 210671/Z/18/Z to AG and the NIH/NIAID grants P01AI083211 and R01AI125588 to PDF

and VCT, respectively. MSZ was supported by a Medical Research Council Centre for

Molecular Bacteriology and Infection (MRC CMBI) studentship. The funders had no role in

the study design, data collection and interpretation, or the decision to submit the work for

publication. The authors have no conflicts of interest to declare. The data supporting the

findings of this study will be openly available.

Author contribution: MSZ, IK and AG designed the study, MSZ, IK and CFS acquired the

data, MSZ, IK, CFS, VCT, PDF and AG, designed experiments, analyzed and interpreted

the data, MSZ, IK and AG prepared the figures and wrote the original draft of the

manuscript. All authors approved the final version of the manuscript.

Graphical abstract:
639

640
641
642

Abbreviated Summary:

A large number of amino acid transporters and oligopeptide permeases are encoded in

bacterial genomes. However, their actual substrate specificity and functions are hard to

predict bioinformatically. In this study, we report that GltS and AlsT are main glutamate and

glutamine transporters in *Staphylococcus aureus*, respectively and show that glutamine and

ammonium uptake inhibits the production of the nucleotide signalling molecule c-di-AMP.

Table 1: Bacterial strains used in this study

Unique ID	Strain name and resistance	Source
Escherichia coli strains		
ANG284	XL1-Blue piTET; AmpR	(Gründling & Schneewind, 2007)
ANG2154	DH10B pIMAY; CamR	(Monk _et al._, 2012)
ANG3724	IM08B	(Monk _et al._, 2015)
ANG3928	IM08B piTET; AmpR	(Zeden _et al._, 2018)
ANG3937	XL1-Blue piTET-_alsT_; AmpR	This study
ANG3955	IM08B piTET-_alsT_; AmpR	This study
ANG5494	XL1-Blue piTET-_gltS_; AmpR	This study
ANG5495	IM08B piTET-_gltS;_ AmpR	This study
	Staphylococcus aureus strains	
AH1263	LAC* Erm sensitive CA-MRSA USA300 strain (ANG1575)	(Boles _et al._, 2010)
ANG1961	LAC*_gdpP::kan;_ KanR	(Corrigan _et al._, 2011)
ANG3301	LAC*Δ_ybbR_	(Bowman _et al._, 2016)
ANG3664	LAC*_dacA_$_{G206S}$; KanR	(Bowman _et al._, 2016)
ANG3666	LAC*_dacA::kan_ (_dacA_) KanR	(Zeden _et al._, 2018)
ANG3835	LAC*_dacA::kan_-S7 (LAC*_dacA/opuD_); KanR	(Zeden _et al._, 2018)
ANG3838	LAC*_dacA::kan_-S10 (LAC*_dacA/alsT_)]; KanR	(Zeden _et al._, 2018)
ANG3940	NE142 (_alsT::tn_) – NMTN strain	(Fey _et al._, 2013)
ANG4054	LAC* piTET; CamR	(Zeden _et al._, 2018)
ANG4803	LAC*_alsT::tn;_ ErmR	This study
ANG4854	LAC*_alsT::tn_ piTET-_alsT ;_ ErmR CamR	This study
ANG4853	LAC*_alsT::tn_ piTE _;_ ErmR CamR	This study
ANG4968	NE1463 (JE2 SAUSA300_0914::tn_) – NMTN strain	(Fey _et al._, 2013)
ANG5070	NE153 (JE2 _glnQ::tn_) – NMTN strain	(Fey _et al._, 2013)
ANG5141	LAC*_0914::tn;_ ErmR	This study
ANG5242	LAC*_glnQ::tn;_ ErmR	This study
ANG5309	NE566 (JE2 _gltT::tn_) – NMTN strain	(Fey _et al._, 2013)
ANG5310	NE560 (JE2 _gltS::tn_) – NMTN strain	(Fey _et al._, 2013)
ANG5366	LAC*_gltT::tn;_ ErmR	This study
ANG5367	LAC*_gltS::tn;_ ErmR	This study
ANG5492	LAC*_gltS::tn_ piTET; ErmR CamR	This study
ANG5493	LAC*_gltS::tn_ piTET-_gltS_; ErmR CamR	This study

653 **Table 2:** Cloning primers used in this study

Primer ID	Name	Sequence
ANG2250	5-AvrII-*alsT*	AGTCCCTAGGCGGTCTAATTTTATAGAAGG
ANG2251	3-SacII-*alsT*	TCCCCGCGGGGTTTATTTGATTTTTATATAATGAATCG
ANG3209	5-AvrII-*gltS*	ATACCTAGGAGGGAGAGGGATATTCAACAAGGGGGATTTG
ANG3210	3-SacII-*gltS*	GCCCGCGGTTTAACTAAACCATTGTATGAATCCCATAATG

 Figures and Legends

Fig. 1. Inactivation of the glycine betaine transporter OpuD rescues the cell size of *S. aureus dacA* mutant bacteria. (A, C, E) Microscopy images of *S. aureus* cells stained with BODIPY-labelled vancomycin. Cultures of *S. aureus* LAC* (WT), LAC*$dacA_{G206S}$ ($dacA_{G206S}$) (panels A and C only), LAC*$dacA::kan$ ($dacA$) (panel E only) and the suppressor strains LAC*$dacA/alsT$ ($dacA/alsT$) and LAC*$dacA/opuD$ ($dacA/opuD$) were grown in (A) TSB or (C and E) TSB 0.4 M NaCl medium and subsequently stained with fluorescently labelled vancomycin. The bacteria were then viewed using a fluorescent microscope and representative images are shown. Scale bars are 1 μm. (B, D, F) Bacterial cell diameter measurements. The diameter of non-dividing bacterial cells was measured as described in the Materials and Method section for *S. aureus* strains grown in (B) TSB or grown in (D and F) TSB 0.4 M NaCl medium. The diameters of 50 cells were determined and the average diameter calculated. The experiment was performed in triplicate (B and F) or quadruplicate (D) and the averages and SDs of the average cell diameters plotted. For statistical analysis, one-way ANOVAs followed by Dunnett's multiple comparison tests were performed (ns = not significant, * = $p<0.01$, ** = $p<0.001$, *** = $p<0.0001$).

Fig. 2. AlsT is a glutamine transporter in *S. aureus*. (A-F) Amino acid uptake assays. (A-C) *S. aureus* strain LAC* (WT), the *alsT* mutant LAC*alsT::tn* piTET (*alsT::tn* piTET) and the complementation strain LAC*alsT::tn* piTET-*alsT* (*alsT::tn* piTET-*alsT*) were grown to mid-log phase in glucose defined media as indicated in the method section and supplemented with 200 ng/ml Atet for the strains containing plasmids. Subsequently radiolabelled (A) serine, (B) glutamate or (C) glutamine was added to culture aliquots, samples removed and filtered at the indicated time points and the radioactivity accumulated in the cells measured. The average values and SDs from three (A, C, D, F) or four (B, E) experiments were plotted. (D-F) The same uptake assay experiment was performed as described in (A-C) but using *S. aureus* strains LAC*dacA::kan* (*dacA*) and LAC*dacA/alsT* (*dacA/alsT*). The amino acid uptake curve for the LAC* (WT) strain is the same as shown in panels A-C, as all strains were grown and processed at the same time.

Fig. 3. LAC*0914::tn* **and LAC***glnQ::tn* **strains do not show a defect in glutamine or glutamate uptake.** Amino acid uptake assays. (A and B) *S. aureus* strains LAC* (WT), LAC*0914::tn* and LAC*alsT::tn* were grown to mid-log phase in GDM+Glu+NH$_3$. Subsequently radiolabelled (A) glutamine or (B) glutamate was added to culture aliquots, samples removed and filtered at the indicated time points and the radioactivity accumulated in the cells measured. The average values and SDs from three experiments were plotted. (C and D) Amino acid uptake assays were performed and the data plotted as described in panels A and B, but using *S. aureus* strains LAC* (WT) and LAC*glnQ::tn* (*glnQ::tn*).

Fig. 4. LAC*_alsT::tn_ shows increased resistance to the toxic glutamine analogue γ-L-glutamyl hydrazide. (A) Bacterial growth curves. _S. aureus_ strains LAC* (WT), LAC*_alsT::tn_, LAC*_0914::tn_ and LAC*_glnQ::tn_ were grown in GDM+NH$_3$ medium in 96-well plates and OD$_{600}$ reading determined over 12 h. The average OD$_{600}$ values and SDs from three independent biological replicates were plotted. (B) γ-L-glutamyl hydrazide susceptibility assay. The same _S. aureus_ strains as in (A) were grown in GDM+NH$_3$ medium in the absence or presence of γ-L-glutamyl hydrazide at a final concentration of 20, 40, 60 or 80 µg/ml. OD$_{600}$ reading were determined over 12 h and the complete growth curves are shown in Figure S3. In this graph, the average OD$_{600}$ values from the 7 h time point (marked with an arrow in (A)) and SDs from three biological replicates were plotted against the γ-L-glutamyl hydrazide concentration in the growth medium.

Fig. 5: GltS is a glutamate transporter in *S. aureus*. Amino acid uptake assays. (A and B)

S. aureus strains LAC* (WT), LAC**gltT::tn* and LAC**gltS::tn* were grown to mid-log phase in

GDM+Glu+NH₃. Subsequently radiolabelled (A) glutamine or (B) glutamate was added to

culture aliquots, samples removed and filtered at the indicated time points and the

radioactivity accumulated in the cells measured. (C) Same as (B) but using *S. aureus* strains

LAC* (WT), LAC**gltS::tn* piTET and the complementation strain LAC**gltS::tn* piTET-*gltS* and

supplementing the GDM+Glu+NH₃ medium with 200 ng/µl Atet. The average values and

SDs from three experiments were plotted.

Fig. 6. Addition of glutamine and ammonium but not glutamate stimulates the growth

of *S. aureus* in glucose-containing defined medium. (A) Bacterial growth curves. The WT

S. aureus strain LAC* was grown in 96-well plates in glucose-containing defined medium

(GDM) or in GDM supplemented with glutamine (Gln), glutamate (Glu), ammonium (NH_3) or

combinations thereof as specified in the legend. OD_{600} readings were determined every 30

min and the average and SDs of three biological replicates plotted. (B) Bacterial growth

curves. *S. aureus* strains LAC* piTET (WT piTET), the *alsT* mutant strain LAC**alsT::tn*

piTET and the complementation strain LAC**alsT::tn* piTET-*alsT* were grown in GDM+Gln

medium supplemented with 200 ng/µl Atet. OD_{600} readings were determined every 30 min

and the average and SDS of three biological replicates plotted.

735
Fig. 7. Ammonium and glutamine uptake inhibit c-di-AMP production in *S. aureus*. (A-

C) Cellular c-di-AMP levels. (A) The WT *S. aureus* strain LAC* was grown in GDM or GDM

supplemented with glutamine (Gln), glutamate (Glu), ammonium (NH_3), or combinations

thereof as indicated on the X-axes. Cell extracts were prepared and c-di-AMP

concentrations measured using a competitive ELISA assay. The average values and SDs

from three biological replicates were plotted as ng c-di-AMP/mg protein. For statistical

analysis, one-way ANOVAs followed by Dunnett's multiple comparison tests were performed

to identify statistically significant differences between the different media as compared to

GDM (ns = not significant, ** = $p<0.001$, *** = $p<0.0001$). (B) *S. aureus* strains LAC* piTET

(WT piTET), LAC**alsT::tn* piTET and the complementation strain LAC**alsT::tn* piTET-*alsT*

were grown in GDM+Gln supplemented with 200 ng/µl Atet and c-di-AMP levels determined

and plotted as described in (A). (C) *S. aureus* strains LAC* piTET (WT piTET), LAC**gltS::tn*

piTET and the complementation strain LAC**gltS::tn* piTET-*gltS* were grown in GDM+Glu

supplemented with 200 ng/µl Atet and c-di-AMP levels determined and plotted as described

in (A).

Fig. 8. The inhibition of c-di-AMP production by glutamine and ammonium is independent of GdpP and YbbR. (A and B) Cellular c-di-AMP levels. (A) *S. aureus* strains LAC* (WT) and LAC*gdpP::kan (*gdpP::kan*) were grown in GDM or in GDM containing glutamine (GDM+Gln) or ammonium (GDM+NH₃). Cell extracts were prepared, and c-di-AMP concentrations measured using a competitive ELISA assay. The average values and SDs from three biological replicates were plotted as ng c-di-AMP/ mg protein. For statistical analysis one-way ANOVAs followed by Dunnett's multiple comparison tests were performed to identify statistically significant differences between the different media as compared to the GDM medium (n.d. = not determined. Statistical analysis was not performed due to values below our detection limit; these values were set to 0, ** = p<0.001, *** = p<0.0001). (B) Same as in (A) but using *S. aureus* strains LAC* (WT) and LAC*ΔybbR (Δ*ybbR*).

References

Anderson, C.B., and Witter, L.D. (1982) Glutamine and proline accumulation by
        Staphylococcus aureus with reduction in water activity. *Appl Environ Microbiol* **43**:
        1501-1503.
Bai, Y., Yang, J., Eisele, L.E., Underwood, A.J., Koestler, B.J., Waters, C.M., Metzger, D.W.,
        and Bai, G. (2013) Two DHH subfamily 1 proteins in *Streptococcus pneumoniae*
        possess cyclic di-AMP phosphodiesterase activity and affect bacterial growth and
        virulence. *J Bacteriol* **195**: 5123-5132.
Bai, Y., Yang, J., Zarrella, T.M., Zhang, Y., Metzger, D.W., and Bai, G. (2014) Cyclic di-AMP
        impairs potassium uptake mediated by a cyclic di-AMP binding protein in
        Streptococcus pneumoniae. *J Bacteriol* **196**: 614-623.
Boles, B.R., Thoendel, M., Roth, A.J., and Horswill, A.R. (2010) Identification of genes
        involved in polysaccharide-independent *Staphylococcus aureus* biofilm formation.
        PLoS One **5**: e10146.
Bowman, L., Zeden, M.S., Schuster, C.F., Kaever, V., and Gründling, A. (2016) New Insights
        into the Cyclic Di-adenosine Monophosphate (c-di-AMP) Degradation Pathway and
        the Requirement of the Cyclic Dinucleotide for Acid Stress Resistance in
        Staphylococcus aureus. *J Biol Chem* **291**: 26970-26986.
Chin, K.H., Liang, J.M., Yang, J.G., Shih, M.S., Tu, Z.L., Wang, Y.C., Sun, X.H., Hu, N.J.,
        Liang, Z.X., Dow, J.M., Ryan, R.P., and Chou, S.H. (2015) Structural Insights into the
        Distinct Binding Mode of Cyclic Di-AMP with SaCpaA_RCK. *Biochemistry* **54**: 4936-
        4951.
Commichau, F.M., Heidemann, J.L., Ficner, R., and Stülke, J. (2019) Making and Breaking
        of an Essential Poison: the Cyclases and Phosphodiesterases That Produce and
        Degrade the Essential Second Messenger Cyclic di-AMP in Bacteria. *Journal of
        Bacteriology* **201**: e00462-00418.
Corrigan, R.M., Abbott, J.C., Burhenne, H., Kaever, V., and Gründling, A. (2011) c-di-AMP is
        a new second messenger in *Staphylococcus aureus* with a role in controlling cell size
        and envelope stress. *PLoS Pathog* **7**: e1002217.
Corrigan, R.M., Bowman, L., Willis, A.R., Kaever, V., and Gründling, A. (2015) Cross-talk
        between two nucleotide-signaling pathways in *Staphylococcus aureus*. *J Biol Chem*
        290: 5826-5839.
Corrigan, R.M., Campeotto, I., Jeganathan, T., Roelofs, K.G., Lee, V.T., and Gründling, A.
        (2013) Systematic identification of conserved bacterial c-di-AMP receptor proteins.
        Proc Natl Acad Sci U S A **110**: 9084-9089.
Crooke, A.K., Fuller, J.R., Obrist, M.W., Tomkovich, S.E., Vitko, N.P., and Richardson, A.R.
        (2013) CcpA-independent glucose regulation of lactate dehydrogenase 1 in
        Staphylococcus aureus. *PLoS One* **8**: e54293.
DeFrancesco, A.S., Masloboeva, N., Syed, A.K., DeLoughery, A., Bradshaw, N., Li, G.W.,
        Gilmore, M.S., Walker, S., and Losick, R. (2017) Genome-wide screen for genes
        involved in eDNA release during biofilm formation by *Staphylococcus aureus*. *Proc
        Natl Acad Sci U S A* **114**: E5969-E5978.
Devaux, L., Sleiman, D., Mazzuoli, M.V., Gominet, M., Lanotte, P., Trieu-Cuot, P., Kaminski,
        P.A., and Firon, A. (2018) Cyclic di-AMP regulation of osmotic homeostasis is
        essential in Group B *Streptococcus*. *PLoS Genet* **14**: e1007342.
Fahmi, T., Faozia, S., Port, G., and Cho, K.H. (2019) The Second Messenger c-di-AMP
        Regulates Diverse Cellular Pathways Involved in Stress Response, Biofilm
        Formation, Cell Wall Homeostasis, SpeB Expression and Virulence in *Streptococcus
        pyogenes*. *Infect Immun* **87**: e00147-00119.

Fey, P.D., Endres, J.L., Yajjala, V.K., Widhelm, T.J., Boissy, R.J., Bose, J.L., and Bayles,
K.W. (2013) A genetic resource for rapid and comprehensive phenotype screening of
nonessential *Staphylococcus aureus* genes. *MBio* **4**: e00537-00512.
Fridkin, S.K., Hageman, J.C., Morrison, M., Sanza, L.T., Como-Sabetti, K., Jernigan, J.A.,
Harriman, K., Harrison, L.H., Lynfield, R., Farley, M.M., and Active Bacterial Core
Surveillance Program of the Emerging Infections Program, N. (2005) Methicillin-
resistant *Staphylococcus aureus* disease in three communities. *N Engl J Med* **352**:
1436-1444.
Fuller, J.R., Vitko, N.P., Perkowski, E.F., Scott, E., Khatri, D., Spontak, J.S., Thurlow, L.R.,
and Richardson, A.R. (2011) Identification of a lactate-quinone oxidoreductase in
Staphylococcus aureus that is essential for virulence. *Front Cell Infect Microbiol* **1**:
19.
Gründling, A., and Schneewind, O. (2007) Genes required for glycolipid synthesis and
lipoteichoic acid anchoring in *Staphylococcus aureus*. *J Bacteriol* **189**: 2521-2530.
Gundlach, J., Commichau, F.M., and Stülke, J. (2018) Perspective of ions and messengers:
an intricate link between potassium, glutamate, and cyclic di-AMP. *Curr Genet* **64**:
191-195.
Gundlach, J., Herzberg, C., Hertel, D., Thurmer, A., Daniel, R., Link, H., and Stülke, J.
(2017a) Adaptation of *Bacillus subtilis* to Life at Extreme Potassium Limitation. *MBio*
8: e00861-00817.
Gundlach, J., Herzberg, C., Kaever, V., Gunka, K., Hoffmann, T., Weiss, M., Gibhardt, J.,
Thurmer, A., Hertel, D., Daniel, R., Bremer, E., Commichau, F.M., and Stülke, J.
(2017b) Control of potassium homeostasis is an essential function of the second
messenger cyclic di-AMP in *Bacillus subtilis*. *Sci Signal* **10**: eaal3011.
Gundlach, J., Krüger, L., Herzberg, C., Turdiev, A., Poehlein, A., Tascon, I., Weiß, M.,
Hertel, D., Daniel, R., Hänelt, I., Lee, V.T., and Stülke, J. (2019) Sustained sensing in
potassium homeostasis: Cyclic di-AMP controls potassium uptake by KimA at the
levels of expression and activity. *J Biol Chem* **294**: 9605-9614.
Gundlach, J., Mehne, F.M., Herzberg, C., Kampf, J., Valerius, O., Kaever, V., and Stülke, J.
(2015) An Essential Poison: Synthesis and Degradation of Cyclic Di-AMP in *Bacillus*
subtilis. *J Bacteriol* **197**: 3265-3274.
Halsey, C.R., Lei, S., Wax, J.K., Lehman, M.K., Nuxoll, A.S., Steinke, L., Sadykov, M.,
Powers, R., and Fey, P.D. (2017) Amino Acid Catabolism in *Staphylococcus aureus*
and the Function of Carbon Catabolite Repression. *MBio* **8**: e01434-01416.
Hengge, R. (2009) Principles of c-di-GMP signalling in bacteria. *Nat Rev Microbiol* **7**: 263-
273.
Huynh, T.N., Choi, P.H., Sureka, K., Ledvina, H.E., Campillo, J., Tong, L., and Woodward,
J.J. (2016) Cyclic di-AMP targets the cystathionine beta-synthase domain of the
osmolyte transporter OpuC. *Mol Microbiol* **102**: 233-243.
Kelly, B., and O'Neill, L.A. (2015) Metabolic reprogramming in macrophages and dendritic
cells in innate immunity. *Cell Res* **25**: 771-784.
Kim, H., Youn, S.J., Kim, S.O., Ko, J., Lee, J.O., and Choi, B.S. (2015) Structural Studies of
Potassium Transport Protein KtrA Regulator of Conductance of K+ (RCK) C Domain
in Complex with Cyclic Diadenosine Monophosphate (c-di-AMP). *J Biol Chem* **290**:
16393-16402.
Kluytmans, J., van Belkum, A., and Verbrugh, H. (1997) Nasal carriage of *Staphylococcus*
aureus: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol*
Rev **10**: 505-520.
Lehman, M.K., Nuxoll, A.S., Yamada, K.J., Kielian, T., Carson, S.D., and Fey, P.D. (2019)
Protease-Mediated Growth of *Staphylococcus aureus* on Host Proteins Is *opp3*
Dependent. *MBio* **10**: e02553-02518.
Mehne, F.M., Gunka, K., Eilers, H., Herzberg, C., Kaever, V., and Stülke, J. (2013) Cyclic di-
AMP homeostasis in *Bacillus subtilis*: both lack and high level accumulation of the
nucleotide are detrimental for cell growth. *J Biol Chem* **288**: 2004-2017.

Monk, I.R., Shah, I.M., Xu, M., Tan, M.W., and Foster, T.J. (2012) Transforming the
untransformable: application of direct transformation to manipulate genetically
Staphylococcus aureus and *Staphylococcus epidermidis*. *MBio* **3**: e00277-00211.
Monk, I.R., Tree, J.J., Howden, B.P., Stinear, T.P., and Foster, T.J. (2015) Complete Bypass
of Restriction Systems for Major *Staphylococcus aureus* Lineages. *MBio* **6**: e00308-
00315.
Moscoso, J.A., Schramke, H., Zhang, Y., Tosi, T., Dehbi, A., Jung, K., and Gründling, A.
(2015) Binding of Cyclic Di-AMP to the *Staphylococcus aureus* Sensor Kinase KdpD
Occurs via the Universal Stress Protein Domain and Downregulates the Expression
of the Kdp Potassium Transporter. *J Bacteriol* **198**: 98-110.
Pham, H.T., Nhiep, N.T.H., Vu, T.N.M., Huynh, T.N., Zhu, Y., Huynh, A.L.D., Chakrabortti,
A., Marcellin, E., Lo, R., Howard, C.B., Bansal, N., Woodward, J.J., Liang, Z.X., and
Turner, M.S. (2018) Enhanced uptake of potassium or glycine betaine or export of
cyclic-di-AMP restores osmoresistance in a high cyclic-di-AMP *Lactococcus lactis*
mutant. *PLoS Genet* **14**: e1007574.
Pham, H.T., and Turner, M.S. (2019) Onwards and [K(+)]upwards: a new potassium
importer under the spell of cyclic-di-AMP. *J Bacteriol* **201**: e00150-00119.
Pham, T.H., Liang, Z.X., Marcellin, E., and Turner, M.S. (2016) Replenishing the cyclic-di-
AMP pool: regulation of diadenylate cyclase activity in bacteria. *Curr Genet* **62**: 731-
738.
Quintana, I.M., Gibhardt, J., Turdiev, A., Hammer, E., Commichau, F.M., Lee, V.T., Magni,
C., and Stülke, J. (2019) The KupA and KupB proteins of *Lactococcus lactis* IL1403
are novel c-di-AMP receptor proteins responsible for potassium uptake. *J Bacteriol*
201: e00028-00019.
Richardson, A.R., Libby, S.J., and Fang, F.C. (2008) A nitric oxide-inducible lactate
dehydrogenase enables *Staphylococcus aureus* to resist innate immunity. *Science*
319: 1672-1676.
Rocha, R., Teixeira-Duarte, C.M., Jorge, J.M.P., and Morais-Cabral, J.H. (2019)
Characterization of the molecular properties of KtrC, a second RCK domain that
regulates a Ktr channel in *Bacillus subtilis*. *J Struct Biol* **205**: 34-43.
Römling, U. (2008) Great times for small molecules: c-di-AMP, a second messenger
candidate in Bacteria and Archaea. *Sci Signal* **1**: pe39.
Schuster, C.F., Bellows, L.E., Tosi, T., Campeotto, I., Corrigan, R.M., Freemont, P., and
Gründling, A. (2016) The second messenger c-di-AMP inhibits the osmolyte uptake
system OpuC in *Staphylococcus aureus*. *Sci Signal* **9**: ra81.
Schuster, C.F., Wiedemann, D.M., Kirsebom, F.C.M., Santiago, M., Walker, S., and
Gründling, A. (2019) High-throughput transposon sequencing highlights the cell wall
as an important barrier for osmotic stress in methicillin resistant *Staphylococcus*
aureus and underlines a tailored response to different osmotic stressors. *Mol*
Microbiol: DOI: 10.1111/mmi.14433.
Schuurman-Wolters, G.K., and Poolman, B. (2005) Substrate specificity and ionic regulation
of GlnPQ from *Lactococcus lactis*. An ATP-binding cassette transporter with four
extracytoplasmic substrate-binding domains. *J Biol Chem* **280**: 23785-23790.
Spahich, N.A., Vitko, N.P., Thurlow, L.R., Temple, B., and Richardson, A.R. (2016)
Staphylococcus aureus lactate- and malate-quinone oxidoreductases contribute to
nitric oxide resistance and virulence. *Mol Microbiol* **100**: 759-773.
Sureka, K., Choi, P.H., Precit, M., Delince, M., Pensinger, D., Huynh, T.N., Jurado, A.R.,
Goo, Y.A., Sadilek, M., Iavarone, A.T., Sauer, J.D., Tong, L., and Woodward, J.J.
(2014) The cyclic dinucleotide c-di-AMP is an allosteric regulator of metabolic
enzyme function. *Cell* **158**: 1389-1401.
Teh, W.K., Dramsi, S., Tolker-Nielsen, T., Yang, L., and Givskov, M. (2019) Increased
Intracellular Cyclic di-AMP Levels Sensitize *Streptococcus gallolyticus* subsp.
gallolyticus to Osmotic Stress and Reduce Biofilm Formation and Adherence on
Intestinal Cells. *J Bacteriol* **201**: e00597-00518.

Tolner, B., Ubbink-Kok, T., Poolman, B., and Konings, W.N. (1995) Characterization of the
925   proton/glutamate symport protein of *Bacillus subtilis* and its functional expression in
926   *Escherichia coli*. *J Bacteriol* **177**: 2863-2869.
927 Tosi, T., Hoshiga, F., Millership, C., Singh, R., Eldrid, C., Patin, D., Mengin-Lecreulx, D.,
928   Thalassinos, K., Freemont, P., and Gründling, A. (2019) Inhibition of the
929   *Staphylococcus aureus* c-di-AMP cyclase DacA by direct interaction with the
930   phosphoglucosamine mutase GlmM. *PLoS Pathog* **15**: e1007537.
931 Underwood, A.J., Zhang, Y., Metzger, D.W., and Bai, G. (2014) Detection of cyclic di-AMP
932   using a competitive ELISA with a unique pneumococcal cyclic di-AMP binding
933   protein. *J Microbiol Methods* **107**: 58-62.
934 Vitko, N.P., Spahich, N.A., and Richardson, A.R. (2015) Glycolytic dependency of high-level
935   nitric oxide resistance and virulence in *Staphylococcus aureus*. *MBio* **6**: e00045-
936   00015.
937 Whiteley, A.T., Garelis, N.E., Peterson, B.N., Choi, P.H., Tong, L., Woodward, J.J., and
938   Portnoy, D.A. (2017) c-di-AMP modulates *Listeria monocytogenes* central
939   metabolism to regulate growth, antibiotic resistance and osmoregulation. *Mol*
940   *Microbiol* **104**: 212-233.
941 Whiteley, A.T., Pollock, A.J., and Portnoy, D.A. (2015) The PAMP c-di-AMP Is Essential for
942   *Listeria monocytogenes* Growth in Rich but Not Minimal Media due to a Toxic
943   Increase in (p)ppGpp. *Cell Host Microbe* **17**: 788-798.
944 Wicke, D., Schulz, L.M., Lentes, S., Scholz, P., Poehlein, A., Gibhardt, J., Daniel, R.,
945   Ischebeck, T., and Commichau, F.M. (2019) Identification of the first glyphosate
946   transporter by genomic adaptation. *Environ Microbiol* **21**: 1287-1305.
947 Witte, C.E., Whiteley, A.T., Burke, T.P., Sauer, J.D., Portnoy, D.A., and Woodward, J.J.
948   (2013) Cyclic di-AMP is critical for *Listeria monocytogenes* growth, cell wall
949   homeostasis, and establishment of infection. *MBio* **4**: e00282-00213.
950 Woodward, J.J., Iavarone, A.T., and Portnoy, D.A. (2010) c-di-AMP secreted by intracellular
951   *Listeria monocytogenes* activates a host type I interferon response. *Science* **328**:
952   1703-1705.
953 Zarrella, T.M., Metzger, D.W., and Bai, G. (2018) Stress Suppressor Screening Leads to
954   Detection of Regulation of Cyclic di-AMP Homeostasis by a Trk Family Effector
955   Protein in *Streptococcus pneumoniae*. *J Bacteriol* **200**: e00045-00018.
956 Zeden, M.S., Schuster, C.F., Bowman, L., Zhong, Q., Williams, H.D., and Gründling, A.
957   (2018) Cyclic di-adenosine monophosphate (c-di-AMP) is required for osmotic
958   regulation in *Staphylococcus aureus* but dispensable for viability in anaerobic
959   conditions. *J Biol Chem* **293**: 3180-3200.
960 Zhu, B., and Stülke, J. (2018) SubtiWiki in 2018: from genes and proteins to functional
961   network annotation of the model organism *Bacillus subtilis*. *Nucleic Acids Res* **46**:
962   D743-D748.
963 Zhu, Y., Pham, T.H., Nhiep, T.H., Vu, N.M., Marcellin, E., Chakrabortti, A., Wang, Y.,
964   Waanders, J., Lo, R., Huston, W.M., Bansal, N., Nielsen, L.K., Liang, Z.X., and
965   Turner, M.S. (2016) Cyclic-di-AMP synthesis by the diadenylate cyclase CdaA is
966   modulated by the peptidoglycan biosynthesis enzyme GlmM in *Lactococcus lactis*.
967   *Mol Microbiol* **99**: 1015-1027.
968 Zhu, Y., Xiong, Y.Q., Sadykov, M.R., Fey, P.D., Lei, M.G., Lee, C.Y., Bayer, A.S., and
969   Somerville, G.A. (2009) Tricarboxylic acid cycle-dependent attenuation of
970   *Staphylococcus aureus* in vivo virulence by selective inhibition of amino acid
971   transport. *Infect Immun* **77**: 4256-4264.
972
973