Decoding Odor Mixtures in the Dog Brain: An Awake fMRI Study

Ashley Prichard[1], Raveena Chhibber[1], Jon King[1], Kate Athanassiades[1], Mark Spivak[2,3], & Gregory
S. Berns[1*]

[1]Psychology Department, Emory University, Atlanta, GA 30322
[2]Comprehensive Pet Therapy, Inc., Sandy Springs, GA 30328
[3]Dog Star Technologies, LLC, Sandy Springs, GA 30328
[*]Corresponding Author: Gregory S. Berns (gberns@emory.edu)

Funding

This work was supported by the Office of Naval Research (N00014-16-1-2276). ONR provided
support in the form of salaries [RC, MS, GSB], scan time, and volunteer payment, but did not
have any additional role in the study design, data collection and analysis, decision to publish, or
preparation of the manuscript.

Competing Interests

G.B. & M.S. own equity in Dog Star Technologies and developed technology used in some of the
research described in this paper. The terms of this arrangement have been reviewed and
approved by Emory University in accordance with its conflict of interest policies.

Acknowledgments

Thank you to all of the owners who trained their dogs to participate in fMRI studies: Lorrie
Backer, Rebecca Beasley, Emily Chapman, Darlene Coyne, Vicki D'Amico, Diana Delatour, Jessa
Fagan, Marianne Ferraro, Anna & Cory Inman, Patricia King, Cecilia Kurland, Claire & Josh
Mancebo, Patti Rudi, Cathy Siler, Lisa Tallant, Nicole & Sairina Merino Tsui, Ashwin
Sakhardande, & Yusuf Uddin.

Author Contributions

24
25 A.P., M.S., and G.B. designed the research; A.P., R.C., K.A. and G.B. collected the data; A.P, M.S.
26 and G.B trained the dogs, A.P., R.C., J.K., and G.B. analyzed data; and A.P., R. C. and G.B. wrote
27 the paper.
28

29   **ABSTRACT**

In working and practical contexts, dogs rely upon their ability to discriminate a target odor from

distracting odors and other sensory stimuli. Few studies have examined odor discrimination

using non-behavioral methods or have approached odor discrimination from the dog's

perspective. Using awake fMRI in 18 dogs, we examined the neural mechanisms underlying

odor discrimination between two odors and a mixture of the odors. Neural activation was

measured during the presentation of a target odor (A) associated with a food reward, a

distractor odor (B) associated with nothing, and a mixture of the two odors (A+B). Changes in

neural activation during the presentations of the odor stimuli in individual dogs were measured

over time within three regions known to be involved with odor processing: the caudate nucleus,

the amygdala, and the olfactory bulbs. Average activation within the amygdala showed that

dogs maximally differentiated between odor stimuli based on the stimulus-reward associations

by the first run, while activation to the mixture (A+B) was most similar to the no-reward (B)

stimulus. To identify the neural representation of odor in the dog brain, we used machine

learning to predict the classification of the odor mixture into its components with 73%

accuracy. Using a random forest classifier, we found that the amygdala, piriform, and posterior

cingulate cortices were most important for the classification of odors in the dog brain.

Confusion matrices suggested that the dogs treated the odor mixture as a new configuration

rather than elemental components.

Keywords

Dog, odor, mixture, fMRI, RFC, MVPA

1. INTRODUCTION

49
50
For many working purposes, trained dogs are generally considered the most practical and
effective means of identifying target substances. In many cases, detection dogs are selectively
bred for olfactory capabilities and behavioral traits that are correlated with their effectiveness
in the field. Given their roles in national security and in detecting different diseases, hunting for
bed bugs and pests, and tracking endangered species for conservation efforts (Bijland, Bomers,
& Smulders, 2013; Cooper, Wang, & Singh, 2014; Davidson, Clark, Johnson, Waits, & Adams,
2014; Gadbois & Reeve, 2014), odor detection dogs remain in high demand. Despite the
growing need for odor detection dogs, current understanding of dogs' abilities falls into three
domains: physiology, rearing and training practices, and chemical detection. Research regarding
dogs' olfactory abilities typically focuses on the improvement of detection behaviors and
trainability. Despite numerous behavioral studies, little is known about the way in which
olfactory information is interpreted by the canine brain. Few studies on canine olfaction
approach the topic from the canine's point of view or without responses mediated by the dog's
handler. While behavior is a necessary measure of a working dog's effectiveness, a dog's
behavior can be biased by unconscious cues given by their handler.
66
Large gaps remain in our understanding of how dogs process odors or discriminate between
pure odors and their mixtures. For instance, it is unknown whether dogs search for the
complete odor signature of a target substance, or if only one or some components serve as a
target odor (Johnen, Heuwieser, & Fischer-Tenhagen, 2017). Despite substantial training on
odor components, a dog's behavioral responses to mixtures often cannot be predicted. This
may be because the detection of individual substances within a mixture depends on chemical
interactions between the different components. Given that most odor discrimination tests for
dogs are behaviorally based and/or unstandardized, it is almost impossible to predict which
components of an odor a particular dog uses to identify the target (Göth, McLean, & Trevelyan,
2003). For example, dogs that were trained to detect pure potassium chlorate failed to reliably
detect potassium chlorate-based explosive mixtures (Lazarowski & Dorman, 2014). This finding,
and others, highlight the potential limitations of training dogs to detect a specific target odor to
then indicate to the target when mixed with distractors (DeGreeff et al., 2017; Hayes,
McGreevy, Forbes, Laing, & Stuetz, 2018). The way in which this information is interpreted by
the canine brain also remains under-researched, but it is likely a complex and contextually
dependent process (Berns, Brooks, & Spivak, 2015; Hayes et al., 2018; Prichard, Chhibber,
Athanassiades, Spivak, & Berns, 2018; Siniscalchi, 2016). Considering that olfactory
neuroanatomy is highly conserved among animals, studies of olfactory processing in dogs may
also shed light on similar mechanisms in humans (Ache & Young, 2005).
86
The brain may have specialized representations for olfactory associations (Yeshurun, Lapid,
Dudai, & Sobel, 2009). In humans, studies of odor perception typically rely on self-report
measures and use suprathreshold odors that are easily detectible. Functional magnetic
resonance imaging (fMRI) of an olfactory matching or identification task has demonstrated
activation in the primary and secondary olfactory regions including: the piriform cortex, insula,
amygdala, parahippocampal gyrus, caudate nucleus, inferior frontal gyrus, middle frontal gyrus,

superior temporal gyrus, and cerebellum (Vedaei et al., 2017). Low level odors that go
unnoticed by participants can also alter brain activation in the piriform cortex and thalamus
(Lorig, 2012). Most of these studies have contributed to the identification of odor processing
regions, but fewer have identified the regions' roles during odor processing or learning during
conditioning to odor stimuli. Regions that are said to support conditioned associations to odor
stimuli include the orbitofrontal and perirhinal cortices when learning odor categories. Even on
the first day of conditioning, activity within the hippocampus predicted the olfactory
associations that would be remembered one week later (Howard, Kahnt, & Gottfried, 2016; Qu,
Kahnt, Cole, & Gottfried, 2016).
102
More recent studies of human olfactory perception have implemented machine learning
strategies to decode odor representation within the brain. FMRI decoding methods can reveal
regions important for coding valence, expected outcomes, or stimulus identity. By using
representational methods, machine learning approaches such as multi-voxel pattern analysis
(MVPA) identify patterns of activation from regions that could fail to show a change in mean
activation measured by a typical univariate approach (Haxby, Connolly, & Guntupalli, 2014;
Kahnt, 2018; Kahnt, Park, Haynes, & Tobler, 2014). For instance, representational similarity
analysis (RSA) suggests that the spatial and temporal pattern of activation within the amygdala
codes for odor valence (Jin, Zelano, Gottfried, & Mohanty, 2015).
112
While these studies report similar regions important for odor discrimination identified in
traditional univariate fMRI analyses, the relationship of odor mixtures to the brain's
representation of odor components remains unknown (Howard & Gottfried, 2014; Howard,
Gottfried, Tobler, & Kahnt, 2015; Howard et al., 2016). Odor mixtures may be represented in
the brain based on their components (elemental) or may be perceived as configural, creating an
odor concept (Thomas-Danguin et al., 2014). FMRI studies on human perception of odor
mixtures shows that activation in the insula increases when the participant experiences the
mixture containing the target odor, even when participants report that they are unable to
distinguish the mixture with and without the target (Hummel, Olgun, Gerber, Huchel, &
Frasnelli, 2013). However, other regions identified in this study included voxel sizes that would
not pass whole brain corrections for multiple comparisons, requiring further study to confirm
these brain regions' roles in the perception of odor mixtures. Despite the need for the study of
human olfactory perception at the neural level, no research has yet investigated similar
considerations in the dog (Hayes et al., 2018).
127
Studies of canine cognition using fMRI are becoming more common, including the adaptation of
human experimental paradigms and analyses. Dogs are willing participants in fMRI studies and
show little anxiety in the testing environment as it is similar to their shared environment with
humans. Due to domestication, dogs are also more likely attuned to stimuli relevant to humans
as opposed to stimuli salient to other model species. Since 2012, dog fMRI studies have
revealed some of the conserved neural mechanisms underlying perception across species
(Berns, Brooks, & Spivak, 2012). Dogs have a region for processing both human and dog faces
similar to that of primates (Cuaya, Hernandez-Perez, & Concha, 2016; Dilks et al., 2015). Dogs
show differential activation in the reward processing regions of the brain such as the caudate

nucleus to social or food rewards (Cook, Prichard, Spivak, & Berns, 2016). And dogs show higher
activation in the amygdala and caudate to odors associated with familiar humans and dogs than
to odors of strangers (Berns et al., 2015). Canine fMRI studies have also revealed neural biases
for stimulus modalities, suggesting that dogs learn visual and odor stimuli at a faster rate than
verbal stimuli, and that differences in activation are most evident in the amygdala and caudate
(Prichard, Chhibber, et al., 2018). Finally, MVPA analysis of dog fMRI data revealed that dogs
and humans have similar brain regions for the representation of semantic knowledge in the
form of words associated with objects (Prichard, Cook, Spivak, Chhibber, & Berns, 2018).
Together, these studies suggest that dogs are not only willing fMRI participants, but that the
existence of functionally similar brain regions shared by dogs and humans make them an
appropriate model species for further research.
148
To examine the neural mechanisms underlying a dog's classification of odor mixtures, in the
current study, we measured the fMRI response to two previously trained odors (Prichard,
Chhibber, et al., 2018) and a mixture of the two odors (A+B). If a dog represents a mixture as
the sum of its parts, then, assuming linearity, the change in activation within the amygdala and
caudate to the mixture (A+B) should be equal to the combined activation to reward (A) and no
reward (B). However, if a dog treats a mixture as a novel substance different from the sum of its
components, or odor "C", then the activation within the two regions will differ from reward and
no reward. Alternatively, rather than resulting in changes in activation level, a mixture might
result in a change in representational pattern. We also used random forest classifiers (RFC) to
identify regions in the brain that contribute to odor identification and whether they classify a
mixture more like its components or as a novel substance.
160

2. MATERIALS AND METHODS

162 *2.1 Participants*
163
Participants were 18 pet dogs volunteered by their Atlanta owners for fMRI training and fMRI
studies (Berns, Brooks, & Spivak, 2013; Berns et al., 2012; Berns & Cook, 2016; Cook et al.,
2016). All dogs had previously completed four or more awake fMRI scans, including previous
training on the two odors used in the current study (Prichard, Chhibber, et al., 2018). No
physical or chemical restraint was implemented. The study utilized odor stimuli that each dog
had previously experienced within the scanner environment. This study was performed in
accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals
of the National Institutes of Health. The study was approved by the Emory University IACUC
(Protocols DAR-4000079-ENTPR-A and PROTO201700572), and all owners gave written consent
for their dog's participation in the study.
174
175 *2.2 Stimuli*
176
Olfactory stimuli were aqueous solutions of isoamyl acetate (IA), hexanol (Hex), and a mixture
of the two calculated to result in approximately 5 ppm in the headspace of the container.
Partial vapor pressures were calculated based on the molecular weight and reported vapor
pressures of 4 mmHg and 0.9 mmHg respectively, obtained from PubChem

(pubchem.ncbi.nlm.nih.gov). The odorants were miscible
with water and the partial pressure of the odorant was
the product of the pure odorant vapor pressure and the
mole fraction of the odorant. The final dilutions in water
were 0.12 mL/L for IA, 0.44 mL/L for Hex.
186
Odorants were delivered using an MRI-compatible
olfactometer used in a previous study and similar to
those constructed for human olfactory imaging studies
(Bestgen et al., 2016; Lowen & Lukas, 2006; Prichard,
Chhibber, et al., 2018; Sezille et al., 2013; Sommer et al.,
2012; Toledano et al., 2012; Vigouroux, Bertrand, Farget,
Plailly, & Royet, 2005). Briefly, odorants were delivered
using a continuous stream of air from an aquarium grade
air pump (EcoPlus Commercial Air Pump 1030 GPH)
through a Drierite filter (drierite.com), and through a 4-
way plastic splitter to three plastic 100 mL jars containing
50 ml of odorant solutions and one jar containing 50 ml
of water to serve as a control. Each solution mixed with a
continuous air stream. The experimenter used plastic
valves to control directional flow of odorized air through
10' of 1/8" ID Teflon tube, where the mixture (air dilution
of the odorant) exited a PVC tube with a 1" diameter

Fig 1. Experimental design with odor stimuli. Three odor stimuli were repeatedly presented during a scan session. One stimulus was associated with food (Reward), while the No Reward and Mixture stimuli were associated with nothing. Presentation of odorants to dog in MRI bore via experimenter-controlled olfactometer during scan session. The owner remained in front of the dog.

204   opening positioned in the MRI bore 12" from the dog's snout (Fig. 1). The fourth tube carrying
205   air from the control jar remained open throughout the presentations of odorized air,
206   maintaining a steady air stream presented to the dog and assisting in the clearing of lingering
207   odor within the magnet bore.
208
2.3 Experimental Design
Dogs entered and stationed themselves in custom chin rests in the scanner bore. All scans took
place in the presence of the dog's primary owner, who stood throughout the scan at the
opening of the magnet bore, directly in front of the dogs, and delivered all rewards (hot dogs)
to the dog. The owner was present to minimize any anxiety that the dog may experience due to
separation, consistent with studies involving pets or human infants. An experimenter was
stationed next to the owner, out of view of the dog. The experimenter controlled the timing
and presentation of odor stimuli to the dogs via a four-button MRI-compatible button box.
Onset of each stimulus was timestamped by the simultaneous press of the button box with the
opening of the appropriate valve. Manual control of the stimuli by the experimenter was
necessary, as opposed to a scripted presentation, because of the variable time it takes dogs to
consume food rewards.
221
In a previous study, dogs were semi-randomly assigned IA or Hex as the reward stimulus such
that roughly half of the dogs were assigned to each group (see Table 1) (Prichard, Chhibber, et

al., 2018). In the current study, the same dogs were presented with the two previously trained
odors, as well as a mixture of the two. An event-based design was used, consisting of reward,
no-reward, and mixture trial types. On reward trials, the odor stimulus was presented for a
fixed duration, which was followed by the delivery of a food reward. During no-reward trials
and mixture trials, the no-reward or mixture odor stimuli were presented for the same fixed
duration and were followed by nothing. Each dog received the same trial sequence. For each
trial type, dogs were presented an odor for an initial 3.6s during a span of 7.2 s, followed by a
reward (hot dog) or nothing, with a 9.6 s inter trial interval between odor presentations.
232
233 **Table 1. Dogs (N=18) and odor stimulus paired with reward.**

Dog	Breed	Sex	Reward Odor
BhuBo	Boxer mix	M	hexanol
Caylin	Border collie	F	hexanol
Daisy	Pitbull mix	F	hexanol
Eddie	Labrador Golden mix	M	isoamyl acetate
Kady	Labrador	F	hexanol
Koda	Pitbull mix	F	isoamyl acetate
Libby	Pitbull mix	F	hexanol
Mauja	Cattle dog mix	F	hexanol
Ninja	Cattle dog mix	F	isoamyl acetate
Ohana	Golden Retriever	F	hexanol
Ollie	Border collie Beagle mix	M	isoamyl acetate
Pearl	Golden Retriever	F	hexanol
Tallulah	Cattle Dog mix	F	hexanol
Truffles	Pointer mix	F	isoamyl acetate
Tug	Portuguese Water dog	M	hexanol
Velcro	Viszla	M	isoamyl acetate
Wil	Australian Shepherd	M	isoamyl acetate
Zen	Labrador Golden mix	M	isoamyl acetate

234 Dog's names, breed, sex, and odor stimuli (S+) are listed
235
236 Each scan session consisted of 4 runs, lasting approximately 9 minutes per run. Each run
237 consisted of 22 trials (~8 reward, ~8 no-reward, ~5 mixture) with a semi-randomized
238 presentation order, for a total of 88 trials per scan session. No trial type was repeated more
239 than 3 times sequentially, as dogs could habituate to the stimulus. Following each run, dogs
240 would exit the scanner and relax, drink water, or stay in the scanner to complete the next run.
241
Scanning was conducted with a Siemens 3 T Trio whole-body scanner using procedures
described previously (Berns et al., 2013; Berns et al., 2012). During the first of the three scans
sessions, a T2-weighted structural image of the whole brain was acquired using a turbo spin-
echo sequence (25-36 2mm slices, TR = 3940 ms, TE = 8.9 ms, flip angle = 131˚, 26 echo trains,
128 x 128 matrix, FOV = 192 mm). The functional scans used a single-shot echo-planar imaging

(EPI) sequence to acquire volumes of 22 sequential 2.5 mm slices with a 20% gap (TE = 25 ms,
TR = 1200 ms, flip angle = 70˚, 64 x 64 matrix, 3 mm in-plane voxel size, FOV = 192 mm). Slices
were oriented dorsally to the dog's brain (coronal to the magnet, as in the sphinx position the
dogs' heads were positioned 90 degrees from the prone human orientation) with the phase-
encoding direction right-to-left. Sequential slices were used to minimize between-plane offsets
from participant movement, while the 20% slice gap minimized the "crosstalk" that can occur
with sequential scan sequences. Four runs of up to 400 functional volumes were acquired for
each subject, with each run lasting about 9 minutes.
255
3. Analyses
3.1 Preprocessing

Preprocessing of the fMRI data included motion correction, censoring, and normalization using
AFNI (NIH) and its associated functions. Two-pass, six-parameter rigid-body motion correction
was used based on a hand-selected reference volume for each dog that corresponded to their
average position within the magnet bore across runs. Aggressive censoring removed unusable
volumes from the fMRI time sequence because dogs can move between trials, when smelling
an odor, and when consuming rewards. Data were censored when estimated motion was
greater than 1 mm displacement scan-to-scan and based on outlier voxel signal intensities
greater than 0.1 percent signal change from scan-to-scan. Smoothing, normalization, and
motion correction parameters were identical to those described in previous studies (Prichard,
Chhibber, et al., 2018). EPI images were smoothed and normalized to %-signal change with
3dmerge using a 6mm kernel at full-width half-maximum. The Advanced Normalization Tools
(ANTs) software was used to spatially normalize the mean of the motion-corrected functional
images (Avants et al., 2011) to the individual dog's structural image.
271
3.2 General Linear Model

Each subject's motion-corrected, censored, smoothed images were analyzed within a general
linear model (GLM) for each voxel in the brain using 3dDeconvolve (part of the AFNI suite).
Motion time courses were generated through motion correction, and constant, linear,
quadratic, cubic, and quartic drift terms were included as nuisance regressors. Drift terms were
included for each run to account for baseline shifts between runs as well as slow drifts
unrelated to the experiment. Task related regressors for each experiment were modeled using
AFNI's dmUBLOCK and stim_times_IM functions and were as follows: (1) reward stimulus, (2)
no-reward stimulus, 3) mixture stimulus. The function created a column in the design matrix for
each of the 88 trials, allowing for the estimation of beta values for each trial. Trials with beta
values greater than an absolute three percent signal change were removed prior to analyses as
described in Prichard et al. (2018) as these were assumed to be beyond the physiologic range of
the BOLD signal and possibly the result of spin-history effects and spurious levels of activation
in specific regions of interest.
286
3.3 Region of Interest (ROI) Analysis

Anatomical ROIs were selected based on
imaging results in canine brain areas
previously observed to be responsive to
olfactory stimuli (Berns et al., 2015; Jia et
al., 2014). Anatomical ROIs of the left and
right caudate nuclei, the left and right
amygdala, and the olfactory bulbs were
defined structurally using each dog's T2-
weighted structural image of the whole
brain (Fig. 2). Beta values for each
presentation of reward stimuli (33 trials),
no-reward stimuli (33 trials), and mixture
stimuli (22 trials) were extracted from and
averaged over the ROIs in the left and
right hemispheres. For each ROI
(amygdala, caudate, olfactory bulb), we
used the mixed-model procedure in SPSS
24 (IBM) with fixed-effects for the
intercept, run number, type (reward, no-
reward, mixture), and hemisphere (left or
right), identity covariance structure, and
maximum-likelihood estimation. Run was
modeled as a fixed effect because it made
no assumptions about the time course. As
hemisphere did not account for a
significant amount of variance, all analyses
removed hemisphere as a factor.
315
316 *3.4 Multivariate Decoding*

Fig 2. Regions of interest (ROIs) defined a priori. ROIs were drawn in individual anatomical space, example ROIs shown in template space here in transverse and dorsal views. **A)** Caudate nuclei have been shown to differentially respond to odor stimuli associated with reward and no-reward. **B)** Amygdalae have shown differential responding to odor stimuli associated with reward and no-reward, as well as arousal. **C)** Olfactory bulbs including olfactory bulbs respond to odor stimuli. ROI is shown here in sagittal and dorsal views in template space.

For this exploratory analysis, our aim was to identify regions in the dog brain that contribute to
the classification of odor stimuli outside of those identified in the univariate analysis. The
question remains whether dogs treat odor mixtures as elemental (A & B) or configural (C).
Univariate analyses may answer the question if odor mixtures result in differences in regional
brain activity, but multivariate methods are required if the identity of odors is distributed in
patterns of neural activity. As in previous decoding human fMRI studies, we used scikit-learn's
random forest classifier (RFC). RFC has previously demonstrated robust performance on human
fMRI data and has the ability to handle complex biological data (Lebedev et al., 2014). RFCs
generally perform better than most linear classifiers and require less parameter tuning (Chollet,
2018). An RFC also allows for mapping of feature importance in the brain without resorting to
searchlight analyses. Thus, in addition to generating whole-brain classification metrics, the
relative importance of individual regions to the classification can be obtained.

The volumes from the current study were concatenated with data from the previous study in
which dogs were presented odors associated with reward and no reward in a classical
conditioning paradigm (Prichard, Chhibber, et al., 2018). As described in the above GLM,
preprocessing included censoring of the unsmoothed volumes for motion and outliers. Using
AFNI's 3dDeconvolve stim_times_IM function, we generated a model of trial-by-trial beta
estimates for each trial type (reward, no reward, and mixture). The anatomical masks from the
ROI analysis described above were used to extract average beta values from the left and right
caudate for each trial. As in the univariate analysis, trials with beta values greater than |3 %|
were removed prior to further analyses. Using AFNI's 3dmerge tool, the remaining whole brain
volumes were smoothed with a kernel of 6 mm to facilitate the identification of regions
important for classification of the odors. The whole brain volumes were used as input for the
classifiers below. To reformat the imaging data for use in the sklearn environment, the volumes
were masked and reshaped using nilearn's NiftiMasker class and split into training and testing
sets using the python library *pandas*.

Two different models were tested. First, to confirm the validity of the MVPA approach, we
trained the classifiers on the data from the original classical conditioning study where dogs
were presented with odors associated with reward or no reward (two-way model). This set was
balanced by selecting a random subset of an equal number of reward and no reward trials to
control for overfitting. The testing set was the current study where dogs were presented with
reward and no reward odors and the mixture. Thus, the training and test sets were obtained
from two completely different scanning sessions, separated by 5 months, yielding the most
stringent test of classification accuracy across potentially varying contexts. Second, to
determine whether the mixture of odors in the new experiment was processed elementally or
configurally, we used the concatenated dataset of both experiments to train and test on three
classes: a) reward odor; b) no reward odor; c) the mixture (three-way model). As above,
training sets were balanced in each dog to have the same number of exemplars. Training was
performed on even number runs, and testing was performed on the odd runs.

3.5 RFC

For both the two- and three-way models, the RFC was instantiated by making 100 forests, each
forest consisting of 100 trees with a max_depth of 5, min_samples_split of .25, bootstrapping
as true, and max_features as log2. We chose to use 100 forests of 100 trees to ensure that all
volumes served as samples. A max_depth of 5, min_samples_split of .25 and max_features of
log2 were included to prevent overfitting to the training set.

For each trial of the test set of the two-way model, the classifier predicted whether the
stimulus presented was reward or no reward. The classifier was not expected to predict mixture
trials because the training set only consisted of reward or no reward. Each forest also produced
a map of feature importances. Briefly, the feature importance is a value scaled between 0 and 1
that reflects how informative a voxel i.e. a larger feature importance corresponds to a voxel
that is more informative in making the final predictions. Higher feature importances are driven
by either voxels that increase accuracy drastically, or by voxels that are present in many trees
within a forest. Sklearn's RFC feature_importances_ method returned feature importances for

each voxel that were subsequently back-mapped into each individual dog's functional space,
generating one map per forest. All 100 maps for each dog were averaged in order to assess
which brain regions contributed to the classification of reward and no reward. Mean images for
each dog were spatially normalized to template space using The Advanced Normalization Tools
(ANTs) software (Avants et al., 2011; Datta et al., 2012). In making the group maps from these
mean images, we excluded dogs where the overall classifier was unable to predict reward and
no reward odors with accuracy above 65%, which corresponds to $p = 0.01$ in a chi-square test
for homogeneity. To test whether dogs learned that A+B = no reward (versus carrying forward a
representation from the first experiment), we measured classification accuracy over the course
of the experiment by run.

For the three-way model, the classifier predicted whether the stimulus presented was reward,
no reward, or mixture. This model was included because the two-way model cannot explain
whether odor is processed configurally or elementally. Maps of feature importance were
produced using the above method. All dogs were included.

3.6 Noise Data Simulation

To determine the significance of any voxels in the group images, we followed the permutation
approach outlined by Stelzer et al. (2013) and used previously to identify the significance of
regions for language processing in dogs (Prichard, Cook, et al., 2018; Stelzer, Chen, & Turner,
2013). For each dog, a random number was appended to the data labels for each trial to
reorder the labels and create one permuted list of labels, whereas the timeseries of fMRI
volumes was kept consistent. The labels were permuted to assess how the RFC performed with
noise data and to determine voxel and cluster significance. The RFC was trained and tested on
this set of permuted labels and the fMRI volumes 100 times, outputting one map of feature
importances per forest for a total of 100 forests and maps, as we did with our real dataset
above. We then averaged across these 100 maps, creating one mean permuted image per set
of permuted labels. We repeated this procedure 100 times to create a distribution of noise
feature importance maps for each dog. To simulate the group image across dogs, we randomly
selected one mean permuted image per dog, normalized that mean image to template space,
and averaged across dogs. This random selection and normalization were repeated 10,000
times. Because each voxel in the brain may have a different distribution given its location in the
brain, we did not assume a canonical distribution across all voxels and opted to make a voxel-
wise distribution. For each voxel in the brain, we created the distribution from the 10,000 noise
maps and determined the values for $p = 0.005$. This map of thresholds was applied to the mean
feature importance map created from the real data and to each of the 10,000 noise maps (Fig.
3).

To determine the significance of any clusters found after thresholding at the voxel-wise level,
we created a distribution of cluster sizes found in the thresholded 10,000 noise maps. All
cluster sizes with more than 2 voxels were used to create a cumulative distribution of possible
cluster sizes. Cluster sizes of 114 voxels and 285 voxels corresponded to $p = 0.05$ and $p = 0.01$
respectively. The final map of thresholded feature importances was converted to a map of p
values.

Fig 3. Schematic diagram of MVPA methods. A random forest classifier (RFC) was trained on a balanced subset of the real data, outputting a map of voxels important for classification. As the balanced subset was randomized, we repeated this process 100 times in order to ensure that all samples were used at least one. The maps from these 100 repetitions were averaged, normalized to group space, and thresholded at a voxel and cluster level to create the final image. To create the voxel and cluster-level thresholds, we created random data by permuting the data labels associated with each volume, then trained as described above for the real data; this process constituted one permutation. The data were permuted 100 times. One map was selected at random to transform into group space. We generated 10,000 random group maps, created a voxel-by-voxel distribution and a further cluster distribution, both applied to the image generated by the real data.

4. RESULTS

4.1 *Univariate*

Changes in neural activation during the presentations of the odor stimuli in individual dogs were measured over time within the three ROIs known to be involved with odor processing. Using the mixed-model procedure in SPSS 24 (IBM) we found neural evidence for differentiation of the three odor stimuli across all ROIs (p = 0.004), which varied significantly by Odor Type (p < 0.001). There was a significant interaction between Odor Type x Run (p = 0.031), suggesting the magnitude of the effect changed over time.

Fig 4. Percent signal change by ROI for odor stimuli. Unadjusted mean values across dogs by ROI and by trial type (*blue* = reward, *red* = no reward, *purple* = mixture of reward and no reward). Error bars denote the standard error of the mean across dogs for each type and ROI. Averaged beta values in the caudate show some differentiation between olfactory stimuli, with greater activation to odor stimuli associated with reward and deactivation to the no reward and mixture odors. Averaged beta values in the amygdala show marked differentiation between odor stimuli with the greatest activation to odor stimuli associated with reward. Averaged beta values in the olfactory bulbs follow a similar pattern of activation to those in the caudate. Across all ROIs, the neural activation to the mixture of odors is most similar to the neural response during the presentation of the no reward odor.

Table 2. Model results for Odor Type, Run, and ROI. Asterisks denote significant results.

ROI	Fixed Effects	Numerator df	Denominator df	*F*	*Sig.*
Caudate	Intercept	1	2580	0.265	0.606
	Run	3	2580	0.607	0.611
	Odor Type	2	2580	2.056	0.128
	Run * Odor Type	6	2580	2.529	0.019*
Amygdala	Intercept	1	2426	12.831	0.000*
	Run	3	2426	2.068	0.102
	Odor Type	2	2426	11.016	0.000*
	Run * Odor Type	6	2426	2.37	0.028*
Olfactory Bulbs	Intercept	1	1296	0.143	0.706
	Run	3	1296	0.592	0.62
	Odor Type	2	1296	3.539	0.029*
	Run * Odor Type	6	1296	0.746	0.613

440
As there was a main effect of ROI, we used post-hoc analyses to examine whether these
differences remained when segregated by ROI (Table 2 & Fig. 4). In the caudate, we found a
significant interaction between Odor Type x Run (p = 0.019) (Fig. 5A), but no main effect of
Odor Type or Run. More robust evidence for the differentiation between odor stimuli was
evident in the amygdala for Odor Type (p < 0.0001) (Fig. 5B), suggesting that the odor-outcome
associations were reinstated from the previous study. There was also an Odor Type x Run
interaction, suggesting a difference in the temporal pattern between odor types (p = 0.028).
Similar to human olfaction studies, we found initial evidence for the differentiation of Odor
Type in the olfactory bulbs (p = 0.029) (Fig. 5C).
450
In sum, the differences in neural activation across regions of the olfactory pathway show that
dogs formed odor stimulus-reward associations. Though the differentiation between the three
odor stimuli was most pronounced in the amygdala, similarity in activation between the no
reward and mixture stimuli across all three ROIs suggested that the mixture was most like the
no reward stimulus. Critically, when we tested whether the sum of activations to reward and no
reward odors was the same as the activation to the mixture, we found significant differences in
the amygdala, such that the sum of activations was greater than activation to mixture (t(17) =

Fig 5. Percent signal change by ROI for each contrast. Unadjusted mean values across dogs by run and by odor (*red* = Reward— No Reward, *purple* = Mixture— No Reward) and averages across all runs. Error bars denote the standard error of the mean across dogs for each contrast and run. Consistent with studies of odor discrimination, there were main effects of odor type across all ROIs (p = 0.004), which was significantly different by odor type (p < 0.001). There was a significant interaction ROI and Run (p = 0.031), suggesting the magnitude of the effect changed over time. **A.)** Averaged beta values in the caudate show a significant interaction between Run and Odor Type (p = 0.036). **B.)** Averaged beta values in the amygdala show marked effects of Odor Type (p = 0.001) and the intercept (p = 0.001). **C.)** Following corrections for multiple comparisons, averaged beta values in the olfactory bulbs were not significantly above chance. While the caudate and olfactory bulbs show little differential activation to olfactory stimuli as an effect of time, the amygdala contributes to the discrimination between reward, no reward, and mixture stimuli over time, beginning after the first run.

458 3.28, *p* = 0.004). This suggests that mixture was, in fact, processed differently than the simple
459 sum of its components. To further test this theory, multivariate decoding was performed.
460
461
462 *4.2 Multivariate Decoding*
463
Because of the limitations and
assumptions of the univariate
analysis, we used MVPA to
explore potential regions in the
whole brain that may code for
different representations of the
odor stimuli. Our classifier was
consistent across most dogs,
however five dogs (Caylin, Libby,
Tallulah, Velcro, & Wil) were
excluded from further group
analyses as the two-way classifier
was unable to predict the two
classes with an accuracy above
65%. In the remaining thirteen
dogs, the two-way model was
able to predict reward and no
reward with accuracy above 65%.
At the voxel-wise threshold of *p* =
0.005 and at the cluster
threshold of *p* = 0.05, we found
three regions with clusters of
informative voxels that decode

Fig 6. Group Mean Thresholded Map of Informative Voxels.
Average feature importance was converted to *p* values for the
thirteen dogs with reward and no reward classification accuracy
above 65%. Color indicates voxel-wise significance for each voxel
(red: *p* < 0.0001 and yellow: *p* < 0.001). The image is clusterized at
114 voxels (*p* = 0.05). This map shows three clusters of informative
voxels found in the posterior cingulate, right piriform and the right
amygdala.

odor representations in the dog brain; posterior cingulate (172 voxels, *p* = 0.026), right piriform
(155 voxels, *p* = 0.031), and right amygdala (130 voxels, *p* = 0.041) (Fig. 6). For this two-way
model, we evaluated performance using accuracy in predicting reward and no reward trials. As
there were no mixture trials in the training set, the classifier did not predict mixture. The
tendency of the RFC to classify mixture trials as no reward suggests that the mixture was
neurally represented more similarly to the no reward odor than the reward odor (Table 3). To
test whether classification accuracy changed over the course of the experiment (suggesting that
the difference was learned during the experiment), we examined the average accuracy in
predicting reward and no reward split by run (Fig. 7). There was not a significant effect of run
on classification rates for average submatrix accuracy (F(54) = 1.196, p = 0.32) or submatrix
classification of no reward (F (54) = 0.238, p = 0.869), suggesting that the RFC was able to
discriminate between the stimuli within the first run and did not change after that.
499

Table 3. RFC Confusion Matrix results for Two Way Model for All Dogs. Trials aggregated over 100 forests (percentages)

		Predictions		
		Mixture	Reward	No Reward
Targets	Mixture	0*	7540 (23.6)	24460(76.4)
	Reward	0*	21358 (56.4)	16542(43.6)
	No Reward	0*	11268(23.2)	37332(76.8)

*RFC never predicts mixture as training only consisted of Reward and No Reward

Given the classifier's good performance in the two-way model, we then created a three-way model to elucidate whether neural processing of odor mixtures was configural or elemental. Here, mixture trials were most often categorized as mixture (Table 4), suggesting that they may be processed configurally. We did not find any additional brain regions that support elemental processing of odors using maps of feature importances.

Table 4. RFC Confusion Matrix results for Three Way Model for All Dogs. Trials aggregated over 100 forests (percentages)

		Predictions		
		Mixture	Reward	No Reward
Targets	Mixture	7516 (47.2)	2479 (15.6)	5905 (37.1)
	Reward	24869(28.4)	36653 (41.9)	25978(29.7)
	No Reward	42686(41)	20611(19.8)	40903(39)

A

B

513
514

5. Discussion

Here, we show fMRI evidence that dogs' brains tended to classify odor mixtures
configurally. To test neural mechanisms of dogs' perception of odors and a mixture, we used
fMRI to examine changes in brain activation to previously trained odors associated with reward
(A) or no reward (B), as well as a mixture of the two (A+B). Our results suggest that while a dog
may be familiar with or have different odor-outcome associations with each individual odor,
dogs perceive the combination of odors as a new odor (C). In reward processing regions of the
brain, we anticipated that if dogs treat mixtures as the sum of their components, then the
neural activation to the mixture (A+B) should be equivalent to the sum of the activation to the
reward (A) and no reward (B) components. However, significant differences in activation within
the amygdala showed that dogs did not treat these as equivalent conditions.
Further, using machine learning, we identified additional regions of the dog brain,
including the posterior cingulate and piriform cortex, that significantly predicted the identity of
the odor beyond the regions specified in *a priori* hypotheses. Given a two-way model with the
classes of reward and no reward, the RFC predicted that a mixture of odors is more like the no
reward odor. However, when three classes were given (reward, no reward, or mixture), the RFC
predicted the mixture predominately as itself, and secondly as no reward. The three-way model
supports configural processing of odor mixtures in the dog brain while the two-way model
supports the univariate analyses that dogs perceive a mixture as no reward.

533   One possible explanation for our results is that the dogs' perception of odor mixtures
534 may depend on the combined ratio of the odor elements. For example, rabbits trained on a
535 target odor B treated the A+B (ratio 68/32) mixture as elemental but the A+B (ratio 30/70)
536 mixture as configural (Schneider et al., 2016). As our study utilized a 50/50 mixture of A+B, we
537 cannot similarly conclude ratio-based differences in elemental or configural processing of odor
538 mixtures in the dog brain. However a second possible explanation is that dogs classify mixtures
539 as themselves as in the 3-way model, but when limited to two classes as in the 2-way model,
540 dogs' neural biases for novelty influences predictions toward the distractor odor (Prichard,
541 Cook, et al., 2018).
542   Because both the univariate and two-way RFC suggested that dogs treat mixtures as
543 more like the no reward stimulus than the reward stimulus, the mechanism underlying dogs'
544 discrimination of odor mixtures may have been a learned association between the mixture with
545 absence of reward. The apparent differences in activation in the caudate nucleus and amygdala
546 to odor stimuli associated with reward or no reward suggested that perception changed over
547 time, consistent with a learned discrimination. The significant differential effect for reward
548 versus no-reward across multiple ROIs is therefore consistent with prior research, as we have
549 shown that reward processing regions of the canine brain such as the striatum change in
550 activation relative to the value of conditioned stimuli regardless of modality (Cook et al., 2016;
551 Prichard, Chhibber, et al., 2018). Further, our lab has previously shown that in an fMRI
552 associative reward-learning paradigm, rapid changes in the neural activation within the caudate
553 and amygdala indicate conditioning to reward and no reward odors within an initial span of 6
554 minutes, such that a mixture-no reward association can form quickly (Prichard, Chhibber, et al.,
555 2018). In contrast to the significant effect of run in the univariate analysis, we did not find
556 significant changes in classification rates with the RFC. This suggest that the identity of the
557 odors and the mixture were separated by the first run. The overall activations in the amygdala
558 and caudate, then, might simply index their relative salience, but not their full identities.
559   The RFC identified regions important for odor processing similar to those in the human
560 studies, including the amygdala, posterior cingulate, and piriform. In human classical
561 conditioning paradigm using odors, MVPA analyses revealed predictive representations of
562 identity-specific reward in OFC and identity general reward in vmPFC. Reward related
563 functional coupling between OFC and piriform cortex and between vmPFC and amygdala
564 further revealed parallel pathways that support identity-specific and general predictive
565 signaling (Howard et al., 2015; Howard et al., 2016; Zelano, Mohanty, & Gottfried, 2011). Our
566 study also mirrors some of the results examining human's perception of odor mixtures. In
567 humans, common neural activation patterns in the superior temporal gyrus, caudate nucleus,
568 and insula occur in response to mixtures containing pleasant and unpleasant odors (Bensafi et
569 al., 2012). Like humans, dogs demonstrated differential activation in the caudate nucleus to a
570 mixture of odor components associated with different outcomes. Given the similar results to
571 human studies, this suggests that shared neural mechanisms may exist across species for odor
572 processing. Further, we show that RFC is a successful classifier for fMRI analyses, though as our
573 study was a simple design, with the caution that specific classifiers may be better suited for
574 some studies over others (Misaki, Kim, Bandettini, & Kriegeskorte, 2010).
575   What does this mean for odor processing in dogs? Understanding how a dog
576 discriminates between odor mixtures can aid in the design of more effective protocols to

increase a dog's performance on odor detection and identification tasks. Protocols designed
based on the dogs' perceptual abilities are less prone to biases inherent to behavioral studies
(e.g. the Clever Hans effect) that require human-reported measures. In addition, the dogs'
perception of the mixture stimulus in our study suggest that dogs perceive the mixture as a new
odor rather than as its individual elements. Consistent with previous behavioral studies, this
may explain why dogs trained on individual target odors have difficulty generalizing to mixtures
(Lazarowski & Dorman, 2014; Lazarowski et al., 2015). Further, dogs' brain activations showed
more similarity between the mixture of odors and a no reward odor, suggesting either a
learned association or a neural bias toward the no reward (distractor) odor. Treating a mixture
as a novel odor or having bias toward the no reward component of a mixture would likely lead
to increased false-negatives during a detection task whereas a learned association for mixtures
may conflict with detection applications. Knowledge of dogs' classification of odor mixtures in
the dog brain should improve training practices for working dogs and highlight the potential
learning aspects inherent in mixture detection tasks. Perceptually driven protocols may
therefore enhance a working dog's detection performance, contributing to the health and
safety of humans.
The opportunity to study the neural mechanisms of odor processing in an awake dog
also offers two clear advantages over the study of odor processing in humans. First, unlike
human studies, dog fMRI offers a unique opportunity to study odor processing in primary
sensory areas like the olfactory bulbs given its large size relative to the rest of the dog brain. In
humans, the olfactory bulbs is proportionately smaller than in canines, making imaging difficult
due to its size and the susceptibility artifact around the sinuses. In our study, the olfactory bulbs
were structurally defined in each dog prior to analysis, allowing us to account for the unique
aspects of brain morphology across individual canines. In dogs, we found a significant main
effect for the differentiation between odor types, similar to human studies of olfactory
processing, but over a much larger region of cortex. In other nonhumans, imaging mammalian
olfactory cortex may prove difficult due to the resulting signal loss from the air-to-tissue
contact in regions near the olfactory bulbs. That said, fMRI of odor processing in canines within
this primary sensory region may offer opportunities to understand the mechanisms of odor
perception above and beyond what is possible in human fMRI.
Second, while humans use language to describe events and percepts, odors are difficult
to describe verbally (Cain, de Wijk, Lulejian, Schiet, & See, 1998; Iatropoulos et al., 2018). When
odors are administered during language-dependent tasks, interference occurs when the odor
and label are simultaneously processed. This difficulty is thought to be due to limitations in
cortical networks, as spatiotemporal patterns produced in neural coding of odors and language
are similar. Additionally, humans' limited language for odors may be a cause for our disregard
of this sense compared to our bias for visual stimuli (Lorig, 1999). Odor naming may also
account for some of the difficulty reported by participants when attempting to evoke images of
the odor objects (Stevenson, Case, & Mahmut, 2007). The inability to name objects based on
their olfactory, as opposed to their visual appearance, may be explained by the brain circuitry
involved in associating olfactory and visual object features to their lexico-semantic
representations (Olofsson & Gottfried, 2015; Olofsson et al., 2014; Olofsson & Wilson, 2018).
Dogs prove to be a valuable model for the study of odor processing because they do not have

the confound of language and have unique brain morphology for imaging of primary olfactory
cortex.
This study also contributes significantly to the existing literature on odor processing in
canines. First, this study replicates findings from our previous odor fMRI study using the same
dogs and the same odor stimuli (Prichard, Chhibber, et al., 2018). Second, ours is the first study
to use data directly from the awake, unanesthetized dog (i.e. brain imaging) as opposed to
behavioral outcomes to assess dogs' perception of odor mixtures. And in contrast, other canine
fMRI studies examining the neural correlates of odor processing have used restrained or
anesthetized subjects (Jia et al., 2014; Siniscalchi, 2016; Thompkins, Deshpande, Waggoner, &
Katz, 2016). Third, we used RFC to perform decoding of the dog brain with awake, unrestrained
dogs. In particular, this study supports the differences inherent in univariate fMRI analyses
compared to MVPA analyses, as the latter do not classify stimuli based on mean activations
(Hebart & Baker, 2018). This allowed us to identify regions supporting classification of stimuli in
addition to those specified in univariate analyses. Further, the RFC proved to be a better
classifier than the popular SVM or searchlight methods. And fourth, ours is the first study to use
RFC in nonhuman fMRI and to back map the feature importances into brain space to identify
regions that contribute to high classification accuracy. Our novel use of RFC can inform future
brain decoding studies as it offers an alternative approach to popular searchlight methods for
localizing important regions.
There are several possible limitations to our study. First, the presence of the human
owner was constant. Because the human was not blind to the nature of the stimuli, they could
have inadvertently influenced the dogs through body language. However, the olfactory stimuli
were least likely to be picked up by the humans and were not communicated by human
owners, so-called 'Clever Hans' effects are unlikely to explain these results. Second, the effects
of habituation counteract those of learning. Habituation was perhaps most evident in the
amygdala, which displayed a generally declining response with run across trial types. There is
ample evidence that the amygdala habituates to repeated presentations of the same stimuli
and specifically to odor stimuli (Gottfried, O'Doherty, & Dolan, 2002; Plichta et al., 2014;
Poellinger et al., 2001; Wright et al., 2001). It would not be surprising that repeated
presentation of the stimuli could lead to decreased physiological response, especially to odors.
Third, with the exception of dogs Pearl and Caylin, the pet dogs that participated in the study
were not previously trained on odor-detection or discrimination. Highly trained working dogs
may perform differently than pets. However, the results were consistent across dogs that
varied in age, breed, and sex, so generalizability to the population is likely. Fourth, the odor
training utilized two component odors and one mixture, so the findings may not generalize to
all odor mixtures or all mixture concentrations (Schneider et al., 2016). Finally, the stimulus-
reward associations were acquired through a passive task in the scanner. No behavioral tests
were conducted to test acquisition of the learned associations or to compare to the neural
activations. This task design was chosen to minimize any additional training required for the
dogs and as a follow-up to our previously published study on odor learning. As in humans,
further study may reveal dissociable neural pathways support the associative and perceptual
representations of sensory stimuli (Howard et al., 2016).
Our results highlight potential neural mechanisms that underly the perception of odors
in dogs. ROI-based analysis of the amygdala and caudate nucleus suggests that learned

associations are important for odor perception across dogs and that these associations are
maintained over time. Further analyses of dogs' perception of an odor mixture suggest that
dogs perceive odor mixtures as new odors rather than as their individual components. This
finding has important implications for the training of odor detection dogs and serves as a
potential mechanism underlying dogs' poor behavioral performance when generalizing from a
target odor to mixture. We also utilized the random forest classifier for decoding fMRI data and
a novel method for mapping the additional regions important for predicting the identity of odor
stimuli. As the use of RFC was exploratory, future more refined decoding studies of the dog
brain may allow us to better understand canine perception and highlight potential neural
mechanisms for olfactory processing conserved across species.

REFERENCES

Ache, B. W., & Young, J. M. (2005). Olfaction: diverse species, conserved principles. *Neuron, 48*(3), 417-
430. doi:10.1016/j.neuron.2005.10.022
Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible evaluation
of ANTs similarity metric performance in brain image registration. *Neuroimage, 54*(3), 2033-
2044. doi:10.1016/j.neuroimage.2010.09.025
Berns, G. S., Brooks, A., & Spivak, M. (2013). Replicability and heterogeneity of awake unrestrained
canine FMRI responses. *PLoS One, 8*(12), e81698. doi:10.1371/journal.pone.0081698
Berns, G. S., Brooks, A. M., & Spivak, M. (2012). Functional MRI in awake unrestrained dogs. *PLoS One,
7*(5), e38027. doi:10.1371/journal.pone.0038027
Berns, G. S., Brooks, A. M., & Spivak, M. (2015). Scent of the familiar: An fMRI study of canine brain
responses to familiar and unfamiliar human and dog odors. *Behav Processes, 110*, 37-46.
doi:10.1016/j.beproc.2014.02.011
Berns, G. S., & Cook, P. F. (2016). Why did the dog walk into the MRI? *Current Directions in Psychological
Science, 25*(5), 363-369. doi:10.1177/0963721416665006
Bestgen, A. K., Schulze, P., Kuchinke, L., Suchan, B., Derdak, T., Otto, T., . . . Sucker, K. (2016). An
extension of olfactometry methods: An expandable, fully automated, mobile, MRI-compatible
olfactometer. *J Neurosci Methods, 261*, 85-96. doi:10.1016/j.jneumeth.2015.12.009
Bijland, L. R., Bomers, M. K., & Smulders, Y. M. (2013). Smelling the diagnosis: A review on the use of
scent in diagnosing disease. *The Netherlands Journal of Medicine*.
Cain, W. S., de Wijk, R., Lulejian, C., Schiet, F., & See, L. C. (1998). Odor identification: Perceptual and
semantic dimensions. *Chemical Senses, 23*(3), 309-326. doi:DOI 10.1093/chemse/23.3.309
Chollet, F. (2018). *Deep learning with python*. NY: Manning.
Cook, P. F., Prichard, A., Spivak, M., & Berns, G. S. (2016). Awake canine fMRI predicts dogs' preference
for praise vs food. *Soc Cogn Affect Neurosci, 11*(12), 1853-1862. doi:10.1093/scan/nsw102
Cooper, R., Wang, C., & Singh, N. (2014). Accuracy of Trained Canines for Detecting Bed Bugs
(Hemiptera: Cimicidae). *J Econ Entomol, 107*(6), 2171-2181. doi:10.1603/EC14195
Cuaya, L. V., Hernandez-Perez, R., & Concha, L. (2016). Our Faces in the Dog's Brain: Functional Imaging
Reveals Temporal Cortex Activation during Perception of Human Faces. *PLoS One, 11*(3),
e0149431. doi:10.1371/journal.pone.0149431
Datta, R., Lee, J., Duda, J., Avants, B. B., Vite, C. H., Tseng, B., . . . Aguirre, G. K. (2012). A digital atlas of
the dog brain. *PLoS One, 7*(12), e52140. doi:10.1371/journal.pone.0052140

Davidson, G. A., Clark, D. A., Johnson, B. K., Waits, L. P., & Adams, J. R. (2014). Estimating Cougar
Densities in Northeast Oregon Using Conservation Detection Dogs. *Journal of Wildlife*
Management, 78(6), 1104-1114. doi:10.1002/jwmg.758
DeGreeff, L. E., Malito, M., Katilie, C. J., Brandon, A., Conroy, M. W., Peranich, K., . . . Rose-Pehrsson, S. L.
(2017). Passive delivery of mixed explosives vapor from separated components. *Forensic*
Chemistry, 4, 19-31. doi:10.1016/j.forc.2017.02.010
Dilks, D. D., Cook, P., Weiller, S. K., Berns, H. P., Spivak, M., & Berns, G. S. (2015). Awake fMRI reveals a
specialized region in dog temporal cortex for face processing. *PeerJ, 3*, e1115.
doi:10.7717/peerj.1115
Gadbois, S., & Reeve, C. (2014). Canine Olfaction: Scent, Sign, and Situation. In *Domestic Dog Cognition*
and Behavior (pp. 3-29).
Göth, A., McLean, I. G., & Trevelyan, J. (2003). How do dogs detect landmines? In.
Gottfried, J. A., O'Doherty, J., & Dolan, R. J. (2002). Appetitive and aversive olfactory learning in humans
studied using event-related functional magnetic resonance imaging. *Journal of Neuroscience,*
22(24), 10829-10837. doi:22/24/10829 [pii]
Haxby, J. V., Connolly, A. C., & Guntupalli, J. S. (2014). Decoding neural representational spaces using
multivariate pattern analysis. *Annu Rev Neurosci, 37*, 435-456. doi:10.1146/annurev-neuro-
062012-170325
Hayes, J. E., McGreevy, P. D., Forbes, S. L., Laing, G., & Stuetz, R. M. (2018). Critical review of dog
detection and the influences of physiology, training, and analytical methodologies. *Talanta, 185*,
499-512. doi:10.1016/j.talanta.2018.04.010
Hebart, M. N., & Baker, C. I. (2018). Deconstructing multivariate decoding for the study of brain
function. *Neuroimage, 180*(Pt A), 4-18. doi:10.1016/j.neuroimage.2017.08.005
Howard, J. D., & Gottfried, J. A. (2014). Configural and elemental coding of natural odor mixture
components in the human brain. *Neuron, 84*(4), 857-869. doi:10.1016/j.neuron.2014.10.012
Howard, J. D., Gottfried, J. A., Tobler, P. N., & Kahnt, T. (2015). Identity-specific coding of future rewards
in the human orbitofrontal cortex. *Proc Natl Acad Sci U S A, 112*(16), 5195-5200.
doi:10.1073/pnas.1503550112
Howard, J. D., Kahnt, T., & Gottfried, J. A. (2016). Converging prefrontal pathways support associative
and perceptual features of conditioned stimuli. *Nat Commun, 7*, 11546.
doi:10.1038/ncomms11546
Hummel, T., Olgun, S., Gerber, J., Huchel, U., & Frasnelli, J. (2013). Brain responses to odor mixtures with
sub-threshold components. *Front Psychol, 4*, 786. doi:10.3389/fpsyg.2013.00786
Iatropoulos, G., Herman, P., Lansner, A., Karlgren, J., Larsson, M., & Olofsson, J. K. (2018). The language
of smell: Connecting linguistic and psychophysical properties of odor descriptors. *Cognition, 178*,
37-49. doi:10.1016/j.cognition.2018.05.007
Jia, H., Pustovyy, O. M., Waggoner, P., Beyers, R. J., Schumacher, J., Wildey, C., . . . Deshpande, G. (2014).
Functional MRI of the olfactory system in conscious dogs. *PLoS One, 9*(1), e86362.
doi:10.1371/journal.pone.0086362
Jin, J., Zelano, C., Gottfried, J. A., & Mohanty, A. (2015). Human Amygdala Represents the Complete
Spectrum of Subjective Valence. *J Neurosci, 35*(45), 15145-15156. doi:10.1523/JNEUROSCI.2450-
15.2015
Johnen, D., Heuwieser, W., & Fischer-Tenhagen, C. (2017). An approach to identify bias in scent
detection dog testing. *Applied Animal Behaviour Science, 189*, 1-12.
doi:10.1016/j.applanim.2017.01.001
Kahnt, T. (2018). A decade of decoding reward-related fMRI signals and where we go from here.
Neuroimage, 180(Pt A), 324-333. doi:10.1016/j.neuroimage.2017.03.067

Kahnt, T., Park, S. Q., Haynes, J. D., & Tobler, P. N. (2014). Disentangling neural representations of value
753   and salience in the human brain. *Proc Natl Acad Sci U S A, 111*(13), 5000-5005.
754   doi:10.1073/pnas.1320189111
755 Lazarowski, L., & Dorman, D. C. (2014). Explosives detection by military working dogs: Olfactory
756   generalization from components to mixtures. *Applied Animal Behaviour Science, 151*, 84-93.
757   doi:10.1016/j.applanim.2013.11.010
758 Lazarowski, L., Foster, M. L., Gruen, M. E., Sherman, B. L., Fish, R. E., Milgram, N. W., & Dorman, D. C.
759   (2015). Olfactory discrimination and generalization of ammonium nitrate and structurally
760   related odorants in Labrador retrievers. *Anim Cogn, 18*(6), 1255-1265. doi:10.1007/s10071-015-
761   0894-9
762 Lorig, T. S. (1999). On the similarity of odor and language perception. *Neurosci Biobehav Rev, 23*(3), 391-
763   398. doi:10.1016/S0149-7634(98)00041-4
764 Lorig, T. S. (2012). Beyond Self-report: Brain Imaging at the Threshold of Odor Perception. *Chemosensory*
765   *Perception, 5*(1), 46-54. doi:10.1007/s12078-012-9116-x
766 Lowen, S. B., & Lukas, S. E. (2006). A low-cost, MR-compatible olfactometer. *Behav Res Methods, 38*(2),
767   307-313. doi:10.3758/BF03192782
768 Misaki, M., Kim, Y., Bandettini, P. A., & Kriegeskorte, N. (2010). Comparison of multivariate classifiers
769   and response normalizations for pattern-information fMRI. *Neuroimage, 53*(1), 103-118.
770   doi:10.1016/j.neuroimage.2010.05.051
771 Olofsson, J. K., & Gottfried, J. A. (2015). The muted sense: neurocognitive limitations of olfactory
772   language. *Trends Cogn Sci, 19*(6), 314-321. doi:10.1016/j.tics.2015.04.007
773 Olofsson, J. K., Hurley, R. S., Bowman, N. E., Bao, X., Mesulam, M. M., & Gottfried, J. A. (2014). A
774   designated odor-language integration system in the human brain. *J Neurosci, 34*(45), 14864-
775   14873. doi:10.1523/JNEUROSCI.2247-14.2014
776 Olofsson, J. K., & Wilson, D. A. (2018). Human Olfaction: It Takes Two Villages. *Curr Biol, 28*(3), R108-
777   R110. doi:10.1016/j.cub.2017.12.016
778 Plichta, M. M., Grimm, O., Morgen, K., Mier, D., Sauer, C., Haddad, L., . . . Meyer-Lindenberg, A. (2014).
779   Amygdala habituation: A reliable fMRI phenotype. *Neuroimage, 103*, 383-390.
780   doi:10.1016/j.neuroimage.2014.09.059
781 Poellinger, A., Thomas, R., Lio, P., Lee, A., Makris, N., Rosen, B. R., & Kwong, K. K. (2001). Activation and
782   habituation in olfaction-an fMRI study. *Neuroimage, 13*(4), 547-560.
783   doi:10.1006/nimg.2000.0713
784 Prichard, A., Chhibber, R., Athanassiades, K., Spivak, M., & Berns, G. S. (2018). Fast neural learning in
785   dogs: A multimodal sensory fMRI study. *Sci Rep, 8*(1), 14614. doi:10.1038/s41598-018-32990-2
786 Prichard, A., Cook, P. F., Spivak, M., Chhibber, R., & Berns, G. S. (2018). Awake fMRI Reveals Brain
787   Regions for Novel Word Detection in Dogs. *Front Neurosci, 12*, 737.
788   doi:10.3389/fnins.2018.00737
789 Qu, L. P., Kahnt, T., Cole, S. M., & Gottfried, J. A. (2016). De Novo Emergence of Odor Category
790   Representations in the Human Brain. *J Neurosci, 36*(2), 468-478. doi:10.1523/JNEUROSCI.3248-
791   15.2016
792 Schneider, N. Y., Datiche, F., Wilson, D. A., Gigot, V., Thomas-Danguin, T., Ferreira, G., & Coureaud, G.
793   (2016). Brain processing of a configural vs elemental odor mixture in the newborn rabbit. *Brain*
794   *Struct Funct, 221*(5), 2527-2539. doi:10.1007/s00429-015-1055-2
795 Sezille, C., Messaoudi, B., Bertrand, A., Joussain, P., Thevenet, M., & Bensafi, M. (2013). A portable
796   experimental apparatus for human olfactory fMRI experiments. *J Neurosci Methods, 218*(1), 29-
797   38. doi:10.1016/j.jneumeth.2013.04.021
798 Siniscalchi, M. (2016). Olfaction and the canine brain. In T. Jezierski, J. Ensminger, & L. E. Papet (Eds.),
799   *Canine Olfaction, Science, and Law*: CRC Press.

Sommer, J. U., Maboshe, W., Griebe, M., Heiser, C., Hormann, K., Stuck, B. A., & Hummel, T. (2012). A
mobile olfactometer for fMRI-studies. *J Neurosci Methods, 209*(1), 189-194.
doi:10.1016/j.jneumeth.2012.05.026
Stelzer, J., Chen, Y., & Turner, R. (2013). Statistical inference and multiple testing correction in
classification-based multi-voxel pattern analysis (MVPA): random permutations and cluster size
control. *Neuroimage, 65*, 69-82. doi:10.1016/j.neuroimage.2012.09.063
Stevenson, R. J., Case, T. I., & Mahmut, M. (2007). Difficulty in evoking odor images: the role of odor
naming. *Mem Cognit, 35*(3), 578-589.
Thomas-Danguin, T., Sinding, C., Romagny, S., El Mountassir, F., Atanasova, B., Le Berre, E., . . .
Coureaud, G. (2014). The perception of odor objects in everyday life: a review on the processing
of odor mixtures. *Front Psychol, 5*, 504. doi:10.3389/fpsyg.2014.00504
Thompkins, A. M., Deshpande, G., Waggoner, P., & Katz, J. S. (2016). Functional Magnetic Resonance
Imaging of the Domestic Dog: Research, Methodology, and Conceptual Issues. *Comp Cogn Behav*
Rev, 11(April), 63-82. doi:10.3819/ccbr.2016.110004
Toledano, A., Borromeo, S., Luna, G., Molina, E., Solana, A. B., Garcia-Polo, P., . . . Alvarez-linera, J.
(2012). Objective assessment of olfactory function using functional magnetic resonance imaging.
Acta Otorrinolaringol Esp, 63(4), 280-285. doi:10.1016/j.otorri.2012.01.010
Vedaei, F., Oghabian, M. A., Firouznia, K., Harirchian, M. H., Lotfi, Y., & Fakhri, M. (2017). The Human
Olfactory System: Cortical Brain Mapping Using fMRI. *Iranian Journal of Radiology, 14*(2).
doi:10.5812/iranjradiol.16250
Vigouroux, M., Bertrand, B., Farget, V., Plailly, J., & Royet, J. P. (2005). A stimulation method using odors
suitable for PET and fMRI studies with recording of physiological and behavioral signals. *J*
Neurosci Methods, 142(1), 35-44. doi:10.1016/j.jneumeth.2004.07.010
Wright, C. I., Fischer, H., Whalen, P. J., McInerney, S. C., Shin, L. M., & Rauch, S. L. (2001). Differential
prefrontal cortex and amygdala habituation to repeatedly presented emotional stimuli.
NeuroReport, 12(2), 379-383. doi:10.1097/00001756-200102120-00039
Yeshurun, Y., Lapid, H., Dudai, Y., & Sobel, N. (2009). The privileged brain representation of first
olfactory associations. *Curr Biol, 19*(21), 1869-1874. doi:10.1016/j.cub.2009.09.066
Zelano, C., Mohanty, A., & Gottfried, J. A. (2011). Olfactory predictive codes and stimulus templates in
piriform cortex. *Neuron, 72*(1), 178-187. doi:10.1016/j.neuron.2011.08.010