Cdc42 promotes Bgs1 recruitment for septum synthesis and glucanase localization for cell separation during cytokinesis in fission yeast

Udo N. Onwubiko, Julie-Rich Robinson, Rose Albu Mustaf, and Maitreyi E. Das*

Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, TN, USA.

Key words: GTPase, Cdc42, cytokinesis, Bgs1, Eng1, Agn1

*Corresponding Author:

Maitreyi Das

University of Tennessee, Knoxville

415, Ken and Blaire Mossman Building

Knoxville, TN, 37996

mdas@utk.edu

Abstract

The conserved small Rho GTPase Cdc42 has recently reported to play a role in cytokinesis in fission yeast, *Schizosaccharomyces pombe.* While the role of Cdc42 in cell polarity has been widely studied, little is known about its role in the regulation of cytokinesis. Here we investigate specific roles of Cdc42 during cytokinesis in fission yeast. Cytokinesis in fission yeast involves the coupling of actomyosin ring constriction to division cell wall (septum) synthesis. Cdc42 activators have been shown to promote timely initiation of septum synthesis and proper septum morphology. However, direct evidence supporting a role for Cdc42 in these events are lacking. Using a thermosensitive *cdc42* allele (*cdc42-1625*), we investigated the direct role of Cdc42 in cytokinesis. We show that Cdc42 is essential for recruitment of the major primary septum synthesizing enzyme Bgs1 during cytokinesis; and that disruption of Cdc42 activity, prevented the initiation of ring constriction. Further, we find that Cdc42 is required for proper localization of the septum digesting enzymes Eng1 and Agn1 at the division site. Thus, we demonstrate that Cdc42 is required during cytokinesis to enable primary septum synthesis, and proper cell separation.

Introduction

Cytokinesis involves the separation of a mother cell into two new daughters. In most eukaryotic cells with the exception of plant cells, cytokinesis initiates with the assembly of a medially positioned actomyosin ring that constricts in a regulated manner to form the cleavage furrow [1, 2, 3]. In the fission yeast *Schizosaccharomyces pombe*, in addition to a medially positioned contractile actomyosin ring, cytokinesis requires the synthesis of a tri-layered cell wall structure called the division septum [4]. Once assembled, the actomyosin undergoes a maturation period where a plethora of proteins necessary for cytokinesis are recruited [5, 6, 7]. Upon completion of maturation, septum synthesis initiates concomitantly with ring constriction. The division septum is a complex structure, consisting of the primary septum which is primarily synthesized by the linear-β-glucan synthase Bgs1, and secondary septum, which is synthesized mainly by the branched B-glucan synthase Bgs4 [8, 9, 10]. Once ring constriction completes, the septum matures, and its middle layer (primary septum) gets degraded by digestive enzymes Eng1 and Agn1 to separate the two new daughter cells, thus completing cytokinesis [11, 12]. Improper regulation of cytokinesis could result in non-viable cells via cell lysis, or stalled cytokinesis.

The small GTPase is a master regulator of cell polarity in fission yeast and has been shown to regulate membrane trafficking [13, 14, 15, 16]. We have previously shown that the small GTPase Cdc42 is activated at the division site in a sequential manner to promote timely initiation of septum ingression and septum formation [17]. Cdc42 is activated by the Guanidine nucleotide exchange factors (GEFs) Gef1 and Scd1 [18, 19]. During cytokinesis, loss of Gef1 delays the initiation of ring constriction and Bgs1 recruitment, and loss of Scd1 results in aberrant septum morphology [17]. It is unclear however whether these roles are specific to the GEFs or Cdc42 activity. Cdc42 has also been implicated in the specific trafficking of Bgs1 containing vesicles to the sites of polarized growth, although the mechanism is not known [16]. Gef1 has also been shown to promote endocytosis and proper organization of the scaffold protein Cdc15 [20].

In this study, we assess the specific role(s) of Cdc42 during cytokinesis using a temperature sensitive Cdc42 allele ─*cdc42-1625* [15, 21]. We find that disruption of Cdc42 activity resulted in defective cytokinesis, impaired Bgs1 localization to the division site, and disrupted localization of the septum remodeling enzymes Eng1 and Agn1. Our results define specific roles of Cdc42 during cytokinesis. We demonstrate that Cdc42 is required for ring constriction by ensuring Bgs1 recruitment and cell separation by ensuring proper localization of Eng1 and Agn1.

Results and Discussion

cdc42-1625 mutants display similar septation index with *cdc42+* cells

To investigate the direct role of Cdc42 in cytokinesis we used the temperature-sensitive Cdc42 allele *cdc42-1625*, to disrupt Cdc42 activity at the non-permissive temperature of 36ºC. Both *cdc42+* and *cdc42-1625* cells were first grown at the permissive temperature of 25ºC, then shifted to 36ºC for 4 hours as previously performed. We observed that *cdc42-1625* mutants appeared wider than *cdc42+* cells even at 25ºC, which became further pronounced and often caused cell death at 36ºC (Fig. 1A). This observation agrees with previous reports that the cdc42-*1625* allele is hypomorphic even at the permissive temperature of 25ºC [21]. We analyzed the number of septated cells in each of the experiments, and find that while *cdc42+*cells displayed an average septation index of ~17% at 25ºC, and ~15% at 36ºC, *cdc42-1625* strains had septation indices of ~22% at 25ºC and ~23% at 36ºC (Fig. 1B). Although we see a mild increase in the septation index both at permissive and non-permissive temperature, the difference is not statistically significant (Fig. 1B).

Cdc42 is required for Bgs1 recruitment to the medial ring during cytokinesis.

To investigate why the septation index of *cdc42-ts* mutants at 25ºC and 36ºC was similar to that of *cdc42+* cells (Fig. 1A), we asked whether these mutant cells displayed fewer septated cells due to an inability recruit the major primary septum synthesizing enzyme Bgs1 and initiate septum synthesis. Bgs1 is recruited to the division site after ring assembly, during maturation [17, 22]. We repeated our experiments in *cdc42+* and *cdc42-1625* cells expressing Rlc1-tdTomato as a ring marker, and GFP-Bgs1. We compared GFP-Bgs1 levels in non-constricting *cdc42+* and *cdc42-1625* cells at 25ºC and 36ºC. We quantified the fluorescence intensities and find that GFP-Bgs1 levels are significantly reduced in *cdc42-1625* mutants at 25ºC and 36ºC (p<0.0001), compared to *cdc42+* cells (Fig. 2B). Thus, we find that irrespective of the shift in temperature, the temperature sensitive mutation is effective enough to disrupt Bgs1 recruitment. We did not observe any differences in GFP-Bgs1 levels in *cdc42+* cells at 25ºC or 36ºC (Fig. 2B). We further assessed the number of cells with actomyosin rings containing Bgs1 in *cdc42+* and *cdc42-1625* cells. We find that ~85% of *cdc42+* cells display actomyosin rings with GFP-Bgs1 at 25ºC and 36ºC (Fig. 2C). Interestingly ~89% of *cdc42-1625* cells at 25ºC, displayed GFP-Bgs1 containing actomyosin rings, and this value was comparable to *cdc42+* cells at 25ºC and 36ºC (Fig. 2C). At 36ºC however, the percentage of GFP-Bgs1 containing actomyosin rings was significantly reduced to ~64%. We find that at 36ºC, fewer actomyosin rings in *cdc42-1625* cells recruit Bgs1. Cdc42 is therefore required for recruitment of Bgs1 to the division site after ring assembly, which further enables the timely initiation of ring constriction.

Next, we probed the percentage of *cdc42-1625* mutants that are able to initiate successful ring constriction. We analyzed the percentage of cells showing constricting rings and found no significant differences in *cdc42+* cells at 25ºC and 36ºC (Fig. 2D). In *cdc42+* cells at 25ºC ~59% of the actomyosin rings appear to be under constriction, while this value is 55% at 36ºC. In *cdc42-1625* mutants, the percentage of constricting rings at 25ºC, was 47%, but decreased to

31%, after 4 hours at 36°C (Fig. 2D). Our data therefore highlight the importance of Cdc42 for Bgs1 recruitment and initiation of ring constriction in cytokinesis.

Cdc42 is required for proper localization of digestive enzymes during cell separation.

While our data indicated that Cdc42 is required for Bgs1 recruitment to the division site, resulting in fewer constricting rings, it fails to explain why the *cdc42-1625* mutants have a septation index similar to the control cells. A recent report indicated that Gef1 has a role in cell separation [23]. Thus, it is possible that the septation index observed in *cdc42-1625* cells is due to a digestion defect of the septa that formed under permissive conditions. To test this, we assessed the role of Cdc42 in septum digestion that leads to cell separation. Cdc42 has also been implicated in endocytosis [20] and polarized trafficking of the exocyst, which is also spatially positioned at the division site during ring constriction and cell separation [24, 25, 26]. To probe the role of Cdc42 in cell separation, we assessed the localization of the septum digesting enzymes Eng1 and Agn1 in *cdc42*+ and *cdc42-1625* strains. Eng1-GFP and Agn1-GFP were simultaneously expressed in *cdc42*+ and *cdc42-1625* strains. Cells were grown at the permissive temperature of 25°C, and imaged to assess Eng1-GFP and Agn1-GFP localization. We observed that these enzymes only localized to septated cells (Fig. 3A), and there were no obvious differences in the fluorescence intensities in both *cdc42*+ and *cdc42-1625* cells. Next, we generated 3D-reconstructions of Eng1-GFP Agn1-GFP localization at the division site, to assess localization patterns. We first looked in *cdc42*+ cells and found three different localization patterns: a ring, a ring and dot, and a disc-like localization (Fig. 2B (i-iii)). For proper septum digestion these enzymes need to be delivered to the outer rim of the membrane barrier in a ring like pattern [25].

3D-reconstructions of Eng1-GFP Agn1-GFP in *cdc42*+ cells mainly displayed 'ring' and 'ring and dot' localization patterns. In *cdc42-1625* mutants at 25 °C, Eng1-GFP Agn1-GFP mostly displayed the 'disc-like' pattern (Fig. 3C), suggesting that these enzymes were not able to localize properly. We quantified the frequency of these patterns within each strain and find that 58% of *cdc42*+ cells, display rings, 33% display a 'ring +dot' and 8% display 'disc-like' localization. In *cdc42-1625* mutants, 0% display any rings, 4% display a 'ring +dot' and 96% display 'disc-like' localization (Fig. 3D). We were unable to observe the enzymes in cells grown at 35°C. The localization of the digestive enzymes Eng1 and Agn1 is thus not polarized, rather appears very diffuse at the division site in *cdc42-1625* cells, suggesting that Cdc42 is required to properly localize these enzymes to enable proper septum digestion.

Conclusions

Cytokinesis is a tightly regulated process, and in fission yeast, septum synthesis and ring constriction are properly coordinated to generate viable daughter cells. We report that Cdc42 is required for proper cytokinesis in fission yeast. We demonstrate that functional Cdc42 is required for the recruitment of the primary septum enzyme Bgs1 and for the proper localization

of the septum synthesizing enzymes Eng1 and Agn1. Cdc42 is an essential protein and hence here we used a conditional mutant to investigate its role in cytokinesis. It is possible that Cdc42 has an essential role in cytokinesis and this has not been elucidated to date due to the nature of *cdc42* mutants available. Cdc42 has been reported to play major roles in membrane trafficking events [15, 16, 20]. However, Cdc42 specifically promotes recruitment of Bgs1 but is not required for the recruitment of a secondary septum synthesizing enzyme Bgs4 [17]. Thus, it is likely that the role of Cdc42 in cytokinesis is to regulate the recruitment of specific cargo required for distinct steps in cytokinesis. Delivery of Bgs1 and the glucanases occur at spatially regions of the division site. Future research will reveal how Cdc42 promotes precise spatial recruitment of these different enzymes.

Materials and methods

Strains and Cell culture

Strains used in this study are listed in Table 1. All *S. pombe* strains used in this study are isogenic to PN972. Unless otherwise mentioned cells were cultured in yeast extract (YE) medium and grown exponentially at 25°C. Genetic manipulations of strains were carried out using standard techniques (Moreno et al., 1991). Cells were grown exponentially for at least 3 rounds of eight generations for each assay. On the day of the experiment, cells were grown to O.D. of 0.1 and a subset of control and *cdc42-1625* cell culture was transferred to another flask, and placed in a 36°C shaking incubator for 4 hours before imaging.

Table 1: Strains list		
Strain	**Genotype**	**Origin**
972	*h-*	*P. Nurse*
527	*h- ade6-704 leu1-32 ura4-D18 his3-D1*	*P. Nurse*
528	*h+ ade6-704 leu1-32 ura4-D18 his3-D1*	*P. Nurse*
PPG5660	*h+ cdc42-1625-kanMX leu1-32 ura4-D18*	*[21]*
YMD1452	*bgs1-GFP:leu1+ rlc1-tdTomato- NATᵣ leu1-32 ura4-D18*	*This study*
YMD825	*h+ eng1-GFP agn1-GFP*	*This study*
YMD1412	*cdc42-1625 GFP-bgs1:leu1 rlc1-tdTomato- NATᵣ leu1-32 ade6-M210 ura4-D18*	*This study*

Microscopy

Image acquisition was performed at room temperature (23-25°C) with the VT-Hawk 2D- array scanning confocal microscope (Visitech intl., Sunderland, UK) using an Olympus IX-83 inverted microscope with a 100x/numerical aperture 1.49 UAPO lens (Olympus, Tokyo, Japan). Cells

were spun down in a mini desk microcentrifuge, mounted directly onto glass slides with a #1.5 coverslip (Fischer Scientific, Waltham, MA) and imaged promptly. Z-series images were acquired with depth interval of 0.4um. All Images were acquired with MetaMorph (Molecular Devices, Sunnyvale, CA). All images were analyzed with Image J (National Institutes of Health, Bethesda, MD).

Statistical analysis was performed using 2-way ANOVA, followed by Tukey's HSD Post Hoc Test where appropriate. Comparisons between experimental groups were considered significant when $p \leq 0.05$.

Calcofluor Staining

Cells were grown as described earlier in the methods section, and stained in yeast extract media (YE) with 50µg/ml Calcofluor White (M2R Sigma-Aldrich, St. Louis, MO) at room temperature. Both controls and mutant strains were washed with fresh YE liquid 1x, and imaged immediately.

Image Analysis

Sum projections of acquired z-series images were performed in Image J. We quantified GFP-Bgs1 levels from sum- projections. To do this, we measured the intensity at the division site in cells with non-constricting actomyosin rings, using a box with an area of 330pixel^2. Background correction was performed with the cytoplasm for each cell.

References:

1. Pollard TD. Mechanics of cytokinesis in eukaryotes. Curr Opin Cell Biol. 2010 Feb;22(1):50-6. doi: 10.1016/j.ceb.2009.11.010. PubMed PMID: 20031383; PubMed Central PMCID: PMC2871152.
2. Li R. Cytokinesis in development and disease: variations on a common theme. Cell Mol Life Sci. 2007 Dec;64(23):3044-58. doi: 10.1007/s00018-007-7285-6. PubMed PMID: 17882379.
3. Guertin DA, Trautmann S, McCollum D. Cytokinesis in eukaryotes. Microbiol Mol Biol Rev. 2002 Jun;66(2):155-78. PubMed PMID: 12040122; PubMed Central PMCID: PMC120788.
4. Cortes JC, Ramos M, Osumi M, et al. Fission yeast septation. Commun Integr Biol. 2016 Jul-Aug;9(4):e1189045. doi: 10.1080/19420889.2016.1189045. PubMed PMID: 27574536; PubMed Central PMCID: PMCPMC4988442.
5. Wu JQ, Kuhn JR, Kovar DR, et al. Spatial and temporal pathway for assembly and constriction of the contractile ring in fission yeast cytokinesis. Dev Cell. 2003 Nov;5(5):723-34. PubMed PMID: 14602073.
6. Roberts-Galbraith RH, Chen JS, Wang J, et al. The SH3 domains of two PCH family members cooperate in assembly of the Schizosaccharomyces pombe contractile ring. J Cell Biol. 2009 Jan 12;184(1):113-27. doi: 10.1083/jcb.200806044. PubMed PMID: 19139265; PubMed Central PMCID: PMC2615086.
7. Ren L, Willet AH, Roberts-Galbraith RH, et al. The Cdc15 and Imp2 SH3 domains cooperatively scaffold a network of proteins that redundantly ensure efficient cell division in fission yeast. Mol Biol Cell. 2015 Jan 15;26(2):256-69. doi: 10.1091/mbc.E14-10-1451. PubMed PMID: 25428987; PubMed Central PMCID: PMCPMC4294673.
8. Cortes JC, Konomi M, Martins IM, et al. The (1,3)beta-D-glucan synthase subunit Bgs1p is responsible for the fission yeast primary septum formation. Mol Microbiol. 2007 Jul;65(1):201-17. doi: 10.1111/j.1365-2958.2007.05784.x. PubMed PMID: 17581129.
9. Munoz J, Cortes JC, Sipiczki M, et al. Extracellular cell wall beta(1,3)glucan is required to couple septation to actomyosin ring contraction. J Cell Biol. 2013 Oct 28;203(2):265-82. doi: 10.1083/jcb.201304132. PubMed PMID: 24165938; PubMed Central PMCID: PMCPMC3812973.
10. Cortes JC, Carnero E, Ishiguro J, et al. The novel fission yeast (1,3)beta-D-glucan synthase catalytic subunit Bgs4p is essential during both cytokinesis and polarized growth. J Cell Sci. 2005 Jan 1;118(Pt 1):157-74. doi: 10.1242/jcs.01585. PubMed PMID: 15615781.
11. Martin-Cuadrado AB, Duenas E, Sipiczki M, et al. The endo-beta-1,3-glucanase eng1p is required for dissolution of the primary septum during cell separation in Schizosaccharomyces pombe. J Cell Sci. 2003 May 1;116(Pt 9):1689-98. doi: 10.1242/jcs.00377. PubMed PMID: 12665550.
12. Dekker N, Speijer D, Grun CH, et al. Role of the alpha-glucanase Agn1p in fission-yeast cell separation. Mol Biol Cell. 2004 Aug;15(8):3903-14. doi: 10.1091/mbc.e04-04-0319. PubMed PMID: 15194814; PubMed Central PMCID: PMCPMC491845.
13. Das M, Verde F. Role of Cdc42 dynamics in the control of fission yeast cell polarization. Biochem Soc Trans. 2013 Dec;41(6):1745-9. doi: 10.1042/BST20130241. PubMed PMID: 24256285.
14. Martin SG. Spontaneous cell polarization: Feedback control of Cdc42 GTPase breaks cellular symmetry. Bioessays. 2015 Nov;37(11):1193-201. doi: 10.1002/bies.201500077. PubMed PMID: 26338468.
15. Estravis M, Rincon SA, Santos B, et al. Cdc42 regulates multiple membrane traffic events in fission yeast. Traffic. 2011 Dec;12(12):1744-58. doi: 10.1111/j.1600-0854.2011.01275.x. PubMed PMID: 21899677.
16. Estravis M, Rincon S, Perez P. Cdc42 regulation of polarized traffic in fission yeast. Commun Integr Biol. 2012 Jul 1;5(4):370-3. doi: 10.4161/cib.19977. PubMed PMID: 23060961; PubMed Central PMCID: PMC3460842.

17.	Wei B, Hercyk BS, Mattson N, et al. Unique spatiotemporal activation pattern of Cdc42 by Gef1 and Scd1 promotes different events during cytokinesis. Mol Biol Cell. 2016 Apr 15;27(8):1235-45. doi: 10.1091/mbc.E15-10-0700. PubMed PMID: 26941334; PubMed Central PMCID: PMCPMC4831878.

18.	Coll PM, Trillo Y, Ametzazurra A, et al. Gef1p, a new guanine nucleotide exchange factor for Cdc42p, regulates polarity in Schizosaccharomyces pombe. Molecular biology of the cell. 2003;14(1):313-23. PubMed PMID: 12529446; eng.

19.	Chang EC, Barr M, Wang Y, et al. Cooperative interaction of S. pombe proteins required for mating and morphogenesis. Cell. 1994 Oct 7;79(1):131-41. doi: 0092-8674(94)90406-5 [pii]. PubMed PMID: 7923372; eng.

20.	Onwubiko UN, Mlynarczyk PJ, Wei B, et al. A Cdc42 GEF, Gef1, through endocytosis organizes F-BAR Cdc15 along the actomyosin ring and promotes concentric furrowing. J Cell Sci. 2019 Feb 28;132(5). doi: 10.1242/jcs.223776. PubMed PMID: 30709916; PubMed Central PMCID: PMCPMC6432710.

21.	Martin SG, Rincon SA, Basu R, et al. Regulation of the formin for3p by cdc42p and bud6p. Molecular biology of the cell. 2007;18(10):4155-67. PubMed PMID: 17699595; eng.

22.	JC GC, Ramos M, Konomi M, et al. Specific detection of fission yeast primary septum reveals septum and cleavage furrow ingression during early anaphase independent of mitosis completion. PLoS Genet. 2018 May;14(5):e1007388. doi: 10.1371/journal.pgen.1007388. PubMed PMID: 29813053; PubMed Central PMCID: PMCPMC5993333.

23.	Wang N, Wang M, Zhu YH, et al. The Rho-GEF Gef3 interacts with the septin complex and activates the GTPase Rho4 during fission yeast cytokinesis. Mol Biol Cell. 2015 Jan 15;26(2):238-55. doi: 10.1091/mbc.E14-07-1196. PubMed PMID: 25411334; PubMed Central PMCID: PMC4294672.

24.	Wang H, Tang X, Liu J, et al. The multiprotein exocyst complex is essential for cell separation in Schizosaccharomyces pombe. Mol Biol Cell. 2002 Feb;13(2):515-29. doi: 10.1091/mbc.01-11-0542. PubMed PMID: 11854409; PubMed Central PMCID: PMC65646.

25.	Martin-Cuadrado AB, Morrell JL, Konomi M, et al. Role of septins and the exocyst complex in the function of hydrolytic enzymes responsible for fission yeast cell separation. Mol Biol Cell. 2005 Oct;16(10):4867-81. doi: E04-12-1114 [pii]10.1091/mbc.E04-12-1114. PubMed PMID: 16079182; eng.

26.	Bendezu FO, Vincenzetti V, Martin SG. Fission yeast Sec3 and Exo70 are transported on actin cables and localize the exocyst complex to cell poles. PLoS One. 2012;7(6):e40248. doi: 10.1371/journal.pone.0040248. PubMed PMID: 22768263; PubMed Central PMCID: PMC3386988.

Figure Legends:

Figure 1. *cdc42-1625* mutants display similar septation index with *cdc42+* cells. A. Representative images of *cdc42⁺* and *cdc42-1625* at permissive (25°C) and non-permissive (36°C) temperatures; [Scale bar: 10μm]. **B.** Septation indices for *cdc42⁺* and *cdc42-1625* at permissive (25°C) and non-permissive (36°C) temperatures. Each point represents the septation index for one image. [Red lines on graph represent the mean; n.s, not statistically significant].

Figure 2. Cdc42 is required for Bgs1 recruitment to the medial ring during cytokinesis. **A.** Sum z-projections of images displaying Bgs1-GFP and Rlc1-tdTomato at permissive (25°C) and non-permissive (36°C) temperatures in control cells and *cdc42-1625* mutants. White arrows point to Bgs1-GFP localization at the ring in cells with non-constricting rings. **B.** Quantification of Bgs1-GFP levels in cells with non-constricting rings in the mentioned strains at permissive and non-permissive temperatures, n=32 cells. **C.** Quantification of the percentage of cells with ring showing Bgs1 localization, n>100 cells, per strain. **D.** Quantification of the percentage of constricting rings observed among cells displaying an actomyosin ring in the mentioned strains, at permissive and non-permissive temperatures. [Red lines on graph represent mean, blue line represents median, for all graphs, *P≤0.05; **P≤0.01; ****P≤0.0001; n.s., not statistically significant; Scale bar: 5μm].

Fig. 3. Cdc42 is required for proper localization of digestive enzymes during cell separation. A. Sum z-projections of images displaying GFP tagged digestive enzymes Eng1 and Agn1 at permissive temperature (25°C) in control cells and *cdc42-1625* mutants, scale bar: 5μm. **B.** Bright-field and fluorescent images of Eng1-GFP and Agn1-GFP in control cells undergoing cell separation. The pink boxes highlight the 3d-reconstructions of the different localization patterns of Eng1-GFP and Agn1-GFP observed: (i) Ring, (ii) Ring + Dot, (iii) Disk. **C.** 3D-reconstructed division site of Eng1-GFP Agn1-GFP localization at the division site during cell separation in the strains listed, at 25°C, Scale bar: 5μm. **D.** Quantification of Eng1-GFP and Agn1-GFP localization patterns at the division site in the strains mentioned. Scale bar for all 3D-reconstructed division site: 2μm.

Figure 1

A

	bright-field	calcofluor

B

Septation Index

Figure 2

Figure 3

A — 25°C, Eng1-GFP Agn1-GFP: cdc42⁺, cdc42-1625

B — Bright-field; Eng1-GFP Agn1-GFP (i), (ii), (iii)

C — 25°C, Eng1-GFP Agn1-GFP: cdc42⁺, cdc42-1625

D — Percentage(%): Ring, Ring+Dot, Disc — cdc42⁺, cdc42-1625