Spherical arena reveals optokinetic response tuning to
stimulus location, size and frequency across entire
visual field of larval zebrafish

Authors

Florian A. Dehmelt[1]*, Rebecca Meier[1]*, Julian Hinz[1]*[#], Takeshi Yoshimatsu[2],
Clara A. Simacek[1], Kun Wang[1], Tom Baden[2], Aristides B. Arrenberg[1]¶

These authors contributed equally to this work.

Affiliations

1: University of Tübingen, Werner Reichardt Centre for Integrative Neuroscience and
Institute of Neurobiology, 72076 Tübingen, Germany

2: Sussex Neuroscience, School of Life Sciences, University of Sussex, UK

[#] Current address: Friedrich Miescher Institute for Biomedical Research (FMI), Basel,
Switzerland

Correspondence

¶Correspondence should be addressed to A.B.A.
(e-mail: aristides.arrenberg@uni-tuebingen.de)

Keywords

optokinetic response, OKR gain, zebrafish, zebrafish larvae, visual stimulus, visual field,
retinal photoreceptors, retina, stimulus tuning, stimulus size, temporal frequency, spatial
frequency, behaviour, stimulus hardware, spherical, asymmetry, area centralis, yoking

Author contributions

A.B.A. conceived of and supervised the study. J.H., A.B.A. and F.A.D. designed the geometry of the arena. J.H. configured circuit boards, CAD-planned and 3D-printed the arena scaffold. J.H. and K.W. assembled LEDs and electronics. R.M. constructed the optical pathway and developed embedding procedures. F.A.D., R.M. and K.W. tested the stimulus arena. F.A.D. developed mathematical procedures and software mapping stimuli onto the arena. F.A.D., R.M. and A.B.A. wrote software for data analysis. F.A.D. and R.M. designed visual stimuli and analysed data. R.M. performed OKR experiments using the spherical LED arena. C.A.S. performed control experiments on the asymmetry of yoking using the quadratic stimulus arena. T.B. and T.Y. contributed the experiments and analysis of retinal photoreceptor density. F.A.D., J.H. and A.B.A wrote the manuscript. F.A.D. created the figures.

Author contributions by category

Conceptualisation. Aristides B. Arrenberg

Data Curation. (Not applicable)

Formal Analysis. Florian A. Dehmelt, Rebecca Meier

Funding Acquisition. Aristides B. Arrenberg

Investigation. Rebecca Meier, Clara A. Simacek

Methodology. Julian Hinz, Rebecca Meier, Florian A. Dehmelt, Kun Wang, Aristides B. Arrenberg

Project Administration. Aristides B. Arrenberg

Resources. Takeshi Yoshimatsu, Tom Baden

Software. Florian A. Dehmelt, Rebecca Meier, Aristides B. Arrenberg

Supervision. Aristides B. Arrenberg, Florian A. Dehmelt

Validation. Rebecca Meier, Clara A. Simacek, Florian A. Dehmelt

Visualisation. Florian A. Dehmelt

Writing – Original Draft Preparation. Florian A. Dehmelt, Julian Hinz, Rebecca Meier, Aristides B. Arrenberg

Writing – Review & Editing. Florian A. Dehmelt, Julian Hinz, Kun Wang, Takeshi Yoshimatsu, Tom Baden, Aristides B. Arrenberg

Acknowledgements

We thank Väinö Haikala and Dierk F. Reiff (University of Freiburg) for sharing their code and design for hardware controllers, Alexander Borst (MPI Neurobiology, Martinsried) for providing the LED panel board design, Thomas Nieß (glassblowing workshop, University of Tübingen) and Klaus Vollmer (precision mechanics workshop, University of Tübingen) for technical support, Prudenter-Agas (Hamburg, Germany) for generating glass bulb illustrations, Andre Maia Chagas for help with 3D printing procedures, and Jan Benda (University of Tübingen) for discussions on visual acuity.

Abstract

Many animals have large visual fields, allowing them to observe almost any stimulus in their surround. The underlying sensory circuits have evolved to sample those visual field regions most informative to the animal. These regions can vary between different visually mediated behaviours, such as stabilisation and hunting behaviour. Despite this, relatively small displays are often used in vision neuroscience, making it difficult to study the tuning of the visual system to specific visual field locations. To overcome these technical limitations and reveal the organisation of motion circuits with respect to visual space, we built a spherical stimulus arena with 14,848 independently controllable LEDs and used it to stimulate almost the entire visual surround of immobilised zebrafish larvae. We measured the gain of the optokinetic response at different stimulus positions relative to the fish, and related behavioural performance to photoreceptor densities in the retina. We report that zebrafish larvae react most strongly and consistently to stimuli located laterally and near the equator of their visual space. The OKR appears to be symmetric between both eyes, although individual animals oftentimes have a dominant eye. For small stimuli, the OKR gain depends on stimulus size in a logarithmic fashion. OKR to our mostly green stimuli was tuned to the higher spatial densities of red, green and blue photoreceptors in the central retina. In addition, experiments in animals mounted upside-down suggest that extra-retinal processing affects the preferred OKR stimulus location. The tuning to stimulus size and spatial frequency was similar across different visual field positions. During monocular motion stimulation, the non-stimulated eye was strongly yoked if a low-contrast stimulus was present, and less so if a high-contrast stimulus was presented. Our results provide a precise analysis of OKR responses across the whole visual field, and relate sensory performance both to the architecture of the retina and to downstream neural pathways. Our results suggest that motion vision circuits in zebrafish are highly anisotropic. We hypothesize that they monitor specific positions in visual space that are relevant for behaviour in nature, and specifically, that the observed variation of OKR performance across visual field locations is caused by retinal and central adaptations of the zebrafish brain to behavioural needs during visual orientation and stabilisation.

Author summary

The visual system of larval zebrafish mirrors many features, present in the visual system of other vertebrates, including its ability to mediate optomotor and optokinetic behaviour. Although the presence of such behaviours and some of the underlying neural correlates have been firmly established, previous experiments did not consider the large visual field of zebrafish, which covers more than 160° for each eye. Given that different parts of the visual field likely carry unequal amount of behaviourally relevant information for the animal, this raises the question whether optic flow is integrated across the entire visual field or just parts of it, and how this shapes behaviour such as the optokinetic response. We constructed a spherical LED arena to present visual stimuli almost anywhere across their visual field, while tracking horizontal eye movements. By displaying moving gratings on this LED arena, we demonstrate that the optokinetic response, one of the most prominent visually induced behaviours of zebrafish, indeed strongly depends on stimulus location and stimulus size, as well as on other parameters such as the spatial and temporal frequency of the gratings. This location dependence is consistent with areas of high retinal photoreceptor densities.

Blurb

Stimulation across entire visual field reveals that zebrafish optokinetic behaviour is most strongly driven by lateral stimulus locations. This anisotropy is a result of retinal and extra-retinal effects.

Introduction

The layout of the retina and the visual system as a whole evolved to serve specific behavioural tasks animals need to perform in order to survive in their respective habitats. A well-known example is the position of the eyes in the head which varies between hunting animals (frontal eyes) and animals that frequently need to avoid predation (lateral eyes) (1). Hunting animals keep the prey within particular visual field regions to maximize behavioural performance (2-4). To avoid predation, however, it is useful to observe a large proportion of visual space, especially those regions in which predators are most likely to occur (5, 6). The ecological significance of visual stimuli thus depends on their location within the visual field, and it is paralleled by non-uniform processing channels across the retina. This non-uniformity manifests as an *area centralis* or a fovea in many species, which is a region of heightened photoreceptor density in the central retina and serves to increase visual performance in the corresponding visual field regions. Photoreceptor densities put a direct physical limit on performance parameters such as spatial resolution (7, 8). In addition to these restrictions mediated by the peripheral sensory circuitry, an animal's use of certain visual field regions is also affected by behaviour-specific neural pathways and orientation behaviour. The resulting combination of retinal and extra-retinal anisotropies affects the behavioural performance in different tasks – such as feeding and stabilisation behaviour – depending on visual field location (2, 3, 9-13).

Investigating behavioural performance limits and non-uniformities can offer insights into the processing capabilities and ecological adaptations of vertebrate brains, especially if they can be studied and quantitatively understood at each processing step. The larval zebrafish is a promising organism for such an endeavour, since its brain is small and a wide array of experimental techniques is available (14, 15). Zebrafish are lateral-eyed animals and have a large visual field, which covers 163° per eye (16). Their retina contains four different cone photoreceptor types (17), each distributed differently across the retina. UV photoreceptors are densest in the ventro-temporal retina (*area temporalis ventralis*), whereas the red, green and blue photoreceptors cover more central retinal regions (12).

Although zebrafish larvae perform a wide range of visually mediated behaviours, ranging from prey capture (18) and escape behaviour (19) to stabilisation behaviour (20, 21), the importance of stimulus location within the visual field is still not well understood in most cases (but see (3) for prey capture). During visually mediated stabilisation behaviours, such as optokinetic and optomotor responses, animals move their eyes and bodies, respectively, in order to stabilize the retinal image and/or the body position relative to the visual surround. The optokinetic response (OKR) consists of reflexively executed stereotypical eye movements, in which phases of stimulus "tracking" (slow phase) are interrupted by quick phases (**S1a Fig**). In the quick phases, eye position is reset by a saccade in the direction opposite to stimulus motion. In humans, optokinetic responses are strongest in the central visual field (22). Furthermore, lower visual field locations of the stimulus evoke stronger OKR than upper visual field locations, which likely represents an adaptation to the rich optic flow information available from the structures on the ground in the natural environments of primates (9, 23).

In zebrafish larvae, OKR behaviour has been used extensively to assess visual function in genetically altered animals (24-26) and OKR tuning to the velocity, frequency, and contrast of grating stimuli has been measured (27-30), and, more recently, as a model for investigating vertebrate sensorimotor transformations (20, 31). While zebrafish can distinguish rotational from translational optic flow to evoke appropriate optokinetic and optomotor responses (20, 32, 33), it is still unclear which regions of the visual field zebrafish preferentially observe in these behaviours. The aquatic lifestyle, in combination with the preferred swimming depths (34), might cause the lower visual field to contain

less relevant information when compared to terrestrial animals. This in turn might have resulted in behavioural biases to other –more informative– visual field regions. A corresponding systematic behavioural quantification in zebrafish, which would relate OKR behaviour to naturally occurring motion statistics and the underlying neuronal representations in retina and retino-recipient brain structures, has been prevented by technical limitations. Specifically, little is known about (i) the dependence of OKR gain on stimulus location or (ii) on stimulus sizes, (iii) possible interactions between stimulus location, size and frequency, (iv) putative asymmetries between the left and right hemispheres of the visual field, and (v) the relationship between a putative dependence of OKR on stimulus location and zebrafish retinal architecture.

In other species with large visual fields, such as Drosophila, full-surround stimulation setups have been used successfully (35-37), but to date, none has been used for zebrafish. This is at least partly due to their aquatic environment and the associated difficulties regarding the refraction of stimulus light at the air-water interface. Such distortions of shape can be partially compensated by pre-emptively altering the shape of the stimulus. However, using regular computer screens or video projection, the resulting luminance profiles remain anisotropic, potentially biasing the response toward brighter locations. Additionally, most stimulus arenas cannot easily be combined with the recording of neural activity, e.g., via calcium imaging, as stimulus light and calcium fluorescence overlap in both the spectral and time domains. These challenges must be overcome to enable full-field visual stimulation in zebrafish neurophysiology experiments (**S1b Fig**).

Here, we present a novel visual stimulus arena for aquatic animals, which covers almost the entire surround of the animal, and use it to characterize the anisotropy of the zebrafish OKR across different visual field locations as well as the tuning to stimulus size, spatial frequency and leftside versus rightside stimulus locations. We find that the OKR is mostly symmetric across both eyes and driven most strongly by lateral stimulus locations. These stimulus locations approximately correspond to a retinal region of increased photoreceptor density. By rotating the experimental setup and/or the animal, our control experiments revealed that additional extra-retinal determinants of OKR drive exist as well. Our characterization of OKR drive across the visual field will help inform bottom-up models of the vertebrate neural pathways underlying optokinetic and other visual behaviour.

Results

Spherical LED arena allows presentation of stimuli across the visual field

By combining 3D printing with electronic solutions developed in *Drosophila* vision research, we constructed a spherical stimulus arena containing 14,848 individual LEDs covering over 90% of the visual field of zebrafish larvae (**Fig 1c, S2 Text**). Using infrared illumination via an optical pathway coupled into the sphere (**Fig 2b**, **Fig 2d**), we tracked eye movements of larval zebrafish during presentation of visual stimuli (38).

To avoid stimulus aberrations at the air-to-water interface, we designed a nearly spherical glass bulb containing fish and medium. With this design, stimulus light from the surrounding arena is virtually not refracted (light is orthogonal to the air-to-water interface), and reaches the eyes of the zebrafish larva in a straight line. Thus, no geometric corrections are required during stimulus design (**S1 Code**), and stimulus luminance is expected to be nearly isotropic across the visual field. We additionally designed the setup to minimise visual obstruction, and developed a new embedding technique to immobilise the larva at the tip of a narrow glass triangle (see **Methods**). In almost all possible positions, fish can thus perceive stimuli without interference. The distance between most

of the adjacent LED pairs is smaller than the photoreceptor spacing in the larval retina (7,
39), resulting in a good spatial resolution across the majority of the spherical arena
surface (see detailed discussion in **S1 Text**). But as flat square LED tiles cannot be
perfectly arranged on a spherical surface, small triangular gaps are unavoidable. More
importantly, several gaps in LED coverage, resulting from structural elements of the
arena, were restricted mainly to the back, the top, and bottom of the animal. The "keel"
behind and in front of the fish supports the horizontal "ribs", and the circular openings in
the top and bottom accommodate the optical path for eye tracking or scanning
microscopy (**S3 Text**).

Stimulus position dependence of the optokinetic response

Horizontally moving vertical bars reliably elicit OKR in zebrafish larvae (40). We used a
stimulus which rotated clock- and counter clockwise with a sinusoidal velocity pattern
(velocity amplitude 12.5 degree/sec, frequency of the velocity envelope 0.1 Hz, spatial
frequency 0.06 cycles/degree, **Fig 3a**). OKR performance was calculated by measuring the
amplitude of the resulting OKR slow-phase eye movements after the saccades had been
removed (**Fig. 3b-d, S3 Code**, **Methods**). The OKR gain then corresponds to the speed of
the slow-phase eye movements divided by the speed of the stimulus (which is equivalent
to the ratio of the eye position and stimulus position amplitudes). To quantify position
tuning, we cropped the presented gratings (**Fig 3a**) to a disk-shaped area of constant size,
centred on one of 38 nearly equidistant parts of the visual field (**Fig 4a**, **Table 1, S1 Video**,
S2 Video). The distribution of positions was symmetric between the left and right, upper
and lower, as well as front and rear hemispheres, with some stimuli falling right on the
edge between two hemispheres. As permanent asymmetries in a stimulus arena or in its
surroundings could affect OKR gain, we therefore repeated our experiments in a second
group of larvae after rotating the arena by 180 degrees (**S1d-e Fig**), then matched the
resulting pairs of OKR gains during data analysis (see **Methods, S1h-i Fig**). Any remaining
asymmetries in the OKR distributions should result from biological lateralisation.

To overcome our spatially discrete sampling, we then fit our data with a symmetric
bimodal function comprised of two Gaussian-like two-dimensional distributions on the
stimulus sphere surface (see **Methods, S4 Code**), to determine the location of highest
OKR gain evoked by ipsilateral stimuli and contralateral stimuli, respectively. We
observed significantly higher OKR gains in response to nearly lateral stimuli, and lower
gains across the rest of the visual field (**Fig 4b-e**). OKR was strongest for stimuli near an
azimuth of 80.3 degrees and an elevation of 6.1 degrees for the left side (in body-centred
coordinates), as well as -77.0 and -2.0 degrees for the right side – slightly rostral of the
lateral meridian, and very close to the equator. Note that due to the fast stimulus speeds,
the absolute slow phase eye velocities were high, while the OKR gain was relatively low.
We chose such high stimulus speeds in order to minimize the experimental recording
time needed to obtain reliable OKR measurements for each visual field location.

As our stimulus arena is not completely covered by LEDs (**Fig 1c, Fig 1d**), some areas
remain permanently dark. These could interfere with the perception of stimuli presented
on adjacent LEDs. This is especially relevant as LED coverage is almost perfect for some
stimulus positions (near the equator), whereas the size of triangular holes increases at
others (towards the poles). We thus performed control experiments comparing the OKR
gain evoked by a stimulus in a densely-covered part of the arena to the OKR gain evoked
by same stimulus, but in the presence of additional dark triangular patches (**S1a Fig**). We
found no significant difference in OKR gain (**S1c Fig**, t-test, p<0.05). Additionally, we
performed another series of control experiments using a dark shape mimicking the dark
structural elements, the front "keel" of the arena (**S1b Fig**). Again, we found no difference
in OKR gain (**S1c Fig**, t-test, p<0.05), and thus ruled out that position dependence data
was corrupted by incomplete LED coverage. Since the eyes were moving freely in our

experiments, the range of eye positions during OKR, or so-called beating field (41), could
have changed with stimulus position. We found that animals instead maintained similar
median horizontal eye positions (e.g., left eye: -83.7±1.8 degrees, right eye: 80.3±1.9
degrees, average median ± standard deviation of medians, n=7 fish, **S2 Fig**) even for the
most peripheral stimulus positions.

A priori, it is unclear whether the sampling preference originates from the peculiarities
of the sensory periphery in the eye, or the behavioural relevance inferred by central brain
processing. The former would prioritise stimulus preference based on its position relative
to the eye and, by extension, its representation on specific parts of the retina. The latter
would prioritise stimulus preference based on its position relative to the environment,
such as a predator approaching from the water surface. To distinguish both possible
effects in the context of OKR, as well as to reveal any stimulus asymmetries accidentally
introduced during the experiment, we performed control experiments with larvae
embedded upside-down (i.e., with their dorsum towards the lower pole of the arena, **S1f**
Fig). Unexpectedly, the elevation of highest OKR gains relative to the eye changed from
slightly above to slightly below the equator of the visual field when comparing upright to
inverted fish (**S1h Fig, S1j Fig**): When upright, azimuths and body-centred elevations of
the peaks of the best fit to data were -67.8° and 8.4° for the left eye, as well as 73.1° and
6.2° for the right eye. When inverted, -88.8° and -1.2° for the left eye, as well as 80.0° and
-12.2° for the right eye. These numbers were obtained from the gains of those eyes to
which any given stimulus was directly visible. Because the set of visual stimuli presented
to inverted fish stemmed from an earlier stimulus protocol with less even sampling of the
visual field, a slight scaling of azimuths and elevations is expected. The consistent sign-
change of the elevation, however, is not. We performed a permutation test in which
embedding-direction labels were randomly swapped while stimulus-location labels were
maintained, and the Gaussian-type fit to data was then repeated on each permuted
dataset. This test confirmed that fish preferred upward (in environmental conditions)
rather than dorsalward elevations (p < 0.05, **S5 Code**).

Adjustment by the fish of its vertical resting eye position between the upright and
inverted body positions would have been a simple potential explanation for this result.
However, time-lapse frontal microscopy images (**Methods**) ruled this out, since for both
upside-up and upside-down embedding the eyes were inclined by an average of about 4
degrees towards the dorsum (3.5±1.0° for the left eye, 4.9±0.8° for the right eye, mean ±
s.e.m., **S3 Fig**). We also tested the influence of camera and infrared light (840 nm)
positions (**S1g, Fig**) – which in either case should have been invisible to the fish (42) –
and found that they could indeed not explain the observed differences. As the body-
centred preferred location in upside-down embedded fish flipped from slightly dorsal to
slightly ventral (**S1j Fig**), and thus remained virtually unchanged in environmental
coordinates, optokinetic stimulus location preference appears to be related to the
behavioural relevance of these stimulus positions, and cannot merely be caused by retinal
feedforward circuitry.

Yoking of the non-stimulated eye

Almost all stimuli were presented monocularly – that is, in a position visible to only one
of the two laterally located eyes. Without exception, zebrafish larvae responded with
yoked movements of both the stimulated and unstimulated eye. To rule out reflections of
stimuli within the arena, we performed a series of experiments in which the unstimulated
side of the glass bulb had been covered with a matte, black sheet of plastic. Reflections on
the glass-air interface would otherwise cause monocular stimuli (that should only be
visible to the ipsilateral eye) to also be seen by the contralateral eye. Yoking indices (YI)
were significantly different between the regular monocular setup (YI≈0.7) and the control
setup (YI≈0.2) containing the black surface on the side of the unstimulated eye,

confirming that yoking indices had been affected by reflections (**S4 Fig**, an index of 1
indicated completely monocular eye movements, an index of 0 perfectly conjugate eye
movements/yoking). This suggests a crucial role for sharp reflections of the stimulus
pattern at the glass-to-air or water-to-air interface (Arrenberg et al., unpublished) in our
spherical setup and other commonly used stimulus arenas. We performed additional
control experiments using a previously described setup (38) with four flat LCD screens
for stimulus presentation in a different room. In these experiments, stimuli were
presented monocularly or binocularly, and the unstimulated eye was either (i) stimulated
with a stationary grating (**S5a-b Fig**), (ii) shielded with a blank, white shield placed
directly in front of the displays (**S5c-d Fig**) or (iii) shielded with a matte, black sheet of
aluminium foil placed inside the petri dish (control for possible reflections on the Petri
dish wall) (**S5e-f Fig**). This experiment showed that yoking was much reduced (YI≈0.3) if
the non-stimulated eye saw a stationary grating (i) instead of the white or black shields
(ii-iii, YI≈0.1) or a binocular motion stimulus (YI≈0) (**S5g-h Fig**, p<0.05).

Spatial asymmetries

As multiple previous studies reported left-right asymmetries in zebrafish visuomotor
processing and behaviour other than OKR (43-45), we computed an asymmetry index B
(**Methods**) to reveal whether zebrafish OKR is lateralised in individuals or across the
population. We did not observe a general asymmetry between the response of the left and
right eyes. Rather, our data is consistent with three distinct sources of asymmetry:
individual bias towards one eye, shared bias across individuals, and asymmetries induced
by the environment (including the experimental setup and stimulus arena). Through
multivariate linear regression, we fit a linear model of asymmetries to our data
(**Methods**), which combined data from fish embedded upside-up (**Fig. 4, S1d Fig, S1h
Fig**), upside-down (**S1f Fig, S1j Fig**) and data obtained with the arena rotated relative to
the fish (**S1e Fig, S1i Fig**). Regression coefficients for external causes of asymmetry were
similar to or smaller than those for biological causes (**S6 Fig**), and individual biases from
fish to fish were broadly and symmetrically distributed from left to right (mean coefficient
$3.7 \cdot 10^{-4} \pm 120.0 \cdot 10^{-4}$ st. dev., n =15), so that no evidence was found for a strong and
consistent lateralisation of OKR behaviour across animals (**S6 Fig**).

Our results show that the OKR behaviour is mostly symmetric across both eyes, with
individual fish oftentimes having a dominant eye due to seemingly random bias for one
eye (lateralisation) across fish. Some of the observed asymmetries are consistent with
external factors. Therefore, the OKR gains presented in **Fig 4** have been corrected in order
to present only biologically meaningful differences (**Methods**).

Spatial frequency dependence of the optokinetic response

We investigated the spatial frequency tuning of OKR behaviour across visual field
positions by presenting 7 different spatial frequencies of the basic stimulus, each cropped
into a planar angle of 40 degrees, at different visual field locations. Because we held the
temporal frequency constant, stimulus velocity decreased whenever spatial frequency
increased. These 7 disk-shaped stimuli were presented while centred on one of 6 possible
locations in different parts of the visual field, with 3 locations on each hemisphere: one
near the location of highest OKR gain as determined in our experiments on position
dependence, one in a nasal location, and one in a lower temporal location. In total, we thus
presented 42 distinct types of stimuli (**Table 2**, **S3 Video**). For each stimulus location and
eye, the highest OKR gain was observed at a spatial frequency of 0.03 to 0.05
cycles/degree (**Fig 5a, Fig 5c-d**). We did not observe any strong modulation of frequency
dependence by stimulus location.

Size dependence of the optokinetic response

It is unclear to what extent small stimuli are effective in driving OKR. We therefore employed a stimulus protocol with 7 OKR stimuli of different covered areas on the sphere. Spatial and temporal frequencies were not altered, so bars appeared with the same width and velocity profile in all cases. These 7 disk-shaped stimuli were presented while centred on one of 6 possible locations, identical to those used to study frequency dependence, again yielding 42 unique stimuli (**Table 3**, **S4 Video**). Stimulus area size was chosen at logarithmic intervals, ranging from stimuli almost as small as the spatial resolution of the zebrafish retina, to stimuli covering the entire arena. In line with many other psychophysical processes, OKR gain increased sigmoidally with the logarithm of stimulus size (**Fig 5b-c, Fig 5e**). Weak OKR behaviour was already observable in response to very small stimulus diameters (e.g. $10.4° − 0.8$ %), and reached half-maximum performance at a stimulus size of roughly 120° (a quarter of the entire surrounding space). As was the case for spatial frequency dependence, we did not observe any strong modulation of size dependence by stimulus location, although OKR gains of the left eye appeared more dependent on stimulus location than those of the right eye.

Optokinetic response gain covaries with retinal density of long-wave sensitive photoreceptors

We hypothesized that the non-uniform distribution of the OKR gain is related to the surface density of photoreceptors and investigate this using data from a recent study (12) on photoreceptor densities in explanted eye cups of 7-8 day old zebrafish larvae. As shown in **Fig 6b**, ultraviolet receptor density exhibits a clear peak in the upper frontal part of the visual field, whereas red, green and blue receptors (**Fig 6a**) are most concentrated across a wider region near the intersection of the equator and lateral meridian, with a bias to the upper visual field (in body coordinates). For comparison, density maps in retinal coordinates, not body coordinates, are shown in **S7 Fig**. To register our OKR gain data onto the photoreceptor density maps, we took the average eye position into account, which was located horizontally at -84.8±6.2 degrees azimuth for the left and 80.1±6.5 deg for the right eye (mean±st.dev., n=7 fish), and vertically at 3.5±3.2 degrees elevation for the left and 4.9±2.7 deg for the right eye (n=10 fish). For green, blue and especially red receptors, the stimulus centred on the position of maximum OKR gain, as inferred from our oculomotor experiments (**Fig 4e**), covers a region of near-maximum photoreceptor density (white ring in **Fig 6**). For ultraviolet receptors, there is no strong correlation between photoreceptor density and OKR gain.

Discussion

The spherical arena introduced here covers a large proportion of the surround and therefore lends itself to many other investigations of zebrafish and other species with limited visual acuity. In comparison to other feasible technical solutions, such as video projections setups, our spherical LED array stimulus setup provides homogeneous light and contrast across the entire stimulation area. Thereby stimulus design becomes much easier since the stimulus warping and conditioning becomes unnecessary. When combined with calcium imaging in a scanning microscope, the use of LED arrays provides the additional advantage that the visual stimulus can be controlled with high temporal precision, fast enough to interlace visual stimuli and line scans.

Despite the common notion that OKR is a whole-field gaze stabilisation behaviour, our results show that the OKR can be driven effectively by moving stimuli that cover only small parts of the spherical surface (with a half-maximum OKR gain at around 25 % of the surface). Our experiment on spatial frequency dependence further demonstrates that the

spatial frequency tuning of the OKR is similar across retinal locations. Here we suggest
two plausible explanations, (1) existing photoreceptor density differences are
compensated for centrally in visual brain areas mediating the OKR, or (2) the
photoreceptor density is simply not the limiting factor for OKR performance in this
frequency range.

Previous reports indicated that the zebrafish visual system is lateralised with the left eye
preferentially assessing novel stimuli, while the right eye being associated with decisions
to respond (43, 46). We therefore investigated whether there are consistent behavioural
asymmetries for the OKR and observed almost no consistent, inter-individual
asymmetries in OKR between the left and right hemispheres of the visual field, other than
those induced by external conditions. Individual fish, however, show a wide and
continuous range of biases towards either hemisphere.

We measured OKR gain in larvae at 5-7 days post fertilisation (dpf) of age, whereas our
data on photoreceptor densities corresponds to slightly older, 7-8 dpf larvae. Owing to
their rapid development, zebrafish undergo noticeable morphological changes on this
timescale, but the zebrafish retina itself is known to be well developed by 5 dpf (47) and
stable OKR behaviour is exhibited from then on. Crucially, we did not observe a salient
age-dependent spatial shift of maximum OKR gain between our 5 dpf and 7 dpf larvae
(data not shown).

The qualitative match between red cone retinal photoreceptor densities and the stimulus
position driving the highest OKR gains may provide a mechanistic bottom-up explanation
of the gradual differences associated with OKR. The correspondence of red photoreceptor
density with the visual field map of OKR gain is consistent with the fact that our LEDs emit
light at 568 nm peak power, which should have activated the red cones most. Our data is
also in agreement with observations in other species, that the OKR drive is strongest
when the moving stimulus covers the central visual field (9, 10, 22). In a simplistic,
additive view of visual processing, increased numbers of receptors would be triggered by
incident light, gradually leading to stronger activation of retinal ganglion cells and
downstream circuits, eventually driving extraocular eye muscles towards higher
amplitudes. Instead, or in addition, the increased resolution offered by denser
distributions of photoreceptors could help reduce sensory uncertainty (and increase
visual acuity). It is unclear however, how more uncertainty would lead to consistently
lower OKR gains instead of a repeated switching between periods of higher and lower
gains. If sensory uncertainty were indeed crucial to OKR tuning, presenting blurred or
otherwise deteriorated stimuli should reduce OKR gain in disfavoured locations more
strongly than those in favoured locations. It is also possible that correlations between
OKR gain and photoreceptor density are entirely coincidental, as our spatial frequency
tuning results for different stimulus locations had implied. Genetic zebrafish variants with
altered photoreceptor distributions would thus be a valuable tool for further studies.

The pronounced increase in OKR gain for nearly lateral stimulus locations raises
questions regarding the top-down behavioural significance of these directions in the
natural habitat of larval zebrafish. While reduced OKR gains near the limits of the visual
field might be expected, we show that gains are also reduced in the frontal binocular area,
as well as in upper and lower visual field locations. Interestingly, when animals were
mounted upside-down, they still prefer stimulus locations just above the equator of the
environment. This result cannot be explained by shifted resting vertical eye positions in
the inverted animal, which we have measured. Instead, it could potentially be explained
by multimodal integration, where body orientation appears to influence the preferred
OKR stimulus locations via the vestibular system (48-50). Furthermore, it seems possible
that the unequal distribution of OKR gains across the visual field is related to the optic
flow statistics that naturally occur in the habitats of larval zebrafish (12, 51-54). For
another stabilisation behaviour of zebrafish, the optomotor response (21), we have

recently shown that the underlying circuits prefer stimulus locations in the lower temporal visual field to drive forward optomotor swimming (55). Therefore, the optokinetic and the optomotor response are preferentially driven by different regions in the visual field, suggesting that they occur in response to different types of optic flow patterns in natural habitats. Both the optokinetic and the optomotor response (OKR, OMR) are thought to be mediated by the pretectum (20, 32), and we therefore hypothesize that circuits mediating OKR and OMR segregate within the pretectum and form neuronal ensembles with mostly different receptive field centre locations. Future studies on pretectal visual feature extraction in the context of naturalistic stimulus statistics are needed in order to establish a more complete picture of the visual pathways and computations underlying zebrafish OKR, OMR and other visually mediated behaviours.

Methods

Animal experiments

Animal experiments were performed in accordance with licenses granted by local government authorities (Regierungspräsidium Tübingen) in accordance with German federal law and Baden-Württemberg state law. Approval of this license followed consultation of both in-house animal welfare officers and an external ethics board appointed by the local government. We used *mitfa-/-* animals (5-7 dpf) for the experiments, because this strain lacks skin pigmentation that could interfere with eye tracking.

Coordinate systems and conventions

To remain consistent with the conventions adopted to describe stimuli and eye positions in previous publications, we adopted an East-North-Up, or ENU, geographic coordinate system. In this system, all positions are relative to the fish itself, and expressed as azimuth (horizontal angle, with positive values to the right of the fish), elevation (vertical angle, with positive values above the fish), and radius (or distance to the fish). The point directly in front of the fish (at the rostrum) is located at [0°, 0°] azimuth and elevation. Azimuth angles cover the range [-180°, 180°] and elevation angles [-90°, 90°]. Azimuth sign is opposite to the conventional mathematical annotation of angles when looking top-down onto the fish. Supplementary materials provide a detailed description of the coordinate systems used, and for transformations between Cartesian and geographic coordinate systems, please consult the supplementary material (**S1 Text**).

Design of the spherical arena

Geometric design of the arena. The overall layout of the spherical arena was optimised to contain the near maximum number of LED tiles that can be driven by our hardware controllers (232 out of a possible 240), and arrange them with minimal gaps in between. Also, care was taken to leave sufficient gaps near the top and bottom poles to insert the optical pathway used to illuminate and record fish behaviour. A further 8 LED tiles could be included as optional covers for the top and bottom poles, bringing the total number to 240 out of 240 possible. A detailed walkthrough of the mathematical planning is found in the supplementary material (**S1 Text**).

Arena elements. The arena consists of a 3D-printed structural scaffold; green light emitting LED tiles (Kingbright TA08-81CGKWA, 20x20 mm each, peak power at 568 nm) hot-glued to the scaffold and connected by cable to a set of circuit boards with hardware

controllers (**Fig 2d**); 8x8 individual LEDs contained in each tile (**Fig 1f**); a nearly spherical
glass bulb filled with water, into which the immobilised larvae are inserted (**Fig 2c**,
middle); a metal rotation mount attached to the scaffold "keel" of the arena (**Fig 2c**, right),
holding the glass bulb in place and allowing corrections of pitch and roll angles; the optical
pathway with an infrared light source to illuminate the fish from below (**Fig 2b**), and a
USB camera for video recording of the transmission image (**Fig 2d**).

Electronics and circuit design. To provide hardware control to the LEDs, we used circuit
boards designs and C controller code provided by Alexander Borst (MPI of Neurobiology,
Martinsried) and Väinö Haikala and Dierk Reiff (University of Freiburg) (56). Any custom
circuit board design and code could be substituted for these, and alternative solutions
exist, e.g., in Drosophila vision research (57). At the front end, these electronics control
the 8x8 LED matrices, which are multiplexed in time to allow control of individual LEDs
with just 8 input and 8 output pins.

Optical pathway, illumination and video recording. A high-power infrared LED was
placed outside the stimulus arena and its light diffused by a sheet of milk glass and then
guided towards the fish through the top hole of the arena (**Fig 2b**, **Fig2d**). Non-absorbed
IR light exits through the bottom hole, where it is focused onto an IR-sensitive camera.
Between the arena and the proximal lens, a neutral density filter (NE13B, Thorlabs, ND
1.3) was inserted half-way (off-axis) into the optic pathway using an optical filter slider
(CFH2/M, Thorlabs, positioned in about 5 cm distance of the camera CCD chip) to improve
image contrast (oblique detection). We used the 840nm, 125 degree IR emitter
Roschwege Star-IR840-01-00-00 (procured via Conrad Electronic GmbH as item 491118-
62) in custom casing, lenses LB1309 and LB1374, mirror PF20-03-P01 (ThorLabs GmbH),
and IR-sensitive camera DMK23U618 (TheImagingSource GmbH). Approximate
distances between elements are 14.5cm (IR source to first lens), 12cm (first lens to centre
of glass bulb), 22cm (bulb centre to mirror centre), 8.5cm (mirror centre to second lens),
28.5 cm (second lens to camera objective).

Fish mounting device. Larvae were mounted inside a custom-built glass bulb (**Fig 2c,
middle**). Its nearly spherical shape minimises reflection and refraction at the glass
surface. It was filled with E3 solution, so there was no liquid-to-air boundary distorting
visual stimuli. Through an opening on one side, we inserted a glass rod, on the tip of which
we immobilise the larva in agarose gel (see description of the embedding procedure
below). The fish was mounted in such a way that the head protruded the tip of the narrow
triangular glass stage, which ensured that visual stimuli are virtually unobstructed by the
glass triangle on their way to the eyes (**Fig 2c, left**). The entire glass structure was held
at the centre of the spherical arena by metal parts attached to the arena scaffold itself (**Fig
2c, right**). Care was taken to remove air bubbles and completely fill the glass bulb with
E3 medium.

Computer-assisted design and 3D printing. To arrange the square LED tiles across a
nearly spherical surface, we 3D-printed a structural scaffold or "skeleton", consisting of a
reinforced prime meridian major circle ("keel") and several lighter minor circles of
latitude (**Fig 2g**). Available hardware controllers allow for up to 240 LED matrices in
parallel, so we chose the exact size of the scaffold (106.5 mm in diameter) to hold as many
of these as possible while minimising gaps in between. As individual LEDs are arranged
in a rectangular pattern on each of the flat LED tiles, and stimuli defined by true meridians
(arcs from pole to pole, or straight vertical lines in Mercator projection), pixelation of the
stimulus is inevitable, and stimulus edges become increasing stair-shaped near the poles.
Because of the poor visual acuity of zebrafish larvae (see **S1 Text**), this should not affect
OKR behaviour. Our design further includes two holes necessary for behavioural
recordings and two-photon imaging, located at the North and South poles of the sphere.
We placed the largest elements of the structural scaffold behind the zebrafish (**Fig 2**).

Given the ~160° azimuth coverage per eye in combination with a slight eye convergence
at rest, this minimises the loss of useful stimulation area.

We printed all structures out of polylactide (PLA) filament using an Ultimaker 2 printer
(Ultimaker B.V.). Parts were assembled using a hot glue gun.

Visual field coverage

561 We can estimate the fraction of the visual field effectively covered by LEDs based on a
562 projection of LED tiles onto a unit sphere. The area A of a surface segment delimited by
563 the projection of the edges of a single tile onto the sphere centre is given by

$$A(S) = \oint dA = \int_{-\lambda}^{\lambda} dx \int_{-\lambda}^{\lambda} dy \| u_\alpha \times u_\beta \|$$

565 where u_α and u_β are the Cartesian unit vectors spanning the tile itself and (±λ, ±λ) is the
566 Cartesian position of the four edges of another rectangle. This smaller rectangle is the
567 straight projection of the sphere segment onto the tile,

$$\lambda = \sin(\tan^{-1}(D/2R_S))$$

where $R_S = 106.5\ mm$ is the sphere radius and $D = 21\ mm$ is the length of the edges of
the tile. Summing over the number of tiles included in the arena, the equations above can
be used to estimate the total coverage of the sphere by its square LED tiles to around
66.5% of the surface area. Using this strict estimate, the small gaps in between LED arrays
are counted as not covered, even though we successfully demonstrated that they are small
enough not to affect OKR performance, likely due to the low visual acuity of zebrafish
larvae. A more meaningful estimate of coverage must take these results into account (**S2
Text**), and in fact reveals that stimuli presented with our LEDs effectively cover 85.6% of
all possible directions. In core parts of the visual field, coverage exceeds 90%.

Stimulus design

We designed visual stimuli, transformed them to geographical coordinates, and mapped
them onto the physical positions of each individual LED with custom MATLAB software.
We have made this code available for free under a Creative Commons NC-BY-SA 4.0
license (**S1 Code**). The mapped stimulus was then uploaded to the hardware controllers
using custom-built C code originally developed by Väinö Haikala.

To investigate OKR gain dependence on stimulus location, we chose to present stimuli
centred on 36 different locations distributed nearly equidistantly across the spherical
arena, as well as symmetrically distributed between the left and right, upper and lower,
front and rear hemispheres (**Fig 4a**). These positions were determined numerically: First,
we populated one eighth of the sphere surface by placing one stimulus centre at a fixed
location at the intersection of the equator and the most lateral meridian (90 degrees
azimuth, 0 degrees elevation), constraining two more stimulus centres to move along this
lateral meridian (90 degrees azimuth, initially random positive elevation), constraining
yet another stimulus centre to move along the equator (initially random positive azimuth,
0 degrees elevation), and allowing three more stimulus centre to move freely across the
surface of this eighth of the sphere (initially random positive azimuth and elevation), for
a total of 7 positions. Second, we placed additional stimulus centres onto all 29 positions
that were mirror-symmetric to the initial 7, with mirror planes placed between the six
hemispheres listed above. We then simulated interactions between all 38 stimulus
centres akin to electromagnetic repulsion, until a stable pattern emerged. Resulting
coordinate values were rounded for convenience (**S2 Code, S1 Video**).

Embedding procedure

To immobilise fish on the glass tip inside the sphere, we developed a novel embedding method. A cast of the glass triangle (and of the glass rod on which it is mounted) was made by placing it inside a Petri dish, which was then filled with a heated 2% agarose solution. After agarose cooled down and polymerised, agarose within a few millimetres of the tip of the glass triangle was manually removed, before removing the triangle itself. The resulting cast was stored in a refrigerator and then used to hold the glass triangle during all subsequent embedding procedures, limiting the freedom of movement of the larva to be embedded. The triangle was stored separately at room temperature. Before each embedding, we coated the glass triangle with polylysine and dried it overnight in an incubator at 29 degrees Celsius to increase the subsequent adhesion of agarose. We then returned the glass triangle into its cast, and constructed a tight, 2 mm high circular barrier around its tip using pieces of congealed agarose. A larva was picked up with as little water as possible using a glass pipette and very briefly placed inside 1 ml of 1.6% low-melting agarose solution at 37 degrees Celsius. Using the same pipette, the larvae was then transferred onto the glass triangle along with the entire agarose. After the larva had been placed a few millimetres away from the tip of the glass triangle, the orientation of the animal could be manipulated with custom-made platinum wire tools without touching its body, as previously described (58). Before the agarose congeals, swimming motions of the animal were exploited to guide it towards the tip and ensure an upright posture. The final position of the fish was chosen as such that its eyes are aligned with the axis of the glass rod, its body is upright without any rotation, and its head protrudes forward from the tip of the glass triangle, maximising the fraction of its field of view unobstructed by glass elements. The agarose was left to congeal, and the Petri dish was filled with in E3 solution. The freshly congealed agarose surrounding the glass triangle was then removed using additional, flattened platinum wire tools, once again separating the glass triangle from the cast. Using the same tools, we finally cut triangular holes into the remaining agarose to completely free both eyes. To ensure free movement of both eyes, we confirmed the presence of large and even optokinetic eye movements using a striped paper drum before the experiment.

We then pick up the glass triangle by the glass rod attached to it, cut off any remaining agarose detritus, and place it inside the E3-filled glass bulb. No air remained in the bulb, and no pieces of detritus were introduced in to the bulb, as these would accumulate near the top and bottom of the bulb, respectively, interfering with the optical pathway and thus reduce image quality.

Data analysis

Video images of behaving zebrafish larvae were processed in real time using a precursor of the *ZebEyeTrack* software (38), available from www.zebeyetrack.com. The resulting traces of angular eye position were combined with analogue output signals from the hardware controllers of the spherical arena to match eye movement to the various stimulus phases. This was achieved using custom-built MATLAB software, which is freely available under a Creative Commons NC-BY-SA 4.0 license (**S3 Code**).

Data was then analysed further by detecting and removing saccades and fitting a piece-wise sinusoidal function to the eye position traces. The parameters of the fit were then compared to the parameters of the equally sinusoidally changing angular positions of the stimulus. For each fish, eye, and stimulus phase, the ratio between the amplitude of the fit to eye position and the amplitude of stimulus position represents one value of the gain of the optokinetic response.

648 For each interval between two subsequent saccades, or inter-saccade-interval (ISI), the
649 fit function to the eye position data is defined by

$$f(t \in ISI_k) = -c_1 \cos(c_2 t + c_3) + c_{k+3}$$

Here, t are the time stamps of data points falling within the k-th ISI, c_1, c_2 and c_3 are the
amplitude, frequency and phase shift of oscillation across all ISIs, and c_{k+3} is a different
constant offset within each ISI, which corrects for the eye position offsets brought about
by each saccade. The best fit value c_1 was taken as an approximation of the amplitude a_E
of eye movement, $a_E \approx c_1$. The process of cropping saccades from the raw data and fitting
a sinusoid to the remaining raw data is demonstrated in **Fig 3**.

The OKR gain g is a common measure of visuomotor function. It is defined as the ratio
between the amplitude a_E of eye movement and the amplitude a_S of the visual stimulus
evoking eye movement,

$$g = \frac{a_E}{a_S} = \frac{c_1}{a_S}$$

In other words, OKR gain indicates the degree to which zebrafish larvae track a given
visual stimulus. For each eye, a single gain value per stimulus phase is computed. While a
value of 1 would indicate a "perfect" match between eye movement and stimulus motion,
zebrafish larvae at 5 dpf often exhibit much lower OKR gains (30). While highest gains are
obtained for very slowly moving stimuli, in our experiments, we chose higher stimulus
velocities. Although these velocities are only tracked with small gains, the absolute
velocities of the eyes are high, which allowed us to collect data with high signal-to-noise
levels and reduce the needed recording time.

To rule out asymmetries induced by the arena itself or by its surroundings, we recorded
two sets of stimulus-position-dependence data, one with the arena in its original
configuration, and another with the arena rotated by 180 degrees (**S1h-i Fig**). Each set
contained data from multiple larvae, and with at least 2 separate presentations of each
stimulus position. For each stimulus position, and separately for both sets of data, we
computed the median OKR gain across fish and stimulus repetitions. We then averaged
between the two datasets, yielding a single OKR gain value per stimulus position. As
asymmetries are less crucial when studying stimulus frequency and size (**Fig 5**), we did
not repeat those with a rotated arena, and could thus omit the final step of the analysis.

Von Mises-Fisher fits to data

Based on the assumption that OKR position tuning could be normally distributed with
respect to each angle, OKR gain would be approximated by a two-dimensional, circular
von Mises-Fisher function centred on the preferred stimulus location. Because the eyes
are yoked, the OKR gain of one eye will be high around its own preferred position, as well
as around the preferred position of the contralateral eye. To account for this, we fit the
sum of two independent von Mises-Fisher functions to our OKR gain data:

$$F(\alpha, \beta) = \frac{C_1 \kappa_1 \exp(\kappa_1 \mu_1{}^T \xi)}{2\pi(\exp(\kappa_1) - \exp(-\kappa_1))} + \frac{C_2 \kappa_2 \exp(\kappa_2 \mu_2{}^T \xi)}{2\pi(\exp(\kappa_2) - \exp(-\kappa_2))} + C_3$$

686 Here, ξ is the Cartesian coordinate vector of a point on the sphere surface, and
687 corresponds to the geographic coordinates azimuth α and elevation β. μ_1 and μ_2 are
688 Cartesian coordinate vectors pointing to the centre of the two distributions, and κ_1 and
689 κ_2 express their respective concentrations, or narrowness. The parameters μ_j, κ_j, the
690 amplitudes C_1, C_2 and the offset C_3 are fit numerically.

Yoking index, asymmetry and mathematical modelling

To quantify asymmetries in the gain between left and right, stimulated and unstimulated eyes, we introduce the yoking index

$$Y = \frac{g_L - g_R}{g_L + g_R}$$

Here, g_L and g_R are the OKR gains of the left eye and right eye, measured during the same stimulus phase. Depending on stimulus phase, only the left eye, only the right eye or both eyes may have been stimulated. If the yoking index is positive, the left eye responded more strongly than the right eye; if it is negative, the amplitude of right eye movement was larger. An index of zero indicates "perfect yoking", i.e. identical amplitudes for both eyes.

In addition, we define a "bias" index to capture innate or induced asymmetries between responses to stimuli presented within the left or right hemisphere of the visual field,

$$B = \frac{m_L - m_R}{m_L + m_R}$$

Here, m_L and m_R are the medians of OKR gains after pooling across either all left-side or all right-side stimulus types (D1-D19 and D20-D38, respectively). Several sources of asymmetry contribute to B: (1) arena- or environment-related differences in stimulus perception, constant across individuals; (2) a biologically encoded preference for one of the two eyes, constant across individuals; (3) inter-individual differences between the eyes, constant across stimulus phases for each individual; (4) other sources of variability unaccounted for, and approximated as a noise term η. We hypothesise that the overall asymmetry observed for each larva k is given by a simple linear combination of these contributions,

$$B_k = \varphi b_1 + b_2 + b_{3,k} + \eta$$

The parameter φ is 1 for the default arena setup, and –1 during control experiments with a horizontally flipped arena setup. To determine b_1, b_2 and b_3, we fit this system of equations by multivariate linear regression to experimentally observed bias indices. The system is initially underdetermined, as it contains $n + 2$ coefficients for every n fish observed. However, if we assume that individual biases average out across the population, we can determine the population-wide coefficients b_1 and b_2 by setting aside the individual $b_{3,k}$ for a first regression. To determine how far each individual deviates from the rest of the population, we then substitute their best regression values of b_1 and b_2 into the full equation, and perform a second regression for the remaining $b_{3,k}$.

Supplementary information

Supplementary Code

S1 Code. (S1Code.zip) MATLAB code to design visual stimuli, convert them to geographic coordinates, and map them onto the actual position of individual LEDs.

S2 Code. (S2Code.zip) MATLAB code to numerically identify a distribution of nearly equidistant stimulus centres that is symmetric between the left and right, upper and lower, as well as front and rear hemispheres.

S3 Code. (S3Code.zip) MATLAB code to read raw eye traces, identify individual stimulus phases, detect and remove saccades, compute piece-wise fits to cropped and pre-processed raw data, and return OKR gains.

S4 Code. (S4Code.zip) MATLAB code to recreate the results figures from data, especially **Fig 4**, **Fig 5**, **Fig 6**.

S5 Code. (S5Code.zip) MATLAB code to assess the significance of differences between the best fits to data for fish embedded upright or upside-down. This code requires access to the raw data repository.

Supplementary Manuals

S1 Text. (S1Text.pdf) Spatial resolution and visual acuity

S2 Text. (S2Text.pdf) Coverage of visual field by stimulus arena

S3 Text. (S3Text.pdf) Mathematical appendix on the design of and stimulus mapping onto the spherical arena, written as a manual.

Supplementary Figures

S1 Fig. Control experiments investigating the effect of artificial triangular holes, an artificial keel, upside-down embedding, and arena effects.

S2 Fig. OKR beating field and average eye position are independent of stimulus location.

S3 Fig. Vertical eye position under upright and upside-down embedding.

S4 Fig. Yoking indices are biased by reflections within the arena.

S5 Fig. Reflections alter perceived yoking indices across arena types.

S6 Fig. Individual fish exhibit weak and broadly-distributed biases towards the left or right half of their visual field.

S7 Fig. Retinal cone densities in retinal coordinates, instead of visual field coordinates.

Supplementary Tables

S1 Table. Arena cross-section.

S2 Table. Absolute positions of fish and setup elements in different experiments.

S3 Table. Stimulus parameters (control experiments).

S4 Table. Stimulus parameters (whole-field and hemispheres).

Supplementary Videos

S1 Video. (stimulusdistribution.avi) Repulsion-based algorithm to numerically distribute stimulus centres across a sphere surface (**S2 Code**), with gradual convergence on a logarithmic timescale (blue to green). A subset of stimulus centres is held on the equator at zero elevation (orange).

S2 Video. (locationvideo.avi) Animation showcasing short samples of all disk stimuli used to study location dependence, as in **Fig 3a**.

S3 Video. (frequencyvideo.avi) Animation showcasing short samples of all disk stimuli used to study frequency dependence, as in **Fig 4a**.

S4 Video. (sizevideo.avi) Animation showcasing short samples of all disk stimuli used to study size dependence, as in **Fig 4b**.

Figures and tables

Figure legends

Figure 1. Presenting visual stimuli across the visual field. (a) When presented with a horizontal moving stimulus pattern, zebrafish larvae exhibit optokinetic response (OKR) behaviour, where eye movements track stimulus motion to minimise retinal slip. Its slow phase is interrupted by intermittent saccades, and even if only one eye is stimulated (solid arrow), the contralateral eye is indirectly yoked to move along (dashed arrow). (b) Often, experiments on visuomotor behaviour such as OKR sample only a small part of the visual field, whether horizontally or vertically. As different spatial directions may carry different behavioural importance, an ideal stimulation setup should cover all or most of the animal's visual field. For zebrafish larvae, this visual field can be represented by an almost complete unit sphere. (c) We arranged 232 LED tiles with 64 LEDs each across a spherical arena, such that 14,484 LEDs (green dots) covered nearly the entire visual field. (d) The same individual positions, shown in geographic coordinates. Each circle represents a single LED. Each cohesive group of eight-by-eight circles corresponds to the 64 LEDs contained in a single tile. (e) To identify LED and stimulus locations, we use Up-East-North geographic coordinates: Azimuth α describes the horizontal angle, which is zero in front of the animal and, when seen from above, increases for rightward position. Elevation β refers to the vertical angle, which is zero throughout the plane containing the animal, and positive above. (f) The spherical arena is covered in flat square tiles carrying 64 green LEDs each. (g) Its structural backbone is made of a 3D-printed keel and ribs. Left and right hemispheres were constructed as separate units. (h) Across 85-90% of the visual field, we can then present horizontally moving bar patterns of different location, frequency and size to evoke OKR.

Figure 2. A spherical LED arena to present visual stimuli across the visual field. (a) LED tiles are arranged in ribbons parallel to the equator, and glued in between structural ribs. Gaps at the top and bottom pole of the sphere allow coupling in an optical pathway for infrared illumination and subsequent video recording of eye movement. (b) Optical pathway for eye movement tracking. (c) To minimise obstruction and refraction, the zebrafish larva is immobilised on the tip of a glass triangle (left) using agarose, which is then inserted into the centre of a spherical glass bulb (middle). This bulb is then mounted into a metal holder (right) and thus placed at the centre of the sphere. (d) Image of the two hemispheres and the camera setup. One hemisphere is mounted on a rail to allow opening and closing the arena.

Figure 3. OKR gain is inferred from a piece-wise fit to the slow phase of tracked eye movements. (a) We present a single pattern of horizontally moving bars to evoke OKR (left). Its velocity follows a sinusoidal time course, repeating every 10 seconds for a total of 100 seconds for each stimulus phase (right). (b) OKR gain is the amplitude of eye movement (grey trace) relative to the amplitude of the sinusoidal stimulus (green trace). The OKR gain is often well below 1, e.g. for high stimulus velocities as used here (up to 12.5°/s). (c) OKR eye movements consist of a slow phase, gradually tracking stimulus motion, and intermittent saccades. (d) After pre-processing data to detect and remove saccades, we fit a piece-wise sinusoidal function with a single amplitude to the remaining slow-phase eye traces. The amplitude of the best fit determines OKR gain.

Figure 4. OKR gain depends on stimulus location. (a) The stimulus is cropped to a disk-shaped area 40 degrees in diameter, centred on one of 38 nearly equidistant locations across the entire visual field (left), to yield 38 individual stimuli (right). (b-d) Dots reveal the location of stimulus centres D1-D38. Their colour indicates the average OKR gain across individuals and trials, corrected for external asymmetries. Surface colour of the sphere displays the best von-Mises Fisher fit to the discretely sampled OKR data. Top row:

OKR gain of the left eye (b), right eye (d), and the merged data including only direct
stimulation of either eye (c), shown from an oblique, rostrodorsal angle. Bottom row:
same, but shown directly from the front. OKR gain is significantly higher for lateral
stimulus locations and lower across the rest of the visual field. The spatial distribution of
OKR gains is well explained by the bimodal sum of two von-Mises Fisher distributions. (e)
Mercator projections of OKR gain data shown in panels (b-d). White and grey outlines
indicate the area covered by each stimulus type. Numbers indicate average gain values
for stimuli centred on this location. Red dots show mean eye position during stimulation.
Dashed outline and white shading on panels (b, d, e) indicate indirect stimulation via
yoking, i.e., stimuli not directly visible to either the left or right eye. Data from n=7 fish for
the original configuration and n=5 fish for the rotated arena,

Figure 5. OKR gain depends on stimulus size and frequency. (a) Patterns with 7
different frequencies were cropped to disks of a single size. These disks were placed in 6
different locations for a total of 42 stimuli. cpd: cycles per degree. (b) Patterns with
identical spatial frequencies were cropped to disks of 7 different sizes. These disks were
also placed in 6 different locations for another set of 42 stimuli. Degrees indicate planar
angles subtended by the stimulus outline, so 360° correspond to whole-field stimulation.
(a, b) Displaying the entire actual pattern at the size of this figure would make the
individual bars hard to distinguish. We thus only show a zoomed-in version of the
patterns in which 45 out of 360 degrees azimuth are shown. (c) Coloured dots indicate
the 6 locations on which stimuli from a and b were centred, shown from above (top), from
front (middle), and from an oblique angle (bottom). (d) OKR gain is unimodally tuned to
a wide range of spatial frequency (measured in cycles per degree). (e) OKR gain increases
sigmoidally as the area covered by the visual stimulus increases logarithmically (a
stimulus size of 1 corresponds to 100% of the spherical surface). (d-e) Colours
correspond to the location of stimulus centres shown in (c). There is no consistent
dependence on stimulus location of either frequency tuning or size tuning. Data from n=7
fish for frequency dependence and another n=7 fish for size dependence.

Figure 6. Maximum OKR gain is consistent with high photoreceptor densities in the
retina. Contour lines show retinal photoreceptor density determined by optical
measurements of explanted eye cups of 7-8 dpf zebrafish larvae, at increments of 10% of
maximum density. Data shown in visual space coordinates relative to the body axis, i.e.,
90° azimuth and 0° elevation corresponds to a perfectly lateral direction. To highlight
densely covered regions, densities from half-maximum to maximum are additionally
shown in shades of colour. Solid circles indicate the location of maximum OKR gain
inferred from experiments of type D in 5-7dpf larvae (**Fig 4**). White outlines indicate the
area that would be covered by a 40° disk-shaped stimulus centred on this location when
the eye is in its resting position. As the eyes move within their beating field during OKR,
the actual, non-stationary retinal coverage extends further rostrally and caudally. For (a)
red, green, and blue photoreceptors, high densities coincide with high OKR gains. (b) For
ultraviolet receptors, there is no clear relationship to the OKR gain. (c) For reference, the
summed total density of all receptor types combined. We did not observe a significant
shift in the position-dependence of maximum OKR gain between groups of larvae at 5 dpf,
6 dpf or 7 dpf of age, consistent with the notion that retinal development is far advanced
and the circuits governing OKR behaviour are stable at this developmental stage.

   Tables

865   **Table 1. Stimulus parameters (position dependence).** These stimuli consisted of a
866   horizontally moving grating, cropped with a disk-shaped stimulus mask, and presented
867   in one of 38 different locations across the visual field. Parameters as in **Table 1**. Results
868   shown in **Fig 4**.
869

type	α	β	δ	SF	TF	v	T
D1	-16	25	64	0.0611	0.7639	12.5	10
D2	-35	53	64	0.0611	0.7639	12.5	10
D3	-90	74	64	0.0611	0.7639	12.5	10
D4	-145	53	64	0.0611	0.7639	12.5	10
D5	-164	25	64	0.0611	0.7639	12.5	10
D6	-60	19	64	0.0611	0.7639	12.5	10
D7	-90	39	64	0.0611	0.7639	12.5	10
D8	-120	19	64	0.0611	0.7639	12.5	10
D9	-30	0	64	0.0611	0.7639	12.5	10
D10	-90	0	64	0.0611	0.7639	12.5	10
D11	-150	0	64	0.0611	0.7639	12.5	10
D12	-60	-19	64	0.0611	0.7639	12.5	10
D13	-90	-39	64	0.0611	0.7639	12.5	10
D14	-120	-19	64	0.0611	0.7639	12.5	10
D15	-16	-25	64	0.0611	0.7639	12.5	10
D16	-35	-53	64	0.0611	0.7639	12.5	10
D17	-90	-74	64	0.0611	0.7639	12.5	10
D18	-145	-53	64	0.0611	0.7639	12.5	10
D19	-164	-25	64	0.0611	0.7639	12.5	10
D20-D38	same as D1-D19, but with positive azimuth α (right hemisphere of arena)						

Table 2. Stimulus parameters (frequency dependence). These stimuli consisted of a horizontally moving grating, cropped with a disk-shaped stimulus mask. At each location, 7 different spatial frequencies and thus velocities were used, while temporal frequency was held constant. Parameters and units as in Table 1. Results shown in **Fig 5**.

type	α	β	δ	SF	TF	v	T
F1	-70	15	40	0.0181	0.7639	42.188	10
F2	-70	15	40	0.0272	0.7639	28.125	10
F3	-70	15	40	0.0407	0.7639	18.750	10
F4	-70	15	40	0.0611	0.7639	12.500	10
F5	-70	15	40	0.0917	0.7639	8.3333	10
F6	-70	15	40	0.1375	0.7639	5.5556	10
F7	-70	15	40	0.2063	0.7639	3.7037	10
F8-F14	same as V1-V7, but with azimuth $\alpha = -110$ and elevation $\beta = -15$						
F15-F21	same as V1-V7, but with azimuth $\alpha = -28$ and elevation $\beta = 15$						
F22-F42	same as A1-A21, but with positive azimuth α (right hemisphere)						

Table 3. Stimulus parameters (size dependence). These stimuli consisted of a horizontally moving grating, cropped with a disk-shaped stimulus mask. At each location, disks with 7 different, logarithmically spaced areas were shown. Parameters and units as in **Table 1**. Results shown in **Fig 5**.

type	α	β	δ	SF	TF	v	T
A1	-70	15	3.58	0.0611	0.7639	12.5	10
A2	-70	15	7.16	0.0611	0.7639	12.5	10
A3	-70	15	14.3	0.0611	0.7639	12.5	10
A4	-70	15	28.7	0.0611	0.7639	12.5	10
A5	-70	15	57.9	0.0611	0.7639	12.5	10
A6	-70	15	120	0.0611	0.7639	12.5	10
A7	-70	15	360	0.0611	0.7639	12.5	10
A8-A14	same as A1-A7, but with azimuth $\alpha = -110$ and elevation $\beta = -15$						
A15-A21	same as A1-A7, but with azimuth $\alpha = -28$ and elevation $\beta = 15$						
A22-A42	same as A1-A21, but with positive azimuth α (right hemisphere)						

Spherical arena reveals optokinetic response tuning to stimulus location, size and frequency across entire visual field of larval zebrafish

Supplementary Figures

S1 Figure. Control experiments. (a) Artificial triangular holes setup. (b) Artificial keel setup. (c) Neither triangular nor elongated gaps result in significantly different OKR gains. (d) Regular arena setup. (e) Arena can be tilted 180 degrees so front and rear, left and right, upper and lower LED positions are swapped. The bulb holder moves accordingly. (f) Upside-down embedding setup. (g) Setup with inverted optical path, including illumination. (h-k) Results in body-centred coordinates, where positive elevations refer to dorsal positions, for the four setups shown in (d-g). (i,j) Experiments with presentation of a less regularly distributed set of stimuli, cropped to disks of 64 degrees polar angle instead of the 40 degrees used in (d,e,h,i). (j) Fish embedded upside-down exhibit a slight preference for stimuli below the body-centred equator, i.e., positions slightly ventral to their body axis. (k) Fish embedded upright, as in (d); this preliminary dataset was acquired during methods development and is more variable than the others. To account for environmental asymmetries such as arena anisotropies, we combined the data underlying (h) and (i) to obtain **Fig 4c-e** (see **Methods**). Data from (h) n=7, (i) n=5, (j) n=3, (k) n=10 fish.

S2 Figure. During OKR, the beating field and average eye position are independent of stimulus location. All data were pooled across fish and trials. One gain value was computed per stimulus presentation. Violin plots show distribution of mean horizontal eye positions across the pooled data; vertical lines indicate 25th percentile, median and 75th percentile of the distribution. Positions are those during presentation of stimulus types (a) D1 to D38 shown in **Fig 4a**, (b) A1 to A42 shown in **Fig 5b**, (c) F1 to F42 shown in **Fig 5a.** Dashed lines in (c) represent axis limits of (a,b).

S3 Figure. Vertical eye position under upright and upside-down embedding. Larvae were embedded in agarose with their eyes free, and placed under a microscope. Using an additional mirror, we recorded image time series from the front, i.e., looking along the anterior-posterior axis. Vertical eye position was determined geometrically for each individual frame. Because some larvae were embedded in such a way that mediolateral axis was not entirely aligned with the true horizon of the environment, we measured vertical eye position (a) relative to both the mediolateral body axis, and (b) the mediolateral body axis, but corrected by its offset from the true horizon. Required corrections were minimal. More importantly, there were no significant differences between the vertical eye positions of larvae embedded upside-up (uu) or upside-down (ud), neither for their left (L) or right eyes (R). Bars indicate mean after pooling across all frames and individuals, error bars show corresponding standard error of the mean (s.e.m.). On all panels, positive angles indicate dorsalward eye positions. In summary, larval eyes were almost always inclined towards the dorsum, irrespective of the direction of embedding. The fish do not appear to compensate for their orientation with respect to the gravitational axis.

S4 Figure. Yoking indices are biased by reflections within the arena. Yoking indices were computed for experiments using the regular setup as in **S1d Fig** (light grey), with a rotated arena as in **S1e Fig** (dark grey), corrected for experimental asymmetries as in Fig4 (green), and with one side of the glass bulb, contralateral to the stimulus centre, painted black (black). Yoking indices from most experiments are close to zero, indicating similar OKR gains for both eyes regardless of stimulus location. In contrast, yoking indices

from the latter control experiment differ markedly from zero, indicating significantly weaker responses by the respectively unstimulated eye. This finding points to reflections at the air-glass interface being visible to the purportedly "unstimulated" eye.

S5 Figure. Across different types of arenas, stimulus reflections affect perceived yoking. Control experiments conducted in a rectangular stimulus arena. OKR-inducing gratings were shown on all four stimulus screens surrounding the larvae, while additional elements were introduced around either the left eye (LE), the right eye (RE) or neither. Specifically, selected eyes were (a,b) shown stationary stimuli of the same frequency and contrast as the moving stimuli, (c,d) shown a blank white surface, or (e,f) shielded with a fully opaque cover. (a,c,e,g) Bars indicate mean OKR gains, and error bars show standard error of the mean. (b,d,f,h) Yoking indices are near zero when both eyes move with identical amplitudes and positive when left eye amplitude exceeds that of the right eye (see **Methods**). (a,b) In the presence of two conflicting stimuli (moving vs. stationary), yoking between the eyes is reduced by almost half, confirming that the unstimulated eye is yoked to the stimulated eye, albeit with a lower OKR gain. (c,d) When there is no conflicting stimulus, yoking drives OKR of the contralateral eyes, albeit with a lower amplitude as if both eyes were stimulated directly with identical stimulus, (e,f) which is equally true in the presence of shielding. (g,h) To assess the effect of reflections on the difference between directly stimulated and purportedly stimulated eyes, we compare blank stimuli (as in c-d, which could diffusely reflect light) to fully shielded eyes (as in e-f, where no reflections should occur). There are no significant differences, indicating that the larger effect of reflections observed in our spherical arena (**S4 Fig**) may be caused specifically by sharp reflections of the stimulus patterns at the air-water interface, instead of more diffuse reflections of light across the background. *Two rectangular arena setups were used for the control experiments. Asterisks indicate data obtained from the second setup, for which balanced illumination was explicitly confirmed via diode photodetector. Data from n=22 fish for initial setup and n=10 fish for second setup.

S6 Figure. Asymmetries between left and right eye are strongly affected by the environment. Differences between the OKR gains of directly stimulated left eyes and directly stimulated right eyes can be explained by a linear combination of biological and environmental factors b_k, e.g., biases of individual animals or asymmetries of the stimulus arena (see **Methods**). Comparing data from the regular and rotated setups (**S1d-e Fig**), as well as data from fish immobilised upside-down (**S1f Fig**), including additional stimuli covering entire hemispheres (**S4 Table**), we can infer the underlying b_k via multivariate regression of our linear model. We find that individual biases (grey) vary strongly from fish to fish, and are broadly distributed from left to right. There are some constant biases across fish (green), both towards the left side of their visual field (b_2) and towards one of the two LED hemispheres (b_1); however, these biases are small and, given the large variability of individual biases, might be a result of the limited number of fish studied.

S7 Figure. Maximum OKR gain compared to photoreceptor densities in retinal coordinates. Same as Figure 6, but showing positions across the retina in Cartesian coordinates, as originally published (Zimmermann et al. 2018 Curr Biol), instead of in geographic visual field coordinates. When plotting photoreceptor densities in Cartesian coordinates, the regions of highest densities appear to be located quite peripheral/eccentric. However, the plot of the densities in visual field coordinates (Figure 6) confirms the coincidence of high densities and high OKR gains. Solid circles indicate the location of maximum OKR gain inferred from experiments of type D in 5-7dpf larvae (**Fig 4**), and corrected by the mean eye position over time.

Supplementary Tables

S1 Table. Arena cross-section. Elevation of LED tile centres, from top ribbon (ribbon
number +5, near the north pole of the sphere) to bottom ribbon (ribbon number -5, near
the South Pole). See **Fig 2c** for a graphical illustration. *As the left and right hemispheres
of the arena are mirror-symmetric to one another, each ribbon contains the same number
of tiles within each of the two hemispheres. **Because the structural scaffold is reinforced
near the bottom of the sphere, ribbons -4 contains one fewer LED tile than ribbon +4.
Other than that, the arrangement of LED tiles is almost mirror-symmetric from top to
bottom as well.

ribbon no. k	tiles per ribbon	elevation β_k of tile centre (degrees)
+5	2* x 5	62.4
+4	2 x 9**	49.9
+3	2 x 11	37.6
+2	2 x 13	25.0
+1	2 x 14	12.5
0 (equator)	2 x 15	0
-1	2 x 14	-12.5
-2	2 x 13	-25.0
-3	2 x 11	-37.6
-4	2 x 8**	-49.9
-5	2 x 5	-62.4

S2 Table. Absolute positions of fish and setup elements in different experiments. All
positions and directions are given in environmental coordinates, i.e. the approximate
cardinal directions (North etc.) of the laboratory, as well as Up and Down (away from or
towards the core of the Earth). Most experiments, including control experiments, are of
type 1. The only exceptions are the position-dependence experiments with rotated arena
(type 2), control experiments with upside-down embedding (type 3), and control
experiments with inverted IR illumination (type 4). Three LED tiles in different parts of
the visual field are included as examples. Positions are given in lab-centred geographic
coordinates as (a, e), where a is the azimuth, e the elevation, and 0° azimuth is arbitrarily
chosen to point South.

experiments	type 1	type 2	type 3	type 4
infrared illumination at	Up (0°,+90°)	Up (0°,+90°)	Up (0°,+90°)	Down (0°,-90°)
camera at	Down (0°,-90°)	Down (0°,-90°)	Down (0°,-90°)	Up (0°,+90°)
flat face of glass triangle facing	Up (0°,+90°)	Up (0°,+90°)	Up (0°,+90°)	Up (0°,+90°)
dorsum facing	Up (0°,+90°)	Up (0°,+90°)	Down (0°,-90°)	Up (0°,+90°)
rostrum facing	South (0°,0°)	North (180°,0°)	South (0°,0°)	South (0°,0°)
arena axis A	South (0°,0°)	North (180°,0°)	South (0°,0°)	South (0°,0°)
arena axis B	West (90°,0°)	West (90°,0°)	West (90°,0°)	West (90°,0°)
arena axis C	Up (0°,+90°)	Down (0°,-90°)	Up (0°,+90°)	Up (0°,+90°)
LED tile #1	Up, West, South (+35°,+53°)	Down, West, North (+145°,-53°)	Up, West, South (+35°,+53°)	Up, West, South (+35°,+53°)
LED tile #2	East (-90°,0°)	East (-90°,0°)	East (-90°,0°)	East (-90°,0°)
LED tile #3	Down, East, South (-16°,-25°)	Up, East, North (+164°,+25°)	Down, East, South (-16°,-25°)	Down, East, South (-16°,-25°)

997
998

S3 Table. Stimulus parameters (control experiments). These stimuli consisted of a horizontally moving grating displayed on four flat, rectangular stimulus screens surrounding the larva. One pair of screens displayed stimuli visible to the left eye only, and the other pair displayed stimuli to the right eye only. Results shown in **S5 Fig**.

type	left eye stimulus	right eye stimulus	shown on panel
C0	moving pattern	moving pattern	**S5a-c Fig**
C1	stationary pattern	moving pattern	**S5a-b Fig**
C2	moving pattern	stationary pattern	**S5a-b Fig**
C3	blank surface	moving pattern	**S5c-d Fig, S5g-h Fig**
C4	moving pattern	blank surface	**S5c-d Fig, S5g-h Fig**
C5	eye shielded	moving pattern	**S5e-f Fig, S5g-h Fig**
C6	moving pattern	eye shielded	**S5e-f Fig, S5g-h Fig**

S4 Table. Stimulus parameters (whole-field and hemispheres). These stimuli consisted of a horizontally moving grating, either covering the entire visual field or cropped to one of the 6 principal hemispheres (front, rear, upper, lower, left, right). The stimulus mask is determined by the azimuth α (degrees) and elevation β (degrees) of its centre, as well as its size, given by the angle δ (degrees) it spans. The moving grating is characterised by its spatial frequency SF (cycles/degree), temporal frequency TF (cycles/sec), peak velocity v (deg/sec), and oscillation period T (sec).

type	α	β	δ	SF	TF	v	T
H0	0	0	360	0.0611	0.7639	12.5	10
H1	0	0	180	0.0611	0.7639	12.5	10
H2	180	0	180	0.0611	0.7639	12.5	10
H3	0	90	180	0.0611	0.7639	12.5	10
H4	0	-90	180	0.0611	0.7639	12.5	10
H5	90	0	180	0.0611	0.7639	12.5	10
H6	-90	0	180	0.0611	0.7639	12.5	10

References

1. Cronin TJ, S.; Marshal, NJ.; Warrant, EJ. Visual Ecology: Princeton University Press; 2014.
2. Hoy JL, Yavorska I, Wehr M, Niell CM. Vision Drives Accurate Approach Behavior during Prey Capture in Laboratory Mice. Curr Biol. 2016;26(22):3046-52.
3. Bianco IH, Kampff AR, Engert F. Prey capture behavior evoked by simple visual stimuli in larval zebrafish. Front Syst Neurosci. 2011;5:101.
4. Yoshimatsu T, Schröder C, Berens P, Baden T. Cellular and molecular mechanisms of photoreceptor tuning for prey capture in larval zebrafish. BioRxiv. 2019.

1023   5.   Smolka J, Zeil J, Hemmi JM. Natural visual cues eliciting predator avoidance in
1024   fiddler crabs. Proc Biol Sci. 2011;278(1724):3584-92.
1025   6.   Zhang Y, Kim IJ, Sanes JR, Meister M. The most numerous ganglion cell type of
1026   the mouse retina is a selective feature detector. Proc Natl Acad Sci U S A.
1027   2012;109(36):E2391-8.
1028   7.   Haug MF, Biehlmaier O, Mueller KP, Neuhauss SC. Visual acuity in larval
1029   zebrafish: behavior and histology. Front Zool. 2010;7:8.
1030   8.   Merigan WH, Katz LM. Spatial resolution across the macaque retina. Vision Res.
1031   1990;30(7):985-91.
1032   9.   Murasugi CM, Howard IP. Human horizontal optokinetic nystagmus elicited by
1033   the upper versus the lower visual fields. Vis Neurosci. 1989;2(1):73-9.
1034   10.   Shimizu N, Tabata H, Wada Y, Sugita Y, Yamanaka T, Hosoi H, et al. Distribution
1035   of optokinetic sensitivity across the retina of mice in relation to eye orientation.
1036   Neuroscience. 2010;168(1):200-8.
1037   11.   Yang X, Guo A. Distinct acute zones for visual stimuli in different visual tasks in
1038   Drosophila. PLoS One. 2013;8(4):e61313.
1039   12.   Zimmermann MJY, Nevala NE, Yoshimatsu T, Osorio D, Nilsson D-E, Berens P, et
1040   al. Zebrafish Differentially Process Color across Visual Space to Match Natural
1041   Scenes. Current Biology. 2018;28:2018-32.e5.
1042   13.   Baden T, Schubert T, Chang L, Wei T, Zaichuk M, Wissinger B, et al. A tale of two
1043   retinal domains: near-optimal sampling of achromatic contrasts in natural
1044   scenes through asymmetric photoreceptor distribution. Neuron.
1045   2013;80(5):1206-17.
1046   14.   Baier H, Scott EK. Genetic and optical targeting of neural circuits and behavior--
1047   zebrafish in the spotlight. Curr Opin Neurobiol. 2009;19(5):553-60.
1048   15.   McLean DL, Fetcho JR. Movement, technology and discovery in the zebrafish.
1049   Curr Opin Neurobiol. 2011;21(1):110-5.
1050   16.   Easter J, Stephen S., Nicola GN. The Development of Vision in the Zebrafish
1051   (Danio rerio). Developmental Biology. 1996;180:646-63.
1052   17.   Nawrocki L, Bremiller R, Streisinger G, Kaplan M. Larval and adult visual
1053   pigments of the zebrafish, Brachydanio rerio. Vision Research. 1985;25:1569-76.
1054   18.   Trivedi CA, Bollmann JH. Visually driven chaining of elementary swim patterns
1055   into a goal-directed motor sequence: a virtual reality study of zebrafish prey
1056   capture. Front Neural Circuits. 2013;7:86.
1057   19.   Heap LAL, Vanwalleghem G, Thompson AW, Favre-Bulle IA, Scott EK. Luminance
1058   Changes Drive Directional Startle through a Thalamic Pathway. Neuron.
1059   2018;99(2):293-301 e4.
1060   20.   Kubo F, Hablitzel B, Dal Maschio M, Driever W, Baier H, Arrenberg AB.
1061   Functional architecture of an optic flow-responsive area that drives horizontal
1062   eye movements in zebrafish. Neuron. 2014;81(6):1344-59.
1063   21.   Orger MB, Kampff AR, Severi KE, Bollmann JH, Engert F. Control of visually
1064   guided behavior by distinct populations of spinal projection neurons. Nat
1065   Neurosci. 2008;11(3):327-33.
1066   22.   Howard IP, Ohmi M. The efficiency of the central and peripheral retina in driving
1067   human optokinetic nystagmus. Vision Res. 1984;24(9):969-76.
1068   23.   Hafed ZM, Chen CY. Sharper, Stronger, Faster Upper Visual Field Representation
1069   in Primate Superior Colliculus. Curr Biol. 2016;26(13):1647-58.
1070   24.   Brockerhoff SE, Hurley JB, Janssen-Bienhold U, Neuhauss SC, Driever W, Dowling
1071   JE. A behavioral screen for isolating zebrafish mutants with visual system
1072   defects. Proceedings of the National Academy of Sciences. 1995;92:10545-9.
1073   25.   Muto A, Orger MB, Wehman AM, Smear MC, Kay JN, Page-McCaw PS, et al.
1074   Forward genetic analysis of visual behavior in zebrafish. PLoS Genet.
1075   2005;1(5):e66.

26. Neuhauss SC, Biehlmaier O, Seeliger MW, Das T, Kohler K, Harris WA, et al.
Genetic disorders of vision revealed by a behavioral screen of 400 essential loci
in zebrafish. J Neurosci. 1999;19(19):8603-15.
27. Clark D. Visual responses in the developing zebrafish (Brachydanio rerio). 1981.
28. Cohen B, Matsuo V, Raphan T. Quantitative analysis of the velocity
characteristics of optokinetic nystagmus and optokinetic after-nystagmus. The
Journal of Physiology. 1977;270:321-44.
29. Huang Y-Y, Neuhauss SCF. The optokinetic response in zebrafish and its
applications. Frontiers in bioscience : a journal and virtual library.
2008;13:1899-916.
30. Rinner O, Rick JM, Neuhauss SCF. Contrast Sensitivity, Spatial and Temporal
Tuning of the Larval Zebrafish Optokinetic Response. Investigative
Opthalmology & Visual Science. 2005;46:137.
31. Portugues R, Feierstein CE, Engert F, Orger MB. Whole-brain activity maps
reveal stereotyped, distributed networks for visuomotor behavior. Neuron.
2014;81(6):1328-43.
32. Naumann EA, Fitzgerald JE, Dunn TW, Rihel J, Sompolinsky H, Engert F. From
Whole-Brain Data to Functional Circuit Models: The Zebrafish Optomotor
Response. Cell. 2016;167(4):947-60 e20.
33. Wang K, Hinz J, Haikala V, Reiff DF, Arrenberg AB. Selective processing of all
rotational and translational optic flow directions in the zebrafish pretectum and
tectum. BMC Biol. 2019;17(1):29.
34. Lindsey BW, Smith FM, Croll RP. From inflation to flotation: contribution of the
swimbladder to whole-body density and swimming depth during development
of the zebrafish (Danio rerio). Zebrafish. 2010;7(1):85-96.
35. Kim SS, Rouault H, Druckmann S, Jayaraman V. Ring attractor dynamics in the
Drosophila central brain. Science (New York, NY). 2017;356:849-53.
36. Maisak MS, Haag J, Ammer G, Serbe E, Meier M, Leonhardt A, et al. A directional
tuning map of Drosophila elementary motion detectors. Nature. 2013;500:212-6.
37. Reiser MB, Dickinson MH. A modular display system for insect behavioral
neuroscience. J Neurosci Methods. 2008;167(2):127-39.
38. Dehmelt FA, von Daranyi A, Leyden C, Arrenberg AB. Evoking and tracking
zebrafish eye movement in multiple larvae with ZebEyeTrack. Nat Protoc.
2018;13(7):1539-68.
39. Tappeiner C, Gerber S, Enzmann V, Balmer J, Jazwinska A, Tschopp M. Visual
acuity and contrast sensitivity of adult zebrafish. Front Zool. 2012;9(1):10.
40. Beck JC, Gilland E, Tank DW, Baker R. Quantifying the ontogeny of optokinetic
and vestibuloocular behaviors in zebrafish, medaka, and goldfish. J
Neurophysiol. 2004;92(6):3546-61.
41. Schaerer S, Kirschfeld K. The role of background movement in goldfish vision. J
Comp Physiol A. 2000;186(6):583-93.
42. Shcherbakov D, Knorzer A, Espenhahn S, Hilbig R, Haas U, Blum M. Sensitivity
differences in fish offer near-infrared vision as an adaptable evolutionary trait.
PLoS One. 2013;8(5):e64429.
43. Sovrano VA, Andrew RJ. Eye use during viewing a reflection: Behavioural
lateralisation in zebrafish larvae. Behavioural Brain Research. 2006;167:226-31.
44. Watkins J, Miklósi A, Andrew RJ. Early asymmetries in the behaviour of zebrafish
larvae. Behavioural Brain Research. 2004;151:177-83.
45. Andrew RJ, Dharmaretnam M, Gyori B, Miklosi A, Watkins JA, Sovrano VA.
Precise endogenous control of involvement of right and left visual structures in
assessment by zebrafish. Behav Brain Res. 2009;196(1):99-105.
46. Miklosi A, Andrew RJ. Right eye use associated with decision to bite in zebrafish.
Behav Brain Res. 1999;105(2):199-205.

47.   Avanesov A, Malicki J. Analysis of the Retina in the Zebrafish Model. Methods in
Cell Biology. 2010;100:153-204.
48.   Lafortune SH, Ireland DJ, Jell RM. Suppression of optokinetic velocity storage in
humans by static tilt in roll. J Vestib Res. 1990;1(4):347-55.
49.   Pettorossi VE, Ferraresi A, Botti FM, Panichi R, Barmack NH. Head position
modulates optokinetic nystagmus. Exp Brain Res. 2011;213(1):141-52.
50.   Zolotilina EG, Eremina SV, Orlov IV. Horizontal optokinetic nystagmus in the
pigeon during static tilts in the frontal plane. Neurosci Behav Physiol.
1995;25(4):300-6.
51.   Arunachalam M, Raja M, Vijayakumar C, Malaiammal P, Mayden RL. Natural
history of zebrafish (Danio rerio) in India. Zebrafish. 2013;10(1):1-14.
52.   Engeszer RE, Patterson LB, Rao AA, Parichy DM. Zebrafish in the wild: a review
of natural history and new notes from the field. Zebrafish. 2007;4(1):21-40.
53.   Parichy DM. Advancing biology through a deeper understanding of zebrafish
ecology and evolution. Elife. 2015;4.
54.   Spence R, Gerlach G, Lawrence C, Smith C. The behaviour and ecology of the
zebrafish, Danio rerio. Biol Rev Camb Philos Soc. 2008;83(1):13-34.
55.   Wang K, Hinz J, Zhang Y, Thiele T, Arrenberg AB. Parallel channels for motion
feature extraction in the pretectum and tectum of larval zebrafish. BioRxiv.
preprint 2019.
56.   Joesch M, Plett J, Borst A, Reiff DF. Response properties of motion-sensitive
visual interneurons in the lobula plate of Drosophila melanogaster. Curr Biol.
2008;18(5):368-74.
57.   Suver MP, Huda A, Iwasaki N, Safarik S, Dickinson MH. An Array of Descending
Visual Interneurons Encoding Self-Motion in Drosophila. J Neurosci.
2016;36(46):11768-80.
58.   Arrenberg AB. Fiber Optic-Based Photostimulation of Larval Zebrafish. Methods
Mol Biol. 2016;1451:343-54.

1157

1159 **Fig 1:**

1160

1161   **Fig 2:**

1162

1163 **Fig 3:**

1164

a sinusoidally moving stimulus pattern

b OKR gain is the ratio of amplitudes

c saccades interrupt slow-phase eye movement

d eye movement amplitude is approximated by piece-wise fit to slow phase

1165

1166   **Fig 4:**
1167

1168

1169   **Fig 5:**

1170

1171
1172
1173
1174   **Fig 6:**
1175

1176

S1 Fig

S2 Fig

1185   **S3 Fig**

1186
1187   **S4 Fig**
1188

1189

S5 Fig

1194   **S6 Fig**

1195

1196
1197
1198   **S7 Fig**

1199

a optokinetic response (OKR)

stim.

10° 10 sec

eyes

b experimental field of view

stimulus screen

natural field of view

out of view

c LED arena

d individual LED positions

elevation

-90

-180 -90 azimuth 90 180

e z axis

+90°

elevation

±180°

-90°

+90°

0° 0°

azimuth

-90°

f 20(.787) mm

14,484 LEDs

8 x 8 LEDs per tile

232 LED tiles

2.48(.098) mm

g keel

ribs

3D-printed structural scaffold

h

a

upper gap

z axis
(β = ±90)

β₁

-β₁

LED tiles

β₅

equator
(β = 0)

sphere
centre

structural
ribs

illumination

lower gap

b

infrared light source

milk glass diffusers

ND filter

lens

embedded larva

mirror

lens

infrared
camera

c

placement of embedded larva

liquid-filled
glass bulb

bulb holder

d

bulb with embedded larva

infrared light source

ND filter

lens

movable
hemisphere

LED cables

infrared
camera

a sinusoidally moving stimulus pattern

b OKR gain is the ratio of amplitudes

stimulus amplitude

eye movement amplitude

5°

3 sec

c saccades interrupt slow-phase eye movement

saccades

slow phase

slow phase

slow phase

d eye movement amplitude is approximated by piece-wise fit to slow phase

amplitude of best fit

saccades cropped before fitting

a stimulus crop

left hemisphere   right hemisphere

stimulus

b left eye gain g_L

c merge stimulated

d right eye gain g_R

OKR gain

0.2

0.1

e

left eye gain

merge stimulated

right eye gain

elevation (deg)

−30

−60

left − azimuth (deg) − right

−180   −90   0   90   180

a

red cones (mm^{-2})   green cones (mm^{-2})   blue cones (mm^{-2})

b

ultraviolet cones

c

all cones combined