Apical length governs computational diversity of L5 pyramidal

neurons

Alessandro R. Galloni[1,2], Aeron Laffere[1,3] and Ede Rancz[1*]

[1] The Francis Crick Institute, London, UK

[2] University College London, UK

[3] Birkbeck College, University of London, UK

*Correspondence to ede.rancz@crick.ac.uk

Abstract

Anatomical similarity across the neocortex has led to the common assumption that the

circuitry is modular and performs stereotyped computations. Layer 5 pyramidal neurons

(L5PNs) in particular are thought to be central to cortical computation because of their

extensive arborisation and nonlinear dendritic operations. Here, we demonstrate that

computations associated with dendritic Ca^{2+} plateaus in L5PNs vary substantially between

the primary and secondary visual cortices. L5PNs in the secondary visual cortex show

reduced dendritic excitability and smaller propensity for burst firing. This reduced excitability

is correlated with shorter apical dendrites. Using numerical modelling, we uncover a

universal principle underlying the influence of apical length on dendritic backpropagation and

excitability, based on a Na^+ channel-dependent broadening of backpropagating action

potentials. In summary, we provide new insights into the modulation of dendritic excitability

by apical dendrite length and show that the operational repertoire of L5 neurons is not

universal throughout the brain.

Introduction

The neocortex is thought to have a modular structure composed of 'canonical circuits' performing stereotyped computations (Harris & Shepherd, 2015; Markram et al., 2015; Miller, 2016). Anatomical evidence supports the existence of repeating circuit architectures that display similar general features across species and brain areas (Carlo & Stevens, 2013; Douglas & Martin, 2004; Mountcastle, 1997). It is generally thought that these architectural motifs serve as a physical substrate to perform a small range of specific, canonical computations (Bastos et al., 2012; Braganza & Beck, 2018).

Pyramidal neurons are the main building blocks of these circuit motifs. Across brain areas and species, their biophysical attributes endow them with non-linear properties that allow them to implement a repertoire of advanced computations at the single cell level (Gidon et al., 2020; London & Häusser, 2005; Spruston, 2008). Layer 5 pyramidal neurons (L5 PNs) in particular provide a striking example of how dendritic properties can underlie circuit-level computations in a laminar circuit. Their dendritic supralinearities enable signal amplification and coincidence detection of inputs — a crucial operation to integrate feedforward and feedback streams that often send projections onto separate dendritic domains. In these cells, a single backpropagating action potential (bAP), when combined with distal synaptic input, can trigger a burst of somatic action potentials. The crucial mechanism underlying this supralinear phenomenon is the all-or-none dendritic Ca^{2+} plateau (M. E. Larkum, Kaiser, & Sakmann, 1999; Matthew E. Larkum, Zhu, & Sakmann, 1999).

Morphology and intrinsic properties have a profound influence on neuronal excitability. Dendritic topology and the electrical coupling between the soma and dendrites is thought to be particularly crucial for determining a neuron's integrative properties (Mainen & Sejnowski, 1996; Schaefer, Larkum, Sakmann, & Roth, 2003; van Ooyen, Duijnhouwer, Remme, & van Pelt, 2002; Vetter, Roth, & Hausser, 2001). Recent experimental work has shown that there can be substantial variation in intrinsic properties of L5 neurons depending on the location within a cortical area or on the species they are recorded from (Beaulieu-Laroche et al., 2018; Fletcher & Williams, 2019). However, it is often assumed that pyramidal neurons have robust enough properties across cortical areas and brain structures to support similar computations (Bastos et al., 2012; Hawkins, Ahmad, & Cui, 2017; M. Larkum, 2013; Shipp, 2016). For instance, analogous to L5 PNs, hippocampal pyramidal neurons also display dendritic Ca^{2+} APs that support coincidence detection of distal and proximal inputs (Jarsky, Roxin, Kath, & Spruston, 2005).

If L5 pyramidal neurons indeed have a common repertoire of operations in support of canonical computations, one would expect the same cell type in adjacent and closely related areas to exhibit the same computational repertoire. Here we have studied the bursting properties of thick-tufted L5 neurons (ttL5) in mouse primary and secondary visual cortices

(V1 and V2m). Through systematic and rigorously standardized experiments, we found

fundamentally different operation patterns linked to morphology in the two brain areas.

Through computational modelling, we reveal new insights into biophysical mechanism

linking excitability to morphology, which is able to account for this difference. Our results

question the notion of a common operational repertoire in pyramidal neurons and thus

cortical canonical computations as well.

Results

Thick-tufted L5 neurons in V2m lack BAC firing

We made whole-cell patch clamp recordings from ttL5 pyramidal neurons in V1 and V2m in

acutely prepared mouse brain slices. To ensure consistency in cell type, recordings were

restricted either to neurons projecting to the lateral posterior nucleus of thalamus, identified

using retrograde labelling with cholera toxin subunit B (**Supplementary Figure 1**), or to

neurons labelled in the Glt25d2-Cre mouse line (Groh et al., 2010). In addition, we were able

to confirm the characteristic morphological features of ttL5 neurons in a subset of the

recorded neurons using biocytin reconstructions. We were thus able to maintain cortical area

as the primary variant when comparing V1 and V2m neurons.

To reproduce the conditions required for triggering BAC firing, we stimulated synaptic inputs

near the distal tuft in L1 using an extracellular electrode in conjunction with somatic

stimulation through the recording electrode (**Fig. 1a**). To avoid recruiting inhibitory inputs

during the extracellular stimulation and create the most favourable conditions to enable BAC

firing (Perez-Garci, Gassmann, Bettler, & Larkum, 2006), we added the competitive GABA$_B$

receptor antagonist CGP52432 (1 µM) to the extracellular solution. Extracellular current

pulses in L1 were adjusted to evoke either a subthreshold EPSP or a single action potential

at the soma. Somatic injection of a 5 ms depolarizing current pulse through the recording

electrode was used to trigger single APs. In V1 neurons, combined stimulation (with the L1

input triggered at the end of the somatic pulse) could evoke a prolonged plateau potential

resulting in a burst of 3 APs. We repeated these experiments in ttL5 pyramidal neurons

located in V2m under the same recording conditions. Upon coincident somatic AP and

extracellular L1 stimulation, BAC firing was almost never observed in V2m, suggesting a

much-reduced dendritic excitability in V2m neurons. For the purposes of these experiments,

we defined as "supralinear" any cell in which three or more APs could be evoked following

combined somatic and L1 stimulation (each evoking no more than one AP individually).

Supralinearity was observed in half the recorded V1 neurons (10/20), while neurons in V2m

showed an almost total lack of supralinearity (1/19, **Fig. 1b**).

Figure 1. V2m neurons are less prone to burst than in V1.

a. *Left:* diagram of experimental configuration. *Right:* example traces during BAC firing protocol,

recorded from V1 (red) and V2m (blue) ttL5 neurons. **b.** Proportion of supralinear cells in V1 and V2m

c. *Left:* diagram of experimental configuration. *Right:* example traces of V1 and V2m ttL5 neurons

stimulated with 50 Hz and 120 Hz AP trains. Note the sustained after-depolarization following the 120

Hz spike train in the V1 neuron. *Inset:* ADP measured as the area between the 50 Hz trace and the

higher frequency trace following the last spike. Inset scale bar: 5 ms x 5 mV. **d.** Quantification of ADP

area at each measured frequency for the example neurons in *c*. The peak integral value is highlighted

in red. **e.** Histogram of results across all recorded cells with defined critical frequency.

Thick-tufted L5 neuron in V2m lack a critical frequency ADP

To further investigate the prevalence of dendritic supralinearities in ttL5 neurons across

visual cortices, we recorded another hallmark of dendritic Ca^{2+} plateaus: a prominent

somatic ADP following a high-frequency train of somatic APs (M. E. Larkum et al., 1999;

Shai, Anastassiou, Larkum, & Koch, 2015). We recorded the somatic membrane potential

from ttL5 neurons and evoked three action potentials using 3 ms pulses of somatic current

injection at frequencies ranging from 50 Hz to 200 Hz in 10 Hz increments (**Fig. 1c**). In V1

neurons, increasing the AP frequency above a critical frequency typically resulted in a

sudden increase in the ADP (**Fig. 1c, middle**). However, when recording in V2m under the

same experimental conditions, there was usually no change in ADP, even at firing

frequencies as high as 200 Hz (**Fig. 1c, right**). To quantify this effect, we aligned the peaks

of the last AP for each frequency and measured the area of the ADP difference between the

50 Hz trace and the higher frequency traces in a 20 ms window (4–24 ms) following the last

AP (**Fig. 1c, inset**). This measure of ADP increased sharply above a critical frequency and

was often largest around the value of this frequency (**Fig. 1d**). The mean critical frequency

across all cells in both V1 and V2m was 110.8 ± 29.6 Hz (SD, n = 37, excluding cells that did

not have a critical frequency, **Fig. 1e**).

Next, we measured the maximal ADP integral value for each cell **(Fig. 2a), r**egardless of the

presence of a critical frequency. Neurons in V2m had significantly smaller ADP area (V1

mean = 91 ± 50 mV*ms, SD, n = 41; V2m mean = 42 ± 33 mV*ms, SD, n = 49; p = 4.54 * 10^{-}

^{6}, two-sample Kolmogorov-Smirnov test), reflecting that most of these cells lacked a critical

frequency altogether. The extracellular artificial cerebrospinal fluid (ACSF) contained either

1.5 or 2 mM $CaCl_2$. As there was no statistically significant difference between the two

conditions in either V1 or V2m (p > 0.05, two-sample Kolmogorov-Smirnov test;

Supplementary Figure 2), we pooled all recordings. Similarly, as there was no significant

difference in the ADP measure across V2m neurons labelled retrogradely or by the Glt25d2-

Cre line (p = 0.617, two-sample Kolmogorov-Smirnov test, **Supplementary Figure 3**), we

pooled these two populations. To obtain an unbiased count of cells showing supralinearity,

we separated the unlabelled maximum ADP values pooled from both V1 and V2m into two

groups using k-means clustering (with k = 2). N.B. that in this experiment the definition of

supralinear classification differed from the experiments of Fig. 1a. The percentage of

neurons classified as supralinear (summarized in **Fig. 2b**) was more than three times higher

in V1 than in V2m, regardless of the specific supralinearity measure.

In both the BAC firing and ADP experiments mentioned above, bursting was typically also

apparent in the spiking response to a long (500 ms) depolarization at the soma. While all

ttL5 neurons are generally characterized by a spike doublet at the beginning of the current

step, in bursting neurons there is also a critical current step above which the initial spike

burst is substantially larger, usually with 3 or 4 spikes followed by a deeper

afterhyperpolarization (**Supplementary Figure 4**).

Figure 2. Excitability correlates with apical trunk length

a. Summary data of peak ADP integral values for all recorded neurons. The dashed line indicates the division between the two groups of cells classified through k-means clustering, drawn halfway between the cell with the lowest maximum ADP in the "supralinearity" cluster and the cell with the highest value in the "no supralinearity" cluster. **b.** Proportion of supralinear cells in V1 and V2m. **c.** Length of the apical trunk (soma to main bifurcation) plotted against the corresponding maximum ADP integral values. Dashed line is a linear fit; curves at the top and right are kernel density plots of the two variables in V1 and V2m.

These results show a much-diminished dendritic excitability, and as such different integrative properties, in V2m ttL5 neurons compared to V1 under the same conditions and in the same operational ranges. Previous research has indicated the length of the apical trunk as a possible factor involved in determining the dendritic excitability of ttL5 neurons in V1 (Fletcher & Williams, 2019). To test this, we reconstructed the apical trunk of 22 V1 and 26 V2m neurons from those recorded. Apical trunk lengths were significantly shorter in V2m than in V1 (V1 mean = 409 ± 64 µm, SD, n = 22; V2m mean = 313 ± 65 µm, SD, n = 26; p = $4.26*10^{-6}$, two-sample t-test, **Fig. 2c**). Additionally, there was a correlation between maximal ADP integral values and apical trunk length across the two populations (p = $5.37*10^{-3}$; F-test). These results suggest that there may be a surprisingly counter-intuitive interaction between apical trunk length and dendritic excitability—the longer the trunk, the more excitable the neuron.

BAC firing is absent in short ttL5 models

To investigate possible mechanisms underlying the dependence of bursting on apical trunk length, we ran numerical simulations in conductance-based compartmental models of ttL5 neurons. We first probed BAC firing in a morphologically detailed model developed by Hay et al. (Hay, Hill, Schürmann, Markram, & Segev, 2011), using the model parameters (biophysical model 3) and morphology (cell #1) favoured for reproducing BAC firing. As in the original paper, BAC firing was triggered by injecting a 0.5 nA current at the apical bifurcation coupled to a somatic action potential evoked by square-pulse current injection at

the soma. Mirroring the responses seen in the subset of strongly bursting ttL5 neurons,

coincident stimulation triggered BAC firing in the detailed model (**Fig. 3a, left**). We then

applied the same model to an example V2m morphology with a shorter apical dendrite. The

Ca^{2+} channel hotspot was moved to the new apical branch point (350–450 µm from the

soma vs 685-885 µm in the long morphology). The amplitude of the dendritic current

injection (0.194 nA) was scaled so as to obtain the same depolarization amplitude at the

bifurcation in both model cells. With this morphology, coincident tuft and somatic stimulation

evoked only a single somatic spike and did not trigger a dendritic Ca^{2+} plateau (**Fig. 3a,**

right). To ensure comparability between the long and short morphology, BAC firing was

probed with both 100 µm and 200 µm Ca^{2+} channel hotspot size. We did not observe any

qualitative effect of hotspot size in either model (**Supplementary Figure 5**). To test if Ca^{2+}

plateaus were at all possible in the short neuron model, we stimulated the short neuron with

a large current injection (0.5 nA) at the dendritic electrode. While the resulting dendritic

potential was substantially larger, showing activation of calcium conductances, it resulted in

only a small depolarization at the soma. Even when combining the large current injection

with a somatic spike, no spike burst could be triggered (**Supplementary Figure 6**).

To explore the sensitivity of Ca^{2+} plateaus to dendritic Ca^{2+} channel density in the long and

short neurons, we scaled the sum Ca^{2+} conductance (g_{Ca}) between 0 and 8 times the

original values. To minimize the number of variables, when scaling the relative g_{Ca} we kept

the ratio of the two channels (low- and high-voltage activated) constant. In the long

morphology the integral of the distal dendritic voltage, acting as an indicator of the large and

sustained depolarization during a Ca^{2+} plateau, increased proportionally to g_{Ca}. In the short

morphology, however, the voltage integral stayed constant across all g_{Ca} values (**Fig. 3b**).

This indicates that, although the size of a Ca^{2+} plateau depends on g_{Ca} in long neurons, in

short neurons there is no Ca^{2+} channel activation and the magnitude of the voltage integral

therefore does not depend on g_{Ca}.

To be able to vary dendritic length across a continuous range of values, we turned to a

reduced ttL5 model based on Bahl et al. (Bahl, Stemmler, Herz, & Roth, 2012). The

simplicity of this model has the added benefit of reducing the number of variables, allowing

us to explore general principles of dendritic voltage propagation with more clarity. As with

the morphologically detailed model, the reduced model with the original published

parameters displayed BAC firing triggered by coincident tuft and somatic stimulation (**Fig.**

3c, left). Shortening the apical trunk was sufficient to eliminate this response (**Fig. 3c,**

right).

Figure 3. Shorter model neurons are less prone to burst.

a. *Left:* detailed morphology of a ttL5 pyramidal neuron from the model favoured by Hay et al. for reproducing BAC firing. *Right:* reconstructed morphology of a ttL5 neuron recorded in V2m. Injected current and recorded voltage traces are shown for the soma (black), the apical trunk (blue, 400 and 200 μm), and the main bifurcation (red, 620 and 370 μm) under three different stimulation paradigms. **b.** Integral of voltage at the branch point during coincident somatic and branch point stimulation, plotted against relative g_{Ca}. **c.** *Left:* diagram of the reduced neuron model. Apical trunk length 800 μm. Injected current and recorded voltage traces as in *a*. *Right:* Same for a version of the reduced model modified to have an apical trunk length of 200 μm. **d.** Heat map representing the normalised tuft voltage integral during combined somatic and tuft stimulation in the reduced model, plotted against the absolute density of Ca^{2+} channels in the tuft compartment and the length of the apical trunk compartment. Default $g_{Ca} \cong 0.45$ mS/cm^2.

We explored the dependence of BAC firing on apical trunk length and g_{Ca} by measuring the time-integral of tuft voltage as an indicator of Ca^{2+} plateau potentials (**Fig. 3d**). The presence of a Ca^{2+} plateau depended strongly on apical trunk length and was only sensitive to g_{Ca} above a critical length of approximately 350 μm ($\cong 0.35$ λ). Below this length, no Ca^{2+} plateaus were triggered regardless of how high g_{Ca} was set to. These experiments show that a reduced model can also reproduce our results, allowing us to explore and dissect the underlying parameters in more detail.

Active propagation enhances voltage in long dendrites

To obtain a mechanistic understanding of what causes the length dependence of bursting, we made recordings from the final segment of the apical trunk as well as the tuft using the reduced model of Fig. 3c. To recreate the experimental conditions of Fig. 1b, we triggered 3 spikes at 100 Hz through a somatic electrode. As with coincident bAP and tuft input, increasing the length of the apical trunk facilitated dendritic Ca^{2+} plateau initiation (**Fig. 4a**). Interestingly, the width and peak voltage in the tuft increased steadily with dendritic length (**Fig. 4b,c**), even in the absence of Ca^{2+} currents (g_{Ca} = 0). In the presence of voltage-gated Ca^{2+} channels, the increased amplitude of bAPs triggered a large all-or-none Ca^{2+} plateau above a certain threshold length.

Figure 4. Tuft voltage increases with trunk length due to the widening of bAPs.
a. Schematic of the simulation: stimulation site in the somatic compartment, and recording sites at the distal end of the apical trunk (blue) and in the centre of the tuft (red). Stimulus shown in black. Solid lines: g_{Ca} = 0; dashed lines: original g_{Ca}. **b.** Peak voltage reached in trunk (blue) and tuft (red) for a range of simulations with different trunk lengths. Length constant λ = 1009 µm. **c.** Same as in *b* but plotting the width of the depolarization measured 2 mV above baseline. **d,e.** Peak voltage, width and integral values measured in the trunk and tuft for dendrites containing different Na^+ channel densities in the apical trunk.

We found that bAP amplitude in the tuft increased as a function of apical trunk length despite a decreasing bAP amplitude in the distal segment of the trunk (**Fig. 4b**). Conversely, the width of the bAP (measured 2 mV above baseline) increased in both the tuft and trunk with length (**Fig. 4c**). While waveform broadening is a natural consequence of passive filtering along dendrites, the sustained voltage in the distal trunk required active dendritic

propagation. In the reduced model, this active propagation in the apical trunk was mediated

primarily by voltage-gated Na^+ channels. Removing these channels caused a substantial

reduction in peak voltage and width of the depolarization in the distal trunk, and importantly

also abolished the trend of increasing tuft voltages with longer dendritic trunks (**Fig. 4d,e**).

More generally, active propagation caused bAPs to be larger and broader at all distances

along a long dendrite compared to the same absolute distances in shorter dendrites

(**Supplementary Figure 7**). Consequently, when comparing the final positions along the

trunk, the peak voltage is only marginally smaller in long dendrites despite the larger

distance from the soma. This is not the case in a passive dendrite, where voltage

attenuation depends on distance alone and is not sensitive to trunk length. We next tested

how the specific distribution of active conductances affected the results. When all

conductances were uniformly distributed along the apical trunk, the waveforms did not

substantially change, and the enhanced voltage continued to trigger Ca^+ plateaus only in

neurons with long apical trunks (**Supplementary Figure 8**). We have also tested the specific

contribution of the H-current (I_h). Reducing I_h in the tuft had no effect on the length

dependence of excitability. Reducing I_h in both the trunk and tuft resulted in the same

qualitative finding, having only a marginal quantitative effect (**Supplementary Figure 9**).

The general phenomenon of enhanced voltage propagation in longer dendrites resulting in

amplification of tuft voltage thus did not depend on any of the particular model parameters

above.

While it might seem counterintuitive that peak tuft voltage is increasing when the trunk

voltage is decreasing, we propose that the temporal broadening of the depolarization can at

least partially account for this via a passive mechanism. Wider depolarizations allow the tuft

compartment to charge to a higher voltage. The rate and peak value of tuft charging

depends on the passive properties of the tuft. The peak value of depolarization and the rate

of voltage change are proportional to membrane resistance (R_m) and membrane capacitance

(C_m), respectively. The product of these two parameters gives the membrane time constant

(τ). To illustrate this, we applied voltage-clamp to the end of the distal segment of the trunk

and delivered 30 mV square voltage pulses of increasing width (**Fig. 5a**). Due to capacitive

filtering, short voltage steps did not fully charge the tuft while wide voltage steps allowed

voltage to reach the steady-state values commanded by R_m. To directly test the hypothesis

that the relationship between trunk depolarization width and tuft membrane time constant

caused the bAP amplitude in the tuft to increase with length, one could vary R_m by changing

g_{leak}. However, this would affect resting membrane potential and consequently alter voltage-

dependent properties in the tuft. In order not to affect other variables in the model, we

therefore chose to vary C_m. For a given value of depolarization amplitude and width,

increasing C_m (and therefore τ) in the tuft caused a reduction in the peak tuft voltage (**Fig.**

5b). These simulations show that the tuft time constant and the width of the bAP interact to

create a higher tuft depolarization with longer apical trunks.

Figure 5. Tuft voltage increases with trunk length due to the widening of bAPs.

a. Different width voltage steps injected into the trunk (blue) under differing tuft membrane

capacitance conditions. Recorded tuft voltage in red. **b.** Peak voltage values reached in the tuft for a

range of trunk step widths and tuft capacitances. In the original model, tuft $C_m \cong 1.75\ \mu F/cm^2$. **c.**

Same as in *a* but showing voltage steps of different width and amplitude with the same voltage

integral in the trunk. For large integrals (top), the tuft voltage crosses the threshold for a Ca^{2+} plateau,

while for smaller integrals (bottom) the voltage remains subthreshold. **d.** Voltage and width

combinations for square voltage steps in the distal trunk which result in a Ca^{2+} plateau in the tuft. Red

points represent the values used in *h.* Grey lines indicate width and amplitude combinations with

equal integral.

It has previously been suggested that axial resistance (R_a) in the apical dendrite may

influence the backpropagation efficiency in dendrites and burstiness of ttL5 neurons

(Fletcher & Williams, 2019). To test this hypothesis, we measured peak voltage and width in

the trunk and tuft for different trunk lengths under different R_a conditions. We found that peak

tuft voltage (and therefore burstiness) increased with increasing trunk R_a, reaching the

highest voltage near the reduced model's original value, and decreasing again for higher

values (**Supplementary Figure 10**). However, in these simulations burstiness always

increased with trunk length regardless of R_a. This indicates that, although important, R_a was

not the primary determinant for generating the length-dependent effect and if R_a indeed

correlates with length, these effects may combine to further enhance the tuft voltage in long

neurons.

Overall, the combination of increased width and a relatively small reduction in amplitude

resulted in a trunk voltage integral that increased with trunk length, thereby passing more

charge to the adjacent tuft compartment. However, if active backpropagation was reduced or

absent, the trunk integral and resulting tuft voltage decreased with length (**Fig.4d,e**). The

peak tuft voltage approximately followed the integral of voltage in the distal trunk. To

illustrate this, we applied voltage-clamp to the end of the trunk and injected square steps

with a range of integrals obtained through various combinations of width and amplitude (**Fig.**

5c). This revealed a zone above a critical trunk integral for which many different width and

depolarization combinations were sufficient to evoke a Ca^{2+} plateau in the tuft (**Fig. 5d**).

Discussion

We made whole-cell patch-clamp recordings from layer 5 pyramidal neurons in primary and

secondary visual cortices of mice. We found that both BAC firing and critical frequency ADP

were almost entirely absent in the V2m neurons. Moreover, we observed that excitability

was positively correlated with the length of the apical dendrite trunk across all neurons. To

investigate the influence of apical trunk length on burstiness, we ran numerical simulations in

compartmental biophysical models. Both morphologically detailed and reduced models

showed that decreasing the apical trunk length resulted in reduced dendritic excitability.

Further simulations revealed that this is due to an interplay between bAP width, amplitude

and tuft impedance that depends critically on the presence of voltage-gated Na^+ channels in

the apical trunk. These results show that the same cell type, in closely related and adjacent

cortical areas, and under the same operating conditions can have a very different

computational repertoire. Our findings are thus inconsistent with the notion of canonical

computations at the single cell level, suggesting they may not exist at the circuit level either.

Contrary to common assumptions, we observed considerable differences in the properties of

ttL5 neurons across different brain regions. BAC firing, a dendritic operation, and critical

frequency ADP, a measure of dendritic excitability, which are both critically dependent on

dendritic Ca^{2+} plateaus, were almost completely absent in V2m. One key factor known to

control dendritic excitability is inhibition (Perez-Garci et al., 2006). To determine the role of

inhibition on the observed differences in excitability, we pharmacologically blocked $GABA_B$

receptors, thus creating more favourable conditions for BAC firing. To make comparisons

rigorous, we have used the same experimental protocols and conditions across all

experiments. Furthermore, to exclude bias in selecting ttL5 neurons, we have recorded only

from two well-defined groups, identified either by their projection target or genetic label. We

found similarly diminished excitability in both groups of V2m neurons.

The extracellular stimulation used to evoke BAC firing could in principle recruit different

polysynaptic circuits in V1 and V2m, which could account for the difference. However, this

would not explain the differences measured with the critical frequency paradigm using

intracellular stimulation only. Another confounding variable affecting excitability could be the

Ca^{2+} concentration in the extracellular solution. We have found the same results under

different physiologically relevant concentrations. Altogether, these results show that

decreased excitability of V2m ttL5 neurons is a robust phenomenon.

Correlated with the differences in excitability, we found ttL5 neurons in V2m to have

significantly shorter apical trunks compared to V1 neurons. This data is consistent with

recent structural MRI data (Fletcher & Williams, 2019) showing a thinner cortical mantle in

more medial and posterior parts of the cortex. Using an existing widely used biophysical

model designed to reproduce ttL5 properties such as BAC firing (Hay et al., 2011), we found

that the same model applied to a shorter morphology resulted in a loss of BAC firing,

independently of Ca^{2+} channel density. This was also true in a reduced ttL5 model with a

simplified morphological structure (Bahl et al., 2012), which allowed for continuous

exploration of apical trunk length. We found a sharp cut-off at a length of 0.35 λ ($\cong$ 350 μm

in model space), below which no BAC firing could be evoked. We note, however, that the

reduced model is based on rat neurons and the apical trunk and oblique dendrites are

pooled into the same compartment. It is also worth noting that, as the reduced model does

not have a distinct compartment to represent the apical bifurcation, all Ca^{2+} channels are

placed in the tuft compartment. Therefore, the numerical values of model length do not

translate directly into apical trunk lengths for real mouse neurons.

Through our simulations, we identified voltage-gated Na^+ channels in the apical dendrite as

a key factor for reproducing our results. The Na^+ channels control dendritic excitability by

supporting active backpropagation, resulting in reduced attenuation over distance.

Combined with a broadening of the bAP proportionally to trunk length (due to capacitive

filtering), this caused longer neurons to have larger voltage integrals in the distal trunk,

leading to greater charging of the tuft. Indeed, the peak tuft voltage depended on the amount

of passive charging it experienced, which was determined by the membrane time constant in

the tuft as well as by bAP width and amplitude. We hypothesised that, above a minimal

threshold for peak trunk voltage, the primary determinant of peak tuft voltage is the time-

averaged voltage in the trunk. Supporting this view, we found that the overall voltage integral

was more important for triggering Ca^{2+} plateaus than the particular combination of

depolarization width and amplitude.

Aside from dendritic length, our results do not exclude the involvement of other mechanisms

to modulate excitability. For example, differences in axial resistance could influence the way

voltage propagates along dendrites. While axial resistance indeed had a marked effect on

the backpropagation of action potentials in our models, this was independent of trunk length

and thus cannot account for our finding. Variations in density and activation properties of

other voltage-gated channels may also influence dendritic excitability. It is interesting to note

that, in the presence of Na^+ channels, the bAP in neurons with longer trunks was larger and

broader at every distance from the soma. This may be due to a cooperative effect of each

trunk section on the sections both up- and downstream, with the voltage at each location

decaying slower because of the more depolarised state of the remaining dendrite. Our data

thus predicts that bAP width and amplitude measured at the same absolute distance from

the soma differ across neurons with different apical trunk lengths.

There are a few notable counterexamples to the principle observed here. For example, the

human ttL5 neuron was recently shown to have greater compartmentalization and reduced

excitability compared to rat neurons despite being substantially longer (Beaulieu-Laroche et

al., 2018). This may still be consistent with our predictions, as human ttL5 neurons also had

reduced ion channel densities, which we show to be crucial for the length-dependent

enhancement. Furthermore, as the boosting effect of a broader depolarization is subject to

saturation (when the depolarization is wide enough to fully charge the tuft), we would expect

the positive effect of length on tuft voltage to not increase monotonically. Consequently,

beyond a certain length the trunk voltage would attenuate to the point where it is no longer

sufficient to trigger a Ca^{2+} plateau. On the other end of the spectrum, layer 2/3 pyramidal

neurons in rat barrel cortex have shorter apical trunks yet do show critical frequency ADP,

although they do not exhibit spike bursts during BAC firing (M. E. Larkum, Waters,

Sakmann, & Helmchen, 2007). It remains to be seen if the principle of length-dependent

excitability generalizes to other species, or across more widespread cortical areas and cell

types.

There may be important clinical implications to gaining a better understanding of how

variations in cortical thickness, and the resulting changes in neuronal morphology, affect the

physiology and computational properties of pyramidal neurons. Indeed, altered cortical

thickness has been implicated in several debilitating neurological diseases and mental

health conditions. For example, cortical thinning is used as a biomarker for Alzheimer's

disease (Dickerson et al., 2009) and correlates strongly with bipolar disorder (Hanford,

Nazarov, Hall, & Sassi, 2016), while increased cortical thickness is present during

development in individuals with autism (Khundrakpam, Lewis, Kostopoulos, Carbonell, &

Evans, 2017). Interestingly, altered dendritic excitability has also been strongly implicated in

several of these diseased conditions (Hall et al., 2015; Nanou & Catterall, 2018; Spratt et al.,

2019). We uncovered a possible mechanistic link between cortical thickness and excitability,

highlighting a new potential avenue of study for understanding the pathophysiology in these

conditions and raising the prospect of identifying intervention targets.

Our findings on dendritic excitability in ttL5 neurons have wide-ranging implications for

cortical computation. Feedback connectivity between cortical areas tends to target

superficial layers while feedforward input favours basal dendrites (Coogan & Burkhalter,

1990; Rockland & Pandya, 1979). BAC firing is believed to play a major role in integrating

these two pathways to modulate sensory perception (Takahashi, Oertner, Hegemann, &

Larkum, 2016) and to enable brain-wide learning algorithms that would otherwise be

intractable (Guerguiev, Lillicrap, & Richards, 2017; Sacramento, Ponte Costa, Bengio, &

Senn, 2018). This is particularly relevant in higher order cortical areas, which are more likely

to process brain-wide feedback to integrate convergent multisensory, motor and cognitive

signals (Freedman & Ibos, 2018). However, we show that, at least within the secondary

visual cortex, different operations must be implementing the multimodal integration of top-

down and bottom-up signals.

Our findings thus challenge the commonly held notion that the neocortex is composed of

canonical circuits performing stereotyped computations on different sets of inputs across the

brain (Douglas & Martin, 2004; Harris & Shepherd, 2015; Markram et al., 2015; Mountcastle,

1997). The heterogeneity of operating modes may expand the ability of cortical areas to

specialize in the computations that are required for processing their particular set of inputs,

at the cost of reduced flexibility in generalizing to other types of input. It may also imply that

the non-linear operations performed through Ca^{2+} plateaus and BAC firing may not be

required outside primary sensory cortices, perhaps because in these regions the input

hierarchy is less defined. It may therefore be more important to maintain equal weighting

between different sensory modalities and rely on other mechanisms to change the weights

according to the reliability of each input. With such simple morphological adjustments

capable of generating a wide range of possible operations, the brain undoubtedly leverages

the array of available computations to improve cognitive processing.

Materials and Methods

Animals

All animal experiments were prospectively approved by the local ethics panel of the Francis
Crick Institute (previously National Institute for Medical Research) and the UK Home Office
under the Animals (Scientific Procedures) Act 1986 (PPL: 70/8935). Wild-type and
transgenic male mice were used. Tg (Colgalt2-Cre)NF107Gsat (MGI:5311719, referred to
as Glt) and Tg (Rbp4-Cre)KL100Gsat (MGI:4367067, referred to as Rbp4) lines created
through the Gensat project (Gerfen, Paletzki, & Heintz, 2013; Groh et al., 2010) were
crossed with the Ai14 reporter line expressing tdTomato (MGI:3809524). Animals were
housed in individually ventilated cages under a normal 12-hour light/dark cycle.

Surgical procedures

Surgeries were performed on mice aged 3–7 weeks using aseptic technique under
isoflurane (2–4%) anaesthesia. Following induction of anaesthesia, animals were
subcutaneously injected with a mixture of meloxicam (2 mg/kg) and buprenorphine (0.1
mg/kg). During surgery, the animals were head-fixed in a stereotactic frame and a small hole
(0.5–0.7 mm) was drilled in the bone above the injection site. Alexa Fluor 488-conjugated
Cholera toxin subunit B (CTB, 0.8% w/v, Invitrogen) was injected using a glass pipette with
a Nanoject II (Drummond Scientific) delivery system at a rate of 0.4 nL/s. Injections of 100–
200 nL were targeted to the lateral posterior (LP) thalamic nucleus, with stereotactic
coordinates: 2.2–2.5 mm posterior to bregma, 1.45 lateral of the sagittal suture, 2.45 mm
deep from the cortical surface. To reduce backflow, the pipette was left in the brain
approximately 5 minutes after completion of the injection before being slowly retracted.

Slice preparation

Male mice (6–12 weeks old) were deeply anaesthetised with isoflurane and decapitated. In
mice that were injected with CTB, this occurred at least 3 weeks after the injection. The brain
was rapidly removed and placed in oxygenated ice-cold slicing ACSF containing (in mM):
125 sucrose, 62.5 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 2 $MgCl_2$, 1 $CaCl_2$, 25
dextrose; osmolarity 340–350 mOsm. Coronal slices (300 μm thick) containing visual cortex
were prepared using a vibrating blade microtome (Leica VT1200S or Campden 7000smz-2).
Slices were immediately transferred to a submerged holding chamber with regular ACSF
containing (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 1 $MgCl_2$, 1.5 or 2 $CaCl_2$,
25 dextrose; osmolarity 308–312 mOsm. The holding chamber was held in a water bath at
35 °C for the first 30–60 min after slicing and was kept at room temperature (22 °C) for the

remaining time (up to 12 hours) after that. All solutions and chambers were continuously

bubbled with carbogen (95% O_2 / 5% CO_2).

Electrophysiology

After the 35 °C incubation period, individual slices were transferred from the holding

chamber to the recording chamber, where they were perfused at a rate of ~6 mL/min with

regular ACSF (see above) continuously bubbled with carbogen and heated to 35 ± 1°C.

Borosilicate thick-walled glass recording electrodes (3-6 MΩ) were filled with intracellular

solution containing (in mM): 115 CH_3KO_3S, 5 NaCl, 3 $MgCl_2$, 10 HEPES, 0.05 EGTA, 3

Na_2ATP, 0.4 NaGTP, 5 K_2-phosphocreatine, 0.5% w/v biocytin hydrochloride (Sigma), 50

µM Alexa Fluor 488 hydrazide (Invitrogen); osmolarity 290-295 mOsm; pH 7.3. Visually

guided whole-cell patch-clamp recordings were targeted to neurons in L5 of medial V2

(V2m) and V1 that were fluorescently labelled with either CTB or with tdTomato (for Glt25d2-

Cre mice), to ensure that the recordings were restricted to ttL5 neurons. Visual areas were

defined based on approximate stereotactic coordinates (Franklin & Paxinos, 2007). All

recordings were made in current-clamp mode. Extracellular monopolar stimulation was

achieved by passing a DC current pulse (0.1-1 ms, 20-320 µA) through a glass patch-clamp

pipette with a broken tip (~20 µm diameter) using a constant current stimulator (Digitimer

DS3). Current was passed between two silver / silver chloride (Ag/AgCl) wires: one inside

the pipette, which was filled with recording ACSF, and the other coiled around the outside of

the pipette. In experiments using extracellular stimulation, 1 µM CGP52432 was added to

the ACSF.

Immunohistochemistry and morphological reconstructions

After recording, slices were fixed overnight at 4 °C in a 4% formaldehyde solution and were

subsequently kept in PBS. For immunohistochemical detection, the fixed slices were first

incubated for 1-2 hours at room temperature in blocking solution containing 0.5% Triton X-

100 and 5% Normal Goat Serum (NGS) in PBS. Slices were then washed twice (10 min

each) in PBS and incubated overnight in a staining solution containing 0.05% Triton X-100,

0.5% NGS, DyLight 594-conjugated streptavidin (2 µg/ml). Slices were then washed in PBS

(3 times, 5 min each) and stained with DAPI (5 µg/ml) for 10 min. After another wash (3

times, 5 min each), slices were mounted on glass slides and images were acquired with a

confocal microscope (Leica SP5, objective: 20x/0.7NA or 10x/0.4NA, pinhole size: 1 airy

unit). The images were used to reconstruct the neurons with Neurolucida 360 (MBF

bioscience).

Data acquisition and analysis

Recorded signals were amplified and low-pass filtered through an 8 kHz Bessel filter using a

MultiClamp 700B amplifier (Molecular Devices). Filtered signals were then digitized at 20

kHz with a National Instruments DAQ board (PCIe-6323). Acquisition and stimulus protocols

were generated in Igor Pro with the Neuromatic software package (Rothman & Silver, 2018).

Further analysis and data visualization were performed with custom macros and scripts

written in Igor Pro and Matlab (MathWorks). Raincloud plots (consisting of a scatter plot, a

box plot, and a kernel density plot) were generated in MATLAB using published scripts

(Allen, Poggiali, Whitaker, Marshall, & Kievit, 2019). All box plots presented show the

median, interquartile range, 2nd and 98th percentile of the dataset. Confocal images were

processed with Fiji (https://fiji.sc/).

Modelling

Simulations were performed with the NEURON (Carnevale & Hines, 2006) simulation

environment (7.7.1) embedded in Python 3.6. To model the consequences of morphological

differences between V1 and V2m ttL5 cells, we utilised existing models of ttL5 pyramidal

cells with either accurate morphological detail (biophysical model 3, cell #1 from Hay et al.

2011 (Hay et al., 2011), referred to as detailed model) or simplified multicompartment

morphologies (Ca^{2+} enriched model 2 from Bahl et al. 2012 (Bahl et al., 2012), referred to

as reduced model). To study the effect of morphology in the detailed model, biophysical

model 3 from Hay et al. 2011 (Hay et al., 2011) was applied to the reconstructed morphology

from one of our recorded ttL5 neurons in V2m (which has a substantially shorter apical trunk

than the morphology used in the original model). Each morphology contained low-voltage-

activated (LVA) and high-voltage-activated (HVA) Ca^{2+} channels located in a 100-200 µm

long region around the main apical bifurcation.

Subsequent simulations using the reduced model were done by modifying only selected

parameters described in the results, such as the length of the apical trunk compartment,

leaving all other parameters unchanged. Briefly, this reduced model (Bahl et al., 2012) is

divided into sections representing the soma, axon (hillock and initial segment, AIS), basal

dendrites, apical trunk, and apical tuft. Active conductances are present in all compartments

and include the following: hyperpolarization-activated cation (HCN) channels (basal

dendrite, apical trunk, tuft), transient voltage-activated Na^+ (Na_t) channels (soma, axon

hillock, AIS, apical trunk, tuft), persistent voltage-activated Na^+ (Na_p) channels (soma), fast

voltage-activated K^+ (K_{fast}) channels (soma, apical trunk, tuft), slow voltage-activated K^+

(K_{slow}) channels (soma, apical trunk, tuft), muscarinic K^+ (K_m) channels (soma), slow Ca^{2+}

(Ca_s) channels (tuft), Ca^{2+} dependent K^+ (K_{Ca}) channels (tuft), and a Ca^{2+} pump (tuft). The

density of the K_{fast} and K_{slow} channels decays exponentially from the soma to the tuft. The

density of Na_t channels decays linearly from the soma to the tuft, while HCN channels

linearly increase in density. N.B. the tuft, but not the trunk, contains Ca^{2+} channels;

consequently, there is no hotspot similar to the apical bifurcation in the detailed model.

When varying trunk length, Na_t, K_{fast}, K_{slow}, and HCN conductances in each trunk segment

were redistributed so as to take into account the new distance of each segment from the

soma (thereby changing the total conductance in the trunk). All code will be available on

GitHub after acceptance.

Acknowledgements

We thank Arnd Roth and Alexandra Tran-Van-Minh for advice on modelling. We are grateful

to Lee Fletcher, Arnd Roth, Alexandra Tran-Van-Minh, Michael Hausser and Anna Cappellini

for helpful comments on the manuscript.

Author contributions

A.R.G.: Designed and performed biological experiments, designed modelling experiments,

analysed data, wrote the paper. A.L.: Designed and performed modelling experiments,

analysed data. E.A.R.: Conceptualized research, designed experiments, acquired funding,

wrote the paper.

All authors discussed the results and implications and commented on the manuscript at all

stages. There are no competing interests.

All authors declare no financial or non-financial competing interest.

Funding

A.R.G. was recipient of a Boehringer Ingelheim Fonds PhD Scholarship and a PhD

scholarship from the Francis Crick Institute. E.A.R. is jointly funded by Wellcome and the

Royal Society through a Sir Henry Dale Fellowship (104285/B/14/Z).

References

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., & Kievit, R. A. (2019). Raincloud plots: a multi-platform tool for robust data visualization. *Wellcome Open Res, 4*, 63. doi:10.12688/wellcomeopenres.15191.1

Bahl, A., Stemmler, M. B., Herz, A. V. M., & Roth, A. (2012). Automated optimization of a reduced layer 5 pyramidal cell model based on experimental data. *Journal of Neuroscience Methods, 210* (1), 22-34. doi:10.1016/j.jneumeth.2012.04.006

Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., & Friston, K. J. (2012). Canonical microcircuits for predictive coding. *Neuron, 76* (4), 695-711. doi:10.1016/j.neuron.2012.10.038

Beaulieu-Laroche, L., Toloza, E. H. S., van der Goes, M.-S., Lafourcade, M., Barnagian, D., Williams, Z. M., . . . Harnett, M. T. (2018). Enhanced Dendritic Compartmentalization in Human Cortical Neurons. *Cell, 175* (3), 643-651.e614. doi:10.1016/j.cell.2018.08.045

Braganza, O., & Beck, H. (2018). The Circuit Motif as a Conceptual Tool for Multilevel Neuroscience. *Trends Neurosci, 41* (3), 128-136. doi:10.1016/j.tins.2018.01.002

Carlo, C. N., & Stevens, C. F. (2013). Structural uniformity of neocortex, revisited. *Proc Natl Acad Sci U S A, 110* (4), 1488-1493. doi:10.1073/pnas.1221398110

Carnevale, N. T., & Hines, M. L. (2006). *The NEURON book*. Cambridge, UK ; New York: Cambridge University Press.

Coogan, T. A., & Burkhalter, A. (1990). Conserved patterns of cortico-cortical connections define areal hierarchy in rat visual cortex. *Experimental Brain Research, 80* (1), 49-53. doi:10.1007/BF00228846

Dickerson, B. C., Bakkour, A., Salat, D. H., Feczko, E., Pacheco, J., Greve, D. N., . . . Buckner, R. L. (2009). The cortical signature of Alzheimer's disease: regionally specific cortical thinning relates to symptom severity in very mild to mild AD dementia and is detectable in asymptomatic amyloid-positive individuals. *Cerebral Cortex, 19* (3), 497-510. doi:10.1093/cercor/bhn113

Douglas, R. J., & Martin, K. A. C. (2004). Neuronal Circuits of the Neocortex. *Annual Review of Neuroscience, 27* (1), 419-451. doi:10.1146/annurev.neuro.27.070203.144152

Fletcher, L. N., & Williams, S. R. (2019). Neocortical Topology Governs the Dendritic Integrative Capacity of Layer 5 Pyramidal Neurons. *Neuron, 101* (1), 76-90.e74. doi:10.1016/j.neuron.2018.10.048

Franklin, K. B. J., & Paxinos, G. (2007). *Paxinos and Franklin's The mouse brain in stereotaxic coordinates* (3rd edition. ed.). Amsterdam: Academic Press, an imprint of Elsevier.

Freedman, D. J., & Ibos, G. (2018). An Integrative Framework for Sensory, Motor, and Cognitive Functions of the Posterior Parietal Cortex. *Neuron, 97* (6), 1219-1234. doi:10.1016/j.neuron.2018.01.044

Gerfen, Charles R., Paletzki, R., & Heintz, N. (2013). GENSAT BAC Cre-Recombinase Driver Lines to Study the Functional Organization of Cerebral Cortical and Basal Ganglia Circuits. *Neuron, 80* (6), 1368-1383. doi:10.1016/j.neuron.2013.10.016

Gidon, A., Zolnik, T. A., Fidzinski, P., Bolduan, F., Papoutsi, A., Poirazi, P., . . . Larkum, M. E. (2020). Dendritic action potentials and computation in human layer 2/3 cortical neurons. *Science, 367* (6473), 83-87. doi:10.1126/science.aax6239

Groh, A., Meyer, H. S., Schmidt, E. F., Heintz, N., Sakmann, B., & Krieger, P. (2010). Cell-Type Specific Properties of Pyramidal Neurons in Neocortex Underlying a Layout that Is Modifiable Depending on the Cortical Area. *Cerebral Cortex, 20* (4), 826-836. doi:10.1093/cercor/bhp152

Guerguiev, J., Lillicrap, T. P., & Richards, B. A. (2017). Towards deep learning with segregated dendrites. *eLife, 6*. doi:10.7554/eLife.22901

Hall, A. M., Throesch, B. T., Buckingham, S. C., Markwardt, S. J., Peng, Y., Wang, Q., . . . Roberson, E. D. (2015). Tau-dependent Kv4.2 depletion and dendritic hyperexcitability in a mouse model of Alzheimer's disease. *J Neurosci, 35* (15), 6221-6230. doi:10.1523/JNEUROSCI.2552-14.2015

Hanford, L. C., Nazarov, A., Hall, G. B., & Sassi, R. B. (2016). Cortical thickness in bipolar disorder: a systematic review. *Bipolar Disord, 18* (1), 4-18. doi:10.1111/bdi.12362

Harris, K. D., & Shepherd, G. M. G. (2015). The neocortical circuit: themes and variations. *Nature Neuroscience, 18* (2), 170-181. doi:10.1038/nn.3917

Hawkins, J., Ahmad, S., & Cui, Y. (2017). A Theory of How Columns in the Neocortex Enable Learning the Structure of the World. *Front Neural Circuits, 11*, 81. doi:10.3389/fncir.2017.00081

Hay, E., Hill, S., Schürmann, F., Markram, H., & Segev, I. (2011). Models of Neocortical Layer 5b Pyramidal Cells Capturing a Wide Range of Dendritic and Perisomatic Active Properties. *PLOS Computational Biology, 7* (7). doi:10.1371/journal.pcbi.1002107

Jarsky, T., Roxin, A., Kath, W. L., & Spruston, N. (2005). Conditional dendritic spike propagation following distal synaptic activation of hippocampal CA1 pyramidal neurons. *Nature Neuroscience, 8* (12), 1667-1676. doi:10.1038/nn1599

Khundrakpam, B. S., Lewis, J. D., Kostopoulos, P., Carbonell, F., & Evans, A. C. (2017). Cortical Thickness Abnormalities in Autism Spectrum Disorders Through Late Childhood, Adolescence, and Adulthood: A Large-Scale MRI Study. *Cerebral Cortex, 27* (3), 1721-1731. doi:10.1093/cercor/bhx038

Larkum, M. (2013). A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex. *Trends in Neurosciences, 36* (3), 141-151. doi:10.1016/j.tins.2012.11.006

Larkum, M. E., Kaiser, K. M. M., & Sakmann, B. (1999). Calcium electrogenesis in distal apical dendrites of layer 5 pyramidal cells at a critical frequency of back-propagating action potentials. *Proceedings of the National Academy of Sciences, 96* (25), 14600-14604. doi:10.1073/pnas.96.25.14600

Larkum, M. E., Waters, J., Sakmann, B., & Helmchen, F. (2007). Dendritic spikes in apical dendrites of neocortical layer 2/3 pyramidal neurons. *J Neurosci, 27* (34), 8999-9008. doi:10.1523/JNEUROSCI.1717-07.2007

Larkum, M. E., Zhu, J. J., & Sakmann, B. (1999). A new cellular mechanism for coupling inputs arriving at different cortical layers. *Nature, 398* (6725), 338-341. doi:10.1038/18686

London, M., & Häusser, M. (2005). Dendritic Computation. *Annual Review of Neuroscience, 28* (1), 503-532. doi:10.1146/annurev.neuro.28.061604.135703

Mainen, Z. F., & Sejnowski, T. J. (1996). Influence of dendritic structure on firing pattern in model neocortical neurons. *Nature, 382* (6589), 363-366. doi:10.1038/382363a0

Markram, H., Muller, E., Ramaswamy, S., Reimann, M. W., Abdellah, M., Sanchez, C. A., . . . Schurmann, F. (2015). Reconstruction and Simulation of Neocortical Microcircuitry. *Cell, 163* (2), 456-492. doi:10.1016/j.cell.2015.09.029

Miller, K. D. (2016). Canonical computations of cerebral cortex. *Current Opinion in Neurobiology, 37*, 75-84. doi:10.1016/j.conb.2016.01.008

Mountcastle, V. B. (1997). The columnar organization of the neocortex. *Brain, 120* (4), 701-722. doi:10.1093/brain/120.4.701

Nanou, E., & Catterall, W. A. (2018). Calcium Channels, Synaptic Plasticity, and Neuropsychiatric Disease. *Neuron, 98* (3), 466-481. doi:10.1016/j.neuron.2018.03.017

Perez-Garci, E., Gassmann, M., Bettler, B., & Larkum, M. E. (2006). The GABAB1b isoform mediates long-lasting inhibition of dendritic Ca2+ spikes in layer 5 somatosensory pyramidal neurons. *Neuron, 50* (4), 603-616. doi:10.1016/j.neuron.2006.04.019

Rockland, K. S., & Pandya, D. N. (1979). Laminar origins and terminations of cortical
connections of the occipital lobe in the rhesus monkey. *Brain Research, 179* (1), 3-20.
doi:10.1016/0006-8993 (79)90485-2

Rothman, J. S., & Silver, R. A. (2018). NeuroMatic: An Integrated Open-Source Software
Toolkit for Acquisition, Analysis and Simulation of Electrophysiological Data. *Front*
Neuroinform, 12, 14. doi:10.3389/fninf.2018.00014

Sacramento, J., Ponte Costa, R., Bengio, Y., & Senn, W. (2018). Dendritic cortical
microcircuits approximate the backpropagation algorithm. In S. Bengio, H. Wallach, H.
Larochelle, K. Grauman, N. Cesa-Bianchi, & R. Garnett (Eds.), *Advances in Neural*
Information Processing Systems 31 (pp. 8721-8732): Curran Associates, Inc.

Schaefer, A. T., Larkum, M. E., Sakmann, B., & Roth, A. (2003). Coincidence Detection in
Pyramidal Neurons Is Tuned by Their Dendritic Branching Pattern. *Journal of*
Neurophysiology, 89 (6), 3143-3154. doi:10.1152/jn.00046.2003

Shai, A. S., Anastassiou, C. A., Larkum, M. E., & Koch, C. (2015). Physiology of Layer 5
Pyramidal Neurons in Mouse Primary Visual Cortex: Coincidence Detection through
Bursting. *PLOS Computational Biology, 11* (3). doi:10.1371/journal.pcbi.1004090

Shipp, S. (2016). Neural Elements for Predictive Coding. *Front Psychol, 7*, 1792.
doi:10.3389/fpsyg.2016.01792

Spratt, P. W. E., Ben-Shalom, R., Keeshen, C. M., Burke, K. J., Jr., Clarkson, R. L., Sanders,
S. J., & Bender, K. J. (2019). The Autism-Associated Gene Scn2a Contributes to
Dendritic Excitability and Synaptic Function in the Prefrontal Cortex. *Neuron, 103* (4),
673-685 e675. doi:10.1016/j.neuron.2019.05.037

Spruston, N. (2008). Pyramidal neurons: dendritic structure and synaptic integration. *Nat Rev*
Neurosci, 9 (3), 206-221. doi:10.1038/nrn2286

Takahashi, N., Oertner, T. G., Hegemann, P., & Larkum, M. E. (2016). Active cortical
dendrites modulate perception. *Science, 354* (6319), 1587-1590.
doi:10.1126/science.aah6066

van Ooyen, A., Duijnhouwer, J., Remme, M. W. H., & van Pelt, J. (2002). The effect of
dendritic topology on firing patterns in model neurons. *Network (Bristol, England), 13*
(3), 311-325.

Vetter, P., Roth, A., & Hausser, M. (2001). Propagation of action potentials in dendrites
depends on dendritic morphology. *J Neurophysiol, 85* (2), 926-937.
doi:10.1152/jn.2001.85.2.926

 Supplementary figures

 Supplementary Figure 1. Confocal images from a Glt25d2-Cre mouse injected with CTB-Alexa Fluor 488 in LP. a. Coronal slice of one hemisphere containing visual cortex, showing the injection site and retrogradely labelled neurons (green), tdTomato-expressing Glt25d2-Cre neurons (magenta), a DAPI stain (blue), and neurons that have been filled with biocytin during intracellular recordings and stained with DyLight 594 (cyan). **b.** Biocytin-filled ttL5 neuron in V2m. **c.** Neighbouring tdTomato-expressing Glt25d2-Cre neurons. **d.** CTB-labelled L5 neurons projecting to LP. **e.** Composite image of the above. Arrowheads highlight two example cells labelled by both tdTomato and CTB.

Supplementary Figure 2. Maximum ADP integral for all cells split by recording ACSF containing either 1.5 or 2 mM $CaCl_2$. p values for two-sample Kolmogorov-Smirnov test.

Supplementary Figure 3. Maximum ADP integral for all cells including V1 (grey), retro-labelled cells recorded from V2m (dark blue), and V2m cells labelled in the Gltd2-Cre mouse line (light blue). p value for two-sample Kolmogorov-Smirnov test.

777
778 **Supplementary Figure 4.** Representative voltage traces for V1
779 and V2m neurons in response to 500 ms wide depolarizing
780 current steps. Stimulation at 60 pA (grey) and 180 pA (coloured
781 or black) above rheobase (200 pA for both).

782
783

784
Supplementary Figure 5. Ca^{2+} hot-spot size does not affect
excitability.
a. Voltage traces recorded from different parts of the long
neuronal morphology containing either a 200 μm (as in the
original), or 100 μm long Ca^{2+} region in the distal apical trunk. **b.**
Same as in a, but for the shorter morphology.

791
792

Supplementary Figure 6. Large current injections to the nexus of short model cells can trigger calcium plateaus.
Same as in *Fig. 3a & b*, but for the short morphology using the same dendritic current injection as in the long morphology (0.5 nA).

802
Supplementary Figure 7. Backpropagation of APs in active
and passive trunks of different length.
a. Backpropagation of a somatic spike elicited through a single 3
ms wide 2 nA square current step at the soma in a model neuron
with 600 µm apical trunk length. Voltage recordings at different
distances along the trunk. **b.** Peak voltage and width measured
at different absolute distances (same relative) for active model
neurons. Width was measured as the interval between the
voltage values 2 mV above baseline membrane potential.
Colours indicate models with different apical trunk lengths. N.B.
At any given absolute distance from the soma, peak voltage and
width of the bAP are larger when the apical trunk is longer. **c.**
Same as in *b* but with all voltage-dependent conductances
removed from the trunk and tuft compartments. N.B. voltage
attenuation is independent of trunk length.

Supplementary Figure 8. Tuft voltage increases with trunk length independently of conductance gradients.
Same experiments as in **Fig. 3a** but with uniform distribution of all active conductances in the apical trunk. Total conductance was maintained for each channel.

Supplementary Figure 9. I_h does not contribute to length dependent increase of excitability.
a. Plots of peak voltage, width and integral of bAPs under different g_{HCN} (shades) in the tuft, both in the presence (solid) or absence (dashed) of Ca^{2+} channels. **b.** Same as *a*, but varying g_{HCN} in both tuft and trunk compartments.

Supplementary Figure 10. Effect of axial resistance on voltage propagation.

Plots of peak voltage, width, and voltage integral reached in the tuft and trunk for varying values of trunk length and axial resistance (R_a). Default $R_a \cong 382.22$ $\Omega \cdot$cm. The stimulus and recording conditions were the same as in **Fig. 2a**, with 3 APs at 100 Hz triggered in the somatic compartment. Solid lines show simulations with $g_{Ca} = 0$, while dashed lines show the same simulations with the original gCa = 0.45 mS/cm^2.