Title: Stronger between-clan than within-clan contests and their ecological correlates in a non-

territorial, fission-fusion species, the Asian elephant

Authors: Hansraj Gautam and T.N.C. Vidya*

Author email address: Hansraj Gautam (gautamhansraj@yahoo.com), T.N.C. Vidya

(Corresponding author, tncvidya@jncasr.ac.in)

Affiliation: Evolutionary and Organismal Biology Unit, Jawaharlal Nehru Centre for Advanced

Scientific Research (JNCASR), Bangalore, India, 560064.

* Corresponding author: T.N.C. Vidya

Address: Evolutionary and Organismal Biology Unit, Jawaharlal Nehru Centre for Advanced

Scientific Research (JNCASR), Bangalore, India, 560064. Email id: tncvidya@jncasr.ac.in,

tncvidya@gmail.com, Phone number: +91-80-22082968.

Abstract

1. Socioecological theory attributes variation in social organization of female-bonded species to

differences in within- and between-group competition, shaped by food distribution. Strong

between-group contests are expected over large, monopolisable resources, and are not generally

expected in species that feed on low quality resources distributed across large, undefended home

ranges. Within groups, frequent contests are expected over discrete feeding sites but not over low-

quality, dispersed resources.

2. We report on the first tests of socioecological theory, largely unexplored in non-primate species,

in female Asian elephants. Asian elephants show graminivory, overlapping home ranges, and high

fission-fusion dynamics, traits that are thought to be associated with infrequent contests.

3. We studied agonistic interactions within and between female elephant clans with respect to food

distribution, food abundance, and competitor density effects of group size and clan density, in a

grassland habitat around the Kabini backwaters, southern India.

4. We found that the Kabini grassland had three times the grass biomass as adjacent forests, and

between-clan encounters were considerably higher than that known from a neighbouring forest.

Individual-level agonism was also more frequent between clans than within clans. Thus, the food-

rich habitat patch probably enabled strong between-clan contest competition under graminivory.

Moreover, the rate of between-clan encounters increased when more clans were present, and the

duration of encounters was positively related to grass biomass at the contested sites. Despite

fission-fusion dynamics, within-clan agonism was also somewhat frequent, but not influenced by

food distribution, in contradiction to classic socioecological predictions, possibly because of

intensified competition due to high density. Interestingly, within-clan agonism increased with

female group size until intermediate group sizes, suggesting that the tension between within-group

and between-group competition might govern group size, since larger groups are advantageous in

this strong between-clan contest regime.

5. Our findings refine the current understanding of female elephant socioecology. Despite

predominant graminivory and fission-fusion dynamics, within-group agonism can be frequent,

especially when large groups face ecological constraints at high density. Further, frequent

between-group contests may arise despite graminivory and non-territoriality when food becomes

patchy and density is high. These changes may be effected by anthropogenic alteration of habitats.

Keywords

Asian elephant, contest competition, food distribution, fission-fusion dynamics, group size,

socioecological theory, within-group and between-group agonism.

Main text

Introduction

Socioecological theory attempts to explain variation in animal societies based on resource-risk

distributions (Crook and Gartlan, 1966; Terborgh and Janson, 1986). The ecological model of

female social relationships (EMFSR), part of the larger socioecological framework, proposes that

feeding competition is pivotal in shaping social organisation and structure in female-bonded

groups (Wrangham, 1980; van Schaik, 1989; Isbell, 1991; Sterck *et al.*, 1997). Food distribution,

abundance, and quality are expected to shape feeding competition regimes. Within groups,

scramble or exploitative competition is expected to predominate when limited food is dispersed

and cannot be usurped or monopolized by individuals, whereas strong contest or interference

competition is expected when high quality food is present in usurpable clumps or patches (Janson,

1985; van Schaik, 1989). Between groups, contest competition is expected when food patches are

large and can be usurped by groups (Wrangham, 1980; Isbell, 1991). Strong and simultaneous

between-group and within-group contest are expected to suppress within-group aggression via

greater tolerance of subordinate group members by the dominants in order to retain the advantages

of larger groups (see Sterck *et al.*, 1997).

Mixed empirical findings from tests of the EMFSR have led to calls for using field-based

quantification of food characteristics rather than indirect indicators such as diet type to test the

ecological basis of contest competition (Snaith and Chapman, 2007; Koenig and Borries, 2009;

Koenig *et al.*, 2013; Wheeler *et al.*, 2013). Contest competition may also be influenced by

competitor density, with group size and group spread affecting contests (Koenig and Borries, 2006;

Wheeler *et al.*, 2013; also see van Schaik *et al.*, 1983). Despite the EMFSR not being conceptually

restricted to specific taxa, it has seldom been examined in non-primate mammals (but see, for

example, Monaghan and Metcalfe, 1985; Wittemyer *et al.*, 2007; Smith *et al.*, 2008). Even in

primates, there has been greater emphasis on the relationship between food distribution and within-

group contest (for example, Janson, 1985; Koenig *et al.*, 1998; Pruetz and Isbell, 2000; Korstjens

et al., 2002; Koenig and Borries, 2006) than with between-group contest (for example, Wilson *et*

al., 2012; Brown, 2013; Roth and Cords, 2016; Pal *et al.*, 2018); many studies of between-group

contest have focused on territorial behaviour or outcomes of contests rather than the link with food

distribution. Moreover, studies of within- and between-group contests have been carried out more

often on frugivores (for example, Janson, 1985; Vogel and Janson, 2011) than on folivores (for

example, Koenig *et al.*, 1998; Harris, 2006), which were initially thought to face reduced feeding

competition because of their seemingly low quality or continuously distributed and abundant diet

(Wrangham, 1980; van Schaik, 1989; Isbell, 1991). In the context of the above, we examined the

relationship between food resources and rates of within- and between-group contests in a non-

primate species, the Asian elephant (*Elephas maximus*), feeding primarily on grass, traditionally

thought of as a low-quality resource, in the Kabini population, southern India.

Asian elephants exhibit female-bonded groups (Sukumar, 1989; Fernando and Lande, 2000; de

Silva *et al.*, 2011), with the most inclusive social unit being the clan (Nandini *et al.* 2018),

equivalent to a social group or community in other fission-fusion species (see Aureli *et al.* 2008).

Females within clans show fission-fusion dynamics, in which clan-members are usually distributed

across multiple groups (or parties), whose group sizes and compositions can change across hours

(Nandini *et al.*, 2017, 2018). We defined a "group" as a "party" of individuals (almost always from

the same clan) seen in the field, with individuals usually within ~50-100 m of one another and

showing coordinated movement (see Nandini *et al.* 2018). Asian elephants are not territorial and

the home ranges of clans may overlap extensively (Baskaran and Desai, 1996), a trait that was

expected to relate to infrequent aggression during between-group encounters (Cheney, 1987, but

see Willems and van Schaik 2015). Fission-fusion dynamics may also decrease competition (see

de Silva *et al.*, 2017) within clans. Adult females are not known to face infanticide risks, and face

negligible predation risk. Therefore, food characteristics are primarily expected to shape group

size and competition regimes in this species. Moreover, as female and male elephant societies are

largely separate (see Keerthipriya *et al.*, 2021), within- and between-group contests do not

typically involve the participation of adult males, unlike that seen in some primates. Thus, the

Asian elephant is an excellent non-primate species that shows female-bondedness and fission-

fusion for examining predictions of the EMFSR.

The Kabini elephant population is part of the largest contiguous population of Asian elephants in

the world, and has been monitored since 2009, with several hundred individuals and their clans

identified (see Methods). Individual-level agonistic contests occur between adult females within,

as well as between, clans (Nandini, 2016). We classified these as individual-level within-clan and

individual-level between-clan agonism, respectively. As females in a group maintain spatial

cohesion during and after encounters with groups from another clan, we also examined agonistic

encounters at the level of groups (parties) belonging to different clans, which we term clan-level

between-clan agonism; this could involve one or more individual-level interactions between

females from the contesting clans. A weak dominance hierarchy had been found in another Asian

elephant population, and had been attributed to fission-fusion dynamics and high habitat

productivity (de Silva *et al.*, 2017); however, food resources had not been directly quantified and

related to contest competition in any elephant species (but see Wittemyer *et al.*, 2007 for relating

remotely-sensed productivity to between-group dominance relations). Here, we directly quantified

food abundance and distribution in a small grassland habitat, and examined their relationship with

the frequencies of agonistic interactions within and between clans.

We addressed the following specific questions to understand how elephant societies are shaped by

socioecological factors.

1) Is grass abundance in the Kabini grassland different from that in the neighbouring forest

habitat, and how is it distributed across and within different areas (focal zones) in the grassland?

Since between-group contests appeared to be more frequent in the Kabini grassland (Nandini,

2016) compared to those in a neighbouring forest habitat of the Nilgiri Biosphere Reserve

(Baskaran, 1998), we wanted to find out whether the grassland was more resource-rich than the

neighbouring forests, possibly explaining the frequent between-group contests. We also quantified

variation within the grassland, for examining its relationship with between- and within-group

agonism (see 3-5 below).

2) Are the rates and intensities of individual-level agonism among females different within and

between clans?

The EMFSR predicts strong between-group contest over large food patches, and predominantly

within-group contest when feeding sites are clumped at the local level (Sterck *et al.*, 1997; Koenig,

2002). Therefore, we expected a higher rate of individual-level agonism (see Methods) during

between-clan encounters than within clans, even after controlling for the number of females

present, if the grassland was a food-rich habitat with grass abundance varying across focal zones.

In contrast, we expected no difference in the rates of individual-level agonism between and within

clans if grass abundance in the grassland and forest was similar and if there was little variation

across zones, since females would not be expected to give up their feeding time to participate in

between-clan agonistic interactions if the patches were not discrete enough to be usurped by their

clan.

3) Is the rate of within-clan agonism explained by variation in grass abundance, grass dispersion,

and group size?

Based on the classical prediction of the EMFSR (Koenig *et al.*, 1998; Pruetz and Isbell, 2000), we

expected the rate of (individual-level) within-clan agonism to be greater when grass was more

clumped at a local scale. We also expected group size, which reflects local competitor density

(Koenig and Borries, 2006), to positively correlate with within-clan agonism.

4) Is the rate of individual-level between-clan agonism explained by variation in grass abundance,

grass dispersion, and group size?

We expected the rate of individual-level between-clan agonism to be positively related to food

abundance at the site of contest and to the total number of females present. Further, we expected

this rate to be higher when the competing clans were evenly matched in group size, a determinant

of resource holding potential of groups (for example, Roth and Cords, 2016).

5) Are the rate and duration of clan-level between-clan agonistic encounters related to grass

abundance/distribution, group size, or the number of clans?

We expected more frequent clan-level between-clan agonistic encounters in focal zones with more

abundant grass and with more heterogeneously distributed grass (see Methods). Further, based on

game theoretical predictions about animal contests (see Smith and Parker, 1976), we expected the

duration of clan-level agonistic encounters to depend positively upon food abundance at the site

of contest, and negatively upon the difference in group sizes of the competing clans.

Methods

Study area

We carried out the study in Nagarahole National Park and Tiger Reserve in the Nilgiris-Eastern

Ghats landscape, southern India (Fig. 1). Nagarahole largely comprises deciduous forest with

grass, herb, and shrub understorey layers, but an open grassland (length <15 km, maximum width

<2 km, SI 1-2) is formed by the receding backwaters of the Kabini reservoir during the dry season.

Hundreds of elephants that use this grassland/reservoir and the surrounding forests have been

identified based on natural physical characteristics and monitored as part of the Kabini Elephant

Project (Vidya *et al*., 2014). Elephants feed almost exclusively on grass in the grassland after the

bamboo die-off following mass-flowering in 2011. The abundance of elephant food plants varies

spatio-seasonally and is lower in the dry than the wet season in the adjacent Nagarahole forest

(Gautam *et al*., 2019). Multiple clans can use the Kabini grassland simultaneously and show

between-clan aggression.

Data collection

Sampling grass abundance and distribution

We sampled six grassland stretches, henceforth called focal zones (Fig. 1, SI 2 Fig. S2), around

the Kabini backwaters during four periods of ~30 days each (referred to as months) in the dry

seasons of 2015 and 2016. In each focal zone, on one day in the middle of each month, we sampled

20 1-m^2 quadrats to visually estimate grass cover (see Gautam *et al.*, 2017), measure grass height

(average of the natural standing heights of grasses at ten locations), and harvest and weigh above-

ground fresh grass biomass in each quadrat (SI 2 Fig. S4). These 20 quadrats were distributed

within the focal zone in 4 clusters (henceforth, plot-clusters) of 5 quadrats each (Fig.1c) to assess

the variability of grass abundance at a local scale. While the centre of each plot-cluster was fixed

across months, we laid the five constituent quadrats each month by choosing random distances up

to 100 m along random directions. Random sampling was chosen because grass was continuous

within focal zones and it was not possible to visually detect distinct patches, unlike individual trees

that might be considered food patches.

We used harvested grass biomass data collected previously from the forest habitat of Nagarahole

National Park (Gautam *et al.*, 2017, see analysis and SI 4) to compare grass abundance between

the grassland and forest habitat. We also confirmed the pattern using grass biomass data collected

simultaneously in the dry season of 2022 from 25 plots in Nagarahole forest and 24 plot-clusters

(4 plot-clusters × 6 zones) in the Kabini grassland, as part of an ongoing study (SI 4 text).

Sampling within-clan and between-clan agonism

We carried out full-day (~6:30 AM to ~6:30 PM) observations of focal zones to quantify elephant

visits and agonistic interactions between elephants. We selected focal zones sequentially, and

sampled each zone on at least three sampling days per month in 2015 and four sampling days per

month in 2016. On each sampling day, the observer (HG) remained in the selected focal zone,

recorded all elephant visits, and noted down the times of arrival and departure of elephant groups

belonging to different clans, group sizes, group compositions, and the identities of all the

individuals based on natural physical characteristics (see Vidya *et al.*, 2014).

Since the grassland had complete visibility, we used focal group sampling (see Altmann, 1974) to

record agonistic interactions between adult females ($\geq$10 years old; Nandini *et al.*, 2018),

henceforth, referred to simply as females. Due to fission-fusion dynamics, groups (parties) of

females from the same or different clans can potentially fight with one another at the group level.

However, since we very rarely saw group-level agonistic interactions between different groups

belonging to the same clan, between-group agonism in this paper refers to agonism between groups

belonging to different clans. The within-clan data came from 17 unique clans and between-clan

agonism was recorded from 39 unique between-clan combinations. We recorded focal

observations (details in SI 3) using a Sony HDR-XR100E video camera. We also noted down the

nearest plot-cluster for each focal group observation (unless there was none within 100 m) to

examine the relationship between grass abundance/distribution and agonism.

Video scoring

We used the video recordings to score agonistic interactions (such as displacements, supplants,

pushing, raising head, etc.; see SI 3 Table S1) between females. We classified focal observations

of individual-level agonistic interactions between females into within-clan and between-clan

agonism. As mentioned above, between-clan interactions could be examined at the level of the

individual females participating, as well as at the level of the entire groups present, the latter

referred to as clan-level interactions/encounters. We recorded the time of the individual

interactions, and the identities of the initiator, recipient, winner, and loser, in within-clan and

between-clan agonistic interactions between females. Additionally, in the case of between-clan

interactions, which could comprise multiple individual-level agonistic interactions, we also

recorded the start and end times of the between-clan encounters, clan identities of the competing

groups, and clan-level outcome of the entire between-clan encounter (a clear resolution consisted

of one group winning by completely displacing the other) (see SI 3 for more details).

Data analysis

1. Grass distribution

We compared the grass biomass from the plot-clusters (average of five 1 m × 1 m quadrats each)

in the grassland habitat during each sampling year (*N*=95 and 96 plot-clusters in 2015 and 2016,

respectively) with those from plots in the forest habitat (*N*=40 plots, averages of three 1 m × 1 m

quadrats per plot) using Welch's test (Welch, 1938; see Fagerland and Sandvik, 2009) since

variances were unequal. Data from the forest had been collected at the end of the wet season in

2013. Although forests and grassland could not be sampled simultaneously due to logistics, our

observations of drying up of grass in forests suggested that any positive difference in the grass

biomass between grassland and forest from the compared datasets would be even larger if the

forest data were from the dry season. To support this assumption, we also compared the dry season

grass biomass sampled from Nagarahole forest and Kabini grassland during February-April 2022

as part of a different ongoing study (SI 4 text). Further, we compared our grass biomass data from

the grassland with that previously estimated from another adjacent forest in a previous study

(Bandipur National Park; Devidas and Puyravaud, 1995, SI 4 text).

We used three measures of grass abundance – biomass, cover, and average height to examine the

distribution of grass abundance in the grassland habitat. We averaged values of each grass

abundance variable over the five 1-m^2 quadrats of each plot-cluster to obtain within-plot-cluster

biomass/cover/height (local abundance). We used fixed-effects ANOVA (*anova()* function in *stats*

package in R, R Core Team 2018) to test the effects of year (while expected to have random effects,

this was a control fixed factor in the analysis as there were only two levels), month, zone, and their

two-way and three-way interactions on within-plot-cluster grass abundance (biomass, cover, and

height). As the five constituent quadrats of each plot cluster were chosen randomly each month

(and the exact same quadrats were not used across months), month was not a repeated-measure.

Data were missing for one data point (out of 192 plot-clusters), and we used the mean of the

neighbouring three plot-clusters for this point in order to obtain a balanced design. We calculated

η^2 (percentage variance explained) to quantify effect size for each term (SS_{effect}/SS_{total}, see Fritz *et*

al., 2012).

Apart from local abundance, we also calculated the within-plot-cluster (using the five quadrats)

coefficient of variation (CV) in grass biomass, cover, and height, as measures of local variability.

Local abundance and variability would be relevant to within-clan feeding competition (see sections

below). We replaced missing data (as one quadrat each from 4 of the 192 plot-clusters, comprising

960 quadrats in all, could not be sampled due to logistical problems) with the average abundance

values of the other quadrats in the plot-cluster. We also averaged grass abundances from all the

four plot-clusters within each zone to obtain within-zone grass abundance, and calculated the

within-zone (across the four plot-clusters) CV of grass abundance. Within-zone abundance and

variability were expected to influence between-clan contest.

2. Rates of individual-level agonism among females within and between clans

Since we wanted to compare the rates of individual-level agonism within and between clans, we

focused on the occurrence of agonistic contests and not their outcomes. We used the rate of

agonistic interactions between females (individual-level agonism, see below), calculated from

focal group observations during foraging, as a measure of contest competition (e.g., Koenig *et al.*,

2013). We considered agonistic interactions involving the same female dyad (within or between

clans) to be independent of each other if the interactions were at least 15 minutes apart (based on

the durations of agonistic interactions, see SI 3 Fig. S6), and non-independent otherwise.

Subsequent within-clan focal observations on the same group were also independent only if there

was a gap of at least 15 minutes. Similarly, we used a 2.5-hour cut-off to define independent

between-clan encounters (based on the durations of such encounters, see SI 3 Fig. S6), and thus

focal group observations on between-clan agonism.

We used only the independent interactions from each independent focal session (either within or

between clans) when the primary group activity was foraging to calculate the rate of individual-

level agonism, as the total agonism experienced per female per hour (see SI 3 Table S2, Fig. S7).

We included agonistic interactions both initiated and received to calculate rates of agonism, thus

reflecting the agonism-related interruptions to feeding faced by an average female (or interference

experienced per capita) in the locality of the focal group observation. Total agonism per female

per hour would reflect contest or interference competition (see Wheeler *et al.*, 2013; Koenig *et al.*,

2013). We used fixed-effects models (*lm* function in *lme4* package in R, Bates *et al.*, 2011) to

investigate the fixed effects of the type of agonism (between-clan or within-clan) and clan identity

on the rate of individual-level agonism. Since both between-clan and within-clan agonism were

not observed in several clans, we initially used data only from the five common clans (at least five

focal observations each of within- and between-clan agonism) for this analysis. Since clan identity

explained very little variation in the rates of agonism, we subsequently included focal group

observations from all clans when foraging was the primary group activity, and compared within-

and between-clan agonism using a fixed-effects model. We also regressed the rate of individual-

level agonism (within and between clans separately) against the number of female competitors

(group size for within-clan, and sum of group sizes for between-clan encounters), and tested for

differences in slopes ($t = (\beta_1 - \beta_2)/\sqrt{s_{\beta 1}^2 + s_{\beta 2}^2}$, where β_1 and β_2 are the slopes and $s_{\beta 1}$ and $s_{\beta 2}$

are the standard errors of β_1 and β_2, respectively) to examine whether the rates of individual-level

agonism within and between clans were differently influenced by the number of female

competitors.

We also tested for the difference between within-clan and between-clan agonism in the ratio of

non-independent to independent interactions (NI/I ratio) based on focal observations that had at

least one agonistic event, using a fixed-effects model that also included clan identity as a fixed

effect. Higher NI/I ratios represent greater engagement between the competitors and thus reflect

greater intensities of interference. To quantify effect sizes, we obtained multiple R^2 from the

summary of *lm* function or calculated η^2 (see Fritz *et al.*, 2012) from the SS values obtained using

the *anova()* function.

3. Ecological correlates of within-clan agonism

To examine what explained within-clan agonism, we included only those independent focal group

observations that were within 100 m of the centre of a plot-cluster, from which we had data on

grass abundance and variability. We used linear fixed-effects models (*lm* function in *lme4* package

in R) to test how the within-clan rate of individual-level agonism (total agonism per female per

hour) was influenced by within-plot-cluster biomass, cover, and height, within-plot-cluster CV in

biomass, cover, and height (all from the nearest plot-cluster), female group size (party size), month,

year, and zone (all fixed effects). We selected the best-subset model based on AICc (*dredge*

function in *MuMIN* package, Barton and Barton, 2012). We subsequently also performed a piece-

wise regression of the within-clan rate of agonism (total agonism per female per hour) on female

group size so that a non-linear relationship, if any, could be visually detected.

4. Ecological correlates of individual-level between-clan agonism

We used independent focal observations of between-clan encounters during foraging, which also

occurred within 100 m of the centre of a plot-cluster, to find out what explained the between-clan

rate of individual-level agonism (total agonism per female per hour). We used linear fixed-effects

models (*lm* function in *lme4* package in R) to examine the effects of within-plot-cluster grass

abundance (biomass, cover, and height), within-zone grass abundance, within-zone grass

variability (CV), month, year, zone, and the sum of and difference in female group sizes of the two

groups (all fixed effects). We chose the best-subset model based on AICc as described in the

previous section.

5. Ecological correlates of clan-level between-clan agonistic encounters

We divided each sampling day in a focal zone into four 2.5-hour intervals (08:30-11:00, 11:00-

13:30, 13:30-16:00, 16:00-18:30; SI 3) and counted the number of clans present and number of

between-clan agonistic encounters (at the clan-level) occurring within each 2.5-hour interval. We

used a generalized linear model (GLM), with the number of between-clan agonistic encounters in

each 2.5-hour interval when at least two clans were observed in the zone as the response variable

(see SI 3). We considered the number of clans (log) and area of zone (log) as offset terms in the

GLM, making it equivalent to a GLM of the rate of agonistic between-clan encounters (encounters

per 2.5 hours) with respect to these offset variables. We used a Poisson link to address the non-

normal error structure of count data. As predictor variables, we included the number of clans (since

it could influence the rate of between-clan encounters, as in gas models of encounters based on

group density; Waser, 1976), number of females in the zone (to examine the effect of local density

as hypothesized in van Schaik 1989), CV in grass biomass, and grass biomass. We used this GLM

rather than the best-subsets approach used in the previous sections because of non-normality of

between-clan agonistic encounter data. The number of clans and zone area had to be included as

offset variables in order to examine the equivalent of the rate of between-clan encounters per clan,

corrected for area.

To analyse the durations of clan-level between-clan agonistic encounters, we included only those

clan-level encounters for which complete durations were known. We used fixed-effects models

with within-plot-cluster grass abundance, within-zone grass abundance and variability (CV)

variables, sum of and difference in female group sizes (of the two competing clans), and selected

best-subsets to find out which of these could explain the duration of between-clan agonistic

encounters. The sum of group sizes was included in the analysis to find out whether the presence

of more females prolonged the duration of between-clan encounters, and difference in group sizes

was included to find out whether between-clan encounters lasted longer when the competing

groups were of similar strength. We did not use month, zone, or year because the sample sizes of

some of their levels were zero or very small. We found that two grass abundance variables in the

best model showed effects in opposite directions (see Results). Therefore, we partitioned the

variances using the ANOVA tables obtained from *anova()* function and obtained η^2 to infer how

strong these effects were.

We used a best subsets approach to model agonism when possible since most analyses had multiple

independent variables, which could lead to overfitting the whole model. Although we drew

inferences largely based on the best model obtained from AICc-based model-selection, we also

provide *P* values for the selected best model for reference. We used Statistica 7.0 (StatSoft Inc.,

2004) and *ggplot* in R to make graphs.

Results

1. Grass abundance and distribution

1a. *Grassland versus forest habitat*

Grass biomass in the Kabini grassland habitat in 2015 (mean=704.78 g/m^2, 95% CI: 646.04—

763.51, *N*=95 plot-clusters) and 2016 (mean=583.46 g/m^2, 95% CI: 546.58—620.34, *N*=96 plot-

clusters) dry seasons were nearly or over three times greater than that in the forest habitat sampled

at the end of the wet season (mean=202.24 g/m^2, 95% CI: 165.22—239.26, *N*=40 plots) (2015:

Welch's *U*=14.187, *df*=132.95, *P*<0.001; 2016: Welch's *U*=14.300, *df*=110.24, *P*<0.001; Fig. 1d).

In the dry season of 2022 also, grass was more abundant in the grassland (mean=762.135 g/m^2,

SD=302.737 g/m^2, N=24) than Nagarahole forest (mean=128.625 g/m^2, SD=143.673 g/m^2, N=25)

(Welch's U=9.296, df=32.56, P<0.001). Grass biomass in the Kabini grassland was also greater

than that reported previously from the adjacent Bandipur National Park (Devidas and Puyravaud,

1995; SI 4 text and Fig. S8). Therefore, there was habitat-level patchiness of food abundance.

1b. *Grass abundance across and within focal zones in the grassland*

The fixed-effects model explained 77.7% of the variation in grass biomass, which was unevenly

distributed across space (zones) and time (Fig. 1e). Zone and year × month interaction had

moderately large effects on grass biomass, while other factors had small to moderate effects when

significant (SI 4 Table S3). Similar trends were also obtained for grass cover (SI 4 Table S4). The

model explained 78.6% of the variation in grass height, with the effect of zone being very large,

explaining 47% of the variation, while other effects were small when significant (SI 4 Table S5).

2. Rates and intensities of individual-level agonism among females within and between clans

During a total of 281 within-clan focal group observations on 17 clans (173.3 hours), we recorded

458 independent and 267 non-independent individual-level interactions. Similarly, during 97

between-clan focal group observations involving 39 between-clan combinations (70.3 hours), we

recorded 377 independent and 578 non-independent individual-level interactions between clans

(see SI 5 Fig. S10). These 97 between-clan focal group observations included 92 independent

between-clan encounters, of which 84 were agonistic between-clan encounters. Of the focal group

observations above, 180 within-clan focals (133.5 hours, containing 345 independent and 188 non-

independent interactions) and 53 between-clan focals (48.2 hours, containing 210 independent and

312 non-independent interactions; no agonism seen in 7 out of 53 between-clan encounters) were

independent observations that were at least 15 minutes long, with all the agonistic interactions

occurring within a focal zone, and with feeding as the primary group activity. The results on the

rate and intensity of individual-level agonism below are based on these focal group observations.

We first compared the rate of individual-level agonism (total agonism per female per hour) during

within-clan and between-clan agonism using data from the five most commonly observed clans

(each with ≥ 5 within- and between-clan focals; 142 within-clan and 51 between-clan focal

observations in all) and a fixed-effects model. The rate of individual-level agonism was

significantly higher between clans than within clans ($F_{1,213}=38.332$, $P<0.001$, $\eta^2=0.148$), with no

appreciable effect of clan identity ($F_{4,213}=1.937$, $P=0.106$, $\eta^2=0.030$, SI 5 Fig. S11, Table S6). The

NI/I ratio was also higher during between-clan than within-clan focals ($F_{1,159}=30.299$, $P<0.001$,

$\eta^2=0.158$, SI 5 Table S6). Similarly, when we included data from all clans, but excluded clan

identity as it did not have an effect above, the rate of individual-level agonism during between-

clan focals (mean=2.514 interactions/female/hour, 95% CI: 1.934—3.093, N=53 focal

observations) was higher than that during within-clan focals (mean=1.152

interactions/female/hour, 95% CI=0.988—1.316, N=180 focal observations; $F_{1,231}=37.569$,

$P<0.001$) (Fig. 2a, see SI 5 Table S7).

There were small but significant effects of local competitor density (sum of female group sizes

during between-clan agonism and female group size during within-clan agonism) on the rates of

individual-level agonism between clans ($R=0.371$, $R^2=0.138$, $F_{1,51}=8.146$, $P=0.006$), as well as

within clans ($R=0.192$, $R^2=0.037$, $F_{1,178}=6.794$, $P=0.010$) (Fig. 2c); although the effect of local

competitor density on the rate of agonism seemed higher in between- than within-clan agonism,

they were not statistically significant (test for differences in slopes: $t_{229}=1.201$, $P=0.231$).

3. Effects of grass abundance, grass dispersion, and group size on the rate of agonistic

interactions within clans

The best-subset model of the rate of individual-level agonism (total agonism per female per hour)

included female group size, within-plot-cluster grass height, and zone as predictor variables, but

explained only 11% (*Multiple R^2=0.111*, $F_{7,\ 166}=2.953$, $P=0.006$) of the variation (Table 1, SI 6

Table S9). There were eight other top models (ΔAICc$\leq$ 2; Table 1), all showing positive effects of

female group size, and some showing effects of zone and within-plot-cluster height (SI 6 Table

S8). We further examined the relationship between total agonism per female per hour and female

group size, by performing a piece-wise regression since the slope appeared nonlinear. We found a

small positive effect up to a group size of five (R^2=0.126, $F_{1,136}=19.651$, $P<0.001$), with the slope

not significantly differing from zero in the remaining segment (R^2=0.001, $F_{1,40}=0.041$, $P=0.841$)

(Fig. 3).

4. Effects of grass abundance, grass dispersion, and group size on the rate of individual-level

agonistic interactions between clans

The best-subset model of the rate of individual-level between-clan agonism (total agonism per

female per hour during between-clan encounters) was well explained (Multiple R^2=0.562,

$F_{10,40}=5.141$, $P<0.001$) by moderate effects of month (η^2=0.291), zone (η^2=0.139), and the sum of

group sizes of the contesting groups (η^2=0.118), while year and grass cover had negligible effects

(Table 2, SI 6 Fig. S13, Table S10-S11). There were seven other top models (ΔAICc $\leq$ 2; Table

2), all showing effects of the sum of group sizes, month and zone, and some having an effect

of within-plot-cluster grass abundance, within-zone CV in grass biomass, and year (Table 2).

5. Rate and duration of clan-level between-clan agonistic encounters

Out of 672 intervals (168 days × 4 2.5-hour intervals), two or more clans were simultaneously

present in the focal zone during 91 intervals (from 56 sampling days), creating the potential for

clan-level agonistic encounters. Agonistic between-clan encounters occurred during 30 of the 91

intervals (62 agonistic between-clan encounters seen; the 84 agonistic between-clan encounters in

section 2 above include those seen on days when entire-day sampling at a zone was not carried

out). Our response variable was the number of between-clan agonistic encounters during each of

the 91 2.5-hour intervals (see SI 3), along with the number of clans and area of zone as offset terms

in the GLM. The model including the effects of number of clans (33.8% deviance explained),

number of females (4.6% deviance explained), grass biomass and CV in grass biomass (≤1%

deviance explained by both) performed better (*AIC*=167.067) than the null model with only the

offset terms (*AIC*=213.63) (Fig. 4a, Table 3a, SI 6 Table S12).

The best-subset model of the duration of clan-level between-clan agonistic encounters explained

33% variation (*Multiple R*^{2}=0.327, $F_{3,41}$=6.647, *P*<0.001), with a positive effect of within-plot-

cluster average grass biomass (η^2=0.191, Fig. 4b), negative effect of within-zone CV of grass cover

(η^2=0.135), and a marginal negative effect of within-plot-cluster average grass cover (η^2<0.001)

(Table 3b, SI 6 Table S13-S14). Three other models were also similar (ΔAICc ≤ 2; Table 3b).

Discussion

Despite the ecological model of female social relationships (EMFSR) conceptually not being

restricted to any taxonomic group, the relationship between food distribution and contest

competition has rarely been studied in non-primate mammals. In this first test of EMFSR in Asian

elephants, we investigated the proximate ecological basis of agonistic interactions among

individually identified females in the Kabini elephant population. This population showed the

occurrence of both between-clan and within-clan agonism despite relying on a graminivorous diet,

which is usually thought of as low-quality subsistence food, widely distributed, and assumed not

to elicit frequent contests (Wrangham, 1980; Archie *et al.*, 2006). Although forest habitats have

high primary productivity and the forest understorey in the wider landscape has some tall-grass

areas, we found that the Kabini grassland had at least three times the grass biomass as the

neighbouring forests of Nagarahole and Bandipur. As expected from EMFSR's prediction of

between-group contest in high quality patches and at high population density, we observed

frequent occurrence of agonistic between-clan encounters, which was rarely seen in forest habitats

previously (Baskaran, 1998; Vidya, personal observation).

Further, consistent with the expectation that high quality resource patches would elicit stronger

between-group contest than within-group contest, we found that individual-level agonism was

more frequent during between-clan agonism than within-clan agonism. However, in contradiction

to EMFSR's prediction of a positive relationship between within-group contest and patchiness of

feeding sites, within-clan rate of agonism was not explained by local distribution of grass, possibly

because high elephant density intensified overall competition, although female group size (up to

intermediate group sizes) positively explained a small amount of variation in within-clan agonism.

Between-clan agonistic encounters, too, were more frequent when the number of clans and female

density were higher, and the duration of between-clan agonistic encounters was longer at contest

sites with greater food abundance. The key findings on the link between resource distribution and

agonistic behaviour are summarized in Fig. 5

Diet and agonism in elephants

Our finding of frequent between-clan and within-clan agonistic contests among females (see also

Nandini, 2016) in the Kabini grassland, where only grasses were available as food, questions the

utility of diet type as a predictor of contest competition (Wrangham, 1980; Isbell, 1991). This has

also been raised by observations of within-group and between-group dominance in folivorous

primates (Koenig *et al*., 1998; Harris, 2006; see also Snaith and Chapman, 2007; Wheeler *et al*.,

2013), whose diet was assumed to result in weak contest competition previously. Strong

dominance was found in African savannah elephants also (Archie *et al*., 2006; Wittemyer and

Getz, 2007). However, Wittemyer and Getz (2007) found that contests among family groups were

common at point resources such as water holes and trees, whereas we recorded frequent agonistic

contests within and between groups in the open grassland.

Individual-level agonism within and between clans

We found strong between-clan contest: individual-level between-clan agonism during foraging

was twice as frequent as within-clan agonism, and between-clan agonistic encounters were

frequent. Since increased agonism may result in the reduction of foraging time and food intake

(Janson, 1985; Vogel, 2005), the more frequent agonism between clans suggests that females

might incur greater costs during between-clan contests than the usual competition faced within

their clans. The slope of the relationship between agonism rates and the number of female

competitors seemed higher between than within clans, but was not statistically significant.

However, since over 50% of the decided between-clan agonistic encounters in Kabini resulted in

the exclusion of the losing group from the contest site (Nandini, 2016; Gautam and Vidya,

unpublished), between-clan agonism had foraging consequences, in addition to time lost from such

interactions. Further, findings on contest outcomes (discussed below) also suggest stronger

operation of between-group than within-group contest competition. Greater agonism among

females from different clans rather than within clans suggests that female Asian elephants may

effectively use agonism and tolerance as a signal of group membership, which could explain why

associations between clans are extremely rare in Kabini (see Nandini *et al.*, 2018).

Does agonism reflect contest competition?

While we have used agonism as a surrogate of contest competition, similar to other studies (e.g.,

Koenig *et al.* 2013), agonism (or higher NI/I ratio) may not always reflect contest competition, as

strictly inferred based on inequality in dominance and its foraging/energetic consequences (Janson

and van Schaik 1988, van Schaik 1989). It is possible that higher agonism reflects greater

uncertainty in the resource-holding potential of the competitors and that greater familiarity results

in lowered agonism within clans. What we have scored as agonistic interactions involve

interference in feeding resulting in a winner and a loser and, to that extent, involve instances of

contests, but not their long-term consequences. According to the EMFSR, an expected

consequence of stronger contest (if inferred from frequent agonism) in the Kabini elephant

population than in Uda Walawe population (studied by de Silva *et al.* 2017) would be strongly

expressed dominance and rank-related skew in foraging success among clan members in Kabini.

Interestingly, weak dominance expression (Nandini 2016, Nandini *et al.* in prep.) and absence of

rank-related skew in feeding success (Gautam 2020), along with our findings of greater agonism

in larger groups suggest that within-group competition may be predominantly of the scramble type,

with classic within-group contest being weak (Nandini *et al.* in prep). However, in contrast, the

advantage to larger groups and exclusion of losing groups from the contested feeding site during

between-clan encounters (Nandini 2016) conform to classical between-group contest (Koenig

2002). Thus, between-clan contest seems to be operating more strongly than within-clan contest

in the Kabini population. It would be interesting to see if these between-clan contests result in

rank-related fitness advantages over the long term. Currently, there are not enough data to perform

such analyses in Asian elephants.

Clan-level between-clan encounters and Kabini as a resource-rich patch

Although opportunities for between-group competition are high in species with extensively

overlapping home ranges, aggression during between-group encounters is thought to be rare due

to large overlaps being associated with low incentives for aggressive defence of territory and

resources (Cheney, 1987; Pisor and Surbeck, 2019) or due to avoidance of groups (Wrangham,

1980). In contrast, species wherein groups show more exclusive ranging show stronger territorial

expression during between-group encounters (Willems and van Schaik, 2015; see also Brown,

2013). Our findings of frequent between-clan agonistic encounters (85 out of 92 encounters were

agonistic) show that the Kabini grassland has an unusual competition regime. Female Asian

elephant clans have highly overlapping home ranges (Baskaran and Desai, 1996) and a previous

long-term study in the neighbouring Mudumalai forest found only a single between-clan agonistic

encounter over a few years (Baskaran, 1998; Baskaran *et al.*, 2018). Between-clan agonistic

encounters have been seen in Uda Walawe, Sri Lanka, but not as frequently (de Silva *et al.*, 2017;

Prithiviraj Fernando, *personal communication*). The strong between-clan contest competition we

find may be explained by the small Kabini grassland being a food-rich habitat compared to the

adjacent forests in Nagarahole and Bandipur (SI 4). This grassland is a result of the receding

backwaters of the Kabini reservoir, formed by the Beechanahalli Dam, built over the River Kabini

in the early 1970s. The concentration of resources (including water) makes Kabini grassland a

highly preferred small habitat patch during the dry season, as evidenced by the exceptionally high

elephant density (median ~26 females/sq.km, SI 2 Fig. S5). Thus, frequent between-clan agonism

is consistent with the expectations of strong between-group contest for high quality resource

patches (Wrangham 1980, Brown 2013) and with the findings of frequent between-group

aggression in many species with extensive home range overlap (e.g., Willems and van Schaik

2015, Garcia *et al.* 2022). We think the anthropogenic creation of new resource has led to the

unusual competition regime that we see in Kabini. It would be interesting to examine the resource

distribution in Uda Walawe, which also has a grassland around a reservoir created in the late 1960s.

Correlates of clan-level between-clan agonism

The observed strong positive effect of the number of clans on the frequency of between-clan

agonistic encounters was expected under gas models of encounter rates based on group density

(Waser, 1976), whereas the overall high frequency of between-clan contests in a habitat with high

elephant density (Fig. S5), as well as the positive effect of the number of females in zones (Fig.

S14d), are broadly consistent with van Schaik's (1989) expectation of stronger between-group

contest at high population density. However, the frequency of agonistic between-clan encounters

was not influenced by either within-zone grass biomass or variability in grass biomass even though

zones differed greatly in grass biomass. Instead, the duration of between-clan agonistic encounters,

which may reflect the intensity of contest, was positively explained by local grass biomass (as

expected when the contest location has high resource value; Smith and Parker, 1976; Wilson *et*

al., 2012; Brown, 2013; Pal *et al.*, 2018; see also Harris, 2006; Roth and Cords, 2016).

The resource holding potential (approximated by group size) of the contestants may also determine

the duration and intensity of contests (Smith and Parker, 1976), as applied to group-level contests

(for example, Roth and Cords, 2016). Data from the Kabini Elephant Project suggest that larger

groups were more likely to win between-clan contests (Nandini, 2016; Gautam and Vidya,

unpublished). Frequent between-group contest would then be expected to result in greater within-

clan tolerance of subordinates by the dominants in order to maintain large group sizes that increase

the probability of winning between-clan contests (EMFSR; Sterck *et al.*, 1997). It would be

interesting to see whether strong between-clan contest leads to greater within-clan tolerance and

socio-spatial cohesion in Kabini than in the neighbouring forests. However, we did not find any

effect of the difference in group size on the duration of between-clan contests. It would be

necessary to examine the participation of sub-adults in between-clan contests and the differential

participation of different adult females, which could affect the collective defence of resources, in

this context. It would also be desirable to follow radiocollared clans to examine the correlates of

agonism in different habitat types.

Correlates of within-clan agonism

Although within-clan agonism occurred frequently, it did not conform to EMFSR's classical

prediction of more frequent agonistic contests in areas with more heterogeneous food distribution

(local variability in grass biomass). We think that resource concentration in the small grassland

habitat and very high female density (SI 2 Fig. S5) with possible habitat saturation may have

intensified competition in Kabini to the point where local variation in food dispersion does not

greatly influence within-clan agonism. Individuals may adopt a tactic of maximising the access to

feeding sites through agonistic contests under high competition. Such high density and competition

in the unusual habitat setting created due to the construction of a reservoir reinforces concerns

about human interference affecting social systems of wild populations (see Sterck *et al.*, 1997;

Janson, 2000; see also Robbins and Robbins, 2018).

Although unlikely, we cannot rule out the possibility that the measure of grass dispersion used

resulted in the absence of a positive association between within-clan agonism and food distribution

in Kabini. The spatial extent of measurement of grass dispersion within plot-clusters (inter-quadrat

distances from random sampling) was of the same order of magnitude as elephant group spread.

However, it is possible that the scale at which feeding competition operates between individual

elephants is different. Consumer-defined measures of food dispersion such as focal tree sampling

(Vogel and Janson, 2011) are challenging to quantify in grasslands with continuously spread food,

although measures such as food-site depletion and residence time can be explored (Chapman *et*

al., 1995; Korstjens *et al.*, 2002).

The observed positive relationship between within-clan rate of agonism and female group size was

expected. Since group size represents local density of competitors for an individual, larger groups

face greater within-group competition when feeding sites are limited (Koenig and Borries, 2006),

as reported frequently in primates (for example, van Schaik *et al.*, 1983; Wittig and Boesch, 2003;

Klass and Cords, 2015; Wheeler *et al*., 2013) but not in elephants. Greater conflict in larger groups

suggests ecological constraints on large group size, as suggested previously in this population

based on observations of average group sizes not differing across clans of different sizes (Nandini

et al., 2018). Thus, although larger group size may confer feeding benefits in this habitat with a

strong between-clan contest regime, a part of such benefits would likely be offset by the greater

costs of within-clan competition. The piece-wise regression of within-clan agonism rate on group

size suggests a non-linear relationship between group size and its cost (as seen, for example, by

Grueter *et al*., 2018 in gorillas), possibly because larger groups occupy better foraging patches in

the strong between-clan competition regime. Such non-linearity in the foraging costs and benefits

associated with group size in elephants will be interesting to explore.

Fission-fusion dynamics and within-clan agonism

Fission-fusion among group members is expected to confer flexibility in grouping to balance the

costs and benefits of large social groups in response to spatiotemporal environmental variability

(Aureli *et al*., 2008; Sueur *et al*., 2011). Thus, de Silva *et al*. (2017), who found less frequent

contests and weaker dominance expression among female Asian elephants in Uda Walawe (Sri

Lanka) compared to African elephants of Samburu, argued that higher resource availability in the

more mesic habitats of Asian elephants, along with their generalist diet may facilitate more flexible

association and result in infrequent agonism. Interestingly, we found frequent within-clan agonism

despite flexibility in associations among females in the Kabini population (Nandini *et al*., 2017;

2018), which may be at least partly explained by high local density and the large group sizes due

to advantage during between-clan contest (Nandini, 2016), as discussed above. It would be

interesting to compare the causes (density, resource distribution), as well as the social and foraging

consequences (dominance structure and effect of dominance on feeding success), of these

differences between Uda Walawe and Kabini. A testable hypothesis to explain the lower agonism

found by de Silva *et al.* (2017) would be the expectation of greater resource availability in the tall-

grass habitat of Uda Walawe and greater similarity between that grassland and the neighbouring

forest, compared to the Kabini grassland and Nagarahole/Bandipur forests.

It is also important to see our findings in the light of the grassland habitats around Kabini and other

reservoirs being recently created habitats that attract high elephant numbers, thus intensifying

competition. Hence, the observed high within-clan and between-clan agonism reflect plastic

responses to such novel environments. Thus, although they may not reflect the basal or adaptive

behaviour in natural habitats, our findings are very important in the context of the current rapid

changes in natural habitats due to human activity. The creation of developmental projects such as

dams and reservoirs, even while seemingly providing resource-rich habitats to elephants and other

animals, can change the fabric of social interactions within species.

Conclusion

We found food distribution to partly explain between-clan contest but not within-clan contest in

female Asian elephants, and competitor density to increase both within-clan and between-clan

contest, in a habitat patch with high food abundance and high competition. The unusual resource

distribution at a larger spatial scale seems to have resulted in high agonism in a species with

flexible associations, highly overlapping home ranges and a broad, graminivorous diet, pointing

to similar possibilities in other species subject to such anthropogenic habitat enrichment, and to

the plasticity of such behaviour. We call for more field-based studies of the proximate ecological

basis of agonistic contests, dominance relationships, and their proximate (food/energy/time) and

reproductive consequences in other elephant populations to understand how socioecological

factors shape group-living in the Proboscidean clade and other mammals.

Acknowledgements and funding sources

This work was funded by the Council of Scientific and Industrial Research, Government of India

(No. 37(1613)/13/EMR-II and partly 37(1375)/09/EMR-II), National Geographic Society, USA

(#9378-13 and partly #8719-09), Department of Science and Technology, Government of India

(Ramanujan Fellowship to TNCV: Grant No. SR/S2/RJN-25/2007 dated 09/06/2008), and

JNCASR. JNCASR provided institutional and logistic support. HG was given a Ph.D. scholarship

by JNCASR and this work is part of HG's Ph.D. thesis. We are grateful to the office of the Principal

Chief Conservator of Forests, Karnataka Forest Department, the office of the Conservator of

Forests and Director, Rajiv Gandhi National Park, Hunsur, and other officials for field permits.

We also thank various officials and staff of Nagarahole National Park for field support. We thank

Revathe T. for help during data collection and Thanikodi for collecting grass biomass data in 2022.

We thank Amitabh Joshi for advice on ANOVAs of grass abundance and Manan Gupta for helping

with MATLAB codes to sort clan-visits data. We thank Shankar, Krishna, Ranga and Pramod for

field assistance and ensuring field safety. We thank two anonymous reviewers for their critical

comments on a previous version of this manuscript.

Conflict of Interest

We declare no potential sources of conflict of interest or competing claims about data ownership,

authorship, or funding associated with this work.

Author contributions:

HG and TNCV conceived this work, designed methodology, and wrote the manuscript. HG collected the field data and carried out data analyses with inputs from TNCV. Both authors gave final approval for submission.

Data availability statement

Data will be made available on a publicly available data repository after the manuscript is accepted for publication.

References

1. Altmann, J. (1974). Observational study of behaviour: sampling methods. *Behaviour, 49*, 227-266. https://doi.org/10.1163/156853974X00534

2. Archie, E.A., Morrison, T.A., Foley, C.A., Moss, C.J. & Alberts, S.C. (2006). Dominance rank relationships among wild female African elephants, *Loxodonta africana*. *Animal Behaviour, 71*, 117-127. https://doi.org/10.1016/j.anbehav.2005.03.023

3. Aureli, F., Schaffner, C.M., Boesch, C., Bearder, S.K., Call, J., Chapman, C.A., Connor, R., Fiore, A.D., Dunbar, R.I., Henzi, S.P. & Holekamp, K. (2008). Fission-fusion dynamics: new research frameworks. *Current Anthropology 49*, 627-654. https://doi.org/10.1086/586708

4. Barton, K. & Barton, M.K. (2012). Package 'MuMIn'. *Model selection and model averaging based on information criteria. R package version 1.1-21.*

5. Baskaran, N. (1998). *Ranging and Resource Utilization by Asian elephant (Elephas maximus*

Linnaeus) in Nilgiri Biosphere Reserve, South India. Ph.D. thesis, Bharathidasan University,

Trichy, India.

6. Baskaran, N. & Desai, A.A. (1996). Ranging behaviour of the Asian elephant (*Elephas*

maximus) in the Nilgiri Biosphere Reserve, South India. *Gajah*, *15*, 41-57.

https://www.asesg.org/PDFfiles/Gajah/15-41-Bskaran.pdf

7. Baskaran, N., Kanakasabai, R., & Desai, A.A. (2018). Ranging and spacing behaviour of Asian

elephant (*Elephas maximus* Linnaeus) in the tropical forests of southern India. In:

Sivaperuman, C. & Venkataraman, K. (eds) *Indian Hotspots*, Springer, Singapore, pp. 295-

315.

8. Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., Dai, B.,

Scheipl, F., & Grothendieck, G. (2011). Package 'lme4'. *Linear mixed-effects models using S4*

classes. R package version 1.43.6.

9. Brown, M. (2013). Food and range defence in group-living primates. *Animal Behaviour*, *85*,

807-816. https://doi.org/10.1016/j.anbehav.2013.01.027

10. Chapman, C.A., Chapman, L.J. & Wrangham, R.W. (1995). Ecological constraints on group

size: an analysis of spider monkey and chimpanzee subgroups. *Behavioural Ecology and*

Sociobiology, *36*, 59-70. https://doi.org/10.1007/BF00175729

11. Cheney, D.L. (1987). Interactions and relationships between groups. In: Smuts, B.B. *et al.*

(eds) *Primate Societies* University of Chicago Press, Chicago, USA, pp. 267-281.

12. Crook, J.H. & Gartlan, J.S. (1966). Evolution of primate societies. *Nature*, *210*, 1200-1203.

https://doi.org/10.1038/2101200a0

13. de Silva, S., Ranjeewa, A.D. & Kryazhimskiy, S. (2011). The dynamics of social networks
among female Asian elephants. *BMC Ecology*, *11*, 17. https://doi.org/10.1186/1472-6785-11-
17

14. de Silva, S., Schmid, V. & Wittemyer, G. (2017). Fission–fusion processes weaken dominance
networks of female Asian elephants in a productive habitat. *Behavioural Ecology*, *28*, 243-
252. https://doi.org/10.1093/beheco/arw153.

15. Devidas, S. & Puyravaud, J. (1995). Primary productivity of the herbaceous layer in a grazed
savannah woodland, Bandipur National Park, southern India. *Acta Oecologica* 16: 491-505.

16. Fagerland, M.W. & Sandvik, L. (2009). Performance of five two-sample location tests for
skewed distributions with unequal variances. *Contemporary Clinical Trials*, *30*, 490-
496.https://doi.org/10.1016/j.cct.2009.06.007

17. Fernando, P. & Lande, R. (2000). Molecular genetic and behavioural analysis of social
organization in the Asian elephant (*Elephas maximus*). *Behavioural Ecology and Sociobiology*,
48, 84-91. https://doi.org/10.1007/s002650000218

18. Fritz, C.O., Morris, P.E. & Richler, J.J. (2012). Effect size estimates: current use, calculations,
and interpretation. *Journal of Experimental Psychology*, *141*, 2-18.
https://doi.org/10.1037/a0024338

19. Garcia, M.G., de Guinea, M., Bshary, R. & van de Waal, E. (2022). Drivers and outcomes of
between-group conflict in vervet monkeys. *Philosophical Transactions of the Royal Society B*,
3772021014520210145. http://doi.org/10.1098/rstb.2021.0145

20. Gautam, H., Arulmalar, E., Kulkarni, M.R. & Vidya, T.N.C. (2019). NDVI is not reliable as a
surrogate of forage abundance for a large herbivore in tropical forest habitat. *Biotropica*, *51*,
443-456. https://doi.org/10.1111/btp.12651.

21. Gautam, H., Potdar, G.G. & Vidya, T.N.C. (2017). Using visual estimation of cover for rapid

assessment of graminoid abundance in forest and grassland habitats in studies of animal

foraging. Phytocoenologia, *47*, 315-327. http://dx.doi.org/10.1127/phyto/2017/0146

22. Gautam, H. (2020). Resource Availability, Within-Clan and Between-Clan Agonistic

Interactions, and Dominance Relationships amongst Female Asian Elephants in Nagarahole

National Park, Southern India. Ph.D. thesis, Jawaharlal Nehru Centre for Advanced Scientific

Research, Bengaluru, India.

23. Grueter, C.C., Robbins, A.M., Abavandimwe, D., Vecellio, V., Ndagijimana, F., Stoinski, T.S.

& Robbins, M.M. (2018). Quadratic relationships between group size and foraging efficiency

in a herbivorous primate. *Scientific Reports*, *8*, 16718. https://doi.org/10.1038/s41598-018-

35255-0

24. Harris, T.R. (2006). Between-group contest competition for food in a highly folivorous

population of black and white colobus monkeys (*Colobus guereza*). *Behavioural Ecology and

Sociobiology*, *61*, 317-329. https://doi.org/10.1007/s00265-006-0261-6.

25. Isbell, L.A. (1991). Contest and scramble competition: patterns of female aggression and

ranging   behaviour   among   primates. *Behavioural   Ecology*, *2*,   143-155.

http://doi.org/10.1093/beheco/2.2.143

26. Janson, C. (1985). Aggressive competition and individual food consumption in wild brown

capuchin monkeys (*Cebus apella*). *Behavioural Ecology and Sociobiology*, *18*, 125-138.

https://doi.org/10.1007/BF00299041

27. Janson, C.H. (2000). Primate socio-ecology: the end of a golden age. Evolutionary

Anthropology,   *9*,   73-86.   https://doi.org/10.1002/(SICI)1520-6505(2000)9:2%3C73::AID-

EVAN2%3E3.0.CO;2-X.

28. Keerthipriya, P., Nandini, S. & Vidya, T.N.C. (2021). Effects of male age and female presence

on male associations in a large, polygynous mammal in southern India: The Asian elephant.

Frontiers in Ecology and Evolution, *9*, 616666. https://doi.org/10.3389/fevo.2021.616666

29. Klass, K. & Cords, M. (2015). Agonism and dominance in female blue monkeys. *American*

Journal of Primatology, *77*, 1299-1315. https://doi.org/10.1002/ajp.22481

30. Koenig, A. (2002). Competition for resources and its behavioural consequences among female

primates. *International   Journal   of   Primatology*, 23,   759-783.

https://doi.org/10.1023/A:1015524931226

31. Koenig, A. & Borries, C. (2006). The predictive power of socioecological models: a

reconsideration of resource characteristics, agonism and dominance hierarchies. In: Hohmann,

G., Robbins, M.M. & Boesch, C. (eds), *Feeding Ecology in Apes and Other Primates:*

Ecological, Physiological and Behavioural Aspects. Cambridge University Press, New York,

USA, pp.263-284.

32. Koenig, A. & Borries, C. (2009). The lost dream of ecological determinism: time to say

goodbye?… or a white queen's proposal? *Evolutionary Anthropology*, 18,166-174.

https://doi.org/10.1002/evan.20225

33. Koenig, A., Beise, J., Chalise, M.K. & Ganzhorn, J.U. (1998). When females should contest

for food–testing hypotheses about resource density, distribution, size, and quality with

Hanuman langurs (*Presbytis entellus*). *Behavioural Ecology and Sociobiology*, *42*, 225-237.

https://doi.org/10.1007/s002650050434

34. Koenig, A., Scarry, C.J., Wheeler, B.C. & Borries, C. (2013). Variation in grouping patterns,

mating   systems   and   social   structure: what   socio-ecological   models   attempt   to

explain. *Philosophical Transactions of the Royal Society B*, *368*, 20120348.

https://doi.org/10.1098/rstb.2012.0348

35. Korstjens, A.H., Sterck, E.H. & Noë, R. (2002). How adaptive or phylogenetically inert is

primate social behaviour? A test with two sympatric colobines. *Behaviour*, *139*, 203-226.

https://doi.org/10.1163/156853902760102654.

36. Monaghan, P. & Metcalfe, N.B. (1985). Group foraging in wild brown hares: effects of

resource distribution and social status. *Animal Behaviour*, *33*, 993-999.

https://doi.org/10.1016/S0003-3472(85)80033-6

37. Nandini, S. (2016). *Social Structure, Genetic Relatedness, and Dominance Relationships in*

Female Asian elephants in Nagarahole and Bandipur National Parks, Southern India. Ph.D.

thesis, Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru, India.

38. Nandini, S., Keerthipriya, P. & Vidya, T.N.C. (2018). Group size differences may mask

underlying similarities in social structure: a comparison of female elephant societies.

Behavioural Ecology, *29*, 145-159. https://doi.org/10.1093/beheco/arx135

39. Nandini, S., Keerthipriya, P. & Vidya, T.N.C. (2017). Seasonal variation in female Asian

elephant social structure in Nagarahole-Bandipur, southern India. *Animal Behaviour*, *134*,

135-145. http://dx.doi.org/10.1016/j.anbehav.2017.10.012

40. Pal, A., Kumara, H.N., Mishra, P.S., Velankar, A.D. & Singh, M. (2018). Between-group

encounters in Nicobar long-tailed macaque (*Macaca fascicularis umbrosus*). *Ethology*

Ecology and Evolution, *30*, 582-599. https://doi.org/10.1080/03949370.2018.1459866

41. Pisor, A.C. & Surbeck, M. (2019). The evolution of intergroup tolerance in nonhuman primates

and humans. *Evolutionary Anthropology*, *28*, 210-223. https://doi.org/10.1002/evan.21793

42. Pruetz, J.D. & Isbell, L.A. (2000). Correlations of food distribution and patch size with

agonistic interactions in female vervets (*Chlorocebus aethiops*) and patas monkeys

(*Erythrocebus patas*) living in simple habitats. *Behavioural Ecology and Sociobiology*, *49*, 38-

47. http://doi.org/10.1007/s002650000272

43. R Core Team (2018). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. 2012. *URL http://www. R-project. org*.

44. Robbins, M.M. & Robbins, A.M. (2018). Variation in the social organization of gorillas: Life

history and socioecological perspectives. *Evolutionary Anthropology*, *27*, 218-233.

https://doi.org/10.1002/evan.21721

45. Roth, A.M. & Cords, M. (2016). Effects of group size and contest location on the outcome and

intensity of intergroup contests in wild blue monkeys. *Animal Behaviour*, *113*, 49-58.

https://doi.org/10.1016/j.anbehav.2015.11.011

46. Smith, J.E., Kolowski, J.M., Graham, K.E., Dawes, S.E. & Holekamp, K.E. (2008). Social and

ecological determinants of fission–fusion dynamics in the spotted hyaena. *Animal

Behaviour*, *76*, 619-636. https://doi.org/10.1016/j.anbehav.2008.05.001

47. Smith, J.M. & Parker, G.A. (1976). The logic of asymmetric contests. *Animal Behaviour*, *24*,

159-175. https://doi.org/10.1016/S0003-3472(76)80110-8

48. Snaith, T.V. & Chapman, C.A. (2007). Primate group size and interpreting socioecological

models: do folivores really play by different rules? *Evolutionary Anthropology*, *16*, 94-106.

https://doi.org/10.1002/evan.20132

49. StatSoft Inc. (2004). STATISTICA (data analysis software system). *Version 7*. StatSoft Inc.,

Tulsa, USA. www.statsoft.com.

50. Sterck, E.H., Watts, D.P. & van Schaik, C.P. (1997). The evolution of female social

relationships in nonhuman primates. *Behavioural Ecology and Sociobiology*, *41*, 291-309.

https://doi.org/10.1007/s002650050390

51. Sueur, C., King, A.J., Conradt, L., Kerth, G., Lusseau, D., Mettke-Hofmann, C., Schaffner,

C.M., Williams, L., Zinner, D. & Aureli, F. (2011). Collective decision-making and fission–

fusion   dynamics: a   conceptual   framework. *Oikos*, *120*,   1608-1617.

https://doi.org/10.1111/j.1600-0706.2011.19685.x

52. Sukumar, R. (1989). *The Asian Elephant: Ecology and Management*. Cambridge University

Press, Cambridge, UK.

53. Terborgh, J. & Janson, C.H. (1986). The socioecology of primate groups. *Annual Review of*

Ecology and Systematics, *17*, 111-136. https://doi.org/10.1146/annurev.es.17.110186.000551

54. van Schaik, C.P. (1989). The ecology of social relationships amongst female primates. In

Standen V and Foley RA (eds) *Comparative Socioecology*. Blackwell, Oxford, UK, pp.195-

218.

55. van Schaik, C.P., van Noordwijk, M.A., de Boer, R.J. & den Tonkelaar, I. (1983). The effect

of group size on time budgets and social behaviour in wild long-tailed macaques (*Macaca*

fascicularis). *Behavioural   Ecology   and   Sociobiology*, *13*,173-181.

http://doi.org/10.1007/BF00299920

56. Vidya, T.N.C., Prasad, D. & Ghosh, A. (2014). Individual identification in Asian

elephants. *Gajah*, *40*,   3-17.   https://www.asesg.org/PDFfiles/2014/Gajah%2040/40-03-

Vidya.pdf

57. Vogel, E.R. (2005). Rank differences in energy intake rates in white-faced capuchin monkeys,

Cebus capucinus: the effects of contest competition. *Behavioural Ecology and Sociobiology*,

58, 333-344. http://doi.org/10.1007/s00265-005-0960-4

58. Vogel, E.R. & Janson, C.H. (2011). Quantifying primate food distribution and abundance for

socioecological studies: an objective consumer-centered method. *International Journal of*

Primatology, *32*, 737-754. http://doi.org/10.1007/s10764-011-9498-7

59. Waser, P.M. (1976). *Cerococebus albigena*: site attachment, avoidance, and intergroup

spacing. *The American Naturalist*, *110*, 911-935.

www.journals.uchicago.edu/doi/abs/10.1086/283117

60. Welch, B.L. (1938). The significance of the difference between two means when the

population variances are unequal. *Biometrika*, *29*, 350-362.

https://doi.org/10.1093/biomet/29.3-4.350

61. Wheeler, B.C., Scarry, C.J. & Koenig, A. (2013). Rates of agonism among female primates: a

cross-taxon perspective. *Behavioural Ecology*, *24*, 1369-1380.

https://doi.org/10.1093/beheco/art076

62. Willems, E.P. & van Schaik, C.P. (2015). Collective action and the intensity of between-group

competition in nonhuman primates. *Behavioural Ecology*, *26*, 625-631.

https://doi.org/10.1093/beheco/arv001

63. Wilson, M.L., Kahlenberg, S.M., Wells, M. & Wrangham, R.W. (2012). Ecological and social

factors affect the occurrence and outcomes of intergroup encounters in chimpanzees. *Animal*

Behaviour, 83, 277-291. https://doi.org/10.1016/j.anbehav.2011.11.004

64. Wittemyer, G. & Getz, W.M. (2007). Hierarchical dominance structure and social organization

in African elephants, *Loxodonta africana*. *Animal Behaviour*, *73*, 671-681.

https://doi.org/10.1016/j.anbehav.2006.10.008

65. Wittemyer, G., Getz, W.M., Vollrath, F. & Douglas-Hamilton, I. (2007). Social dominance,

seasonal movements, and spatial segregation in African elephants: a contribution to

conservation behaviour. *Behavioural Ecology and Sociobiology*, *61*, 1919-1931.

http://doi.org/10.1007/s00265-007-0432-0

66. Wittig, R.M. & Boesch, C. (2003). Food competition and linear dominance hierarchy among

female chimpanzees of the Tai National Park. *International Journal of Primatology 24*, 847-

867. https://doi.org/10.1023/A:1024632923180

67. Wrangham, R.W. (1980). An ecological model of female-bonded primate groups. *Behaviour*,

75, 262-300. https://doi.org/10.1163/156853980X00447

Figure Legends

Figure 1. Upper panels: illustration of the sampling regime around the Kabini backwaters (inset in

a), showing b) focal zones in the grassland habitat, and c) plot-clusters within a zone. Lower

panels: Grass abundance comparison between d) the forest (Forest-N: Nagarahole National Park,

Forest-B: Bandipur National Park) and grassland habitats, and e) the different focal zones across

different months (mean ± 95% CI obtained from 20 1-m^2 quadrats from four plot-clusters).

Figure 2. a) Rate of individual-level agonism (total agonism per female per hour), b) intensity of

inter-individual agonism (number of non-independent agonistic interactions per independent

agonistic interaction) observed during between-clan and within-clan focal group observations of

all clans, and c) overlaid scatterplots (simple correlation) and slopes of rate of individual-level

agonism during within-clan (open circles, dashed line) and between-clan (filled diamonds, solid

line) focal group observations with respect to the number of female competitors (group size in the

case of within-clan and sum of group sizes in the case of between-clan agonism). Data are from

all the observed clans.

Figure 3. Piece-wise regression to show the relationship between female group size and within-

clan rate of agonism (total agonism per female per hour) for group size segments 2-5 and greater

than five. Each data point in the graph is a focal group observation. Since many data points overlap

on the graph, the median within-clan rate of agonism corresponding to each group size is also

shown.

Figure 4. Scatterplots showing the effects of a) the number of clans on the number of clan-level

between-clan agonistic encounters (per 2.5-h interval), and b) within-plot-cluster grass biomass on

the duration of clan-level between-clan agonistic encounters. Larger bubble size represents count

of multiple observations as shown in the legend.

Figure 5. Summary of the key findings on the effects of food distribution (first column) and

competitor density (second column) on within-clan (lower row) and between-clan (upper row)

agonistic contests. 1: Between-clan agonism showed mixed trends with respect to food variability,

with higher between-clan agonism in the Kabini grassland with respect to that reported earlier in

neighbouring forests, in keeping with the grassland being a high-quality habitat patch (3a), but

between-clan agonism within the Kabini grassland not varying across zones based on variability

in food abundance (3b). 2: Between-clan agonism increased with clan density as expected. 3:

Contrary to expectation, within-clan agonism did not increase with local food variability. 4:

Within-clan agonism increased with group size, as expected.

Tables

Table 1. Top fixed-effects models explaining within-clan rate of individual-level agonism per

female per hour. Variables that feature in each best-subset model are listed, followed by *AICc*,

Δ*AICc*, and model weight.

Model of within-clan rate of agonism/female/h	*df*	*AICc*	Δ*AICc*	w_i
group size + grass height (plot) + zone	9	534.602	0	0.037
group size + CV biomass (plot)	4	535.035	0.433	0.030
group size + grass height (plot) + grass biomass (plot) + zone	10	535.718	1.115	0.021
group size + biomass (plot) + zone	9	535.730	1.127	0.021
group size + grass cover (plot) + grass height (plot) + zone	10	535.934	1.331	0.019
group size	3	535.979	1.377	0.019
group size + grass height (plot) + CV grass cover (plot) + zone	10	536.037	1.434	0.018
group size + grass height (plot) + CV grass biomass (plot) + zone	10	536.080	1.477	0.018
group size + grass height (plot) + zone + year	10	536.442	1.839	0.015

974   Table 2. Top best-subset fixed-effects models explaining the rate of between-clan individual-level

975   agonism per female per hour. Variables that feature in each best-subset model are listed, followed

976   by *AICc*, *ΔAICc*, and model weight.

Model of rate of between-clan agonism/female/h	*df*	*AICc*	*ΔAICc*	w_i
sum of group sizes + cover (plot) + month + zone + year	12	207.358	0	0.022
sum of group sizes + grass height (plot) + month + zone	11	207.938	0.581	0.017
sum of group sizes + grass height (plot) + biomass (plot) + CV biomass (zone) + month + zone	13	208.427	1.069	0.013
sum of group sizes + grass height (plot) + CV biomass (zone) + month + zone	12	208.764	1.407	0.011
sum of group sizes + grass height (plot) + biomass (plot) + month + zone	12	208.794	1.436	0.011
sum of group sizes + grass height (plot) + cover (plot) + CV biomass (zone) + month + zone	13	209.086	1.728	0.009
sum of group sizes + cover (plot) + CV biomass (zone) + month + zone + year	13	209.283	1.925	0.008
sum of group sizes + biomass (plot) + month + zone + year	12	209.330	1.973	0.008

980   Table 3. a) Generalized linear fixed-effects model (with Poisson error structure) explaining the

981   number of clan-level between-clan agonistic encounters (in 2.5-hour intervals), with the number

982   of clans and area of zone included as offset variables, and b) top best-subset fixed-effects general

983   linear models explaining the duration of between-clan agonistic encounters. Variables that feature

984   in each best-subset model are listed, followed by *AICc*, $\Delta AICc$, and model weight.

a) Number of clan-level between-clan agonistic encounters		*AIC*	Null deviance (*df*)	Residual deviance (*df*)
no. of clans + no. of females + CV of grass biomass (zone) + grass biomass (zone)		167.06	138.42 (90)	83.852 (86)
b) Duration of between-clan agonistic encounters	*df*	*AICc*	$\Delta AICc$	w_i
grass biomass (plot) + grass cover (plot) + CV of grass cover (zone)	5	456.980	0	0.048
grass biomass (plot) + grass cover (plot)	4	457.594	0.614	0.035
grass biomass (plot) + grass cover (plot) + CV of grass cover (zone) + grass biomass (zone)	6	458.539	1.560	0.022

Figures

Figure 1.

Figure 2.

Figure 3

Figure 4.

Figure 5.

1001

1002