Simultaneously vocalizing Asian barbets adopt different frequencies without

coordinating temporal rhythms

Anand Krishnan[1]

1- Indian Institute of Science Education and Research (IISER) Pune, Pashan Road, Pune

411008, India. Email: anandk@iiserpune.ac.in

Running title: Vocal rhythms of Asian barbets

Keywords: Barbet, duet, frequency, sound stream segregation, vocal coordination

Abstract

Sound stream segregation is an important challenge faced by simultaneously vocalizing

animals. Chorusing birds, for instance, coordinate vocal timing to minimize overlap.

Alternatively, other birds may use frequency differences to segregate sound streams, and

vocalizing at different frequencies may enable them to remain distinct from each other. Here, I

show that conspecific Asian barbets vocalize at distinctly different peak frequencies from each

other. Additionally, they also differ in repetition rate, as measured by the inter-phrase interval.

However, conspecific individuals across species do not temporally coordinate with each other

during vocal interactions, maintaining independent and highly stereotyped individual rhythms

together with different peak frequencies. Frequency differences between individuals may

facilitate sound stream segregation when calls overlap in time. I propose that frequency

differences between conspecifics may be widespread among birds possessing stereotyped,

repetitive calls such as those found in barbets. This may enable segregation of competing

sound streams both during cooperative duets and competitive singing during territorial

interactions.

Introduction

Animal choruses represent a striking natural example of the 'cocktail party problem', where individuals must segregate relevant information from competing streams of sound, both conspecific and heterospecific (Bee and Micheyl, 2008; Greenfield, 1994). This applies both to territorially singing males, and birds singing agonistically together in duets (Hall, 2004; Hall, 2009). In both scenarios, conspecific individuals must emit distinct signals in order to communicate with each other. Vocalizing at differing frequencies (Krishnan and Tamma, 2016; Narins, 1995; Nityananda and Bee, 2011), or altering vocal timing (Brumm, 2006; Cody and Brown, 1969; Fleischer et al., 1985; Luther, 2008) may minimize masking interference from overlapping sounds. Some birds are known to simply sing together without temporal coordination, and their vocalizations drift in and out of phase with each other (Hall, 2009; Payne and Skinner, 1970). As a result, simultaneously vocalizing conspecifics may repeatedly overlap in time, depending on the differences in repetition rate. In such cases, frequency differences may assume greater importance in sound stream segregation (MacDougall-Shackleton et al., 1998). By recording animal choruses and isolating the frequencies and repetition rates of each individual, we may obtain an indication into how each individual may be able to distinguish themselves over a chorus of conspecifics.

Here, I describe the vocal strategies employed by Asian barbet (Piciformes: Megalaimidae) (Short and Horne, 2001) choruses in India. Across Asia, multiple species of sympatric barbet co-occur, each vocalizing in a species-specific frequency band (Krishnan and Tamma, 2016). In addition to these interspecific differences, multiple individuals of the same species frequently vocalize together in choruses (Ali and Ripley, 1997; Short and Horne, 2001). However, the mechanisms that barbets employ to segregate their signals from conspecifics remain poorly understood. By studying simultaneously vocalizing conspecifics of four different species, I aimed to preliminarily understand whether vocal patterns were similar or diverse across the family. My

study examined whether simultaneously vocalizing conspecific barbets of each species

segregate from each other either in vocal peak frequencies, or by coordination of vocal timing.

For this, I examined whether conspecifics vocalizing together differed in peak frequency and

repetition rates, and then examined the inter-phrase intervals between two conspecific

individuals to elucidate whether, across species, there was any evidence that barbets

coordinated their calls with respect to vocal conspecifics. Understanding the vocal strategies

employed by these non-passerine birds enables a broader understanding of the diversity of

animal strategies to minimize acoustic masking.

Materials and Methods

Recording

I recorded barbet choruses in the city of Pune in Maharashtra (Peninsular India), and the village

of Mandal in Uttarakhand (Western Himalayas) in March-April 2018, early in the breeding

season. Each site houses two species of barbet, *Psilopogon viridis/P.haemacephalus* in Pune,

and *P.virens/P.asiaticus* in Mandal (Figure 1A. For recordings, I used Sennheiser (Wedemark,

Germany) ME62 omnidirectional microphones connected to a Zoom H6 (Tokyo, Japan) recorder

sampling at 44.1KHz, making note of multiple simultaneously vocalizing conspecific individuals.

The recorder and microphones were stationary on the ground to avoid movement noise. The

overall dataset consisted of approximately seven hours of barbet chorus recordings.

Analysis

Using Raven Pro 1.5 (Cornell Laboratory of Ornithology, Ithaca, NY, USA), I digitized

vocalizations of barbets from each recording and calculated peak frequencies of each note (PF,

using an FFT window size of 2048 for higher frequency resolution of small differences) as well

as the inter-phrase interval (IPI, the time gap between the end of one phrase and the beginning

of the next, using a window size of 512 for higher resolution of small timing differences). For

each species, I digitized approximately 2500 individual phrases (1-5 notes/phrase depending on

the species). I listened to all recordings using headphones while labeling vocalizations,

differentiating conspecific individuals by the differences in relative amplitude (birds that were

further from the microphone were lower-amplitude in recordings, and distinguishable in Raven

Pro). First, I examined whether two simultaneously vocalizing conspecific individuals differed in

PF and IPI, using paired Wilcoxon signed-rank tests. Next, I measured the interquartile range or

IQR of IPI (for 108 total instances of a vocal barbet across all four species), to quantify

stereotypy in temporal rhythms. Secondly, to examine whether simultaneously vocalizing

conspecifics (36 instances across four species, an instance defined as bouts of vocalizations

from the same individual without long gaps; this was the most conservative way to identify

behavioral instances given that I could not identify individual birds in the field) were temporally

synchronous or asynchronous with each other, I calculated the time lag between the beginning

of each phrase of one individual and the closest call of the other using MATLAB (Mathworks

Inc., Natick, MA, USA). If birds were synchronous, I predicted that the distribution of time lags

across a bout should show a clear peak and low coefficient of variation (i.e. a stereotypical time

lag between individuals) (Taylor et al., 2019). However, an asynchronous bout would imply

independent rhythms from each other; the two individuals would thus drift in and out of phase,

resulting in a uniform distribution of time lags. For each instance with 5 or more measured time

lags (27 in total), I performed two-sample Kolmogorov-Smirnov tests against 100 randomly

generated uniform distributions spanning the same range of values. I scored the results as 0 if

they did not differ significantly from uniform ($P>0.05$), or 1 if they did ($P<0.05$), and measured

the percentage of total 1's (out of 100 tests) for each of the 27 instances. A score of 0 for an

instance indicated a uniform distribution of time lags consistent with independent temporal

rhythms, and a score of 1 indicated a constant time lag consistent with two birds singing in

coordination.

Results

Conspecific barbets sing at different frequencies and repetition rates

PF-IPI probability density plots revealed intraspecific variation in PF and IPI (Figure 1B, see

supplementary videos for 3D rotations). For example, in *P. haemacephalus*, two distinct peaks

were consistent with two distinct types of song, one at about 0.7KHz with an IPI at

approximately 0.5s, and a slower song at approximately 0.95KHz with an IPI over 1s (total of 27

measured instances). For *P. viridis,* I observed a similar bimodal distribution with variation

(difference between minimum and maximum) of over 130Hz in PF and over 1.3s in IPI (total of

65 measured instances). For *P.virens* and *P.asiaticus*, my data likely involved a long series of

measurements from relatively few individuals. Thus, although I digitized a similar number of

phrases in these species, these were separable into only 9 and 7 separate behavioral

instances, respectively. Although this sample size was too small to construct probability density

distributions, the measured PF and IPI exhibited high intraspecific variation (maximum-

minimum) for both species: 116Hz PF and 0.23s IPI variation for *P.virens*, and 148Hz PF and

0.04s IPI variation for *P.asiaticus* (although IPI variation was small in the latter species, there

was a consistent difference in the few cases when two individuals were vocalizing together,

even across very long bouts, as discussed in the next section). Intraspecific PF and/or IPI

variation was thus broadly consistent across all four species, suggesting that simultaneously

vocal conspecifics may adopt different frequencies and/or repetition rates from each other.

To statistically test this assertion, I compared the PFs and IPIs of simultaneously vocal barbets

using paired Wilcoxon signed-rank tests. Because the data for *P.virens* and *P.asiaticus*

consisted mainly of very long interactions across relatively few individuals (therefore, a very

large number of labeled notes, but few individual instances of simultaneously vocal barbets), I

performed this analysis only for *P. viridis* (N=22 instances of two barbets vocalizing together)

and *P.haemacephalus* (N=8). In both species, tests revealed significant (*P. viridis*: IPI: signed

rank= 253, P<0.001, PF: signed rank= 256, P<0.001, *P.haemacephalus*: IPI: signed rank= 36,

P<0.01, PF: signed rank= 36, P<0.01). This suggests that intraspecific variation in the

measured time and frequency parameters resulted from conspecific individuals adopting

different frequencies and vocal rhythms from each other. This can be seen in Figure 2A, where

the spectrograms clearly show conspecifics vocalizing at different frequencies and repetition

rates.

Simultaneously vocalizing conspecifics sing at independent rhythms without synchronization

I next sought to establish whether vocally interacting conspecific barbets adjusted their

repetition rates to maintain synchrony with their vocal neighbors. Across 36 instances of

simultaneously vocalizing conspecifics, I calculated the time lags between the calls of two

individuals (Figure 2A). For this, I used data from all four species, as there were enough

individual vocalizations digitized for the sample size of total time lag values to be sufficient for

statistical analysis. The time lags between the calls of two individuals had very high coefficients

of variation for each species (*P.haemacephalus:* 144.13%*, P.viridis:* 164.52%*, P.virens:*

182.55%*, P.asiaticus:* 197.11%), supporting a lack of vocal timing coordination (Hall, 2009).

Time lags exhibited a range of values across all 4 species, as opposed to a single value that

would be expected if two individuals coordinated vocal timing by phase-locking (Figure 2B).

When compared to 100 randomized uniform distributions (this was done for each of 27

instances, see Methods), time lags fit a uniform distribution 92% of the time on average

(average score 0.08, P>0.05, Supplementary Data, Figure 3A, also see Methods). This uniform

distribution of time lags, together with high CVs, thus supports barbets vocalizing with

independent temporal rhythms rather than synchronizing with each other. In further support of

this, each barbet maintained its individual rhythm, whether vocalizing alone or with conspecifics.

The probability density plot of IQRs for the IPI of each individual barbet (all individual behavioral

instances measured, either solo or with another barbet) peaked at 47ms (Figure 3B), indicating

<50ms jitter in vocal timing across all species (including measurement-related variation). This

analysis included every individual instance of a vocal barbet across the dataset (solo and with

other conspecifics), in spite of which I observed a sharp peak indicating a precise vocal rhythm.

This further supports the idea that barbets maintain stable, independent vocal rhythms

regardless of whether they vocalize solo or with conspecifics.

Discussion

Avian vocal interactions have received much study for the precisely coordinated vocal timing

between individuals; many of these studies have focused on duetting birds (Fortune et al., 2011;

Hall, 2009), but others have also focused on temporal asynchrony of songs to minimize overlap

(Cody and Brown, 1969; Fleischer et al., 1985). However, other birds exhibit independent

temporal rhythms with no phase-locking or coordination between simultaneously vocalizing

individuals (Hall, 2009). My data suggests that Asian barbets do not coordinate vocal timing with

respect to other vocal conspecifics, and I also find that simultaneously vocalizing individuals

tend to differ in the peak frequencies of their vocalizations. If two barbets vocalize at

independent and different rhythms, their vocalizations will drift in and out of phase with each

other (resulting in a uniform distribution of time lags between the two individuals, as opposed to

the single peak one might expect if they were coordinated with each other).

Dilger (Dilger, 1953) described simultaneous singing in *P.haemacephalus* involving both

members of a pair vocalizing at different frequencies. My dataset, consisting of broad passive

recordings of barbet choruses, presumably consists of a mix of vocal individuals, both vocal

pairs and territorial countersinging birds. This, of course, assumes that both members of a pair

sing across species as per Dilger's observations, which requires further study. Because barbets

are often difficult to locate when singing, usually high up in the canopy, and the sexes are alike,

it is extremely challenging to identify individual birds. However, at the species-level, I find that

conspecific individuals vocalize together with differences both in peak frequency and the interval

between phrases. Additionally, in no instance do I find evidence of birds changing their temporal

rhythms to minimize overlap. Instead, barbets appear to simply adopt different temporal rhythms

from each other, which may involve paying attention only to the start of another bird's bout.

Although different repetition rates may reduce temporal overlap to some extent, some

vocalizations of two individuals will still overlap in time. In this case, frequency differences

between individuals may support sound stream segregation (MacDougall-Shackleton et al.,

1998; Nityananda and Bee, 2011). Frequency differences between conspecifics are smaller

than those between heterospecifics (Krishnan and Tamma, 2016), suggesting that they serve to

differentiate individuals within each species' frequency band.

Some species of the related African barbets (Lybiidae:*Trachyphonus*) pair-duet, and certain

species appear to exhibit vocal timing coordination (Horne and Short, 1983; Thorpe, 1963).

However, duetting partners in other species may exhibit independent rhythms (Payne and

Skinner, 1970). It is thus possible that Asian barbets may sometimes coordinate their rhythms

over short time scales, although my study does not uncover evidence of this. By and large, the

evidence I present suggests that frequency differences between conspecifics are more

important than temporal mechanisms to minimize vocal overlap. Similar mechanisms may also

operate in other non-passerine birds such as the pheasant-coucal, where simultaneously

vocalizing members of a pair exhibit different frequencies (Maurer et al., 2008). Comparative

study may thus help us understand the diversity of strategies employed by birds to minimize

masking interference when singing in a chorus.

Acknowledgments

I thank Rohit Chakravarty, Zareef Khan, Baseer Baniya and Arun Varghese for assistance

during recordings, Raghav Rajan and his lab for inputs on the data, Samira Agnihotri and

Shivam Chitnis for discussions on analysis.

Funding

215 My research is funded by an INSPIRE Faculty Award from the Department of Science and

216 Technology, Government of India and an Early Career Research (ECR/2017/001527) Grant

217 from the Science and Engineering Research Board (SERB), Government of India.

References

Ali, S. and Ripley, S. D. (1997). *Handbook of the Birds of India and Pakistan, Volume 4:*

Frogmouths to Pittas. 2nd Editio. New Delhi: Oxford University Press.

Bee, M. A. and Micheyl, C. (2008). The "Cocktail Party Problem": What is it? How can it be

solved? And why should animal behaviorists study it? *J. Comp. Psychol* **122**, 235–251.

Brumm, H. (2006). Signalling through acoustic windows: Nightingales avoid interspecific

competition by short-term adjustment of song timing. *J. Comp. Physiol. A Neuroethol.*

Sensory, Neural, Behav. Physiol. **192**, 1279–1285.

Cody, M. L. and Brown, J. H. (1969). Song asynchrony in neighbouring bird species. *Nature*

222, 778–780.

Dilger, W. C. (1953). Duetting in the Crimson-breasted barbet. *Condor* **55**, 220–221.

Fleischer, R. C., Boarman, W. I. and Cody, M. L. (1985). Asynchrony of song series in the

Bewick's wren and wrentit. *Anim. Behav.* **33**, 674–676.

Fortune, E. S., Rodríguez, C., Li, D., Ball, G. F. and Coleman, M. J. (2011). Neural

mechanisms for the coordination of duet singing in wrens. *Science (80-.).* **334**, 666–670.

Greenfield, M. D. (1994). Synchronous and alternating choruses in insects and anurans:

Common mechanisms and diverse functions. *Integr. Comp. Biol.* **34**, 605–615.

Hall, M. L. (2004). A review of hypotheses for the functions of avian duetting. *Behav. Ecol.*

Sociobiol. **55**, 415–430.

Hall, M. L. (2009). Chapter 3 A Review of Vocal Duetting in Birds. *Adv. Study Behav.* **40**, 67–

121.

Horne, J. F. M. and Short, L. L. (1983). A Review of Duetting, Sociality and Speciation in

Some African Barbets (Capitonidae). *Condor* **85**, 323–332.

Krishnan, A. and Tamma, K. (2016). Divergent morphological and acoustic traits in sympatric

communities of Asian barbets. *R. Soc. Open Sci.* **3**, 160117.

Luther, D. A. (2008). Signaller: receiver coordination and the timing of communication in

Amazonian birds. *Biol. Lett.* **4**, 651–654.

MacDougall-Shackleton, S. A., Hulse, S. H., Gentner, T. Q. and White, W. (1998). Auditory

scene analysis by European starlings (Sturnus vulgaris): perceptual segregation of tone

sequences. *J. Acoust. Soc. Am.* **103**, 3581–3587.

Maurer, G., Smith, C., Süsser, M. and Magrath, R. D. (2008). Solo and duet calling in the

pheasant coucal: sex and individual call differences in a nesting cuckoo with reversed size

dimorphism. *Aust. J. Zool.* **56**, 143–149.

Narins, P. M. (1995). Frog Communication. *Sci. Am.* **273**, 78–83.

Nityananda, V. and Bee, M. A. (2011). Finding your mate at a cocktail party: Frequency

separation promotes auditory stream segregation of concurrent voices in multi-species frog

choruses. *PLoS One* **6**,.

Payne, R. B. and Skinner, N. J. (1970). Temporal patterns of duetting in african barbets. *Ibis*

(Lond. 1859). **112**, 173–183.

Short, L. . and Horne, J. F. M. (2001). *Toucans, Barbets and Honeyguides.* Oxford Univ.

Press.

Taylor, C. J., Hall, M. L., Cain, K. E. and Langmore, N. E. (2019). A superb solo, or a deviant

duet? Overlapping songs in superb fairy-wrens. *Behav. Ecol.* **30**, 1076–1086.

Thorpe, W. H. (1963). Antiphonal singing in birds as evidence for avian auditory reaction time.

Nature **197**, 774–776.

Figure legends

Figure 1: A. Spectrograms of barbet vocalizations from Peninsular India (left) and the Western

Himalayas (right). **B.** Three-dimensional probability density distributions indicating the

occurrence of different mean PF and repetition rates (IPI) for *P.haemacephalus* and *P.viridis*.

Figure 2: A. Spectrograms of two simultaneously vocalizing *P.viridis* (left) and

P.haemacephalus (right), demonstrating intraspecific differences in frequency and repetition

rate. The white bars represent the time lag between two individuals; note how it changes with

each repetition. **B.** Graphs of all measured values of time lag between two individuals for each

species of barbet. The longer the horizontal bar, the longer the time lag; the y-axis represents

the number of such measurements made per species. Values vary from approximately 100ms

to over a second, indicating a lack of temporal synchrony between individuals.

Figure 3: A. Proportion of 100 Kolmogorov-Smirnov tests in which the observed distribution of

time lags differed from a random uniform distribution (for 27 instances of two conspecific

barbets vocalizing together). In 18/27 cases, the observed distribution fit >90% to a uniform

distribution (x-axis values <0.1, indicating independent temporal rhythms without synchrony).

Almost all other instances also fit well to a uniform distribution (the lowest being a 59% fit to

uniform, x-axis value 0.41), again supporting independent vocal rhythms. **B.** Distribution of IQRs

for the IPI of each individual instance of a vocalizing barbet across the entire dataset (both solo

and simultaneous calling, N=108). The peak at 47ms indicates very low IQRs, and therefore a

highly stereotyped temporal rhythm regardless of whether the barbet is vocalizing solo or with

conspecifics.

**Time lag between simultaneously
vocalizing individuals (s)**

A

Uniformly distributed
time lags.
**INDEPENDENT
RHYTHMS**

Non-uniform
time lags.
**SYNCHRONOUS
RHYTHMS**

Frequency (number of instances where
two birds sang together)

Proportion of 100 Kolmogorov-Smirnov tests where
observed distribution of time lags differed from a
uniform distribution (P<0.05)

B

N=108

Probability density

Peak at 47 ms: High degree of stereotypy in
the interval between calls of individual barbets
regardless of whether they vocalize solo or
simultaneously

Interquartile range of inter-phrase intervals (ms)-
Lower IQR indicates more stereotyped vocal rates