Size-dependent increase in RNA Polymerase II initiation rates mediates gene expression scaling with cell size

Xi-Ming Sun[1,2]*, Anthony Bowman[3]*, Miles Priestman[1,2,3], Francois Bertaux[1,2,3, §], Amalia Martinez-Segura[1,2], Wenhao Tang[3], Dirk Dormann[1,2], Vahid Shahrezaei[3,4] & Samuel Marguerat[1,2,4]

[1] MRC London Institute of Medical Sciences (LMS), Du Cane Road, London W12 0NN, UK

[2] Institute of Clinical Sciences (ICS), Faculty of Medicine, Imperial College London, Du Cane Road, London W12 0NN, UK

[3] Department of Mathematics, Faculty of Natural Sciences, Imperial College London, London SW7 2AZ, UK

[4] Correspondence to v.shahrezaei@imperial.ac.uk or samuel.marguerat@imperial.ac.uk

* These authors contributed equally to this work.

§ Present address: Institut Pasteur, 28 rue du Docteur Roux, 75015 Paris, France.

ABSTRACT

Cell size varies during the cell cycle and in response to external stimuli. This requires the tight coordination, or "scaling", of mRNA and protein quantities with the cell volume in order to maintain biomolecules concentrations and cell density. Evidence in cell populations and single cells indicates that scaling relies on the coordination of mRNA transcription rates with cell size. Here we use a combination of single-molecule fluorescence *in situ* hybridisation (smFISH), time-lapse microscopy and mathematical modelling in single fission yeast cells to uncover the precise molecular mechanisms that control transcription rates scaling with cell size. Linear scaling of mRNA quantities is apparent in single fission yeast cells during a normal cell cycle. Transcription rates of both constitutive and regulated genes scale with cell size without evidence for transcriptional bursting. Modelling and experimental data indicate that scaling relies on the coordination of RNAPII transcription initiation rates with cell size and that RNAPII is a limiting factor. We show using real-time quantitative imaging that size increase is accompanied by a rapid concentration independent recruitment of RNAPII onto chromatin. Finally, we find that in multinucleated cells, scaling is set at the level of single nuclei and not the entire cell, making the nucleus the transcriptional scaling unit. Integrating our observations in a mechanistic model of RNAPII mediated transcription, we propose that scaling of gene expression with cell size is the consequence of competition between genes for limiting RNAPII.

INTRODUCTION

Gene expression is coordinated with cell size in order to maintain biomolecule concentrations. Understanding the mechanisms that mediate this coordination, called hereafter "scaling", is a fundamental and intriguing problem in cell biology [1,2]. Messenger RNAs (mRNA) and proteins are synthesised from the cell DNA genome, which is one of the few cellular components that do not scale with size. Because cell volume increases exponentially and mRNA half-lives are typically short, constant rates of mRNA or protein production cannot lead to gene expression scaling. Recent work, in yeast [3], animal [4,5] and plant cells [6] has shown that mRNA synthesis rates instead are coordinated globally with cell size and are a major mechanism of scaling. Conversely, mRNA degradation seems to be mostly unconnected to scaling [3,4,6], although evidence suggests that degradation rates are adjusted early after budding yeast asymmetric division [7] and when growth rate changes [8,9]. Scaling is pervasive and only few mRNAs are able to deviate from its regulation [10–12]. Interestingly, two of them have been found to participate in the control of size homeostasis [10,11].

What could be the molecular mechanism behind transcription scaling with cell size? For a gene with an active promoter mRNA numbers follow a Poisson distribution [13]. Transcription is however often discontinuous and periods of RNA synthesis or 'bursts" alternate with periods of promoter inactivity [14]. Work in single mammalian cells has shown that scaling of mRNA numbers results from a coordination of the size of the transcription bursts with cell volume and not from their frequency [4]. This is compatible with the increased RNAPII occupancy observed genome-wide in large fission yeast cell cycle mutants [3]. This also indicates that the mechanism behind scaling may not be related to activation of transcription but rather to the efficiency of an active promoter. Critically, transcription is a complex process and is regulated at many levels including, RNAPII initiation, pause/release, elongation, and termination [15–18]. Which of these processes is coordinated with cell size to mediate scaling remains unclear.

How could a complex set of molecular reactions such as transcription become more efficient as cell size increases? In an elegant experiment Padovan-Merhar and colleagues fused cells of small and large size and found that the number of mRNAs from a gene encoded in the small cell genome would increase in response to the large cell environment, however its concentration was almost halved. This suggests that scaling responds to both cell volume and DNA content [4]. This is consistent with the observation that the cell synthesis capacity is split between genome copies in diploid budding yeast cells [10]. Importantly, changes in

gene numbers, in the case of increased ploidy for instance, are associated with overall cell

size increase in many organisms [1]. This indicates that the number of genes present in a

cell is linked to its volume and number of macromolecules. This also suggests that the cell's

overall synthetic capacity could be limiting and have a determining role in setting its size

[19]. It is therefore likely that scaling depends on a limiting factor involved in transcription but

its identity and regulation with the cell volume is not known [19,20].

In this study, we measured gene expression in over 20,000 single cells of the fission yeast

Schizosaccharomyces pombe by single molecule *in situ* hybridisation (smFISH, **Tables S1-**

3) [21]. We combined these data with agent-based models of growing and dividing cells [22–

24], stochastic models of gene expression [13,25,26] and Bayesian inference [27–33] to

investigate the quantitative parameters of gene expression that mediate scaling. This

integrative approach enabled us to determine which part of the transcription process is

scaling with cell size and which molecular event connects transcription with the cell volume.

RESULTS

Gene expression scaling is a single cell attribute of constitutive and inducible gene

expression.

We first confirmed that scaling of gene expression with cell size was an attribute of single

fission yeast cells during rapid proliferation. To do this, we measured mRNA levels of 7

constitutively expressed genes in wild-type haploids (*wt*), and in conditional mutants of the

Wee1 (*wee1-50*) and Cdc25 (*cdc25-22*) cell cycle regulators by smFISH (**Figure 1A, Table**

S1). At semi-permissive temperature, mutant cells divide at smaller and larger sizes than *wt*

respectively (**Figure S1A**) [3]. Consistent with population microarray data, mean numbers of

the 7 mRNAs measured in single cells scale with the average size of the three strains

(**Figure 1B**) [3].

We then wondered whether scaling could be detected as cells elongate and progress

through a normal cell cycle. The fission yeast elongation phase is restricted to G2.

Confounding effects arising from changes in genome content during DNA replication are

therefore unlikely. Using cell length as a measure of size (**methods**), we analysed

expression of the *rpb1* mRNA as a function of single cells' sizes in *wt*, mutant and diploid

cells and observed linear scaling within each genotype across a wide range of sizes (**Figure**

1C and 1F). Linear scaling during the cell cycle was apparent for all the constitutive genes

analysed in this study (**Figure S1B-C**). This indicates that gene expression scaling is not an

artefact of mutations in the *wee1* and *cdc25* cell-cycle regulators. Moreover, the mutant data

demonstrate that gene expression scales over a dynamic range of sizes larger than that
observed in *wt* cells. Together with earlier studies in cell populations this analysis
establishes fission yeast as a powerful model system for studying scaling [3].

We next analysed the relation of scaling with ploidy using diploid cells. Diploids follow
scaling parameters (slope, intercept) similar to smaller haploids (**Figure 1C, 1F**).
Interestingly, for a given size, mRNA numbers in diploids and in *cdc25-22* haploids were
similar (**Figure S1D-F**). Moreover, when analysing 3 diploid strains bearing heterozygous
deletions of single genes we observed significant linear scaling of the corresponding mRNA
but decreased concentrations compared to both diploid and *cdc25-22* cells. This suggests
that scaling of single gene copies in fission yeast is coordinated with the cell genome content
and that the machinery behind scaling may be limiting (**Figure S1D-F**).

At the level of single genes, is scaling a property of constitutive expression or do genomes of
larger cells have a globally higher gene expression potential? To answer this question, we
analysed three mRNA that are induced during specific phases of the cell cycle (**Figure
S1G**). All three mRNA showed stronger induction levels in larger cells when comparing *wt*
with *wee1-50* and *cdc25-22* mutants (**Figure 1D, 1G**, **S1I and S1J**). This indicates that
larger cells have an increased gene expression potential in order to support scaling of
constitutive mRNA and inducible transcripts, which were not expressed in smaller cells at the
beginning of the cycle.

To investigate this further, we analysed the induction of two genes, which respond to acute
changes in external conditions. The uracil regulated gene *urg1* responds to changes in uracil
concentration and *sib1* to addition of 2,2-dipyridyl (DIP) [34,35]. Induction of the *urg1* mRNA
was heterogeneous and scaled with cell size across size mutants (**Figure 1E, 1H, and S1H**).
The *sib1* mRNA showed a much more homogeneous response which also showed clear
scaling within and across cell types (**Figure S1K**). This indicates that higher gene
expression capacity in larger cells supports scaling in response to unexpected changes in
external conditions. Taken together, these data indicate that scaling is universal and does
not depend on the mode of gene regulation. Scaling does not result from a passive
accumulation of mRNA during the cell cycle but from a change in the cell gene expression
capacity that is coordinated with cell size.

Coupling of mRNA decay rates with cell size is not a mechanism of scaling

Messenger RNA quantities are regulated at the level of transcription but also post-
transcriptionally through modulation of degradation rates [36]. Three studies have reported

that mRNA degradation rates are not regulated as a function of cell size in fission yeast,
plant and mammalian cells [3,4,6]. To confirm these observations and extend them to single
fission yeast cells, we analysed expression of 3 genes by smFISH after transcription
inhibition with Thiolutin in *wt*, *wee1-50*, and *cdc25-22* cells. Thiolutin has been shown to
inhibit transcription in *S. pombe* efficiently (**Figure S2A**) [11]. We observed mRNA half-lives
of around 30-40min for the *rpb1* and *rpb2* mRNAs consistent with previous observations
(**Figure S2B**) [37]. In *wee1-50* and *cdc25-22* mutants both mRNAs showed half-lives similar
to *wt*, consistent with an absence of scaling of mRNA degradations rates. We also measured
mRNA half-lives as a function of the cell cycle using cells binned by size. This analysis did
not show consistent positive or negative coordination of degradation kinetics with cell size
(**Figure S2B, left**). The absence of scaling of mRNA degradation rates was further
confirmed using an orthogonal promoter switch-off approach for the *rbp1*, *rbp2* and *rtp1*
genes (**Figure S2B, right**). Finally, as discussed in the next section mathematical modelling
and inference do not support scaling of degradation rate. Overall, in agreement with previous
studies, our analysis indicates that mRNA degradation is not a major mediator of scaling.

Coupling of transcription rates to cell size and not burst frequency mediates scaling.

We next explored the contribution of transcription rates to scaling using measurements of
mRNA quantities and mathematical modelling. Using this combined approach allowed us to
study dynamic transcription rates from static smFISH measurements of single cells. We
developed agent-based models that incorporate the two-state model of gene expression
inside growing and dividing cells, which are themselves described by phenomenological
models of cell growth and size control (**Figure 2A, methods**). We used an Approximate
Bayesian Computation (ABC) approach on cell size and smFISH measurements to infer
mechanism of scaling. This inference approach determines the simplest model that captures
the statistics of the experimental measurements and returns posteriors of the parameters
and model probabilities for all models (**methods**). We used 2 classes of gene expression
models that are the limiting cases of the two-state model shown in **Figure 2A**. The first class
describes transcriptional bursts explicitly, while the second assumes only non-bursty
transcription kinetics with a simple birth-death process that produces Poisson distributions
(**Figure 2A**). Each class is in turn composed of two models the first assuming constant
transcription rates during the cell cycle and the second assuming transcription rates that
increase linearly ("scale") with cell size (**methods**).

We performed model selection between these four models on the smFISH data for 7
constitutively expressed genes in the *wt*. This analysis generated two clear conclusions.

First, models assuming bursty transcription were strongly penalised and supported by small
model probabilities (**Figure 2B**, compare proportion of blue and red colours). Second,
among the Poisson models, the one assuming transcription scaling with size was preferred
for all but one mRNA (*rpb3* in **Figure 2B**, compare orange and dark red colours). From this
we conclude that transcription rates of constitutively expressed genes in fission yeast are not
bursty and scale with cell size.

We then asked whether transcription regulated by the cell cycle or by external cues followed
the same paradigm. We found that, as constitutively expressed genes, cell-cycle regulated
mRNAs do not follow a bursty transcription dynamics (**Figure 2B, middle**). This indicates
that transcription, even when regulated in defined sections of the cell cycle falls into the
Poisson regime. In terms of scaling, model selection is, as for the constitutive case, in overall
support of scaling of transcription rate with cell size (**Figure 2B, middle**). Messenger RNAs
that respond to external stimuli showed a different picture. Model selection favoured the
bursty model with support for burst size scaling. This could point to bursty transcription of
inducible genes, to the presence of strong extrinsic noise, or to a scenario where cells
respond heterogeneously to the external signals.

In order to validate the model selection analysis, we generated simulations for all constitutive
and cell-cycle regulated genes from **Figure 2B**, using models where transcription rates scale
with cell size and follow a Poisson regime. This analysis reproduced quantitatively the mean
and coefficient of variation (sd/mean, CV) of all experimental smFISH measurements, which
illustrated the non-bursty nature of transcription for these genes further (**Figure 2C**). We
then used the inferred parameters for the Poisson models to simulate gene expression of
multiple genes in single cells and compared them to experimental data where 3 different
mRNAs were measured in each cell (**methods**). This analysis suggests that most gene-to-
gene correlations in expression are explained by scaling of transcription rates with cell size
and may not require extensive additional regulation (**Figure 2D**). However, our data also
suggest that for some gene pairs (e.g. Rpb1-Rpb2), correlations can be explained better by
including some additional extrinsic variability.

We finally asked whether our modelling approach was also in support of a negligible role of
mRNA degradation in scaling. We performed further model selection and simulation
analyses on the transcription shut-off experiment from **Figure S2B** using models that
consider different possible scalings of transcription and degradation rates (**methods**). This
analysis shows that the model assuming transcriptional scaling with constant rates of mRNA
degradation with cell size is overwhelmingly chosen (**Figure 2E**) and could capture
experimental data quantitatively confirming our initial analysis (**Figure S2B**).

In summary, we show that, as in metazoans, scaling occurs through coordination of transcriptional rates with cell size in a rapidly growing unicellular organism. However, we find that transcription during the fission yeast rapid cell cycle is mainly Poissonian except during acute response to external changes where we detect signs of bursty expression. This is consistent with previous observation in the budding yeast *Saccharomyces cerevisae* [38]. Finally, we find that scaling of transcription rates explains most of the expression correlation of multiple mRNAs.

Coordination of RNAPII initiation rates with cell size as the main mechanism of scaling.

We then wondered which specific aspect of the transcription process was directly coordinated with cell size to mediate scaling. To investigate this and to confirm model predictions from **Figure 2**, we measured transcription rates experimentally. Single cells' transcription rates can be estimated from smFISH images by measuring intensities of nuclear transcription sites [38]. As a model, we used probes directed against the 5' region of the *rbp1* mRNA. This probe design provides strong sensitivity for detection of nascent transcripts (**Figure S3A, S3B**). Transcription site intensities and cell size were overall positively correlated when comparing *wt*, *wee1-50* and *cdc25-22* cells confirming modelling results and previous observation in a metazoan cell line (**Figure 3A**, **3B**, **S3C**) [4]. We then investigated the impact of increased ploidy on transcription rates scaling. For this we compared nascent site intensities of 4 mRNA in haploid cells with those of diploid strains. We used probes against the 5' end of *rpb1* as above and against three mRNA induced in specific phases of the cell cycle, since inducible expression increases the sensitivity of nascent site analysis. Intensities of individual transcription sites were similar between haploid and diploid cells (**Figure 3C**, green and yellow boxes). However, when considering the total intensity per cell, diploids showed increased rates and scaling was apparent again (**Figure 3C**, orange boxes). This indicates that the cell transcriptional capacity is limiting and distributed between the two gene copies of diploids. This was confirmed by analysis of the nascent intensity of the *rbp1* mRNA in a heterozygous deletion strain which showed no increased rates in the remaining copy (**Figure 3C**).

To investigate further the mechanism behind scaling of transcription rates, we designed an orthogonal modelling approach where transcription is modelled as RNAPII particles hopping on a gene represented by a lattice (**Figure 3D**). This approach, which is based on a totally asymmetric simple exclusion process (TASEP), has been used successfully to study transcription and translation [6,39–42]. In our model, a gene is represented by a lattice of

length L_G and non-bursty transcription is modelled using three rates: i) the transcription

initiation rate α is the rate at which RNAPII molecules enter the first site of the lattice (gene);

ii) the elongation rate β is the rate at which RNAPII molecules hop one site forward on the

lattice (gene); and iii) the termination rate γ is the rate at which the RNAPII molecule which

sits on the last site of the gene leaves the lattice and produces a full length mRNA (**Figure**

3D). We incorporated this model in the agent-based framework from **Figure 2** assuming

each rate could be linearly coupled to cell size.

By sampling the rates α, β and γ over physiological time scales estimated from previous

studies (**Methods**) we found that coupling of initiation rates with size produced the most

robust linear scaling (**Figure 3E**) and the strongest positive correlation of nascent intensities

with size (**Figure S3D**). Although a model coupling transcription elongation rates with cell

size could also generate linear scaling in some parameter regions (**Figure 3E, S3D**), these

required elongation rates to be much slower than experimental measurements either in

yeast or metazoans which report elongation rates of around 2kb/min (**Figure S3F**) [38,43–

47]. Importantly, linear scaling could only be observed in regimes with slow initiation rates

relative to elongation and termination indicating that initiation is rate limiting (**Figure S3E**).

Interestingly, this also suggests that fast, non-limiting initiation rates could be a mechanism

by which some genes escape scaling (e.g. *rpb3*; **Figure 2B**). Finally, these results were

confirmed by ABC inference analysis of nascent sites intensities data using the same

TASEP model for 3 genes in different strains which showed clear preference for the initiation

model (**Figure S3G**). This *in silico* analysis indicates that scaling of initiation rates with cell

size is the likely mechanism of scaling.

The initiation model generated two important predictions. First, the model predicts that, even

for non-bursty genes, cells in a population are not all actively transcribing at all times and the

frequency of active transcription should increase with cell size. To test this prediction, we

compared the fraction of cells with a nascent transcription site ("active cell fraction") in *wee1-*

50, *wt*, *cdc15-22* and diploid cells. As predicted by the model, a clear increase in the fraction

of transcriptionally active cells with size could be observed (**Figure 3F**). Moreover, a strong

positive correlation of the active cells fraction with cell length was apparent when calculated

in sliding windows of increasing cell numbers during the normal cell cycle (**Figure S3I**). The

second prediction of the initiation scaling model is a strong positive correlation between the

number of transcribing RNAPII and cell size (**Figure S3D**). To test this, we analysed RNAPII

occupancy across the genomes of *wt*, *wee1-50* and *cdc25-22* cells by chromatin

immunoprecipitation followed by next generation sequencing (ChIP-seq). We used three

antibodies against total RNAPII and Serine 5 phosphorylation of its carboxy terminal domain

(CTD, **methods**). Using data normalised to occupancy at histone genes (**methods**) we
could observe a significant increase in overall RNAPII occupancy with size consistent with
previous observation and confirming model predictions (**Figure 3G**) [3]. Importantly, we did
not find evidence for a redistribution of RNAPII from the 5' to the 3' of genes in larger cells
which makes regulation of scaling at the level of RNAPII pause/release unlikely (**Figure 3H,**
S3J). This profile was also reproduced quantitatively in our minimal TASEP model (**Figure**
3D, data not shown). In summary, our *in silico* and experimental data indicate that
transcriptional scaling is mediated by an increase in RNAPII initiation rates coordinated with
cell size. In addition, our modelling data together with the increase in RNAPII occupancy
observed in larger cells suggest that RNAPII could be a limiting factor for transcription as cell
size increases.

RNAPII amounts on chromatin increases with cell size.

If scaling of initiation rates is the mechanism behind scaling and RNAPII is limiting, the
amount of RNAPII complexes on the genome of single cells should increase during the cell
cycle. To test this hypothesis, we measured localisation of RNAPII in single cells by live-cell
imaging. To do this, we tagged components of the RNAPII complex with green fluorescent
protein (GFP) and imaged them during the cell cycle (**Figure 4A, S4A**). First, we observed
that the cellular concentration of Rpb1 and Rpb9, two RNAPII subunits, remained constant
during the fission yeast G2 growth phase (**Figure 4B, S4B**). This indicates that scaling of
transcription initiation is not controlled by regulation of the cellular concentration RNAPII. Our
image data show that a very large fraction of the tagged RNAPII subunits localises in the
nucleus in the DAPI-positive area (**Figure 4A**). We therefore asked whether the amount of
RNAPII on chromatin changes with cell size during the cell cycle. To assess this, we
measured fluorescence intensities of the nuclear region occupied by RNAPII subunits
(**Figure 4C, S4C**). Strikingly, the signal for both Rpb1 and Rpb9 increases steadily during
fission yeast G2 phase where most cell size increase occurs (**Figure 4C, S4C**). Importantly
this was not the case for another DNA bound protein the histone Hta2 (H2A beta), (**Figure.**
4D-E). This is consistent with the prediction of the initiation scaling model in the previous
section and indicates that RNAPII quantities are likely to be limiting for transcription. In
summary, these experiments indicate that increased initiation rates that mediate scaling are
sustained by efficient import of RNAPII in the nucleus together with rapid recruitment onto
chromatin.

The nucleus is the scaling unit

In fission yeast and other organisms, nuclear and cytoplasmic volume are intimately connected [48]. As scaling of initiation rates depends on RNAPII levels in the nucleus we wondered whether nuclear size rather than cell size itself could be the quantitative determinant of scaling.

To test this idea, we analysed nascent site intensities of *cdc11-119* mutant cells cultivated at non-permissive temperature. Under these conditions, cells elongate, undergo mitosis and nuclear division but do not divide (**Figure 5A**) [49]. Strikingly, scaling of nascent site intensities of individual nuclei with cell size was not apparent in this system ($R_{Pearson}$ = 0.06, **Figure 5B, right**). Scaling was only restored when all nascent sites present in a cell were added together ($R_{Pearson}$ = 0.46, **Figure 5B, left**). This mirrors data from multinucleated cells showing that while the overall nuclear volume scales with cell volume, the volume of individual nuclei is proportional to their surrounding cytoplasmic volume [48]. Consistent with this, the ratios of nascent site intensities between nuclei of *cdc11-119* cells were weakly correlated with their immediate cytoplasmic volumes (**Figure S5A**). This indicates that the cell "scaling unit" could be the nucleus and not the whole cell.

We next analysed scaling in conditions where the correlation between cell and nuclear size is compromised. Pom1 is a regulator of cell polarity and division which when deleted leads to increased size variability at cell birth due to cell partitioning errors [50]. We analysed expression of three mRNA by smFISH in *pom1Δ* mutant cells expressing a marker of the nuclear envelope to allow measurement of nuclear size (**Figure 5C**). As expected cell and nuclear size show smaller correlation in the *pom1Δ* mutants compared to *wt* cells (**Figure S5B**). Scaling in this system was comparable to *wt* cells with the exception that *pom1Δ* mutants showed a higher y-axis intercept when mRNA numbers are plotted as a function of cell size (**Figure S5C**). This indicates that mRNA concentrations in *pom1Δ* cells are higher in smaller cells after birth and negatively correlated with cell size (**Figure 5D**). This deviation from perfect concentration homeostasis is also observed but to a lower degree in *wt* cells (**Figure 5D**) as reported for mammalian cells [4]. Moreover, mRNA numbers divided by nuclear volume show a negative correlation with nuclear volume (**Figure S5D**). The modelling approach described in **Figure 2** which is based on transcription scaling with cell size and binomial partitioning of mRNAs based on daughter cell sizes failed to capture this behaviour (**Figure 5E, magenta line**). However, a modified model coupling transcription rates to nuclear size instead produced a good fit to the data (**Figure 5E, S5F green line**). Together, this analysis suggests that beside the nucleus being the scaling unit, nuclear size could be an important determinant of scaling.

A mechanistic model of scaling

We used the results from this study to develop a mechanistic model of scaling centred around RNAPII mediated transcription (see also [19,20]). (**Figure 5F**, **methods**). In this model the rates of RNAPII complex synthesis and maturation scale with cell size. This is consistent with our smFISH and live-cell imaging data showing that RNAPII subunits have a constant cellular concentration during the cell cycle (**Figure 1 and 4**). RNAPII is then transported to the nucleus with a rapid rate that is depleting cytosolic RNAPII (as observed in live-cell imaging data; **Figure 4)**. Once in the nucleus, RNAPII binds to DNA with a constant high affinity and transcription rates are proportional to the numbers of DNA-RNAPII complexes present on each gene at any given time consistent with scaling of initiation rates (**Figure 3**). Finally, RNAPII levels is set to be limiting in line with the initiation scaling model (**Figure 3**), the behaviour of diploid and heterozygous mutants (**Figure 3**) [10] and with the fact that the cell synthetic capacity is titrated against the number of genes in heterokaryons [4]. Moreover, this assumption fits the observation that many RNAPII subunits are limiting for growth in fission yeast [51–53]. The high affinity and the limiting amount of RNAPII ensures that the majority of RNAPII molecules are bound to DNA and increase with cell size consistent with our imaging data (**Figure 4**) and with biochemical evidence from mammalian cells [4]. Finally, DNA replication occurs close to cell division and each daughter cell inherits about half of the DNA-bound RNAPII mostly independent of its size as observed in our live cell imaging experiment (**Figure 4C, S4C**). This simple model is able to capture the different features of our data. First, it retrieved the scaling of DNA-bound RNAPII with cell size while keeping the overall cellular concentration constant as observed in **Figure 4** (**Figure S5E**). Second, it explained the scaling of mRNA numbers with cell size including the higher mRNA concentration observed in smaller cells (**Figure 5E**). As this mechanistic model relies on both cell size mediated production of RNAPII and nuclear mediated partitioning of RNAPII it captures better the negative correlation between mRNA numbers over nuclear area and cell size than a purely nuclear size mediated transcriptional scaling (**Figure S5E**, compare blue and green line). Overall, this simple mechanistic model of RNAPII transcription by integrating the experimental findings from this study, explains the origin of transcriptional scaling and its subtler features and identifies the competition between genes for the limited pool of transcriptional machinery as central to the phenomenon of transcriptional scaling.

CONCLUSIONS

We performed an extensive experimental and modelling study of gene expression scaling with cell size in single fission yeast cells. We found that scaling is a pervasive feature of

gene expression that impinges on constitutive and regulated expression. We then showed
that scaling relies on an increase in RNAPII initiation rates with cell size and a concentration
independent recruitment of a limiting RNAPII on the genome. Finally, we propose that the
nucleus is the scaling unit and that nuclear size may participate in setting scaling levels.

Our work supports a simple and robust model for the scaling of gene expression with cell
size in which the competition between promoters for a limited pool of RNAPII determines
their relative strength. Because RNAPII maintains a constant concentration as cell size
increases (as proteins do in general), the number of RNAPII complexes increase linearly
with cell size. cell size increase will not affect the relative strength of promoters, but will
cause their absolute rate of transcriptional initiation to scale linearly with cell size, exactly as
required to produce the observed gene expression scaling.

Our model assumes that the general and specific transcription factors that regulate relative
promoter strength are in excess even in small cells, and thus are not affected by cell size. To
gain a deeper mechanistic understanding of scaling, it will be important to determine whether
this assumption is met for most regulators or if some escape the rule. It could also inform
about mechanisms through which some mRNA escape regulation by scaling as our
modelling results suggest that non-limiting initiation rates can produce this behaviour [10–
12]. Another interesting question will be to determine the role of chromatin remodellers in
facilitating transcription initiation in larger cells. It is possible that a more permissive
chromatin environment in large cells synergises with increased RNAPII local concentrations
to support higher transcription rates of inducible genes.

Recent evidence suggests that, in mammals, burst size is regulated at the level of the
proximal promoter sequence, while distal enhancers are involved in setting burst frequency
[54]. Moreover, burst initiation and RNAPII pause/release but not RNAPII recruitment have
been shown to be regulated in response to biological perturbations [55]. Our model of
scaling through initiation of RNAPII transcription fits well with these data as this process is
independent of both gene activation and control of burst frequency. It is also consistent with
the observation that promoters maintain their relative expression levels in changing growth
conditions [56].

Our observation that scaling is regulated at the level of single nuclei in multinucleated cells
and may be linked to nuclear size is interesting. RNA synthesis levels have been connected
to nuclear size in other systems such as multinucleated muscle cells [57], or for the HTLV-1
mRNA [58]. It could reflect a higher availability of RNA polymerases around larger nuclei as
these tend to be surrounded by larger cytoplasmic volume (**Figure S5A**) [48,57].
Interestingly, the nucleus was also found to be an independent transcriptional unit in mature

osteoclasts [59] and in multinucleated fungi where nuclei retain local control of cell cycle
periodic transcription [60,61]. This suggests that more complex feedback and molecular
mechanisms may also be at play.

An important result from our study, which is not directly related to scaling, is the absence of
bursty transcription for most fission yeast genes tested. This is in line with previous
observations in budding yeast and plants [29,38,62]. Transcriptional bursts result in high
gene expression noise and are associated with Fano factors ($\frac{\sigma^2}{\mu}$ of mRNA numbers) greater
than 1, whereas Poissonian birth-death processes have a Fano factor = 1. Our finding that
most transcription followed a non-bursty regime relied on our modelling taking cell size and
the cell cycle into account explicitly. Without doing so, all genes in this study have been
thought to have bursty expression as Fano factors calculated on the raw counts were well
above 1 (not shown). This reiterates the importance of studying gene expression considering
potential confounding effects of morphological features such as cell size and the cell cycle
[23,24,29,63,64].

Finally, in addition to progressing our understanding of the mechanisms behind scaling, this
study provides a large quantitative dataset of gene expression and cell size measurement in
over 20000 cells in various conditions. This will support future modelling efforts aimed at
understanding regulation of gene expression.

ACKNOWLEDGEMENTS

We would like to thank Nick Rhind for stimulating discussion and his help with formulating
our model of scaling. We are grateful to Kurt Schmoller, Philipp Thomas, Marc Sturrock and
Nick Rhind for critical reading of the manuscript. We thank Snezhana Oliferenko and Jürg
Bähler for sharing strains. This research was supported by the UK Medical Research
Council, a Leverhulme Research Project Grant (RPG-2014-408). It used the computing
resources of the UK Medical Bioinformatics partnership (UK MED-BIO), which is supported
by the UK Medical Research Council (grant MR/L01632X/1) and the Imperial College High
Performance Computing Service. ChIP-seq data have been deposited in ArrayExpress,
accession number XXXXXX.

METHODS

Strains and culture conditions

The strains and their genotypes that were used in this study are listed in **Table S1.** Genetic crossing confirmed by polymerase chain reaction was used for strains generated unless otherwise specified. Strains were revived from glycerol frozen stocks on solid yeast extract agar (YE agar), or YE agar supplemented with 25 mg l^{-1} adenine, L-histidine, L-leucine, uracil, L-lysine, and L-arginine (Sigma), and with or without appropriate antibiotics for selection. YE agar plates were incubated for approximately 48 h at 32°C in a static incubator until visible large colonies could be observed. Single colonies were transferred into liquid yeast extract medium (YE), in YE supplemented as above (YES), Edinburgh minimal medium (EMM), or EMM supplemented as above (EMMS), unless otherwise indicated in figure legends, and incubated at 170 rpm in a shaking incubator. Temperature sensitive strains were grown at 32°C and shifted to 36.5°C for the time indicated in figure legends. For the induction of *sib1* expression, the strains were grown at 25°C to an optical density at 600 nm (OD_{600}) of ≈0.4 and treated with 2,2-dipyridyl (DIP; ACROS) at a final concentration of 250 μM for the time indicated in figure legends, or left untreated. For measuring mRNA decay rates, cells were grown in YE at 25°C to OD_{600} ≈0.4; cells were treated with thiolutin (AXXORA) for the time indicated at a final concentration of 15 μg/ml, or left untreated. For *urg1* induction, cells were grown in EMM supplemented with or without 0.25 mg/ml uracil for the time indicated. For transcription inhibition, log phase cultures (OD600~0.5) were treated with thiolutin (15 ug/ml) and same volume of DMSO (used for dissolve thiolutin) was added to thiolutin untreated culture. Samples were taken at 0, 25, 35, 45 mins and processed as for smFISH. For live-cell experiments, cells were grown in EMMS in syphonstats – chemostat-like devices (http://klavinslab.org/hardware.html) which maintain the turbidity of liquid cultures by diluting with fresh medium appropriately – and maintained at OD_{600} 0.4 at 32°C by frequent dilution [65].

RNA single molecule fluorescence in situ hybridisation (smFISH)

All smFISH datasets are described in **Table S2**. All the mRNA counts, nascent site intensities and cell size measurements are available in **Table S3**. smFISH samples were prepared according to a method modified from published protocols [66,67]. Briefly, cells were fixed in 4% formaldehyde and the cell wall partially digested using zymolyase. Cells were permeabilised in 70% ethanol, pre-blocked with bovine serum albumin and salmon sperm DNA, and incubated overnight with custom Stellaris oligonucleotide sets (Biosearch Technologies) labelled with CAL Fluor Red 610, Quasar 670, or Quasar 570 (probe sequences are listed in **Table S4**). Cells were mounted in ProLong™ Gold antifade

mountant with DAPI (Molecular Probes), and imaged on a Leica TCS Sp8 confocal
microscope, using a 63x oil objective (NA 1.40). Optical z sections were acquired (0.3 µm
step size) for each scan to cover the entire depth of cells. Cell boundaries were outlined
manually and single mRNA molecules were identified and counted using the FISH-quant
MATLAB package [68]. Cell area, length, and width were quantified using custom ImageJ
macros. The technical error in FISH-quant detection was estimated at 6–7% by quantifying
rpb1 mRNA foci with two sets of probes labelled with different dyes. The nascent mRNA foci
were identified and quantified using intensity information from the above-mentioned FISH
quantification where an intensity 2.5 to 3-fold above the modal intensity within the same cell
was chosen as a threshold for nascent mRNA. The quality of the identification of nascent
sites was validated manually by visualising high intensity foci in the nucleus, with an
accuracy of over 90% in all three strains (wild-type, *cdc25-22*, and *wee1-50*).

ChIP-seq

Chromatin immunoprecipitation (ChIP) assays was carried out essentially according to
published methods [69]. In brief, cells were grown in YES to an OD600 of ~ 0.8 and fixed
with formaldehyde solution (1% final) and then quenched with glycine. After washing twice
with cold PBS (phosphate-buffed saline), cells were re-suspended in lysis buffer containing
proteases inhibitors and disrupted vigorously with acid-washed glass beads 8-11 times for
20 sec in a FastPrep instrument. Samples were then sonicated in Bioruptor (at High setting
and 6 times 5 mins with 30 secs ON/30 secs OFF). Chromatin were immunoprecipitated with
antibody against Rpb1 (ab817) or Rpb1 CTD-ser 5 (ab5408 or sigma 04-1572), which were
coupled to Dynabead protein-G and protein-A and Dynabead sheep anti-mouse or rat IgG
(Invitrogene). DNA was purified from immunoprecipitated samples using MinElute Qiagen
kit. Quantification of the DNA was done using QuDye dsDNA HS assay kit and quality was
verified using Bioanalyzer.

For sequencing DNA from immunoprecipitated samples, the libraries were made using the
NEBNext ChIP-Seq Library Prep Master Mix Set for Illumina (E6240S) with the indexes
provided in NEBBext Multiplex Oligos for Illumina (Index Primers Set1,2 and 3). Negative
control DNA are those from the same chromatin extracts without going through
immunoprecipitation steps. Pools of libraries were sequenced on an Illumina HiSeq 2500
instrument at the MRC LMS genomics facility. Paired-end reads (100 nt) were generated
from two pools of 12 or 18 samples per sequencing lane. Data were processed using RTA
1.18.64, with default filter and quality settings. The reads were de-multiplexed with bcl2fastq-
1.8.4 (CASAVA, allowing 0 mismatches).

ChIP-seq analysis

A description of ChIP-seq libraries can be found in **Table S5**. Sequencing reads were
aligned to the fission yeast genome as available in PomBase in July 2019 using BWA
[70,71]. For figure 3G, RNAPII occupancy counts were extracted for each transcript using
HTseq [72] and the fission yeast annotation available in PomBase in July 2019 [71]. Data
were normalised using DESeq2 and scaled using the mean counts of fission yeast histone
genes as a scaling factor to allow comparison of global RNAPII occupancy between size
mutants [73]. Amounts of histone proteins are thought to scale with DNA content rather than
cell volume and are commonly used as normalisation factors for absolute proteomics
measurements [74]. Moreover, average synthesis rates of histones were found to remain
constant across a wide range of sizes in budding yeast (Kurt Schmoller personal
communication). For figures 3H and S3I, RNAPII immunoprecipitation data were normalised
with their respective input and average gene analysis was performed using the deeptools
analysis suite [75].

Live-cell microscopy and analysis
Strains of interest and wild-type ySBM2 were grown from single colonies in 5 ml YES before
they were transferred into syphonstats and maintained at OD_{600} 0.4 overnight in EMMS.
Prior to microscopy, ySBM2 cells were mixed at a 1:10 ratio with each strain of interest and
diluted to a final OD_{600} of 0.3 in fresh EMMS. Cells were loaded directly into a CellASIC®
ONIX Y04C-02 microfluidic plate (EMD Millipore) according to the manufacturer instructions.
Fresh EMMS was continually perfused through the growth chamber with a constant pressure
of 6.9 kPa (approximately 3 µl/h). Cells were imaged on an Olympus IX70 inverted widefield
fluorescence microscope with an environmental chamber maintained at 32°C, with a high
precision motorised XYZ stage (ASI), controlled with µManager version 1.4.22 [76]. Cells
were continually imaged at a 10 minute interval with a 40× objective (NA 0.95, UPlanSApo;
Olympus) with brightfield (30 ms exposure), GFP (250 ms exposure, emission filter Semrock
514/30 nm), and dsRed (500 ms exposure, emission filter Semrock 617/73 nm) channels
captured by a Hamamatsu Orca Flash 4.0 V2 sCMOS camera, with illumination provided by
a Lumencor Spectra X LED light source set to 20% power. For each of the four growth
chambers within the microfluidic plate, three positions were defined, each of which was
focussed using the software autofocus using the brightfield channel; since this feature was
generally inaccurate, a 3 µm Z-stack was used around the autofocus position to ensure that
at least one Z-position was in focus for each position.

Initial analysis was performed with Fiji – a derivative of ImageJ – with time-lapses assembled
if necessary, and in-focus slices for each time-point selected using a custom macro (written
by Stephen Rothery, Facility for Imaging by Light Microscopy, Imperial College London),
resulting in a 4-dimensional OME-TIFF file for each field-of-view (XYCT). Each file was

subsequently analysed using a series of custom Python scripts utilising the scikit-image,
SciPy, NumPy, and pandas packages extensively among others. The scripts permit the
semi-automated definition of cell boundaries (segmentation) within the brightfield channel,
followed by the quantification of fluorescence within cell boundaries, identification of nuclei,
further quantification within nuclear boundaries, and assignment of cells into lineages. Cell
segmentation was effected using a custom 'balloon-filling' algorithm, in which a connected
series of nodes is 'inflated' from its centre inducing an outwards force on all nodes; this
outward force is counteracted by an 'image force', which applies an opposing force inversely
proportional to the intensity of pixels neighbouring the node, this has the effect of preventing
nodes from expanding through areas of low light intensity – which generally surrounds the
cell boundary; finally each node is also affected by its direct neighbouring nodes which pull
each node sideways according to their position, ensuring smooth contours. Together, after
multiple iterations, nodes migrate from a central location until the forces equilibrate at areas
of low light intensity (generally the edge of the cell). This procedure can be performed in an
automated manner in which iterations cease when the area contained within the nodes does
not significantly change, or in a supervised manner, in which further iterations are prompted
manually via a keyboard command, with progress displayed via a graphical interface. Initial
centres of cells are defined either by manually clicking, or by the centre of the cell in the
previous frame. Nuclei are defined as areas within the cell boundary which have red
fluorescence pixel intensity values greater than 1.1 standard deviations above the mean
intensity within the whole cell boundary. Time from mitosis is defined as the number of hours
from the point at which the number of detected nuclei increases from one to two.
Fluorescence intensity is adjusted for uneven illumination according to a series of empty
fields imaged using the same settings. Background and autofluorescence is determined from
wild-type cells cultured within the same field-of-view, with the mean level of fluorescence
within these cells subtracted from measurements. Fluorescence is normalised by cell or
nuclear area by dividing total fluorescence of all pixels within their respective boundary by
the area of that boundary. Scripts are available upon request.

For live cell imaging in Figure S4, cells were imaged in ibidi microfluidic channel slides
(µslide VI 0.4, #80601) instead of a CellASIC ONIX microfluidic plate. 30 µL of cells sampled
from syphonstat cultures maintained at OD600=0.5 for at least 15 generations were loaded
into channels of a pre-warmed slide. 40 µL of pre-warmed EMMS was then added to each
reservoir of channels, followed by 10 µL of mineral oil (Sigma, M5904) to prevent
evaporation during imaging. Live-cell microscopy was performed as described above.

Image analysis was semi-automated and performed using custom interactive MATLAB
scripts available upon request. Tracks corresponding to individual cell cycles are recorded

by user clicking. Only cell cycles for which cells remained in in focus for at least the 2 hours

preceding mitosis were collected. Local background subtraction was automatically applied to

fluorescence images. Cell segmentation was performed automatically based on thresholding

of the brighfield images. Nuclear segmentation was performed automatically based on

thresholding of the uch2-mCherry fluorescence images.

Cell size measurements

We extracted both cell area and cell length measurements form the smFISH images as

proxies for cell volume. We observed that both measurements support robustly data

characteristics such as scaling of mRNA numbers and positive intercepts. We used cell

length as a proxy for cell volume throughout the manuscript as it proved to be a simpler and

more consistent measure. Importantly, as fission yeast has a cylindrical shape, its length is

directly proportional to its volume. For the nucleus, we acquired area measurements only. As

the nucleus is spherical, area and volume are not proportional. We have therefore derived

volume estimates from area measurements assuming a perfect sphere using:

$$\text{volume} = 4/3* \pi *(\text{area}/\pi)^{3/2}$$

Mathematical modelling

We use agent-based simulations of stochastic gene expression coupled to cell size in

growing and dividing cells (**Figure 2A**) [24]. We assume cells grow exponentially with a

constant growth rate from birth to division that is sampled from a truncated Gaussian

distribution with mean m_g and standard deviation σ_g. A cell that is born with birth length L_i

grows until it reaches the division length L_f. We use a phenomenological model of cell size

control that relates the final size to initial size through a noisy linear map, which captures

experimentally observed variability and correlations in cell size [77–79].

$$L_f = aL_i + b + \eta_b \qquad (1)$$

where a and b are size control parameters ($a = 0$ denotes a sizer mechanism and $a = 1$ an

adder mechanism) and η_b is a truncated Gaussian with mean zero and standard deviation

σ_b. The dividing cell of length L_f produces two daughter cells with birth sizes L'_i and $L_f - L'_i$,

where $L'_i = L_f \eta_d$ and η_d is another truncated Gaussian with mean 0.5 and standard

deviation σ_d. The biomolecules such as mRNA molecules (except for DNA) are binomially

partitioned in the daughter cells with a probability proportional to the daughter cell size

L'_i. As shown in **Figure 2A**, we simulate a fixed number of cells, so upon cell division one of

the existing cells (including one of the newly born daughter cells) is chosen randomly and

taken out of the simulation. This ensures we are simulating a constant number of cells in

time and can produce snap-shot data with the correct cell age and size distribution as

observed in the experimental data. This has been used in the simulations used in the ABC

inference (**Figure 2**). For the modelling results shown in **Figure 3 and 5**, where the results

are conditioned on cell size, we have used a simpler scheme [24], where upon cell division

only one of the daughter cells is followed modelling a single lineage (similar to the data

generated in a mother machine) [80]. The simulations are performed using a simple

algorithm that uses discretised time steps to simulate exactly the Gillespie method [81] with

time-dependent parameters [82]. The simulation code for inference is written in the Julia

programming language and the rest of simulations are performed in R. The codes are

available upon request.

Our main gene expression model is the so-called telegraph model or the two-state model

[83] where the gene can be in an 'off' or an 'on' state and transcription can only occur when

the gene is on (**Figure 2A**). If the gene is always on then this model reduces to a simple

non-bursty birth-death process with parameters transcription rate v and mRNA decay rate d.

Here the mRNA counts per cell have a Poisson distribution (in the absence of cell cycle

effects). In the limit where the duration of the promotor on-state is shorter than the mRNA life

time we have the bursty limit characterised by 3 parameters of average burst frequency k_{on},

average burst size v/k_{off} and mRNA decay rate d. In this model, birth events are

simulated as geometrically distributed increases in mRNA numbers [84]. For **Figure 2B**

model selection, we used four variants of this model including the Poisson and bursty limits

with or without transcriptional dependence to cell size. Transcriptional scaling is modelled as

linear dependence of transcription rate or burst size to cell size ($v(L) = v_0 L$, where, v_0 is a

constant and L,is the cell length). The models for the cell cycle regulated genes, assume that

there is a point in the cell cycle, where gene expression increases from a basal level to an

active level. ABC model selection in **Figure 2E** is performed on the data from the

transcription shut-off experiments of the 3 strains of *wt*, *wee1-50*, and *cdc25-22.* The 3

models included are all the Poisson limit with different scaling assumptions for transcription

and decay rates. Model one assumes transcriptional scaling with a single constant decay

rate across the 3 strains, the second model assumes also transcriptional scaling but with 3

different constant decay rates within each strain. And the third model assumes constant

transcription and decay rate that is proportional to inverse cell size across the 3 strains. The

priors used in the model selection, are wide over a physiological range. The model selection

results were not too senstivie to the choice of the priors.

In **Figure 3D** a more detailed model of transcription is illustrated, which is based on the

totally asymmetric simple exclusion process (TASEP) [6,39–41]. Here, the promoter is

assumed to be always active, i.e. we are modelling a non-bursty gene. The gene is modelled

as a lattice of size L_G. Transcription starts by initiation through binding of a RNAPII molecule

to the first site on the gene with rate α if the site is empty. Elongation is modelled as hopping

of RNAPII forward with rate β if the next site is available. Termination is modelled with

RNAPII leaving the last site on the gene with rate γ which gives rise to a fully transcribed

mRNA. In our model, we have ignored pausing, backtracking and incomplete termination. In

Figure 3E, we compare 3 variants of this model, where size scaling is through linear

coupling of initiation, elongation or termination rate to cell size. We chose $L_G = 20$ for

computational efficiency and as it is larger than the typically observed number of RNAPIIs on

the genes, which is related to nascent site intensity (**Figure 3A**). In each model, we

randomly picked the initiation time scale τ_I, elongation time scale through the whole gene

τ_E, and termination time scale τ_T between 0.001-0.1 hours that produces average mRNA

numbers of between 20-30 for a moderately expressed gene. The lower limit on the time

scale is significantly shorter than the mRNA life time and the upper limit represents very slow

steps to achieve moderate transcription, given the mRNA life time, and is also slower than

the time-scales reported in the literature [cite]. The rates are inversely proportional to the

time-scales as $\alpha = 1/\tau_I$, $\beta = L_G/\tau_E$ and $\gamma = 1/\tau_T$. Note that as β is the rate of hopping per

site, it is also proportional to the number of sites on the gene L_G. We also performed an ABC

model selection using an implementation of our TASEP model in Julia on several datasets

(**Figure S3G**). We used the same prior as discussed above.

The nuclear scaling model and the RNAPII model in **Figure 5E** rely on both cell size and

nuclear size dynamics. It is known that nuclear size scales closely with cell size [48]. There

has not been much modelling of nuclear size control in the literature. We introduce a

phenomenological and passive model of cell and nuclear size control, extending the noisy

linear map of cell size control (**Equation 1**). We assume cellular exponential growth, cell size

control and division as before. We assume nuclear size also grows exponentially and follows

its own noisy linear map:

$$L_{Nf} = a_N \, L_{Ni} + b_N + \eta_{bN}$$

Cell division time is determined when cells reach their final size (L_f). For simplicity, we

assume mitosis is taking place at cell division and the size of the newly divided daughter

cells and their nucleus is determined by $L'_i = L_f \eta_d$ and $L'_{Ni} = L_{Nf} \eta_{Nd}$, where η_d and η_{Nd} are

truncated correlated Gaussian noise with mean equal 0.5, standard deviations σ_d and σ_{Nd}

and correlation coefficient of ρ_d. We choose $\rho_d = 0.5$ based on analysis of our time-lapse

imaging data (**Figure 4**) and the rest of the parameters of our dual noisy linear map model of

cell and nuclear control were fitted on the static pom1 mutant size data using the ABC

inference.

Given, our dual noisy linear map model discussed above, in the nuclear scaling model

(**Figure 5E**), we assume transcription rate v to be linearly dependent on the nuclear size L_N

but mRNAs are partitioned upon division based on the size of the daughter cells (not nuclear

size of the daughter cells). In this model a small daughter cell is likely to inherit a nucleus of

average size, with transcription rates higher than expected from cell size, resulting in an

increase in mRNA concentration for small cells.

The RNAPII model (**Figure 5F**) provides a mechanistic RNAPII based model of

transcriptional scaling. In this hybrid deterministic and stochastic model, transcription,

translation, complex formation and maturation of RNAPII molecules are modelled as simple

cell size dependent production steps. The RNAPII is then transported to the nucleus by a

nuclear size dependent rate and it binds to DNA with high affinity with a rate that is

dependent on the concentration of DNA in the nucleus (inversely proportional to nuclear

size). In this model transcription rate of a gene is assumed to be proportional to the amount

of DNA-bound RNAPII.

In this hybrid deterministic and stochastic model, RNAPII dynamics are modelled

deterministically by a series of ODEs inside growing and dividing cells, while transcription of

mRNA is modelled stochastically. Upon cell division, we assume mRNA are partitioned

binomially according to the size of the daughter cells. The free cytosolic and nuclear RNAPII

are portioned binomially according to the size of the daughter cells and their nucleus. The

scaling in this model comes about from sequestration of RNAPII on the DNA. The model is

very robust to different model parameters and assumptions as long as the level of free

cytosolic and nuclear RNAPII is much smaller than the DNA bound RNAPII. The qualitative

model results for Rpb1 shown in **Figure 5E**, are obtained by using the parameters of the

dual noisy linear map model discussed above for the pom1 mutant, tuning RNAPII

parameters to obtain about 10% free RNAPII and linear scaling of DNA bound RNAPII, as

well as setting the transcription rate to match expression levels of Rpb1. The model without

any further tuning recovers deviation from concentration homeostasis observed at small cell

sizes, which is observed for the different genes in the WT and Pom1 mutant strains.

ABC inference and model selection

In this study we have used Approximate Bayesian Computation (ABC) for inference. When

the likelihood function is intractable, we require a tool for carrying out inference without it.

ABC is precisely such a tool. The algorithm originated in the 1980s and 90s (see e.g. [85]).

For review of more recent developments, see [15]. ABC aims to carry out Bayesian

inference without having to evaluate the likelihood function. Given data D and model with

parameter set θ, this is done by approximating the posterior distribution:

$$P(\theta|D) = \frac{P(D|\theta)P(\theta)}{P(D)} \approx P(\theta|\rho(D,D_\theta) < \varepsilon)$$

Where D_θ is a set of data generated from the model with parameters θ, sampled from the

prior $P(\theta)$, $\rho(D,D_\theta)$ is a distance measure that is defined on the set of such datasets (or their

summary statistics) and ε is a tolerance, representing the degree of approximation we are

willing to accept. The simplest ABC algorithm, that is based on sampling θ repeatedly and

rejecting the ones that produce data with larger distance than our tolerance (which is called

ABC rejection sampling [86]), is too inefficient. Much work has been carried out over the past

decade in this area, leading to a variety of different implementations with much more

favourable scaling of computation time with the dimensionality of the parameter space [86].

For the purpose of this project we will use a Sequential Monte Carlo implementation, based

on the implementations of Toni et. al [87] (ABC-SMC) and Lenormand et.al [88] (APMC). In

the ABC-SMC, one fixes the size of the posterior sample, N, and a finite sequence of

decreasing tolerances, $\{\varepsilon_t\}$, a priori. The primary differences between APMC and ABC-SMC

are firstly that the sequence of epsilons is not determined a priori; it is dynamically

determined from the previous iteration's distribution of errors until a stopping criterion

$(pacc_{min})$ is fulfilled and secondly that simulations from earlier iterations are not discarded.

ABC lends itself very naturally to model selection [86,87]. In essence, all we have to do is to

extend our priors to one extra dimension, representing different models. Formally, we require

a joint prior distribution over models and parameters, $P(M,\theta)$. We have combined the model

selection aspects of ABC-SMC implementation and adaptive aspect of APMC to obtain our

APMC with model selection algorithm.

In order to apply our APMC algorithm, we need to choose an appropriate distance $\rho(D,D_\theta)$.

As the problem at hand is stochastic in nature, we have chosen to use sum of square

differences of summary statistics of the data and simulated data in the distance measure:

$$\rho(D,D_\theta) = \sqrt{\sum_i \left(\frac{m_i(D) - m_i(D_\theta)}{sd(m_i(D))}\right)^2}$$

761 where, we used central sample moments and cross moments $m_i(D)$ of our data up to order

762 3. With our data sample sizes, moments beyond the first three are usually too noisy to be

763   useful. Also, each term in the distance measure are weighted by the bootstrap estimates of

764   standard deviation of the central moments. This rescales the terms in the sum appropriately

765   and downweighs the noisier moments, helping to prevent overfitting of the data.

    FIGURE LEGENDS

Figure 1: Gene expression scaling is a single cell attribute of constitutive and
inducible gene expression. A. Representative images of an smFISH experiment. The
rpb1, *rpb2*, and *rpb3* mRNA are labelled in *wee1-50*, *cdc25-22*, and *wt* cells. Overlay of the
three channels and DAPI staining of DNA are shown in the last column. The white scale bar
represents 5μm. **B.** $\log_2$ average copies/cell of 7 mRNAs in wee1-50 (green) and cdc25-22
(red) cells plotted against $\log_2$ average copies/cell of the same mRNAs in *wt* cells. mRNA
common names are shown on the figure. Plain line shows equality and dotted lines values 2
fold up or down. **C.** *rpb1* mRNA copies/cell plotted against cell length for *wee1-50* (green), *wt*
(blue), *cdc25-22* (red) and *wt* diploid cells (orange). **D.** *ace2* mRNA copies/cell plotted
against cell length for *wee1-50* (green), *wt* (blue), *cdc25-22* (red) and wt diploid cells
(orange). **E.** *urg1* mRNA copies/cell plotted against cell length for *wee1-50* (green), *wt* (blue),
cdc25-22 (red) cells. Cells were analysed 3h after addition of uracil to the medium. **F.** same
as C. with all the cells in grey. The solid line shows median counts in running windows
sampled from 100 bootstrap samples of the experimental data and the shaded area
represents 95% confidence intervals. Only cells above the dotted lines were considered as
expressing the mRNA and were used for the running window analysis. **G.** same as D. with
all the cells in grey. The solid line shows median counts in running windows sampled from
100 bootstrap samples of the experimental data and the shaded area represents 95%
confidence intervals. Only cells above the dotted lines were considered as expressing the
mRNA and were used for the running window analysis. **H.** same as E. with all the cells in
grey. The solid line shows median counts in running windows sampled from 100 bootstrap
samples of the experimental data and the shaded area represents 95% confidence intervals.
Only cells above the dotted lines were considered as expressing the mRNA and were used
for the running window analysis.

Figure 2: Coordination of transcription rates to cell size and not burst frequency nor
mRNA decay rates mediates scaling. A. Cartoon of the modelling strategies used in this
section (**methods**). Domino-like shapes represent cells/agents with mRNA pictured as
yellow dots. Modelling of the cell cycle uses a noisy linear map framework and population
size is kept constant by removing a random cell for every division event. The left part of the
figures represents the two states model of gene expression. The transcription rate v and
degradation rate d can be coupled to cell length (L). **B.** Bar chart showing the sum of particle
weights for four models of transcription scaling after model selection analysis. Results for

genes with three types of regulation are shown. Constitutive: constitutively expressed genes,
Periodic: cell cycle regulated genes, Regulated: genes regulated by external stimuli. For
regulated genes, times after addition of 2,2-dipyridyl (sib1) or uracil (urg1) are shown. The
four types of model used in this analysis are marked on the right. Poisson: non-bursty
transcription, Bursty: bursty transcription, Constant: constant transcription rates during the
cell cycle, Scaling: transcription rates scaling with cell length (**methods**). **C.** Simulation of
mRNA numbers during the cell-cycle using a model where transcription is scaling under a
poisson regime. Coefficients of variation (CV) are plotted as a function of mean simulated
mRNA numbers for each particle of each mRNA. Yellow circles denote the median of
simulated data for all particles of a given mRNA. Red circles denote median for all
experimental data used for parameter inference. **D.** Expression correlation between pairs of
mRNA in single cells. smFISH experimental measurements are marked by orange dots with
the 95% confidence interval shown with orange bars. 95% confidence intervals for
correlations obtained from model simulations as in C are shown with blue bars. Red bars
show the 95% confidence interval for simulations including 20% additional extrinsic noise
(**methods**). **E.** Bar chart showing the sum of particle weights for three models of mRNA
degradation after model selection analysis. Decay rate scaling: transcription constant and
degradation rates scale with cell length, Constant single: *wt*, *wee1-50*, and *cdc25-22* cells
have an identical constant degradation rate during the cell cycle with transcriptional scaling.
Constant multiple: *wt*, *wee1-50*, and *cdc25-22* cells have distinct constant degradation rates
during the cell cycle with transcriptional scaling.

Figure 3: Coordination of RNAPII initiation rates with cell size as the main mechanism
of scaling. A. *rpb1* normalised nascent sites intensities plotted against their respective cell
length for *wee1-50* (green), *wt* (blue), *cdc25-22* (red). **B.** same as A. with all the cells in grey.
The solid line shows median counts in running windows sampled from a count distribution
identical to the experimental data. Shaded area represents 95% confidence intervals
(methods). Cells with no nascent sites are excluded. Pearson correlation coefficient of
nascent site intensity and length is shown. **C.** Boxplots of normalised nascent intensities for
rpb1 and the cell cycle regulated genes *ace2*, *mid2* and *fkh2*. Intensities are compared
between haploid (green), diploid (yellow, orange), and *rpb1* heterozygote deletion cells
(purple). For diploid cells the yellow boxplot shows intensities of single nascent sites, and
orange boxplots show the total nascent intensities per cell. **D.** Cartoon of the TASEP
modelling strategy of RNAPII transcription. A gene is represented by a lattice of length L_G.
RNAPII molecules enter the lattice at a rate α (Initiation), hop through the lattice at a rate β
(Elongation) and exit the lattice at a rate γ (Termination, **methods**). **E.** Transcription linear
scaling captured by the TASEP model where rates α (Initiation), β (Elongation) or γ

(Termination) are linked to cell length. Simulations where run with each model with 500
random sets of TASEP parameters α, β and γ over times scales of 0.001 to 0.1 hours
(**methods**). Deviation from linearity is shown for parameter samples in each of the 3 model
variants. For each parameter set 1500 cells are simulated and two linear regressions are
done on the mRNA numbers vs cell length data for the smaller and larger half of simulated
cells. The deviation from linearity is estimated as the difference between the linear
regression coefficients of small vs large cells. Values close to zero indicate a linear scaling
and larger values indicate saturation of mRNA numbers at large cells. The dashed line
shows the error in estimating the slope of the linear fit. Note that the initiation model shows
the most robust linearity. **F.** Fraction of cells with a nascent site ("Transcribing fraction")
plotted as a function of mean cell length for the *rpb1* mRNA in a series of strains of different
average length. Strains names are labelled next to the data points. Red line shows linear
regression and R^2 is shown in the top left corner. **G.** Chromatin immunoprecipitation analysis
of RNAPII occupancy in wee1-50 (green), wt (blue), and cdc25-22 (red) cells. RNAP ChIP-
seq data for antibodies against Serine 5 CTD phosphorylation (left, middle) and total
Rbp1(right) are shown. RNAP occupancy values were normalised using DEseq2 and scaled
using occupancy at histone genes (**methods**). **H.** RNAPII occupancy 5'/3' ratio in wee1-50
(green), wt (blue), and cdc25-22 (red) cells. RNAP ChIP-seq data for antibodies against
Serine 5 CTD phosphorylation (left, middle) and total Rbp1(right) are shown. RNAP
occupancy values were normalised to input samples and each gene was divided into 100
bins using the deeptools package. Ratios between mean occupancy in bins 1-50 compared
to bins 51-100 are shown (**methods**). Note that ratios are not changing with size and are
consistently lower than 1.

Figure 4: Nuclear RNA Polymerase II concentration increases with cell size. A.
Confocal images of Rpb1, Rpb9, and Hta2 proteins tagged with GFP in *wt* fission yeast cells.
DAPI straining for DNA and overlay of both channels are shown. The white scale bar
represents 5µm. **B.** Widefield fluorescence data from live-cell imaging RNAPII subunit Rpb9.
Normalised intensity/cell as a proxy for total cellular protein concentration is plotted along
time relative to the mitotic phase of the cell cycle. **C.** Widefield fluorescence data from live-
cell imaging RNAPII subunit Rpb9. Total pixel intensity in the area of strong fluorescence
signal as a proxy for chromatin bound amounts is plotted along time relative to the mitotic
phase of the cell cycle. **D.** As in B. but for the histone protein Hta2. **E.** As in C. but for the
histone protein Hta2.

Figure 5: The nucleus is the scaling unit. A. Representative smFISH images for the rpb1
and rpb2 mRNA in cdc11-119 grown at non-permissive temperature. The white scale bar
represents 5µm. **B.** rpb1 normalised nascent sites intensities per nucleus (blue) or per cell

(orange) plotted as a function of each cell respective size. Red lines are regression lines
from linear models. The white scale bar represents 5µm. **C**. Representative smFISH images
for the *rpb1*, *rpb2*, and *shd1* mRNA in *wt* and *pom1* deletion mutants carrying the Cut11-
GFP nuclear marker (left column). Overlay is shown on the right. The white scale bar
represents 5µm. **D.** mRNA concentration (mRNA/cell length) for the *rpb1*, *rpb2*, and *shd1*
mRNA in *wt* (black) and *pom1* deletion mutants (red). The solid line shows median counts in
running windows sampled from a count distribution identical to the experimental data.
Shaded area represents 95% confidence intervals. **E.** *rpb1* concentrations measured by
smFISH are plotted as a function of cell length. Running average for the experimental data
(grey) or derived from model predictions assuming transcription rates scaling with cell length
(magenta), nuclear area (green), or predicted from a mechanistic model of RNAPII prediction
are shown (light blue). **F.** Cartoon representation of a mechanistic model of RNAPII
transcription (see Methods for details and parameters).

Figure S1 related to Figure 1. A. Cell length distribution for the 111 datasets described in
this study. Colours represent different genotypes. Datasets are described in **Table S2**.
Coloured lines show the interquartile range for all the *wee1-50* (green), *wt* (blue), *cdc25-22*
(red), diploid and heterozygote deletions (orange) and *pom1∆* (light blue) cells. **B.** Top:
Pooled count distribution of 7 constitutively expressed genes in *wee1-50* (green), *wt* (blue),
and *cdc15-22* (red) cells. Bottom: Median counts of 7 constitutively expressed genes as a
function of cell length in running windows sampled from a count distribution identical to the
experimental data. Shaded area represents 95% confidence intervals (**methods**). Data from
wee1-50, *wt*, *cdc15-22* cells are pooled and gene names are indicated next to their
respective data line. **C.** Deviation from linearity is plotted as a function of linearity P-values
(see Figure 3E, **methods**). Low P-values mean significant deviation from linearity. Colours
are as in panel A. P-values equal 0.05 and 0.001 are shown with dashed lines and
annotated. **D.** *rpb1* mRNA copies/cell plotted against cell length for diploid (orange), *cdc25-*
22 (red) and *rpb1* heterozygote diploid (purple). The solid line shows median counts in
running windows sampled from a count distribution identical to the experimental data.
Shaded area represents 95% confidence intervals (**methods**). **E.** *myo1* mRNA copies/cell
plotted against cell length for diploid (orange), *cdc25-22* (red) and *myo1* heterozygote diploid
(purple). The solid line shows median counts in running windows sampled from a count
distribution identical to the experimental data. Shaded area represents 95% confidence
intervals (**methods**). **F.** *pan1* mRNA copies/cell plotted against cell length for diploid
(orange), *cdc25-22* (red) and *pan1* heterozygote diploid (purple). The solid line shows
median counts in running windows sampled from a count distribution identical to the
experimental data. Shaded area represents 95% confidence intervals (**methods**). **G.**

Representative images of an smFISH experiment for cell cycle regulated genes. The *mid2*,
fkh2, and *ace2* mRNA are labelled in *wee1-50*, *cdc25-22* and *wt* cells. Overlay of the three
channels and DAPI staining of DNA is shown on the last column. The white scale bar
represents 5µm. **H.** Representative images of an smFISH experiment for a gene regulated
by external conditions (*urg1*). The *urg1*, *ace2*, and *rpb1* mRNA are labelled in *wt* cells
before, 3 hours and 12 hours after addition of uracil to the culture. Overlay of the three
channels and DAPI staining of DNA is shown on the last column. The white scale bar
represents 5µm. **I.** *mid2* mRNA copies/cell plotted against cell length for *wee1-50* (green), *wt*
(blue), cdc25-22 (red). The solid line shows median counts in running windows sampled
from a count distribution identical to the experimental data. Shaded area represents 95%
confidence intervals (**methods**). Only cells above the dotted lines were considered as
expressing the mRNA and were used for the running window analysis. **J.** *fkh2* mRNA
copies/cell plotted against cell length for *wee1-50* (green), *wt* (blue), cdc25-22 (red). The
solid line shows median counts in running windows sampled from a count distribution
identical to the experimental data. Shaded area represents 95% confidence intervals
(**methods**). Only cells above the dotted lines were considered as expressing the mRNA and
were used for the running window analysis. **K.** *sib1* mRNA copies/cell plotted against cell
length for wt cells before (light blue), 20min after (blue) or 45min (pink) after addition of 2,2-
dipyridyl (DIP). The solid line shows median counts in running windows sampled from a
counts distribution identical to the experimental data. Shaded area represents 95%
confidence intervals (**methods**).

Figure S2 related to Figure 2. A. Representative images of smFISH experiments for the
rpb1 mRNA in cells before (t = 0) and after a 45min (t = 45) treatment with 15 µg/ml of
Thiolutin. Note the number of mRNA dots decreases with time and that the transcription foci
disappear after treatment. The white scale bar represents 5µm. **B. Left:** mRNA half-lives
were measured after Thiolutin treatment in bins of increasing cell length for the *rpb1* (2
repeats) and *rpb2* mRNA in *wee1-50*, *wt*, and *cdc25-22* cells (left). **Right:** mRNA half-lives
were measured after nmt41 promoter switch-off in bins of increasing cell length for the *rpb1*,
rpb2 and *rtp1* mRNA in *wt* cells.

Figure S3 related to figure 3. A. Representative images of smFISH experiments for the
rpb1 mRNA in haploid and diploid cells. Nascent transcription sites are indicated with white
arrows. The white scale bar represents 5µm. **B.** Fraction of cells with 1, 2, or 3 nascent sites
(as in A) in haploid and diploid cells. Note the diploid cells have mostly 2 sites validating our
approach specificity and sensitivity to detect nascent transcription sites. **C.** Boxplot of
nascent sites intensities for the *rpb1* gene in *wee1-50* (green), *wt* (blue) and *cdc25-22* (red)
(related to **Figure 3A**). **D.** Pearson correlation between mRNA numbers and cell length for

the simulation data from **Figure 3E**. **E.** Histogram of ratio of initiation time scale $(1/\alpha)$ and

the sum of elongation time scale and termination time scale $(\frac{L_G}{\beta} + \frac{1}{\gamma})$ for parameter sets

resulting in linear scaling and parameter sets resulting in saturating transcription scaling

indicating that initiation rate should be limiting for linear transcriptional scaling. **F.** Elongation

time of a 2kb gene obtained from the simulations from **Figure 3E** for the models with

Initiation rates (α, green) and elongation rates (β, red) coupled with size. Only simulation that

achieved linearity are shown. Elongation time from published experimental measurements of

elongation rates (1-2kb/min) are shown. Note that the elongation scaling simulations require

elongation rates much slower than experimentally measured in order to achieve linearity. **G.**

ABC model selection results between the initiation scaling and elongation scaling TASEP

models for 3 genes across several strains. **H.** Distribution of the correlations of frequencies

of actively transcribing cells with cell length for different parameter sets from the **Figure 3E**

simulations. Note the strong positive correlations as observed in experimental data (**Figure**

3F). **I.** Boxplot of correlations between normalised intensities of the rpb1 mRNA and cell

length for all cell genotypes in sliding windows or containing different number of cells. Each

box represents a window of a given width. Sliding windows are shifted with increment of 1

cell along all cells ordered by length. Note that correlations are mainly positive. **J.** Average

gene analysis for the RNAPII ChIP-seq data from Figure 3H. Data were normalised to input

samples and split in 200 bins (50: upstream, 100: gene, 50: downstream) using the

deeptools package.

Figure S4 related to figure 4. A. Representative image of a single cell followed for a full

cell-cycle using ibidi slides. Brightfield (top), the nuclear marker Uch2 (middle) and the

RNAPII subunit Rpb1 bottom. **B.** Rpb1 normalised intensity/cell as a proxy for total cellular

protein concentration is plotted along time relative to the mitotic phase of the cell cycle. **C.**

Rpb9 total pixel intensity in the area of strong fluorescence signal (green mask in A) as a

proxy for chromatin bound amounts is plotted along time relative to the mitotic phase of the

cell cycle.

Figure S5 related to figure 5. A. Ratios of nascent site intensities of bi-nucleated cdc11-

119 cells plotted as a function of the ratio of local cytoplasmic volumes (**methods**). The

black line is the regression line from a linear model**. B.** Correlation between cell length and

nuclear area in pom1Δ (red) and wt (black) cells. Pearson corretions are shown. **C.** As in

Figure 5D for mRNA numbers. **D.** As in **Figure 5D** for mRNA numbers/nuclear volume

plotted as a function of nuclear volume. **E.** Simulation data from the model described in

Figure 5F. DNA bound RNAPII is plotted as a function of cell length. Magenta lines show a

running average. **F.** As in **Figure 5E** for mRNA numbers divided by nuclear area.

REFERENCES

1. Marguerat, S., and Bähler, J. (2012). Coordinating genome expression with cell size.
Trends Genet. *28*, 560–5.

2. Schmoller, K.M., and Skotheim, J.M. (2015). The Biosynthetic Basis of Cell Size
Control. Trends Cell Biol.

3. Zhurinsky, J., Leonhard, K., Watt, S., Marguerat, S., Bähler, J., and Nurse, P. (2010).
A coordinated global control over cellular transcription. Curr. Biol. CB *20*, 2010–2015.

4. Padovan-Merhar, O., Nair, G.P., Biaesch, A.G., Mayer, A., Scarfone, S., Foley, S.W.,
Wu, A.R., Churchman, L.S., Singh, A., and Raj, A. (2015). Single Mammalian Cells
Compensate for Differences in Cellular Volume and DNA Copy Number through
Independent Global Transcriptional Mechanisms. Mol. Cell *58*, 339–52.

5. Kempe, H., Schwabe, A., Crémazy, F., Verschure, P.J., and Bruggeman, F.J. (2015).
The volumes and transcript counts of single cells reveal concentration homeostasis
and capture biological noise. Mol. Biol. Cell *26*, 797–804.

6. Ietswaart, R., Rosa, S., Wu, Z., Dean, C., Howard, M., Ietswaart, R., Rosa, S., Wu, Z.,
Dean, C., and Howard, M. (2017). Cell-Size-Dependent Transcription of FLC and Its
Antisense Long Non-coding RNA COOLAIR Explain Article Cell-Size-Dependent
Transcription of FLC and Its Antisense Long Non-coding RNA COOLAIR Explain Cell-
to-Cell Expression Variation. Cell Syst. *4*, 622-635.e9.

7. Mena, A., Medina, D.A., García-Martínez, J., Begley, V., Singh, A., Chávez, S.,
Muñoz-Centeno, M.C., and Pérez-Ortín, J.E. (2017). Asymmetric cell division requires
specific mechanisms for adjusting global transcription. Nucleic Acids Res. *45*, 12401–
12412.

8. García-Martínez, J., Delgado-Ramos, L., Ayala, G., Pelechano, V., Medina, D.A.,
Carrasco, F., González, R., Andrés-León, E., Steinmetz, L., Warringer, J., *et al.*
(2016). The cellular growth rate controls overall mRNA turnover, and modulates either
transcription or degradation rates of particular gene regulons. Nucleic Acids Res. *44*,
3643–3658.

9. García-Martínez, J., Troulé, K., Chávez, S., and Pérez-Ortín, J.E. (2016). Growth rate
controls mRNA turnover in steady and non-steady states. RNA Biol. *13*, 1175–1181.

10. Schmoller, K.M., Turner, J.J., Kõivomägi, M., and Skotheim, J.M. (2015). Dilution of
the cell cycle inhibitor Whi5 controls budding-yeast cell size. Nature *526*, 268–72.

11. Keifenheim, D., Sun, X.-M., D'Souza, E., Ohira, M.J., Magner, M., Mayhew, M.B.,
Marguerat, S., and Rhind, N. (2017). Size-Dependent Expression of the Mitotic
Activator Cdc25 Suggests a Mechanism of Size Control in Fission Yeast. Curr. Biol.
27, 1491-1497.e4.

12. Saint, M., Bertaux, F., Tang, W., Sun, X.-M., Game, L., Köferle, A., Bähler, J.,
Shahrezaei, V., and Marguerat, S. (2019). Single-cell imaging and RNA sequencing
reveal patterns of gene expression heterogeneity during fission yeast growth and
adaptation. Nat. Microbiol. *4*, 480–491.

13. Swain, P.S., Elowitz, M.B., and Siggia, E.D. (2002). Intrinsic and extrinsic

contributions to stochasticity in gene expression. Proc. Natl. Acad. Sci. U. S. A. *99*,
12795–800.

14. Sanchez, A., and Golding, I. (2013). Genetic determinants and cellular constraints in
1029 noisy gene expression. Science *342*, 1188–93.

15. Coulon, A., Chow, C.C., Singer, R.H., and Larson, D.R. (2013). Eukaryotic
1031 transcriptional dynamics: From single molecules to cell populations. Nat. Rev. Genet.
1032 *14*, 572–584.

16. Adelman, K., and Lis, J.T. (2012). Promoter-proximal pausing of RNA polymerase II:
1034 emerging roles in metazoans. Nat. Rev. Genet. *13*, 720–731.

17. Chen, F.X., Smith, E.R., and Shilatifard, A. (2018). Born to run: Control of
1036 transcription elongation by RNA polymerase II. Nat. Rev. Mol. Cell Biol. *19*, 464–478.

18. Proudfoot, N.J. (2016). Transcriptional termination in mammals: Stopping the RNA
1038 polymerase II juggernaut. Science (80-.). *352*.

19. Heldt, F.S., Lunstone, R., Tyson, J.J., and Novák, B. (2018). Dilution and titration of
1040 cell-cycle regulators may control cell size in budding yeast. PLoS Comput. Biol. *14*, 1–
1041 26.

20. Lin, J., and Amir, A. (2018). Homeostasis of protein and mRNA concentrations in
1043 growing cells. Nat. Commun. *9*, 4496.

21. Raj, A., van den Bogaard, P., Rifkin, S.A., van Oudenaarden, A., and Tyagi, S.
1045 (2008). Imaging individual mRNA molecules using multiple singly labeled probes. Nat.
1046 Methods *5*, 877–879.

22. Ciechonska, M., Sturrock, M., Grob, A., Larrouy-Maumus, G., Shahrezaei, V., and
1048 Isalan, M. (2019). Ohm's Law for emergent gene expression under fitness pressure.
1049 bioRxiv, 693234.

23. Thomas, P. (2019). Intrinsic and extrinsic noise of gene expression in lineage trees.
1051 Sci. Rep. *9*, 474.

24. Bertaux, F., Marguerat, S., and Shahrezaei, V. (2018). Division rate, cell size and
1053 proteome allocation: impact on gene expression noise and implications for the
1054 dynamics of genetic circuits. R. Soc. Open Sci. *5*, 172234.

25. McAdams, H.H., and Arkin, A. (1997). Stochastic mechanisms in gene expression.
1056 Proc. Natl. Acad. Sci. U. S. A. *94*, 814–9.

26. Thattai, M., and van Oudenaarden, A. (2001). Intrinsic noise in gene regulatory
1058 networks. Proc. Natl. Acad. Sci. U. S. A. *98*, 8614–9.

27. Munsky, B., Trinh, B., and Khammash, M. (2009). Listening to the noise: random
1060 fluctuations reveal gene network parameters. Mol. Syst. Biol. *5*, 318.

28. Zechner, C., Ruess, J., Krenn, P., Pelet, S., Peter, M., Lygeros, J., and Koeppl, H.
1062 (2012). Moment-based inference predicts bimodality in transient gene expression.
1063 Proc. Natl. Acad. Sci. U. S. A. *109*, 8340–5.

29. Zechner, C., Unger, M., Pelet, S., Peter, M., and Koeppl, H. (2014). Scalable
1065 inference of heterogeneous reaction kinetics from pooled single-cell recordings. Nat.
1066 Methods *11*, 197–202.

30. Fritzsch, C., Baumgärtner, S., Kuban, M., Steinshorn, D., Reid, G., and Legewie, S.
(2018). Estrogen-dependent control and cell-to-cell variability of transcriptional
bursting. Mol. Syst. Biol. *14*, e7678.

31. Suter, D.M., Molina, N., Gatfield, D., Schneider, K., Schibler, U., and Naef, F. (2011).
Mammalian genes are transcribed with widely different bursting kinetics. Science *332*,
472–4.

32. Komorowski, M., Costa, M.J., Rand, D.A., and Stumpf, M.P.H. (2011). Sensitivity,
robustness, and identifiability in stochastic chemical kinetics models. Proc. Natl. Acad.
Sci. U. S. A. *108*, 8645–50.

33. Tiberi, S., Walsh, M., Cavallaro, M., Hebenstreit, D., and Finkenstädt, B. (2018).
Bayesian inference on stochastic gene transcription from flow cytometry data.
Bioinformatics *34*, i647–i655.

34. Watt, S., Mata, J., López-Maury, L., Marguerat, S., Burns, G., and Bähler, J. (2008).
urg1: a uracil-regulatable promoter system for fission yeast with short induction and
repression times. PLoS One *3*, e1428.

35. Mercier, A., and Labbé, S. (2010). Iron-Dependent remodeling of fungal metabolic
pathways associated with ferrichrome biosynthesis. Appl. Environ. Microbiol. *76*,
3806–3817.

36. Wilusz, C.J., and Wilusz, J. (2004). Bringing the role of mRNA decay in the control of
gene expression into focus. Trends Genet. *20*, 491–497.

37. Amorim, M.J., Cotobal, C., Duncan, C., and Mata, J. (2010). Global coordination of
transcriptional control and mRNA decay during cellular differentiation. Mol. Syst. Biol.
6, 380.

38. Zenklusen, D., Larson, D.R., and Singer, R.H. (2008). Single-RNA counting reveals
alternative modes of gene expression in yeast. Nat. Struct. Mol. Biol. *15*, 1263–1271.

39. Zhao, Y.-B., and Krishnan, J. (2014). mRNA translation and protein synthesis: an
analysis of different modelling methodologies and a new PBN based approach. BMC
Syst. Biol. *8*, 25.

40. Sahoo, M., and Klumpp, S. (2013). Backtracking dynamics of RNA polymerase:
pausing and error correction. J. Phys. Condens. Matter *25*, 374104.

41. Voliotis, M., Cohen, N., Molina-París, C., and Liverpool, T.B. (2008). Fluctuations,
pauses, and backtracking in DNA transcription. Biophys. J. *94*, 334–48.

42. Dobrzynski, M., and Bruggeman, F.J. (2009). Elongation dynamics shape bursty
transcription and translation. Proc. Natl. Acad. Sci. U. S. A. *106*, 2583–8.

43. Darzacq, X., Shav-Tal, Y., de Turris, V., Brody, Y., Shenoy, S.M., Phair, R.D., and
Singer, R.H. (2007). In vivo dynamics of RNA polymerase II transcription. Nat. Struct.
Mol. Biol. *14*, 796–806.

44. Edwards, A.M., Kane, C.M., Young, R.A., and Kornberg, R.D. (1991). Two dissociable
subunits of yeast RNA polymerase II stimulate the initiation of transcription at a
promoter in vitro. J. Biol. Chem. *266*, 71–5.

45. Mason, P.B., and Struhl, K. (2005). Distinction and relationship between elongation
rate and processivity of RNA polymerase II in vivo. Mol. Cell *17*, 831–40.

46. Pelechano, V., Chávez, S., and Pérez-Ortín, J.E. (2010). A Complete Set of Nascent
Transcription Rates for Yeast Genes. PLoS One *5*, e15442.

47. Swinburne, I.A., and Silver, P.A. (2008). Intron Delays and Transcriptional Timing
during Development. Dev. Cell *14*, 324–330.

48. Neumann, F.R., and Nurse, P. (2007). Nuclear size control in fission yeast. J. Cell

Biol. *179*, 593–600.

49.   Verde, F., Mata, J., and Nurse, P. (1995). Fission yeast cell morphogenesis:
identification of new genes and analysis of their role during the cell cycle. J. Cell Biol.
131, 1529–38.

50.   Bahler, J., and Pringle, J.R. (1998). Pom1p, a fission yeast protein kinase that
provides positional information for both polarized growth and cytokinesis. Genes Dev.
12, 1356–1370.

51.   Hayles, J., Wood, V., Jeffery, L., Hoe, K.-L., Kim, D.-U., Park, H.-O., Salas-Pino, S.,
Heichinger, C., and Nurse, P. (2013). A genome-wide resource of cell cycle and cell
shape genes of fission yeast. Open Biol. *3*, 130053.

52.   Kim, D.-U., Hayles, J., Kim, D., Wood, V., Park, H.-O., Won, M., Yoo, H.-S., Duhig, T.,
Nam, M., Palmer, G., *et al.* (2010). Analysis of a genome-wide set of gene deletions in
the fission yeast Schizosaccharomyces pombe. Nat. Biotechnol. *28*, 617–623.

53.   Weston, L., Greenwood, J., and Nurse, P. (2017). Genome-wide screen for cell
growth regulators in fission yeast. J. Cell Sci. *130*, 2049–2055.

54.   Larsson, A.J.M., Johnsson, P., Hagemann-Jensen, M., Hartmanis, L., Faridani, O.R.,
Reinius, B., Segerstolpe, Å., Rivera, C.M., Ren, B., and Sandberg, R. (2019).
Genomic encoding of transcriptional burst kinetics. Nature *565*, 251–254.

55.   Bartman, C.R., Hamagami, N., Keller, C.A., Giardine, B., Hardison, R.C., Blobel, G.A.,
and Raj, A. (2018). Transcriptional Burst Initiation and Polymerase Pause Release
Are Key Control Points of Transcriptional Regulation. Mol. Cell.

56.   Keren, L., Zackay, O., Lotan-Pompan, M., Barenholz, U., Dekel, E., Sasson, V.,
Aidelberg, G., Bren, A., Zeevi, D., Weinberger, A., *et al.* (2013). Promoters maintain
their relative activity levels under different growth conditions. Mol. Syst. Biol. *9*, 701.

57.   Windner, S.E., Manhart, A., Brown, A., Mogilner, A., and Baylies, M.K. (2019).
Nuclear Scaling Is Coordinated among Individual Nuclei in Multinucleated Muscle
Fibers. Dev. Cell *49*, 48-62.e3.

58.   Billman, M.R., Rueda, D., and Bangham, C.R.M. (2017). Single-cell heterogeneity and
cell-cycle-related viral gene bursts in the human leukaemia virus HTLV-1. Wellcome
open Res. *2*, 87.

59.   Youn, M., Takada, I., Imai, Y., Yasuda, H., and Kato, S. (2010). Transcriptionally
active nuclei are selective in mature multinucleated osteoclasts. 1025–1035.

60.   Roberts, S.E., and Gladfelter, A.S. (2015). Nuclear autonomy in multinucleate fungi.
Curr. Opin. Microbiol. *28*, 60–65.

61.   Dundon, S.E.R., Chang, S.-S., Kumar, A., Occhipinti, P., Shroff, H., Roper, M., and
Gladfelter, A.S. (2016). Clustered nuclei maintain autonomy and nucleocytoplasmic
ratio control in a syncytium. Mol. Biol. Cell *27*, 2000–2007.

62.   Wu, Z., Ietswaart, R., Liu, F., Yang, H., Howard, M., and Dean, C. (2016). Quantitative
regulation of FLC via coordinated transcriptional initiation and elongation. Proc. Natl.
Acad. Sci. U. S. A. *113*, 218–23.

63.   Battich, N., Stoeger, T., and Pelkmans, L. (2015). Control of Transcript Variability in
Single Mammalian Cells. Cell *163*, 1596–1610.

64.   Soltani, M., Vargas-Garcia, C.A., Antunes, D., and Singh, A. (2016). Intercellular
Variability in Protein Levels from Stochastic Expression and Noisy Cell Cycle
Processes. PLoS Comput. Biol. *12*, e1004972.

65. Takahashi, C.N., Miller, A.W., Ekness, F., Dunham, M.J., and Klavins, E. (2015). A
Low Cost, Customizable Turbidostat for Use in Synthetic Circuit Characterization.
ACS Synth. Biol. *4*, 32–38.

66. Heinrich, S., Geissen, E.-M., Kamenz, J., Trautmann, S., Widmer, C., Drewe, P.,
Knop, M., Radde, N., Hasenauer, J., and Hauf, S. (2013). Determinants of robustness
in spindle assembly checkpoint signalling. Nat. Cell Biol. *15*, 1328–39.

67. Trcek, T., Chao, J.A., Larson, D.R., Park, H.Y., Zenklusen, D., Shenoy, S.M., and
Singer, R.H. (2012). Single-mRNA counting using fluorescent in situ hybridization in
budding yeast. Nat. Protoc. *7*, 408–19.

68. Mueller, F., Senecal, A., Tantale, K., Marie-Nelly, H., Ly, N., Collin, O., Basyuk, E.,
Bertrand, E., Darzacq, X., and Zimmer, C. (2013). FISH-quant: automatic counting of
transcripts in 3D FISH images. Nat. Methods *10*, 277–278.

69. Rodríguez-López, M., Gonzalez, S., Hillson, O., Tunnacliffe, E., Codlin, S., Tallada,
V.A., Bähler, J., and Rallis, C. (2019). The GATA transcription factor Gaf1 represses
tRNA genes, inhibits growth, and extends chronological lifespan downstream of
fission yeast TORC1. bioRxiv, 700286.

70. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics *25*, 1754–1760.

71. Lock, A., Rutherford, K., Harris, M.A., Hayles, J., Oliver, S.G., Bähler, J., and Wood,
V. (2019). PomBase 2018: user-driven reimplementation of the fission yeast database
provides rapid and intuitive access to diverse, interconnected information. Nucleic
Acids Res. *47*, D821–D827.

72. Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with
high-throughput sequencing data. Bioinformatics *31*, 166–169.

73. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

74. Wiśniewski, J.R., Hein, M.Y., Cox, J., and Mann, M. (2014). A "proteomic ruler" for
protein copy number and concentration estimation without spike-in standards. Mol.
Cell. Proteomics *13*, 3497–506.

75. Ramírez, F., Dündar, F., Diehl, S., Grüning, B.A., and Manke, T. (2014). deepTools: a
flexible platform for exploring deep-sequencing data. Nucleic Acids Res. *42*, W187–
W191.

76. Edelstein, A.D., Tsuchida, M.A., Amodaj, N., Pinkard, H., Vale, R.D., and Stuurman,
N. (2014). Advanced methods of microscope control using µManager software. J.
Biol. Methods *1*, 10.

77. Amir, A. (2014). Cell size regulation in bacteria. Phys. Rev. Lett. *112*.

78. Tanouchi, Y., Pai, A., Park, H., Huang, S., Stamatov, R., Buchler, N.E., and You, L.
(2015). A noisy linear map underlies oscillations in cell size and gene expression in
bacteria. Nature *523*, 357–360.

79. Jun, S., and Taheri-Araghi, S. (2014). Cell-size maintenance: universal strategy
revealed. Trends Microbiol. *23*, 4–6.

80. Wang, P., Robert, L., Pelletier, J., Dang, W.L., Taddei, F., Wright, A., and Jun, S.
(2010). Robust growth of Escherichia coli. Curr. Biol. *20*, 1099–103.

81. Gillespie, D.T. (1977). Exact stochastic simulation of coupled chemical reactions. J.
Phys. Chem. *81*, 2340–2361.

82. Shahrezaei, V., Ollivier, J.F., and Swain, P.S. (2008). Colored extrinsic fluctuations
and stochastic gene expression. Mol. Syst. Biol. *4*, 196.

83. Peccoud, J., and Ycart, B. (1995). Markovian Modeling of Gene-Product Synthesis.
Theor. Popul. Biol. *48*, 222–234.

84. Shahrezaei, V., and Swain, P.S. (2008). Analytical distributions for stochastic gene
expression. Proc. Natl. Acad. Sci. U. S. A. *105*, 17256–61.

85. Rubin, D.B. (1984). Bayesianly Justifiable and Relevant Frequency Calculations for
the Applied Statistician. Ann. Stat. *12*, 1151–1172.

86. Marin, J.-M., Pudlo, P., Robert, C.P., and Ryder, R.J. (2012). Approximate Bayesian
computational methods. Stat. Comput. *22*, 1167–1180.

87. Toni, T., Welch, D., Strelkowa, N., Ipsen, A., and Stumpf, M.P.. (2009). Approximate
Bayesian computation scheme for parameter inference and model selection in
dynamical systems. J. R. Soc. Interface *6*, 187–202.

88. Lenormand, M., Jabot, F., and Deffuant, G. (2015). Adaptive approximate bayesian
computation for complex models. Comput. Stat. *2013*, 2777–2796.

1219

Figure 1 Sun, Bowman et al

Figure 4 Sun, Bowman et al

Figure 5 Sun, Bowman et al

Figure S1 Sun, Bowman et al

Figure S2 Sun, Bowman et al

A

B

Figure S4 Sun, Bowman et al

A

Bright field

Uch2-mCherry

Rpb1-GFP

Time

B

Normalised intensity

1 nucleus   2 nuclei
Time from mitosis

C

Total pixel intensity

1 nucleus   2 nuclei
Time from mitosis

Figure S5 Sun, Bowman et al