Title: Anatomical features, efficient photosystem, and Rubisco activity contribute to high leaf photosynthesis in wild rice

Running Title: Efficient Leaf Photosynthesis in Wild Rice

Jyotirmaya Mathan[#], Anuradha Singh[#], Vikram Jathar, Aashish Ranjan*

National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-110067, India.

[#]These authors contributed equally to the work

*Corresponding Author

email: aranjan@nipgr.ac.in

Phone no. 91-11-26735117

Authors' email

Jyotirmaya Mathan: jyotirmaya@nipgr.ac.in

Anuradha Singh: annusingh1206@gmail.com

Vikram Jathar: vikra.jathar@gmail.com

Date of submission: September 24, 2019

Number of figures: 6 figure (all colour)

The word count: 6,418

Supplementary data: 7 supplementary figures and 6 supplementary tables

Title: Anatomical features, efficient photosystem, and Rubisco activity contribute to
high leaf photosynthesis in wild rice

Running Title: Efficient Leaf Photosynthesis in Wild Rice

Highlight

Distinct leaf developmental, photochemical and biochemical features contribute to efficient
photosynthesis in wild rice that could potentially be exploited for increasing rice leaf
photosynthesis.

Abstract

The importance of increasing photosynthetic efficiency for sustainable crop yield
increases to feed the growing world population is well recognized. The natural genetic
variation for leaf photosynthesis in crop plants is an overlooked and untapped resource. The
genus *Oryza*, including cultivated rice and wild relatives, offers tremendous genetic
variability to explore photosynthetic differences, and underlying developmental,
photochemical, and biochemical basis. We quantified leaf photosynthesis and related
physiological parameters for ten cultivated and wild rice genotypes to identify
photosynthetically efficient wild rice species. Wild rice species with high leaf photosynthesis
per unit area had striking anatomical features, such as larger mesophyll cells with more
chloroplasts, larger and closer veins, and a smaller number of mesophyll cells between two
consecutive veins. In addition, photosynthetically efficient wild rice species showed an
efficient Photosystem II as well as higher carboxylation activity of Rubisco compared to less
efficient cultivated varieties, such as IR 64. Our results show the existence of desirable
variations in mesophyll and vein features, Photosystem II efficiency, and Rubisco activity in
the rice system itself that could possibly be targeted for higher leaf photosynthesis. Detailed
genetic and molecular understanding of these traits shall be instrumental in increasing
photosynthetic efficiency of cultivated rice.

Keywords: Chlorophyll fluorescence, Leaf anatomy, Photosynthesis, Photosystem II (PSII),
Rubisco, Wild rice.

Introduction

A remarkable increase in food grain production is needed to meet the demand of the growing world population that is expected to reach 10 billion by 2050 (Alexandratos and Bruinsma, 2012). Crop yield increases due to extensive breeding and an agronomic practice is approaching a plateau over the last decade (Lawson *et al.,* 2012). Yield increases due to the breeding for semi-dwarf varieties, resulting in optimization of photosynthates' allocation, during the green revolution have also largely been saturated (Jansson *et al.,* 2018). The further increases in crop yield and productivity, therefore, requires improving photosynthetic efficiency of crop plants (Lawson *et al.,* 2012; Ort *et al.,* 2015; Simkin *et al.,* 2019).

Photosynthesis, the key biochemical process fixing atmospheric CO_2 in the form of carbohydrates, is a multifaceted process that has contributions from leaf physiological, developmental, and biochemical features (Long *et al.,* 2006; Mathan *et al.,* 2016). Stomatal conductance (g_s), carboxylation efficiency (CE), transpiration (E), and water-use efficiency (WUE) are critical physiological parameters determining leaf gaseous exchange and water relationship that eventually affect the leaf photosynthesis (Farquhar and Sharkey, 1982). Extrinsic factors, such as light and CO_2 concentration, influence photosynthetic performance by regulating the activity of light-harvesting complexes (LHCs) and Rubisco as well as stomatal opening (Eichelmann and Laisk, 2002; Murchie and Niyogi, 2011; Hiyama *et al.,* 2017). In addition, the diurnal timing of the day influences the photosynthesis by mid-day stomatal closure and photorespiration (Zhao *et al.,* 2010).

Developmental characteristics of leaf are strong determinants of photosynthesis (Mathan *et al.,* 2016). Flag leaf width, flag leaf thickness, and specific leaf area showed a positive correlation with photosynthetic capacity, thus effecting overall crop performance (Liu *et al.,* 2015). Anatomical features such as size, lobbing, and surface area of mesophyll cells, are critical for proper light distribution and CO_2 diffusion within a leaf (Sage and Sage, 2009; Tholen *et al.,* 2012). Similarly, leaf vasculature serves as a channel for distribution of water and minerals for photosynthesis and mobilizes photoassimilated CO_2 from leaves to sink organs (Sack and Scoffoni, 2013). Vein density and interveinal distance have been shown to influence CO_2 fixation efficiency (Amiard *et al.,* 2005; Brodribb *et al.,* 2007). Thus, investigating relationships between leaf anatomy and photosynthetic features, and identification of genetic regulators of desirable anatomical features could help in optimizing crop photosynthetic efficiency (Evans, 2013).

Photochemical features, such as organization of light-harvesting complexes, quantum efficiency of photosystem, electron transport rate, and photochemical and non-photochemical

quenching are key factors for photosynthetic efficiency (Murchie and Lawson, 2013; Wu *et*

al., 2019). Maintaining a higher level of chlorophyll content for longer duration in a leaf is

one of the suggested strategies for increasing net photosynthesis and crop production (Huang

et al., 2013). Quantification of chlorophyll fluorescence is used to differentiate

photochemical quenching and non-photochemical quenching, and to evaluate the efficiency

of Photosystem II (PSII) reaction centres (Kalaji *et al.,* 2016). Approaches to increase light

harvesting by PSII reaction centres and photochemical quenching shall be effective in

increasing crop photosynthetic efficiency (Cardona *et al.,* 2018).

Different steps of CO_2 photoassimilation are under complex genetic and

environmental regulation. Transcriptional control of photosynthesis and associated leaf

features is suggested to have potential for improving photosynthetic assimilation (Wang *et*

al., 2017). Transcription factors, such as GLK1, HY5, ATAF1, ORE1, and PIF, have been

shown to activate the photosynthetic gene expression in crop plants (Toledo-Ortiz *et al.,*

2014; Wang *et al.,* 2017). Golden2-like (GLK) transcription factor positively regulates the

chloroplast development and also binds directly to the promoter of LHC genes (Waters *et al.,*

2009). Similarly, HY5 and PIF4/5 are important regulators controlling the regulatory network

governing photosynthetic gene expression and senescence, respectively (Kindgren *et al.,*

2012; Toledo-*Ortiz et al.,* 2014). However, genetic rewiring of transcription factors towards

improving crop photosynthetic performance remains elusive.

Over the past years, the biochemical understanding of photosynthesis has been

extensively elaborated. There have been examples for increasing photosynthetic efficiency,

resulting in increased yield and/or biomass. One such example is the evolution of C_4 plants,

where specific developmental and biochemical features promote efficient photoassimilation

of CO_2 (Kajala *et al.,* 2011; Koteyeva *et al.,* 2011). Free air CO_2 enrichment increases leaf

photosynthesis in C_3 plants (Rogers *et al.,* 2004). Optimization of additional parameters, such

as decreasing Rubisco oxygenase activity, increasing RuBP regeneration, suppressing

photorespiration, and increasing photoprotection of the photosynthesis have also been

suggested to improve photosynthetic efficiency (Long *et al.,* 2006; Murchie and Niyogi,

2011).

Engineering C_4 photosynthesis in C_3 plants has been envisioned as a potential strategy

towards enhancing crop photosynthetic performance. However, this is challenging due to

evolutionary, developmental, and biochemical constraints (von Caemmerer *et al.,* 2012;

Schuler *et al.,* 2016; Reeves *et al.,* 2017). The natural genetic variation in leaf photosynthesis,

associated developmental, photochemical, and biochemical features, and underlying genetic

regulation in crop plants is an overlooked and untapped resource (Flood *et al.,* 2011; van
Bezouw *et al.,* 2019). Rice, with two cultivated species: *Oryza sativa* (Asian rice with two
subspecies: *indica and japonica*) and *Oryza glaberrima* (African rice), and many related wild
rice species, provides an excellent system to investigate photosynthetic differences, and
underlying genetic basis. Many of the wild rice species have been used in crop breeding
programs towards improving biotic and abiotic stress tolerance of the cultivated rice
(Sanchez *et al.,* 2013; Atwell *et al.,* 2014; Wing *et al.,* 2018). Interestingly, some of the wild
rice species (*O. rufipogon*, *O. latifolia*, and *O. australiensis*) are reported to accumulate
higher biomass than cultivated varieties, and introgression from a wild *Oryza* species to
cultivated rice has also been shown to increase biomass (Yeo *et al.,* 1994; Zhao *et al.,* 2008).
However, a systematic effort to identify the photosynthetic differences, linking the
differences to developmental, photochemical, and biochemical features, and investigating the
underlying genetic and molecular basis is warranted.

We examined a set of cultivated (including both subspecies of *Oryza sativa* and *Oryza*
glaberrima) and wild rice species, differing in biomass accumulation, to find out strong
differences in leaf photosynthesis. We, further, identified the developmental, photochemical,
and biochemical differences among the selected photosynthetically different accessions.
Expression pattern of genes encoding photosynthetic components, chlorophyll-biosynthetic
enzymes and transcription factors, and Rubisco enzyme activity corroborated the leaf
photosynthesis differences across cultivated and wild rice.

Materials and methods
Plant material
Ten rice genotypes, consisting of four varieties of *Oryza sativa* ssp. *indica* (IR 64, Swarna,
Nagina 22, and Vandana), a variety from *Oryza sativa* ssp. *japonica* (Nipponbare), African
cultivated rice *Oryza glaberrima*, and four wild rice species, *Oryza rufipogon, Oryza*
punctata, Oryza latifolia, and *Oryza australiensis* were grown in Kharif 2016 and 2017 at
NIPGR research farm (Supplementary Table S1). These genotypes were also grown in pots
under controlled growth chambers with the temperature (28 ℃), humidity (70-90 %), and day
length (14h: 10h, light: dark). Physiological, developmental, photochemical, and biochemical
traits were evaluated at heading stage in the widest part of the flag leaves.

Photosynthesis rate and related physiological trait measurements

Flag leaf photosynthesis was measured at the full heading stage with a portable photosynthesis system (LI-6400XT equipped with a 6400-40 Leaf Chamber Fluorometer (LI-COR Biosciences). Measurement was conducted on a clear day between 9.00 h to 11.00 h under ambient CO_2 level (400 µmol mol^{-1}), air temperature (30 ℃), leaf temperature (27 ℃), and PAR (1,000 µmol m^{-2} s^{-1}) conditions. Net photosynthesis rate (P_N), and related physiological parameters such as stomatal conductance (g_s), intercellular CO_2 concentration (Ci), transpiration rate (E), water use efficiency (WUE; ratios P_N : E), and carboxylation efficiency (CE; ratios P_N : Ci) were recorded once the expanded leaf was enclosed in the LI-COR leaf chamber. In order to examine the effect of light intensity and CO_2 concentration, leaf photosynthesis was measured at different photosynthetically active radiations (PAR; 0, 300, 500, 700, 1000, 1200, and 1500 µmol m^{-2} s^{-1}), and at intercellular CO_2 concentration (Ci; 0, 400, 600, 800, 1000, 1200, and 1400 µmol mol^{-1}). Further, leaf photosynthesis was measured at every hour from 9.00 h to 15.00 h to examine the diurnal effect on P_N for the selected cultivated and wild rice as described in Feldman et al. (2017).

Visualization and quantification of leaf developmental traits

Morphological trait such as flag leaf width (FLW), leaf thickness (FLT) was measured from the widest part of the leaves from ten individual rice plants for each accession. For anatomical traits visualization, transverse sections were cut from the mid-portion of four leaves of different plants, and stained with toluidine blue with some modification of the method as described in Lux et al. (2005). Briefly, thin sections were mounted with 50 % glycerol on a tissue culture plate, chlorophyll pigments were cleared using 85 % (w/v) lactic acid saturated with chloral hydrate, heated on a water bath at 70 ℃ for 1 hour and washed several times with distilled water. To visualize mesophyll cells, sections were incubated in absolute alcohol for overnight, stained with 1% toluidine blue in 1% (w/v) disodium tetraborate for 15 sec. Finally, thin sections were visualized using a bright field microscope (LMI, UK). The quantification of leaf anatomical features related to minor vein, mesophyll cell and, bundle sheath cells were performed at 10X and 100X magnification (ten images per leaf) using Fiji-Image J software (Wayne Rasband, National Institute of Health, USA) as described in Chatterjee et al. (2016) and Ivanova et al. (2017).

Chloroplast ultrastructure

Freshly harvested flag leaf segments (1 × 1 mm; four leaves from four different plants) were fixed under fixative buffer (3% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.2) for 2 hours and then treated with 1% osmium tetroxide overnight at 4 ℃. The fixed segments were dehydrated in a graded acetone series and embedded in epoxy resin. Ultrathin sections were cut with diamond knives, stained with uranyl acetate and lead citrate double staining procedure as described by Reynolds (1963). Chloroplasts of the uppermost part of the leaf sections were viewed using a digital camera (BH-2, Olympus) equipped in transmission electron microscopy at 1000X magnification (JEM 1230; JEOL).

Leaf chlorophyll measurement

Total chlorophyll content (TCC) were measured in 80% acetone following the procedure described by Kiran *et al.* (2013). Briefly, flag leaf (50 mg) was incubated in 5 ml ice-cold 80 % acetone for 48 h, centrifuged at 5,000 × g for 5 min and absorbance was recorded at 663 nm and 645 nm. The pigment concentration was calculated using the method described by van Campen *et al.* (2016) and expressed in mg g^{-1} FW.

Leaf Chlorophyll Fluorescence Measurements

Leaf chlorophyll (chl) fluorescence was measured using the LI-6400XT fluorometer (LI-COR Biosciences) according to the method described in van Campen *et al.* (2016). Plants were grown under controlled growth conditions, dark-adapted for at least 45 min before the measurement of F_v/F_m (ratio of the variable to maximum fluorescence - the quantum efficiency of open PSII centers). After the initial or minimum fluorescence (F_o) was measure, plants were illuminated with a 0.8 sec saturated pulse of light provided by an array of light-emitting diodes (LEDs) [mixture of red (90 %) and blue (10 %) lights at 650 nm peak wavelength, 1000 μmol m^{-2} s^{-1}] focused on a circular 2.0 cm^2 leaf surface area. Maximum fluorescence (F_m) was measured, and F_v/F_m calculated as (F_m-F_o/F_m). Plants were then placed under natural irradiance, and the steady-state of fluorescence (F_s) was recorded. A second saturating pulse at 1000 μmol m^{-2} s^{-1} was imposed to determine the maximal fluorescence level in light-adapted leaves ($F_m´$). Then, the actinic light was removed, and the minimal fluorescence level in the light-adapted state ($F_o´$) was determined through of illuminating the leaf with 1 sec of far-red light. Using light fluorescence parameters, we calculated: (a) ratio of variable to maximum fluorescence - the quantum efficiency of open PSII centers (F_v/F_m) in dark-adapted state, (b) the actual PSII efficiency (φPSII), (c) photochemical quenching

203 coefficient (qP), and (d) the photosynthetic electron transport rate (ETR) in light-adapted
204 state as described in Murchie and Lawson (2013).
205
RNA isolation and expression analysis
Total RNA was isolated from flag leaf and leaf 4 at primordial 4 stages (P4) (van Campen *et*
al., 2016). Tissues were harvested, immediately frozen in liquid nitrogen. Total RNA
extraction (TRIzol reagent), DNase treatment (DNase enzyme), and cDNA synthesis
(RevertAid First-strand cDNA synthesis kit) were performed according to the manufacturer
protocol (Thermo Fisher Scientific, USA). The PowerUp™ SYBR® Green fluorescence dye
and specific primer sets (Supplementary Table S6) were used for quantitative gene
expression analysis on CFX connect Real-time PCR detection system (Bio-Rad, CA, USA).
The expression analysis was confirmed by analyzing two biological replicates, with three
technical repeats each. Rice actin was used as an internal control gene for normalization of
gene expression. The relative change or fold change was calculated following the method of
(Livak and Schmittgen, 2001).
218
Rubisco activity assay
Rubisco extraction and activity (Vcmax) was performed following the procedure described in
Kubien *et al.* (2011) and Scales *et al.* (2014). Briefly, leaf tissue (3.6×1.0 cm^2) from
illuminated leaves were excised from five plants of each species and homogenized in
extraction buffer [EB, 5 mM DTT, 10 mg ml^{-1} PVPP, 2 mg ml^{-1} BSA, 2 mg ml^{-1} PEG, 2%
(v/v) Tween-80, 12 mM 6-aminocaproic acid, 2.4 mM benzamidine in HEPES stock) at 4 ℃.
The homogenate centrifuged for 1 min at ~16,000 $\times$ g and pellet was discarded. To assess the
activity of Rubisco from leaf extract, 50 µl of leaf extract was incubated with 1,550 ml of the
EME buffer, 40 µl 10 mM of NADH (final concentration 200 µM), 80 µl of 250 mM
NaH^{12}CO$_3$ (10 mM), 200 µl of ATP/phosphocreatine (1 and 5 mM, respectively), 40 µl of
coupling enzymes (Carbonic anhydrase 500U, Glyceraldehyde-3-P dehydrogenase 50U, 3-
PGA kinase 50U, Triose-P Isomerase/Glycerol-3-P dehydrogenase 400/40U), 40 µl of 25
mM RuBP (500 µM), mixed gently, and decrease of absorbance was monitored at 340 nm
continuously for 0 to 180 sec using a UV–Vis spectrophotometer. Linear region of
spectrophotometer absorbance were used to determine rate of RuBP-use using equations
given below:

$$RuBP - use \left(\frac{\mu mol}{s} \right) = \frac{A(t0 - t1) * vol}{6.18 * 4 * (t0 - t1)}$$

Where A is absorbance (340 nm) at time 0 or time 1, vol (assay volume in ml), t time in sec,

and 6.18 is an extinction coefficient in $\mu mol\ mL^{-1}cm^{-1}$ for NADH oxidation at 340 nm.

Further, the maximum rate of RuBP-use was also used to calculate Rubisco activity in terms

of Vcmax.

Statistical analysis

One-way analysis of variance (ANOVA) followed by statistical significance subscripts were

grouped according to post-hoc Tukey HSD calculation ($*P < 0.05$, $**P < 0.01$). Least

significant difference was calculated by Fisher's test ($\alpha = 0.05$). Pearson correlation matrices

were calculated based on mean values of each leaf developmental and physiological trait for

each accession to evaluate the trait-to-trait associations. The level of significance (α) of

correlation coefficients was determined at $*P < 0.05$ (± 0.666; degree of freedom (df; n-2; 9-

2 = 7).

Results

Quantification of leaf photosynthesis and physiological parameters

Cultivated and wild rice species, showing remarkable differences for biomass

accumulation for the same crop growth period, were examined for leaf photosynthesis (P_N)

and related physiological features (Supplementary Fig. S1). The mean P_N value for the wild

rice species, *O. latifolia*, *O. australiensis*, and *O. rufipogon*, and African cultivated rice

Oryza glaberrima were found to be significantly higher than the cultivated rice varieties,

such as IR64, Nagina 22 and Vandana (Fig. 1A). Among the selected accessions, *O.*

australiensis and *O. latifolia* showed highest P_N, 23.50 and 24.18 $\mu mol\ CO_2\ m^{-2}\ s^{-1}$,

respectively. Also, we observed remarkable variations for P_N among the cultivated varieties.

For instance, one of the high yielding *indica* variety Swarna and *japonica* variety Nipponbare

displayed higher P_N compared to other cultivated varieties.

We further examined other important physiological traits of the selected accessions.

As expected, wild rice species, *O. latifolia* and *O. australiensis* with very high P_N, showed

higher stomatal conductance (g_s), and intercellular CO_2 concentration (Ci) (Fig. 1B, C). In

addition, these species along with *O. rufipogon* displayed higher carboxylation efficiency

(CE) than those of cultivated rice (Fig. 1F). Interestingly, *O. australiensis*, one of the two

wild species with highest P_N, had relatively lower transpiration rate (E), resulting in highest

water use efficiency (WUE) for the species (Fig. 1D, E). Wild species *O. punctata* and *O.*

sativa ssp *indica* cv. Vandana also exhibited lower E compared to other genotypes (Fig. 1D),

and corresponding higher WUE of Vandana (Fig. 1E). WUE of *O. punctata* was still

relatively lower due to lower P_N of the species (Fig. 1A, E).

Pearson correlation analysis showed a strong positive correlation between P_N with g_s

(*r = 0.665*), Ci (*r = 0.632*), and CE (*r = 0.656*) (Supplementary Fig. S2). Higher WUE

results from higher P_N coupled with lower E, which is evident from the strong positive

correlation between WUE and P_N (*r = 0.618*) and a negative correlation between WUE and E

(r = *-0.589*). A poor correlation was observed between P_N and E (*r = 0.248*) (Supplementary

Fig. S2). We, further, quantified leaf photosynthesis and related physiological traits at early

vegetative stage under controlled conditions in two independent experiments. Similar to our

observations at full heading stage under field conditions, wild rice species (excluding *O.*

punctata) and *O. glaberrima* showed higher P_N and g_s compared to cultivated rice since the

beginning of vegetative stage (Supplementary Table S2). Taken together, these results

showed significant differences in photosynthetic traits across cultivated and wild rice, with *O.*

australiensis showing higher WUE along with remarkably higher leaf photosynthesis than

cultivated rice varieties. We then selected representative cultivated rice accessions (IR 64,

Nipponbare, and *O. glaberrima*) and wild rice species (*O. rufipogon, O. latifolia,* and *O.*

australiensis) for detailed analyses.

Effects of light intensity, CO_2 concentration, and diurnal time on photosynthesis and

related physiological traits

Light intensity and CO_2 concentration are the two most important factors determining

leaf photosynthesis. Therefore, we examined the effect of different light intensity and CO_2

concentration on P_N, g_s, E, WUE, P_N (max), Ci and CE of selected wild and cultivated rice

accessions (Fig. 2 and Supplementary Fig. S3). All the species showed a steeper initial slope,

instantaneous enhancement followed by a plateau for leaf photosynthesis per unit area with

an increasing photosynthetically active radiation (PAR) and CO_2 concentration (Fig. 2A and

2B-I). Wild rice species, *O. latifolia* and *O. australiensis*, together with *O. glaberrima* and *O.*

sativa cv. Nipponbare exhibited higher P_N at the beginning, with clearly higher P_N at PAR >

1000 μmol m^{-2} s^{-1} (Fig. 2A-I). Similarly, P_N in response to increasing CO_2 concentration

segregated the selected genotypes under two distinct groups: *O. australiensis, O. latifolia* and

O. glaberrima with higher P_N, and cultivated varieties with lower P_N. The genotypes showed

the pattern of g_s similar to that of P_N in response to changing light intensity (Fig. 2A-II). The

g_s, however, showed a decreasing trend over increasing CO_2 concentration, likely due to the

closure of stomata (Fig. 2B-II). Nonetheless, P_N showed an increasing trend at higher
concentration of CO_2. Interestingly, wild rice species were able to maintain higher g_s at
different light intensity and CO_2 concentration. Cultivated varieties showed increased E, thus
reduced WUE, in response to increasing light intensity (Fig. 2A, III –IV). In contrast, the
wild species, *O. australiensis* and *O. latifolia*, showed lower E, and hence higher WUE,
compared to cultivated varieties at different light intensities. Reducing E, and hence
increased WUE, was detected for all the genotypes beyond 400 µmol mol^{-1} CO_2
concentration (Fig. 2B, III –IV).

In order to investigate the effect of diurnal timing, P_N and other physiological traits
were quantified for the selected genotypes at different times of the day (Fig. 2C). P_N
increased for all the genotypes until mid-day (9.00 h to 12.00 h) and decline thereafter (12.00
h to 15.00 h) (Fig. 2C-I). However, the extent of the decline was lesser in wild rice species,
indicating that the wild species were able to maintain higher photosynthetic activity
throughout the day. The g_s also followed the similar pattern to that of P_N for the selected
species at different times of a day (Fig. 2C-II). Interestingly, E was found to be higher for the
wild species, with an associated decrease in WUE in the immediate afternoon (13.00-14.00
h), likely due to leaf and stomatal features (Fig. 2C, D-III).

Taken together, all these results showed that the selected wild rice species were
photosynthetically more efficient than cultivated rice varieties at different light intensities,
CO_2 concentrations, and time points of a day.

Leaf developmental traits underpinning photosynthetic variations

Leaves are the prime photosynthetic organs, and expectedly the selected rice
accessions exhibited striking differences in overall leaf architecture (Supplementary Fig. S1).
At the morphological level, cultivated and wild rice species showed considerable variation
for flag leaf width (FLW) and flag leaf thickness (FLT) (Supplementary Table S3). Wild rice
species, except *O. punctata* that showed lower P_N, had thicker and wider leaves. A strong
positive association of P_N were found with FLW (*r = 0.829*) and FLT (*r = 0.806)* (Fig. 3A).

At the anatomical level, remarkable differences in the vascular architecture and
mesophyll cell features were observed. Wild rice species, with higher net photosynthesis, had
larger and closely spaced veins with fewer number of mesophyll cells between two
consecutive veins (Fig. 3B, Supplementary Table S3, S4). Those wild species also had larger
mesophyll cell size, and thus lesser number of mesophyll cells per unit leaf area (Fig. 3C,
Supplementary Fig. S4A-B and Supplementary Table S4). We, then, quantified total

mesophyll cell surface area per unit leaf area of the selected accessions. Wild rice species
with higher P_N, *O. australiensis* and *O. latifolia*, showed remarkably higher mesophyll cell
surface area per unit leaf area as compared to cultivated varieties (Fig. 3D). We also
investigated the organization of chloroplasts in mesophyll cells of the representative wild and
cultivated leaf. Photosynthetically efficient wild rice exhibited larger and numerous
chloroplasts, systematically distributed along the mesophyll cell walls (Supplementary Fig.
S5). Furthermore, wild species with higher P_N exhibited a larger vein height and width (Fig.
3E and Supplementary Table S3). Nipponbare and Swarna that showed relatively higher P_N
compared to other cultivated varieties also showed larger vein height and width
(Supplementary Table S3). Swarna, further, had wider leaf and higher vein number. Pearson
correlation coefficient analysis showed a strong positive correlation of P_N with mesophyll
area ($r = 0.905$), mesophyll length ($r = 0.874$), mesophyll width ($r = 0.930$), and mesophyll
lobbing ($r = 0.771$) (Fig. 3A). The P_N was also strongly correlated with total number of veins
($r = 0.857$), minor veins height ($r = 0.906$), and minor vein width ($r = 0.758$) (Fig. 3A).
However, P_N was poorly correlated with bundle sheath features, number, and area (Fig. 3A).
352

Photochemical differences among the selected rice accessions

Most of the accessions with higher photosynthesis also had higher total chlorophyll
content (TCC; $r = 0.54$) (Fig. 4A). The *O. australiensis, O. latifolia* and *O. rufipogon* among
wild rice species, and *O. glaberrima*, *O. sativa* cv. *Swarna*, and *O. sativa* cv. *Nipponbare*
among cultivated rice had higher leaf chlorophyll content. The *O. punctata*, despite having
highest chlorophyll content, showed lower P_N, suggesting developmental and/or biochemical
constraints for efficient photosynthesis in the species. We, then, quantified chlorophyll
fluorescence (F_v/F_m) of dark-adapted leaves to determine the differences in quantum
photochemical efficiency of Photosystem II (PSII). Significant variations for F_v/F_m, which
ranged from 0.73 to 0.83, were found across the selected cultivated and wild rice species
(Fig. 4B). Wild rice species, *O. australiensis*, and *O. latifolia*, and *O. glaberrima* had higher
F_v/F_m, indicating the better quantum yield of PSII chemistry in those species. *O. latifolia* and
O. australiensis showed a significantly higher value of efficiency of PSII (□PSII),
photochemical quenching (qP), and electron transport rate (ETR) compared to other
accessions (Fig. 4C-E). *O. glaberrima*, the African cultivated rice with reasonably higher P_N,
also showed relatively higher □PSII, qP, and ETR than *O. sativa* cultivated varieties.
Cultivated accessions Nipponbare and Swarna didn't show significantly different □PSII, qP,
and ETR compared to IR64, despite showing relatively higher P_N, suggesting contribution of

anatomical features and/or efficiency of photoassimilation reaction towards higher P_N in the

two accessions (Fig. 1A, 4C-E). A strong correlation was observed between P_N with F_v/F_m (*r*

= 0.789), ΦPSII (*r =0.981*), qP (*r = 0.980*), and ETR (*r = 0.981*) (Supplementary Fig. S6).

These results showed that photochemical features of wild rice species, as well as *O.*

glaberrima, contribute to the higher leaf photosynthesis in those species.

Expression pattern of genes encoding core photosynthetic and chlorophyll biosynthesis

enzymes

To envisage any differences in photosynthetic gene expression across the genotype,

we checked the expression pattern of representative genes encoding key photosynthetic

components, LHC1 (Light harvesting chlorophyll protein complex), Rubisco LS (Large

Subunit), Rubisco SS (Small Subunit), and TPT (Triose phosphate translocator)

(Supplementary Table S5). The expression of the selected photosynthetic genes was

significantly higher in wild rice species (Fig. 5A), suggesting higher photosynthesis in those

species could, at least in part, be due to the higher abundance of these gene products. These

genes could possibly increase net photosynthetic efficiency by increasing the efficiency of

light capture, CO_2 assimilation, and transport of triose phosphate from chloroplast to

cytoplasm. Since wild rice species with higher photosynthesis also showed higher

chlorophyll content, we checked the expression level of chlorophyll biosynthesis genes. The

expression of chlorophyll biosynthesis genes, such as *GluTR*, *PPOX*, *Mg-CHLD*, *POR*, and

CHLSYN, were found to be significantly higher in *O. latifolia* and *O. australiensis* (Fig. 5B).

O. rufipogon also had significantly higher expression of selected photosynthetic and

chlorophyll-biosynthetic genes compared to Nipponbare and IR 64.

We then investigated the involvement of transcription factors in the regulation of

expression of the selected photosynthetic and chlorophyll-biosynthetic genes. A

comprehensive list of transcription factors regulating photosynthetic and chlorophyll

biosynthetic-genes has been reported in Wang *et al.* (2017). The differentially expressed

photosynthetic and chlorophyll biosynthetic-genes were hierarchically clustered with the

expression of potential transcription factors in rice using Genevestigator

(https://genevestigator.com/gv/). We found a number of transcription factors co-expressed

with the selected differentially expressed genes in flag leaves (Supplementary Fig. S7). We,

then, selected ten such transcription factors and profiled their gene expression at primordia 4

(P4) stage of leaf 4 (Supplementary Table S5). At least six of those ten genes, including

GLK1, *GLK2*, and *SIGMA* factors, were expressed at higher levels in wild rice with higher

leaf photosynthesis (Fig. 5C). *O. glaberrima* and *O. sativa cv. Nipponbare*, with moderately

high P_N, also showed higher expression of some of these transcription factors. Differential

expression of these transcription factors suggested that differences in the transcriptional

module regulating photosynthetic gene expression could be critical for determining leaf

photosynthetic efficiency.

Rubisco activity differences across the selected rice genotypes

The catalytic properties of Rubisco vary considerably from diverse plant sources,

suggesting that differences in the Rubisco activity might contribute to differences in

photosynthetic activity (Parry *et al.,* 2013). To test the possible Rubisco activity differences,

we quantified the RuBP-use and Rubisco activity (Vcmax) across selected cultivated and

wild rice species. The CO_2-saturated RuBP-use was found to be significantly higher for wild

rice species, *O. latifolia*, *O. australiensis*, and *O. rufipogon*, compared to *Oryza sativa cv.* IR

64 (Fig. 5D). Interestingly, *Oryza sativa cv. Nipponbare*, a *japonica* variety, also had higher

RuBP-use compared to *indica* variety IR 64. The photosynthesis rate per unit area, as

discussed earlier, was found to be higher in wild rice species as well as Nipponbare compared

to IR 64 (Fig. 1A). Regression analysis between P_N and Rubisco enzyme activity (Vcmax at

180 sec.) showed strong association between P_N and Vcmax ($R^2 = 0.835$, *P < 0.001*) (Fig.

5E). These results indicated the existence of variation in Rubisco carboxylation activity

within the rice system, cultivated and wild rice, which could possibly be exploited for

increasing CO_2 assimilation.

Discussion

The importance of investigating variations in leaf photosynthesis within and across

species to be exploited for sustainable increases in crop yield and productivity has been

emphasized (Ort *et al.,* 2015; Simkin *et al.,* 2019; van Bezouw *et al.,* 2019). A number of

studies have shown the variation in photosynthesis and related traits within species, such as

maize (Yabiku and Ueno, 2017), wheat (Driever *et al.,* 2014), rice (Giuliani *et al.,* 2013; Qu

et al., 2017), and even C_4 species *Gynandropsis gynandra* (Reeves *et al.,* 2018). Extensive

comparative genetic and genomic studies have been undertaken to investigate the variation in

photosynthetic efficiency across C_3 and C_4 species (Külahoglu *et al.,* 2014; Wang *et al.,*

2014). Based on these studies, global efforts to engineer C_4 photosynthesis in rice are being

made, which is challenging due to extensive genetic, developmental, and biochemical

differences between C_4 plants and rice. An alternative and efficient strategy would be

exploiting the photosynthetic variations within rice system itself, including cultivated and

wild rice. There had been a few studies highlighting the photosynthetic differences within

cultivated, and among cultivated and wild rice accessions (Kiran *et al., 2013;* Kondamudi *et*

al., 2016; Haritha *et al.,* 2017). In this study, we not only quantified leaf photosynthetic

differences across cultivated and wild rice accessions, but also identified the underlying

anatomical, biochemical, and photochemical traits mediating higher leaf photosynthesis in

rice. In addition, we evaluated the relationship between photosynthesis and other

physiological traits, such as gas exchange, carboxylation efficiency, transpiration, and water

use efficiency for the wild and cultivated rice accessions.

Wild relatives of rice, *Oryza australiensis*, *Oryza latifolia*, and *Oryza rufipogon*, as

well as African domesticated rice *Oryza glaberrima* showed higher leaf photosynthesis per

unit area (P_N) compared to cultivated varieties of *Oryza sativa* (Fig. 1A). Stomatal

conductance (g_s) is known to be an important factor determining the leaf photosynthesis rate

(Gago *et al.,* 2016). Two wild rice species, *O. australiensis* and *O. latifolia*, with highest P_N

also had highest values of g_s, which in turn resulted in higher internal CO_2 concentration (Ci)

in those species (Fig. 1B, C), supporting a strong correlation between P_N and g_s (Kanemura *et*

al., 2007; Giuliani *et al.,* 2013). *O. rufipogon* with reasonably higher P_N, however, had lower

g_s and Ci. The higher P_N in *O. rufipogon* was likely due to highest carboxylation efficiency

(CE) in the species among the selected accessions (Fig. 1F). Further, *O. australiensis* and *O.*

latifolia also had reasonably higher CE. The P_N has been shown to be positively associated

with CE in rice (Yeo *et al.,* 1994). Indeed, quantification of photosynthesis and related

physiological parameters for introgressions from *O. rufipogon* in a cultivated background

showed the influence of both CE as well as g_s on higher leaf photosynthesis (Haritha *et al.,*

2017). Furthermore, higher g_s and CE could also explain higher P_N across different PAR in

the wild rice (Fig. 2A-II and Supplementary Fig. S3A-III). Higher P_N in the wild species at

different times of a day, even in the afternoon when stomata tend to close, compared to

cultivated varieties could also be attributed to relatively higher g_s and CE in those species

(Fig. 2C and Supplementary Fig. S3C). Notably higher photosynthetic efficiency in the wild

rice species suggests that leaf photosynthesis in cultivated rice has not yet reached the

saturation level, and therefore, can possibly be achieved by targeting higher g_s and/or higher

CE. Moreover, increments in g_s might also help in optimizing photosynthetic performance

under environmental stress conditions. *O. australiensis*, interestingly, appeared to be the most

water-use efficient among the selected accessions due to high P_N coupled with low

transpiration rate (E), making it an ideal genotype to investigate for maintaining higher leaf

photosynthesis under drought conditions (Fig. 1A, D, and E). Many of the wild rice species,

including *O. australiensis*, have been marked for their potential usage towards tolerance to

heat and drought (Sanchez *et al.,* 2013; Menguer *et al.,* 2017). The differences in g_s and E are

likely related to differences in the stomatal features or regulation of stomatal opening across

the genotypes, as stomatal features are critical for photosynthesis, gas exchange, and water

use efficiency (Farquhar and Sharkey, 1982). Kondamudi *et al.* (2016) showed that leaves

with larger-sized stomata had higher photosynthesis rate than the leaves with smaller-sized

stomata due to higher g_s. Similarly, rice species with broader leaves have been shown to

exhibit higher P_N due to increased g_s than species with narrow leaves (Giuliani *et al.,* 2013).

Differences in internal CO_2 concentration and/or biochemical differences, such as Rubisco

activity, may mediate the observed differences in the CE (Eichelmann and Laisk, 2002).

Nonetheless, efforts to increase g_s and CE in the cultivated rice varieties comparable to the

high photosynthetic wild species shall help in increasing photosynthetic efficiency.

The strong correlation of leaf developmental traits with P_N is not surprising

considering the strong influence of leaf traits on light perception, gas exchange, CO_2

assimilation, and downstream photoassimilate partitioning. We observed a strong positive

association of P_N with leaf width and thickness (Fig. 3A). Leaf thickness has been suggested

to influence photosynthesis through efficient absorption and utilization of light by

homogenizing the leaf internal light distribution as well as maintaining leaf temperature

(Tholen *et al.,* 2012). The association of leaf width with photosynthetic performance and

yield in rice has been observed earlier. *WUSCHEL-RELATED HOMEOBOX* (*WOX*) genes

NAL1, *NAL2*, and *NAL3* have been shown to be associated with increased leaf width, which

in turn results in increased photosynthetic efficiency and yield (Fujita *et al.,* 2013; Ishiwata *et*

al., 2013). Introgression of a natural variant of *NAL1* from a *japonica* to *indica* cultivar led to

thicker and wider leaves with higher leaf photosynthesis rate and yield (Takai *et al.,* 2013).

Rice system also displays significant inter-species variation for anatomical traits,

suggesting strong genetic control over leaf anatomical features (Chatterjee *et al.,* 2016).

Photosynthetically efficient wild rice species have mesophyll cell features that promote both

higher accesses to available CO_2 as well as efficient light harvesting. Larger sized and lobed

mesophyll cells in wild rice species likely provides increased surface area to access more

internal CO_2 (Fig. 3C, D), whereas the organized pattern and systematic distribution of

chloroplasts in the mesophyll cells of those species may promote efficient light capture to

enhance photosynthesis (Supplementary Fig. S5 and Fig. 6). Interestingly, *NAL1* (discussed

above) promotes leaf width and thickness by increasing mesophyll cell number and area. An
increased vein density has been suggested to facilitate higher leaf photosynthesis by
promoting efficient export of photosynthates (Amiard *et al.,* 2005; Feldman *et al.,* 2017).
Lesser number of mesophyll cells between two consecutive veins in photosynthetically
efficient wild species, a feature similar to C_4 plants, should be promoting efficient transport of
photoassimilate. Larger vein size in the wild rice species shall, further, be adding to efficient
transport, thus increasing the leaf photosynthesis. The larger mesophyll cells with bigger
chloroplasts and larger veins in photosynthetically efficient wild rice species are reminiscent
of "Proto-kranz anatomy," an evolutionary step towards C_4 photosynthesis and have been
suggested to be an essential step towards increasing photosynthetic efficiency (Osborne and
Sack, 2012). Photosynthetically efficient wild rice species with desirable anatomical features
provide a potential genetic resource towards the identification of genes for desirable
manipulation in mesophyll and vein features.

Wild species with higher P_N showed significantly higher F_v/F_m compared to cultivated
varieties, indicating better quantum efficiency of Photosystem II in those species (Fig. 4B).
The F_v/F_m is used as the most common parameter for monitoring 'stress' in leaves as stress
results in inactivation damage of PSII by lowering of F_v/F_m (Murchie and Niyogi, 2011). The
comparative higher values for quantum efficiency of PSII ($\Box$PSII), photochemical quenching
(qP), and Electron Transport Rate (ETR) in the wild rice indicate PSII reaction centres are
more receptive for electron transport from PSII to PSI, making an efficient link between light
and light-independent reactions to increase P_N (Fig. 4C-E) (Kramer and Evans, 2011). A
positive correlation has been reported between chlorophyll fluorescence in leaf and grain
yield (von Kroff *et al.,* 2008). The wild species with better photochemical features and higher
P_N accumulates more biomass, and not grain yield, likely due to defects in photoassimilate
partitioning to the grain. Nonetheless, the desirable photochemical features, at least in part,
contribute to higher photosynthetic efficiency of the wild species. Gene expression studies
also showed higher expression of *LHC1* gene, which is involved in organization of reaction
centre, in photosynthetically efficient wild species.

Rubisco is the rate-limiting enzyme for photoassimilation of CO_2, and Rubisco
abundance and activity strongly determines the photosynthetic performance of a plant (Parry
et al., 2013). High photosynthetic wild species had higher expression of genes encoding
Rubisco large subunit and small subunit, indicating the higher abundance of Rubisco in those
species (Fig. 5A). Those wild species also had higher expression of chlorophyll biosynthetic
genes (Fig. 5B). The photosynthetic process as such, and genes encoding key components of

photosynthesis are under strong regulation of transcription factors (Wang *et al.,* 2017).

Differences in the expression of transcription factors, such as *GLKs* involved regulation of

chlorophyll biosynthesis and *SIGMA* factors regulating the chloroplastic gens *Rubisco LS*,

suggested the rewiring of transcriptional module regulating photosynthesis as a potential

target to increase rice leaf photosynthesis. Moreover, we also observed a higher RuBP-use

and Rubisco carboxylation activity (Vcmax) in photosynthetically efficient wild species (Fig.

5D, E). The prospects of improving Rubisco catalysis to increase photosynthetic efficiency

have been explored. It has been suggested that engineering maize Rubisco in rice would

improve CO_2 assimilation (Parry *et al.,* 2013; Sharwood *et al.,* 2016). This study suggests a

possible existence of an efficient Rubisco in photosynthetically efficient wild rice. Further

studies are needed to investigate whether Rubisco abundance or Rubisco kinetic properties

contribute to differences in Rubisco carboxylation activity across cultivated and wild rice.

In summary, we characterized photosynthetically efficient wild rice species for leaf

developmental, photochemical, and biochemical differences. Leaf developmental features

(larger and lobbed mesophyll cells, larger veins, and lesser number of mesophyll cells

between two consecutive veins), photochemical features (quantum efficiency of PSII,

Photochemical quenching, and Electron Transport Rate), and Rubisco activity contribute to

higher leaf photosynthesis in wild rice (Fig. 6). The study highlights the existence of

significant exploitable variation in leaf photosynthesis and underlying traits across cultivated

and wild rice. Future efforts to dissect the genetic and molecular basis of the underlying

developmental, photochemical, and biochemical traits shall be instrumental in tailoring

photosynthetically efficient rice.

Supplementary data

Fig. S1. Representative images of cultivated and wild rice species showing remarkable

differences in plant architectural traits, leaf features, and biomass. Shown are the field grown

plants (A), and controlled growth chambers grown plants (B) of the selected genotypes.

Fig. S2. Pair-wise Pearson correlation coefficient (*r*) analysis among net photosynthetic rate

(P_N) and related physiological traits.

Fig. S3. Quantification of P_N (max), Ci and CE in response to increasing PAR (A, I-III),

increasing CO_2 concentration (B, I-III), and at different diurnal time (C, I-III) for selected

cultivated and wild rice species.

Fig. S4. Quantification of average mesophyll cell area (A) and mesophyll cell number per

unit leaf area (cm^{-2}) (B) of the selected genotypes.

Fig. S5. TEM images of organization of chloroplasts in mesophyll cells.

Fig. S6. Pair-wise Pearson correlation coefficient (r) analysis among net photosynthetic rate

(P_N), chlorophyll fluorescence traits, and photochemical features.

Fig. S7. An *in silico* generated hierarchical clustering dendrogram for the expression pattern

of key photosynthetic and chlorophyll-biosynthetic genes, and transcription factor regulating

photosynthetic machinery across nine anatomical parts of rice plants (seedling, coleoptile,

seedling leaf, stem, vegetative leaf, blade, collar, sheath, flag leaf). The data were collected

and dendrograms was generated using Genevestigator (https://genevestigator.com/gv/).

Table S1. Details of rice accessions used to investigate the leaf photosynthesis and related

physiological features.

Table S2. Variation in leaf photosynthesis and related physiological traits among cultivated

and wild rice species under controlled conditions.

Table S3. Quantitative measurement of flag leaf width (FLW, cm), flag leaf thickness (FLT,

mm), Minor vein width (µm), Minor vein height (µm), and total veins (n) in the fully

expanded flag leaf of cultivated and wild rice species.

Table S4. Quantitative measurement of mesophyll lobes (n), length (µm), width (µm), and

area ($µm^2$) and bundle sheath number (n), and area ($µm^2$) in the fully expanded flag leaf of

cultivated and wild rice species.

Table S5. Details of genes and transcription factors used for gene expression analysis.

Table S6. List of primers for qRT-PCR analyses.

Acknowledgments

This work was supported by the core funding from National Institute of Plant Genome
Research as well as Ramalingaswamy Re-entry Fellowship (BT/RLF/Re-entry/05/2013) and
Innovative Young Biotechnologist Award (BT/09/IYBA/2015/01) from the Department of
Biotechnology, Ministry of Science and Technology, India. JM, AS, and VJ acknowledge
their CSIR-JRF, SERB-NPDF, and UGC-JRF, fellowships, respectively. Seeds of wild rice
species and *Oryza glaberrima* were kindly provided by Dr. Kuldeep Singh and Dr. Kumari
Neelam, Punjab Agricultural University, Ludhiana, India.

References

Alexandratos N, Bruinsma J. 2012. World agriculture towards 2030/2050, the 2012 revision. ESA Working paper, Rome, FAO, No. 12–03.

Amiard V, Mueh KE, Demmig-Adams B, Ebbert V, Turgeon R, Adams WW III. 2005. Anatomical and photosynthetic acclimation to the light environment in species with differing mechanisms of phloem loading. *Proceedings of the National Academy of Sciences, USA* **102**, 12968–12973.

Atwell BJ, Wang H, Scafaro AP. 2014. Could abiotic stress tolerance in wild relatives of rice be used to improve Oryza sativa? *Plant Science* **215–216**, 48–58.

Brodribb TJ, Feild TS, Jordan GJ. 2007. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiology* **144**, 1890–1898.

Cardona T, Shao S, Nixon PJ. 2018. Enhancing photosynthesis in plants, the light reactions. *Essays In Biochemistry* **62**, 85–94.

Chatterjee J, Dionora J, Elmido-Mabilangan A, Wanchana S, Thakur V, Bandyopadhyay A, Brar DS, Quick WP. 2016. The evolutionary basis of naturally diverse rice leaves anatomy. *PLoS ONE* **11**, 1–25.

Driever SM, Lawson T, Andralojc PJ, Raines CA, Parry MAJ. 2014. Natural variation in photosynthetic capacity, growth, and yield in 64 field-grown wheat genotypes. *Journal of Experimental Botany* **65**, 4959–4973.

Eichelmann H, Laisk A. 2002. Ribulose-1,5-bisphosphate carboxylase/oxygenase content, assimilatory charge, and mesophyll conductance in leaves. *Plant Physiology* **119**, 179–190.

Evans JR. 2013. Topical reviews on photosynthesis improvement, Improving photosynthesis. *Plant Physiology* **162**, 1780–1793.

Farquhar GD, Sharkey TD. 1982. Stomatal conductance and photosynthesis. Annu Rev *Plant Physiology* **33**, 317–345.

Feldman AB, Leung H, Baraoidan M, Elmido-mabilangan A. 2017. Increasing leaf vein density via mutagenesis in rice results in an enhanced rate of photosynthesis, smaller cell

sizes and can reduce interveinal mesophyll cell number. *Frontiers in Plant Science* **8**, 1–10.

Flood J, Harbinson J, Aarts MGM. 2011. Natural genetic variation in plant photosynthesis. *Trends in Plant Science* **16**, 327–335.

Fujita D, Trijatmiko KR, Tagle AG, *et al.* 2013. *NAL1* allele from a rice landrace greatly increases yield in modern indica cultivars. *Proceedings of the National Academy of Sciences, USA* **110**, 20431–20436.

Gago J, Daloso Dde M, Figueroa CM, Flexas J, Fernie AR, Nikoloski Z. 2016. Relationships of leaf net photosynthesis, stomatal conductance, and mesophyll conductance to primary metabolism, a multispecies meta-analysis approach. *Plant Physiology* **171**, 265–279.

Giuliani R, Koteyeva N, Voznesenskaya E, Evans MA, Cousins AB, Edwards GE. 2013. Coordination of leaf photosynthesis, transpiration, and structural traits in rice and wild relatives (Genus *Oryza*). *Plant Physiology* **162**, 1632–1651.

Haritha G, Vishnukiran T, Yugandhar P, Sarla N, Subrahmanyam D. 2017. Introgressions from *Oryza rufipogon* increase photosynthetic efficiency of KMR3 rice lines. *Rice Science* **24**, 85–96.

Hiyama A, Takemiya A, Munemasa S, Okuma E, Sugiyama N, Tada Y, Murata Y, Shimazaki KI. 2017. Blue light and CO_2 signals converge to regulate light-induced stomatal opening. *Nature Communications* **8**, 1284.

Huang M, Jiang L, Xia B, Zou Y, Jiang P, Ao H. 2013. Yield gap analysis of super hybrid rice between two subtropical environments. *Australian Journal of Crop Science* **7**, 600–608.

Ishiwata A, Ozawa M, Nagasaki H, et al. 2013. Two WUSCHEL-related homeobox genes, narrow leaf2 and narrow leaf3, control leaf width in rice. *Plant and Cell Physiology* **54**, 779–792.

Ivanova LA, Yudina PK, Ronzhina DA, Ivanov LA, Hölzel N. 2018. Quantitative mesophyll parameters rather than whole-leaf traits predict response of C_3 steppe plants to aridity. *New Phytologist* **217**, 558–570.

Jansson C, Vogel J, Hazen S, Brutnell T, Mockler T. 2018. Climate-smart crops with enhanced photosynthesis. *Journal of Experimental Botany* **69**, 3801–3809.

Kajala K, Covshoff S, Karki S, *et al.* 2011. Strategies for engineering a two-celled C_4 photosynthetic pathway into rice. *Journal of Experimental Botany* **62**, 3001–3010.

Kalaji HM, Jajoo A, Oukarroum A, Brestic M, Zivcak M, Samborska IA, Cetner MD, Łukasik I, Goltsev V, Ladle RJ. 2016. Chlorophyll a fluorescence as a tool to monitor physiological status of plants under abiotic stress conditions. *Acta Physiologiae Plantarum* **38**, 102.

Kanemura T, Homma K, Ohsumi A, Shiraiwa T, Horie T. 2007. Evaluation of genotypic variation in leaf photosynthetic rate and its associated factors by using rice diversity research set of germplasm. *Photosynthesis Research* **94**, 23–30.

Kindgren P, Norén L, Barajas López JDD, Shaikhali J, Strand Å. 2012. Interplay between HEAT SHOCK PROTEIN 90 and HY5 controls PhANG expression in response to the GUN5 plastid signal. *Molecular Plant* **5**, 901–913.

Kiran TV, Rao YV, Subrahmanyam D, Rani NS, Bhadana VP, Rao PR. 2013. Variation in leaf photosynthetic characteristics in wild rice species. *Photosynthetica* **51**, 350–358.

Kondamudi R, Swamy KN, Rao YV, *et al.* 2016. Gas exchange, carbon balance and stomatal traits in wild and cultivated rice (*Oryza sativa L.*) genotypes. *Acta Physiologiae Plantarum* **38**, 160.

Koteyeva NK, Voznesenskaya EV, Berry JO, Chuong SDX, Franceschi VR, Edwards GE. 2011. Development of structural and biochemical characteristics of C_4 photosynthesis in two types of Kranz anatomy in genus Suaeda (family Chenopodiaceae). *Journal of Experimental Botany* **62**, 3197–3212.

Kramer DM, Evans JR. 2011. The importance of energy balance in improving photosynthetic productivity. *Plant Physiology* **155**, 70–78.

Kubien DS, Brown CM, Kane HJ. 2011. Quantifying the amount and activity of rubisco in leaves. *Methods in Molecular Biology* **684**, 349–362.

Külahoglu C, Denton AK, Sommer M, *et al.* 2014. Comparative transcriptome atlases reveal altered gene expression modules between two Cleomaceae C_3 and C_4 plant species. *The Plant Cell* **26**, 3243-60.

Lawson T, Kramer DM, Raines CA. 2012. Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. *Current Opinion in Biotechnology* **23**, 215–220.

Liu L, Sun G, Ren X, Li C, Sun D. 2015. Identification of QTL underlying physiological and morphological traits of flag leaf in barley. BMC Genetics **16**, 29.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **25**, 402–408.

Long SP, Zhu X, Naidu SL, Ort DR. 2006. Can improvement in photosynthesis increase crop yields ? Plant, Cell & Environment **29**, 315–330.

Lux A, Morita S, Abe J, Ito K. 2005. An improved method for clearing and staining free-hand sections and whole-mount samples. *Annals of Botany* **96**, 989–996.

Mathan J, Bhattacharya J, Ranjan A. 2016. Enhancing crop yield by optimizing plant developmental features. *Development* **143**, 3283–3294.

Menguer PK, Sperotto RA, Ricachenevsky FK. 2017. A walk on the wild side, Oryza species as source for rice abiotic stress tolerance. *Genetics and Molecular Biology* **40**, 238–252.

Murchie EH, Niyogi KK. 2011. Manipulation of Photoprotection to Improve Plant Photosynthesis. *Plant Physiology* **155**, 86–92.

Murchie EH, Lawson T. 2013. Chlorophyll fluorescence analysis , a guide to good practice and understanding some new applications. *Journal of Experimental Botany* **64**, 3983–3998.

Ort DR, Merchant SS, Alric J, *et al.* 2015. Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proceedings of the National Academy of Sciences, USA* **112**, 8529–8536.

Osborne CP, Sack L. 2012. Evolution of C_4 plants, A new hypothesis for an interaction of CO_2 and water relations mediated by plant hydraulics. *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**, 583–600.

Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM. 2013. Rubisco activity and regulation as targets for crop improvement. *Journal of Experimental Botany* **64**, 717–730.

Qu M, Zheng G, Hamdani S, Essemine J, Song Q, Wang H. 2017. Leaf photosynthetic parameters related to biomass accumulation in a global rice diversity survey. *Plant Physiology* **175**, 248–258.

Reeves G, Grangé-Guermente MJ, Hibberd JM. 2017. Regulatory gateways for cell-specific gene expression in C_4 leaves with Kranz anatomy. *Journal of Experimental Botany* **68**, 107-116.

Reeves G, Singh P, Rossberg TA, Sogbohossou EOD, Schranz ME, Hibberd JM. 2018. Natural variation within a species for traits underpinning C_4 photosynthesis. *Plant Physiology* **177**, 504–512.

Reynolds ES. 1963. The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *Journal of Cell Biology* **17**, 208–212.

Rogers A, Allen DJ, Davey PA, *et al.* 2004. Leaf photosynthesis and carbohydrate dynamics of soybeans grown throughout their life-cycle under Free-Air Carbon dioxide Enrichment. *Plant, Cell & Environment* **27**, 449–458.

Sack L, Scoffoni C. 2013. Leaf venation, structure, function, development, evolution, ecology and applications in the past, present and future. *New Phytologist* **198**, 983–1000.

Sage TL, Sage RF. 2009. The functional anatomy of rice leaves, implications for refixation of photorespiratory CO_2 and efforts to engineer C_4 photosynthesis into rice. *Plant and Cell Physiology* **50**, 756–772.

Sanchez PL, Wing RA, Brar DS. 2013. The wild relative of Rice, Genomes and genomics. In, Zhang Q, Wing RA (eds) Genetics and genomics of rice, plant genetics and genomics, crops and models 5. Springer, New York, 9–25.

Scales JC, Parry MAJ, Salvucci ME. 2014. A non-radioactive method for measuring Rubisco activase activity in the presence of variable ATP: ADP ratios , including modifications for measuring the activity and activation state of Rubisco. *Photosynthsis Research* **119**, 355–365.

Schuler ML, Mantegazza O, Weber APM. 2016. Engineering C_4 photosynthesis into C_3 chassis in the synthetic biology age. *Plant J* **87**, 51–65.

Sharwood RE, Ghannoum O, Whitney SM. 2016. Prospects for improving CO_2 fixation in C_3-crops through understanding C_4-Rubisco biogenesis and catalytic diversity. *Current Opinion in Plant Biology* **31**, 135–142.

Simkin AJ, López-calcagno PE, Raines CA. 2019. Feeding the world, improving photosynthetic efficiency for sustainable crop production. *Journal of Experimental Botany* **70**, 1119–1140.

Takai T, Adachi S, Taguchi-shiobara F, *et al.* 2013. A natural variant of NAL1, selected in high-yield rice breeding programs, pleiotropically increases photosynthesis rate. *Scientific Reports* **3**, 1–11.

Tholen D, Ethier G, Genty B, Pepin S, Zhu XG. 2012. Variable mesophyll conductance revisited, theoretical background and experimental implications. *Plant, Cell & Environment* **35**, 2087– 2103.

Toledo-Ortiz G, Johansson H, Lee KP, Bou-Torrent J, Stewart K, Steel G, Rodríguez-Concepción M, Halliday KJ. 2014. The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. *PLoS Genetics* **10**, e1004416.

van Bezouw RFHM, Keurentjes JJB, Harbinson J, Aarts MGM. 2019. Converging phenomics and genomics to study natural variation in plant photosynthetic efficiency. *Plant J* **97**, 112-133.

von Caemmerer S, Quick WP, Furbank RT. 2012. The development of C$_4$ rice, Current progress and future challenges. *Science* **336**, 1671-1672.

van Campen JC, Yaapar MN, Narawatthana S, *et al.* 2016. Combined chlorophyll fluorescence and transcriptomic analysis identifies the P3/P4 transition as a key stage in rice leaf photosynthetic development. *Plant Physiology* **170**, 1655–1674.

von Korff M, Grando S, This D, Baum M, Ceccarelli S. 2008. Quantitative trait loci (QTL) associated with agronomic performance of barley under drought. *Theoretical and Applied Genetics* **117**, 653–669.

Wang L, Czedik-Eysenberg A, Mertz RA, *et al.* 2014. Comparative analyses of C$_4$ and C$_3$ photosynthesis in developing leaves of maize and rice. *Nature Biotechnology* **32**, 1158-65.

Wang P, Hendron R, Kelly S. 2017. Transcriptional control of photosynthetic capacity, conservation and divergence from Arabidopsis to rice. *New Phytologist* **216**, 32–45.

Waters MT, Wang P, Korkaric M, Capper RG, Saunders NJ, Langdale JA. 2009. GLK transcription factors coordinate expression of the photosynthetic apparatus in Arabidopsis. *The Plant Cell* **21**, 1109–1128.

Wing RA, Purugganan MD, Zhang Q. 2018. The rice genome revolution: from an ancient grain to Green Super Rice. *Nature Reviews Genetics* **19**, 505-517.

Wu A, Hammer GL, Doherty A, von Caemmerer S, Farquhar GD. 2019. Quantifying impacts of enhancing photosynthesis on crop yield. *Nature Plants* **5**, 380-388.

Yabiku T, Ueno O. 2017. Variations in physiological, biochemical, and structural traits of photosynthesis and resource use efficiency in maize and teosintes (NADP-ME-type C$_4$). *Plant Production Science* **20**, 448–458.

Yeo ME, Yeo AR, Flowers TJ. 1994. Photosynthesis and photorespiration in the genus Oryza. *Journal of Experimental Botany* **45**, 553–560.

Zhao M, Lafitte HR, Sacks E, Dimayuga G, Acun TLB. 2008. Perennial *O. sativa* $\times$ *O. rufipogon* interspecific hybrids, I. Photosynthetic characteristics and their inheritance. *Field Crops Research* **106**, 203–213.

Zhao M, Ding Z, Lafitte R, Sacks E, Dimayuga G, Holt D. 2010. Photosynthetic characteristics in *Oryza* species. *Photosynthetica* **48**, 234–240.

Figure Legends

Fig. 1. Variations in leaf photosynthesis and related physiological parameters across cultivated and wild rice species. Quantification of net photosynthesis per unit area of flag leaf surface area (P_N, A), stomatal conductance (g_s, B), intercellular CO_2 concentration (Ci, C), transpiration rate (E, D), water use efficiency (WUE, E), and carboxylation efficiency (CE, F) for six cultivated rice accessions and four wild rice species. Cultivated rice accessions and wild species are represented with blue and green shading box plots, respectively. Asterisks indicate significant differences between genotypes (one-way ANOVA, n = 15, *P < 0.05*). Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Fig. 2. Effects of PAR intensity, CO_2 concentration, and diurnal time on leaf photosynthesis (P_N), stomatal conductance (g_s), transpiration (E), and water use efficiency (WUE) of different rice genotypes.

(A) Quantification of P_N (I), g_s (II), E (III), and WUE (IV) in response to increasing PAR from 0 to 1500 $\mu mol\, m^2\, s^{-1}$.

(B) Quantification of P_N (I), g_s (II), E (III), and WUE (IV) in response to increasing CO_2 concentration from 0 to 1400 $\mu mol\, mol^{-1}$.

(C) Quantification of P_N (I), g_s (II), E (III), and WUE (IV) at different time points of a day from 9.00 h to 15.00 h

Each value represents the mean of 5 data points ± SD from different plants.

Fig. 3. Leaf anatomical traits effecting photosynthesis rate and related physiological traits of rice.

(A) Pearson correlation analysis of net photosynthesis rate (P_N) and related physiological traits with leaf developmental features. The level of significance (α) of correlation coefficients at *P < 0.05 (± 0.666; degree of freedom (df; n-2): 9-2 = 7).

(B) Bright-field images of a flag leaf of the selected genotypes showing variations in mesophyll cells and vasculature. Arrows indicate bulliform cells (bc), bundle sheath (bs),

epidermis (ep), mesophyll cells (mc), and minor veins (miv). Scale bars represent 100 µm at 10X magnification.

(C) Bright-field microscopy images showing variations in mesophyll cell features, including lobbing. Scale bar represents 100 µm at 100X magnification.

(D-E) Quantification of total mesophyll cell surface area per unit leaf area (D) and minor vein height (E) of selected cultivated and wild rice species. Cultivated rice accessions and wild species are represented with blue and green shading box plots, respectively. Asterisks indicate significant differences between genotypes [one-way ANOVA, n = 4 leaves (10 images per leaf), $*P < 0.05$]. Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Fig. 4. Quantification of photochemical features of selected cultivated and wild rice species. Total chlorophyll content (TCC) per gram of fresh weight (A), F_v/F_m in dark-adapted leaves (B), Efficiency of PSII (ΦPSII, C), Photochemical quenching (qP, D), and Electron transport rate (ETR, E) of the selected genotypes. Cultivated rice accessions and wild species are represented with blue and green shading box plots, respectively. All the measurements were performed in fully developed vegetative stage leaves of controlled growth chambers grown plants. Asterisks indicate significant differences between genotypes (one-way ANOVA, n = 7, $*P < 0.05$). Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Fig. 5. Expression analysis of genes and transcription factors regulating photosynthesis, and differences in Rubisco activity for selected cultivated and wild rice species.

(A) Expression levels of genes encoding key component of light harvesting complex, Rubisco, and Triose phosphate/phosphate translocator.

(B) Expression levels of chlorophyll biosynthetic genes in flag leaves.

(C) Expression pattern of transcription factors regulating photosynthetic and chlorophyll-biosynthetic genes in primordia 4 stage (P4) of leaf 4.

(D) Rate of RuBP-use (ribulose-1, 5-bisphosphate) (nmol s^{-1}) from the conversion of RuBP to 3-PGA (3-phosphoglycerate). The rate of 3-PGA production was determined from oxidation of NADH during assays by continuously monitoring of decrease in absorbance at 340 nm.

(E) Regression analysis between P_N and Vcmax at 180 sec of the assays (µmol m^2 s^{-1}) generated from the maximum rate of RuBP-use of the selected genotypes.

Asterisks indicate significant differences between genotypes [one-way ANOVA, n = 6 (two biological with three technical replications, *P < 0.05, **P < 0.01, ***P < 0.001]*. Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Gene abbreviations: GluTR, *Glutamyl-tRNA reductase*; PPOX, *Protoporphyrinogen oxidase*; Mg-CHL, *Magnesium chelatase*; CHLSYN, *Chlorophyll synthase*; CAO, *Chlorophyllide a oxygenase;* LHCI, *Light-harvesting chlorophyll protein complex I*; Rubisco SS, *Ribulose bisphosphate carboxylase small subunit*; Rubisco LS, *Ribulose bisphosphate carboxylase large subunit*; TPT, *Triose phosphate/phosphate translocator*; GLK, *Golden like* (1 and 2); SIG, *Sigma factors* (1, 2, 5, and 6); ZML2, *ZINC FINGER PROTEIN 2*; ORE1, *ORESARA1*; CGA1, *Cytokinin responsive GATA Factor*; GRF5, *Growth regulating factor 5*.

Fig. 6. Compilation of developmental, photochemical, and biochemical traits contributing towards leaf photosynthetic differences between two extreme genotypes: *Oryza sativa* ssp. *indica* cv. IR 64 and *Oryza australiensis*. Wider leaf, larger mesophyll cells, larger veins, fewer number of mesophyll cells between two consecutive veins, more chloroplasts, efficient light harvesting and Electron Transport Rate along with more active Rubsico facilitate higher leaf photosynthesis in wild rice species *Oryza australiensis* compared to cultivated *Oryza sativa* ssp. *indica* cv. IR64.

Fig. 1. Variations in leaf photosynthesis and related physiological parameters across cultivated and wild rice species. Quantification of net photosynthesis per unit area of flag leaf surface area (P_N, A), stomatal conductance (g_s, B), intercellular CO_2 concentration (Ci, C), transpiration rate (E, D), water use efficiency (WUE, E), and carboxylation efficiency (CE, F) for six cultivated rice accessions and four wild rice species. Cultivated rice accessions and wild species are represented with blue and green shading box plots, respectively. Asterisks indicate significant differences between genotypes (one-way ANOVA, n = 15, *$P < 0.05$*). Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Fig. 2. Effects of PAR intensity, CO_2 concentration, and diurnal time on leaf photosynthesis (P_N), stomatal conductance (g_s), transpiration (E), and water use efficiency (WUE) of different rice genotypes.
(A) Quantification of P_N (I), g_s (II), E (III), and WUE (IV) in response to increasing PAR from 0 to 1500 µmol m² s⁻¹.
(B) Quantification of P_N (I), g_s (II), E (III), and WUE (IV) in response to increasing CO_2 concentration from 0 to 1400 µmol mol⁻¹.
(C) Quantification of P_N (I), g_s (II), E (III), and WUE (IV) at different time points of a day from 9.00 h to 15.00 h
Each value represents the mean of 5 data points ± SD from different plants.

Fig. 3. Leaf anatomical traits effecting photosynthesis rate and related physiological traits of rice.
(A) Pearson correlation analysis of net photosynthesis rate (P_N) and related physiological traits with leaf developmental features. The level of significance (α) of correlation coefficients at *$P < 0.05$ (± 0.666; degree of freedom (df; n-2): 9-2 = 7).
(B) Bright-field images of a flag leaf of the selected genotypes showing variations in mesophyll cells and vasculature. Arrows indicate bulliform cells (bc), bundle sheath (bs), epidermis (ep), mesophyll cells (mc), and minor veins (miv). Scale bars represent 100 μm at 10X magnification.
(C) Bright-field microscopy images showing variations in mesophyll cell features, including lobbing. Scale bar represents 100 μm at 100X magnification.
(D-E) Quantification of total mesophyll cell surface area per unit leaf area (D) and minor vein height (E) of selected cultivated and wild rice species. Cultivated rice accessions and wild species are represented with blue and green shading box plots, respectively. Asterisks indicate significant differences between genotypes [one-way ANOVA, n = 4 leaves (10 images per leaf), *$P < 0.05$]. Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Fig. 4. Quantification of photochemical features of selected cultivated and wild rice species. Total chlorophyll content (TCC) per gram of fresh weight (A), F_v/F_m in dark-adapted leaves (B), Efficiency of PSII (ΦPSII, C), Photochemical quenching (qP, D), and Electron transport rate (ETR, E) of the selected genotypes. Cultivated rice accessions and wild species are represented with blue and green shading box plots, respectively. All the measurements were performed in fully developed vegetative stage leaves of controlled growth chambers grown plants. Asterisks indicate significant differences between genotypes (one-way ANOVA, n = 7, *P < 0.05). Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Fig. 5. Expression analysis of genes and transcription factors regulating photosynthesis, and differences in Rubisco activity for selected cultivated and wild rice species.

(A) Expression levels of genes encoding key component of light harvesting complex, Rubisco, and Triose phosphate/phosphate translocator.

(B) Expression levels of chlorophyll biosynthetic genes in flag leaves.

(C) Expression pattern of transcription factors regulating photosynthetic and chlorophyll-biosynthetic genes in primordia 4 stage (P4) of leaf 4.

(D) Rate of RuBP-use (ribulose-1, 5-bisphosphate) (nmol s^{-1}) from the conversion of RuBP to 3-PGA (3-phosphoglycerate). The rate of 3-PGA production was determined from oxidation of NADH during assays by continuously monitoring of decrease in absorbance at 340 nm.

(E) Regression analysis between P_N and Vcmax at 180 sec of the assays (µmol m^2 s^{-1}) generated from the maximum rate of RuBP-use of the selected genotypes.

Asterisks indicate significant differences between genotypes [one-way ANOVA, n = 6 (two biological with three technical replications, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$]. Different letters indicate significant groupings according to post-hoc Tukey HSD calculation.

Gene abbreviations: GluTR, *Glutamyl-tRNA reductase*; PPOX, *Protoporphyrinogen oxidase*; Mg-CHL, *Magnesium chelatase*; CHLSYN, *Chlorophyll synthase*; CAO, *Chlorophyllide a oxygenase;* LHCI, *Light-harvesting chlorophyll protein complex I*; Rubisco SS, *Ribulose bisphosphate carboxylase small subunit*; Rubisco LS, *Ribulose bisphosphate carboxylase large subunit*; TPT, *Triose phosphate/phosphate translocator*; GLK, *Golden like* (1 and 2); SIG, *Sigma factors* (1, 2, 5, and 6); ZML2, *ZINC FINGER PROTEIN 2*; ORE1, *ORESARA1*; CGA1, *Cytokinin responsive GATA Factor*; GRF5, *Growth regulating factor 5*.

Fig. 6. Compilation of developmental, photochemical, and biochemical traits contributing towards leaf photosynthetic differences between two extreme genotypes: *Oryza sativa* ssp. *indica* cv. IR 64 and *Oryza australiensis*. Wider leaf, larger mesophyll cells, larger veins, fewer number of mesophyll cells between two consecutive veins, more chloroplasts, efficient light harvesting and Electron Transport Rate along with more active Rubsico facilitate higher leaf photosynthesis in wild rice species *Oryza australiensis* compared to cultivated *Oryza sativa* ssp. *indica* cv. IR64.