Cariogenic *Streptococcus mutans* produces strain-specific

antibiotics that impair commensal colonization

Xiaoyu Tang[a,b]*, Yuta Kudo[b]*, Jonathon Baker[a], Sandra LaBonte[a,c], Peter A. Jordan[b], Shaun M.

K. McKinnie[b], Jian Guo[d], Tao Huan[d], Bradley S. Moore[b,e] and Anna Edlund[a]

[a]Genomic Medicine Group, J. Craig Venter Institute, La Jolla, California, USA. [b]Scripps institution

of oceanography, University of California, La Jolla, California, USA. [c]Department of Biochemistry

and Biophysics, Texas A&M University and Texas AgriLife Research, College Station, Texas,

USA. [d]Department of Chemistry, University of British Columbia, Vancouver, Canada. [e]Skaggs

School of Pharmacy and Pharmaceutical Sciences, University of California, La Jolla, California,

USA. *These authors contributed equally to this work.

Address correspondence to Bradley S. Moore, bsmoore@ucsd.edu or Anna Edlund

aedlund@jcvi.org

Streptococcus mutans is a common constituent of dental plaque and an etiologic agent of dental caries (tooth decay). Here we elucidate a biosynthetic pathway, encoded by globally distributed strains of *S. mutans,* which produces a series of bioactive small molecules including reutericyclin and two *N*-acyl tetramic acid analogues active against oral commensal bacteria. This pathway may provide *S. mutans* with a competitive advantage, promoting dysbiosis and caries pathogenesis.

The human microbiota consists of trillions of symbiotic microbial cells that not only help its host to digest dietary components(1-3), metabolize drugs(4-6), regulate immune system(7-9) but also produce complex small molecules such as antimicrobial nonribosomal peptides (NRPs) and polyketides (PKs)(10-12). Recent computational mining efforts of genomes and metagenomes of the human microbiome revealed a vast diversity (~14,000) of putative biosynthetic gene clusters (BGCs) encoding small molecules across all human body sites(13), of which many represented NRP, PK and hybrid NRP-PK small molecules. Typically, a typical gut harbors 599 BGCs, while a typical oral cavity harbors 1061 BGC(13). Thus far, most research efforts have focused on characterization of BGCs and small molecules from the gut microbiome, leaving large knowledge gaps of crucial signaling molecules of the oral cavity.

The oral cavity harbors a high species diversity with over 700 bacterial species, which mainly colonize four physically distinct niches including dental plaque, tongue dorsum, buccal mucosa, and saliva(14). Residents of the dental plaque have been implicated in a variety of diseases, including dental caries, which affects more than a third of the world's population and results in approximately $300 billion in direct treatment costs to the global economy annually(15-17). Although caries is a polymicrobial disease caused by a dysbiosis in the dental plaque microbial community, *Streptococcus mutans*, with its copious acid production and prodigious biofilm formation, is still considered a primary etiologic agent(18, 19). To persist in the dental

plaque community and cause disease, *S. mutans* must be able to outcompete commensal

bacteria directly.

Small molecules produced by BGCs are increasingly recognized to play major roles in

species-species communication and interactions(10, 13), and a recent study predicted 355 strain-

specific BGCs across 169 *S. mutans* genomes(20). Although the production of mutactins (a group

of bacteriocins) has been recognized for contribution to the colonization and establishment of *S.*

mutans in the dental biofilm(21), the roles of other genetically encoded small molecules in *S.*

mutans is barely explored, with exception of the mutanobactins. Mutanobactins are compounds

of hybrid polyketide synthase and nonribosomal peptide synthetase (PKS/NRPS) origin that

inhibit the morphological transition of *Candida albicans*(22). Previous bioinformatics efforts

identified an orphan hybrid PKS/NRPS BGC (recently designated *muc*)(20, 23), that is distributed

among a subset of *S. mutans* strains. Within *muc*, five biosynthetic proteins are highly

homologous (48%-69%) to cognates involved in the biosynthesis of reutericyclin (RTC)(24)

(**Supplementary Fig. 1** and **Fig. 1a**). RTC, which originated from the sourdough isolates of

Lactobacillus reuteri, acts as a proton ionophore antibiotic that modulates the microbial

community of sourdough(25, 26). Interestingly, we found that *S. mutans* strains encoding *muc*

were dispersed geographically and frequently associated with severe dental caries

(**Supplementary Table 1**). The goal of this study was to determine whether *muc* produces RTC

or RTC-like molecules, and if these molecules can affect inter-species competition in the oral

cavity.

To determine the product of *muc*, we utilized homologous recombination to delete the

gene *mucD*, which encodes an assembly line NRPS tridomain protein (**Fig. 1a**). We conducted

all our biosynthetic experiments in *S. mutans* B04Sm5, a strain bearing *muc*, which was isolated

from a child with severe early childhood caries(27). The wild type (WT) strain and Δ*mucD* mutant

were cultured and extracted for HPLC analyses. The results showed that the WT strain produced

four metabolites not present in the Δ*mucD* mutant (**Fig. 1b**). These molecules were purified via

preparative HPLC (**Supplementary Note 1**) and characterized by High-Resolution Mass Spectrometry (HR-MS/MS), chemical synthesis, and/or one- and two-dimensional Nuclear Magnetic Resonance (NMR) experiments to yield a group of tetramic acids, including RTC (renamed RTC A, **1**), two new RTC analogues (RTC B (**2**) and RTC C (**3**)), and a new organic acid (**4**) (**Fig. 1c**, **Supplementary Note 1**, **Supplementary Tables 2-3**, **Supplementary Figs. 2-22**). During preparation of this manuscript, Chen and colleagues published the identification of **4** (designated mutanocyclin (MUC)) using a new heterologous genetic system(23); however, production of **1**-**3** in both heterologous host and wild type producers was not reported.

Muc is a ~13 kb hybrid NRPS-PKS pathway encoding nine proteins (**Supplementary Table 5**). *In silico* analysis revealed that *mucD* to *mucE* encode the core assembly line protein machinery (**Fig. 1a** and **Fig. 2a**). While MucD is a C-A-T tridomain protein, with specificity for adenylating leucine, MucE contains an KS-T-TE module, as commonly present in the termination modules of PKS assembly lines. Based on the enzymatic logic of thiotemplate-mediated assembly line biosynthesis, we propose that **1**-**3** are assembled, respectively, from *trans*-2-decenoyl-ACP, decenoyl-ACP, and *trans*-2-dodecenoyl-ACP starter units through elongation with leucine, followed by elongation with a malonyl-CoA extender unit (**Fig. 2a**). Inspection of structures of **1**-**3** suggested that the A domain of MucD appears to install a *D*-leucine residue into the final product. To explore this hypothesis, we fed both [$^{13}C_1$] *L*- and *D*-leucine to cultures of *S. mutans* B04Sm5. MS analyses of the purified **1** and **3** only revealed the incorporation of [$^{13}C_1$] *L*-leucine (**Supplementary Figs. 23-24**). The same result was observed by feeding the original RTC producer *L. reuteri* with the same isomers (**Supplementary Fig. 25**). These results indicate that an unrecognized epimerization reaction is involved in **1**-**3** biosynthesis; however, no standard epimerization (E) domain or dual functioning C/E domains could be found either in the assembly line or encoded elsewhere in the BGC. Additionally, although a dual-function TE/E domain has been characterized from the nocardicin (NOC) biosynthetic assembly line(28), MucTE shows very low homology (20%/35%, identity/similarity) to the dual functioning NocTE domain.

The first three genes (*mucA-C*) encode a hydroxymethylglutaryl-CoA synthase (MucA), a thiolase (MucB), and a hypothetical protein (MucC) (**Supplementary Table 5**), which show homology to the three submits (PhlA, PhlB and PhlC, respectively) of a multicomponent C-acetyltransferase involved in the acetylation of the type III PKS product phloroglucinol from *Pseudomonas fluorescens* Q2-87(29, 30). As the combination of the three genes was also identified in *rtc* (*rtcA*, *rtcC* and *rtcB*) from *L. reuteri* (24), the function of MucA–C is consistent with introducing the acetyl group to the pyrrolidine ring of **1**-**3** (**Fig. 2a**). We additionally annotated four genes downstream of the *mucA-E* operon encoding a small HXXEE domain-containing membrane-protein (MucF) of unknown function, two TetR/AcrR family transcriptional regulators (MucG and MucH), and one multidrug efflux pump (MucI) (**Supplementary Table 5**). Presumably, they are not involved in the direct synthesis of **1**-**3**. To verify this hypothesis, we cloned the operon from *mucA* to *mucE* into the pACYCDuet-1 vector to generate the plasmid pEXT06, in which the operon is exclusively under control of a T_7 promoter. As expected, expression of *mucA-E* in *Escherichia coli* BAP1 strain resulted in the production of at least four products, including **1**-**3** and new compound **5** (**Fig. 2b** and **Supplementary Fig. 26**). **5** was purified via preparative HPLC and its structure was further confirmed as a new RTC analogue (RTC D) possessing a saturated C-12 fatty acid side chain by MS and NMR analyses (**Supplementary Table 4**, **Supplementary Fig. 26** and **Supplementary Figs. 27-30**). This result indicates that the first six genes *mucA-E* indeed compose the minimal BGC for **1**–**3** production (**Fig. 2a**).

As the structure of **4** is consistent with the RTC core lacking the fatty acyl chain, we first set out to detect whether the free fatty acid, *trans*-2-decenoic acid per compound **1**, is present in the extract of *S. mutans* B04Sm5. HPLC analyses confirmed that *S. mutans* B04Sm5 readily produced *trans*-2-decenoic acid (**Supplementary Fig. 31**). In contrast, it was not detected in the pathway-deficient mutant *S. mutans* B04Sm5/Δ*mucD*. These findings suggested that **4** may be derived from **1**-**3** via deacylation by an unknown enzyme. Interestingly, *trans*-2-decenoic acid is a known *Streptococcus* diffusible signal factor (SDSF) isolated from many *Streptococcus*

species(31), which inhibits the hyphal formation of the opportunistic fungus *Candida albicans*.

Among the annotated pathway enzymes, only the function of MucF was unassigned. The MucF

protein sequence was subjected to a secondary structure prediction-based homology search

(Phyre2), which suggested it is a polytopic (five) transmembrane α-helical protein

(**Supplementary Fig. 32**) with low similarity (15%) to a viral protein (PDB 3LD1) with putative

hydrolase activity. To explore whether MucF might be involved in the deacylation of **1**-**3**, we

generated a *mucF* deletion mutant in *S. mutans* B04Sm5. HPLC analysis of the extract of mutant

cultures showed that the Δ*mucF* mutant not only increased the production of **1**-**3** by ~3-5 fold, but

also lost the ability to produce **4** (**Fig. 1b**). These findings strongly suggested that MucF is

essential for converting **1**-**3** to **4**. To further evaluate the function of MucF, we cloned and

expressed *mucF* in *E. coli* and incubated **1** with the *E. coli*/*mucF* cell lysate, leading to the *in vitro*

production of the deacylated **4** (**Fig. 2c**). In contrast, no conversation was detected in the control

experiment (*E. coli* carrying empty vector) (**Fig. 2c**). To further support this observation, we

inserted a copy of *mucF* into the secondary expression site of pEXT06, resulting in the plasmid

pEXT07. Its expression in *E. coli* BAP1 further led to the formation of **4** (**Fig. 2b**). Collectively,

these mutations, *in vitro*, and *in vivo* expression studies support the hypothesis that MucF is a

newly recognized deacylase responsible for converting **1**-**3** to **4**. Notably, MucF showed sequence

similarity to a large group of hypothetical proteins from the genomes of bacteria associated with

the human gut and skin (**Supplementary Fig. 33**). We therefore speculate MucF joins a large

family of unrecognized deacylases that may play important roles within the human microbiota.

Next, we took two approaches to determine whether RTCs and MUC play roles in

mediating inter-species bacterial competition. Frist, we used a plate-based competition assay, in

which a colony of *S. mutans* UA159 (a model organism for caries disease), *S. mutans* B04Sm5

(producing **1**-**4**), *S. mutans* B04Sm5/Δ*mucD* (**1**-**4** deficient strain), or *S. mutans* B04Sm5/Δ*mucF*

(producing **1**-**3** exclusively) was plated next to nascent colonies of other oral bacteria, including

Rothia mucilaginosa, *S. sanguinis*, *S. gordonii*, *S. mitis*, *S. pneumoniae*, and *S. salivarius* (**Fig.**

3). In general, *S. mutans* B04Sm5 exhibited greater inhibition of neighbors than the *S. mutans* model strain UA159. In contrast, *S. mutans* B04Sm5/Δ*mucD*, which lacks the production of compounds **1**-**4**, dramatically exhibited reduced inhibition of its neighbors. For instance, *S. mutans* B04Sm5 impaired the growth of *S. sanguinis* completely, while *S. mutans* B04Sm5/Δ*mucD* only showed slight inhibition of *S. sanguinis*. Furthermore, although the growth of *S. mutans* B04Sm5/Δ*muc*F was impaired by overproducing **1**-**3**, it exhibited the strongest inhibition of neighbors among all tested strains. *S. sanguinis* is one of the predominant species of the indigenous oral biota colonizing dental plaque, which is normally associated with healthy dental biofilm(32). The antagonistic relationship between *S. sanguinis* and *S. mutans* is well-characterized, and plays an important role in caries development(33, 34). Therefore, we determined the Minimum Inhibitory Concentration (MIC) of isolated **1** and **4** against the *S. sanguinis* ATCC 49296. Remarkably, we observed significant antibacterial activity of **1** against the *S. sanguinis* (MIC=3.1 µM), However, **4** did not show any antibacterial activities against *S. sanguinis* up to a concentration of 2 mM. These results provide compelling support that the *muc* pathway, through compounds **1**-**3,** provides a competitive advantage for *S. mutans* B04Sm5 by inhibiting the growth of its competitors. The increased competitive fitness conferred by *muc* is likely to increase the virulence *S. mutans* strains bearing the gene cluster. As *S. mutans* is an exceptionally productive biofilm-former, higher numbers of *S. mutans* are likely to increase plaque biofilm formation and promote the dysbiosis which leads to the formation of caries lesions. Interestingly, Chen and colleagues showed that **4** can significantly suppress the infiltration of leukocytes (CD45[+] cells) into the Matrigel plug in a mice model, suggesting an anti-inflammatory activity(23).

In summary, we describe a versatile biosynthetic pathway from an oral pathogen *S. mutans* B04Sm5, which can produce three types of compounds with divergent biological activities. These include three *N*-acyl tetramic acids (**1**-**3**) that display antibacterial properties against oral commensal bacteria, a new organic acid (**4**) with a reported anti-inflammatory activity in a mice

model(23), and a previously characterized SDSF with the ability to interact with pathogenic oral

fungi(31). Although two antibiotics have been discovered from the commensal bacteria of human,

including lactocillin(13) and lugdunin(35), RTCs, to our knowledge, are the first group of low

molecular weight antibacterial molecules identified from a human opportunistic pathogen. While

this study merely scrapes the "tip of the iceberg" of the recently identified oral biosynthetic

potential(13, 36), these findings clearly exemplify that deeper exploration of leads provided by

genome mining studies will help elucidate the complex ecological underpinnings of the human

microbiome and its relationship to disease.

METHODS

General methods. A complete list of the primers, plasmids, and strains used in this study can be

found in **Supplementary Table 6**. PCR products were amplified with PrimeSTAR HS DNA

polymerase (Clontech Laboratories, Inc., USA). DNA isolations and manipulations were carried

out using standard protocols. *Escherichia coli* strains were cultivated in LB medium (Thermo

Fisher Scientific, USA) supplemented with appropriate antibiotics. *S. mutans* B04Sm5 and its

respective derivatives were all grown on Brain Heart Infusion (BHI) agar or liquid medium (BD

Biosciences, USA) at 37 °C in a CO_2 incubator (5% CO_2/95% air). *Lactobacillus reuteri* LTH2584

was also grown on MRS medium (BD Biosciences, USA) or agar at 37 °C in a CO_2 incubator (5%

CO_2).

Production, extraction, and detection of reutericyclins (1-3) and mutanocyclin (4) from *S.

mutans. 50 milliliters of BHI medium were inoculated with a loop of glycerol stock of *S. mutans*

or a mutant thereof overnight. Thirty milliliters of preculture was inoculated into 3 liters of BHI

medium containing 1% glucose. After 12 hours incubation, 60 g of autoclaved Amberlite XAD7-

HP resin (Sigma-Aldrich, USA) was added to the cultures. The cultures were incubated for

another 36 hours at 37 °C in a CO_2 incubator (5% CO_2). The resin was recovered and washed

twice with water (200 mL) and the extract was subsequently extracted twice with ethyl acetate

(400 mL in total). The organic phase was evaporated, and extracts were dissolved in 2 mL

methanol. Each extract was monitored at 280 nm during separation by HPLC using a Kinetex®

C18 100 Å, LC Column (5 μm, 150 × 2.1 mm; Phenomenex, US) as follows: 0-15 min, 30% B;

15-16 min, 30%-100% B; 16-25 min, 100% B; 26-27 min, 100%-30% B; 28-35 min, 30% B (solvent

A: H_2O/TFA (999:1, v/v); solvent B: CH_3CN/TFA (999:1)).

Construction of *S. mutans* knockout plasmids. A 1010-bp fragment containing the

spectinomycin resistance gene (*spec*[R]) was amplified from pCAPB2(37) with primers spec_fwd

and spec_rev (**Supplementary Table 6**). The left (532 bp) and right (555 bp) flanks of *mucD* were

amplified from the genomic DNA of *S. mutans* B04Sm5 with the primer pairs of mucD_KO_L-

fwd/mucD_ KO_L-rev and mucD_ KO_R-fwd/mucD_ KO_R-rev (**Supplementary Table 6**),

respectively. These three PCR products were assembled with a double digested pUC19 (PstI and

EcoRI) using a NEBuilder HiFi DNA Assembly kit (New England Biolabs, USA), which resulted in

the vector pEXT01. Amplification of the left (602 bp) and right (611 bp) homology arms for

knocking out *mucF* were accomplished with primer pairs mucF_ KO_L-fwd/mucF_ KO_L-rev and

mucF_ KO_R-fwd/mucF_ KO_R-rev (**Supplementary Table 6**), respectively. These two PCR

products and *spec*[R] cassette were further cloned into pUC19 to give pEXT02 using the method

described above. Vector clones were verified by restriction analysis and sequencing.

Generation of Δ*mucD* and Δ*mucF* mutants. The disruption cassettes were amplified from

pEXT01 (2159 bp) and pEXT02 (2159 bp) using primer pairs mucD_KO_L-fwd/mucD_ KO_R-rev

and mucF_ KO_L-fwd/mucF_ KO_R-rev (**Supplementary Table 6**), respectively. PCR products

were digested by DpnI and then purified using the QIAquick PCR Purification Kit (Qiagen, USA).

The disruption cassettes were transferred to *S. mutans* B04Sm5 by a previously reported

protocol(38). Spectinomycin resistance clones were selected by growth on BHI agar

supplemented with 500 µg/mL spectinomycin, confirmed by PCR and sequencing, and

designated as *S. mutans* B04Sm5/Δ*mucD* and *S. mutans* B04Sm5/Δ*mucF*.

Generation of *muc* expression plasmid. The 8.6-kb DNA region containing *mucA-E* was PCR

amplified from the genomic DNA of *S. mutans* B04Sm5 in two fragments (each approximately 4

kb) with primer pairs mucA-E_ fwd1/ mucA-E_rev1 and mucA-E_fwd2/ mucA-E_rev2

(**Supplementary Table 6**). These fragments were cloned into the XhoI site of pACYCDuet-1 by

a NEBuilder HiFi DNA Assembly kit (New England Biolabs, USA), resulting in the plasmid pEXT06.

To construct pEXT07, *mucF* was amplified with primers mucF-coexp_fwd and mucF_coexp_rev.

PCR product and pEXT06 were digested with the restriction enzyme pair NcoI/BamHI and ligated

with T4 DNA ligase (New England Biolabs, USA). The resulting vectors were verified by restriction

analysis and sequencing. pEXT06 and pEXT07 were transformed into *E. coli* BAP1, respectively.

Expression, extraction, and detection of *muc* expression in *E. coli* BAP1. *E. coli* BAP1(39)

containing pEXT06 or pEXT07 were cultivated on LB plates supplemented with 1% glucose and

50 µg/mL chloramphenicol at 37 °C. The following day, a loop of *E. coli* cells was transferred for

precultures grown at 37 °C in 10 mL LB medium supplemented with 1% glucose and 50 µg/mL

chloramphenicol for 4-5 h. One microliter of each preculture was transferred to 50 mL of fresh LB

with the same supplements and grown at 37 °C to an OD_{600} of 0.4 to 0.6. Cultures were induced

with 200 µM IPTG and incubated for an additional 12-14 h at 30°C with shaking (220 rpm).

Cultures were harvested and 1 mL of H_2O supplemented with 0.5 mg/mL lysozyme was added to

the pellets. Cells were disrupted by sonication at room temperature. The lysates were acidified

with acetic acid (1% final concertation) and extracted by twice with an equal volume of EtOAc.

The organic phase was evaporated, resuspended in MeOH (0.2 mL), and filtered through

Acrodisc MS PTFE Syringe filters (Pall Inc., Ann Arbor, MI, USA) prior to HPLC analysis. Each

extract was monitored at 280 nm during separation by HPLC using a Kinetex® C18 100 Å, LC

Column (5 μm, 150 × 2.1 mm; Phenomenex, US) as follows: 0-10 min, 30% B; 10-11 min, 30%-

100% B; 11-25 min, 100% B; 26-27 min, 100%-30% B; 28-35 min, 30% B (solvent A: H_2O/TFA

(999:1, v/v); solvent B: CH_3CN/TFA (999:1)).

Feeding experiments for biosynthetic pathway study. *S. mutans* B04Sm5 was cultivated in

50 mL BHI medium supplemented with 1% glucose and 200 mg/L isotope-labeled [$^{13}C_1$] *L-* or *D-*

leucine (Sigma-Aldrich, USA). The compounds were extracted and isolated using the method

described above. Each extract prepared was dissolved in 100 μL MeOH for MS analysis using

the method described in the supplementary information (**Supplementary Note 1**).

Expression and activity mensuration of MucF in *E. coli*. Primer pair mucF_pET_fwd/mucF_

pET_rev (**Supplementary Table 6**) was used for amplification of *mucF* from the genomic DNA of

S. mutans B04Sm5. The PCR product was cloned into the NcoI and XhoI sites of pET28a to

obtain pEXT26 (with a C-terminal His-tag). Next, pET28a and pEXT26 were transferred into *E.*

coli Rosetta2™ (DE3)pLys, respectively. Single clones were picked for precultures grown

overnight at 37°C in TB broth (Thermo Fisher Scientific, USA) with 50 μg/mL kanamycin and

50μg/mL chloramphenicol at 37°C. One microliter of preculture was transferred to 1 L of fresh TB

broth with the same antibiotics and grown at 37°C to an OD600 of 0.4 to 0.6. Cultures were

induced with 500 μM IPTG and incubated for an additional 16 h at 18°C with shaking (220 rpm).

Cultures were harvested and 10 mL of buffer (50 mM Tris-HCl, pH 8, 150 mM NaCl, 10% glycerol)

supplemented with 0.5 mg/mL lysozyme and 0.5 mM PMSF was added to the pellets. Cells were

disrupted by sonication at 4°C. The lysate was used for MucF activity testing. The assay mixture

for the reaction (100 μL) consisted of 96 μL *E. coli* lysate (both carrying empty pET28a or pEXT26)

and 4 μl reutericyclin A (**1**) solution (6.6 mM, 80% EtOH). The reaction solutions were prepared

on ice and incubated at 37°C for 10 min, 30 min and 60 min. Reactions were terminated by the

addition of 1 μL acetic acid and then extracted twice with 200 μL EtOAc. After centrifugation of

the assay at 12,000 *g* for 10 min, the organic phase was evaporated and resuspended in 100 μL

MeOH (0.2 mL). The extracts were monitored by HPLC.

Agar plate-based assays. The interspecies competition assays with *S. mutans* (UA159,

B04Sm5, Δ*mucD,* Δ*mucF*), *R. mucilaginosa, S. gordonii*, and *S. sanguinis, S. mitis, S.*

pneumoniae, and *S. salivarius* were performed as described previously(40), with some

modifications. 8 μl of overnight cultures of *S. mutans* strains were inoculated onto BHI agar and

incubated at 37°C under a 5% CO_2/95% air atmosphere (vol/vol). After a 24 h incubation period,

an overnight culture of the indicated competing species was inoculated next to the *S. mutans* and

the plates were incubated for an additional 24 h and subsequently photographed.

Acknowledgments

The authors thank Y. Li (NYU College of Dentistry, USA) for providing *S. mutans* strain B04Sm5,

M.G. Gänzle (University of Alberta, Canada) for providing *L. reuteri* LTH2584 strain, C. Khosla

(Stanford University, USA) for *E. coli* BAP1, and J.J. Zhang (Massachusetts Institute of

Technology, USA), M.S. Donia (Princeton University, USA) and M.A. Fischbach (Stanford

University, USA) for valuable discussions. This work was supported by NIH grants R00-

DE0245543 to A.E., R01-GM085770 to B.S.M., and F32-DE026947 to J.B., and the Japan

Society for Promotion of Science Overseas Research Fellowship to Y.K.

Author contributions

X.T. designed the research and analyzed the *muc* pathway. X.T. generated and analyzed the

mutants, performed the biochemical experiments and the heterologous expression experiments.

Y. K. and X.T. purified the compounds and elucidated the structures of all compounds. X.T., P.A.J.

and S.M.K.M. performed the chemical synthesis. X.T., Y.K., J.G. and T.H. performed mass

spectrometry experiments and analyzed mass spectrometry data. X.T., J.B. and S.L. designed

and performed the agar plate-based assays. X.T., Y.K., J.B., A.E. and B.S.M. wrote the

manuscript. All authors analyzed and discussed the data and contributed to the writing of the

manuscript.

Competing financial interests

315   The authors declare no competing financial interests.

References

1.   Valdes AM, Walter J, Segal E, Spector TD. 2018. Role of the gut microbiota in nutrition
and health. BMJ 361:k2179.
2.   Gentile CL, Weir TL. 2018. The gut microbiota at the intersection of diet and human health.
Science 362:776-780.
3.   Singh RK, Chang HW, Yan D, Lee KM, Ucmak D, Wong K, Abrouk M, Farahnik B,
Nakamura M, Zhu TH, Bhutani T, Liao W. 2017. Influence of diet on the gut microbiome
and implications for human health. J Transl Med 15:73.
4.   Spanogiannopoulos P, Bess EN, Carmody RN, Turnbaugh PJ. 2016. The microbial
pharmacists within us: a metagenomic view of xenobiotic metabolism. Nat Rev Microbiol
14:273-287.
5.   Koppel N, Maini Rekdal V, Balskus EP. 2017. Chemical transformation of xenobiotics by
the human gut microbiota. Science 356.
6.   Lam KN, Alexander M, Turnbaugh PJ. 2019. Precision Medicine Goes Microscopic:
Engineering the Microbiome to Improve Drug Outcomes. Cell Host Microbe 26:22-34.
7.   Round JL, Mazmanian SK. 2009. The gut microbiota shapes intestinal immune responses
during health and disease. Nat Rev Immunol 9:313-323.
8.   Hooper LV, Littman DR, Macpherson AJ. 2012. Interactions between the microbiota and
the immune system. Science 336:1268-1273.
9.   Belkaid Y, Hand TW. 2014. Role of the microbiota in immunity and inflammation. Cell
157:121-141.
10.   Donia MS, Fischbach MA. 2015. HUMAN MICROBIOTA. Small molecules from the
human microbiota. Science 349:1254766.
11.   Garg N, Luzzatto-Knaan T, Melnik AV, Caraballo-Rodriguez AM, Floros DJ, Petras D,
Gregor R, Dorrestein PC, Phelan VV. 2017. Natural products as mediators of disease. Nat
Prod Rep 34:194-219.
12.   Wilson MR, Zha L, Balskus EP. 2017. Natural product discovery from the human
microbiome. J Biol Chem 292:8546-8552.

13. Donia MS, Cimermancic P, Schulze CJ, Wieland Brown LC, Martin J, Mitreva M, Clardy J, Linington RG, Fischbach MA. 2014. A systematic analysis of biosynthetic gene clusters in the human microbiome reveals a common family of antibiotics. Cell 158:1402-1414.

14. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. 2005. Defining the normal bacterial flora of the oral cavity. J Clin Microbiol 43:5721-5732.

15. Listl S, Galloway J, Mossey PA, Marcenes W. 2015. Global economic impact of dental diseases. J Dent Res 94:1355-1361.

16. Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, Ramos-Gomez F, Tagami J, Twetman S, Tsakos G, Ismail A. 2017. Dental caries. Nat Rev Dis Primers 3:17030.

17. Disease GBD, Injury I, Prevalence C. 2018. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet 392:1789-1858.

18. Bowen WH, Burne RA, Wu H, Koo H. 2018. Oral biofilms: pathogens, matrix, and polymicrobial interactions in microenvironments. Trends Microbiol 26:229-242.

19. Banas JA, Drake DR. 2018. Are the mutans streptococci still considered relevant to understanding the microbial etiology of dental caries? BMC Oral Health 18:129.

20. Liu L, Hao T, Xie Z, Horsman GP, Chen Y. 2016. Genome mining unveils widespread natural product biosynthetic capacity in human oral microbe Streptococcus mutans. Sci Rep 6:37479.

21. Merritt J, Qi F. 2012. The mutacins of *Streptococcus mutans*: regulation and ecology. Mol Oral Microbiol 27:57-69.

22. Joyner PM, Liu J, Zhang Z, Merritt J, Qi F, Cichewicz RH. 2010. Mutanobactin A from the human oral pathogen Streptococcus mutans is a cross-kingdom regulator of the yeast-mycelium transition. Org Biomol Chem 8:5486-5489.

23. Hao T, Xie Z, Wang M, Liu L, Zhang Y, Wang W, Zhang Z, Zhao X, Li P, Guo Z, Gao S, Lou C, Zhang G, Merritt J, Horsman GP, Chen Y. 2019. An anaerobic bacterium host system for heterologous expression of natural product biosynthetic gene clusters. Nat Commun 10:3665.

24. Lin XB, Lohans CT, Duar R, Zheng J, Vederas JC, Walter J, Ganzle M. 2015. Genetic determinants of reutericyclin biosynthesis in *Lactobacillus reuteri*. Appl Environ Microbiol 81:2032-2041.

25. Holtzel A, Ganzle MG, Nicholson GJ, Hammes WP, Jung G. 2000. The first low molecular weight antibiotic from lactic acid bacteria: reutericyclin, a new tetramic acid. Angew Chem Int Ed Engl 39:2766-2768.

26. Ganzle MG. 2004. Reutericyclin: biological activity, mode of action, and potential applications. Appl Microbiol Biotechnol 64:326-332.

27. Argimon S, Konganti K, Chen H, Alekseyenko AV, Brown S, Caufield PW. 2014. Comparative genomics of oral isolates of Streptococcus mutans by in silico genome subtraction does not reveal accessory DNA associated with severe early childhood caries. Infect Genet Evol 21:269-278.

28. Gaudelli NM, Townsend CA. 2014. Epimerization and substrate gating by a TE domain in beta-lactam antibiotic biosynthesis. Nat Chem Biol 10:251-258.

29. Hayashi A, Saitou H, Mori T, Matano I, Sugisaki H, Maruyama K. 2012. Molecular and catalytic properties of monoacetylphloroglucinol acetyltransferase from *Pseudomonas sp.* YGJ3. Biosci Biotechnol Biochem 76:559-566.

30. Pavkov-Keller T, Schmidt NG, Zadlo-Dobrowolska A, Kroutil W, Gruber K. 2019.
Structure and catalytic mechanism of a bacterial friedel-crafts acylase. Chembiochem
20:88-95.
31. Vilchez R, Lemme A, Ballhausen B, Thiel V, Schulz S, Jansen R, Sztajer H, Wagner-
Dobler I. 2010. *Streptococcus mutans* inhibits *Candida albicans* hyphal formation by the
fatty acid signaling molecule trans-2-decenoic acid (SDSF). Chembiochem 11:1552-62.
32. Zhu B, Macleod LC, Kitten T, Xu P. 2018. *Streptococcus sanguinis* biofilm formation &
interaction with oral pathogens. Future Microbiol 13:915-932.
33. Lamont RJ, Koo H, Hajishengallis G. 2018. The oral microbiota: dynamic communities
and host interactions. Nat Rev Microbiol 16:745-759.
34. Kreth J, Giacaman RA, Raghavan R, Merritt J. 2017. The road less traveled - defining
molecular commensalism with *Streptococcus sanguinis*. Mol Oral Microbiol 32:181-196.
35. Zipperer A, Konnerth MC, Laux C, Berscheid A, Janek D, Weidenmaier C, Burian M,
Schilling NA, Slavetinsky C, Marschal M, Willmann M, Kalbacher H, Schittek B, Brotz-
Oesterhelt H, Grond S, Peschel A, Krismer B. 2016. Human commensals producing a novel
antibiotic impair pathogen colonization. Nature 535:511-516.
36. Aleti G, Baker JL, Tang X, Alvarez R, Dinis M, Tran NC, Melnik AV, Zhong C, Ernst M,
Dorrestein PC, Edlund A. 2019. Identification of the bacterial biosynthetic gene clusters of
the oral microbiome illuminates the unexplored social language of bacteria during health
and disease. MBio 10:e00321-19.
37. Li Y, Li Z, Yamanaka K, Xu Y, Zhang W, Vlamakis H, Kolter R, Moore BS, Qian PY.
2015. Directed natural product biosynthesis gene cluster capture and expression in the
model bacterium *Bacillus subtilis*. Sci Rep 5:9383.
38. Perry D, Kuramitsu HK. 1981. Genetic transformation of *Streptococcus mutans*. Infect
Immun 32:1295-1297.
39. Pfeifer BA, Admiraal SJ, Gramajo H, Cane DE, Khosla C. 2001. Biosynthesis of complex
polyketides in a metabolically engineered strain of *E. coli*. Science 291:1790-2.
40. Kreth J, Zhang Y, Herzberg MC. 2008. *Streptococcal antagonism* in oral biofilms:
Streptococcus sanguinis and *Streptococcus gordonii* interference with *Streptococcus*
mutans. J Bacteriol 190:4632-40.

Figure legends

Figure 1 | Identification of an orphan gene cluster from *S. mutans* and its metabolites. (**a**) Mutanocyclin gene cluster (*muc*) annotation. (**b**) HPLC profile of extracts from wild type (WT) *S. mutans* B04Sm5 (i), *S. mutans* B04Sm5/Δ*mucD* (ii), and *S. mutans* B04Sm5/Δ*mucF* (iii). (**c**) Structures of metabolites identified from *S. mutans* in this study, including reutericyclin A (**1**), reutericyclin B (**2**), reutericyclin C (**3**), and mutanocyclin (**4**). C, condensation; A, adenylation; T, thiolation; KS, ketosynthase; TE, thioesterase.

Figure 2 | Model for 1-4 biosynthesis and characterization of MucF as deacetylase. (**a**) Model for the biosynthesis of **1-4**. (**b**) HPLC profiles of extracts from *E. coli* BAP1/pEXT06 (*mucA-D*) (i) and *E. coli* BAP1/pEXT07 (*mucA-D+mucF*) (ii). (**c**) HPLC analysis of (i) isolated **4** as a standard, (ii) compound **1** incubated with *E. coli* Rosetta2™ (DE3)pLys/pET28a (empty vector) cell lysate for 60 min, compound **1** incubated with *E. coli* Rosetta2™ (DE3)pLys/pEXT26 (carrying *mucF*) for 10 min (i), 30 min (ii), and 60 min (iii).

Figure 3 | Interspecies competition assay. Overnight cultures of oral commensal bacteria: *R. mucilaginosa, S. gordonii*, and *S. sanguinis, S. mitis, S. pneumoniae,* and *S. salivarius* were plated next to the indicated *S. mutans* strain (UA159, B04Sm5, Δ*mucD,* Δ*mucF*)).

a

The *muc* from *Streptococcus mutans* B04Sm5

Tailoring enzymes

Non-ribosomal peptide synthetase

Polyketide synthase

Hypothetical protein

Regulators and transporters

b

Retention time (min)

iii) Δ*mucF*

ii) Δ*mucD*

i) WT

c

a

MucD

MucE

R= 1

*= L or D

MucA
MucB
MucC

MucF

1-3

4

Free fatty acids

b

(ii)

(i)

Retention time (min)

c

(v) MucF_cell lysate_60min

(iv) MucF_cell lysate_30min

(iii) MucF_cell lysate_10min

(ii) empty vector_60min

(i) 4_std

Retention time (min)