Isolates from ancient permafrost help to elucidate species boundaries in *Acanthamoeba castellanii* complex (Amoebozoa: Discosea)

Stas Malavin[1], Lyubov Shmakova[2]

Institute of Physicochemical and Biological Problems in Soil Science RAS, Soil Cryology

Lab. Russia, 142290, Pushchino, Institutskaya 2.

[1] Corresponding author. E-mail: stas.malavin@gmail.com

[2] E-mail: lushmakova@gmail.com

Abstract

Acanthamoeba castellanii species complex (genotype T4) comprises of more than ten species with unclear synonymy. Its molecular phylogeny has several conflicts with published morphological data. In this paper, we analyze morphometric traits and temperature preferences in six new strains belonging to *A. castellanii* complex isolated from Arctic permafrost in the framework of molecular phylogeny. This integrative approach allows us to cross-link genotypic and phenotypic variability and identify species-level boundaries inside the complex. We also analyze previously known and newly found discrepancies between the nuclear and mitochondrial gene-based phylogenies. We hypothesize that one reason for these discrepancies may be the intragenomic polymorphism of ribosomal RNA genes.

Key words: Acanthamoeba, genotype T4, permafrost, Pleistocene, species concept, species delimitation, rDNA, 18S SSU, intragenomic polymorphism, morphometrics.

1 Introduction

Acanthamoeba is a protist genus belonging to the supergroup Amoebozoa. It has very

characteristic angular, or stellate, cysts with an envelope consisting of two distinct layers

(Fig. 1K, L, M) and fine tapered hyaline projections in the active forms – the so-called

acanthopodia (Fig. 1A, B, D, E). It is common in soils and aquatic environments all around

the world (Geisen et al., 2014). The genus is also known as an opportunistic pathogen causing

amoebic keratitis and encephalitis in humans (Khan, 2005; Maycock and Jayaswal, 2016).

Due to a well-established method of axenic cultivation, it has become a popular model in

biochemical and cell structure studies (Siddiqui and Khan, 2012). *Acanthamoeba* was among

the first organisms whose nuclear small subunit ribosomal RNA gene (18S rDNA) sequence

was determined (Gunderson and Sogin, 1986). Its genome is assembled and annotated

(Clarke et al., 2013), and several other draft assemblies are present in GenBank.

Despite such a rich history of study (or partially because of it), the system of the genus is

surprisingly tangled. The list of described species includes around 30 names. For several of

them, the synonymy is apparent from allozyme and molecular data and could also be

suspected for some others. Since the beginning of rDNA phylogeny era, several ribotypes

(traditionally called "genotypes" in *Acanthamoeba* literature) have been described based on a

threshold in genetic distances and characteristic signatures in genus-specific hypervariable

regions of the 18S gene. These types were designated T1–T22 (Corsaro and Venditti, 2018;

Tice et al., 2016). While some of the types directly correspond to formally described species,

others include type strains of several species. This particularly concerns type T4, the most

frequently isolated one in both clinics and nature (Maciver et al., 2013). Internally, type T4 is

also heterogeneous, which has been noted since the beginning of comparative studies of the

18S rDNA in *Acanthamoeba* (Fuerst et al., 2015; Gast et al., 1996), though genetic distances

between clusters inside T4 are smaller than those between this and the other types (Stothard

et al., 1998). Here belong the type strains of most of the species described on the basis of

morphology, including the generic type, *Acanthamoeba castellanii* (Douglas, 1930) and

thoroughly redescribed by Page (1967) *Acanthamoeba polyphaga* (Puschkarew, 1913). Due

to the plastic nature of characters by which T4 species are differentiated and unclear

synonymy, some authors prefer to designate this type "*A. castellanii* species complex"

(Corsaro et al., 2017; Kong, 2009). Therefore, reconciling molecular with morphological data

inside T4 is essential for the system of the genus as a whole, which means establishing a

genus-specific species concept integrating phenotypic and genotypic taxonomic markers

(Boenigk et al., 2012).

We isolated six new *Acanthamoeba* strains from Arctic permafrost sediments, a specific

environment, where they have been buried and trapped in a dormant but viable state. It is

shown that in cases when freezing occurs in parallel to sedimentation (which can be proven

by geological data), the age of embedded cells corresponds to the age of the permafrost

sediments, as both transportation and vital activity is impossible at temperatures below 10°C

(Rivkina *et al.*, 2000; Gilichinsky & Rivkina, 2011). Our strains were isolated from the

sediments dated from the late Pleistocene to early Holocene. Previously, other eukaryotes,

including ciliates (Shatilovich et al., 2015), amoebozoans (Shmakova et al., 2016, 2018),

flagellates (Shatilovich et al., 2009; Stoupin et al., 2012), green algae (Vishnivetskaya, 2009),

fungi (Kochkina et al., 2012), and nematodes (Shatilovich et al., 2018) were isolated from the

sediments of similar age. Even a higher plant belonging to the genus *Silene* was regenerated

from the tissue of a seed found in the excavated borrowing of a ground squirrel, although the

attempts to grow those seeds failed (Yashina et al., 2012).

All isolated *Acanthamoeba* strains belong to ribotype T4. To precisely identify the taxonomic

position of the strains, we additionally sequenced nearly full-length 16S (mitochondrial) gene

and the barcode region of cytochrome oxidase subunit I (Cox1) gene. Further, we performed

morphometrics of all life cycle stages, including cysts formed in different conditions. We also

studied the growth of isolated strains at different temperatures. By integrating these data, we

identified a species-level boundary within the studied range of variation. Putting

morphometrics and ecophysiology of several strains into a broader (almost genus-wide)

phylogenetic framework allowed us to figure out which level of genetic heterogeneity

corresponds to the species-level difference in phenotypes. Additionally, we analyzed

previously known and newly identified discrepancies between the phylogenies based on

different genes and hypothesized their origin.

10 Material and Methods

11 Sampling and strain isolation

Samples for subsequent enrichment cultivation of amoebae were taken from the intact inner

part of the permafrost cores. The cores were obtained in the Gydan Peninsula (Western

Siberia), Kolyma Lowland, and Bykovsky Peninsula (Eastern Siberia) by a special drilling

technique allowing for the sterility of the sample, as described elsewhere (Gilichinskiy et al.,

1989; Shi et al., 1997). For the map of the borehole locations see Supplementary Fig. 1. The

cores were kept frozen during all steps of transportation and laboratory processing, which

completely prevents the contamination of the inner material by modern microorganisms (Juck

et al., 2005; Gilichinsky & Rivkina, 2011).

Samples of about 1 cm^3 were taken from the cores in a laminar flow hood using a sterile

scalpel and put into sterile Petri dishes filled with autoclaved mineral Prescott and James

medium (Page, 1988). No bacterial culture was added, as reproducing viable bacteria from

permafrost soil formed sufficient food supply for the emerging amoebae. The plates have

been checked daily under a phase-contrast inverted microscope, and observed amoebas

transferred individually by a tapering Pasteur pipette into new plates, from which clonal

cultures were subsequently established.

Culturing and encystment

Strains were maintained as monoclonal bacterized cultures (the bacterial community

originated from initial permafrost sample) on 1.5% non-nutrient agar plates with 0,01%

cerophyl infusion overlay (Page, 1988) at 18°C. Cultures were passaged monthly. Cysts

formed in such cultures without any special induction one to two weeks after plating.

For axenization, cyst-rich cultures were washed from the agar layer, collected by

centrifugation at 1,000×g for 1 min, and treated with 1M HCl for 24 hours. The remaining

cysts were then washed thrice with autoclaved distilled water and incubated in sterile

proteose-yeast-glucose (PYG) medium (Neff et al., 1964) with 100 U mL^{-1} of penicillin and

0.1 mg mL^{-1} of streptomycin (Sigma-Aldrich). Axenic cultures were maintained in 25 cm^2

culture flasks in the same medium at 18°C and passaged every two weeks.

Three conditions of encystment were tested: "bacterized culture," "sparse axenic culture,"

and "dense axenic culture." A typical bacterized culture represented the first. Axenic cysts

were obtained from cultures grown in 3 mL PYG on 60 mm plates to the monolayer stage.

The "sparse axenic culture" conditions were achieved as follows: a subculture of 200 μL was

transferred into 3 mL Prescott and James (PJ) mineral medium (Page, 1988) and left for

several days. The rest of the cells were scraped and centrifuged at 1,000×g for 1 min, washed

with PJ, and left in 3 mL PJ for several days representing the "dense axenic culture"

conditions. From now on, we refer to the cysts formed in "bacterized culture" conditions as

"bacterized cysts"; to those formed in "sparse axenic culture" conditions as "sparse axenic

cysts"; to the cysts formed in "dense axenic culture" conditions as "dense axenic cysts."

Pseudocysts were obtained from axenic cultures. Cells grown at 18°C in 60 mm Petri dishes (3 mL PYG) up to the monolayer stage were washed once with PJ, collected by centrifugation, and put into 3 mL 5% dimethyl sulfoxide (DMSO) solution in PJ (Kliescikova et al., 2011). To test the viability of the obtained pseudocysts, they were put into fresh PYG medium, either immediately or after being completely dry during a week.

Microscopy and measurements

Living trophozoites and cysts were observed and filmed either on slides with petroleum jelly sealing using Leica DM2500 microscope with phase and differential contrast optics or directly in culture using Zeiss Axiostar Plus microscope with phase contrast optics and a water-immersion objective. Nikon DS-Fi3 and Watec WAT-221S digital cameras were used for filming in the first and the second case, correspondingly.

Measurements were performed on a monitor with a calibrated ruler using pictures obtained with a Zeiss Axiostar Plus microscope equipped with a Watec WAT-221S camera. In the case of trophozoites, locomotor forms and rounded cells were measured. The measured parameters in locomotor forms included (1) the longest dimension ("the length"), (2) the dimension perpendicular to the longest and going through the nucleus ("the width"), (3) the diameter of the nucleus, (4) the diameter of the nucleolus, (5) the maximum width of the hyaloplasm crescent at the anterior edge of the cell. For the rounded cells, cysts, and pseudocysts, the diameter was registered. For the cysts, we also counted the number of ostioles. The counting was done without any special staining using phase contrast optics caused the ostiole complex to appear glowing, which in the case of our strains was found to allow practically unambiguous identification of the ostiole number. The cyst measurements were performed across all three conditions tested (see Culturing and encystment).

1 Statistical analysis of morphometric characters

All calculations were performed in R v. 3.5.2 (R Core Team, 2019). We assumed no

normality assumptions for all variables. For a measure of central tendency, we calculated

medians and expressed variation as a median absolute deviation (as implemented in package

stats; scale factor 1.4826) or a range. Values exceeding three interquartile range around the

median were removed as outliers. The statistical significance of median differences was

assessed with Kruskal-Wallis rank-sum test (package *stats*) and non-parametric

("permutational") analysis of variance (PERMANOVA), as implemented in the function

adonis, package *vegan* v. 2.5-3 (Oksanen et al., 2019). Between-groups pairwise Wilcoxon

rank-sum tests with Benjamini & Yekutieli correction for multiple comparisons (package

stats) was used for the post-hoc comparison. Clustering of the morphometric data was done

using the neighbor-joining algorithm on euclidean distances, as implemented in *dist* and

hclust functions.

14 Sequence acquisition

DNA was extracted from axenic cultures grown in 60-mm Petri dishes to the stationary

phase. Cells were scraped from the plastic and collected by centrifugation as described above.

The pellets were washed thrice with autoclaved distilled water and finally resuspended in 100

μL of it. This suspension was subjected to the DNA isolation by ExtractDNA Blood kit

(Proteinase K plus SDS based lysis and silica-membrane purification; Evrogen, Russia),

according to the manufacturer protocol.

The nuclear SSU rRNA gene (18S), mitochondrial SSU rRNA gene (16S), and the

mitochondrially encoded cytochrome c oxidase subunit I gene (Cox1) were amplified,

purified by agarose gel electrophoresis, and sequenced on an automated Sanger sequencer in

Evrogen Co. (Russia). Details of PCR and sequencing may be found in the Supplementary Material.

GenBank accession numbers for the newly obtained sequences are as follows. 16S: MK100243–MK100248. Cox1: MK105577–MK105582. 18S: MK124583–MK124588.

Phylogenetic analysis

For phylogenetic inference, all available good-quality (nearly) full-length sequences were downloaded from GenBank. MAFFT v. 7.307 L-INS-i algorithm (Katoh and Standley, 2013) was used to align sequences. For 18S, hypervariable regions of the alignment were allocated into a separate partition, where gaps were coded as the fifth state (Simmons and Ochoterena, 2000). A phylogeny was reconstructed by Bayesian analysis of posterior probabilities (Bayesian inference, BI) in MrBayes v.3.2.6 (Ronquist et al., 2012) and maximum likelihood (ML) approach in IQTree v.1.6.6 (Nguyen et al., 2015). The detailed description of the phylogenetic analyses may be found in Supplementary Material.

Growth at different temperatures

We tested the growth of the isolated strains at 4°C, 11°C, 18°C, 25°C, 32°C, 37°C, 40°C, and 42°C ($\Delta = 1$°C for all). At 4–32°C, only axenic cultures were tested, while to 37–42°C, both axenic and bacterized cultures were exposed. Preliminarily, all temperatures were checked with 40-mm plates (1 mL PYG) in several replicates, to test if growth is possible at all. Axenic cultures for the tests were obtained by transferring 50 µL of a 5-day culture. In the case of bacterized cultures, test plates were inoculated with 50 µL of a two-week culture. This assay was carried out for one week; the cultures were checked daily.

For the temperatures, at which the strains have demonstrated growth or at least persisted successfully, the growth rate was estimated quantitatively. 12-well culture plates, 2 mL of

PYG per well, were inoculated with a growing axenic culture as described above. Each strain

was tested in two to three replicates per temperature level. The assays were conducted until

detached rounded cells started to prevail in the culture. Cells were counted in a

hemocytometer each 24 h (each 12 h at 32°C). At 4°C, the experiment lasted for a month, and

samples for counting were taken weekly, as observations showed no discernible increase of

the cell number. The culture growth was approximated by a logistic curve $N = \frac{L}{1+e^{-k(x-x_0)}}$,

where L is the asymptote and k – the logistic growth rate. The approximation was done by

Levenberg-Marquardt nonlinear least-squares algorithm as implemented in MINIPACK

library interfaced through *minpack.lm* package v.1.2-1 (Elzhov et al., 2016) for R v.3.5.2.

Results

Isolated strains

12 In total, we have obtained ten *Acanthamoeba* isolates from around 400 studied permafrost

13 sediment samples from East and West Siberia. We have succeeded in cloning and axenization

14 of six out of ten isolates. The location of the sampling sites is shown in Supplementary Fig. 1.

15 For the essential characteristics of the samples, see Tab. 1.

Microscopy observations

17 All obtained strains feature typical morphological characters of the genus, i.e., acanthopodia

18 in trophozoites and angular shape of the endocyst, and belong to the morphotype II following

19 Pussard and Pons (1977).

Trophozoites

21 Trophozoites of all strains readily develop locomotor forms when transferred to fresh

medium or the slide. The locomotor form is flattened, with a distinct frontal hyaline area

usually occupying about one-fourth of the cell length (Fig. 1A–E). Typically, it is elongated

in the direction of movement, with a width-to-length ratio of about 0.6 (Fig. 1A, B).

Occasionally, we have observed locomotor forms with the width equal to or even greater than

length (Fig. 1C, D). Such forms have been encountered more often in strains SCL-14-2, SCL-

14-3, SCL-14-12, and are generally more typical for larger cells. Acanthopodia are absent on

the leading lobopodium in locomotor forms, though the hyaline crescent edge has rarely been

even for a long time, regularly demonstrating undulations and protrusions. In cells moving

irregularly, acanthopodia are present on lobopodia also. At the beginning of an active

movement or while changing its way, trophozoites demonstrate sudden extensions of the

lobopodium in a new direction similar to the eruptive movements of heterolobose amoebae.

Generally, one vesicular nucleus with a single nucleolus is situated in the central part of the

cell in locomotor forms. At temperatures 25 to 32 °C, we have observed multinucleate

trophozoites (Fig. 1G). Those have always been bigger than the median, while nucleoli have

been smaller than in normal cells (Fig. 1G).

A single contractile vacuole is usually situated in the rear part of the moving cell (Fig. 1A,

B). In axenic cultures, the cytoplasm is strongly vacuolated, while in bacterized cultures, the

number of vacuoles is found to be considerably lower. A bulbous uroid often appeared in

locomotor forms (Fig. 1A, C, D), sometimes with adhesive filaments (Fig. 1D). Sporadically,

we observed a sizeable goblet-like food cup at the rear end of the moving cell (Fig. 1E),

identically to what was described and figured by Page (1967). This formation was

considerably more often found in strain SCL-14-12.

Floating forms have been rarely observed in bacterized cultures. The floating form represents

a central body mass with 15–20 radiating pseudopodia up to 15 μm long (Fig. 1F).

In axenic cultures grown in plates or culture flasks without agitation, amoebae form a

characteristic monolayer on the surface of the substrate (Fig. 1H). The "closeness" of the

monolayer depends on the temperature and adhesive properties of the plastic and differs

across the vessel area (Fig. 1H demonstrates the "ideal" case). After the monolayer reaches

its closeness limit, cells start to appear above it massively. This scenario is followed by all

studied strains but SCL-14-12, in which cells begin to form compact clumps above the

surface from the early stages of the culture growth, in parallel to the monolayer formation.

When the culture is roughly shaken or centrifuged, trophozoites appear as rounded cells

without lobopodia and acanthopodia. Such cells demonstrate the internal movement of

organelles, though no cell movements can be observed for some time. Similar rounded cells

also start to appear soon after an axenic culture reaches its growth limit. In this case, most of

them probably die in some time, as the final number of cysts in such cultures is much lower

than the number of rounded cells observed at the end of growth.

Cysts

All isolated strains readily encyst in laboratory cultures. In bacterized cultures, all or at least

most of the amoebae form cysts in about two weeks after transfer. In axenic cultures, we

obtained cysts by transferring amoebae into either the Neff's encystment medium (Neff et al.,

1964) or merely starving PJ medium. When starting to encyst, cells form plaques.

Mature cysts of all strains have well-defined endo- and ectocyst, the former folded and

rippled, the latter smoothly polyhedral. Beneath the endocyst, the characteristic layer of

oriented granules is always apparent (Fig. 1I, K, L, M). The number of endocyst corners

usually vary from 3 to 7 (Fig. 1I, K, M, O), but cysts with 8 and more corners have also been

encountered (Fig. 1L). The majority of ostioles lay in one plane. Opperculi do not protrude

above the ectocyst level. Dense axenic cysts usually have less prominent endocyst corners

than sparse axenic cysts and bacterized culture cysts. In the latter two, a considerable share of cysts has smooth endocyst without any defined corners (Fig. 1I).

Pseudocysts

The pseudocyst formation is total only at DMSO concentrations not lower than 5% and when the DMSO-to-cells ratio is not less than the PYG-to-cells ratio during culture growth. In such conditions, all cells transform into pseudocysts in 12 hours. Both dried and non-dried pseudocysts demonstrate almost total viability being put into PYG; in both cases, trophozoites appear in no more than a day.

Pseudocysts are rounded cells with a fine smooth envelope well seen under both phase- and DIC-contrast (Fig. 1J). The nucleus is situated in the central area of the cell; the nucleolus is slightly smaller than in trophozoites. The cytoplasm is evenly filled with refractile granules. The size of the granules in pseudocysts is bigger than that in cysts. No granular layer beneath the envelope characteristic for cysts may be observed.

Molecular phylogeny

18S

The inspection of the preliminary automatic alignment of the T4 18S sequences has revealed six hypervariable regions corresponding to the parts of the third, fifth, sixth, ninth, and eleventh hypervariable ("expansion") regions identified by Gast et al. (1996). Two of those six regions are parts of the diagnostic fragment 3 (DF3) (Booton et al., 2002). The DF3 region is used to distinguish sequence types (Booton et al., 2009, 2005, 2002), but we have found it sufficient to discriminate between T4 subtypes also, except for some rare cases of closely related sequences. The alignment of concatenated hypervariable regions has yielded easily discernible patterns (Supplementary Fig. 3) and has, first of all, demonstrated many

several-nucleotide-long indels present in many sequences which by any chance could not be

attributed to the alignment method.

Both BI and ML analyses produce seven subtypes, T4$_A$ through T4$_E$ and T4$_{Neff}$, previously

mentioned by Fuerst (2014) and Fuerst et al. (2015), further referred to as A–E and Neff. All

but A and B receive maximum support in both analyses; B is highly supported (0.9) and A

moderately supported (0.78) in the BI analysis. At the same time, the ML and BI trees display

one difference in their topologies (Fig. 2). While in both trees the major division lay between

the clades A+B and C+F+E, in the ML tree, subtype D appear sister to the A+B clade and

subtype Neff sister to subtype E. In the BI tree, vice versa, subtype D is sister to subtype E

and subtype Neff sister to the A+B clade.

Several subclusters have got high support in both ML and BI analyses inside clades A and B

(Fig. 2). From now on, we refer to them using well-known and long-studied strains they

include, for which 16S sequences are also available. This is done for the sake of

descriptiveness and does not imply that these strains are somehow "characteristic" of the

clusters being described. In clade A, those are "Castellani," "Galka," "Haas," "Rawdon,"

"Garcia" clades, and one other, not discussed further. In clade B, those are "Ma," "Diamond,"

"V017" clades, and one other.

In T4$_A$, the topology of the Bayesian tree differs from that of the ML tree (Fig. 3) chiefly by

the position of "Haas" clade, which is sister to the cluster containing "Galka" and

"Castellani" clades instead of being the deepest T4$_A$ branch as in the ML tree, and the

position of "Garcia" clade placed inside "Rawdon" clade instead of being the second deepest

branch as in the ML tree.

Strains SCL-am8, SCL-14-2, and SCL-14-3 branch together in the ML tree (Fig. 3) and

group with strain COHH 43 from marine sediments of Mt Hope Bay, MA, USA (Gast et al.,

2011) as a distinct clade sister to "Ma" clade of T4$_B$. In the BI tree, the permafrost strains and

COHH 43 branch before "Ma" clade. Other differences between the BI and ML tree

topologies in T4$_B$ involved terminal branching only. Three other permafrost strains grouped

within T4$_A$ clade. Strain SCL-14-9 is relatively distant from the other strains except for one

clinical isolate KA/E6 from Korea (Liu et al., 2006); their position in the tree is unstable.

Strain SCL-am9 is firmly placed into "Galka" clade, and strain SCL-14-12 – into "Castellani"

clade.

16S

The analysis of the 16S alignment confidently recovers all the previously established 18S

types for which 16S sequences are available (Fig. 4). We thus refer to them further without

mentioning in the framework of which analysis are they inferred. Type T4 is moderately

supported (94%) in ML-analysis but has posterior probability of 1 in BI. T2/6 type is sister to

T4 type; the next-closest is T5 followed by T3/T11 clade, though this topology as a whole

appears moderately supported. Types T7–T9 (morphogroup I by Pussard and Pons (1977)

form a very distinct clade with full support.

The internal structure of 16S-T4 clade partially reflects that of 18S-T4. D, E, and Neff

subtypes appear monophyletic with full support. At the same time, subtypes A and B, as well

as C and F, are not separate. Subtype E is the most distant from A/B clade, followed by C/F

clade, then subtype Neff. Subtype D is the closest to A/B clade. The 16S tree thus

corroborates the 18S one inferred with the ML approach, besides the A/B and C/F fusion.

The blended 16S A/B clade, like its 18S separate counterparts, is internally structured

(Fig. 4). Galka, Rawdon, and Garcia strains from 18S-T4$_A$ also hit highly supported clusters.

Strain Haas does not fall into any cluster but is itself steadily separated. At the same time,

strains Castellani and Ma have very similar 16S sequences and form a supported cluster,

together with some other strains. The same concerns strains Diamond and CDC:V017.

Surprisingly, all strains from permafrost except SCL-14-12 have identical 16S sequences, also identical to those of strains AC11 and AC16 from India (Megha et al., 2018). In addition to AC15, AC20, AC28, and CDC:V168 (18S-T4$_A$), they form a highly-supported cluster inside T4. SCL-14-12 groups together with Indian AC4 (Megha et al., 2018) and JPH24 from Japan (Rahman et al., 2013) (18S-T4$_B$) as a sister clade to a group containing strains Galka, HC-2, CDC:V390, Liu-E1 (all 18S-T4$_A$), with absolute support. Indian strains AC13 and AC25 reveal rather divergent sequences, probably corresponding to a not yet recovered T4 subtype. Unfortunately, their 18S sequences in GenBank are not full, so it is not possible to either confirm or deny this hypothesis by now.

Cox1

As with the 16S gene, strains belonging to Rns D, E, and Neff subtypes fall into different highly supported clusters, while those within Rns A and B, as well as C and F, form fused clusters (Fig. 5). Clade D groups together with clades C/F, E, and Neff. This subdivision is not supported by ML analysis, though gained 0.97 posterior probability. Inside the C/F/Neff half of the tree, no subtype relationships are confidently resolved.

As with 16S, all permafrost strains, except SCL-14-12, possess identical sequences. Together, they form a supported clade with three sequences from the Ohio State University collection (OSU) 07-033, 07-003 (unfortunately, lacking available 18S sequences) and 07-027 (18S-T4$_A$). A sequence of unknown strain identified as *Acanthamoeba quina* Pussard et Pons, 1977 (genome CDFN) appears sister to that clade with high support. Another highly supported cluster, forming a sister to the just described clade, contains the sequence of strain V017 (Rns T4$_B$), sequences from genomes CDFL, CDFK, and CDFJ (probably misidentified as *Acanthamoeba pearcei* Nerad, Sawyer, Lewis et McLaughlin, 1993), and a set of OSU

sequences belonging to A and B 18S subtypes. Permafrost strain SCL-14-12 groups together

with strains OSU 06-024 and 08-016 (both 18S-T4$_A$). One more firm cluster includes the

sequence from genome CDFB, identified as *Acanthamoeba lugdunensis* Pussard et Pons,

1977. Other strains in this cluster are Linc Ap-1 (genome LQHA) and two OSU strains, both

attributed to 18S-T4$_A$. The last cluster in T4$_{A/B}$ is that encompassing sequences of strain

LVPEI 1060/96 (18S-T4$_B$) together with OSU sequences 02-031 (18S-T4$_A$), 05-023 and 07-

019 (18S-T4$_B$).

Morphometrics

Trophozoites

The length of the measured trophozoites of all strains is between 15 and 47 μm, and the width

between 5 and 27 μm (Fig. 6, Tab. 2). Locomotor forms of strain SCL-14-12 are considerably

shorter (median 23 μm; $\square X^2(1, 5) = 92.765$, $p < 2.2 \times 10^{-16}$). The difference in length in all

other strains is not significant. At the same time, the most narrow locomotor forms are those

of strain SCL-am9 (median 12 μm; $\square X^2(1, 5) = 51.785$, $p = 5.971 \times 10^{-10}$). This leads to the

significant difference in width-to-length ratio in the two strains, which is the biggest in SCL-

14-12 (median 0.69) and the smallest in SCL-am9 (median 0.46) ($X^2(1, 5) = 72.762$, $p =$

2.725×10^{-14}). This distinction in shape could be clearly observed by eye when inspecting a

series of locomotor forms (Fig. 1 A, B, C demonstrates the typical variants).

Some slight differences in the diameter of the nucleus and nucleolus have been observed. The

median nucleus diameter in locomotor forms of strains SCL-14-12 and SCL-am9 (4 μm) are

slightly lower than in other strains (5 μm) ($\square X^2(1, 5) = 62.971$, $p = 2.952 \times 10^{-12}$). The

difference is marginally significant between SCL-14-12 and SCL-14-9, SCL-am9 and SCL-

am8, and not significant between SCL-14-12 and SCL-am8. At the same time, the nucleolus

in SCL-14-12 is significantly bigger (median diameter 3 μm) than in other strains (2 μm)

($X^2(1, 5) = 46.089$, $p = 8.712 \times 10^{-9}$). This results in significantly higher nucleolus-to-nucleus

diameter ratio in strains SCL-14-12 and SCL-am9 (not different from SCL-14-9), and also

partially in strain SCL-14-9. In strain SCL-14-12, the median of the hyaline crescent breadth

is the lowest (4 μm) differing significantly from all other strains ($X^2(1, 5) = 128.58$, $p <$

2.2×10^{-16}). Strain SCL-14-9 has the next lowest breadth, also differing significantly from

other strains at 0.05 level.

Rounded cell diameter is between 15 μm in SCL-14-12 and 19 μm in SCL-14-9 and well

correlates to the cell length (Spearman's $\rho = 0.87$, $S = 4.6324$, $p = 0.025$).

Cysts

We have found the cyst diameter a good character in terms of dispersion and inter-group

differences. Not surprisingly, the dispersion of the cyst diameter in all estimated factor

groups ("Strains" per "Culture type" design) is lower than that of the length and width in

locomotor forms (MAD not greater than 12% of the median). The range of the measured cyst

diameters is 10 to 24 μm (medians from 13 to 17 μm in bacterized cultures, from 14 to 17 μm

in dense axenic cultures, and 15 to 19 μm in sparse axenic cultures) (Tab. 3). Permutational

analysis of variance shows the significance of the effect of "Strain" ($F(1, 5) = 263.6$,

$p < .001$), "Culture type" ($F(1, 2) = 420.4$, $p < .001$), and their interaction ($F(1, 10) = 29.09$,

$p < .001$). Generally, the cysts formed in bacterized cultures are the smallest followed by the

cysts from dense axenic cultures and then by the cysts from sparse axenic cultures, which in

all strains are about the same size than the "normal" rounded trophozoites (Fig. 7), indicating

almost no shrinkage during the cyst formation. In strains SCL-am8 and SCL-14-9, these

differences in diameter are the most prominent, while SCL-14-12 has the smallest range (the

difference is still significant).

No variation in the number of operculi is observed between cysts formed in different

conditions.

Pseudocysts

The diameter of the pseudocysts in the studied strains varies from 11 to 16 μm (Tab. 4). As

with cysts, SCL-14-12 is the smallest. Also, in all strains except SCL-14-9 and SCL-am8,

pseudocysts are smaller than any type of cysts. In the two mentioned strains, cysts formed in

bacterized cultures are slightly but significantly smaller than pseudocysts (Fig. 7).

Growth at different temperatures

The results of the preliminary test demonstrate that none of the isolated strains can maintain

the stable growth at temperatures equal to or higher than 37°C, whether as axenic cultures in

PYG medium or bacterized cultures on agar plates with WG medium overlay. At 40 and

42°C, cells die entirely in about a day. At 37°C, some cells remain alive for several days

(more in bacterized than in axenic cultures), but their number declined to zero to the end of

the week.

32°C is not favorable for all cultures, although at this temperature, they exhibit the fastest

growth (Fig. 8, Tab. 5). In strains SCL-am8, SCL-am9, SCL-14-2, and SCL-14-3, cells do

not attach to the substratum and always appear roundish. A typical monolayer has never been

observed. Multinucleate cells start to appear from the first counting; subsequently, their

amount and size increase. The population density curve reaches stationarity very quickly. The

maximum culture density at this temperature does not exceed 8×10^5 cells mL^{-1} for SCL-am8,

5×10^5 cells mL^{-1} for SCL-14-2 and SCL-14-2, 2×10^5 cells mL^{-1} for SCL-am9. Strain SCL-

14-12 attaches to the bottom and grows better than all in terms of the culture density (up to

2×10^6 cells mL^{-1} have been recorded), though this constitutes a considerable decrease

compared to the densities at lower temperatures. Strain SCL-14-9 demonstrates a behavior

somewhat intermediate between SCL-14-12 and others. During the first two days, the culture

grows well, and cells attach to the substratum. Later, most of them are found roundish and

detached, and the monolayer is never formed. Multinucleate cells also appear, but later than

in others and lower quantity. The density reaches 6×10^5 cells mL^{-1}.

At 25°C, all strains grow well and form a monolayer in three to four days. Generally, the

culture density increases faster than at 18°C but stops growing earlier, reaching the lower

maximum. In strains SCL-am8, SCL-14-2, SCL-14-3 multinucleate cells also appear.

18°C is the optimal temperature for all strains. The maximal recorded densities are up to

8×10^5 cells mL^{-1} in SCL-14-2 and SCL-14-3; 1.5×10^6 cells mL^{-1} in SCL-am8, SCL-am9, and

SCL-14-9; 2.8×10^6 cells mL^{-1} in SCL-14-12) since the active growth continues longer,

although the growth rate is slightly lower than at 25°C. Cultures form a monolayer in four to

seven days. Cells form pseudopodia and actively move during a long time, even at the

stationary phase of growth.

At 11°C, all the cultures demonstrate a considerably lower rate of growth (Fig. 4). SCL-14-12

reaches a plateau in about 600 h at a density of more than 3×10^6 cells mL^{-1}. Other strains

show similar dynamics and reach a plateau in 400 h at a density of about 1.5×10^6 cells mL^{-1}

in SCL-am8 and about 1.0×10^6 cells mL^{-1} in others.

At 4°C, all the cultures have been in a good state during the whole period of observation,

judging by the cell and overall culture appearance. However, counting has shown that the cell

number has been slightly declining, although dividing cells have been observed during the

first two weeks.

The modeling of the logistic growth demonstrates three differences of SCL-14-12 in

comparison to other strains. First, its density asymptote is higher than in other strains at any

tested temperature, including 32°C. Second, its growth rate is about the same at 18 and 25°C,

whereas in other strains, the growth rate at 25°C is considerably higher than at 18°C.

Consequently, at 25°C, the growth rate of SCL-14-12 is lower than that of other strains. At

the same time, at 32°C, the growth rate of SCL-14-12 is strikingly higher, while in others it is

lower, the same, or (in SCL-14-3) just a little bit higher than at 25°C.

Discussion

Molecular phylogeny of *A. castellanii* complex (type T4)

18S

The distinctiveness of different clades inside type T4 has begun to manifest itself since the

first works on *Acanthamoeba* rDNA diversity (Booton et al., 2002; Chung et al., 1998; Fuerst

et al., 2015; Gast et al., 1996; Kong, 2009; Martín-Pérez et al., 2017; Stothard et al., 1998).

Subtypes $T4_C$, $T4_D$, $T4_E$, $T4_F$, and $T4_{Neff}$, when represented by several sequences, were

usually well-supported, while $T4_B$ was sometimes mixed with $T4_A$, and the latter mostly

paraphyletic. At the same time, the relationship between the subtypes was never well-

supported and appeared in part unstable. The deepest branching clade was either $T4_D$, $T4_E$, or

a clade containing both of them. $T4_{Neff}$ appeared sister to either $T4_{A/B}$ or $T4_{D/E}$ mixed clades.

$T4_{C/F}$ clade always branched after $T4_{D/E}$, except for the phylogeny using GTR model, where

$T4_D$ was robustly sister to $T4_{A/B/Neff}$ cluster (Martín-Pérez et al., 2017).

In our case, automatic alignment with subsequent filtering yields robust clades (subtypes)

$T4_C$ through $T4_{Neff}$, moderately supported $T4_B$, and paraphyletic $T4_A$ (tree not shown). All

seven subtypes as monophyletic clades could only be obtained with Jukes-Cantor substitution

model, the least-likely one in MrModelTest assessment. Relationships between the subtypes

are poorly supported. However, the fifth-state-coding approach demonstrates the

distinctiveness of all the T4 subtypes. The fact that $T4_C$—$T4_F$ subtypes may be obtained from

the masked alignment demonstrates the adequateness of the fifth-state-coding method used.

Moreover, monophyletic and fully-supported $T4_B$, $T4_D$—$T4_F$, and $T4_{Neff}$ may also be

obtained using the alignment of conservative regions only.

Thus, our 18S phylogeny corroborates and validates the previously obtained results. At the

same time, the method bears some possible sources of bias: (i) Indels of more than one

nucleotide are treated as separate events, while they may actually represent a single event, (ii)

Inversions are treated as several events, (iii) The amount of transversions is probably

underestimated, (iv) Unresolvable position of indels in single-nucleotide repeats. This

probable bias may explain differences in topology between BA and ML trees.

16S

As demonstrated by Ledee et al. (2003), 16S and 18S-based phylogenies of *Acanthamoeba*

corroborate each other in terms of identified sequence types (ribotypes). This holds for T4

type as well. Further, even a restriction analysis of the 16S gene (Byers et al., 1983) has

revealed remarkable heterogeneity inside 16S-T4 comparable to that found in 18S-T4. At the

same time, this study has first discovered the major inconsistency between 18S and 16S

phylogenies – Castellani and Ma strains (18S-$T4_A$ and $T4_B$, correspondingly) appeared

closely related and showed differences at the level of intraspecific variation in other

organisms. Other 18S-T4 type strains included in this study, namely Neff, Singh, and Page-

23, were found to be as distant as other ribotypes.

The sequence-based 16S phylogeny (Ledee et al., 2003) demonstrated the identity of

Castellani and Ma (as well as CDC:V042) sequences. Here, about 15% bases were removed

as ambiguously aligned, which explains why differences found by Byers et al. (1983) were

not observed. The rest of the sequences attributed to 18S-T4$_A$ and T4$_B$ subtypes were

intermixed. Nine new clades were identified inside 16S-T4. Sequences of the strains

attributed to 18S subtypes T4$_C$, T4$_D$, T4$_E$, T4$_{Neff}$ formed corresponding clades on the 16S tree

(T4$_F$ sequences were not included). The deepest branching clade was T4$_E$ following by T4$_C$,

then T4$_{Neff}$ clade. T4$_D$ sequences formed a clade sister to mixed T4$_{A/B}$ clade. Kong (2009)

using a smaller dataset identified five 16S clades, namely the deepest branching T4$_C$, the next

branching T4$_D$, then T4$_{Neff}$, and the last branching T4$_{A/B}$, which in turn consisted of two

clusters: the one containing, among others, Ma and Castellani strains and another one

containing *A. lugdunensis* L3a. Finally, Rahman et al. (2013) adding 13 T4 rns sequences

obtained generally the same nine clades as Ledee et al. (2003) inside T4 subtype.

We have identified around 13 well-supported clusters inside 16S-T4 subtype. In addition to

the proximity of Castellani and Ma strains, our study reveals other discrepancies between the

18S and 16S phylogenies. Permafrost strains SCL-am9 and SCL-14-9 belonging to 18S-T4$_A$

subtype and SCL-am8, SCL-14-2, SCL-14-3, belonging to 18S-T4$_B$ subtype, all have

identical 16S sequences (even before alignment masking). Strains CDC:V042 (ATCC 50493)

and CDC:V390 get into same "Rawdon" clade in 18S tree, but appear all distant in 16S tree.

Strain CDC:V042 is close to Castellani strain while being placed in a different highly

supported subcluster of T4$_A$ by 18S phylogeny. Galka strain (ATCC 50496) groups together

with SCL-am9 on the 18S tree, but is steadily apart from it on the 16S tree. By contrast, strain

CDC:V390 robustly branches apart from Galka on the 18S tree but has identical 16S

sequence. At the same time, strain Liu-E1 (ATCC 50709) is close to Galka in terms of 18S,

and 16S sequences of these two strains are identical too. Strain SCL-14-12, which robustly

clusters together with Castellani strain in absolutely all 18S analyses, steadily falls apart from

it on the 16S tree. Finally, strains Rawdon (BCM:37, ATCC 50497) and CDC:V390 cluster

together with high support on the 18S tree, but are steadily apart on the 16S tree.

To summarize, some of the strains demonstrate proximity of their 18S sequences and have

distant 16S sequences, some, *vice versa*, have distant 18S sequences and similar 16S

sequences, and other ones demonstrate similarity in both. To explain such complex

dissimilarity between 18S and 16S phylogenies, three hypotheses may be involved. The first

implies different rates of gene evolution in different strains, which is unlikely regarding their

taxonomic proximity. As the second, the improper alignment of indels may be invoked (see

the consideration of possible sources of bias above). Indeed, in single-nucleotide stretches in

an alignment, indels may occur in any position and ideally should not be aligned (which is

nonetheless done for the sake of parsimony). When indels are treated as the fifth state and

included in the calculation, their alignment may force sequences to appear more similar than

they are. This possibility cannot be completely ruled out here, but against it is the fact that

only A and B and C and F 18S subtypes became intermingled in 16S tree, while the others

(D, E, and Neff) stayed separate. Also, this hypothesis may explain why sequences that are

close to each other in 18S phylogeny appear distant in 16S one, but not the opposite cases (as

our 16S phylogeny did not account for indels).

The third hypothesis maintains that 18S subtypes $T4_A$ and $T4_B$, as well as $T4_C$ and $T4_F$,

actually represent intra-genomic variants of the gene present in the cell in a different number

of copies, with only the dominant one revealed by the PCR. Different strains may bear

different dominant variants and thus be attributed to either A or B (C or F) subtype while

sharing the same (or closely related) 16S gene sequences. Intragenomic variation in 18S gene

sequence is known in different protist supergroups including other Amoebozoa (Amaral-

Zettler et al., 2006; Gong et al., 2013; Gribble and Anderson, 2007; Kudryavtsev and

Gladkikh, 2017; Nassonova et al., 2010; Ntakou et al., 2019; Pillet et al., 2012) and reported

for several *Acanthamoeba* strains (Nuprasert et al., 2010; Qvarnstrom et al., 2013; Rahman et

al., 2013; Stothard et al., 1998). Furthermore, among 18S intragenomic variants found by

Rahman et al. (2013) in two strains, one belongs to $T4_A$ and another to $T4_B$ (the alignment is

available upon request). Also, this hypothesis may explain common difficulties with

obtaining a full 18S sequence from *Acanthamoeba* strains (the picture of contamination in

chromatograms from a clonal culture) and rarely reported cases when different 18S variants

were obtained from the same lens of a patient (Booton et al., 2002); $T4_A$ and $T4_C$) or from the

right and left lens cases (Ledee et al., 2009); both $T4_B$).

Cox1

The strains for which Cox1 sequences are available mostly have no full 18S sequences in the

public domain. Our Cox1 phylogeny fully corroborated the 16S phylogeny in the proximity

of $18S-T4_{A/B}$, as well as $18S-T4_{C/F}$, strains, the distinctiveness of other subtypes, and the

internal heterogeneity of $T4_{A/B}$ clade. Here again, strain SCL-14-12 was firmly placed into

different cluster than all other permafrost strains, which suggests the correspondence of the

Cox1 clusters to the 16S ones.

Morphometrics

Traditional taxonomy of *Acanthamoeba* has been primarily based on metric (overall

diameter, nucleus diameter, nucleolus diameter) and meristic (number of "angles", or

"corners", or "arms", i.e. protrusions of endocyst towards operculi) characters of cysts since

Page's redescription of *A. castellanii*, *A. polyphaga*, and *Acanthamoeba astronyxis* (Page,

1967). Pussard and Pons emphasized the number of angles and properties of endo- and

ectocyst (smoothness, distinctiveness), which they have studied in a considerable amount of

strains using different staining techniques (Pussard and Pons, 1977). However, with the

number of strains growing, the variability in these characters also grew and has become

overlapping. Also, as a rule, no statistical comparison of studied characters has been

conducted those times, although the number of measured individuals was usually sufficient.

The use of molecular methods has shown striking similarity of some of the described species,

e.g., *Acanthamoeba terricola* Pussard, 1964 and *A. castellanii* Neff strain (Visvesvara and

Balamuth, 1975); *Acanthamoeba divionensis* Pussard et Pons, 1977, *Acanthamoeba*

paradivionensis Pussard et Pons, 1977, *Acanthamoeba mauritaniensis* Pussard et Pons, 1977,

and *Acanthamoeba rhysodes* Singh, 1952 (De Jonckheere, 1983; Liu et al., 2005). Since the

beginning of the DNA sequencing era, the attention of investigators has been concentrated on

molecular phylogeny, and little attempts have been made to correspond morphometric

features to molecular clusters. As a result, no integrated species concept for *Acanthamoeba*

that would include both molecular and morphological characters seems to exist by now.

All our strains exhibit differences in the measured morphometric characters. The

morphometrics of locomotor forms demonstrates the relative stability of metric characters.

For the width, the most variable trait, the median absolute deviation (MAD) is in the range

from 17.4% (SCL-14-2) to 29.7% (SCL-am8) of the median (standard deviation (SD) from

16.5% to 25.5%). The MAD of the length is between 9.9% (SCL-14-3 and SCL-14-9) and

16.5% (SCL-am9) of the median (SD 12.1% to 19.4%). The level of variability found in the

measured characters of cysts is even lower (not surprisingly). With the number of measured

cells around 50, this level of variability allow to register significant differences (at $p \leq .05$)

even between genetically closely related strains (Tab. 2, 3).

It should be stressed that the culture conditions among the strains during the taking of

measurement were held essentially the same to assure that the observed differences may be

attributed to genetics. Unsurprisingly, we have found significant variability associated with

the type and density of the culture in cysts (Tab. 3, Fig. 7). This indicates that the obtained

results are valid for the comparison of the studied strains but may not be used to compare

with literature data unless the conditions of culturing coincide exactly.

The classifications of the strains based upon different subsets of the morphometric data

demonstrates different levels of similarity with molecular classification (Fig. 9). As it was

discussed previously, five out of six strains are identical in mitochondrial gene sequences, so

the only topological feature shared by all molecular trees is the most distant position of SCL-

14-12. Not all morphological trees share this feature as well. Specifically, this is the case for

locomotor form and sparse axenic cyst measurements (Fig. 9F, G), which also drives the

overall configuration (Fig. 9D). The dense axenic cyst and bacterized cyst data (Fig. 9H, I),

as well as the pseudocyst data, do not show this topology. Notably, monoclonal bacterized

culture on an agar plate is the most common setting in which measurements have been taken

from different *Acanthamoeba* strains, partially because not in every case a culture has been

axenized, and partially because this has been (probably rightly) believed to most closely

resemble the natural conditions. Our results suggest that measurements taken in axenic

conditions may better reflect genetic differences.

Thus, the most distant position of strain SCL-14-12 is not absolutely stable when different

sets of features and different culture conditions are compared. At the same time, we tend to

attach greater importance to the measurements of locomotor forms as to embracing more

different cell features. In addition to significant differences in linear dimensions, strain SCL-

14-12 also differs from others by its proportions, specifically the width to length, nucleolus to

nucleus diameter, and hyaloplasm width to cell length ratios. It should be noted that not only

SCL-14-12 is the most distant strain in the locomotor form-based classification, but also the

further branching (SCL-am9, then SCL-14-9) partially follows the order of the 18S tree (Fig.

9C).

So, to reliably compare morphometric data between strains in *Acanthamoeba* (and other

amoebae), the establishment of standardized culture conditions is absolutely necessary.

Axenic culture in a defined medium is the best candidate for such a standard, but the

temperature should obviously be considered as well.

2   Growth at different temperatures

As with the morphometric data, we observe differences in the culture growth are between all

studied strains and temperatures. Generally, the higher the temperature, the higher the logistic

growth rate, and the lower the culture density asymptote. Also, at temperatures from 25°C

and higher, multinucleate cells appear, prevailing over the normal mononucleate cells at 32°C

in most strains. James and Byers (1967) also observed multinucleates in aging cultures; their

amount was increased by agitation or air bubbling. Corsaro et al. (2015) observed

multinucleates in a culture of *Acanthamoeba micheli* Corsaro, Walochnik, Köhsler et Rott,

2015. Most likely, multinucleate cells result from incomplete cell division (karyokinesis not

followed by cytokinesis), but the possibility of cell fusion also cannot be ruled out (Pigon,

1972). Our data strongly supports the first way.

In the growth data, strain SCL-14-12 is also obviously different from others. At 25°C, it

shows approximately the same growth rate as at 18°C, while other strains increase it. At

32°C, no multinucleate cells appear in this strain, and the cells are attached to the substratum.

Also, the growth rate is the highest. Thus, in addition to higher growth limits at all

temperatures tested (a feature stemming probably from its smaller size), SCL-14-12 is

characterized by a broader range of temperatures suitable for normal culture growth.

The potential pathogenicity of *Acanthamoeba* strains is often addressed by testing their

possibility to grow at 34°C, a temperature of the human cornea (Walochnik et al., 2000), and

37°C. All permafrost strains except SCL-14-12 demonstrate clear signs of dysfunction

already at 32°C, and may thus be considered non-pathogenic. For SCL-14-12, the potential to

cause keratitis cannot be excluded.

Conclusion

Although a complex comparative study of strains inside *A. castellanii* complex (ribotype T4) remains a future perspective, our investigation demonstrates that the results of consistently applied morphometrics and ecophysiological studies (characteristics of growth at different temperatures) are in line with molecular variability of several kinds of genes (nuclear and mitochondrial) and may thus form a basis for *Acanthamoeba*-specific species concept. Amid the diversity of the entire genus, all studied strains are closely related to each other. However, strain SCL-14-12 is unique in all three sequenced genes, while the others possess the same sequences of the mitochondrial genes. Phylogenies built with all available sequences firmly place SCL-14-12 apart from the other strains. At the same time, it differs by many morphometric characters of both trophozoites and cysts, including proportions, as well as growth characteristics at different temperatures. Besides, directly observable features of cultures also distinguish this strain from the others. Other strains also exhibit some variation (and are obviously different in their 18S sequences), but this variation does not involve all studied features and is smaller in magnitude. Taking all this into account, we consider strain SCL-14-12 and other permafrost strains as two separate species. This conclusion may also be extrapolated onto other clusters obtained in mitochondrial phylogenies inside T4$_{A/B}$ clade, if only morphological and ecophysiological differences of strains between those clusters are consistent, which needs further investigation. Until such investigation placing phenotypic divergence in the framework of steady multigene phylogeny is done, a revision of *A. castellanii* species complex and thus the genus as a whole seems not possible.

Although most of *Acanthamoeba* 18S rDNA types are conceivably real taxa, in *A. castellanii* complex, the phylogenetic signal of this gene is biased, and thus it should not be used for species delimitation in this group. Mitochondrial gene phylogenies seem to be more suitable

for this purpose.

Acknowledgements

This study partially utilized microscopy equipment of the Core Facility Centres "Development of molecular and cell technologies" and "Culture Collection of Microorganisms" of Saint Petersburg State University (SPSU). We thank Dr. Alexei Smirnov for providing the possibility to use these facilities. We are obliged to all Soil Cryology lab members who participated in field sampling. We would also like to thank Prof. Dr. Jean-Michel Claverie, Laboratoire Information Génomique et Structurale, Aix Marseille Université, for his valuable comments on the manuscript.

This study was conducted in the framework of the Governmental Program 0191-2019-0044 and supported by Russian Foundation for Basic Research (RFBR) grants 18-04-00824 (isolation and culture establishment) and 17-54-150003 (description of the strains).

References

Amaral-Zettler, L.A., Cole, J., Laatsch, A.D., Nerad, T.A., Anderson, O.R., Reysenbach, A.-L., 2006. *Vannella epipetala* n. sp. isolated from the leaf surface of *Spondias mombin* (Anacardiaceae) growing in the dry forest of Costa Rica. J. Eukaryot. Microbiol. 53, 522–530. https://doi.org/10.1111/j.1550-7408.2006.00143.x

Boenigk, J., Ereshefsky, M., Hoef-Emden, K., Mallet, J., Bass, D., 2012. Concepts in protistology: Species definitions and boundaries. Eur. J. Protistol. 48, 96–102. https://doi.org/10.1016/j.ejop.2011.11.004

Booton, G.C., Joslin, C.E., Shoff, M., Tu, E.Y., Kelly, D.J., Fuerst, P.A., 2009. Genotypic identification of *Acanthamoeba sp.* isolates associated with an outbreak of

Acanthamoeba keratitis (AK). Cornea 28, 673–676.

https://doi.org/10.1097/ICO.0b013e31819342a7

Booton, G.C., Kelly, D.J., Chu, Y.-W., Seal, D.V., Houang, E., Lam, D.S.C., Byers, T.J., Fuerst, P.A., 2002. 18S ribosomal DNA typing and tracking of *Acanthamoeba* species isolates from corneal scrape specimens, contact lenses, lens cases, and home water supplies of *Acanthamoeba* keratitis patients in Hong Kong. J. Clin. Microbiol. 40, 1621–1625. https://doi.org/10.1128/JCM.40.5.1621-1625.2002

Booton, G.C., Kelly, D.J., Fuerst, P.A., 2005. Genotypic identification and species names in *Acanthamoeba*: correlation and conflict. J. Eukaryot. Microbiol. 52, 7S-27S. https://doi.org/10.1111/j.1550-7408.2005.05202003_1_13.x

Byers, T.J., Bogler, S.A., Burianek, L.L., 1983. Analysis of mitochondrial DNA variation as an approach to systematic relationships in the genus *Acanthamoeba*. J. Protozool. 30, 198–203. https://doi.org/10.1111/j.1550-7408.1983.tb02903.x

Chung, D.I., Yu, H.S., Hwang, M.Y., Kim, T.H., Kim, T.O., Yun, H.C., Kong, H.H., 1998. Subgenus classification of *Acanthamoeba* by riboprinting. Korean J. Parasitol. 36, 69–80. http://dx.doi.org/10.3347/kjp.1998.36.2.69

Clarke, M., Lohan, A.J., Liu, B., Lagkouvardos, I., Roy, S., Zafar, N., Bertelli, C., Schilde, C., Kianianmomeni, A., Bürglin, T.R., others, 2013. Genome of *Acanthamoeba castellanii* highlights extensive lateral gene transfer and early evolution of tyrosine kinase signaling. Genome Biol. 14, R11. https://doi.org/10.1186/gb-2013-14-2-r11

Corsaro, D., Köhsler, M., Montalbano Di Filippo, M., Venditti, D., Monno, R., Di Cave, D., Berrilli, F., Walochnik, J., 2017. Update on *Acanthamoeba jacobsi* genotype T15, including full-length 18S rDNA molecular phylogeny. Parasitol. Res. 116, 1273–1284. https://doi.org/10.1007/s00436-017-5406-1

Corsaro, D., Venditti, D., 2018. An apparent *Acanthamoeba* genotype is the product of a

chimeric 18S rDNA artifact. Parasitol. Res. 117, 571–577.

https://doi.org/10.1007/s00436-017-5690-9

Corsaro, D., Walochnik, J., Köhsler, M., Rott, M.B., 2015. *Acanthamoeba* misidentification

and multiple labels: Redefining genotypes T16, T19, and T20 and proposal for

Acanthamoeba micheli sp. nov. (genotype T19). Parasitol. Res. 114, 2481–2490.

https://doi.org/10.1007/s00436-015-4445-8

De Jonckheere, J.F., 1983. Isoenzyme and total protein analysis by agarose isoelectric

focusing, and taxonomy of the genus *Acanthamoeba*. J. Protozool. 30, 701–706.

https://doi.org/10.1111/j.1550-7408.1983.tb05346.x

Demidov, N.E., Baranskaya, A.V., Durdenko, E.V., Zanina, O.G., Karaevskaya, E.S.,

Pushina, Z.V., Rivkina, E.M., Spirina, E.V., Spenser, R., 2016. [Biogeochemistry of

the permafrost of the Arctic coast of the Gydan Peninsula]. Probl. Arct. Antarct. 3,

34–49. (In Russian)

Elzhov, T.V., Mullen, K.M., Spiess, A.-N., Bolker, B., 2016. Minpack.lm: R interface to the

Levenberg-Marquardt nonlinear least-squares algorithm found in MINPACK, plus

support for bounds. URL https://CRAN.R-project.org/package=minpack.lm (accessed

8.8.19).

Fuerst, P.A., 2014. Insights from the DNA databases: Approaches to the phylogenetic

structure of *Acanthamoeba*. Exp. Parasitol. 145, S39–S45.

https://doi.org/10.1016/j.exppara.2014.06.020

Fuerst, P.A., Booton, G.C., Crary, M., 2015. Phylogenetic analysis and the evolution of the

18S rRNA gene typing system of *Acanthamoeba*. J. Eukaryot. Microbiol. 62, 69–84.

https://doi.org/10.1111/jeu.12186

Gast, R.J., Ledee, D.R., Fuerst, P.A., Byers, T.J., 1996. Subgenus systematics of

Acanthamoeba: Four nuclear 18S rDNA sequence types. J. Eukaryot. Microbiol. 43,

498–504. https://doi.org/10.1111/j.1550-7408.1996.tb04510.x

Gast, R.J., Moran, D.M., Dennett, M.R., Wurtsbaugh, W.A., Amaral- Zettler, L.A., 2011. Amoebae and *Legionella pneumophila* in saline environments. J. Water Health 9, 37– 52. https://doi.org/10.2166/wh.2010.103

Geisen, S., Fiore-Donno, A.M., Walochnik, J., Bonkowski, M., 2014. *Acanthamoeba* everywhere: High diversity of *Acanthamoeba* in soils. Parasitol. Res. 113, 3151– 3158. https://doi.org/10.1007/s00436-014-3976-8

Gilichinskiy, D.A., Khlebnikova, G.M., Zvyagintsev, D.G., Fedorov-Davydov, D.G., Kudryavtseva, N.N., 1989. Microbiology of sedimentary materials in the permafrost zone. Int. Geol. Rev. 31, 847–858. https://doi.org/10.1080/00206818909465938

Gilichinsky, D.A., Rivkina, E.M., 2011. Permafrost microbiology, in: Reitner, J., Thiel, V. (Eds.), Encyclopedia of Geobiology, Encyclopedia of Earth Sciences Series. Springer Netherlands, pp. 726–732. https://doi.org/10.1007/978-1-4020-9212-1_162

Gong, J., Dong, J., Liu, X., Massana, R., 2013. Extremely high copy numbers and polymorphisms of the rDNA operon estimated from single cell analysis of oligotrich and peritrich ciliates. Protist 164, 369–379. https://doi.org/10.1016/j.protis.2012.11.006

Gribble, K.E., Anderson, D.M., 2007. High intraindividual, intraspecific, and interspecific variability in large-subunit ribosomal DNA in the heterotrophic dinoflagellates *Protoperidinium*, *Diplopsalis*, and *Preperidinium* (Dinophyceae). Phycologia 46, 315–324. https://doi.org/10.2216/06-68.1

Gunderson, J.H., Sogin, M.L., 1986. Length variation in eukaryotic rRNAs: Small subunit rRNAs from the protists *Acanthamoeba castellanii* and *Euglena gracilis*. Gene 44, 63–70. https://doi.org/10.1016/0378-1119(86)90043-0

James, T.E., Byers, T.J., 1967. The induction of multinuclearity in agitated and in aging

cultures of *Acanthamoeba sp.* Neff. J. Cell. Physiol. 70, 53–61.

https://doi.org/10.1002/jcp.1040700108

Juck, D.F., Whissell, G., Steven, B., Pollard, W., McKay, C.P., Greer, C.W., Whyte, L.G.,

2005. Utilization of fluorescent microspheres and a green fluorescent protein-marked

strain for assessment of microbiological contamination of permafrost and ground ice

core samples from the Canadian High Arctic. Appl. Environ. Microbiol. 71, 1035–

1041. https://doi.org/10.1128/AEM.71.2.1035-1041.2005

Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30, 772–780.

https://doi.org/10.1093/molbev/mst010

Khan, N.A., 2005. Granulomatous amoebic encephalitis: Clinical diagnosis and management.

Am. J. Infect. Dis. 1, 79–83. https://doi.org/10.3844/ajidsp.2005.79.83

Kliescikova, J., Kulda, J., Nohynkova, E., 2011. Stress-induced pseudocyst formation – a

newly identified mechanism of protection against organic solvents in acanthamoebae

of the T4 genotype. Protist 162, 58–69. https://doi.org/10.1016/j.protis.2010.03.006

Kochkina, G., Ivanushkina, N., Ozerskaya, S., Chigineva, N., Vasilenko, O., Firsov, S.,

Spirina, E., Gilichinsky, D., 2012. Ancient fungi in Antarctic permafrost

environments. FEMS Microbiol. Ecol. 82, 501–509. https://doi.org/10.1111/j.1574-

6941.2012.01442.x

Kong, H.H., 2009. Molecular phylogeny of *Acanthamoeba*. Korean J. Parasitol. 47, S21–S28.

https://doi.org/10.3347/kjp.2009.47.S.S21

Kudryavtsev, A., Gladkikh, A., 2017. Two new species of *Ripella* (Amoebozoa, Vannellida)

and unusual intragenomic variability in the SSU rRNA gene of this genus. Eur. J.

Protistol. 61, 92–106. https://doi.org/10.1016/j.ejop.2017.09.003

Ledee, D.R., Booton, G.C., Awwad, M.H., Sharma, S., Aggarwal, R.K., Niszl, I.A., Markus,

M.B., Fuerst, P.A., Byers, T.J., 2003. Advantages of using mitochondrial 16S rDNA

sequences to classify clinical isolates of *Acanthamoeba*. Investig. Opthalmology Vis.

Sci. 44, 1142. https://doi.org/10.1167/iovs.02-0485

Ledee, D.R., Iovieno, A., Miller, D., Mandal, N., Diaz, M., Fell, J., Fini, M.E., Alfonso, E.C.,

2009. Molecular identification of T4 and T5 genotypes in isolates from

Acanthamoeba keratitis patients. J. Clin. Microbiol. 47, 1458–1462.

https://doi.org/10.1128/JCM.02365-08

Legendre, M., Bartoli, J., Shmakova, L., Jeudy, S., Labadie, K., Adrait, A., Lescot, M.,

Poirot, O., Bertaux, L., Bruley, C., others, 2014. Thirty-thousand-year-old distant

relative of giant icosahedral DNA viruses with a pandoravirus morphology. Proc.

Natl. Acad. Sci. 111, 4274–4279. https://doi.org/10.1073/pnas.1320670111

Liu, H., Ha, Y.-R., Lee, S.-T., Hong, Y.-C., Kong, H.-H., Chung, D.-I., 2006. Genetic

diversity of *Acanthamoeba* isolates from ocean sediments. Korean J. Parasitol. 44,

117. https://dx.doi.org/10.3347/kjp.2006.44.2.117

Liu, H., Moon, E.-K., Yu, H.-S., Jeong, H.-J., Hong, Y.-C., Kong, H.-H., Chung, D.-I., 2005.

Evaluation of taxonomic validity of four species of *Acanthamoeba*: *A. divionensis*, *A.*

paradivionensis, *A. mauritaniensis*, and *A. rhysodes*, inferred from molecular

analyses. Korean J. Parasitol. 43, 7–13. https://doi.org/10.3347/kjp.2005.43.1.7

Maciver, S.K., Asif, M., Simmen, M.W., Lorenzo-Morales, J., 2013. A systematic analysis of

Acanthamoeba genotype frequency correlated with source and pathogenicity: T4 is

confirmed as a pathogen-rich genotype. Eur. J. Protistol. 49, 217–221.

https://doi.org/10.1016/j.ejop.2012.11.004

Martín-Pérez, T., Criado-Fornelio, A., Martínez, J., Blanco, M.A., Fuentes, I., Pérez-Serrano,

J., 2017. Isolation and molecular characterization of *Acanthamoeba* from patients

with keratitis in Spain. Eur. J. Protistol. 61, 244–252.

https://doi.org/10.1016/j.ejop.2017.06.009

Maycock, N.J.R., Jayaswal, R., 2016. Update on *Acanthamoeba* keratitis: diagnosis,

treatment, and outcomes. Cornea 35, 713.

https://doi.org/10.1097/ICO.0000000000000804

Megha, K., Sharma, M., Gupta, A., Sehgal, R., Khurana, S., 2018. Protein profiling of

Acanthamoeba species using MALDI-TOF MS for specific identification of

Acanthamoeba genotype. Parasitol. Res. 117, 729–736.

https://doi.org/10.1007/s00436-017-5743-0

Nassonova, E., Smirnov, A., Fahrni, J., Pawlowski, J., 2010. Barcoding amoebae: comparison

of SSU, ITS and COI genes as tools for molecular identification of naked lobose

amoebae. Protist 161, 102–115. https://doi.org/10.1016/j.protis.2009.07.003

Neff, R.J., Ray, S.A., Benton, W.F., Wilborn, M., 1964. Induction of synchronous

encystment (differentiation) in *Acanthamoeba sp.*, in: Methods in Cell Biology. pp.

55–83.

Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol.

Biol. Evol. 32, 268–274. https://doi.org/10.1093/molbev/msu300

Ntakou, E., Siemensma, F., Bonkowski, M., Dumack, K., 2019. The dancing star:

reinvestigation of *Artodiscus saltans* (Variosea, Amoebozoa) Penard 1890. Protist

170, 349–357. https://doi.org/10.1016/j.protis.2019.06.002

Nuprasert, W., Putaporntip, C., Pariyakanok, L., Jongwutiwes, S., 2010. Identification of a

novel T17 genotype of *Acanthamoeba* from environmental isolates and T10 genotype

causing keratitis in Thailand. J. Clin. Microbiol. 48, 4636–4640.

https://doi.org/10.1128/JCM.01090-10

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,

P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E.,
Wagner, H., 2019. Vegan: community ecology package. URL https://CRAN.R-project.org/package=vegan (accessed 7.8.19).

Page, F.C., 1988. A new key to freshwater and soil Gymnamoebae: with instructions for culture. Freshwater Biol. Assoc., Ambleside.

Page, F.C., 1967. Re-definition of the genus *Acanthamoeba* with descriptions of three species. J. Protozool. 14, 709–724. https://doi.org/10.1111/j.1550-7408.1967.tb02066.x

Pigon, A., 1972. Inhibition of movement, attachment, and cytokinesis by autogenous substances in the amoeba *Hartmannella*. Exp. Cell Res. 73, 170–176. https://doi.org/10.1016/0014-4827(72)90115-2

Pillet, L., Fontaine, D., Pawlowski, J., 2012. Intra-genomic ribosomal RNA polymorphism and morphological variation in *Elphidium macellum* suggests inter-specific hybridization in Foraminifera. PLOS ONE 7, e32373. https://doi.org/10.1371/journal.pone.0032373

Pussard, M., Pons, R., 1977. Morphologie de la paroi kystique et taxonomie du genre *Acanthamoeba* (Protozoa, Amoebida). Protistologica 13, 557–598.

Qvarnstrom, Y., Nerad, T.A., Visvesvara, G.S., 2013. Characterization of a new pathogenic *Acanthamoeba* species, *A. byersi* n. sp., isolated from a human with fatal amoebic encephalitis. J. Eukaryot. Microbiol. 60, 626–633. https://doi.org/10.1111/jeu.12069

R Core Team, 2019. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/ (accessed 7.8.19).

Rahman, M.M., Yagita, K., Kobayashi, A., Oikawa, Y., Hussein, A.I.A., Matsumura, T., Tokoro, M., 2013. Genetic characterization of clinical *Acanthamoeba* isolates from

Japan using nuclear and mitochondrial small subunit ribosomal RNA. Korean J. Parasitol. 51, 401–411. https://doi.org/10.3347/kjp.2013.51.4.401

Rivkina, E.M., Friedmann, E.I., McKay, C.P., Gilichinsky, D.A., 2000. Metabolic activity of permafrost bacteria below the freezing point. Appl. Environ. Microbiol. 66, 3230–3233. https://doi.org/10.1128/AEM.66.8.3230-3233.2000

Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol 61, 539–542. https://doi.org/10.1093/sysbio/sys029

Schirrmeister, L., Kunitsky, V., Grosse, G., Wetterich, S., Meyer, H., Schwamborn, G., Babiy, O., Derevyagin, A., Siegert, C., 2011. Sedimentary characteristics and origin of the Late Pleistocene Ice Complex on north-east Siberian Arctic coastal lowlands and islands – A review. Quat. Int., Timing and Vegetation History of Past Interglacials in Northern Eurasia 241, 3–25. https://doi.org/10.1016/j.quaint.2010.04.004

Shatilovich, A., Stoupin, D., Rivkina, E., 2015. Ciliates from ancient permafrost: Assessment of cold resistance of the resting cysts. Eur. J. Protistol. 51, 230–240. https://doi.org/10.1016/j.ejop.2015.04.001

Shatilovich, A.V., Shmakova, L.A., Mylnikov, A.P., Gilichinsky, D.A., 2009. Ancient Protozoa isolated from permafrost, in: Margesin, R. (Ed.), Permafrost Soils. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 97–115. https://doi.org/10.1007/978-3-540-69371-0_8

Shatilovich, A.V., Tchesunov, A.V., Neretina, T.V., Grabarnik, I.P., Gubin, S.V., Vishnivetskaya, T.A., Onstott, T.C., Rivkina, E.M., 2018. Viable nematodes from Late Pleistocene permafrost of the Kolyma River Lowland. Dokl. Biol. Sci. 480, 100–

102. https://doi.org/10.1134/S0012496618030079

Shi, T., Reeves, R.H., Gilichinsky, D.A., Friedmann, E.I., 1997. Characterization of viable

    bacteria from Siberian permafrost by 16S rDNA sequencing. Microb. Ecol. 33, 169–

    179. https://doi.org/10.1007/s002489900019

Shmakova, L., Bondarenko, N., Smirnov, A., 2016. Viable species of *Flamella* (Amoebozoa:

    Variosea) isolated from ancient Arctic permafrost sediments. Protist 167, 13–30.

    http://dx.doi.org/10.1016/j.protis.2015.11.001

Shmakova, L.A., Karpov, S.A., Malavin, S.A., Smirnov, A.V., 2018. Morphology, biology

    and phylogeny of *Phalansterium arcticum* sp. n. (Amoebozoa, Variosea), isolated

    from ancient Arctic permafrost. Eur. J. Protistol. 63, 117–129.

    https://doi.org/10.1016/j.ejop.2018.02.002

Siddiqui, R., Khan, N.A., 2012. Biology and pathogenesis of *Acanthamoeba*. Parasit. Vectors

    5, 6–6. https://doi.org/10.1186/1756-3305-5-6

Simmons, M., Ochoterena, H., 2000. Gaps as characters in sequence-based phylogenetic

    analyses. Syst. Biol. 49, 369–381.

Stothard, D.R., Schroeder-Diedrich, J.M., Awwad, M.H., Gast, R.J., Ledee, D.R., Rodriguez-

    Zaragoza, S., Dean, C.L., Fuerst, P.A., Byers, T.J., 1998. The evolutionary history of

    the genus *Acanthamoeba* and the identification of eight new 18S rRNA gene sequence

    types. J. Eukaryot. Microbiol. 45, 45–54.

Stoupin, D., Kiss, A.K., Arndt, H., Shatilovich, A.V., Gilichinsky, D.A., Nitsche, F., 2012.

    Cryptic diversity within the choanoflagellate morphospecies complex *Codosiga*

    botrytis – Phylogeny and morphology of ancient and modern isolates. Eur. J. Protistol.

    48, 263–273. https://doi.org/10.1016/j.ejop.2012.01.004

Tice, A.K., Shadwick, L.L., Fiore-Donno, A.M., Geisen, S., Kang, S., Schuler, G.A., Spiegel,

    F.W., Wilkinson, K.A., Bonkowski, M., Dumack, K., Lahr, D.J.G., Voelcker, E.,

Clauß, S., Zhang, J., Brown, M.W., 2016. Expansion of the molecular and morphological diversity of Acanthamoebidae (Centramoebida, Amoebozoa) and identification of a novel life cycle type within the group. Biol. Direct 11. https://doi.org/10.1186/s13062-016-0171-0

Vishnivetskaya, T.A., 2009. Viable cyanobacteria and green algae from the permafrost darkness, in: Margesin, R. (Ed.), Permafrost Soils. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 73–84. https://doi.org/10.1007/978-3-540-69371-0_6

Visvesvara, G.S., Balamuth, W., 1975. Comparative studies on related free-living and pathogenic amebae with special reference to *Acanthamoeba*. J. Protozool. 22, 245–256. https://doi.org/10.1111/j.1550-7408.1975.tb05860.x

Walochnik, J., Obwaller, A., Aspöck, H., 2000. Correlations between morphological, molecular biological, and physiological characteristics in clinical and nonclinical isolates of *Acanthamoeba* spp. Appl. Environ. Microbiol. 66, 4408–4413. http://doi.org/10.1128/AEM.66.10.4408-4413.2000

Yashina, S., Gubin, S., Maksimovich, S., Yashina, A., Gakhova, E., Gilichinsky, D., 2012. Regeneration of whole fertile plants from 30,000-y-old fruit tissue buried in Siberian permafrost. Proc. Natl. Acad. Sci. U. S. A. 109, 4008–4013. https://doi.org/10.1073/pnas.1118386109

Zanina, O.G., Gubin, S.V., Kuzmina, S.A., Maximovich, S.V., Lopatina, D.A., 2011. Late-Pleistocene (MIS 3-2) palaeoenvironments as recorded by sediments, palaeosols, and ground-squirrel nests at Duvanny Yar, Kolyma lowland, northeast Siberia. Quat. Sci. Rev., Beringia and Beyond: Papers Celebrating the Scientific Career of Andrei Vladimirovich Sher, 1939–2008 30, 2107–2123. https://doi.org/10.1016/j.quascirev.2011.01.021

1   Figure captions

2   **Figure 1 A–E** Locomotor forms: A – SCL-14-3; B – SCL-14-3; C – SCL-14-12, D – SCL-

3   am9; E – SCL-14-3; **F** Floating form, SCL-am8; **G** – multinucleate trophozoite, SCL-

4   am9; **H** Monolayer, SCL-14-12; **I, K–N** Cysts, various strains; **J** Pseudocysts, SCL-

5   am8; **O** Empty cyst, SCL-am8.

6   **Figure 2** ML (**A**) and BI (**B**) 18S SSU trees of *Acanthamoeba* T4 genotype showing seven

7   subtypes. Branch supports are given as bootstrap percentage / Bayesian posterior

8   probability. The trees are rooted at midpoint. Clades with variable position are

9   dashed. **Scale bar**: 0.01 substitution per site for both A and B.

10   **Figure 3** 18S SSU ML tree of T4A and T4B clades (a subtree of Fig. 5A, expanded).

11   Bayesian tree gave similar topology (see text). Branch supports are given as bootstrap

12   percentage / posterior probability. • – 100% or 1.00 support; ∗ – clade not observed.

13   Support values inside the clades are omitted. The tree is rooted at midpoint.

14   **Figure 4** 16S SSU ML tree of *Acanthamoeba* T4 genotype. BA tree gave the same topology.

15   Branch supports are given as ML bootstrap percentage / Bayesian posterior

16   probability. • – 100% or 1.00 support. Support values near tree tips are omitted. The

17   tree is rooted at midpoint.

18   **Figure 5** Cox1 ML-tree of *Acanthamoeba* T4 genotype rooted at midpoint. BA-tree gave the

19   same topology. Branch supports are given as ML-bootstrap percentage / Bayesian

20   posterior probability. • – 100% or 1.00 support. * – Identification is dubious. Support

21   values lower than 95/.75 are not shown.

Figure 6 Locomotor form length in studied strains.

Figure 7 Diameter of rounded cell, cyst, and pseudocyst of the studied strains in different

conditions.

Figure 8 A – Scatter-plot representing modeled parameters of the logistic growth of the

studied strains at different temperatures. Abscissa – L, ordinate – k. Lines connect

points belonging to the same strain. **B** – Cluster diagram of all three logistic model

parameters (scaled). Euclidean distance, Ward's clustering method.

Figure 9 Classification of the strains based on different data. **Gene trees: A** – Cox1 NJ tree,

observed distances; **B** – 16S NJ tree, observed distances; **C** – 18S tree extracted from

the complete tree of T4 type. **Morphometric trees: D** – All features included; **E** –

Cysts and pseudocysts only; **F** – Trophozoites only; **G** – "Sparse axenic" cysts; **H** –

"Dense axenic" cysts; **I** – "Bacterized culture" cysts. All trees are rooted at midpoint.

Not to scale.

Figure 2

Figure 3

0.01

Figure 4

0.05

Figure 5

Figure 8

Table 1

Click here to access/download
Table
Tab1.docx

dí\ ÿSupplementaryPlaceholder R   ÿÿ EÏlé

Table 2

dí\ ÿSupplementaryPlaceholder R   ÿÿ EÏlé

Table 3

dí\ ÿSupplementaryPlaceholder R   ÿÿ EÏlé

Table 4

dí\ ÿSupplementaryPlaceholder R   ÿÿ EÏlé

Table 5

Click here to access/download
Table
Tab5.docx

dí\ ÿSupplementaryPlaceholder R   ÿÿ EÏlé

Supplementary Material

‹

Click here to access/download
e-Component
Supplementary material.pdf