**No division of labour, and subfertile foundresses, in a phyllode-gluing Acacia thrips**

James D. J. Gilbert[1,2]

[1]School of Environmental Sciences, University of Hull, Cottingham Road, Hull HU6 7RX, UK.

Correspondence: james.gilbert@hull.ac.uk

[2]Fowlers Gap Arid Zone Research Station, School of Biological, Earth & Environmental

Sciences, University of New South Wales, NSW 2052, Australia

Keywords: Pleometrosis, joint nesting, cooperation, quasisocial, parasocial, division of labour

ABSTRACT

Behavioural variation is a hallmark of animal societies, which commonly contain breeders and

nonbreeders, and helpers and nonhelpers. In some cases labour is divided with nonbreeders

"helping" – gaining indirectly, via genetic benefits, or directly, e.g. by augmenting group size.

Conversely, they may benefit by not helping, conserving energy for breeding later. However,

subordinate behaviour after inheriting a breeding position is rarely evaluated.

In the Australian interior, Acacia thrips *Dunatothrips aneurae* (Thysanoptera) glue *Acacia*

phyllodes together into "domiciles". Foundresses, usually sisters, build domiciles singly or

communally. Some co-foundresses are nonreproductive, and their role is currently unknown. I

experimentally rejected the idea that they substantially "help" by contributing to domicile repair.

Nonreproductives were less likely to repair damage than reproductives. Alternatively, they may

be waiting to inherit the domicile, or simply of too poor quality to reproduce or help. To test

these alternatives, in the field, I allowed repairer or nonrepairer females to "inherit" a domicile by

removing their nestmate(s). Thus isolated, "nonrepairer" females took much longer to repair

domiciles than "repairers", control singletons or pairs. Although ovarian condition was equivalent

across groups, after 21 days nonrepairers actually laid fewer eggs compared to other groups.

Thus, labour was not divided: instead reproduction and helping covaried, probably depending

on female quality and the outcome of intra-domicile competition. Nonreproductive nonhelpers

were not waiting to breed. Their role, and their net effect on colony productivity, remains to be

shown. They are likely subfertile, and may make the "best of a bad job" by gaining indirect

benefits to the best of their limited ability.

**INTRODUCTION**

Social groups are composed of members that often vary in their contribution to necessary tasks.

In cooperative or eusocial societies, at any given time individuals typically partially or completely

specialize in particular tasks, especially reproduction ("reproductive skew"; Vehrencamp 1979;

Vehrencamp 1983a; Vehrencamp 1983b), but also other tasks (Beshers and Fewell 2001), for

example, foraging (Johnson 2010), parental care (Browning et al. 2012), nest defence (Gerber

et al. 1988), nest homeostasis (Jandt et al. 2009), and others (see Komdeur 2006). Within these

societies, costly non-reproductive tasks such as foraging and defence are typically performed by

"helpers" who are completely or partially reproductively suppressed, while breeding is performed

by one or a few reproductive individuals (Solomon and French 1997; Koenig and Dickinson

2004).

In some species, however, non-reproductive putative "helpers" do not appear to engage in

helping (Cant and Field 2001; Korb 2007; Browning et al. 2012). This can sometimes be

explained if the helper in question is waiting to inherit a breeding position and would incur

reproductive costs later by helping now (Field and Foster 1999; Cant and Field 2001;

Leadbeater et al. 2011). Alternatively, a helper may appear "lazy" compared to other helpers if it

has less genetic stake in current offspring (e.g. Curry 1988), may function as an insurance

workforce in case other helpers are injured (Baglione et al. 2010) or may be helping in subtle

ways that may not have yet been measured (Turnbull et al. 2012; Frank et al. 2018). Perhaps

more parsimoniously, one simple and long-standing explanation for non-reproductive, non-

helping individuals is that they are simply of poor quality, or subfertile - and, as such, may or

may not help depending on the extent of their limited ability, making the "best of a bad job"

(Eberhard 1975; Craig 1983). Perhaps surprisingly, there is as yet no unequivocal empirical

evidence for the "subfertility hypothesis" (Strassmann and Queller 1989), although there are

several cases where it has been shown not to apply (Sullivan and Strassmann 1984;

Leadbeater et al. 2011). In one example, Field & Foster (1999) found that nonreproductive

worker hover wasps given the chance to "inherit" their nest subsequently developed ovaries and

began reproducing.

In contrast to cooperative species in which contributions to different tasks are typically sharply

divided, some other groups are characterized by less pronounced behavioural and reproductive

specialization, e.g. lions (Packer et al. 2001), rodents (Hayes 2000), burying beetles (Trumbo

and Wilson 1993) and some sweat bees (although see Abrams and Eickwort 1981; Kukuk

1992). Where breeding is thus shared, individuals may help according to their genetic stake in a

brood (Hoogland 1983; Curry 1988), or conserve their helping effort for a time when they are

themselves reproducing (Packer et al. 2001).

*Dunatothrips aneurae* Mound (Thysanoptera:Phlaeothripidae), are tiny thrips that live in the arid

Australian Outback, where they can be solitary or social (Morris et al. 2002). Females construct

"domiciles" by gluing together terminal phyllodes of *Acacia aneura* trees into a completely

enclosed nest, within which they produce a single generation of offspring (Gilbert and Simpson

2013). Offspring develop entirely within the domicile, whose integrity is maintained by resident

adults. Domiciles are built from a silklike anal secretion (henceforth "silk") and function to

maintain humidity for developing larvae (Gilbert 2014). Domiciles are initiated by solitary

females or by groups of 2-5 (rarely up to 20), typically but not always related (Bono and Crespi

2008). The factors promoting nest co-founding are only partially understood, partly because

individual roles within groups are unknown. Co-founding carries survival advantages to all

females, especially when invaded by kleptoparasites (Bono and Crespi 2008), but also entails

reproductive competition for space which is intensified in small domiciles (Gilbert et al. 2018).

There is a low and variable degree of reproductive competition and skew within cofounded *D.*

*aneurae* domiciles. Cofoundresses vary in ovarian development, with a proportion of females

nonreproductive, especially in small domiciles (Gilbert et al. 2018). However, whether

reproduction co-varies with any form of "helping" is unknown. Females are uniformly pacifist

towards intruders (Gilbert et al. 2012; Gilbert and Simpson 2013) suggesting a lack of any

defence-based helping behaviour. On the other hand, the production of silk by nonreproductive

females may constitute altruistic behaviour (West et al. 2007). Silk is necessary for domicile

function (Gilbert 2014) and its production is potentially costly (see Craig et al. 1999 for a review

of "real" silk production in animals).

Any trade-off between silk and egg production would create at least the potential for selection

for division of labour within domiciles (Gilbert 2014). During fieldwork, I noticed that cohabiting

female *D. aneurae* varied in their latency to begin repairing experimental damage to domiciles -

variation that may potentially reveal just such a division of labour.  If good repairers are bad

reproducers, and *vice versa*, this would be good evidence of cooperative division of labour

within *D. aneurae* domiciles. In contrast, if good repairers are good reproducers, and *vice versa*,

*D. aneurae* is more likely to be a communal breeder, where subordinate females either (a) help

in proportion to their genetic stake in offspring or, similarly, are waiting to inherit the breeding

position; or (b) are of too poor quality either to breed or help at the same level as breeders.

These two alternatives can be separated by the behaviour of subordinates that are

experimentally allowed to inherit the breeding position: if subordinates fail to begin to breed

upon inheritance of the nest, alternative (b) is more likely. I asked two questions:

1.  Are reproduction and domicile repair divided among foundresses?

2.  Do nonreproductive foundresses breed upon inheritance of the domicile?

To address these questions, I performed two experiments.  First, in the lab, I identified "repairer"

and "nonrepairer" females by their latency to begin repairing experimental damage, and then

investigated whether repair tendency was associated with ovary development and egg

maturation, assessed via dissection. Second, in the field I conducted a removal experiment to

test whether the tendency to repair the nest was associated with reproduction when females

were given the chance to breed within the domicile - both potential reproduction (ovarian status)

and realized reproduction (eggs laid).  I did this by removing all other females, allowing the focal

female to "inherit" the domicile.

I had no particular *a priori* expectations with which to form strong predictions. On the one hand,

in species that form associations of foundresses from the same generation, which is almost

certainly true for *D. aneurae,* strong reproductive skew and division of labour is rarely found

(being more typical of matrifilial associations; Reeve and Keller 1995). Supporting this,

morphological castes do not appear to occur in *D. aneurae*. On the other hand, well-studied

relatives of *D. aneurae* within the Phlaeothripinae are eusocial (*Kladothrips* spp.; Crespi 1992),

although, being gall-inducers and with matrifilial societies, their ecology is somewhat different

(Crespi et al 2004).  Yet even morphologically similar individuals within groups can often show

naturally occurring behavioural variation in task performance, which can self-organize into

eusocial-like patterns of division of labour (Page 1997; Jeanson et al. 2005). I therefore made a

tentative prediction that I would find at least incipient division of labour within *Dunatothrips*

domiciles, such that breeding and repairing would be negatively associated.

METHODS

Fieldwork was conducted between January and October 2013 at a large patch of *A. aneura* at

the Bald Hills paddock (S 30° 57′ 39″; E 141° 42′ 18″) near the University of New South Wales

Arid Zone Research Station on the Fowlers Gap property, approx. 110km N of Broken Hill,

Australia (see Gilbert and Simpson 2013 for details).

*Are reproduction and repair divided?*

For lab studies, I used a subset of the *D. aneurae* domiciles collected from the field and

dissected as described in Gilbert & Simpson (2013).  In 48 nests, during nest dissections I

experimentally damaged the nest with watchmaker's forceps, and assessed their repair

behaviour before removing them from the nest and dissecting them to assess their ovarian (i.e.

reproductive) condition. Only dealate (reproductively mature) individuals were included in

experimental observations, as typically only these individuals contribute to repair (Gilbert and

Simpson 2013). I made a small tear in the nest, roughly equivalent to about 20% of the silk

area, and folded back the silk. Typically at least one foundress would thereupon begin repairing

immediately (within 30 seconds), which I scored as 2 ("Immediate repair"). I continued these

initial observations for 1 h. During this time, individuals were brushed lightly on the pronotum

with a few grains of a coloured fluorescent micronized paint powder (US Radium Co®), applied

using a fine paintbrush, to create a temporary distinctive ID that allowed me to follow individuals

within the nest. After 1 h I removed the first immediate repairer from the nest, using a probe

made from the hair of a shaving brush, and observed the behaviour of the remaining female(s)

over the subsequent 1 h period. One of more of these females typically began repairing upon

removal of a nestmate. I scored these individuals as 1 ("Delayed repair"). Over successive 1 h

periods I removed one-by-one the immediate/delayed repairers in the sequence that they began

repairing, and observed any changes in behaviour in remaining females. Some individuals

remained inactive within the nest for the duration of the experiment, even if another female was

repairing damage only a few millimetres away, and never began repairing. I scored these as 0

("Never repaired").

Removed individuals were dissected, and the extent of development of their ovaries was

examined, according to protocols described in Gilbert et al (2018). I recorded the number and

volume ($\pi \times length \times (width/2)^2$) of any developing oocytes, and also of any mature, chorionated

eggs that were ready to lay.  I also measured female body size (pronotum width).  I tested for

differences in ovarian development according to repair tendency using Kruskal-Wallis tests.

*Do nonreproductives breed when they inherit the domicile?*

Our removal experiment can be summarized briefly as follows: In domiciles with 2 or 3

individuals, I first categorised all females as "repairer" or "nonrepairer" according to their

tendency to repair an experimentally damaged nest. Then I experimentally removed a subset of

females from each nest (1 or 2 females per nest, so as to leave 1 behind) and dissected them to

assess the reproductive status of their ovaries.  I then assigned nests randomly to treatment

groups in which I replaced either a repairer (+R group), or a nonrepairer (+NR group), thereby

allowing (or forcing) this female to "inherit" her domicile. This remaining female I left in the nest

for 21 days to assess (1) her efficacy in repairing the damage, and (2) her realized reproduction

over this period. After this period, the remaining female was removed and dissected. The two

treatment groups, +R and +NR, were compared to control groups consisting of singly built nests

(CS group) and nests built by female pairs (CP group).

I identified *Dunatothrips aneurae* domiciles on thin-phyllode variants of *A. aneura* trees in the

field containing either 2 or 3 dealate females, and which were at a stage of near-complete

construction (i.e. without eggs or larvae).  Foundress females were initially counted within intact

domiciles non-invasively, using a powerful LED (LED Lenser®) shone through the silk from

behind; these preliminary counts and the absence of eggs were confirmed following

experimental nest damage (see below). Where possible, nests were chosen that were simple in

shape without convolutions that might complicate experimental removals, or which might

prevent individuals from detecting experimental damage to the domicile silk.

To test individual contributions to repair, I slightly damaged the domicile wall by peeling away

approximately 10-20% of its surface area using watchmaker's forceps. I then observed the

behaviour of the inhabitants for the next 3 h or until both (or all) individuals had responded by

repairing the domicile. Repair behaviour is very obvious in this tiny insect, even in the field, as

the female prepares to lay down a strand of silk by vigorously waving her abdomen; the function

of this waving is probably either mechanical (pumping the abdomen) or sensory (locating with

abdominal hairs the surfaces to be tied together). During this 3 h period, near-constant

observation (at least every 30 seconds) allowed me to keep track of individuals without marking

them. At the low humidities typical of the Outback environment, 6 h is long enough for small

larvae to die of desiccation when exposed outside a domicile (Gilbert 2014), so 3 h represents

an amount of time within which repairing a nest can have meaningful effects upon larval

survival.

Reflecting my lab classification of repair behaviour, I classified individuals into repairers (R) and

nonrepairers (NR). An individual was classified as a repairer if, during the observation period, it

added more than a negligible amount of new silk to the damaged area (I disregarded a minority

of individuals that added one or a few silk strands and then became quiescent). A two-female

domicile could thus be classified as (R,R), (NR, R) or (NR, NR). In practice (NR, NR) domiciles

were rare; I only encountered 2 during my fieldwork.

I randomly allocated the domiciles to three experimental groups. In all cases, I initially removed

both (or all) individuals from the domicile, and then replaced different individuals depending on

the experimental group.  Both removal and replacement were done using a coarse hair probe

and replacement was completed within 30 minutes of removal.  In the "Leave Repairer" group

(+R, n=11) I removed both (or all) individuals from the domicile and then replaced one repairer.

In the "Leave Nonrepairer" group (+NR, n=8) I instead replaced one nonrepairer. In the control

pairs group (CP, n=9) I replaced both (or all) individuals.  In the control singletons group (CS,

n=46), I identified early-stage colonies that were singly founded and performed the same

experimental damage procedure, before removing and replacing the foundress female.  All

permanently removed individuals were taken at this point to the lab for dissection to determine

ovarian development (see above).

Experimental domiciles were monitored each day in the field for 21 days. I noted the time taken

to repair the experimental damage by visually categorising the state of repair as "no repair",

"incomplete repair" or "complete repair".  "No repair" was where a negligible number of silk

strands had been added to the damaged area and the area remained substantially open to the

environment.  "Incomplete repair" was judged to be where repair had begun but strands were

sparse and the damaged area was either (i) not completely covered by silk strands, or (ii)

covered very loosely by silk such that any perturbation of the phyllodes (e.g. by mild wind)

would likely reopen the hole in the silk. "Complete repair" was where the new silk completely

covered the damaged area to the extent that gentle perturbation of the phyllodes would be

unlikely to reopen the hole in the domicile wall. I also recorded any instances of naturally

occurring domicile damage (recording the repair of this damage in the same way as above),

domicile destruction, or abandonment during the experimental period.

After a period of 21 days, experimental domiciles were collected and taken to the lab. 21 days is

the maximum time between domicile completion and egg-laying seen previously in the lab

(Gilbert and Simpson 2013), and I assumed an individual's reproduction in the field during that

time would be a reliable, standardized snapshot of its egg-laying rate, as a proxy for fitness. The

number of eggs and larvae in each domicile was counted and the latest stage of larvae attained

was noted. I excluded from analysis domiciles that were blown down, predated or destroyed

over the experimental period.

To analyse both repair times and numbers of offspring, I used generalised linear models (GLM)

with poisson errors and "treatment" as a single predictor variable. I compared the treatment

groups using planned orthogonal contrasts:

1.  Control pairs (CP group) against all singletons (pooled mean of CS, +R and +NR

259           groups).

260       2.  Experimental nonrepairers (+NR group) against all other singletons (pooled mean of CS

261           and +R groups)

3.  Experimental repairers (+R group) against control singletons (CS group).

I dissected all removed individuals as described above, recording number and volume of

developing oocytes. All individuals remaining in domiciles after the 21 d experimental period

were also dissected. Owing to time constraints, only a subset of the CS group (n=8) was used

for this part of the experiment.

I asked whether R and NR individuals from experimental domiciles differed in reproductive

status, both before (i.e. individuals removed immediately) and after the experiment (individuals

left in the domicile for 21 days).

I used linear models of the total volume of developing oocytes (log-transformed after adding the

minimum nonzero value in the dataset to all values; the precise value of this small added

increment did not affect the results). As predictor variables, I included main effects of repair

tendency (R versus NR) and treatment (before versus after), plus their interaction.

The treatment term in this analysis was perfectly confounded by the number of females

cohabiting in the domicile at the time of removal ("before" individuals all had >1 females in each

domicile at the time of removal, while "after" individuals uniformly had only 1 individual per

domicile). My experiment was unable to separate these two confounded variables, but I was

able to shed partial light on the problem by conducting a second analysis that also included

individuals in the CS and CP groups.  All of these individuals had been "left" for the entire

experimental period, but they differed in the number of females in the nest: 1 for the CS group

and >1 for the CP group.  For this second analysis, instead of treatment ("removed" versus

"left"), I instead used the number of females in the domicile at the time of removal ("1" versus

">1") as a predictor variable.

RESULTS

*Are reproduction and repair divided?*

24 females in the lab experiment (14 repairers and 10 nonrepairers) had no developing oocytes

and were assigned an oocyte volume of zero. Among the remaining 118 females dissected, the

volume of developing oocytes varied from $6.0 \times 10^{-4}$ to $3.0 \times 10^{-2}$ mm$^3$.  65 females had no mature

eggs; among the remainder, the volume of chorionated oocytes varied from $1.6 \times 10^{-3}$ to $2.8 \times 10^{-2}$

mm$^3$.  Pronotum width varied from 240 to 360 μm; body length varied from 1625 to 2422 μm. As

in Gilbert et al (2018), body size was not related to reproductive state (Kruskal-Wallis test, H =

0.64, df = 1, p-value = 0.72).

I found that females were more likely to repair (whether immediate or delayed) if they had

developing eggs than if they did not, and were much more likely to repair if they had mature,

chorionated eggs (X-squared = 9.52, df = 4, p-value = 0.049, Figure 1). Body size was not

related to repair tendency (Kruskal-Wallis test, H = 0.40, df = 1, p-value = 0.53).

Females that participated in nest repair had only a marginally (nonsignificantly) higher volume of

developing oocytes than those that never participated in repair (Kruskal-Wallis test, H=3.41,

df=1, p=0.06), and the number of developing oocytes was not associated with repair (Kruskal-

Wallis test, H=1.55, df=1, p=0.21).  In contrast, females that participated in nest repair had both

a higher number (Kruskal-Wallis test, H = 5.24, df = 1, p-value = 0.022) and volume of mature

eggs (Kruskal-Wallis test, H = 4.285, df = 1, p-value = 0.038) than those that did not.  This

pattern was mainly driven by the frequency of females with zero mature eggs: when I restricted

the dataset to only those females carrying mature eggs, there was then no difference between

repairers and nonrepairers in either number of mature eggs (Kruskal-Wallis test, H= 0.036, df =

1, p-value = 0.85) or volume  (H= 0.17466, df = 1, p-value = 0.676).

*Do nonreproductives breed when they inherit the domicile?*

Of 21 experimental domiciles with 2 females, 8 were categorised as (R, R), 11 were (NR, R)

and 2 were (NR, NR). Of 9 domiciles with 3 females, 1 was (R, R, R), 3 were (R, R, NR) and 5

were (R, NR, NR). All 8 singleton nests in the CS group whose repair activity I assessed in the

field were (R). Among repairers, latency to repair varied from <30 seconds to 126 m within the 3

h observation period, while "nonrepairers" were those that failed to repair, or repaired negligibly,

within this period.

*Time to repair damage*

There were significant differences in repair time among groups (GLM, poisson errors, dropping

"group", Δdeviance=43.76, Δdf=3, p=0.004). These differences were driven entirely by the +NR

group (Figure 2a). Domiciles in the control (CP and CS) and +R groups were completely

repaired after a median of 2 d following damage, with some completed by the next day. In

contrast, the +NR group took a median of 4 d to complete repair (range 2 – 7 d). Contrast 1 was

marginally significant (control pairs against all singleton groups: CP against pooled mean of CS,

+R and +NR; z=1.651, p=0.099). Contrast 2 was not significant (Experimental repairers, +R

group, against control singletons, CS group; z=1.022, p=0.307). Contrast 3 was significant

(Experimental nonrepairers, +R group, against pooled mean of CS and +R groups; z=-3.048,

p=0.002).

*Frequency of domicile damage*

There were no differences in the frequency of domiciles destroyed or abandoned during the

experimental period: 16 out of 46 domiciles in the CS group, 3 out of 9 domiciles in the CP

group, 3 out of 11 domiciles in the +R group and 4 out of 9 in the +NR group ($\chi^2$=0.625,

p=0.884).

*Number of offspring*

There were significant differences among groups in the numbers of offspring present after 21

days (GLM with poisson errors, dropping "group", Δdeviance=49.09, Δdf=3, p<0.0001; Figure

2b). CP had more offspring than all other groups (Contrast 1, CP against pooled mean of CS,

+R and +NR; z=5.360, p<0.0001). Experimental repairers were similar to control singletons in

productivity(Contrast 2, +R against CS; z=-1.383, p=0.167). The +NR group had substantially

fewer offspring than all other groups (Contrast 3, +NR group against pooled mean of CS and +R

groups; z=-4.225, p<0.0001).

*Reproductive status*

Out of the dissected females in my field experiment, 4 females (3 repairers and 1 nonrepairer)

had no developing oocytes and were assigned an oocyte volume of zero. Among the remainder

(n=64), the volume of developing oocytes varied from $5.3 \times 10^{-4}$ to $3.4 \times 10^{-2}$ mm$^3$.  22 had no

mature eggs, and among the remainder (n=41; 5 females belonging to control groups were not

assessed for mature eggs) varied from $5.3 \times 10^{-4}$ to $3.4 \times 10^{-2}$ mm$^3$.

Individuals of different repair tendency (R versus NR) did not significantly differ in ovarian

status, either in the first analysis (using "treatment" as a co-predictor) or in the second analysis

(incorporating the CS and CP groups and using "number of females in domicile" as a co-

predictor). In the first analysis, there was no interaction between repair tendency and treatment

(dropping this interaction, $F_{1,29}$=0.01, p=0.929), nor any statistical effect of repair tendency

(dropping repair tendency, $F_{1,30}$=0.179, p=0.676). However, treatment had a significant effect

(dropping treatment, $F_{1,30}$=5.99, p=0.020). In the second analysis, again there was no interaction

between repair tendency and female number (dropping interaction, $F_{1,47}$=0.217, p=0.643), nor

any effect of repair tendency (dropping repair tendency, $F_{1,48}$=0.01, p=0.923). As for "treatment"

in the first analysis, dropping female number had a significant upon model performance

(dropping female number, $F_{1,48}$=6.55, p=0.014). Individuals sharing domiciles with other females

(or which were removed at the beginning of the experiment) had a smaller volume of developing

oocytes than females alone in a domicile (or which were removed at the end of the experiment)

(Figure 3a). There were no significant differences in volume of mature eggs, whether among

treatments or repair tendencies (linear models, all NS; Figure 3b).

DISCUSSION

In multi-female domiciles of *D. aneurae*, helping effort and realized reproductive output were

positively associated: helpers were reproducers, and non-helpers were non-reproducers.

Although repairers and non-repairers had statistically similar amounts of developing oocytes

both in the lab or in the field, in the lab repairers had more mature eggs, and in the field they

ended up actually laying more eggs. A negative association between reproduction and helping

would have indicated reproductive division of labour, characteristic of a typically eusocial colony

(e.g. Vespid wasps; O'Donnell 1996) or of cooperative breeding vertebrates (Clutton-Brock et al.

2004; Russell et al. 2008).

Instead, with a positive association between realized reproduction and helping, joint-founded *D.*

*aneurae* domiciles resembled low-skew cooperative or communal societies in which group

members may help in proportion to their (i) genetic stake in current offspring (Curry 1988;

Packer et al. 2001) or probability of future breeding (Hartley et al. 1995; Gilchrist and Russell

2007; Leadbeater et al. 2011), or (ii) quality or condition (as proposed by West-Eberhard 1978;

Craig 1983). Between these alternatives, I found little evidence that nonreproductive females

were conserving energy for later breeding, suggesting instead that female quality may mediate

both helping and breeding effort in this system. Although repairer and nonrepairer females had

similar numbers/sizes of developing oocytes (Figure 1b), any given costly task (e.g. domicile

repair) may be more costly for females of lower quality, requiring them to resorb oocytes, and

consequently they may actually mature and lay fewer eggs (Figure 1c).

Reproductive competition is likely to exacerbate differences among females in realized

reproductive output. Competition occurs among *D. aneurae* females, especially in smaller

domiciles containing more females (Gilbert et al. 2018) and those which are occupied by

inquilines (Gilbert et al. 2012). In smaller domiciles, competition (for example from ecological

constraints preventing dispersal, or from successive joining of the domicile by additional

females) may force females of poorer quality into a nonreproductive or less reproductive state.

For these females, remaining in (or joining) an established domicile may be "the best of a bad

job", so that any small contribution stands a chance of being translated into offspring.

However, breeders gained no apparent advantage by sharing a domicile with nonrepairers.

Consistent with Gilbert et al (2018), reproductive females in this study had *fewer* developing

oocytes in cofounded domiciles than in singleton domiciles (Figure 3a), suggesting that

competition over resources within the domicile reduces fertility of all individuals, rather than

nonreproductives being of benefit to breeders, as occurs in some cooperative breeders (Nelson-

Flower Martha J. et al. 2013). Furthermore, control pairs and singletons repaired damage

equally quickly (Figure 2a). All experimental groups suffered similar rates of wind damage,

suggesting that domiciles do not differ in their vulnerability to strong wind, whether maintained

by one or two females, or by good versus bad repairers.

It therefore remains unclear whether subfertile females actually help, or whether they reduce

group fitness via competition. Research should now focus on testing between these two

alternatives. The subfertility hypothesis holds that naturally poor quality females can be selected

to invest exclusively in helping, even if their help is less efficient than that of breeders (Eberhard

1975; Craig 1983). Traditionally, the critical test of this hypothesis has been whether putative

helpers breed upon experimentally inheriting their natal nest or breeding resource. To my

knowledge, there has been no study in which subordinates failed to breed when given this

opportunity (e.g. (Field and Foster 1999; Smith et al. 2009; Rehan et al. 2014). If true, this study

would provide the first such case. However, previous studies have all been in taxa where

subordinates unequivocally act as helpers.

The question of why nonreproductive, apparently costly females should be tolerated by

reproductive females, posed and discussed initially by Gilbert et al (2018), therefore remains

open. FIrst, it is possible subfertile females may contribute to group fitness in subtle ways not

measured here. For example, Bono and Crespi (2006) found that joint-nesting females enjoyed

a survival advantage when invaded by kleptoparasitic thrips, providing a possible defensive role

for nonreproductive females. Nonreproductives may also participate more actively at the

building stage of the domicile than in ongoing maintenance, or may contribute to maintenance of

middens in mature domiciles (Gilbert and Simpson 2013). In other social insects, subfertile

individuals can have subtle but important effects upon colony function; for example, "soldiers"

have been recently discovered to play a primary role in combatting pathogens in both thrips

(Turnbull et al. 2012) and ants (Frank et al. 2018). Subordinates may also act via "load

lightening", extending longevity of breeders, even though immediate effects are not obvious

(Russell et al. 2007), or via "assured fitness returns" (Gadagkar 1990; Lucas and Field 2011)

whereby the effect of a given unit of help in a well-established group, however small, if positive,

is worth more to a subordinate's inclusive fitness than it would have been had she been

breeding on her own and therefore likely to fail.

A second alternative is that nonreproductive, nonrepairing female *D. aneurae* may be

reproductively suppressed, either behaviourally through aggression (e.g. Kolmer and Heinze

2000) or via pheromonal control or signalling, as in molerats and many social insects (e.g. Keller

and Nonacs 1993; Bennett et al. 1996). In some cooperative societies, putative "helpers"

actually perform less work than reproductives (Korb 2007; Browning et al. 2012), consistent with

the pattern I show here. This hypothesis is also consistent with the fact that nonreproductives

tend only to occur as a result of reproductive competition in small domiciles, as shown in Gilbert

et al (2018). However, the data shown here suggest this is unlikely. Theoretically, reproductively

suppressed individuals should withhold help only to conserve energy if they may inherit a

breeding position later (part of the "future fitness" hypothesis; Cant and Field 2001). However, in

*D. aneurae*, nonrepairing individuals neither bred nor repaired the domicile after removal of their

nestmate (Figure 2). Moreover, aggressive reproductive suppression in *D. aneurae* is unlikely

as I have never observed aggression apart from ejecting males after mating (Gilbert and

Simpson 2013), although chemical suppression is still possible.

A third and untested possibility remains that nonrepairing females were in 'akinesis' or

temporary diapause (Varadarasan and Ananthakrishnan 1982), i.e. waiting to reproduce *much*

later. This might be the case, for example, if females take a very long time to accumulate

resources for producing their relatively very large eggs (suggested by Gilbert et al 2018). If not

collected for dissection, nonrepairers may have "come online" at a point more than 21 days after

manipulation, and subsequently either dispersed and bred independently or bred within their

existing domicile. I regard these possibilities as unlikely. Regarding dispersal, reproductively

mature female *D. aneurae* are dealate (i.e. having lost their wings) (Gilbert and Simpson 2013),

reducing their capabilities of independent dispersal - all females involved in this study were

dealate. Regarding breeding within the domicile, it is very rare to observe females breeding

among the remains of an old brood (JDJG, pers. obs.); there is also as yet no evidence for

multiple generations of offspring within *D. aneurae* domiciles (see Gilbert and Simpson 2013) so

it is equally unlikely that nonreproductives were offspring of foundresses waiting to breed.

Finally, theoretically, public good games models suggest that "laggards" can coexist peacefully

(without punishment) with producers in situations where selection acts at the level of the group

as well as the individual (Boza and Számadó 2010), an idea with potential in the *Dunatothrips*

system where the likelihood of whole-group mortality is likely to be high relative to individual

mortality.

Primitively social or communal societies such as I have demonstrated here in *D. aneurae* have

sometimes been thought of as an intermediate step in the evolution of eusociality (Lin and

Michener 1972) although it has turned out that the two strategies are often phylogenetically

distinct (Kukuk 1992; Danforth and Eickwort 1997; but see Richards et al. 2003). Phylogenetic

separation appears also to be the rule in Acacia thrips, since the phyllode-tying habit occurs in a

clade separate from all other eusocial thrips (Crespi et al. 2004), but to confirm this idea, we

would need to investigate the relationship between breeding and helping in relatives of *D.*

*aneurae*, which are almost entirely unstudied. Domicile size and social tendencies vary widely

within the genus, with some being obligately solitary (D. gloius, Bono 2007), and others

moderately social such as *D. armatus* (Gilbert, unpublished data). The species that most

urgently requires study is *D. vestitor*, which is reported to form much larger societies with up to

70 individuals within domiciles that are loosely merged into "supercolonies" (Crespi et al. 2004

and JDJG, pers. obs.).

ACKNOWLEDGEMENTS

I thank Prof. S. J. Simpson for mentorship, guidance and use of his lab and facilities, Dr K

Leggett and G. and V. Dowling for helpful discussions, help during fieldwork and use of facilities,

Dr LA Mound and Dr A Wells for assistance with fieldwork and helpful discussions, Drs T and M

Flower, Dr LE Browning and Dr A Russell for helpful discussions and Dr P. Bierzychudek for

supplying micronized paint.

FIGURE CAPTIONS

Figure 1 (a) frequency of participation in domicile repair (black = no repair, grey = repair) by

female *D. aneurae* in different reproductive condition; (b) Volume of developing oocytes in

females according to participation in repair; (c) Volume of mature eggs in females according to

participation in repair.

Figure 2 (a) Time to repair damage in *D. aneurae* domiciles from different treatment groups. (b)

Number of eggs in the domicile 21 d after treatment. Bars with different letters are statistically

different. Sample sizes (domiciles) are given in parentheses. Boxes show median ± IQR,

whiskers show range, dashed line shows mean.

Figure 3 (a) Developing oocyte volume and (b) mature egg volume in females from different

treatment groups. Bars with different letters are statistically different. Sample sizes (females) are

given in parentheses. Boxes show median ± IQR, whiskers show 95% CI, dashed line shows

mean.

532    Figure 1

538    Figure 2

Figure 3

REFERENCES

Abrams J, Eickwort GC. 1981. Nest switching and guarding by the communal sweat
beeAgapostemon virescens (Hymenoptera, Halictidae) Le changement des nids et le
gardiennage par l'abeille Agaspostemon virescens (Hymenoptera, Halictidae). Insectes Soc.
28(2):105–116.

Baglione V, Canestrari D, Chiarati E, Vera R, Marcos JM. 2010. Lazy group members are
substitute helpers in carrion crows. Proc Biol Sci. 277(1698):3275–3282.

Bennett NC, Faulkes CG, Molteno AJ. 1996. Reproductive suppression in subordinate, non-
breeding female Damaraland mole-rats: two components to a lifetime of socially induced
infertility. Proc Biol Sci. 263(1376):1599–1603.

Beshers SN, Fewell JH. 2001. Models of division of labor in social insects. Annu Rev Entomol.
46:413–440.

Bono JM. 2007. Patterns of kleptoparasitism and inquilinism in social and non-social
Dunatothrips on Australian Acacia. Ecol Entomol. 32(4):411–418.

Bono JM, Crespi BJ. 2006. Costs and benefits of joint colony founding in Australian Acacia
thrips. Insectes Soc. 53(4):489–495. [accessed 2015 Sep 4].

Bono JM, Crespi BJ. 2008. Cofoundress relatedness and group productivity in colonies of social
Dunatothrips (Insecta: Thysanoptera) on Australian Acacia. Behav Ecol Sociobiol. 62(9):1489–
1498.

Boza G, Számadó S. 2010. Beneficial laggards: multilevel selection, cooperative polymorphism
and division of labour in threshold public good games. BMC Evol Biol. 10:336.

Browning LE, Young CM, Savage JL, Russell DJF, Barclay H, Griffith SC, Russell AF. 2012.
Carer provisioning rules in an obligate cooperative breeder: prey type, size and delivery rate.
Behav Ecol Sociobiol. 66(12):1639–1649. [accessed 2016 Dec 7].

Cant MA, Field J. 2001. Helping effort and future fitness in cooperation animal societies. Proc
Biol Sci. 268(1479):1959–1964.

Clutton-Brock TH, Russell AF, Sharpe LL. 2004. Behavioural tactics of breeders in cooperative

meerkats. Anim Behav. 68(5):1029–1040.

Craig CL, Hsu M, Kaplan D, Pierce NE. 1999. A comparison of the composition of silk proteins

produced by spiders and insects. Int J Biol Macromol. 24(2-3):109–118.

Craig R. 1983. Subfertility and the evolution of eusociality by kin selection. J Theor Biol.

100(3):379–397.

Crespi BJ, Morris DC, Mound LA, Others. 2004. Evolution of ecological and behavioural

diversity: Australian Acacia thrips as model organisms. Australian Biological Resources Study.

Curry RL. 1988. Influence of kinship on helping behavior in Galápagos mockingbirds. Behav

Ecol Sociobiol. 22(2):141–152.

Danforth BN, Eickwort GC. 1997. The evolution of social behavior in the augochlorine sweat

bees (Hymenoptera: Halictidae) based on a phylogenetic analysis of the genera. The evolution

of social behavior in insects and arachnids.:270–292.

Eberhard MJW. 1975. The Evolution of Social Behavior by Kin Selection. Q Rev Biol. 50(1):1–

33.

Field J, Foster W. 1999. Helping behaviour in facultatively eusocial hover wasps: an

experimental test of the subfertility hypothesis. Anim Behav. 57(3):633–636.

Frank ET, Wehrhahn M, Linsenmair KE. 2018. Wound treatment and selective help in a termite-

hunting ant. Proc Biol Sci. 285(1872). doi:10.1098/rspb.2017.2457.

Gadagkar R. 1990. Evolution of Eusociality: The Advantage of Assured Fitness Returns. Philos

Trans R Soc Lond B Biol Sci. 329(1252):17–25. [accessed 2016 Jan 14].

Gerber C, Badertscher S, Leuthold RH. 1988. Polyethism in Macrotermes bellicosus (Isoptera).

Insectes Soc. 35(3):226–240.

Gilbert JDJ. 2014. Thrips domiciles protect larvae from desiccation in an arid environment.

Behav Ecol. 25(6):1338–1346.

Gilbert JDJ, Mound LA, Simpson SJ. 2012. Biology of a new species of socially parasitic thrips

(Thysanoptera: Phlaeothripidae) inside Dunatothrips nests, with evolutionary implications for

inquilinism in thrips. Biol J Linn Soc Lond. 107(1):112–122.

Gilbert JDJ, Simpson SJ. 2013. Natural history and behaviour of Dunatothrips aneurae Mound
(Thysanoptera: Phlaeothripidae), a phyllode-gluing thrips with facultative pleometrosis. Biol J
Linn Soc Lond. 109(4):802–816.

Gilbert JDJ, Wells A, Simpson SJ. 2018. Skew in ovarian activation depends on domicile size in
phyllode-glueing thrips. Sci Rep. 8(1):3597.

Gilchrist JS, Russell AF. 2007. Who cares? Individual contributions to pup care by breeders vs
non-breeders in the cooperatively breeding banded mongoose (Mungos mungo). Behav Ecol
Sociobiol. 61(7):1053–1060.

Hartley IR, Davies NB, Hatchwell BJ, Desrochers A, Nebel D, Burke T. 1995. The
polygynandrous mating system of the alpine accentor, Prunella collaris. II. Multiple paternity and
parental effort. Anim Behav. 49(3):789–803.

Hayes LD. 2000. To nest communally or not to nest communally: a review of rodent communal
nesting and nursing. Anim Behav. 59(4):677–688.

Hoogland JL. 1983. Nepotism and alarm calling in the black-tailed prairie dog (Cynomys
ludovicianus). Anim Behav. 31(2):472–479.

Jandt JM, Huang E, Dornhaus A. 2009. Weak specialization of workers inside a bumble bee
(Bombus impatiens) nest. Behav Ecol Sociobiol. 63(12):1829–1836.

Jeanson R, Kukuk PF, Fewell JH. 2005. Emergence of division of labour in halictine bees:
contributions of social interactions and behavioural variance. Anim Behav. 70(5):1183–1193.

Johnson BR. 2010. Division of labor in honeybees: form, function, and proximate mechanisms.
Behav Ecol Sociobiol. 64(3):305–316.

Keller L, Nonacs P. 1993. The role of queen pheromones in social insects: queen control or
queen signal? Anim Behav. 45(4):787–794.

Koenig WD, Dickinson JL. 2004. Ecology and evolution of cooperative breeding in birds.
Cambridge University Press.

Kolmer K, Heinze J. 2000. Rank orders and division of labour among unrelated cofounding ant
queens. Proc Biol Sci. 267(1454):1729–1734.

Komdeur J. 2006. Variation in individual investment strategies among social animals. Ethology.
112(8):729–747.

Korb J. 2007. Workers of a drywood termite do not work. Front Zool. 4:7.

Kukuk PF. 1992. Social interactions and familiarity in a communal halictine bee Lasioglossum
(Chilalictus) hemichalceum. Ethology. 91(4):291–300.

Leadbeater E, Carruthers JM, Green JP, Rosser NS, Field J. 2011. Nest inheritance is the
missing source of direct fitness in a primitively eusocial insect. Science. 333(6044):874–876.

Lin N, Michener CD. 1972. Evolution of Sociality in Insects. Q Rev Biol. 47(2):131–159.

Lucas ER, Field J. 2011. Assured fitness returns in a social wasp with no worker caste. Proc
Biol Sci. 278(1720):2991–2995.

Morris DC, Schwarz MP, Crespi BJ. 2002. Pleometrosis in phyllode-glueing thrips
(Thysanoptera: Phlaeothripidae) on Australian Acacia. Biol J Linn Soc Lond. 75(4):467–474.

Nelson-Flower Martha J., Hockey Philip A. R., O'Ryan Colleen, English Sinead, Thompson Alex
M., Bradley Katharine, Rose Rebecca, Ridley Amanda R. 2013. Costly reproductive competition
between females in a monogamous cooperatively breeding bird. Proceedings of the Royal
Society B: Biological Sciences. 280(1762):20130728.

O'Donnell S. 1996. Reproductive potential and division of labor in wasps: are queen and worker
behavior alternative strategies? Ethol Ecol Evol. 8(3):305–308.

Packer C, Pusey AE, Eberly LE. 2001. Egalitarianism in female African lions. Science.
293(5530):690–693.

Page RE Jr. 1997. The evolution of insect societies. Endeavour. 21(3):114–120.

Reeve HK, Keller L. 1995. Partitioning of Reproduction in Mother-Daughter Versus Sibling
Associations: A Test of Optimal Skew Theory. Am Nat. 145(1):119–132.

Rehan SM, Richards MH, Adams M, Schwarz MP. 2014. The costs and benefits of sociality in a

facultatively social bee. Anim Behav. 97:77–85.

Richards MH, von Wettberg EJ, Rutgers AC. 2003. A novel social polymorphism in a primitively

eusocial bee. Proc Natl Acad Sci U S A. 100(12):7175–7180.

Russell AF, Langmore NE, Cockburn A, Astheimer LB, Kilner RM. 2007. Reduced egg

investment can conceal helper effects in cooperatively breeding birds. Science. 317(5840):941–

944.

Russell AF, Langmore NE, Gardner JL, Kilner RM. 2008. Maternal investment tactics in superb

fairy-wrens. Proc Biol Sci. 275(1630):29–36.

Smith AR, Kapheim KM, O'Donnell S, Wcislo WT. 2009. Social competition but not subfertility

leads to a division of labour in the facultatively social sweat bee Megalopta genalis

(Hymenoptera: Halictidae). Anim Behav. 78(5):1043–1050.

Solomon NG, French JA. 1997. Cooperative breeding in mammals. Cambridge University

Press.

Strassmann JE, Queller DC. 1989. Ecological determinants of social evolution. WESTVIEW

STUD INSECT BIOL.:81–101.

Sullivan JD, Strassmann JE. 1984. Physical variability among nest foundresses in the

polygynous social wasp, Polistes annularis. Behav Ecol Sociobiol. 15(4):249–256.

Trumbo ST, Wilson DS. 1993. Brood discrimination, nest mate discrimination, and determinants

of social behavior in facultatively quasisocial beetles (Nicrophorus spp.). Behav Ecol. 4(4):332–

339. [accessed 2018 Mar 19].

Turnbull C, Caravan H, Chapman T, Nipperess D, Dennison S, Schwarz M, Beattie A. 2012.

Antifungal activity in thrips soldiers suggests a dual role for this caste. Biol Lett. 8(4):526–529.

Varadarasan S, Ananthakrishnan TN. 1982. Biological studies on some gall thrips. In: Proc.

Indian Acad. Sci. B. Vol. 48. insa.nic.in. p. 35–43.

Vehrencamp SL. 1979. The Roles of Individual, Kin, and Group Selection in the Evolution of

Sociality. In: Marler P, Vandenbergh JG, editors. Social Behavior and Communication. Boston,

MA: Springer US. p. 351–394.

Vehrencamp SL. 1983a. Optimal Degree of Skew in Cooperative Societies. Integr Comp Biol.
23(2):327–335. [accessed 2018 Mar 19].

Vehrencamp SL. 1983b. A model for the evolution of despotic versus egalitarian societies. Anim
Behav. 31(3):667–682.

West-Eberhard MJ. 1978. Temporary queens in metapolybia wasps: nonreproductive helpers
without altruism? Science. 200(4340):441–443.

West SA, Griffin AS, Gardner A. 2007. Social semantics: altruism, cooperation, mutualism,
strong reciprocity and group selection. J Evol Biol. 20(2):415–432.

685