

## **Prevalence of Stunting and Associated Factors among public Primary School pupils of Bahir Dar city, Ethiopia: School-Based Cross-Sectional Study**

<sup>1</sup>\*Getasew Mulat Bantie, <sup>2</sup>Kidist Hailu Akenew, <sup>2</sup>Mahlet Tilahun Belete, <sup>2</sup>Eyerusalem Teshome Tena, <sup>2</sup>Genet Gebreselasie Gebretsadik, <sup>2</sup>Aynalem Nebebe Tsegaw, <sup>2</sup>Tigist Birru Woldemariam, <sup>3</sup>Ashenafi Abate Woya, <sup>4</sup>Amare Alemu Melese

<sup>1</sup>\*GMB: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

**\*corresponding author**

Email: [getasewmulat@gmail.com](mailto:getasewmulat@gmail.com)

Cell phone: +251-939699185

Po.box: +251-209

<sup>2</sup>KHA: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [kidisthole@gmail.com](mailto:kidisthole@gmail.com)

<sup>2</sup>MTB: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [mahtilahun26@gmail.com](mailto:mahtilahun26@gmail.com)

<sup>2</sup>ETT: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [jesus.jerry5@gmail.com](mailto:jesus.jerry5@gmail.com)

<sup>2</sup>GGG: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [genetgbreslase@gmail.com](mailto:genetgbreslase@gmail.com)

<sup>2</sup>ANT: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [aynalemnebebe@gmail.com](mailto:aynalemnebebe@gmail.com)

<sup>2</sup>TBW: Department of Nursing, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [tigbirru@gmail.com](mailto:tigbirru@gmail.com)

<sup>3</sup>AAW: Department of Statistics, Bahir Dar University, Bahir Dar, Ethiopia

Email: [ashu.abate@gmail.com](mailto:ashu.abate@gmail.com)

<sup>4</sup>AAM: Department of Medical Laboratory Science, GAMBY Medical and Business College, Bahir Dar, Ethiopia

Email: [Amare.vip@gmail.com](mailto:Amare.vip@gmail.com)

## ABSTRACT

**Background:** Stunting is a well-established pupils-health indicator of chronic malnutrition which reliably gives a picture of the past nutritional history and the prevailing environmental and socioeconomic circumstances.

**Objective:** The prevalence of stunting and associated factors among public primary school pupils of Bahir Dar city.

**Method:** A cross-sectional study was carried out from March to June 2019. Data were coded and entered into Epi-Data and exported to SPSS. Then, Anthropometric data were converted into height for age Z-scores to determine the pupils stunting outcomes using WHO Anthro-Plus software. Then, the final analysis was done by SPSS version 20 software. Anthropometric measurements determined the proportion of stunting (z-score of height for age less than minus two standard deviations from WHO Anthro-plus software output). A simple logistic regression model was fitted to identify factors associated between the independent variables and the dependent variable at a 95% confidence level and p-value <0.05.

**Results:** 370 primary school pupils were included in the study with the mean age of 121.84(± 26.67) months. About 51.6% of the pupils were females. The total prevalence of stunting was 15.13% (95%CI; 11%, 19%). The burden of stunting was higher in the age group of greater than 132 months. Pupil's age  $\geq 132$  months (AOR=15.6; 95%CI; 3.31, 73.45; p-value<0.001) and male pupils (AOR=7.07; 95%CI: 2.51, 19.89; p-value<0.0002) were significantly associated with stunting.

**Conclusion:** The prevalence of stunting was relatively lower than the regional estimated stunting level. However, this result is also very significant figure to get critical attention. Pupil's age  $\geq 132$  months and male sex were significantly associated with stunting.

**Keywords:** Stunting, Anthropometric measurement, school-age pupils, malnutrition, Bahir Dar city.

## INTRODUCTION

School pupils pass through great physical and mental changes, which affect both their growth and school performance(1) and nutrition is one of the many key factors affecting mental development of children (2).

Undernutrition is a condition that results from eating a diet which has either inadequate nutrient to the extent that causes health problems(3). Under nutrition is widespread among school children particularly in low income countries particularly in Africa and their nutritional status often worsens during their school years. It greatly affects both the cognitive and physical development of school age children(4). It is known that it may increase susceptibility to infection(5) undernutrition among children has lifelong impacts; its effect could also transfer from the current generation to future.

Under nutrition related health challenges in school-age children are among the most frequent reasons of low school enrolment, high absenteeism, early dropout, poor academic performance, delayed cognitive development, short stature, reduced work capacity, and poor reproductive performance(4) According to some studies done in rural Ethiopia, the extent of stunting in school-age children ranges from 26.5% to 42.7%(6). Among the urban dwellers of this group, the range of stunting was 5.4% to 29.2(7).

Different studies have been done on factors determining undernutrition, but still, there are inconsistencies across the studies especially primary school children.

Therefore, the main aim of this study is to determine prevalence of stunting and associated factors among public primary school children of Bahir Dar city. The results of this study may reveal the existing prevalence and determinants of stunting among public primary school children of Bahir Dar city.

## **METHODS AND MATERIALS**


### **Study design, setting and period**

A school-based cross-sectional study was conducted in Bahir Dar city public primary school pupils from March 10 to June 10, 2019. Bahir Dar city is located 565 km away from Addis Ababa, the capital city of Ethiopia. The city is administratively had six sub-cities. There are two governmental and two private hospitals, eight governmental health centers, thirty-six private clinics and one governmental University, four private and two governmental colleges are found in the city. The total population of the city is estimated to be 249,851 (124,426 females and 125,425 males), according to the population and household survey. According to Bahir Dar city education department of 2018/19 report, there are 19 public primary schools (1 up to 8 graded).

### **Population, sample and sampling procedure**

The source population for this study consisted of all pupils attending primary schools of Bahir Dar city. The study population for this study was pupils attending the selected primary schools of the city were included in the study. Those with drop out or absent during the data collection period were excluded. The sample size was determined using a single population proportion formula by considering; 95% confidence level, 5% margin of error, stunting level of 18.3% in Bahir Dar city(8). Taking the design effect 1.5 and 10% non-response rate and considering the correction formula ( $N < 10,000$ ), the final sample size was 375. Then, multistage sampling was used to select eligible pupils. Five primary schools were recruited for the study from 19 total primary schools (1 up to 8 graded) by lottery method. From the five selected schools, primary cycle (1 up to 4 graded) pupils were recruited for the study. Then, systematic sampling technique with proportional allocation to size was applied from the registration list to select 375 samples.

**(Figure 1)**



**Figure 1: Schematic presentation of the sampling procedure in Bahir-Dar city, 2019**

### **Data collection procedures and measurement of variables**

Data collection tools initially prepared in English then translated to local Amharic language. Six BSC nurses (four for data collection and two for supervision) were recruited. Two days training was given for data collectors and supervisors by the principal investigator about the objective of the study. Socio-demographic characteristic, health care characteristic, maternal awareness on dietary diversification, and household food security status-related data were collected using a pretested structured interviewer-administered questionnaire. Anthropometric data such as weight, height, and body mass measurements were taken from each study participants. After anthropometric measurements were taken, study participants' parents were interviewed with face to face based on the objective of the study. Participants were classified as stunted if they had z-score of height for age less than minus two standard deviations from WHO Anthro-plus software output.

## Data Processing and analysis

Data were coded and entered into Epi-Data and exported to SPSS version 20 software and then, to WHO Anthro-Plus software. Again back to SPSS for analysis. Descriptive statistics were used to summarize characteristics of study participants and presented using text and table. Bivariate and multivariate logistic regression analysis was employed to assess the association between the exploratory variables and stunting. The strength of the association was measured using the adjusted odds ratio (AOR) and 95% confidence interval (CI). A p-value < 0.05 was considered as a statistically significant predictor of stunting.

## RESULT

### Socio-demographic characteristics of the participants

A total of 370 primary school pupils were recruited for the study with a response rate of 98.6%. The mean age of the respondents was 121.84(± 26.67) months. About 51.6% of the stunted pupils were females. About 35% of households with more than ten family size got their child stunted. 20% of the merchant parents had a stunted child. About 15% of the mothers were not able to read and write and 20% had family monthly income of fewer than 500 birrs. Similarly, more than 15% of the child was underweight (15.5%) and wasted (17.9%), respectively. (Table 1)

**Table 1: Socio-demographic related characteristics of the participants, Bahir Dar city, North West Ethiopia, 2019**

Variable	Category	Stunted	Not-stunted
		N(%)	N(%)
Age of the pupil (in months)	72-96	90(92.8)	7(7.2)
	108-120	116(89.2)	14(10.8)
	≥ 132	108(75.5)	35(24.5)
Sex of the pupil	Female	22(11.5)	169(88.5)
	Male	34(19.0)	145(81.0)
Religion	Orthodox	49(14.7)	283(85.3)
	Islam	7(22.6)	24(77.4)
	Catholic	0(0)	2(100)

	Protestant	0	5(100)
Family size of the household	1-5	27(13.6)	172(86.4)
	6-10	22(14.6)	129(85.4)
	11-15	7(35)	13(65)
Mother's occupation	Merchant	98(20)	36(80)
	Government employee	5(11.4)	39(88.6)
	Self-employed	26(18.3)	116(81.7)
	Housewife	15(11.1)	120(88.9)
Mother's educational status	Can't read and write	18(15.5)	98(84.5)
	Can read and write	23(17.8)	106(82.2)
	Primary school complete	7(11.9)	52(88.1)
	Secondary school and above	7(11.3)	55(88.7)
Monthly income( in Birrs)	<500	12(20.3)	47(79.7)
	500-1000	13(14.1)	79(85.9)
	1000-1500	8(13.6)	51(86.4)
	> 1500	23(14.4)	137(85.6)
Body mass index	Under weight	42(15.3)	232(84.7)
	Normal	9(15.2)	50(84.8)
	Over weight	1(50)	1(50)
Middle upper arm circumference	Wasting	27(17.9)	124(82.1)
	Normal	29(13.2)	190(86.8)

### Household food security, dietary diversification, and health care

22.2% of the parents who were worried for fear of not having enough food in the household had their child stunted. Similarly, those parents who shared foods from relatives when a shortage of food happens were their child stunted. Family members who didn't get enough food were 19% their child stunted. 15% of stunted children seen in households faced a shortage of food. The main reason, (24.1%) for this shortage was due to low family income were stunted.

24.5% of the stunted children mothers' did not participate in their child feeding. The families who didn't hear about the variety of food were 21.1% their child stunted. 17.6% of stunting was observed in the family members who had a chronic disease. Similarly, 16.7% of currently ill study children were stunted (**Table 2**)

**Table 2: Household food security, dietary diversification, and healthcare-related characteristics of the participants, Bahir Dar city, North West Ethiopia, 2019**

Variable	Category	Stunted N(%)	Not-stunted N(%)
Did you worry for fear of not having enough* household food	Yes	35(22.2)	123(77.8)
	No	21(9.9)	191(90.1)
If the family get a shortage of food what is your measurement	Share with relatives	12(22.2)	42(77.8)
	Debt	14(13.2)	92(86.8)
	Work hard	20(16.1)	104(83.9)
	Do nothing	4(9.3)	39(90.7)
	Asking for donation	6(14)	37(86)
In your judgment is there anyone who didn't get enough food* in your family	Yes	23(19)	98(81)
	No	33(13.7)	216(86.3)
Do you face shortage of food	Yes	22(15)	125(85)
	No	34(15.2)	189(84.8)
Reason for food shortage	Divorce	4(10.9)	33(89.1)
	Fool about	2(8)	23(92)
	In come shortage	14(24.1)	44(75.9)
	Lack of peace in the county	0(0)	13(100)
	Economical in stability	2(14.3)	12(85.7)
The mother participate in child feeding	Yes	44(13.7)	277(86.3)
	No	12(24.5)	37(75.5)
Do you hear about variety of food	Yes	52(14.8)	299(85.2)
	No	4(21.1)	15(78.9)
Source of information	Health care provider	22(13.8)	138(86.3)
	Family members	5(8.1)	57(91.9)
	Television	17(23.0)	57(77.0)
	Radio	3(21.4)	11(78.6)
Variety of food is good for your children	Yes	55(15.1)	309(84.9)
	No	1(16.7)	5(83.3)
Where will you go when your child gets sick	Health institution	39(14.8)	225(85.2)
	Traditional medicine	4(10.5)	34(89.5)
	Holly water	13(19.7)	55(80.3)
Family member who had any chronic disease**	Yes	6(17.6)	30(83.3)
	No	50(15.0)	284(85.0)
Current health status of this child	Healthy	4(15.0)	20(85.0)


	Ill	52(16.7)	294(83.3)
--	-----	----------	-----------

Enough food\*(eating food at least three times per day); chronic disease\*\* (diabetes mellitus, hypertension and/or asthma)

### The prevalence of stunting among primary school pupils

This study revealed that 15.13% of the pupils of public primary school of the city were stunted.

(Figure 2)



**Figure 2: the prevalence of stunting among primary school pupils of Bahir-Dar city, 2019**

### Factors associated with stunting among primary school children

Multivariable logistic regression analysis was conducted to identify independent predictors of stunting among public primary school pupils. On multivariable logistic regression model, age and sex of the pupil were significantly associated with stunting at a p-value of 0.05.

Accordingly, those primary school children in the age group of > 132 months were about 15 times, (AOR=15.6; 95%CI: 3.31, 73.45) more likely to be stunted compared to 72-98 months' of age group. The odds of stunting among male pupils were about 7 times (AOR=7.07; 95%CI: 2.51, 19.89) higher compared to those female pupils. (Table 3)

**Table 3: Factors associated with stunting among primary school children in Bahir Dar city, North West Ethiopia, 2109**

Variable	Stunting		COR (95%CI)	AOR (95%CI)	P-value
	Yes	No			
<b>Age of the child (in months)</b>					
72-96	90	7	1.00	1.00	
108-120	116	14	1.55(0.6,4.0)	4.44(0.88,22.34)	0.070
≥ 132	108	35	4.17(1.76,9.83)	15.6(3.31,73.45)	0.001
<b>Family size</b>					
1-5	27	172	1.00	1.00	
6-10	22	129	1.08(0.59,1.99)	1.83(0.72,4.67)	0.199
11-15	7	13	3.43(1.26,9.36)	2.95(0.43,19.81)	0.266
<b>Worrying due to fear of not having enough household food</b>					
Yes	35	123	2.58(1.44, 4.65)		
No	21	191	1.00	1.00	
<b>Frequency of worrying due to fear of not having enough household food</b>					
Always	10	27	1.00	1.00	
Usually	7	57	0.33(0.11,0.96)	0.45(0.13,1.25)	0.116
Sometimes	35	123	1.24(0.49,3.11)	1.75(0.64,4.79)	0.274
<b>Current health status of the pupils</b>					
Healthy	4	20	1.00	1.00	
Ill	52	294	1.13(0.37,0.44)	1.83(0.41,8.08)	0.426
<b>Sex of the pupil</b>					
Female	22	169	1.00	1.00	
Male	34	145	1.80(1.00,3.21)	7.07(2.51,19.89)	0.0002

## DISCUSSION

The present study revealed that the prevalence of stunting among primary school pupils in Bahir Dar City Administration was 15.13% (95%CI; 11, 19). This finding was consistent with studies conducted at Rabwah, Pakistan, 11.5%(9) and Baghdad, Iraq, 18.7%(10), Southern Pakistan,16.5%(11), in Abeokuta, Southwest Nigeria,17.4%(12)

But the present result showed a lower prevalence than a study conducted at Jimma Zone, southwest Ethiopia,24.1% (13), and Urban Communities in Obafemi Owode, Southwestern Nigeria ,33.8%(14), Gondar town, Ethiopia, 46.1% (15), Humbo district, Southern Ethiopia, 57%(16), west Bengal,23.0%(17), Western Kenya,36% (18), Mecha District, Amhara Regional State, 37.9%(19), southern Ethiopia,28%(20),southern Angola, 41.5%(5), in Cambodian,40.0%(2), in Arba Minch Health and Demographic Surveillance Site, Southern Ethiopia,41.9%(4)

On the contrary, the current study finding was higher than the study finding from Northcentral Nigeria 10.5%,(21), Tehran, Iran,3.7%(22), in China, 1.0-1.9%(23), Bogota, Colombia,19.8%(24), northern region of Sudan,7.1%(25), Argo city, Northern Sudan, 3.8%(1), in Lahore, Pakistan, 8%(26), in Jiangsu Province, China, 4.7%(27), in Beijing, China, 2.6%(28). The possible explanation might be the difference in socio-demographic characteristics like study setting and lifestyle and health information dissemination.

The current study also identified the risk factors associated with stunting among public primary school pupils in Bahir Dar city.

Pupils in the age group of 132 months and above were sixteen times more likely stunted than the age group of 72-96 months. This finding was supported by a study done in Southern highlands of Tanzania (29), Southwestern Nigeria(14), Baghdad, Iraq(10), Humbo district, Southern Ethiopia(16), Gondar town, Ethiopia(15), Southern Pakistan(11), Mecha District, Amhara Regional State(19), Lahore, Pakistan(26), Southern Angola(5),southwest Nigeria(12), and Arba Minch, Ethiopia(4). This might be due to the fact older children may suffer from the presence of younger siblings.

Male primary school pupils were sevenfold at higher risk for stunting than females; which was supported by a study finding from Southern highlands of Tanzania(29), and Arba Minch, Ethiopia (4) and EDHS 2016(30). In contrast, the study finding from Beijing-China(28) and Southern Pakistan,(11) revealed that females were more likely at risk of being stunted than males. This might be because males' growth and development is more influenced by environmental and nutritional stress than females and thus, making males more likely to be affected by stunting (4).

### **Limitation of the study**

There might be a potential bias among respondents answering questions relating to events happening in the past such as food availability. Measurement bias might also occur during data collection.

### **Conclusion**

The prevalence of stunting was relatively lower than the regional estimated stunting level. However, this result was a very significant figure to get critical attention. Pupil's age  $\geq 132$  months and male sex were significantly associated with stunting.

### **Acronyms/Abbreviations**

AOR: Adjusted Odds Ratio; BMI: Body Mass Index; CI: Confidential Interval; EDHS: Ethiopian Demographic and Health Survey; SD: Standard Deviation; WHO: World Health Organization

## **Ethical considerations**

Ethical clearance was obtained from GAMBY Medical and Business College, Research and Publication Office with the reference number of GC-221/2011. The support letter was obtained from Bahir Dar city health and education department before the beginning of data collection. The directors of each primary school was informed about the purpose of the study that it will contribute for the health promotion and intervention of the primary school children of the study area and in the nation at large. Before collecting the data, parents of the students were informed and written consent was obtained from them and finally the parents were interviewed. Names of the students and/ or parents did not use to ensure anonymity and confidentiality.

## **Availability of data and materials**

The data can be accessed from the corresponding author

## **Consent to publish**

Written consent was obtained that the interview will be included in publications

## **Competing interests**

Authors declare that there are no competing interests.

## **Funding**

No fund was obtained

## **Authors' Contributions**

GMB, KHA, MTB, ETT, GGG, ANT, TBW AAW, and AAM involved in the design, data collection, statistical analysis and interpretation, and manuscript drafting and critical revision. All authors have an equal contribution.

## **Acknowledgments**

We would like to thank our data collectors and the supervisor for their invaluable effort; our deep gratitude also goes to our study participants who volunteered and took their time to give us all the relevant information for the study. Last but not least, we would like to thank all Bahir Dar city primary school directors for their cooperation and support during the data collection.

## REFERENCE

1. Hussein MD, Alonazi NA, Mohamed S. Prevalence of obesity, overweight, underweight, and stunting among school children in Argo city, Northern Sudan. *Sudan J Paediatr*. 2018;18(2):15-9.
2. Perignon M, Fiorentino M, Kuong K, Burja K, Parker M, Sisokhom S, et al. Stunting, poor iron status and parasite infection are significant risk factors for lower cognitive performance in Cambodian school-aged children. *PLoS One*. 2014;9(11):e112605.
3. WHO. Malnutrition. WHO. 2018;1-5.
4. Tariku EZ, Abebe GA, Melketsedik ZA, Gutema BT. Prevalence and factors associated with stunting and thinness among school-age children in Arba Minch Health and Demographic Surveillance Site, Southern Ethiopia. *PLoS One*. 2018;13(11):e0206659.
5. Oliveira D, Ferreira FS, Atouguia J, Fortes F, Guerra A, Centeno-Lima S. Infection by Intestinal Parasites, Stunting and Anemia in School-Aged Children from Southern Angola. *PLoS One*. 2015;10(9):e0137327.
6. Mekonnen A, Tefera B, Woldehanna T, Jones N, Seager J, Alemu T, et al. Child nutritional status in poor Ethiopian households. 2005.
7. Tadesse G. The prevalence of intestinal helminthic infections and associated risk factors among school children in Babile town, eastern Ethiopia. *Ethiopian Journal of Health Development*. 2005;19(2):140-7.
8. Hailegebriel T. Undernutrition, intestinal parasitic infection and associated risk factors among selected primary school children in Bahir Dar, Ethiopia. *BMC infectious diseases*. 2018;18(1):394.
9. Ahmad MS, Husain Zaidi SA, Medhat N, Farooq H, Ahmad D, Nasir W. Frequency of underweight and stunting among children entering school in a small urban locality and their association with academic performance. *J Pak Med Assoc*. 2018;68(1):28-32.
10. Al-Saffa AJ. Stunting among primary-school children: a sample from Baghdad, Iraq. *East Mediterr Health J*. 2009;15(2):322-9.
11. Khuwaja S, Selwyn BJ, Shah SM. Prevalence and correlates of stunting among primary school children in rural areas of southern Pakistan. *J Trop Pediatr*. 2005;51(2):72-7.
12. Senbanjo IO, Oshikoya KA, Odusanya OO, Njokanma OF. Prevalence of and risk factors for stunting among school children and adolescents in Abeokuta, southwest Nigeria. *J Health Popul Nutr*. 2011;29(4):364-70.

13. Abate KH, Belachew T. Care and not wealth is a predictor of wasting and stunting of 'The Coffee Kids' of Jimma Zone, southwest Ethiopia. *Nutr Health*. 2017;23(3):193-202.
14. Adenuga WU, Obembe TA, Odebunmi KO, Asuzu MC. Prevalence and Determinants of Stunting among Primary School Children in Rural and Urban Communities in Obafemi Owode Local Government Area, Southwestern Nigeria. *Ann Ib Postgrad Med*. 2017;15(1):7-15.
15. Getaneh Z, Melku M, Geta M, Melak T, Hunegnaw MT. Prevalence and determinants of stunting and wasting among public primary school children in Gondar town, northwest, Ethiopia. *BMC Pediatr*. 2019;19(1):207.
16. Bogale TY, Bala ET, Tadesse M, Asamoah BO. Prevalence and associated factors for stunting among 6-12 years old school age children from rural community of Humbo district, Southern Ethiopia. *BMC Public Health*. 2018;18(1):653.
17. Bose K, Bisai S. Prevalence of underweight and stunting among school children in West Bengal. *Indian J Pediatr*. 2008;75(12):1272.
18. Friedman JF, Phillips-Howard PA, Mirel LB, Terlouw DJ, Okello N, Vulule JM, et al. Progression of stunting and its predictors among school-aged children in western Kenya. *Eur J Clin Nutr*. 2005;59(8):914-22.
19. Lisanu Mazengia A, Andargie Biks G. Predictors of Stunting among School-Age Children in Northwestern Ethiopia. *J Nutr Metab*. 2018;2018:7521751.
20. Grimes JET, Tadesse G, Gardiner IA, Yard E, Wuletaw Y, Templeton MR, et al. Sanitation, hookworm, anemia, stunting, and wasting in primary school children in southern Ethiopia: Baseline results from a study in 30 schools. *PLoS Negl Trop Dis*. 2017;11(10):e0005948.
21. Adedeji IA, Bashir MF, Shwe DD, John C. Prevalence and correlates of stunting among the school-age population in North-Central Nigeria. *Pan Afr Med J*. 2018;31:170.
22. Esfarjani F, Roustae R, Mohammadi F, Esmailzadeh A. Determinants of stunting in school-aged children of tehran, iran. *Int J Prev Med*. 2013;4(2):173-9.
23. Dong Y, Bennett K, Jan C, Dong B, Zou Z, Hu P, et al. Subnational variation of stunting, wasting and malnutrition in Chinese primary-school children between 2010 and 2014: urban-rural disparity. *Public Health Nutr*. 2019;22(11):2043-54.
24. McDonald CM, Baylin A, Arsenault JE, Mora-Plazas M, Villamor E. Overweight is more prevalent than stunting and is associated with socioeconomic status, maternal obesity, and a snacking dietary pattern in school children from Bogota, Colombia. *J Nutr*. 2009;139(2):370-6.

25. Mohamed S, Hussein MD. Prevalence of Thinness, Stunting and Anemia Among Rural School-aged Sudanese Children: A Cross-sectional Study. *J Trop Pediatr.* 2015;61(4):260-5.
26. Mushtaq MU, Gull S, Khurshid U, Shahid U, Shad MA, Siddiqui AM. Prevalence and socio-demographic correlates of stunting and thinness among Pakistani primary school children. *BMC Public Health.* 2011;11:790.
27. Qin Y, Melse-Boonstra A, Zhao J, Wu M, Hu X, Kok FJ. Stunting and zinc deficiency among primary school children in rural areas with low soil zinc concentrations in Jiangsu Province, China. *Asia Pac J Clin Nutr.* 2009;18(1):15-21.
28. Wu H, Li H, Zong X. The prevalence of overweight, obesity and stunting in school children aged 6-19 years in Beijing, China. *Ann Hum Biol.* 2016;43(6):505-9.
29. Altare C, Delbiso TD, Mutwiri GM, Kopplow R, Guha-Sapir D. Factors Associated with Stunting among Pre-school Children in Southern Highlands of Tanzania. *J Trop Pediatr.* 2016;62(5):390-408.
30. Ethiopian demographic and health survey. Ethiopian Statistical agency. 2016.