Cortical neuronal activity determines the dynamics of local sleep homeostasis

Christopher W. Thomas[1], Mathilde C. C. Guillaumin[2], Laura E. McKillop[1],

Peter Achermann[3], Vladyslav V. Vyazovskiy[1]*

[1] Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, UK; [2] Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK; [3] The KEY Institute for Brain-Mind Research, Department of Psychiatry, Psychotherapy and Psychosomatics, University Hospital of Psychiatry, Zurich, Switzerland

* Corresponding author. Email: vladyslav.vyazovskiy@dpag.ox.ac.uk

Funding acknowledgements: BBSRC (BB/K011847/1, BB/M011224/1), MRC (MR/L003635/1), Wellcome Trust (098461/Z/12/Z), John Fell OUP Research Fund (131/032).

Abstract

The homeostatic regulation of sleep manifests as a relative constancy of its total daily amount, and the compensation of sleep loss by an increase in its subsequent duration and intensity. Theoretical descriptions of this phenomenon define "Process S", a variable with dynamics dependent only on sleep-wake history and whose levels are reflected in EEG slow wave activity. While numerous hypotheses have been advanced regarding the substrate and role of Process S, such as synaptic or energy homeostasis, it remains unclear whether these dynamics are fundamentally driven by a need to homeostatically regulate specific variables, or by an unknown innate process which enforces that a certain daily sleep quota is obtained. Sleep is typically defined based on brain-derived criteria, such as behaviour or EEG power spectra, and variation in brain activity during wakefulness has been linked to variation in Process S accumulation. We therefore hypothesised that Process S dynamics might be related to the quantity and characteristics of spiking activity in cortical neurones. Specifically, we assumed that Process S changes as a function of the deviation of neuronal firing rate from a locally defined set point. To relate these dynamics explicitly to patterns of spiking activity, we incorporated the occurrence of network spiking off periods as both the defining measure of Process S and as the determinant of its rate of decay. This approach was able to describe the time course of Process S, crucially without explicit knowledge of the animal's global sleep-wake state. This work provides a conceptual advance in our understanding of the substrate of sleep homeostasis and provides important links between local and global aspects of sleep regulation.

Introduction

According to traditional theory, the need for sleep accumulates during wakefulness and dissipates during sleep. Despite decades of research, it is still uncertain precisely which biological variables form the substrate of sleep need, what characteristics of wake challenge their stability and how sleep mediates the restoration of homeostasis. More fundamentally, it remains unclear whether homeostatic sleep regulation reflects an active process, dynamically shaping daily sleep architecture in response to a physiological need for the homeostatic regulation of specific variables, or whether it corresponds instead to an unknown innate process which ensures only that a certain daily quota of sleep is obtained.

The earliest theories of sleep homeostasis supposed the existence of a single variable, termed Process S, which describes sleep drive at the global level (Borbély, 1982). This variable is assumed to always increase during wakefulness, independently of its content, and to decline during sleep. It is widely acknowledged that homeostatic sleep pressure is reflected in the levels of slow wave activity (SWA, 0.5-4 Hz spectral power) observable during NREM sleep in neurophysiological field potentials, such as electroencephalogram (EEG) or local field potential (LFP).

Current views on the origin of sleep homeostasis emphasise its local and activity-dependent component (Krueger & Tononi, 2011; Rattenborg et al., 2012; Tononi & Cirelli, 2014). It was shown that SWA is far from uniform across the brain, and that the behavioural and cognitive content of waking, beyond merely its duration, influences subsequent SWA magnitude (Huber et al., 2004; Vyazovskiy & Tobler, 2008; Rector et al., 2009; Murphy et al. 2011; Fisher et al., 2016). Indeed, many candidate mechanisms for the substrate of sleep homeostasis implicate processes occurring at a cellular and network level. These include the maintenance of cellular homeostasis (Reimund 1994; Mackiewicz et al., 2007; Vyazovskiy & Harris, 2013; Bellesi et al., 2016), the replenishment of energy stores (Scharf et al., 2008), the influence of sleep-related signalling molecules such as adenosine or cytokines (Krueger et al., 2008), and the regulation of imbalanced synaptic strength (Tononi & Cirelli 2003; Vyazovskiy et al., 2008; Liu et al., 2010). Although the equivalence of these processes with Process S has not been conclusively demonstrated (Frank & Heller, 2019), the existing evidence supports the relevance of sleep-wake dependent differences in neuronal activity for understanding the regulation of sleep.

Cortical neuronal activity is generally higher during waking compared to sleep (Vyazovskiy et al., 2009; Hengen et al., 2016; McKillop et al., 2018). Lower spike firing typical of sleep is due, at least in part, to regular periods of widespread synchronous network silence, termed off periods, intruding on ongoing activity (Steriade et al., 1993a; Steriade et al., 2001; Sanchez-Vives & Mattia, 2014). Importantly, off periods in neural populations are thought to underpin slow wave dynamics at the level of the field potential (Steriade et al., 1993b; Massimini et al., 2004; Buzsáki et al., 2012) and their properties reflect homeostatic sleep need (Vyazovskiy et al., 2009; McKillop et al., 2018; Saberi-Moghadam et al., 2018).

Neuronal firing rates typically fluctuate around a homeostatic set point, which is characteristic for individual cells and variable across the population (Turrigiano, 2011; Hengen et al., 2013; O'Leary et al., 2014; Styr et al., 2019). The homeostatic regulation of firing rates may depend on processes occurring specifically in sleep and wakefulness (Grosmark et al., 2012; Hengen et al, 2016) and evidence suggests that firing rates change as a function of time spent awake, conditional on wake's behavioural quality (Vyazovskiy et al., 2009; Fisher et al., 2016). Additionally, the magnitude and direction of state-dependent changes in firing rate differs between neurons, depending on brain region and their individual firing rate set point (Miyawaki & Diba 2016; Watson et al., 2016; Miyawaki et al., 2019). Firing rate homeostasis may occur above the individual cell level (Slomowitz et al., 2015), as network level mechanisms dependent on the NREM sleep slow oscillation constrain the distribution of population firing rates within an optimal functional range (Levenstein et al., 2017).

Overall, evidence suggests that the homeostatic regulation of sleep and of neuronal firing rates may be intrinsically related, however, this functional link remains incompletely defined. Mathematical modelling approaches present an opportunity to begin addressing this problem. To this end, we develop quantitative models of Process S, describing its temporal dynamics on a local level in terms of spiking activity and off periods, instead of global sleep-wake history. We find that the magnitude of the deviation of multi-unit firing rate from a locally specified set point carries sufficient information to account for empirically derived patterns of SWA. We then introduce the total time spent in off periods as an alternative measure for Process S with more local origins than SWA, and account for the time course of this variable in terms of two opponent processes dependent on spiking rate and off period occurrence.

Materials & Methods

Animals, surgery & husbandry

Chronic electrophysiological recordings from six male young adult (4.8 – 5.7 months old, mean 5.2 months) C57BL/6J mice were analysed here. This data set is a subset of that used in a previous study (McKillop et al., 2018). The animals were surgically implanted with electrodes for the continuous recording of electroencephalography, electromyography and cortical neuronal activity. EEG screw electrodes (Fine Science Tools) were inserted into the skull unilaterally above the frontal cortex (primary motor area: anteroposterior 2mm, mediolateral 2mm) and bilaterally above the occipital (primary visual area: anteroposterior 3.5mm, mediolateral 2.5mm) cortex. One occipital screw (contralateral to the frontal screw) served as the ground electrode and an additional screw located above the cerebellum served as the reference. A pair of stainless steel wires were inserted into nuchal muscle for the recording of electromyogram (EMG). A polyimide-insulated tungsten microwire array (Tucker-Davis Technologies) was implanted through a craniotomy window into the frontal cortex (primary motor area: anteroposterior 2mm, mediolateral -2mm), contralateral to the EEG screw. The array comprised 16 wire channels of 33µm diameter, arranged in 2 rows of 8, with columnar separation of 250µm, row separation of 375µm and tip angle of 45 degrees. One row of wires was 250µm longer than the other to account for cortical curvature. A silicone gel (KwikSil, World Precision Instruments) was used to seal the craniotomy, and dental acrylic cement used to stabilise all the implanted electrodes. Surgeries were performed under isoflurane anaesthesia (4% induction, 1-2% maintenance). Analgesics were given immediately before surgery (1-2 mg/kg metacam and 0.08 mg/kg vetergesic, subcutaneous injection) and for at least 3 days during recovery following surgery (1-2 mg/kg metacam, oral). In addition, immunosuppressant (0.2 mg/kg dexamethasone) was given the day before surgery (oral), immediately before surgery (subcutaneous injection) and during recovery for at least 2 days (oral). Animal wellbeing was closely monitored during recovery until a stable return to baseline was observed for at least 2 days. All procedures were performed under a UK Home Office Project License and conformed to the Animal (Scientific Procedures) Act 1986.

Mice were housed individually following surgery. Two weeks after surgery, mice were transferred, still

individually, to custom made Plexiglas cages (20.3x32x35 cm), containing a running wheel (Campden

Instruments), which were placed within ventilated sound-attenuated Faraday chambers (Campden

Instruments). The animals were exposed to a standard 12hr-12hr light dark cycle, with food and water

available *ad libitum*. Mice were allowed to habituate to the recording chamber and to attachment of the

recording cables for a minimum of three days before recordings began.

Experimental design

All the data analysed here were collected over two days of experimental recording. The first day served

as a baseline, while the animals were completely undisturbed. A sleep deprivation protocol was

enforced at light onset on the second day, immediately after the baseline day, and lasted 6 hours. Sleep

deprivation was performed using novel object presentation. During this period, experimenters

constantly monitored both the behaviour and ongoing neurophysiological recordings of the mice. As

soon as any animal showed signs of sleepiness (such as stillness with eyes closed, or slow waves in the

EEG), novel objects were introduced to the cage (such as cardboard, colourful plastic and tissue paper)

in order to encourage wakefulness. During the 6-hr sleep deprivation period, these mice slept 6.0 ± 3.1

minutes (mean ± sd) only.

Data collection & pre-processing

Data acquisition was performed using a Multichannel Neurophysiology Recording System (Tucker Davis

Technologies). EEG, EMG and microwire array LFP signals were filtered (0.1-100 Hz), amplified (PZ5

NeuroDigitizer preamplifier, TDT) and stored locally (256.9 Hz sampling rate). Custom written Matlab

scripts were used for signal conversion and data pre-processing. The LFP, EMG and EEG signals were

filtered again offline between 0.5 - 100 Hz (4th order Type II Chebyshev filter) and resampled at 256 Hz.

Extracellular multi-unit spiking was additionally obtained from each microwire array channel, recorded

at 25 kHz and filtered 300 Hz – 5 kHz. An amplitude threshold (at least 2 standard deviations, minimum

-25µV) was used to identify putative spikes. Individual spikes were saved as a voltage waveform

comprising 46 data samples (0.48ms before to 1.36ms after threshold crossing) plus a time stamp of

occurrence. Spiking activity from each channel was cleaned offline for artefacts using the Matlab spike sorting software Wave_clus (Quiroga et al., 2004). All putative single unit clusters identified by the algorithm from the same channel were merged, excluding only noise spikes. MUA firing rate for each channel was calculated in 4-second epochs as the mean number of spikes per second.

Vigilance states were scored manually by visual inspection with a 4-second epoch resolution (using the software SleepSign, Kissei Comtec). Vigilance states were classified as waking (low amplitude but high frequency EEG with high or phasic EMG activity), NREM sleep (presence of EEG slow waves, a signal of a high amplitude and low frequency, and a low level of EMG activity) or REM sleep (low amplitude, high frequency EEG, and low EMG). For this analysis, over the six mice, a total of 78 out of 96 channels could be analysed (minimum of 10 out of 16 from one mouse). The excluded channels were characterised by an unstable MUA firing rate, including a large drift in firing which was persistent and sleep-wake state independent. For the off occupancy models described below, 75 out of 96 channels were used; a further 3 channels had to be excluded because multi-unit firing rates, while stable, were too low to yield a reasonable estimate of the occurrence of off periods.

Slow wave activity

Each EEG and LFP signal was processed to extract a measure of the SWA. Signal segments were extracted within windows of 4-second duration and 1-second spacing (giving 3-second overlap), and Hann tapered. A Fourier Transform was applied to each signal segment and the mean power in the slow wave range (frequencies from 0.5 to 4 Hz) was calculated. This measure was smoothed by finding the median over five temporally adjacent overlapping segments, yielding a SWA measurement for each sequential 4 second epoch. For normalisation within each channel, SWA values were then expressed as a percentage of the mean SWA calculated over all artefact-free epochs scored as NREM sleep in the baseline 24 hours. The median value of the SWA during each continuous NREM sleep episode served as an estimate of Process S. NREM sleep episodes shorter than 1 minute (15 epochs) were excluded, as in a previous study modelling Process S in mice (Guillaumin et al., 2018). Brief awakenings (short arousals accompanied by movement, lasting ≤20 seconds) were excluded from analysis but were not considered to be the ending of a NREM sleep episode.

Off periods detection and definition of "off occupancy"

One aim of this study was to develop measures of Process S independent of the EEG, and so we focused on off periods which are the neuronal counterpart of slow waves. Off periods refer to brief interruptions of spiking activity which occur synchronously across many recording sites, last approximately 70-100ms (Vyazovskiy et al., 2009), and are coincident with the positive deflection of LFP slow waves (McKillop et al., 2018). In these studies, off periods were detected by pooling spikes over all channels and identifying inter-spike intervals (ISIs) that exceed some long duration threshold. However, pooling spikes removes the ability to compare local differences, and if an off period does not involve all channels it will go undetected. To overcome this limitation, off periods were defined separately for each channel by looking at the co-occurrence of local slow waves and spiking silence.

There is no universally accepted method for slow wave detection, and a recent comparison suggests that a simple amplitude threshold based approach, while in theory adequate, may underperform due to channel differences in overall LFP amplitude and large amplitude fluctuations of higher frequencies (Bukhtiyarova et al., 2019). For this reason, the LFP was first filtered from 0.5 to 6 Hz (4th order Butterworth filter), and a threshold defined individually for each channel, using the median plus one median absolute deviation of the peak amplitude of all positive half waves (including all vigilances states, but excluding epochs with artefacts). All positive half waves with peak amplitude above threshold were then considered to be slow waves.

Next, for each LFP slow wave, the multi-unit spike preceding and following the slow wave peak was identified and the corresponding ISI determined. The distribution of these ISIs which coincide with slow waves was often (in 64 out of 75 channels) unambiguously bimodal, allowing the threshold to be selected at the local minimum between these two modes. When there was no evidence of bimodality a value of 120ms was chosen, corresponding to the maximum value selected for the other channels. All inter-spike intervals aligned to slow wave peaks with duration exceeding the threshold were considered off periods. Finally, the metric termed off occupancy was defined, for each 4-second epoch, as the percentage of time spent in a detected off period during that epoch. Just as with SWA (see above), the median value of off occupancy over NREM sleep episodes was used to represent the level of Process S.

Model fitting and parameter optimisation

Three theoretical models were used to describe the time course of Process S (see Results). The model equations were solved using a discrete time approximation, iteratively updating the value of modelled Process S in time steps of 4 seconds (Euler method). The fitting of various models to a particular channel of data is equivalent to finding the optimal choice of parameters to achieve the closest match between simulation and empirical SWA/off occupancy. In all models, the initial value of Process S was included as an additional free parameter. The selection of parameter values for model fitting was achieved using a semi-automated approach. First, an algorithmic methodology was established, which depends primarily on the definition of an error metric to assess fit quality between modelled Process S and empirical data. For each animal, NREM sleep episodes were identified with a duration of at least 1 minute. For each NREM sleep episode ($n=1:N$), the median empirical SWA/off occupancy (X_n), and similarly the median modelled value of Process S (S_n), were computed. The error metric (E) is defined as the sum of absolute differences between model and data, weighted by the relative episode duration (w_n). This weight was defined as the absolute duration (d_n) of the NREM sleep episode in seconds, normalised by the total duration of all episodes.

$$E = \sum_{n=1:N} w_n.|X_n - S_n|$$

$$w_n = d_n / \sum_{i=1:N} d_i$$

Algorithmic parameter optimisation was performed separately for each channel, aiming to minimise this error metric. This was achieved using the Matlab function *fminsearch*, which uses the Nelder-Mead simplex algorithm (Lagarias et al., 1998). Many parameter combinations produce very similar dynamics. Therefore, there exist many possible optimal (or near-optimal) parameter regimes and so algorithmic optimisation is sensitive to initial values. For this reason, initial values for the parameters were first set manually, aided with the use of a custom-made Matlab graphical user interface. Manually selected values were then fed into the algorithmic optimisation. Final parameter values were visually inspected to ensure that this optimisation produced an improvement of fit.

The final fit quality of a model to the data was expressed as the median percent error (E^*). This is

calculated by finding the absolute difference between median empirical (X_n) and simulated (S_n) SWA/off

occupancy for each NREM sleep episode, expressed as a percentage of the empirical SWA/off occupancy

in that episode. E^* is then defined as the median over all NREM sleep episodes of these difference values.

This alternative error metric is used for presentation of results because it has more comprehensible

units (percentage of empirical SWA/off occupancy) compared to the error metric, with arbitrary units.

221
$$E^* = \operatorname*{median}_{n=1:N} \left[\frac{|X_n - S_n|}{X_n} \cdot 100\% \right]$$

Statistical analyses

The correlation between wake duration, wake firing rates and changes in SWA was calculated separately

for each LFP channel. We first identified episodes of NREM sleep of at least 1-minute duration (exactly

as for modelling) and obtained the median SWA in each episode. We then identified which intervening

time periods, between two NREM episodes, comprised at least 80% wake and lasted at least 5 minutes.

We then calculated Pearson correlation coefficients between i) the duration of these wake periods, ii)

the mean multi-unit firing rate during these periods, and iii) the change in median SWA from preceding

to the following NREM sleep episode.

Analysis of variance was performed to explore factors influencing model parameters and fit quality

using the Matlab functions *anova1* (one-way) and *anovan* (two-way with unequal group size). For the

rate parameters, ANOVA was calculated after applying a log transform. The effect size (η^2) is calculated

for a factor as its sum of squares divided by the total sum of squares in the ANOVA and reflects the

fraction of the variance accounted for by that factor.

To summarise the results in figures, boxplots were included alongside individual data points. These

indicate the median, 25th and 75th percentile, with whiskers extending to the most extreme value which

falls within 150% of the interquartile range of the box. For these plots, results from individual channels

were typically pooled across animals. In some cases, where indicated, channels from the same animal

were presented as separate populations.

 Results

We selected a dataset of chronic recordings of frontal electroencephalogram (EEG) alongside local field

potential (LFP) and multi-unit activity (MUA) spiking from primary motor cortex from six mice. These

recordings were made continuously over 48 hours, starting at light onset, while the mice were freely

behaving within their cage and exposed to a standard 12hr-12hr light-dark cycle. At light onset of the

second day a sleep deprivation protocol was enforced for 6 hours, involving the presentation of novel

objects. Outside of this sleep deprivation period, mice were undisturbed and able to sleep at will.

Process S dynamics can be described as a function of vigilance state history

NREM sleep can be defined and distinguished from waking by the presence of high amplitude slow (0.5

- 4Hz) waves in the EEG. The average EEG spectral power in the slow wave range (termed slow wave

activity; SWA) in NREM sleep varies as a function of the animal's recent sleep-wake history, and this

relationship has been captured in a classical quantitative theory using the concept of "Process S"

(Borbély, 1982; Daan et al., 1984). Process S describes a variable whose magnitude can be estimated

using the level of SWA during NREM sleep, reflecting the intensity of sleep, and which is interpreted as

corresponding to the homeostatic component of sleep drive (Franken et al., 1991a; Achermann et al.,

1993; Huber et al., 2000a; Vyazovskiy et al., 2007; Guillaumin et al., 2018). Theoretically, this Process S

follows simple dynamics; during wake it increases according to a saturating exponential function

towards an upper asymptote, and similarly during NREM sleep it decays exponentially towards a lower

asymptote. There are many published variants of the precise equations for this model, but crucially, all

these variants use the sleep-wake state history of the animal as the key predictive variable for Process

S. Here, the specific equations used are:

$$\text{Wake/REM Sleep:} \quad dS/dt = \alpha(S_{max} - S(t))$$

$$\text{NREM Sleep:} \quad dS/dt = -\beta(S(t) - S_{min})$$

Where S represents the level of Process S, S_{max} and S_{min} are upper and lower asymptotes, and α and β

are rate parameters. The first equation is applied when the animal is scored to be awake or in REM sleep

and the second equation is applied when it is scored to be in NREM sleep. Here, we use the typical

approach of applying the wake dynamics equation to REM sleep (Franken et al., 1991a; Huber et al.,

2000a; Vyazovskiy et al.; 2007), but for simplicity do not set separate parameters for wake vs. REM

sleep. For convenience, S is expressed in units equivalent to those of SWA. This classical formulism is

later abbreviated as model Cl-SWA. Process S as described by these simple dynamics accounts for the

time course of empirical SWA with high accuracy. We applied this model to SWA derived from the frontal

EEG of all animals, and as expected obtained a high quality fit (Figure 1A).

Neuronal firing rates are associated with Process S dynamics and vigilance state distributions

Whichever specific processes within the brain underpin these state-dependent dynamics for Process S,

it is likely associated in some way to neuronal spiking activity. Spiking patterns differ characteristically

between wake and NREM sleep, due to the presence of off periods in NREM sleep which typically

coincide with slow waves in both EEG and LFP (Figure 1B). The occurrence of such spiking off periods

cause firing rates to be typically much lower during NREM sleep compared to wake and REM sleep. In

this dataset, the mean multi-unit firing rate averaged over all periods of wake was larger than the firing

rate averaged over all periods of NREM sleep in every recording channel and every animal (Figure 1C).

Because the slow oscillation is underpinned by local neuronal dynamics, it is expected to be highly

heterogeneous across the neocortex. Regional differences in SWA dynamics have been previously

described, for example between frontal and occipital EEG derivations (Werth et al., 1996; Huber et al.,

2000b), and can be accounted for within the classical Process S model through the selection of locally

variable values for model rate parameters (Zavada et al., 2009; Rusterholz & Achermann, 2011;

Guillaumin et al., 2018). To explore whether spike firing rates might account for some of the variation

in the rate of Process S increase, we correlated the change in LFP SWA from one NREM sleep episode to

the next, when separated by wakefulness lasting at least 5 minutes, with the mean spike firing rate

during this intervening wake period, and with the duration of that wake period. Correlation coefficients

were obtained separately for each recording channel, and pooled across animals. As expected, we

obtained large positive correlation coefficients between the duration of a period of wakefulness and the

change in LFP SWA in all channels (mean=0.66, sd=0.13; Figure 1D). Importantly however, the change

in LFP SWA was on average also positively correlated with firing rate (mean=0.27, sd=0.30; Figure

1D,E), meaning that, generally, a higher firing rate during waking was associated with a larger increase

in SWA in subsequent NREM sleep. Interestingly, these phenomena were not independent, as a positive

correlation was also found between wake episode duration and firing rate (mean=0.35, sd= 0.33; Figure

1D). These results further support the possibility that neuronal activity is associated with Process S

dynamics and vigilance state distributions. To test this hypothesis we next turned to a quantitative

modelling approach.

Process S dynamics can be described as a function of neuronal firing rates

We first sought to determine whether a model expressing Process S dynamics solely as a function of

local multi-unit neuronal firing rates might describe the levels of sleep SWA in the corresponding LFP

with comparable accuracy to the classical model dependent on vigilance states at the global level. The

classical Process S model was used as a starting point for the development of a novel firing rate

dependent alternative. To do this, the equations of the classical model were adapted in two ways. Firstly,

an instantaneous firing rate threshold (F_θ) was introduced as a new model parameter to replace the

wake vs. sleep criterion, assuming an increase in Process S when the threshold is exceeded and decrease

when firing is below. Conceptually, this firing rate threshold resembles a set point; a target firing rate at

which the dynamics are stable. Secondly, we assumed that the rate of change of Process S is proportional

to the difference between firing rate and this threshold. Introducing this change to the equations ensures

that the rate of change of S is equal to zero exactly at the set point, and a continuous function of firing

rate around this value. This version of the model is later abbreviated as Fr-SWA. The equations are:

$$F(t) > F_\theta: \quad dS/_{dt} = \ \alpha(S_{max} - S(t))(F(t) - \ F_\theta)$$

$$F(t) < F_\theta: \quad dS/_{dt} = \ -\beta(S(t) - S_{min})(F_\theta - F(t))$$

$$F(t) = F_\theta: \quad dS/_{dt} = \ 0$$

310   We applied this novel model, alongside the classical model, to describe SWA dynamics at the LFP level,

311   by finding parameter values that would minimise the difference between empirical SWA and modelled

312   Process S. The two models were fit to the SWA from each LFP channel separately, using the multi-unit

313   firing rate from the same channel for the firing-rate-dependent model. The models were also fit to the

SWA obtained by averaging the LFP (and firing rate) over the whole population of channels within the same mouse.

Figure 2 shows two examples of the fit of both models, first to SWA derived from the average LFP, and also to SWA derived from a single LFP channel (these examples are from the same animal as in Figure 1A). The overall pattern of Process S dynamics is similar at both recording levels, and to the corresponding EEG, and is well described by both the classic and novel firing-rate-based model. Throughout the dataset, this purely firing-rate-dependent model described the overall dynamics of LFP SWA during NREM sleep to a comparable accuracy as the classical model, as reflected in the median percent error deviation between modelled and empirical SWA (Figure 3B). Both the model type and animal have a highly significant effect on the median percent error of the model fit to individual LFP channels, although there is no significant interaction (Model: $F_{(1, 144)}$ = 25.9, p = $1.1x10^{-6}$; Animal: $F_{(5, 144)}$ = 30.2, p = $6.3x10^{-21}$; Model x Animal: $F_{(5, 144)}$ = 1.65, p = 0.15; two-way ANOVA unequal groups). Errors were higher for the novel model, importantly however, the differences in fit quality due to the model type are small relative to the effect size of the particular animal and channel, on which fit quality depends much more strongly (Model: η^2 = 0.079; Animal: η^2 = 0.459; Model x Animal: η^2 = 0.025; Channel (residuals): η^2 = 0.437). The errors of the model fit on the averaged LFP and on the EEG (classical model only) are also shown in Figure 3B, and are more explicitly compared in Figure 3C. We do not find any significant effect of the field potential level (EEG, average LFP, single LFP) on the model error ($F_{(2,171)}$ = 0.2, p = 0.82, one-way ANOVA).

The distributions of final optimised parameters are shown in Figure 3D-G for the classic model and Figure 3H-L for the firing rate based model. Most parameters in both models were significantly different between animals, with the exception of S_{min} in the firing rate dependent model (Cl-SWA α: $F_{(5,77)}$ = 4.03, p = $2.8x10^{-3}$; Cl-SWA β: $F_{(5,77)}$ = 8.65, p = $1.9x10^{-6}$; Cl-SWA S_{max}: $F_{(5,77)}$ = 14.35, p = $9.8x10^{-10}$; Cl-SWA S_{min}: $F_{(5,77)}$ = 12.42, p = $1.1x10^{-8}$; Fr-SWA α: $F_{(5,77)}$ = 8.33, p = $3.0x10^{-6}$; Fr-SWA β: $F_{(5,77)}$ = 9.27, p = $7.5x10^{-7}$; Fr-SWA F_θ: $F_{(5,77)}$ = 14.3, p = $1.0x10^{-9}$; Fr-SWA S_{max}: $F_{(5,77)}$ = 6.64, p = $3.9x10^{-5}$; Fr-SWA S_{min}: $F_{(5,77)}$ = 2.16, p = 0.07; one way ANOVA). The optimised rate parameters for EEG and averaged LFP data consistently fall within the range of the single channel population (Figure 3D-L). The relationship between the optimal

firing set point and state specific firing rates is shown in Figure 3M. Firing rate threshold was z-

normalised (subtract mean, divide by standard deviation) separately with respect to the distribution of

firing rates in wake, NREM and REM sleep. This shows, as expected, that the set point is typically below

mean firing in wake (-1.0 ± 0.6; mean ± sd), well above mean firing in NREM sleep (1.6 ± 1.1), and slightly

above mean firing of REM sleep (0.7 ± 1.1). These were all significantly different from zero (wake: p =

$1.6x10^{-13}$, NREM: p = $1.9x10^{-14}$, REM: p = $3.4x10^{-7}$, two-sided Wilcoxon signed rank test). Figure 3N

further shows the state dependent distribution of average firing rate, expressed as a percentage of the

chosen firing set point parameter. Mean firing was 182 ± 57.6% of the firing rate set point during wake,

61.8 ± 17.6 % during NREM sleep, and 84.2 ± 31.8 % during REM sleep. Again, these distributions were

all significantly different from 100% (Wake: p = $6.7x10^{-14}$, NREM: p = $2.0x10^{-14}$, REM: p = $5.4x10^{-5}$). Note

that, in a few channels, the optimal firing rate threshold is actually above the mean firing rate during

waking. This occurs when the waking firing rate distribution overlaps substantially with the NREM sleep

firing rate distribution but has a heavier tail. REM sleep, in this dataset, is typically associated with firing

below the set point, and therefore Process S decrease, albeit at a reduced rate compared to NREM sleep

(Figure 3O). Interestingly, there are no significant correlations between the model fit error and the

threshold normalisation relative to any one vigilance state (Wake: p=0.06, r=0.21; NREM: p=0.35, r=-

0.11; REM: p=0.94, r=-0.01), suggesting that there is no clear relationship between the firing rate set

point and firing rate distribution of any one particular state.

Process S can be defined in terms of neuronal spiking off periods

When considering the relationship between multi-unit firing rates and LFP slow waves, a conceptual

complication arises due to the different origins of the LFP and MUA from within the same channel. While

the MUA firing rate represents the activity of only a few individual neurones, factors such as volume

conduction result in spatial smoothing of the LFP signal, and as such it represents the combined activity

of neurones covering a cortical area potentially on the order of several millimetres (Kajikawa &

Schroeder, 2011). This means that when a slow wave is detected in the LFP, it is not guaranteed that

local neurones which contribute to the MUA are necessarily in an off period. Similarly, not every long

interspike interval occurs during a slow wave. An estimation of the occurrence of off periods may be

obtained by combining LFP and spiking data (Figure 4A). Slow wave detection was performed on each LFP over the whole 48hrs, including all vigilance states (0.5-6 Hz filter followed by an amplitude threshold, values shown in Figure 4C). All multi-unit inter-spike intervals (ISIs) which coincide with the peak of detected slow waves were identified. The distribution of the duration of these ISIs was often (64 out of 75 channels) unambiguously bimodal (Figure 4B). This could be interpreted as evidence of the existence of two spiking states occurring locally during the more widespread network slow oscillation; high frequency spiking (on period) and extended silence (off period). Note that these two distributions do not simply correspond exactly to sleep vs. wake conditions, because bimodality is often evident in the distribution of slow wave coincident ISIs from NREM sleep only, or even from REM sleep only (Figure 4B). The distribution of slow wave coincident ISIs (over all vigilance states) was used to define an ISI duration threshold for the detection of off periods, separately for each channel. The values used for this threshold are shown in Figure 4D. The average multi-unit firing rate aligned to the peak of slow waves in detected off periods reveals a clear suppression of firing, consistent with expectations (Figure 4E). A rebound increase in average firing is visible in this example channel immediately after the off period, as has been previously documented in some cortical neuronal populations (Chauvette et al., 2010). The total fraction of time each channel spends in off periods was calculated over all 4-second epochs, and termed the "off occupancy". Off occupancy defined in this way is high in NREM sleep and low in both wake and REM sleep (Figure 4F). The existence of a non-zero frequency of local cortical off periods has been previously reported during wakefulness (Vyazovskiy et al., 2011; Vyazovskiy et al., 2014; Fisher et al., 2016), and REM sleep (Funk et al., 2016). The off occupancy measure displays similar temporal dynamics to LFP (and EEG) SWA over both sleep deprivation and spontaneous sleep and wake (Figure 4G).

Process S dynamics, defined using off periods, can be described as a function of vigilance states or neuronal firing rates

In order to investigate whether the off occupancy measure reflects Process S, we applied the classical state based model to the time course of off occupancy, exactly as was done with single channel SWA. The classical model was applied with its equations unchanged, and is abbreviated as Cl-Off. Figure 5A shows

an example of the resulting Process S time course obtained in this way with a high quality of fit. In contrast, some changes were introduced to the firing-rate-based model in order to describe off occupancy dynamics. We considered that the different dynamics above vs. below a particular firing rate set point may be due to two opponent processes simultaneously active in dynamic opposition but with differential magnitude in wake vs. NREM sleep. Specifically, we consider a Process S increasing term which is proportional to instantaneous firing rate, and a Process S decreasing term which is proportional to the time spent in off periods (off occupancy). The equation used is:

$$dS/dt = \alpha F(t)(S_{max} - S(t)) - \beta X(t)(S(t) - S_{min})$$

In this model, one term drives S towards an upper asymptote (S_{max}) in proportion to firing rate (F), while the other drives S towards a lower asymptote (S_{min}) in proportion to the off occupancy (X). Again, two rate parameters α & β are required. This model behaves similarly to previous models because firing is high in wake and low in NREM sleep, whereas off occupancy is high in NREM sleep and low in wake. This variant of the model is abbreviated as Fr-Off. Figure 5A includes also the fit from this model, demonstrating a high level of agreement between the two models and an accurate fit to the data.

The distribution over all animals and channels of the median percent errors for the fits from both models is shown in Figure 5C. As before, the model type and animal has a significant effect (Model: $F_{(1,138)} = 8.06$, p = 5.2x10^{-3}; Animal: $F_{(5,138)} = 17.2$, p = 3.2x10^{-13}; Model x Animal: $F_{(5,138)} = 0.52$, p = 0.76; two-way ANOVA with unequal groups) and the classic model achieved a slightly lower median percent error. However, this effect was again very weak compared with the variation in fit quality between animals and channels (Model: $\eta^2 = 0.034$; Animal: $\eta^2 = 0.367$; Model x Animal: $\eta^2 = 0.011$; Channel (residuals): $\eta^2 = 0.588$). Figure 5D shows the distribution of values for the change in modelled Process S from one time step to the next resulting from the firing-rate-based model (Fr-Off), in wake, NREM sleep and REM, pooling all animals, channels and time. Unlike in the previous firing-rate-based model, REM sleep is now typically associated with Process S increase. The distributions of final optimised parameters are shown in Figure 5E-H for the classic model and Figure 5I-L for the firing rate and off occupancy model. Most parameters in both models were different between animals to a high significance level (Cl-Off α: $F_{(5,74)} = 2.4$, p = 0.047; Cl-Off β: $F_{(5,74)} = 2.29$, p = 0.055; Cl-Off S_{max}: $F_{(5,74)} = 9.49$, p = 6.4x10^{-7}; Cl-Off S_{min}: $F_{(5,74)} = 16.79$, p

422 = 8.0×10^{-11}; Fr-Off $\boldsymbol{\alpha}$: $F_{(5,74)}$ = 9.86, p = 3.9×10^{-7}; Fr-Off $\boldsymbol{\beta}$: $F_{(5,74)}$ = 5.14, p = 4.6×10^{-4}; Fr-Off $\boldsymbol{S_{max}}$: $F_{(5,74)}$ =

423 10.08, p = 2.8×10^{-7}; Fr-Off $\boldsymbol{S_{min}}$: $F_{(5,74)}$ = 10.25, p = 2.3×10^{-7}; one way ANOVA). Notably, the weakest

424 evidence for inter-animal differences were for $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ in the classic model.

Modelling identifies local variation in Process S dynamics

All four modelling approaches described here suggest the existence of variability in Process S between

recording channels, indicating that a local component determines its dynamics. Figure 6A-D illustrates

this diversity, showing the overlaid time courses of Process S for all channels within a single

representative animal, expressed as a percentage of their individual maximum value for normalisation

purposes. Although the models all fit empirical SWA or off period occupancy with generally high

accuracy, there are nonetheless differences between models in the precise shape of Process S. The

average Process S over all channels was also calculated for each model and compared with the Process

S derived from applying the classical model to EEG SWA (Figure 6E). This reveals that Process S derived

from modelling LFP SWA more closely resembles Process S derived from EEG SWA than Process S

derived from off occupancy. This suggests that global Process S calculated at a higher spatial scale might

reflect an averaging across a manifold of local Processes S that exist on a finer spatial level.

Discussion

Here, we show that Process S can be quantitatively described entirely in terms of measures based on

neuronal spiking activity, without reference to global vigilance states. A model was outlined whereby

the integrated history of multi-unit firing rates, relative to a firing rate set point, predicted the temporal

dynamics of LFP SWA. The accuracy of this model was demonstrated in a dataset of recordings from

mouse frontal cortex over 48 hours, including both voluntary sleep and wake, and sleep deprivation. A

novel metric for Process S was then presented, termed off occupancy, which measures the fraction of

time a neural population spends in off periods, defined by the coincidence of LFP slow waves and multi-

unit spiking silence. The modelling approach was combined with the off occupancy metric and tested on

the same dataset to present a quantitative framework for understanding dynamics of sleep pressure at

a highly local level in terms of neural spiking and off periods.

Central to this modelling perspective is the assumption that the generation of spikes by a neural network

is in some way correlated with an increase in sleep pressure, and that this manifests in the subsequent

expression of slow waves and off periods, reflecting Process S. The energetic cost of spiking is high

(Attwell & Laughlin 2001), the regulation of a neurone's firing rate set point is linked to cellular

energetics (Styr et al., 2019; Vergara et al., 2019), and neurones are susceptible to cellular stresses that

can result from sustained metabolic load, such as oxidative stress (Wang & Michaelis 2010; Cobley et al.,

2018; Kempf et al., 2019). Furthermore, spiking activity may be mechanistically associated with synaptic

plasticity, and it was suggested that firing rates are an important determinant of overall changes in

synaptic strength, with higher spiking leading to greater changes (Graupner et al., 2016; Lappalainen et

al., 2019). Compensatory processes exist within neurones to oppose cellular stress and related

homeostatic challenges (Kültz 2005) and off periods could provide the opportunity for neurones to

prioritise such processes, therefore mediating the restorative benefits of sleep (Vyazovskiy & Harris,

2013). Similarly, off periods are associated with distinct synaptic plasticity rules (Gonzalez-Rueda et al.,

2018), and so their prevalence and patterning is likely also related to whatever regulation of synaptic

strength occurs during sleep (Tononi & Cirelli 2014; Timofeev & Chauvette 2017; Seibt & Frank, 2019).

According to this view of sleep homeostasis, in principle, regulation can occur entirely at the local level, within cortical networks or perhaps even single neurones. Indeed, this is reflected in our results through the differences between channels both with respect to the accuracy of the model and in the values of optimised parameters. Why then is global sleep preferred over a hypothetical state including asynchronous local off periods, which presumably could be used to sustain longer periods of behavioural wakefulness? It has been argued previously that it is ecologically optimal to synchronise off periods and undergo dedicated periods of total behavioural shutdown, because local off periods during waking impair behaviour (Vyazovskiy et al., 2011; Rattenborg et al., 2012). Mechanistically, the occurrence of off periods might be obstructed by strong synaptic coupling and shared neuromodulatory tone. Indeed, it has been observed that the degree of coupling between an individual neurone's firing and the population firing rate, is variable between cells but characteristic to an individual neurone and likely reflects total synaptic strength with its neighbours (Okun et al., 2015). Some neurones may therefore be less able to express asynchronous off periods than others.

The preference for global sleep, despite its fundamentally local mechanisms, may be evidence that sleep homeostasis ultimately does not serve a single specific local function. Instead, the recent history of local activities, by alteration of the local propensity to generate off periods (Process S), may be integrated over neuronal populations through intrinsic network mechanisms, in order to produce a sleep propensity signal that estimates the total time spent awake with great accuracy. The brain would thereby aim to enforce a daily quota of sleep, which could have many physiological and ecological benefits, rather than initiating sleep in response to the homeostatic need of one specific regulated variable (Vyazovskiy, 2015). The overall accuracy of both firing-rate-based and vigilance-state-based models supports this possibility, as does the evidence that homeostatically regulated cellular variables can actually be stable during extended wakefulness and that maintenance processes in sleep are ultimately prophylactic (Vyazovskiy & Harris, 2013). In this case, the sleep quota typically required by the brain would be determined by the neural population(s) that regularly accumulate sleep pressure the fastest. In order to maximise the efficiency of global sleep, single neuronal activities may be modulated such that sleep pressure accumulates, on average, as uniformly as possible. This is consistent with recent reports that sleep regulates the population level firing rate distribution (Watson et al., 2016;

Levenstein et al., 2017; Miyawaki & Diba, 2019) and could account for sleep's link with synaptic and firing rate homeostasis, reconciling the local origins of sleep pressure accumulation with the global level of its dissipation.

Here, spike firing rate is used to represent the level of neural activity because it is convenient to record, locally variable, and directly linked to neuronal functionality. However, no strong claim is made that firing rates are necessarily causally responsible for the accumulation of sleep pressure. Indeed, it would likely be possible to obtain a reasonable quantitative account of sleep homeostasis using any physiological variable (or set of variables) that are consistently higher in either the wake or sleep state. For example, a model assuming that Process S increases in proportion to local cortical temperature, which drops by ~2°C when falling asleep (Franken et al., 1991b) and which has been mechanistically implicated in sleep regulation (Hoekstra et al., 2019), might also provide a plausible description of Process S dynamics. Importantly, these results demonstrate that firing rates are a useful measurable correlate of the processes that directly underpin Process S, and therefore firing rate variance resulting from differences in experience and behaviour may well account for the variance in Process S accumulation in normal individuals, between waking periods and between cortical regions.

It should be addressed explicitly that the novel firing rate based models typically slightly under-perform, relative to the classic model, in terms of minimising the error between simulation and empirical data. There are a number of reasons why a limit on the accuracy of the model is to be expected, related to technical restrictions rather than conceptual ones. In this approach, each MUA channel records only a few randomly sampled nearby neurones, which may have very different spiking properties, and so grouping these as a single measure of local network activity is somewhat artificial. It may be valuable to explore genetic and pharmacological manipulations of firing rate in order to further test this model's validity. Unfortunately, the effects of any manipulation on the accumulation, expression and dissipation of sleep pressure, separately to the effects on firing, is unknown, and so this may be hard to interpret. For example, a recent study found that systemic atropine administered during behavioural wakefulness produces slow wave activity, reducing spiking (as measured by c-Fos) yet increasing the duration of subsequent NREM sleep (Qiu et al., 2015). While this was interpreted as evidence that spiking activity

is not related to sleep pressure accumulation, the direct effects of atropine on Process S and the

functional significance of the resulting induced slow oscillation are unclear. Similarly, another study

found that optogenetic activation of cortex during sleep, despite raising firing rates to waking levels,

was accompanied with decreased Process S (lower SWA and fewer off periods) during subsequent

NREM sleep (Rodriguez et al., 2016). Again, it is difficult to disentangle the direct effects of this

stimulation, which is not physiologically realistic, on the mechanisms surrounding the accumulation,

expression and dissipation of sleep pressure. On the other hand, these results might be interpreted as

evidence that the expression of the level of Process S involves an integration of homeostatic sleep need

across neural populations and therefore local firing rate manipulations are not able to substantially

influence Process S dynamics at a global level.

It is reasonable to assume that sleep homeostasis unfolds on multiple time scales and Process S as

defined by these models describes a relatively fast one, approaching its upper asymptote after

continuous wakefulness on a time scale of hours. The inclusion of processes acting over longer or

shorter time scales might explain discrepancies in all these models, however, the challenge remains to

identify what these could be. Furthermore, the role of REM sleep in Process S dynamics has not been

explicitly addressed or considered in the construction of these models. Depending on the model variant,

REM sleep is associated either with small increase or small decrease in Process S, because firing rates

are low in REM sleep (closer to NREM sleep than waking) and yet off period occupancy is also low (closer

to wake levels than NREM sleep). It is possible that REM sleep might represent a homeostatically neutral

state, in which the level of Process S changes little, or not at all (Vyazovskiy & Delogu, 2014).

In summary, Process S is reflected in both slow wave activity and the occurrence of local off periods, and

its dynamics can be quantitatively described using information derived from local spike firing rates and

off periods. Such a description has comparable accuracy to the classical model of Process S, dependent

only on global sleep-wake history. This result has important implications for our understanding of the

nature of sleep homeostasis and these novel models provide a methodology to quantify and compare

Process S dynamics, potentially between different cell types and brain regions.

References

Achermann, P., Dijk, D.-J., Brunner, D.P., and Borbély, A.A. (1993). A model of human sleep homeostasis

based on EEG slow-wave activity: Quantitative comparison of data and simulations. Brain Research

Bulletin *31*, 97–113.

Attwell, D., and Laughlin, S.B. (2001). An energy budget for signaling in the grey matter of the brain. J.

Cereb. Blood Flow Metab. *21*, 1133–1145.

Bellesi, M., Bushey, D., Chini, M., Tononi, G., and Cirelli, C. (2016). Contribution of sleep to the repair of

neuronal DNA double-strand breaks: evidence from flies and mice. Scientific Reports *6*, 36804.

Borbély, A.A. (1982). A two process model of sleep regulation. Hum Neurobiol *1*, 195–204.

Bukhtiyarova, O., Soltani, S., Chauvette, S., and Timofeev, I. (2019). Slow wave detection in sleeping

mice: Comparison of traditional and machine learning methods. Journal of Neuroscience Methods *316*,

35–45.

Buzsáki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and currents —

EEG, ECoG, LFP and spikes. Nature Reviews Neuroscience *13*, 407–420.

Chauvette, S., Volgushev, M., and Timofeev, I. (2010). Origin of Active States in Local Neocortical

Networks during Slow Sleep Oscillation. Cereb Cortex *20*, 2660–2674.

Cirelli, C., Faraguna, U., and Tononi, G. (2006). Changes in brain gene expression after long-term sleep

deprivation. Journal of Neurochemistry *98*, 1632–1645.

Cobley, J.N., Fiorello, M.L., and Bailey, D.M. (2018). 13 reasons why the brain is susceptible to oxidative

stress. Redox Biology *15*, 490–503.

Daan, S., Beersma, D.G., and Borbély, A.A. (1984). Timing of human sleep: recovery process gated by a

circadian pacemaker. Am. J. Physiol. *246*, R161-183.

Fisher, S.P., Cui, N., McKillop, L.E., Gemignani, J., Bannerman, D.M., Oliver, P.L., Peirson, S.N., and

Vyazovskiy, V.V. (2016). Stereotypic wheel running decreases cortical activity in mice. Nature

Communications *7*, ncomms13138.

569 Frank, M.G., and Heller, H.C. (2019). The Function(s) of Sleep. Handb Exp Pharmacol.

570 Franken, P., Dijk, D.J., Tobler, I., and Borbély, A.A. (1991a). Sleep deprivation in rats: effects on EEG

571 power spectra, vigilance states, and cortical temperature. American Journal of Physiology-Regulatory,

572 Integrative and Comparative Physiology *261*, R198–R208.

573 Franken, P., Tobler, I., and Borbély, A.A. (1991b). Sleep homeostasis in the rat: simulation of the time

574 course of EEG slow-wave activity. Neurosci. Lett. *130*, 141–144.

575 Funk, C.M., Honjoh, S., Rodriguez, A.V., Cirelli, C., and Tononi, G. (2016). Local Slow Waves in Superficial

576 Layers of Primary Cortical Areas during REM Sleep. Curr. Biol. *26*, 396–403.

577 González-Rueda, A., Pedrosa, V., Feord, R.C., Clopath, C., and Paulsen, O. (2018). Activity-Dependent

578 Downscaling of Subthreshold Synaptic Inputs during Slow-Wave-Sleep-like Activity In Vivo. Neuron

579 *97*, 1244-1252.e5.

580 Graupner, M., Wallisch, P., and Ostojic, S. (2016). Natural Firing Patterns Imply Low Sensitivity of

581 Synaptic Plasticity to Spike Timing Compared with Firing Rate. J Neurosci *36*, 11238–11258.

582 Grosmark, A.D., Mizuseki, K., Pastalkova, E., Diba, K., and Buzsáki, G. (2012). REM Sleep Reorganizes

583 Hippocampal Excitability. Neuron *75*, 1001–1007.

584 Guillaumin, M.C.C., McKillop, L.E., Cui, N., Fisher, S.P., Foster, R.G., de Vos, M., Peirson, S.N., Achermann,

585 P., and Vyazovskiy, V.V. (2018). Cortical region specific sleep homeostasis in mice: effects of time of

586 day and waking experience. Sleep.

587 Hengen, K.B., Lambo, M.E., Van Hooser, S.D., Katz, D.B., and Turrigiano, G.G. (2013). Firing Rate

588 Homeostasis in Visual Cortex of Freely Behaving Rodents. Neuron *80*, 335–342.

589 Hengen, K.B., Pacheco, A.T., McGregor, J.N., Van Hooser, S.D., and Turrigiano, G.G. (2016). Neuronal

590 firing rate homeostasis is inhibited by sleep and promoted by wake. Cell *165*, 180–191.

591 Hoekstra, M.M., Emmenegger, Y., Hubbard, J., and Franken, P. (2019). Cold-inducible RNA-binding

592 protein (CIRBP) adjusts clock-gene expression and REM-sleep recovery following sleep deprivation.

593 ELife *8*.

Huber, R., Deboer, T., and Tobler, I. (2000a). Effects of sleep deprivation on sleep and sleep EEG in

three mouse strains: empirical data and simulations. Brain Res. *857*, 8–19.

Huber, R., Deboer, T., and Tobler, I. (2000b). Topography of EEG dynamics after sleep deprivation in

mice. J. Neurophysiol. *84*, 1888–1893.

Huber, R., Ghilardi, M.F., Massimini, M., and Tononi, G. (2004). Local sleep and learning. Nature *430*,

78–81.

Kajikawa, Y., and Schroeder, C.E. (2011). How local is the local field potential? Neuron *72*, 847–858.

Kempf, A., Song, S.M., Talbot, C.B., and Miesenböck, G. (2019). A potassium channel β-subunit couples

mitochondrial electron transport to sleep. Nature *568*, 230–234.

Krueger, J.M., and Tononi, G. (2011). Local Use-Dependent Sleep; Synthesis of the New Paradigm. Curr

Top Med Chem *11*, 2490–2492.

Krueger, J.M., Rector, D.M., Roy, S., Dongen, H.P.A.V., Belenky, G., and Panksepp, J. (2008). Sleep as a

fundamental property of neuronal assemblies. Nature Reviews Neuroscience *9*, 910.

Kültz, D. (2005). Molecular and Evolutionary Basis of the Cellular Stress Response. Annual Review of

Physiology *67*, 225–257.

Lagarias, J., Reeds, J., Wright, M., and Wright, P. (1998). Convergence Properties of the Nelder--Mead

Simplex Method in Low Dimensions. SIAM J. Optim. *9*, 112–147.

Lappalainen, J., Herpich, J., and Tetzlaff, C. (2019). A Theoretical Framework to Derive Simple, Firing-

Rate-Dependent Mathematical Models of Synaptic Plasticity. Front. Comput. Neurosci. *13*.

Levenstein, D., Watson, B.O., Rinzel, J., and Buzsáki, G. (2017). Sleep regulation of the distribution of

cortical firing rates. Current Opinion in Neurobiology *44*, 34–42.

Liu, Z.-W., Faraguna, U., Cirelli, C., Tononi, G., and Gao, X.-B. (2010). Direct evidence for wake-related

increases and sleep-related decreases in synaptic strength in rodent cortex. J. Neurosci. *30*, 8671–

8675.

Mackiewicz, M., Shockley, K.R., Romer, M.A., Galante, R.J., Zimmerman, J.E., Naidoo, N., Baldwin, D.A.,

Jensen, S.T., Churchill, G.A., and Pack, A.I. (2007). Macromolecule biosynthesis: a key function of sleep.

Physiol. Genomics *31*, 441–457.

Massimini, M., Huber, R., Ferrarelli, F., Hill, S., and Tononi, G. (2004). The sleep slow oscillation as a

traveling wave. J. Neurosci. *24*, 6862–6870.

McKillop, L.E., Fisher, S.P., Cui, N., Peirson, S.N., Foster, R.G., Wafford, K.A., and Vyazovskiy, V.V. (2018).

Effects of Aging on Cortical Neural Dynamics and Local Sleep Homeostasis in Mice. J. Neurosci. *38*,

3911–3928.

Miyawaki, H., and Diba, K. (2016). Regulation of hippocampal firing by network oscillations during

sleep. Curr Biol *26*, 893–902.

Miyawaki, H., Watson, B.O., and Diba, K. (2019). Neuronal firing rates diverge during REM and

homogenize during non-REM. Scientific Reports *9*, 689.

Murphy, M., Huber, R., Esser, S., Riedner, B.A., Massimini, M., Ferrarelli, F., Ghilardi, M.F., and Tononi, G.

(2011). The cortical topography of local sleep. Curr Top Med Chem *11*, 2438–2446.

Okun, M., Steinmetz, N.A., Cossell, L., Iacaruso, M.F., Ko, H., Barthó, P., Moore, T., Hofer, S.B., Mrsic-

Flogel, T.D., Carandini, M., et al. (2015). Diverse coupling of neurons to populations in sensory cortex.

Nature *521*, 511–515.

O'Leary, T., Williams, A.H., Franci, A., and Marder, E. (2014). Cell types, network homeostasis, and

pathological compensation from a biologically plausible ion channel expression model. Neuron *82*,

809–821.

Qiu, M.-H., Chen, M.C., and Lu, J. (2015). Cortical neuronal activity does not regulate sleep homeostasis.

Neuroscience *297*, 211–218.

Quiroga, R.Q., Nadasdy, Z., and Ben-Shaul, Y. (2004). Unsupervised spike detection and sorting with

wavelets and superparamagnetic clustering. Neural Comput *16*, 1661–1687.

Rattenborg, N.C., Lima, S.L., and Lesku, J.A. (2012). Sleep Locally, Act Globally. Neuroscientist *18*, 533–

546.

Rector, D.M., Schei, J.L., Van Dongen, H.P.A., Belenky, G., and Krueger, J.M. (2009). Physiological

Markers of Local Sleep. Eur J Neurosci *29*, 1771–1778.

Reimund, E. (1994). The free radical flux theory of sleep. Medical Hypotheses *43*, 231–233.

Rodriguez, A.V., Funk, C.M., Vyazovskiy, V.V., Nir, Y., Tononi, G., and Cirelli, C. (2016). Why Does Sleep

Slow-Wave Activity Increase After Extended Wake? Assessing the Effects of Increased Cortical Firing

During Wake and Sleep. J. Neurosci. *36*, 12436–12447.

Rusterholz, T., and Achermann, P. (2011). Topographical aspects in the dynamics of sleep homeostasis

in young men: individual patterns. BMC Neurosci *12*, 84.

Saberi-Moghadam, S., Simi, A., Setareh, H., Mikhail, C., and Tafti, M. (2018). In vitro Cortical Network

Firing is Homeostatically Regulated: A Model for Sleep Regulation. Sci Rep *8*, 6297.

Sanchez-Vives, M., and Mattia, M. (2014). Slow wave activity as the default mode of the cerebral cortex.

Archives Italiennes de Biologie *152*, 147–155.

Scharf, M.T., Naidoo, N., Zimmerman, J.E., and Pack, A.I. (2008). The energy hypothesis of sleep

revisited. Prog. Neurobiol. *86*, 264–280.

Seibt, J., and Frank, M.G. (2019). Primed to Sleep: The Dynamics of Synaptic Plasticity Across Brain

States. Front Syst Neurosci *13*, 2.

Slomowitz, E., Styr, B., Vertkin, I., Milshtein-Parush, H., Nelken, I., Slutsky, M., and Slutsky, I. (2015).

Interplay between population firing stability and single neuron dynamics in hippocampal networks.

Elife *4*.

Steriade, M., Nuñez, A., and Amzica, F. (1993a). A novel slow (< 1 Hz) oscillation of neocortical neurons

in vivo: depolarizing and hyperpolarizing components. J. Neurosci. *13*, 3252–3265.

Steriade, M., Nuñez, A., and Amzica, F. (1993b). Intracellular analysis of relations between the slow (<

1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. J. Neurosci. *13*,

3266–3283.

Steriade, M., Timofeev, I., and Grenier, F. (2001). Natural waking and sleep states: a view from inside

neocortical neurons. J. Neurophysiol. *85*, 1969–1985.

Styr, B., Gonen, N., Zarhin, D., Ruggiero, A., Atsmon, R., Gazit, N., Braun, G., Frere, S., Vertkin, I., Shapira,

I., et al. (2019). Mitochondrial Regulation of the Hippocampal Firing Rate Set Point and Seizure

Susceptibility. Neuron *102*, 1009-1024.e8.

Timofeev, I., and Chauvette, S. (2017). Sleep slow oscillation and plasticity. Curr. Opin. Neurobiol. *44*,

116–126.

Tononi, G., and Cirelli, C. (2003). Sleep and synaptic homeostasis: a hypothesis. Brain Research Bulletin

62, 143–150.

Tononi, G., and Cirelli, C. (2014). Sleep and the Price of Plasticity: From Synaptic and Cellular

Homeostasis to Memory Consolidation and Integration. Neuron *81*, 12–34.

Turrigiano, G. (2011). Too Many Cooks? Intrinsic and Synaptic Homeostatic Mechanisms in Cortical

Circuit Refinement. Annual Review of Neuroscience *34*, 89–103.

Vergara, R.C., Jaramillo-Riveri, S., Luarte, A., Moënne-Loccoz, C., Fuentes, R., Couve, A., and Maldonado,

P.E. (2019). The Energy Homeostasis Principle: Neuronal Energy Regulation Drives Local Network

Dynamics Generating Behavior. Front Comput Neurosci *13*, 49.

Vyazovskiy, V.V. (2015). Sleep, recovery, and metaregulation: explaining the benefits of sleep. Nat Sci

Sleep *7*, 171–184.

Vyazovskiy, V.V., and Delogu, A. (2014). NREM and REM Sleep: Complementary Roles in Recovery after

Wakefulness. Neuroscientist *20*, 203–219.

Vyazovskiy, V.V., and Harris, K.D. (2013). Sleep and the single neuron: the role of global slow

oscillations in individual cell rest. Nat Rev Neurosci *14*, 443–451.

Vyazovskiy, V.V., and Tobler, I. (2008). Handedness leads to interhemispheric EEG asymmetry during

sleep in the rat. J. Neurophysiol. *99*, 969–975.

Vyazovskiy, V.V., Achermann, P., and Tobler, I. (2007). Sleep homeostasis in the rat in the light and

dark period. Brain Res. Bull. *74*, 37–44.

Vyazovskiy, V.V., Cirelli, C., Pfister-Genskow, M., Faraguna, U., and Tononi, G. (2008). Molecular and

electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. Nat

Neurosci *11*, 200–208.

Vyazovskiy, V.V., Olcese, U., Lazimy, Y.M., Faraguna, U., Esser, S.K., Williams, J.C., Cirelli, C., and Tononi,

G. (2009). Cortical firing and sleep homeostasis. Neuron *63*, 865–878.

Vyazovskiy, V.V., Olcese, U., Hanlon, E.C., Nir, Y., Cirelli, C., and Tononi, G. (2011). Local sleep in awake

rats. Nature *472*, 443–447.

Vyazovskiy, V.V., Cui, N., Rodriguez, A.V., Funk, C., Cirelli, C., and Tononi, G. (2014). The dynamics of

cortical neuronal activity in the first minutes after spontaneous awakening in rats and mice. Sleep *37*,

1337–1347.

Wang, X., and Michaelis, E.K. (2010). Selective neuronal vulnerability to oxidative stress in the brain.

Front. Aging Neurosci. *2*.

Watson, B.O., Levenstein, D., Greene, J.P., Gelinas, J.N., and Buzsáki, G. (2016). Network Homeostasis

and State Dynamics of Neocortical Sleep. ResearchGate.

Werth, E., Achermann, P., and Borbély, A.A. (1996). Brain topography of the human sleep EEG: antero-

posterior shifts of spectral power. Neuroreport *8*, 123–127.

Zavada, A., Strijkstra, A.M., Boerema, A.S., Daan, S., and Beersma, D.G.M. (2009). Evidence for

differential human slow-wave activity regulation across the brain. J Sleep Res *18*, 3–10.

 Figure Legends

Figure 1. Cortical spike firing patterns are associated with the dynamics of Process S. **A)** An example of

the classical state-based Process S model (blue) describing the dynamics of frontal EEG SWA (median

per NREM sleep episode, black bars) over 48 hours in one representative animal. Sleep deprivation

occurred as indicated at light onset of the second day and lasted 6 hours. Scored vigilance states are also

shown. **B)** An example of frontal electroencephalogram (EEG), primary motor cortical local field

potentials (LFP), corresponding raw signal with multi-unit activity (MUA), and detected spikes, in

representative segments of waking and NREM sleep. Slow waves and synchronous spiking off periods

are visible in NREM sleep but not wakefulness. **C)** The distribution of mean firing rate during wake,

NREM and REM sleep over the whole dataset, in addition to the difference in mean firing rate in wake

compared to NREM sleep (all are positive, reflecting higher firing in wake). Points indicate channels

grouped by animal (left to right), but boxplots reflect channels from all animals treated as a single

population. **D)** Distribution of correlation coefficients, calculated within each single channel, between

wake episode duration (Duration), the change in slow wave activity (dSWA), and mean firing rate (mean

FR). Points indicate channels grouped by animal (left to right), but boxplots reflect channels from all

animals treated as a single population. **E)** An example scatter plot of the correlation between the change

in median SWA from one NREM episode to the next and the mean firing rate during the intervening

period of wakefulness. This channel is representative because it has the median correlation coefficient

of all channels.

Figure 2. Slow wave activity dynamics at the LFP level can be modelled using multi-unit spiking

information. **A)** An example from one representative animal modelling the SWA averaged over all LFP

channels, of both the classical model (blue) and novel firing-rate-based model (orange), calculated from

the firing rate also averaged over all LFP channels (brown). **B)** An example of both models applied to

the SWA of a single LFP channel, which came from the same animal as used in **A**.

Figure 3. The fit quality and parameters for both classical and firing-rate-based models of LFP SWA. **A)**

Equations for the classic state-based model (blue) and novel firing-rate-based model (orange). **B)** For

each animal, the distribution over channels of the median difference between the model and empirical

SWA, expressed as a percentage of empirical SWA, for both classic and firing-rate-based models. **C)** The

same median percent error, grouped over animals and models, separately showing errors at the level of

single LFP, averaged LFP and EEG. **D-G)** The distribution of values used for each parameter in the classic

model and **H-L)** in the firing-rate-based model, with boxplots plotted separately for each animal. The

black lines indicate the values obtained from modelling the averaged LFP SWA and firing rate over all

channels within an animal. Additionally, the red lines in **D-G** give the parameter value obtained from the

model of the frontal EEG SWA of that animal. **M)** The distribution of the final optimised value for the

firing rate set point parameter (F_θ) of the firing-rate-based model, z-normalised to the distribution of

firing rates within wake, NREM and REM sleep. Points indicate channels grouped by animal (left to

right), but boxplots reflect all channels treated as a single population. **N)** The distribution of mean firing

rate in wake, NREM and REM sleep, expressed as a percentage of the firing rate set point parameter (F_θ).

Points indicate channels grouped and coloured by animal, but boxplots reflect all channels treated as a

single population. **O)** The distribution of values of the change in Process S ($\Delta S / \Delta t$) from one 4-second

time step to the next derived from the Fr-SWA model in wake, NREM sleep and REM sleep. All mice,

channels and time are pooled.

Figure 4. The definition of off periods and off occupancy. **A)** An example section of LFP (raw in grey,

0.5-6Hz filtered in black) and simultaneous MUA spikes. During this time window, the filtered LFP

crosses the amplitude threshold (265 µV for this channel, red line) five times. The multi-unit inter-spike

interval aligned to two of these peaks exceeds the duration threshold (85ms for this channel) and so

two off periods are detected (grey boxes). ISIs aligned to the other three out of five crossings (asterisks)

are too short to be considered off periods. **B)** Histograms of multi-unit inter-spike intervals aligned with

detected slow waves for this example channel. The four plots show, from left to right, ISIs over the whole

recording and separately ISIs from wake, NREM and REM sleep only. The ISI duration threshold (red

line) is selected using the histogram of all ISIs (leftmost) at the minimum between the two modes. **C)**

The distribution of LFP amplitude and **D)** ISI duration threshold values used for definition of off periods

for each channel, with boxplots plotted separately for each animal. **E)** The mean multi-unit firing rate

over a 1-second period centred on the peak of detected slow waves, calculated over all slow waves

within one example channel with 1-millisecond resolution. **F)** Distributions of mean off occupancy (%)

for all channels averaged over wake, NREM and REM sleep. Points indicate channels grouped by animal (left to right), but boxplots reflect all channels treated as a single population. **G)** Off occupancy is shown alongside EEG and LFP SWA for an example channel over 48 hours. Traces represent these values calculated at 4-second resolution (light grey), in addition to the median value per NREM sleep episode, as used for model fitting (black bars). Firing rate (brown) and scored vigilance states are also shown.

Figure 5. Process S is reflected in an LFP channel's off occupancy and its dynamics are described well by both state-based and firing-rates-based models. **A)** An example of the novel model based on firing rates and off occupancy (purple), and the classic state-based model (blue), with optimised parameters describing the dynamics of off occupancy (median per NREM episode, black bars) over 48 hours. Sleep deprivation occurred as indicated at light onset of the second day and lasted 6 hours. Firing rate (brown), off occupancy (value per 4-second epoch, grey) and scored vigilance states are also shown. **B)** Equations for the classic state based model (blue) and firing-rate-and-off-occupancy-based model for off occupancy (purple). **C)** For each animal, the distribution over channels of the median difference between the model and empirical off occupancy, expressed as a percentage of the off occupancy, for both classic and firing-rate-based models. **D)** The distribution of values of the change in Process S ($\Delta S/\Delta t$) from one 4-second time step to the next derived from the Fr-SWA model in wake, NREM sleep and REM sleep. All mice, channels and time are pooled. **E-H)** The distribution of optimised values used for each parameter in the classic model and **I-L)** the firing rate model, with boxplots plotted separately for each animal.

Figure 6. The time course of Process S is similar between models and individual channels. Process S time courses, expressed for each channel as a percentage of the maximum value, overlaid for all channels within a single representative animal obtained from **A)** the classic state-based model applied to LFP SWA, **B)** the firing-rate-based model applied to LFP SWA, **C)** the classic state-based model applied to off occupancy, **D)** the firing-rate-based model applied to off occupancy. In these panels the black line indicates the mean Process S over all channels. **E)** The mean Process S calculated over all channels is now plotted in colour (Cl-SWA light blue, Fr-SWA orange, Cl-Off dark blue, Fr-Off purple), and the red line indicates the Process S obtained by applying model Cl-SWA to the EEG derived SWA.

Figure 3

Figure 4

Figure 5

Figure 6