# Loss of *AUXIN RESPONSE FACTOR 4* function alters plant growth, stomatal functions and improves tomato tolerance to salinity and water deficit

Sarah BOUZROUD[1,2], Maria Antonia Machado BARBOSA[3], Karla GASPARINI[3], Mouna FAHR[1], Najib BENDAOU[1], Mondher BOUZAYEN[2], Agustin ZSÖGÖN[3], Abdelaziz SMOUNI[1], Mohamed ZOUINE[2]

[1]*Laboratoire de Biotechnologie et Physiologie Végétales, Centre de biotechnologie végétale et microbienne biodiversité et environnement, Faculté des Sciences, Université Mohammed V de Rabat – Maroc*

[2]*GBF, Université de Toulouse, INRA, Castanet-Tolosan, France*

[3]*Departamento de Biologia Vegetal, Universidade Federal de Viçosa, 36570-900, Viçosa, MG, Brazil*

*Corresponding author: Abdelaziz SMOUNI (azizsmouni@gmail.com)*

## Summary

Understanding physiological and molecular basis of plant response and tolerance to environmental stresses is important to improve plant survival, crop yield and quality. Auxin controls many aspects of plant growth and development. However, its role in stress responses remains so far poorly studied. Auxin acts on the transcriptional regulation of target genes, mainly through Auxin Response Factors (*ARF*). The current study aims to study the involvement of *SlARF4* in tomato tolerance to salinity and water deficit. Using a reverse genetic approach, we found that the down-regulation of *SlARF4* improves tomato tolerance to salt and drought stress by promoting root development and density, increasing soluble sugars content and maintaining chlorophyll content at high levels in stress conditions. Besides that, *ARF4*-as plants displayed a leaf curl phenotype, a low stomatal conductance coupled with an increase in leaf relative water content and ABA content under normal and stressful conditions. This increase in ABA content was correlated with the activation of ABA biosynthesis genes

25    and the repression of ABA catabolism genes. *cat1*, *Cu/ZnSOD* and *mdhar* genes were up-

26    regulated in *ARF4*-as plants suggesting that *ARF4*-as mutant is more tolerant to salt and water

27    stress.The data presented in this work strongly support the involvement of ARF4 as a key

28    actor in tomato tolerance to salt and drought stresses.

29    **Keywords:** *ARF4*, Auxin, salt, drought, tomato, tolerance.

**Significance statement**

31    Down regulation of *Auxin Response Factor* 4 modify leaf structure, alters stomatal responses

32    under normal conditions, and improves tomato tolerance to salt and drought stress.

**Introduction**

Auxin is a key regulator of many plant growth and development processes throughout the

plant life cycle (Vanneste and Friml, 2009). Auxins regulate diverse cellular and

developmental responses in plants via three types of transcriptional regulators, auxin response

factors (ARFs), Aux/IAA and TOPLESS (TPS)  proteins (Reed, 2001) Szemenyei, Hannon &

Long 2008). ARFs play a key role in regulating the expression of auxin response genes

(Liscum and Reed, 2002). 23 *ARF* genes have been isolated from *Arabidopsis thaliana*

(Liscum and Reed, 2002; Tom J Guilfoyle and Hagen, 2007). Most of them share a structure

with conserved domains: an  N-terminal DNA-binding domain (DBD), a variable central

transcriptional regulatory region (MR), which can function as an activator or repressor

domain, and a carboxy-terminal dimerization domain (CTD) that contributes to the formation

of either ARF/ARF homo- and hetero-dimers or ARF/Aux/IAA hetero-dimers (Ulmasov *et*

*al.*, 1999; Tom J. Guilfoyle and Hagen, 2007; Zouine *et al.*, 2014).

Since *ARF1*, the first *Arabidopsis* ARF gene, was cloned and its function investigated

(Ulmasov *et al.*, 1997), the ARF gene family has been identified and well characterized in

many crop species, including tomato (*Solanum lycopersicum*) (Wu et al., 2011; Zouine et al.,

2013), maize (*Zea mays*) (Xing et al., 2011), rice (*Oryza sativa*) (Wang et al., 2007; Jain and

Jitendra P Khurana, 2009), poplar (*Populus trichocarpa*) (Kalluri et al., 2007), Chinese

cabbage (*Brassica rapa*)(Mun et al., 2012), sorghum (*Sorghum bicolor*) (Ha et al., 2013),

banana (Hu et al., 2015) and physic nut (*Jatropha curcas* L.) (Tang et al., 2018).

Genetic and molecular studies conducted in *Arabidopsis* and other plant species showed that

ARF proteins are key factors in regulatory networks controlling plant growth and

development (Chandler, 2016). Indeed, ARF mutations can affect plant morphology and

physiology, and thus can affect plant development and interaction with the environment. In

*Arabidopsis*, a loss-of-function allele of *AtARF2*, has pleiotropic effects on vegetative organ

growth (Schruff *et al.*, 2005). An *Atarf8* mutation affects hypocotyl elongation and auxin

homeostasis  (Goetz *et al.*, 2006) while *AtARF7* loss of function mutation impairs the

hypocotyl response to blue light and reduces the auxin response (Harper *et al.*, 2000). The

*Atarf7/Atarf19* double mutant shows aberrant lateral root formation and abnormal

gravitropism in both hypocotyls and roots (Narise *et al.*, 2010). Double mutant *Atarf3/Atarf4*

plants display adaxialized tissues in the shoot, resembling mutations in the *KANADI*

transcriptional repressor (Pekker *et al.*, 2005). In tomato, down-regulation of *ARF3* decreased

the density of epidermal cells and trichomes (Zhang *et al.*, 2015). Transcriptional down-

regulation of *ARF4* expression leads to severe leaf curling along the longitudinal axis (Jones

*et al.*, 2002). The *ARF* gene family is also involved in the control of many physiological

processes. In tomato, the *Slarf2A/B* mutation leads to a severe ripening inhibition with

dramatically reduced ethylene production, while the over-expression of *ARF2A* resulted in a

blotchy ripening pattern in fruit as a result of a significant accumulation of ripening-related

genes and metabolites (Hao *et al.*, 2015; Breitel *et al.*, 2016). Down-regulation of *SlARF6* and

*SlARF8* in transgenic tomato plants through the overexpression of *miR167a* may lead to floral

development defects and female sterility (N., Liu *et al.*, 2014). The down-regulation of

*SlARF4* resulted in ripening-associated phenotypes such as enhanced firmness, sugar and chlorophyll content leading to dark green fruit and blotchy ripening (Jones *et al.*, 2002; Sagar *et al.*, 2013). This phenotype was also reported in *SlARF10* gain of function mutant (Mei *et al.*, 2018). *ARF* genes also seem to be implicated in plant responses to environmental stresses. It has been reported that both *OsARF11* and *OsARF15* show differential expression in salt stress conditions, suggesting that they are involved in the rice response to stress (Jain and Khurana, 2009). In banana, the expression of many *ARF* genes was altered in response to salinity and osmotic stresses (Hu *et al.*, 2015). In tomato, many *SlARF* genes are responsive to a wide range of abiotic stress conditions (Bouzroud *et al.*, 2018). Among them, *SlARF4* expression was significantly regulated by salinity and water deficit. *SlARF4* regulatory region was enriched in cis-*acting* elements specific to salinity and water deficit response (Bouzroud *et al.*, 2018). Moreover, *SlARF4* loss of function mutation induced modification in plant morphology and physiology. Indeed, *ARF4* down-regulated plants showed a severe upward curling along the longitudinal axis of leaves with an increase in chlorophyll content (Jones *et al.*, 2002). We carried out morphological, anatomical, physiological and molecular analyses to approach the possible involvement of *SlARF4* in tomato response to salinity and drought stress.

**Results and discussion**

**An *SlARF4* downregulated line shows altered anatomical, morphological and physiological parameters**

We analyzed previously published *AUXIN RESPONSE FACTOR 4* antisense RNA post-transcriptionally silenced lines (*ARF4*-as) (Sagar *et al.*, 2013). The *ARF4*-as plants exhibit a wide range of morphogenic phenotypes, namely delayed flowering, increased height and leaf curling as compared to their isogenic Micro-Tom (wild-type, WT) counterparts (Figure 1). Stem and root dry weight, on the other hand, showed a significant reduction in *ARF4*-as plants

compared to WT, along with an insignificant increase in leaf area and specific leaf area

(Figure 2).

*ARF4*-as plants analyzed in this work displayed a severe upward curling along the

longitudinal axis of leaves (Figure 1c), consistent with those observed in DR12-ASL lines

generated in the Kemer cultivar by Jones *et al.* (2002) and those obtained in WT by Sagar *et*

*al.* (2013). The curled leaf morphology observed in *ARF4*-as plants could result from

modification in cell division or elongation of leaf tissues (Volkenburgh, 1999). The extent of

the changes detected in leaf morphology prompted us to check leaf ultrastructure in WT and

*ARF4*-as plants. Leaf thickness was clearly reduced in *ARF4*-as plants as compared to wild-

type (Figure 3a, b). In fact, the average of leaf blade thickness was 209 ± 8 µm in WT

compared to 181 ± 7 µm in *ARF4*-as (Figure 3c). The palisade parenchyma (PP) and spongy

parenchyma (SP) tissue layers were thinner in *ARF4*-as plants than in WT plants (Figure 3d-

e). The palisade:spongy mesophyll ratio was significantly higher in WT than in *ARF4*-as

(Figure 3f). Meanwhile, the intracellular air spaces (IAS) were more conspicuous in *ARF4*-as

compared to WT, while no visible difference was observed in adaxial or abaxial epidermis

between genotypes (Figure 3g-h).

Consistent with the presence of thinner leaves in the *ARF4*-as mutant plants, photosynthetic

performance estimated through photosynthetic $CO_2$ assimilation rate (*A*) was significantly

lower than in the WT (Table 1). It is well known that *A* can be limited by the slowest of two

biochemical processes: (1) the maximum rate of ribulose 1,5-biphosphate (RuBP)

carboxylase/oxygenase (Rubisco) catalyzed carboxylation ($V_{cmax}$) and (2) regeneration of

RuBP controlled by electron transport rate ($J_{max}$) (Bernacchi *et al.*, 2009). Consistently, $V_{cmax}$

and $J_{max}$ exhibited a significant decrease in *ARF4*-as plants, indicating that photosynthesis is

reduced due to the silencing of *SlARF4* (Figure 4a).

Stomatal conductance is one of the main parameters that affect photosynthesis under drought

stress (Jiang *et al.*, 2006; Siddique *et al.*, 1999). Because stomatal behavior is closely linked

to $CO_2$ uptake (Wong *et al.*, 1979), it is considered to be one of the main reasons for reduced

photosynthesis (Ashraf and Harris, 2013; Medrano *et al.*, 2002). The response of stomatal

conductance ($g_s$) to changing air vapor pressure deficit (*VPD*) was determined in *ARF4*-as and

WT plants. At low *VPD*, wild-type plants exhibited high $g_s$ and tended to reduce $g_s$ as *VPD*

increased from 1.25 kPa to 2 kPa and maintained a moderate $g_s$ values at higher *VPD*s.

Meanwhile, *ARF4*-as plants presented a relatively low $g_s$ values at both low and high *VPD*

compared to wild-type plants (Figure 4b). We verified a similarly impaired stomatal response

in *ARF4*-as plants in response to step changes in irradiance and $CO_2$ levels (Figure S1),

suggesting that *ARF4* could be an important player in the regulation of stomatal movements.

Furthermore, intrinsic water-use efficiency was notably higher in *ARF4*-as plants (Table 1)

suggesting that the down-regulation of *SlARF4* has the potential to improve the ration of

carbon assimilation to transpirational water loss. Plants' tolerance to abiotic stress depends on

their ability to adjust the relationship between water, transpiration, photosynthesis and water

use efficiency through stomatal changes in order to maximize $CO_2$ assimilation (Zhang *et al.*,

2006). In fact, the decrease in stomatal conductance can enhance plant tolerance to drought

stress of many plant species such as chickpea (*Cicer arietinum*), durum wheat (*Triticum*

*turgidum*), Rice (*Oryza sativa*) and cotton (*Gossypium hirsutum*) (Flowers and Yeo, 1981)

(Brugnoli and Lauteri, 1991) (Mafakheri *et al.*, 2010) (Rahnama *et al.*, 2011). *ARF4*-as plants

showed a decrease in stomatal conductance coupled with an increase in water-use efficiency,

however, this can generally lead to a yield penalty through stomatal limitation to carbon

assimilation (Blum, 2009; Blum, 2005). Given this potential impact on carbon assimilation,

we next sought to examine whether *SlARF4* under-expression might affect plant agronomic

parameters. Yield, number of fruits and fruit fresh weight were not altered between WT and

*ARF4*-as plants, whereas fruit shape was altered in *ARF4*-as plants (Figure S3), as expected,

given the well-known role of auxin in fruit development (De Jong *et al.*, 2009; Hendelman *et*

*al.*, 2012; Chaabouni *et al.*, 2009; Su *et al.*, 2014; Hao *et al.*, 2015; Breitel *et al.*, 2016).

Successful growth of plants depends mostly on plasticity in leaf anatomical characteristics,

which enables plants to cope with diverse stress environments (Pereira-Netto *et al.*, 1999).

Leaf architecture plays a vital role in resisting a variety of stresses including drought stress

(Balsamo *et al.*, 2006). Structural modifications in leaf anatomy contribute significantly to

increasing drought tolerance, particularly under limited moisture conditions (Rhizopoulou and

Psaras, 2003). Leaf rolling is a one of the common adaptation to water stress (Kadioglu *et al.*,

2012; Kadioglu and Terzi, 2007). It has been reported that greater leaf rolling may be an

important indicator linked to drought tolerance and may have a positive impact on crop yield

under water stress conditions (Fen *et al.*, 2015). Plants having leaf rolling mechanism exhibit

a resistance to drought and high temperature and had higher water use efficiency (Kadioglu *et*

*al.*, 2012). This was explained by the fact that leaf rolling reduced transpiration rate (Kadioglu

and Terzi, 2007; Kadioglu *et al.*, 2012). The combination of a marked leaf curling phenotype

associated with decreased $g_s$ in *ARF4*-as plants suggests that these plants might tolerate better

water deficit. We thus endeavored to assess the influence of *ARF4* on plants responses to

water and salt stress.

**_SlARF4_ expression is altered in response to salt and drought stresses**

The expression pattern of *SlARF4* was analyzed by real-time PCR in leaves and roots exposed

to salt or drought stress. *ARF4* gene expression was significantly regulated by salt and drought

stresses. *SlARF4* expression was significantly repressed in leaves and roots after 2hours of salt

treatment while its expression was significantly induced in leaves and roots after 24 hours

(Figure 5a-b). During drought treatment, *SlARF4* gene was slightly induced after 48 hours of

stress application in leaves (Figure 5c). In drought stressed roots, *SlARF4* was although

significantly induced after 5 days of stress exposure (Figure 5d). Tomato lines harboring

*pARF4*::*GUS* constructs were generated and used to assay *in planta* analysis of *ARF4*

expression under stress conditions. We observed strong expression in leaves, primary root and

root tip after 24 hours of salt stress exposure (Figure 5e). After five days of drought treatment,

GUS activity was detected in the root system and in different parts of the leaf (Figure 5f).

Extensive research has shown that various environmental signals are associated with changes

in auxin homeostasis, redistribution, and signaling (Navarro *et al.*, 2006; Park *et al.*, 2007).

Accumulating evidence indicates that auxin plays a role in plant responses to abiotic stresses

through complex metabolic and signaling networks. Genome-wide expression analyses have

suggested that the expression of numerous *ARF* genes change when plants respond to abiotic

stresses in various species including rice (Jain and Khurana, 2009), sorghum (Wang *et al.*,

2010), soybean (Van Ha *et al.*, 2013) and banana (Hu *et al.*, 2015). In tomato, expression

profiling of the *ARF* family under abiotic stress conditions showed that most of the tomato

*ARF*s were responsive and some of them were significantly regulated (Bouzroud *et al.*, 2018).

Among them, *SlARF4* expression was significantly induced in tomato plants in response to

salt (24 hours) or drought (five days). Additionally, *SlARF4* regulatory region was enriched in

*cis*-acting elements specific to salinity and water deficit response (Bouzroud *et al.*, 2018).  In

this work, *ARF4* expression assessed both *in vitro* and *in planta* was affected by either salt or

drought treatment which suggest that this gene might be involved in tomato response to those

stresses. We thus evaluated the broad consequences of drought and salt stress in transgenic

downregulated (*ARF4*-as) or constitutively expressing (*ARF*-ox) plants.

***ARF4* alters the plants response to salt and drought stress**

***Shoot and root fresh weight are differentially altered in ARF4 transgenic lines***

Shoot fresh weight was investigated in WT and *ARF4* transgenic plant lines under salt and

drought stress conditions. In the absence of stress, fresh weight was significantly higher in the

*ARF4*-ox as compared to the other lines. The average of shoot fresh weight was 1,5 and 2

times higher than that of *ARF4*-as and WT plants respectively (Figure 6a). Two levels of

NaCl (100 mM and 150 mM) were applied, and both led to reductions in shoot fresh weight in

all genotypes. In response to 150 mM NaCl, shoot fresh weight decreased by 60% and 70% in

WT  and *ARF4*-ox plants respectively and only by 30% in *ARF4*-as plants (Figure 6a). In

roots, the reduction in the fresh weight was around 55%, 28% and 65% for WT, *ARF4*-as and

*ARF4*-ox plants lines respectively (Figure 6c). Similarly to salt stress, drought stress affected

the growth of the three plant lines. Shoot fresh weight decreased by 66% and 82%

respectively for WT and *ARF4*-ox plants and by 44% for *ARF4*-as plants when exposed to

15% of PEG 20000 (Figure 6b). Root fresh weight deceases significantly with the increase of

PEG concentration. The decrease reached 57% respectively in WT and *ARF4*-ox plants while

the reduction was around 40% in *ARF4*-as in response to 15% of PEG20000 (Figure 6d). The

reduction in fresh weight was reported in many plant species which includes barley, sugar

beet, cabbage and *sorghum*  in response to salinity or water deficit (Jamil *et al.*, 2007; Farooq

*et al.*, 2009;  Fayez and Bazaid, 2014; Saddam *et al.*, 2014). The decrease in seedling growth

(evaluated through the measurement of fresh weight) is due to restricted cell division and

enlargement, as salt and drought stress directly reduces growth by decreasing cell division and

elongation (Akıncı and Lösel, 2012). However, the reduction in fresh weight was less

pronounced in the *ARF4*-as plants which suggest that *ARF4*-as plants might tolerate better salt

and drought stress than the other plant lines.

***Root development and density are less affected in ARF4-as plants***

Root development was investigated in *ARF4* transgenic lines in response to different

concentrations of NaCl or PEG. Our results showed a strong reduction in primary root length

in *ARF4*-ox plants (by 65% in salt stress condition and by 42% in drought stress condition)

while the decrease was nearby 40% and 15% in WT and *ARF4*-as plants in response to salt

and drought stress respectively (Figure 6e-f).

Root density was also investigated in the three lines under salt and drought stress conditions.

Our results showed that root density significantly increased with the increase of NaCl or PEG

concentrations in WT and in *ARF4*-as plants (Figure 6g-h). Root density increases by 57% in

WT and 120% in *ARF4*-as, respectively while the *ARF4*-ox showed a significant reduction of

in root density (10%) when exposed to 150 Mm of NaCl. In response to drought stress, the

three plant lines showed a significant increase in root density. At 5% of PEG 20000, root

density increased by 50%, 130% and 43% in WT, the *ARF4*-as and the *ARF4*-ox plants

respectively.

Root system is the first organ to encounter salinity and drought stress. Root development can

severely be affected by environmental stresses (Kurth *et al.*, 1986). In *Arabidopsis*, root

length and development was significantly reduced in response to salinity (Burssens *et al.*,

2000). In this study, we had noticed a significant decrease in primary root length in WT and in

*ARF4* transgenic lines in response to salt or drought stress conditions as compared with

normal conditions. However, less reduction was reported in the *ARF4*-as. Deeper rooting has

been shown to be beneficial for plant production and survival as it increases water uptake

which confer the advantage to support plant growth during adverse conditions (Comas *et al.*,

2013). Numerous studies have linked plant stress tolerance with the increase in root length

and density for several plant species such as barley, sunflower, wheat, rice and cotton (Munns

*et al.*, 2006). *ARF4*-as plants showed less reduction in primary root length coupled with an

increase in root density. *ARF4* was found to be implicated in defining root architecture in

*Arabidopsis* under optimal growth conditions through the control of lateral root emergence

(Marin *et al.*, 2010) which suggest that this gene might play an important role in root

architecture in stress conditions and thus contribute to the improvement tomato tolerance to

salinity and water deficit.

***ARF4-as plants are less affected by salt and drought stress***

***Photosynthesis is less affected in ARF4-as plants***

Photosynthesis is among the primary processes to be affected by salinity and drought stress

(Chaves, 1991; Munns *et al.*, 2006). Our results showed that total chlorophyll content

decreased significantly in WT and *ARF4*-ox plants exposed to different concentrations of

NaCl while no significant changes in chlorophyll content was detected in *ARF4*-as plants

(Figure S4). At 150 mM of NaCl, the total chlorophyll content decreased only by 4% in the

*ARF4*-as plants as compared to the control. Meanwhile, the diminution of total chlorophyll

content was around 53,2% and 62,8% respectively in the *ARF4*-ox line and in WT at 150 mM

of NaCl compared to the control (Figure S4). The drastic decline in the total chlorophyll

content was also observed as a result of drought stress application on WT plants and *ARF4*-ox

plants. The reduction in the total chlorophyll content averaged the 37% and the 15% in WT

and *ARF4*-ox plants cultivated at 15% of PEG 20000 respectively (Figure S4). Meanwhile,

total chlorophyll content increased significantly*ARF4*-as in response to drought stress. In fact,

the average of chlorophyll content was 1,2 times higher in *ARF4*-as at 15% PEG 20000 than

at 0% PEG 20000. Stress-induced decrease in chlorophyll content have been reported in

several plant species including tomato and wheat (Dogan *et al.*, 2010; Marcińska *et al.*, 2012;

Saeidi and Abdoli, 2015). Zarafshar *et al.*, (2014) explained this decrease as the result of

pigment photo-oxidation and chlorophyll degradation. *ARF4*-as plants showed although a

slight decrease in total chlorophyll content in response to salt stress and a slight increase in

drought stress conditions. tomato and Pea tolerant genotypes showed less losses in

photosynthetic pigments (Hernandez *et al.*, 2000; Furdi *et al.*, 2013). The lack of changes in

chlorophyll content in the *ARF4*-as plants shows the capacity to preserve the photosynthetic

apparatus and thus indicates their better tolerance to salinity and drought stress.

### *Sugars are highly accumulated in ARF4-as plants in stress conditions*

Soluble carbohydrates content was next investigated in tomato *ARF4* transgenic and WT lines

in response to different concentrations of NaCl or PEG.  In response to salt stress, leaf soluble

sugar content was decreased by 57, 5% and 69, 3% in WT and *ARF4*-ox lines respectively

and was increased by 146% in *ARF4*-as line exposed to 150 mM of NaCl for 2 weeks (Figure

S4). In roots, the soluble sugar content was also decreased in WT and *ARF4*-ox lines by 49,

4% and 51, 46% respectively, while an increase around the 191% in soluble sugar content was

detected in *ARF4*-as line. PEG induced drought stress imposed to plants significantly

decreased soluble sugar contents at all the stress levels in WT, *ARF4*-as and *ARF4*-ox lines. In

fact, leaf soluble sugar content decreased with the increase of PEG concentration. Leaf

soluble sugar content significantly decreased by 42,8%, 43% and 18% respectively in WT,

*ARF4*-ox and *ARF4*-as plants exposed to 15% of PEG 20 000. In roots, PEG application

induced a significant decrease in soluble sugar content by 41, 1% and 62, 27% (at 15% of

PEG 20 000) in WT plants and in *ARF4*-ox plants respectively. Meanwhile, a significant

increase in soluble carbohydrates content was although reported in *ARF4*-as plants and

reached the 67% at the high level of the PEG as compared to the control (Figure S4).

Accumulation of soluble carbohydrates has been associated with drought and salt tolerant

mechanisms in many plant species. Sugar accumulation help plants to tolerate abiotic stresses

by improving their ability to preserve osmotic balance within the cells (Atkinson & Urwin

2012). Tolerant species  and varieties accumulate more soluble sugars in leaves and growing

tissues than sensitive ones (Pattanagul and Thitisaksakul, 2008). Soluble sugars lead to buffer

cellular redox potential, help to stabilize membranes and protein structures and also provide

possible energy source under severe constraints (Keunen *et al.*, 2013; Rosa *et al.*, 2009; Sami

*et al.*, 2016). Sugar accumulation in the *ARF4*-as plants might improve tolerance to salinity

and water stress by increasing cellular osmolarity which promotes the influx of water into

cells, or at least a reduced water efflux providing the turgor necessary for cell expansion.

Carbohydrates produced in leaves during photosynthesis are transported throughout the plant

by SUTs in order to support plant growth. The *SUT1* proteins are involved in the movement

of sucrose into and out of the source and sink tissues and through the phloem via apoplastic

pathways. Thus, it is accepted that SUTs are important for plant growth and stress tolerance

(Jia *et al.*, 2015). Investigating the expression of the sucrose transporter *SlSUT1* under salt and

drought stresses had revealed that its expression was induced in WT and the *ARF4*-as leaves

and roots exposed to salt stress while *SlSUT1* was significantly repressed in the *ARF4*-ox

plant line (Figure S5). These findings were also detected in response to drought stress. In

drought stressed leaves, *SlSUT1* was significantly induced in WT and *ARF4*-as plants whereas

its expression was significantly down-regulated in *ARF4*-ox plants. In roots, the expression of

*SlSUT1* was also up-regulated significantly. The sucrose transporters have also been shown to

be regulated by abiotic stresses such as salt stress in potato celery and rice (Kühn *et al.*, 1997;

Noiraud *et al.*, 2000). *SUT1* transcript increased by 40% in sugarcane leaves after 24 hours

PEG treatment (Patade *et al.*, 2012). An increase in *SUT1* expression was also reported in

sweet sorghum salt tolerant genotype  (Sui *et al.*, 2015). Thus, the up-regulation of *LeSUT1* in

*ARF4*-as might improve its tolerance to drought and salinity through the stimulation of

sucrose transport required for the osmoregulation and for the cellular energy demands during

drought and salinity stresses.

***ARF4-as plants showed lower stomatal conductance***

Stomata play an imminent role in plant response to environmental changes as they control

both water losses and $CO_2$ uptake (Damour *et al.*, 2010). Stomatal conductance was estimated

in WT and *ARF4*-as plants in salt or drought stress (Figure 7a-b). Under normal conditions,

*ARF4*-as plants exhibited low stomatal conductance as compared to WT plants. Salt and

drought stress application induces a significant decrease in stomatal conductance in WT. In

fact, the reduction in stomatal conductance observed in the WT was around 32% and 46% in

response to salt and drought stress respectively. Meanwhile, no significant changes in

stomatal conductance were detected in *ARF4*-as plants that still maintained the same values of

stomatal conductance observed in normal conditions Salt and drought tolerance was

correlated with a decline in stomatal conductance in many plant species such as chickpea,

maize, common bean and cotton (Brugnoli and Lauteri, 1991; Heschel and Riginos, 2005;

Miyashita *et al.*, 2005; Mafakheri *et al.*, 2010; Anjum *et al.*, 2011) .

Stomatal closure can be trigged by various internal and external factors. Schultz, (2003) have

shown that the decline in leaf water potential can trigger stomatal closure under stressful

conditions. Relative water content (RWC) was investigated in WT and *ARF4*-as plants. A

remarkable decrease in RWC was observed in the three plant lines (Figure 7c). Up to 25% and

15% of water losses was recorded during the first 30 minutes of dehydration in WT and

*ARF4*-as plants respectively. RWC reached 72% in WT and after 4 hours of dehydration

while a 50% decrease of water losses was found in *ARF4*-as plants. The maintenance of a

relatively high RWC during mild drought is indicative of drought tolerance in wheat, barley,

sugarcane, rice and maize (Roy and Basu, 2009). The maintenance of higher RWC in *ARF4*-

as under stress conditions can possibly be the result of the stomatal closure and suggest that

these plants might better tolerate dehydration stress than WT.

**_ARF4_-as plants exhibited a high ABA content**

Abscisic acid is identified as the key hormone that mediates plant responses to adverse

environmental stimuli. Stomatal conductance can also be trigged by the subsequent

accumulation of ABA (Tombesi *et al.*, 2015). Endogenous ABA content of 5 weeks *ARF4*-as

and WT plants in normal or under salt and drought conditions were then measured in leaf

tissues. Our results showed that the ABA content in *ARF4*-as increased by 120% and 101%

under salt and drought treatment respectively whereas in WT plants increased by only 20% in

salt conditions and decreased by 26% in drought treatment (Table 2). The lowest stomatal

conductance observed in *ARF4*-as as compared with WT plants may be associated with the

accumulation of relatively more ABA in *ARF4*-as stressed leaves and suggests that *ARF4*-as

plants might tolerate better salt and drought stresses than WT plants.

Endogenous ABA content is related to the balance between its biosynthesis and its

degradation. In tomato, *SlNCED1* and *SlNCED2* encoding 9-cis-epoxycarotenoid dioxygenase

are the primary genes responsible for ABA biosynthesis, whereas *SlCYP707A1*, *SlCYP707A2*,

*SlCYP707A3* and *SlCYP707A4* encoding ABA 8'-hydroxylase are main genes for ABA

catabolism (Yang *et al.*, 2014). Investigating the expression of these genes in WT and the

*ARF4*-as plants had revealed that their expressions were significantly regulated by salt or

drought stress application (Figure 8). Indeed, *SlNCED1* expression was significantly induced

in *ARF4*-as leaves after 2 hours and 24 hours of salt stress exposure while no significant

changes was observed in roots. In WT, the expression of *SlNCED1* was significantly up-

regulated in leaves and roots after 24 hours of salt treatment. In response to drought stress, the

expression of *SlNCED1* was significantly up-regulated in *ARF4*-as roots while no significant

change in *SlNCED1* expression was detected in WT plants. *SlNCED2* expression was

significantly induced in the *ARF4*-as leaves after 2 hours and 24 hours of salt application and

in WT after 24 hours. A significant increase in the expression of *SlNCED2* was detected in

*ARF4*-as plants in response to drought stress. Water stress is known to induce the expression

of *NCED* genes in several plant species which include tomato, maize, common bean, maize,

Arabidopsis, cowpea and avocado (Tan *et al.*, 1997; Burbidge *et al.* 1999; Qin & Zeevaart

1999; Chernys and Zeevaart, 2000; Iuchi, Kobayashi, Yamaguchi-Shinozaki & Shinozaki

2000; Iuchi *et al.* 2001). Moreover, transgenic plants overexpressing the *NCED* gene

accumulated large amounts of ABA and were more resistant to drought stress (Thompson *et*

*al.* 2000; Wan & Li 2006).

The expression pattern of *SlCYP707A1*, *SlCYP707A2* and *SlCYP707A3*, key ABA

degradation genes was also assessed in *ARF4*-as and WT plants under salt and drought stress

conditions (Figure 8). Quantitative RT-PCR results showed that the expression of the ABA

metabolism genes was up-regulated in WT leaves after 2 hours and 24 hours of salt stress

exposure. In roots, the expression of *SlCYP707A1* and *SlCYP707A3* was up-regulated in WT

plants in response to salt stress while a significant repression was reported in the expression of

*SlCYP707A2*. Although, the expression of the ABA metabolism genes expression was

repressed in the *ARF4*-as salt stressed roots. In response to drought stress, the expression of

*CYP707A1*, *CYP707A2* and *CYP707A3* was repressed in *ARF4*-as plants while their

expression was up-regulated in WT roots and down-regulated in leaves. An increase in the

transcript of *CYP707A* genes was reported in response to salt or drought stress in many plant

species including common bean, Arabidopsis and Peanut (Saito *et al.* 2004; Yang & Zeevaart

2006; Liu *et al.* 2014). *Arabidopsis* knock-out mutant *cyp707a3-1* accumulated higher

endogenous ABA levels coupled with a reduced transpiration rate, thereby resulting in a

phenotype exhibiting enhanced tolerance to drought stress (Umezawa *et al.* 2006). Down-

regulation of *CYP707A* genes in *ARF4*-as plants in response to salt and drought stress might

prevent ABA degradation. The significant increase in *SlNCED1* expression in *ARF4*-as leaves

and roots along with the repression of the three ABA catabolism genes in normal and salt

stress conditions might explain the increase in the concentration of ABA in leaves.

**Antioxidant genes expression is altered in response to salinity and drought stress**

In plants, abiotic stresses induce the overproduction  of reactive oxygen species (ROS), a

highly  reactive and toxic molecules that cause damage to proteins, lipids, carbohydrates and

DNA and  results ultimately in oxidative stress (Das and Roychoudhury, 2014). The ability of

plants to mitigate the negative effects of abiotic stresses relays on the efficiency of the

antioxidant defense systems to protect plant cells from oxidative damage by scavenging ROS

accumulation The key enzymatic antioxidants are catalase, superoxide dismutase (SOD),

monodehydroascorbate    reductase    (MDHAR),    dehydroascorbate    reductase    (DHAR)

(Choudhury *et al.* 2013).

***Cat1 (Catalase) expression in response to salt and drought stress***

Catalases are tetrameric heme-containing enzymes with the potential to directly dismutate

$H_2O_2$ into $H_2O$ and $O_2$ and essential for ROS detoxification (Das and Roychoudhury, 2014).

In angiosperms, catalase is encoded by three genes known as *Cat1*, *Cat2* and *Cat3*

(Choudhury *et al.*, 2013). We investigated *Cat1* expression in WT and *ARF4* transgenic lines

exposed to salt and drought stresses. Our results showed that *Cat1* expression was induced in

*ARF4*-as plants exposed to salt stress while its expression was slightly repressed in the *ARF4*-

ox plants. A high up-regulation in *Cat1* expression was although detected in the wild type

plant roots in response to salt stress (Figure S6). In drought stressed leaves, *Cat1* expression

was significantly induced in WT while a remarkable but not significant repression was

observed in *ARF4* transgenic lines. Meanwhile, *Cat1* gene expression was highly up-regulated

in WT and *ARF4*-as plant roots (Figure S6). Numerous studies have linked abiotic stress

tolerance to an overproduction of catalase in many plant species such maize, potato and rice

(Moriwaki *et al.*, 2008; M'Hamdi *et al.*, 2009; Gondim *et al.*, 2012). An increase in *Cat*

transcript and activity was reported in wheat in response to drought stress and sweet potato

exposed to salt and drought stresses (Luna *et al.* 2005; Yong *et al.* 2017). The overproduction

of a bacterial *cat1* gene improved tolerance to salinity in tomato and tobacco (Al-Taweel *et al.*

2007; Mohamed *et al.* 2003). Introduction of the same catalase gene into Rice was reported to

increase resistance to salt (Moriwaki *et al.*, 2008). *Cat1* gene was significantly up-regulated in

*ARF4*-as plants might contribute to the improvement of their tolerance to salinity and water

stress.

***SOD (Superoxide dismutase) expression in response salt and drought stress***

Superoxide dismutase (SOD) is an enzyme that belongs to the family of metalloenzymes

ubiquitous in all aerobic organisms. It is one of the first defense against ROS induced

damages by catalyzing the removal of $O_2^-$ by dismutating it into $O_2$ and $H_2O_2$ (Choudhury *et*

*al.*, 2013; Das and Roychoudhury, 2014). Based on the metal ion it binds, SOD are classified

into three isozymes; Mn-SOD (in mitochondria), Fe-SOD (in chloroplasts) and Cu/ZnSOD (in

cytosol, peroxisomes and chloroplasts) (Das and Roychoudhury, 2014). Investigating the

expression of *Cu/ZnSOD* had revealed that its expression was remarkably repressed in WT

plants and *ARF4*-ox plants subjected to salt and drought stresses. Meanwhile, *Cu/ZnSOD* gene

expression was induced in *ARF4*-as leaves in response to salt and drought stresses while its

expression was repressed in the roots (Figure S6).

Abiotic stresses such as salinity and water deficit induce the generation of superoxide in plant

cells because of impaired  electron transport in the chloroplasts due in part to the

accumulation of high concentrations of iron (Wang, Wang, Kwon, Kwak & Su 2005). Several

studies had stated a significant increase in SOD activity in response to osmotic stress in many

plant species such as tomato (Gapińska *et al.*, 2007), Chickpea (*Cicer arietinum*) (Kukreja *et*

*al.*, 2005), potato (Aghaei *et al.*, 2009) and mulberry (Ahmad *et al.*, 2014). Pea (*Pisum*

*sativum*) tolerant genotypes responded to NaCl stress by increasing mitochondrial Mn-SOD

and chloroplastic CuZn-SOD and ascorbate peroxidase activities (Hernández *et al.*, 1993).

Szőllősi, (2014) reported an increase in the activity of total SOD in *Olea europaea* leaves or

in shoots of *Oryza sativa* cultivars in response to drought stress. An increase in SOD activity

was also detected in both tolerant and non-tolerant tomato species subjected to water deficit

(Rahman *et al.* 2004). This finding was also demonstrated in *P. vulgaris* (Zlatev *et al.*, 2006),

*Trifolium repens* (Chang-Quan and Rui-Chang, 2008) and *Picea asperata* (Yang *et al.*, 2008).

Genetic manipulation of genes encoding SOD isoforms increased plant tolerance to abiotic

stress conditions. In *Arabidopsis*, the constitutive expression of *MnSOD* gene improved salt

tolerance in transgenic plants (Wang *et al.*, 2004). The introduction of tomato Cu/ZnSOD in

potato plants showed increased tolerance to paraquat (Perl *et al.*, 1993). The overexpression

of Rice cytosolic Cu/Zn-SOD in chloroplasts of tobacco plant improved their photosynthetic

performance during photooxidative stresses such as high salt or drought stresses (Badawi *et*

*al.* 2004). Knowing that SOD iso-enzymes act as an early scavenger, keeping the superoxide

radical at low level, preventing oxidative damage and higher photosynthetic rate which

improves plant growth in limited conditions. The up-regulation of Cu/ZnSOD in the *ARF4*-as

might reduce oxidative stress caused by abiotic stress and might increase tomato survival

under salt and drought stress conditions.

***mdhar (Monodehydroxyascorbate reductase) expression in response to salt and drought***

***stress***

Monodehydroxyascorbate reductase (MDHAR) is another enzymatic antioxidant indirectly

involved in the ROS scavenging through regenerating Ascorbic acid, indispensable for the

dismutation of $H_2O_2$ by APX enzyme (Das and Roychoudhury, 2014). Investigating the

expression pattern of *mdhar* gene in salt stress conditions had revealed a significant up-

regulation in *ARF4*-as leaves and roots while its expression was repressed in *ARF4*-ox and

WT plants (Figure 18). In response to drought stress, the expression of *mdhar* was

significantly up-regulated and repressed respectively in WT and the *ARF4*-ox plant leaves. In

roots, *mdhar* is significantly up-regulated and down-regulated in the *ARF4*-as and *ARF4*-ox

plant line respectively. Several studies had underlined the protective role of *mdhar* in plant

tolerance to various abiotic stresses. (Sharma and Shanker Dubey, 2005) had reported an

increase in MDHAR activity in rice seedlings exposed to drought stress. This finding was also

stated in wheat (*Triticum aestivum*) and in mulberry (*Morus alba*) when subjected to drought

stress (Reddy *et al.*, 2004). The overexpression of *mdhar* in transgenic tobacco increased the

tolerance against salt and osmotic stresses (Eltayeb *et al.*, 2006). MDHAR plays an important

role in the Ascorbate- Glutathione ROS detoxification pathway through restoring ascorbate

(AsA) used as an electron donor to scavenger $H_2O_2$ molecules (Anjum *et al.*, 2010). Thus, the

up-regulation of *mdhar* gene in the ARF4-as plants might increase their tolerance to salinity

and water deficit by providing the necessary substrate for the degradation of $H_2O_2$.

**Conclusion**

In the face of a global scarcity of water resources and the increased water and soil

salinization, abiotic stresses present major challenges in sustaining crop yield. Plant tolerance

to these stresses relays on the implementation of molecular mechanisms for cellular

adjustments, including signal perception and transduction cascades, transcriptional networks

and adaptive metabolic pathways. Plant hormones such as ABA, ethylene and SA play an

important role in mediating plant responses to stresses. Auxin, the key regulator of many

aspects of plant growth and development, was identified as an actor in plant response to biotic

and abiotic stress. ARFs play crucial role in auxin signaling. *ARF4* is involved in the

regulation of several processes such as lateral root development, fruit quality and

development (Marin *et al.*, 2010; Jones *et al.*, 2002; Sagar *et al.*, 2013). In this work, we

investigated the involvement of this gene in the acquisition of salt and drought tolerance in

tomato. Using a reverse genetic approach, we found that the down-regulation of *SlARF4*

increases tomato tolerance to salinity and drought stress. Indeed, the down-regulation of

*SlARF4* increases root length and density resulting in a better development of root system

which might increase water acquisition and minimize water losses and thus stimulate plant

growth under adverse conditions. Furthermore, *ARF4*-as plants displayed a severe leaf

curling, a lower stomatal conductance, a higher water use efficiency under normal conditions

and maintained high level of chlorophyll content even in stress conditions suggesting that

their photosynthetic activity was less affected by the stress application. Total carbohydrates

were accumulated in high proportion in photosynthetic tissues of *ARF4*-as plants. At the

molecular level, the expression of the sucrose transporter *LeSUT1* was significantly up-

regulated in *ARF4*-as which might explain carbohydrates accumulation in root tissues. *cat1*,

*Cu/ZnSOD* and *mdhar* genes were found to be up-regulated in *ARF4*-as plants suggesting that

*ARF4*-as mutant is more tolerant to salt and water stress. Furthermore, *SlARF4*-as plants

showed a significant increase in ABA content associated with a low stomatal conductance

under stressful conditions. This increase in ABA content is due to the activation of ABA

biosynthesis genes and the repression of ABA catabolism genes. Taking together, the data

presented in this work brings new elements on auxin involvement in stress tolerance in tomato

and underline the role of *ARF4* in this process and provides new insights into tomato selection

and breeding for environmental stress-tolerant.

**Experimental procedure**

**Plant material**

Wild-type tomato (*Solanum lycopersicum*, L.) cv Micro-Tom (WT), and the following *AUXIN*

*RESPONSE FACTOR 4* lines were used in this study: an *ARF4* RNAi antisense silenced line

(*ARF4*-as), a line with *ARF4* expression driven by the constitutive cauliflower mosaic virus

(CaMV) promoter 35S (*ARF4*-ox), and a reporter line with the β-glucoronidase (GUS) gene

driven by the native *ARF4* promoter region (pARF4::GUS). *ARF4*-as plant lines and *ARF4*-ox

plant lines were previously generated and well-characterized by Sagar *et al.*, (2013). For each

mutant, the most representative plant line was chosen.

**Characterization of *SlARF4*-as plants**

*Plant growth conditions*

Seeds of WT and *ARF4*-as transgenic plants were sown in germination trays with commercial

substrate (Tropstrato HT Hortaliças, Vida Verde) and supplemented with 1g L$^-$1 10:10:10

NPK and 4 g L$^{-1}$ dolomite limestone ($MgCO_3$ + $CaCO_3$). After the emergence of the first pair

of true leaves, the seedlings were transferred to plastic pots (350mL) with the same substrate

described above, supplemented with 8 g L$^{-1}$ 10:10:10 nitrogen: phosphorus: potassium

(N:P:K) and 4 g L$^{-1}$ lime. The experiment was conducted in greenhouse localized at

Universidade Federal de Viçosa (642 m asl, 20$^o$45' S; 42$^o$51' W), Minas Gerais, Brazil, under

semi-controlled conditions (mean temperature of 28°C and 450-500 μmol m$^{-2}$ s$^{-1}$ PAR

irradiance). Irrigation was performed twice a day, where each plastic pot received the same

volume of water.

***Growth analyses***

Plant height and internode length were measured 35 days after germination (DAG). At 45

DAG the plants were harvest and divided into leaves, stem and root.  Leaf area was measured

using leaf area meter (Li-Cor Model 3100 Area Meter, Lincoln, NE, USA). Leaf, stem and

root were then packed separately in paper bags and oven dried at 70°C for 72h until they

reached constant weight. Shoot and root biomass were measured from the dry weight of

leaves, stem and root. Specific Leaf Area (SLA) was determined as described by Hunt,

(1982).

***Microscopy***

For anatomical analysis, leaf discs were collected from the medium point of the fifth lateral

leaflet and fixed in FAA50 (Formaldehyde, acetic acid and ethanol 50%) for 48 h, and then

stored in ethanol 70%. Samples were infiltrated with historesin (Leica Microsystems, Wetzlar,

Germany) and cut in cross sections with ~5 μm (RM2155, Leica Microsystems, Wetzlar,

Germany). Leaf cross sections were mounted in water on glass slides and stained with

toluidine blue. Histological sections were observed using an optic microscope (AX-70 TRF,

Olympus Optical, Tokyo, Japan) and then photographed using digital photo camera (Zeiss

AxioCam HRc, Göttinger, Germany). Anatomical features, as leaf thickness and thickness

cell layers, were analyzed using Image-Pro Plus® software (version 4.5, Media Cybernetics,

Silver Spring, USA).

***Measurements of Photosynthetic Parameters***

Photosynthetic parameters were performed using an open-flow infrared gas exchange

analyzer system (Li 6400XT, Li-Cor, Lincoln, USA) equipped with an integrated

fluorescence chamber (Li-6400-40; Li-Cor Inc.). All measurements were made on terminal

leaflets of intact and completely expanded leaves. The analysis was conducted under common

conditions for photon flux density (1000 μmol m-2 s -1), leaf temperature (25 ± 0.5ºC), leaf-

to-air vapor pressure difference (16.0 ± 3.0 mbar), air flow rate into the chamber (500 μmol s-

1), reference $CO_2$ concentration of 400 ppm using an area of 2 cm$^2$ in the leaf chamber.

$A$/Ci curves were determined initiated at an ambient $CO_2$ concentration of 400 μmol mol$^{-1}$

under a saturating photosynthetic Photon Flux Density (PPFD) of 1000 μmol m$^{-2}$ s $^{-1}$ at 25°C

under ambient $O_2$ supply. $CO_2$ concentration was decreased to 50 μmol mol$^{-1}$ of air in step

changes. Upon the completion of the measurements at low $C_a$, $C_a$ was returned to 400 μmol

mol$^{-1}$ of air to restore the original $A$. Next, $CO_2$ was increased stepwise to 1600 μmol mol$^{-1}$ of

air. The maximum rate of carboxylation ($V_{cmax}$), maximum rate of carboxylation limited by

electron transport ($J_{max}$) and triose-phosphate utilization (TPU) were estimated by fitting the

mechanistic model of $CO_2$ assimilation proposed by Farquhar *et al.*, (1980). Corrections for

the leakage of $CO_2$ into and out of the leaf chamber of the LI-6400 were applied to all gas-

exchange data as described by  Rodeghiero *et al.*, (2007).

***Kinetics of stomatal conductance***

The evaluation of stomatal behavior in response to light and $CO_2$ variation was performed in

WT and *ARF4*-as plants 40 DAG, under laboratory conditions (temperature 22-25°C; relative

humidity 50-60% and radiation of 150-200 µmol m$^{-2}$ s$^{-1}$), using an open-flow infrared gas

exchange analyzer system (Li 6400XT, Li-Cor, Lincoln, USA). The variation of $g_s$ was

performed on the fifth or sixth fully expanded leaf. For step-change evaluation of $g$s in

response to light intensity (Lawson *et al.*, 2010), leaves were stabilized for 30 min in the dark

and then radiation was increased to 1000 µmol m$^{-2}$ s$^{-1}$, and allowed to stabilized for 240 min.

Subsequently, light was turned off and observations recorded for another 100 min. The

alternation in the amount of light was based on the results presented by. The response of $g_s$ to

$CO_2$ was performed in a similar manner to those described above for light response. The

analysis was performed in a range of 240 minutes, with changes in $CO_2$ concentration (400-

800-400 µmol $CO_2$ m$^{-2}$ s$^{-1}$).

*Productivity traits*

The productivity parameters were measured in eight plants per genotype (WT and *ARF4*-as).

The average fruit weight was determined after individual weighing of each fruit, using a semi

analytical balance with a sensitivity of 0.01 g (Shimadzu® AUY220 model). Equatorial and

polar diameter was measured using a digital pachymeter (Jomarca *STAINLESS HARDENED*).

The determination of the soluble solids content (°Brix) was performed in 10 fruits per plant

using a digital temperature-compensated refractometer, model RTD 45 (Instrutherm®, São

Paulo, SP).

**Salt and drought stress analyses**

*Plant growth and stress conditions*

WT tomato seeds were sterilized for 10 min in 50% sodium hypochlorite, rinsed four times

with sterile distilled water and sown in pots containing peat. Then they were incubated in a

culture room with 16h light/ 8h dark photoperiod and 25± 2°C temperature. After 3 weeks,

plants were subjected to salt and drought stresses. Salt stress was performed by watering daily

the plants with 250mM of NaCl solution. Control plants were daily watered with distilled

water. Leaves and roots samples were harvested after 2 and 24 hours of salt stress application.

Drought stress was performed on three week-old plants by water holding for 48 hours and for

5 days. Watering continued normally throughout for control plants. Leave and root samples

were collected after drought stress application. Three biological replicates were done for each

condition.

***RNA extraction***

Total RNA was extracted from leaves and roots samples by using The Plant RNeasy

extraction kit (RNeasy Plant Mini Kit, Qiagen, Valencia, CA, USA). To remove any residual

genomic DNA, the RNA was treated with an RNase-Free DNase according to the

manufacturer's instruction (Ambion® DNA-*free*$^{TM}$DNase). The concentration of RNA was

accurately quantified by spectrophotometric measurement and 1μg of total RNA was

separated on 2% agarose gel to monitor its integrity. DNase-treated RNA (2μg) was then

reverse-transcribed in a total volume of 20μl using the Omniscript Reverse Transcription Kit

(Qiagen,).

***Real time PCR***

The real-time quantification of cDNA corresponding to 2μg of total RNA was performed in

the ABI PRISM 7900HT sequence detection system using the QuantiTech SYBR Green RT-

PCR kit (Qiagen). The Gene-specific primers used are listed in Supplementary Table 1. The

reaction mixture (10μl) contained 2μg of total RNA, 1,2 μM of each primer and appropriate

amounts of enzymes and fluorescent dyes as recommended by the manufacturer. Actin gene

was used as reference. Real-Time PCR conditions were as follow: 50°C for 2 min, 95°C for

10 min, then 40 cycles of 95°C for 15 s and 60°C for 1 min, and finally one cycle at 95°C for

15 s and 60°C for 15 s. Three independent biological replicates were used for real-time PCR

analysis. For each data point, the $C_T$ value was the average of $C_T$ values obtained from the

three biological replicates.

*Histochemical analysis of GUS expression*

Transgenic plants expressing pARF4::GUS were generated by *A. tumefaciens*-mediated

transformation according to H., Wang *et al.*, (2005). For that, PCR was performed on the

genomic DNA of tomato 'Micro-Tom' (10 ng.ml$^{-1}$) using specific primers.    The

corresponding amplified fragment was cloned into the pMDC162 vector (Curtis and

Grossniklaus, 2003)containing the GUS reporter gene using Gateway technology (Invitrogen).

The cloned *SlARF* promoter was sequenced from both sides using vector primers in order to

see whether the end of the promoter is matching with the beginning of the reporter gene.

Sequencing results analyses were carried out using the Vector NTI (Invitrogen) and

ContigExpress software by referring to ARF promoter sequences.

After being surface sterilized, pARF4::GUS seeds were cultivated in Petri dishes containing

half strength Murashige & Skoog medium for 7 days in a growth chamber at 25°C with 16h

light/ 8h dark cycle. One week plants were then grown hydroponically for two weeks in

Broughton & Dillworth (BD) liquid medium (Broughton and Dilworth, 1971). Three week-

old plants were subjected to salt and drought treatment. Salt stress was performed by adding

250 mM of NaCl to the culture medium. After 24 hours of the salt stress application, plants

were incubated overnight at 37°C in GUS buffer (3 mM 5-bromo-4-chloro-3-indolyl-β-D-

glucuronide [Duchefa Biochemie, Haarlem, The Netherlands], 0.1% v/v Triton X-100 [Sigma,

Steinhaim, Germany] 8 mM β-mercaptoethanol, 50 mM $Na_2HPO_4/NaH_2PO_4$ [pH 7.2]) then

followed by a destaining in 70% EtOH.. Drought stress was conducted by adding 15%  PEG

20000 to the liquid culture solution. Plants were collected after five days of stress application

and were GUS-stained as described above. For each stress condition, control plants were

cultivated in BD liquid medium for the same period.

*Stress tolerance assays in the transgenic tomato plants*

WT, *ARF4*-as and *ARF4*-ox seeds were first surface-sterilized for 10 min in 50% sodium

hypochlorite, rinsed four times with sterile distilled water. They were then cultured in petri

dishes containing half strength Murashige & Skoog (MS) medium for 7 days in a growth

chamber at 25° with 16h light/ 8h dark cycle. One week plants were then grown

hydroponically in pots containing 1L of BD medium  for two weeks  in the same growth

conditions (25° with 16h light/ 8h dark cycle) (Broughton and Dilworth, 1971). Three weeks

plants were then subjected to salt and drought stresses. Salt stress was performed by adding

100 mM or 150 mM NaCl to the liquid BD medium. Leaves and roots samples were harvested

after 2 weeks of treatment. Drought treatment was conducted as follows:  three weeks plants

were subjected to drought stress by adding 5% or 15% PEG 20 000 corresponding to final

osmotic potentials of -0,09 MPa and -0,28 MPa  to the hydroponic solution for 2 weeks.

Meanwhile, the control plants were grown normally in BD medium. Leaves and roots samples

were collected after 2 weeks of culture for both stressed and unstressed plants. For each

treatment, the hydroponic solution was renewed each 3 days.

***Determination of shoot and root fresh weights***

Leaf and root fresh weights were determined as followed: stressed and unstressed plants from

each line were harvested and rinsed thoroughly with distilled water (DW). The plants were

blot dried on blotting sheet and cut into shoots and roots. The fresh weight of each part of the

plant was measured and the mean of shoots and roots weights was determined based on at

least twelve plants per line in each condition.

***Determination of primary root length and lateral root density***

Primary root length was determined was determined as followed. Pictures for each plant were

analyzed using ImageJ software in order to determinate the root length. Twelve independent

plants from each line under control and stress conditions were used to calculate the mean.

The lateral root number was determined by counting the number of emerging roots and the

mean was calculated based on at least twelve plants per line in each condition. The lateral root

density (LR density) was determined using the following equation: number of LRs / the

length of the root.

### *Determination of chlorophyll content*

Determination of chlorophyll content of each line under stressful and normal conditions was

performed as described in Bassa *et al.*, (2012). A 100 mg aliquot of leaves collected from

stressed and unstressed plants was weighted and ground with 1 ml of 80% acetone. The liquid

obtained was centrifuged for 1 min at 10000 rpm to remove any remaining solid tissue.

Samples were analyzed by spectrophotometry at two wavelengths, 645 and 663 nm, using

80% acetone as the blank. The total chlorophyll content was determined using the following

equations: Total Chlorophyll Content $= 20.2 \times$ Chl a $+ 8.02 \times$ Chl b ( Chl a $= 0.999A663 -$

$0.0989A645$ and    Chl b $= -0.328A663 + 1.77A645$)

### *Determination of soluble sugar content*

Total soluble sugar content was determined as in Dubois *et al.*, (1956). 100 mg of fresh leaves

and roots of each sample were homogenized in 5 ml of 80 % ethanol in test tubes. Test tubes

were kept in water bath of 80 °C for 1 hour, and sample extracts were transferred to another

test tube, and 0.5 ml distilled water and 1 ml of 5 % phenol then added and allowed to

incubate for 1 hour. Finally after 1 hour, 2.5 ml sulphuric acid was added to the test tubes, and

shaken well on an orbital shaker. Absorbance was read at 485 nm on Spectrophotometer.

Sucrose was used as standard.

### *Determination of leaf stomatal conductance*

688 Leaf abaxial stomatal conductance of the youngest fully expanded leaf (usually the fifth leaf,

689 counting from the base) of stressed and well-drained plants of each studied line were

690 determined after two weeks of stress treatment using a hand-held leaf diffusion poromoter

691  (SC-1 LEAF POROMETER, Decagon, USA). For each line, three biological replicates were

692  conducted at each condition.

### Determination of ABA content

ABA measurement assays were performed as described (Forcat *et al.*, 2008). Briefly, 110 mg

of frozen tissue were extracted at 4°C for 30 min with 400 µl of $H_2O$ with 10% methanol +

1% acetic acid. The internal standard was 2H6 ABA. The extract was centrifuged at 13,000 g

for 10 min at 4°C. The supernatant was carefully removed and the pellet re-incubated for 30

min with 400 µl of methanol-acetic acid mix. Following the centrifugation, the supernatants

were pooled. Extracts were then analyzed by LC-MS using an Acquity UPLC coupled to a

XevoQtof (Waters, Massachusetts, USA). Analysis parameters were described in (Jaulneau *et*

*al.*, 2010).

### Determination of Relative water content (RWC)

703  Tomato fully expanded leaves were excised and fresh weight (FW) was recorded every 30

704  minutes. The excised leaves were then allowed to float on deionised water for about 24 h and

705  turgid weight (TW) was recorded. Leaves were dried at 80 °C for 24 h and dry weight (DW)

706  recorded. Finally, RWC was calculated according to Smart and Bingham, (1974).

### Quantitative expression assays

Three weeks WT, *ARF4*-as and *ARF4*-ox plants grown hydroponically in BD medium were

subjected to either 150 mM of NaCl or 15% of PEG. Leaves and root samples were harvested

after 2 hours and 24 hours for salt stress and after 48 hours for drought. RNA extraction,

cDNA synthesis and real time PCR were performed as previously described in paragraph

"ARF4 gene expression under salt and drought stress conditions". The Gene-specific primers

used are listed in Supplementary Table1.

**Acknowledgments**

This research was supported by Hubert-Curien partnership program Volubilis (MA/ 12/280-

27105PL) and benefited from networking activities within the European COST Action

FA1106. The authors are grateful to L. Lemonnier, D. Saint-Martin and O. Berseille for

genetic transformation and culture of tomato plants.

**Conflict of interest**

All Authors have read, edited and approved the final version of the manuscript. They have

declared that no competing interests exist.

**Supporting information**

**Figure S1. stomatal responses to irradiance and $CO_2$ levels in Micro-Tom (WT) and**

***ARF4*-as transgenic line.** Stomatal conductance ($g_s$) response in tomato plants cv. Micro-

Tom (WT) and isogenic *ARF4* antisense transgenic line (*ARF4*-as) from dark-adapted (0 µmol

$m^{-2}$ $s^{-1}$) leaves exposed to light (1000 µmol $m^{-2}$ $s^{-1}$) and on subsequent transfer back to

darkness (0 µmol $m^{-2}$ $s^{-1}$) in 350 minutes interval (n=5). Stomatal conductance in response to

$CO_2$ elevation and subsequent decrease to ambient $CO_2$ (400-800-400 µmol $CO_2$ $m^{-2}$ $s^{-1}$) in

240 minute interval (n=4). Measurements were performed using a LI-6400; LI-COR gas

exchange chamber in plants aged 40 days after germination. Data presented are mean ± SE

obtained using the 5th leaf totally expanded.

**Figure S2. Net $CO_2$ assimilation rate (*A*) and stomatal conductance ($g_s$) in Micro-Tom**

**(WT) and isogenic *ARF4* antisense transgenic line (*ARF4*-as).** (a) $CO_2$ assimilation rate

(*A*) as a function of stomatal conductance (*g*s), each point represents one measurement on an

individual plant. Line fitted by linear regression. (b) net $CO_2$ assimilation rate (*A*). (c)

stomatal conductance ($g_s$). (d) intrinsic water efficiency ($A/g_s$). Values are means ± s.e.m

(n=8). Asterisks indicate values that were determined by Student's *t* test to be significantly

different (P < 0.05) between genotypes.

**Figure S3. Productive parameters in tomato Micro-Tom (WT) and *ARF4*-as transgenic**

**line.** (a) and (b) representative tomato plants in reproductive stage. (c) representative fruits of

a single plant. (d) Number of fruit. (e) Fruit fresh weight. (f) and (g) equatorial and polar

diameter of the fruit, respectively Values are means $\pm$ s.e.m (n=8 plants).Asterisks indicate

values that were determined by Student's *t* test to be significantly different (P < 0.05) from

Micro-tom (WT). Scale bar = 1cm.

**Figure S4. Photosynthesis and sugar accumulation in WT, *ARF4*-as and *ARF4*-ox plant**

**lines exposed to different concentrations of NaCl or PEG.** (a) and (b) total chlorophyll

content in salt and drought stress conditions respectively, (c) and (d) leaf soluble sugars in salt

and drought stress conditions respectively, (e) and (f) root soluble sugars content in salt and

drought stress conditions respectively. Salt and drought stresses were performed on three

weeks tomato plants for two weeks by adding 100mM of NaCl or 150 mM of NaCl for salt

stress or 5% or 15% of PEG 20 000 for drought. Values are mean $\pm$ SD of three biological

replicates. Bars with different letters indicate the statistical significance (p<0,05) according to

Student Newman-Keuls test.

**Figure S5. Expression of sucrose transporter *LeSUT1* in WT, *ARF4*-as and *ARF4*-ox**

**plants exposed to salt or drought stresses.** (a) gene expression in leaves and roots  exposed

to 150mM of NaCl, (b) gene expression in leaves and roots leaves exposed to 15% PEG.

$\Delta\Delta$Ct refers to fold differences in gene expression relative to untreated plants. Values are

mean $\pm$ SD of three biological replicates. Stars (*) indicate the statistical significance (p<0,05)

according to Student's t-test.

**Figure S6. Expression of *Cat1*, *mdhar* and *SOD* in WT, *ARF4*-as and *ARF4*-ox plants**

**exposed to salt or drought stresses.** (a) gene expression in leaves and roots  exposed to

150mM of NaCl, (b) gene expression in leaves and roots leaves exposed to 15% PEG. $\Delta\Delta$Ct

refers to fold differences in gene expression relative to untreated plants. Values are mean ±

SD of three biological replicates. Stars (*) indicate the statistical significance (p<0,05) using

Student's t-test.

**Supplementary Table S1:** Gene ID and quantitative RT-PCR primers

**References**

**Aghaei, K., Ehsanpour, A.A. and Komatsu, S.** (2009) Potato responds to salt stress by
increased activity of antioxidant enzymes. *J. Integr. Plant Biol.*, **51**, 1095–1103.

**Ahmad, P., Ozturk, M., Sharma, S. and Gucel, S.** (2014) Effect of sodium carbonate-
induced salinity–alkalinity on some key osmoprotectants, protein profile, antioxidant
enzymes, and lipid peroxidation in two mulberry (Morus alba L.) cultivars. *J. Plant*
*Interact.*, **9**, 460–467.

**Akıncı, Ş. and Lösel, D.M.** (2012) Plant water-stress response mechanisms. In *Water stress*.
InTech.

**Al-Taweel, K., Iwaki, T., Yabuta, Y., Shigeoka, S., Murata, N. and Wadano, A.** (2007) A
Bacterial Transgene for Catalase Protects Translation of D1 Protein during Exposure
of Salt-Stressed Tobacco Leaves to Strong Light. *Plant Physiol.*, **145**, 258–265.

**Anjum, N.A., Umar, S. and Chan, M.-T.** (2010) *Ascorbate-glutathione pathway and stress*
*tolerance in plants*, Springer Science & Business Media.

**Anjum, S., Wang, L., Farooq, M., Hussain, M., Xue, L. and Zou, C.** (2011) Brassinolide
application improves the drought tolerance in maize through modulation of enzymatic
antioxidants and leaf gas exchange. *J. Agron. Crop Sci.*, **197**, 177–185.

**Ashraf, M. and Harris, P.** (2013) Photosynthesis under stressful environments: an overview.
*Photosynthetica*, **51**, 163–190.

**Atkinson, N.J. and Urwin, P.E.** (2012) The interaction of plant biotic and abiotic stresses:
from genes to the field. *J. Exp. Bot.*, ers100.

**Badawi, G.H., Yamauchi, Y., Shimada, E., Sasaki, R., Kawano, N., Tanaka, K. and**
**Tanaka, K.** (2004) Enhanced tolerance to salt stress and water deficit by
overexpressing superoxide dismutase in tobacco (Nicotiana tabacum) chloroplasts.
*Plant Sci.*, **166**, 919–928.

**Balsamo, R., Willigen, C. Vander, Bauer, A. and Farrant, J.** (2006) Drought tolerance of
selected Eragrostis species correlates with leaf tensile properties. *Ann. Bot.*, **97**, 985–
991.

**Bassa, C., Mila, I., Bouzayen, M. and Audran-Delalande, C.** (2012) Phenotypes associated
with down-regulation of Sl-IAA27 support functional diversity among Aux/IAA
family members in the tomato. *Plant Cell Physiol.*, pcs101.

**Bernacchi, C.J., Rosenthal, D.M., Pimentel, C., Long, S.P. and Farquhar, G.D.** (2009) Modeling the temperature dependence of C 3 photosynthesis. In *Photosynthesis in silico*. Springer, pp. 231–246.

**Blum, A.** (2005) Drought resistance, water-use efficiency, and yield potential - are they compatible, dissonant, or mutually exclusive? *Aust. J. Agric. Res.*, **56**, 1159–1168.

**Blum, A.** (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. *Field Crops Res.*, **112**, 119–123.

**Bouzroud, S., Gouiaa, S., Hu, N., Bernadac, A., Mila, I, Bendaou, N., Smouni, A., Bouzayen, M. and Zouine, M.** (2018) Auxin Response Factors (ARFs) are potential mediators of auxin action in tomato response to biotic and abiotic stress (Solanum lycopersicum). *PLOS ONE*, **13**, e0193517.

**Breitel, D.A., Chappell-Maor, L., Meir, S., et al.** (2016) AUXIN RESPONSE FACTOR 2 Intersects Hormonal Signals in the Regulation of Tomato Fruit Ripening G. P. Copenhaver, ed. *PLOS Genet.*, **12**, e1005903.

**Broughton, W.J. and Dilworth, M.J.** (1971) Control of leghaemoglobin synthesis in snake beans. *Biochem. J.*, **125**, 1075–1080.

**Brugnoli, E. and Lauteri, M.** (1991) Effects of Salinity on Stomatal Conductance, Photosynthetic Capacity, and Carbon Isotope Discrimination of Salt-Tolerant (Gossypium hirsutum L.) and Salt-Sensitive (Phaseolus vulgaris L.) C3 Non-Halophytes. *Plant Physiol.*, **95**, 628–635.

**Burssens, S., Himanen, K., Cotte, B. van de, Beeckman, T., Montagu, M. Van, Inze, D. and Verbruggen, N.** (2000) Expression of cell cycle regulatory genes and morphological alterations in response to salt stress in Arabidopsis thaliana. *Planta*, **211**, 632–640.

**Chaabouni, S., Jones, B., Delalande, C., et al.** (2009) Sl-IAA3, a tomato Aux/IAA at the crossroads of auxin and ethylene signalling involved in differential growth. *J. Exp. Bot.*, **60**, 1349–1362.

**Chandler, J.W.** (2016) Auxin response factors. *Plant Cell Environ.*, **39**, 1014–1028.

**Chang-Quan, W. and Rui-Chang, L.** (2008) Enhancement of superoxide dismutase activity in the leaves of white clover (Trifolium repens L.) in response to polyethylene glycol-induced water stress. *Acta Physiol. Plant.*, **30**, 841.

**Chaves, M.M.** (1991) Effects of Water Deficits on Carbon Assimilation. *J. Exp. Bot.*, **42**, 1–16.

**Chernys, J.T. and Zeevaart, J.A.** (2000) Characterization of the 9-cis-epoxycarotenoid dioxygenase gene family and the regulation of abscisic acid biosynthesis in avocado. *Plant Physiol.*, **124**, 343–354.

**Choudhury, S., Panda, P., Sahoo, L. and Panda, S.K.** (2013) Reactive oxygen species signaling in plants under abiotic stress. *Plant Signal. Behav.*, **8**, e23681.

**Comas, L.H., Becker, S.R., Cruz, V.M. V., Byrne, P.F. and Dierig, D.A.** (2013) Root traits contributing to plant productivity under drought. *Front. Plant Sci.*, **4**.

**Curtis, M.D. and Grossniklaus, U.** (2003) A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiol.*, **133**, 462–9.

**Damour, G., Simonneau, T., Cochard, H. and Urban, L.** (2010) An overview of models of stomatal conductance at the leaf level. *Plant Cell Environ.*, **33**, 1419–38.

**Das, K. and Roychoudhury, A.** (2014) Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Environ. Toxicol.*, **2**, 53.

**De Jong, M., Mariani, C. and Vriezen, W.H.** (2009) The role of auxin and gibberellin in tomato fruit set. *J. Exp. Bot.*, **60**, 1523–1532.

**Dogan, M., Tipirdamaz, R. and Demir, Y.** (2010) Salt resistance of tomato species grown in sand culture. *Plant Soil Environ. - UZEI Czech Repub.*

**Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P. t and Smith, F.** (1956) Colorimetric method for determination of sugars and related substances. *Anal. Chem.*, **28**, 350–356.

**Eltayeb, A.E., Kawano, N., Badawi, G.H., Kaminaka, H., Sanekata, T., Shibahara, T., Inanaga, S. and Tanaka, K.** (2006) Overexpression of monodehydroascorbate reductase in transgenic tobacco confers enhanced tolerance to ozone, salt and polyethylene glycol stresses. *Planta*, **225**, 1255–1264.

**Farooq, M., Wahid, A., Kobayashi, N., Fujita, D. and Basra, S.** (2009) Plant drought stress: effects, mechanisms and management. In *Sustainable agriculture*. Springer, pp. 153–188.

**Farquhar, G.D., Caemmerer, S. von and Berry, J.A.** (1980) A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. *Planta*, **149**, 78–90.

**Fayez, K.A. and Bazaid, S.A.** (2014) Improving drought and salinity tolerance in barley by application of salicylic acid and potassium nitrate. *J. Saudi Soc. Agric. Sci.*, **13**, 45–55.

**Fen, L.L., Ismail, M.R., Zulkarami, B., Rahman, M.S.A. and Islam, M.R.** (2015) Physiological and molecular characterization of drought responses and screening of drought tolerant rice varieties. *Biosci. J.*, **31**.

**Flowers, T. and Yeo, A.** (1981) Variability in the resistance of sodium chloride salinity within rice (Oryza sativa L.) varieties. *New Phytol.*, **88**, 363–373.

**Forcat, S., Bennett, M.H., Mansfield, J.W. and Grant, M.R.** (2008) A rapid and robust method for simultaneously measuring changes in the phytohormones ABA, JA and SA in plants following biotic and abiotic stress. *Plant Methods*, **4**, 16.

**Furdi, F., Velicevici, G., Petolescu, C. and Popescu, S.** (2013) The effect of salt stress on chlorophyll content in several Romanian tomato varieties. *J. Hortic. For. Biotechnol.*, **17**, 363–367.

**Gapińska, M., Skłodowska, M. and Gabara, B.** (2007) Effect of short- and long-term
salinity on the activities of antioxidative enzymes and lipid peroxidation in tomato
roots. *Acta Physiol. Plant.*, **30**, 11–18.

**Goetz, M., Vivian-Smith, A., Johnson, S.D. and Koltunow, A.M.** (2006) AUXIN
RESPONSE FACTOR8 Is a Negative Regulator of Fruit Initiation in Arabidopsis.
*Plant Cell Online*, **18**, 1873–1886.

**Gondim, F.A., Gomes-Filho, E., Costa, J.H., Mendes Alencar, N.L. and Prisco, J.T.**
(2012) Catalase plays a key role in salt stress acclimation induced by hydrogen
peroxide pretreatment in maize. *Plant Physiol. Biochem. PPB Société Fr. Physiol.*
*Végétale*, **56**, 62–71.

**Guilfoyle, T.J. and Hagen, G.** (2007) Auxin response factors. *Curr. Opin. Plant Biol.*, **10**,
453–460.

**Guilfoyle, T.J. and Hagen, G.** (2007) Auxin response factors. *Curr. Opin. Plant Biol.*, **10**,
453–460.

**Hao, Y., Hu, G., Breitel, D., et al.** (2015) Auxin Response Factor SlARF2 Is an Essential
Component of the Regulatory Mechanism Controlling Fruit Ripening in Tomato G. P.
Copenhaver, ed. *PLOS Genet.*, **11**, e1005649.

**Harper, R.M., Stowe-Evans, E.L., Luesse, D.R., Muto, H., Tatematsu, K., Watahiki,**
**M.K., Yamamoto, K. and Liscum, E.** (2000) The NPH4 locus encodes the auxin
response factor ARF7, a conditional regulator of differential growth in aerial
Arabidopsis tissue. *Plant Cell*, **12**, 757–770.

**Hendelman, A., Buxdorf, K., Stav, R., Kravchik, M. and Arazi, T.** (2012) Inhibition of
lamina outgrowth following Solanum lycopersicum AUXIN RESPONSE FACTOR
10 (SlARF10) derepression. *Plant Mol. Biol.*, **78**, 561–576.

**Hernandez, J., Jimenez, A., Mullineaux, P. and Sevilia, F.** (2000) Tolerance of pea (Pisum
sativum L.) to long‐term salt stress is associated with induction of antioxidant
defences. *Plant Cell Environ.*, **23**, 853–862.

**Hernández, J.A., Corpas, F.J., Gómez, M., Río, L.A. del and Sevilla, F.** (1993) Salt-
induced oxidative stress mediated by activated oxygen species in pea leaf
mitochondria. *Physiol. Plant.*, **89**, 103–110.

**Heschel, M.S. and Riginos, C.** (2005) Mechanisms of selection for drought stress tolerance
and avoidance in Impatiens capensis (Balsaminaceae). *Am. J. Bot.*, **92**, 37–44.

**Hu, W., Zuo, J., Hou, X., Yan, Y., Wei, Y., Liu, J., Li, M., Xu, B. and Jin, Z.** (2015) The
auxin response factor gene family in banana: genome-wide identification and
expression analyses during development, ripening, and abiotic stress. *Crop Sci.*
*Hortic.*, 742.

**Hunt, R.** (1982) Plant Growth Analysis: Second derivatives and compounded second
derivatives of splined plant growth curves. *Ann. Bot.*, **50**, 317–328.

**Iuchi, S., Kobayashi, M., Taji, T., et al.** (2001) Regulation of drought tolerance by gene manipulation of 9⬚cis⬚epoxycarotenoid dioxygenase, a key enzyme in abscisic acid biosynthesis in Arabidopsis. *Plant J.*, **27**, 325–333.

**Iuchi, S., Kobayashi, M., Yamaguchi-Shinozaki, K. and Shinozaki, K.** (2000) A stress-inducible gene for 9-cis-epoxycarotenoid dioxygenase involved in abscisic acid biosynthesis under water stress in drought-tolerant cowpea. *Plant Physiol.*, **123**, 553–562.

**Jain, M. and Khurana, J.P.** (2009) Transcript profiling reveals diverse roles of auxin-responsive genes during reproductive development and abiotic stress in rice. *FEBS J.*, **276**, 3148–3162.

**Jamil, M., Rehman, S. and Rha, E.** (2007) Salinity effect on plant growth, PSII photochemistry and chlorophyll content in sugar beet (Beta Vulgaris L.) and cabbage (Brassica Oleracea Capitata L.). *Pak J Bot*, **39**, 753–760.

**Jaulneau, V., Lafitte, C., Jacquet, C., Fournier, S., Salamagne, S., Briand, X., Esquerré-Tugayé, M.-T. and Dumas, B.** (2010) Ulvan, a Sulfated Polysaccharide from Green Algae, Activates Plant Immunity through the Jasmonic Acid Signaling Pathway. *J. Biomed. Biotechnol.*, **2010**.

**Jia, W., Zhang, L., Wu, D., et al.** (2015) Sucrose transporter AtSUC9 mediated by a low sucrose level is involved in Arabidopsis abiotic stress resistance by regulating sucrose distribution and ABA accumulation. *Plant Cell Physiol.*, **56**, 1574–1587.

**Jiang, Q., Roche, D., Monaco, T. and Hole, D.** (2006) Stomatal conductance is a key parameter to assess limitations to photosynthesis and growth potential in barley genotypes. *Plant Biol.*, **8**, 515–521.

**Jones, B., Frasse, P., Olmos, E., Zegzouti, H., Li, Z.G., Latché, A., Pech, J.C. and Bouzayen, M.** (2002) Down-regulation of DR12, an auxin-response-factor homolog, in the tomato results in a pleiotropic phenotype including dark green and blotchy ripening fruit. *Plant J.*, **32**, 603–613.

**Kadioglu, A. and Terzi, R.** (2007) A Dehydration Avoidance Mechanism: Leaf Rolling. *Bot. Rev.*, **73**, 290–302.

**Kadioglu, A., Terzi, R., Saruhan, N. and Saglam, A.** (2012) Current advances in the investigation of leaf rolling caused by biotic and abiotic stress factors. *Plant Sci.*, **182**, 42–8.

**Keunen, E., Peshev, D., Vangronsveld, J., Ende, W. Van Den and Cuypers, A.** (2013) Plant sugars are crucial players in the oxidative challenge during abiotic stress: extending the traditional concept. *Plant Cell Environ.*, **36**, 1242–1255.

**Kühn, C., Franceschi, V.R., Schulz, A., Lemoine, R. and Frommer, W.B.** (1997) Macromolecular trafficking indicated by localization and turnover of sucrose transporters in enucleate sieve elements. *Science*, **275**, 1298–1300.

**Kukreja, S., Nandwal, A.S., Kumar, N., Sharma, S.K., Sharma, S.K., Unvi, V. and Sharma, P.K.** (2005) Plant water status, H2O2 scavenging enzymes, ethylene

evolution and membrane integrity of Cicer arietinum roots as affected by salinity. *Biol. Plant.*, **49**, 305–308.

Kurth, E., Cramer, G.R., Läuchli, A. and Epstein, E. (1986) Effects of NaCl and CaCl2 on Cell Enlargement and Cell Production in Cotton Roots. *Plant Physiol.*, **82**, 1102–1106.

Lawson, T., Caemmerer, S. von and Baroli, I. (2010) Photosynthesis and Stomatal Behaviour. In Springer Berlin Heidelberg, pp. 265–304.

Liscum, E. and Reed, J. (2002) Genetics of Aux/IAA and ARF action in plant growth and development. *Plant Mol. Biol.*, **49**, 387–400.

Liu, N., Wu, S., Houten, J. Van, Wang, Y., Ding, B., Fei, Z., Clarke, T.H., Reed, J.W. and Knaap, E. Van Der (2014) Down-regulation of AUXIN RESPONSE FACTORS 6 and 8 by microRNA 167 leads to floral development defects and female sterility in tomato. *J. Exp. Bot.*, **65**, 2507–2520.

Liu, S., Lv, Y., Wan, X.-R., Li, L.-M., Hu, B. and Li, L. (2014) Cloning and Expression Analysis of cDNAs Encoding ABA 8'-Hydroxylase in Peanut Plants in Response to Osmotic Stress. *PLOS ONE*, **9**, e97025.

Luna, C.M., Pastori, G.M., Driscoll, S., Groten, K., Bernard, S. and Foyer, C.H. (2005) Drought controls on H2O2 accumulation, catalase (CAT) activity and CAT gene expression in wheat. *J. Exp. Bot.*, **56**, 417–423.

Mafakheri, A., Siosemardeh, A., Bahramnejad, B., Struik, P. and Sohrabi, Y. (2010) Effect of drought stress on yield, proline and chlorophyll contents in three chickpea cultivars. *Aust. J. Crop Sci.*, **4**, 580.

Marcińska, I., Czyczyło-Mysza, I., Skrzypek, E., et al. (2012) Impact of osmotic stress on physiological and biochemical characteristics in drought-susceptible and drought-resistant wheat genotypes. *Acta Physiol. Plant.*, **35**, 451–461.

Marin, E., Jouannet, V., Herz, A., Lokerse, A.S., Weijers, D., Vaucheret, H., Nussaume, L., Crespi, M.D. and Maizel, A. (2010) miR390, Arabidopsis TAS3 tasiRNAs, and their AUXIN RESPONSE FACTOR targets define an autoregulatory network quantitatively regulating lateral root growth. *Plant Cell*, **22**, 1104–1117.

Medrano, H., Escalona, J.M., Bota, J., Gulías, J. and Flexas, J. (2002) Regulation of Photosynthesis of C3 Plants in Response to Progressive Drought: Stomatal Conductance as a Reference Parameter. *Ann. Bot.*, **89**, 895–905.

Mei, L., Yuan, Y., Wu, M., et al. (2018) SlARF10, an auxin response factor, is required for chlorophyll and sugar accumulation during tomato fruit development. *bioRxiv*, 253237.

M'Hamdi, M., du Jardin, P., Bettaieb, T., Harbaoui, Y. and Aziz Mougou, A. (2009) Insight into the role of catalases in salt stress in potato (Solanum tuberosum L.). *Base*.

Miyashita, K., Tanakamaru, S., Maitani, T. and Kimura, K. (2005) Recovery responses of photosynthesis, transpiration, and stomatal conductance in kidney bean following drought stress. *Environ. Exp. Bot.*, **53**, 205–214.

**Mohamed, E.-A., Iwaki, T., Munir, I., Tamoi, M., Shigeoka, S. and Wadano, A.** (2003) Overexpression of bacterial catalase in tomato leaf chloroplasts enhances photo-oxidative stress tolerance. *Plant Cell Environ.*, **26**, 2037–2046.

**Moriwaki, T., Yamamoto, Y., Aida, T., Funahashi, T., Shishido, T., Asada, M., Prodhan, S.H., Komamine, A. and Motohashi, T.** (2008) Overexpression of the Escherichia coli catalase gene, katE, enhances tolerance to salinity stress in the transgenic indica rice cultivar, BR5. *Plant Biotechnol. Rep.*, **2**, 41–46.

**Munns, R., James, R.A. and Läuchli, A.** (2006) Approaches to increasing the salt tolerance of wheat and other cereals. *J. Exp. Bot.*, **57**, 1025–1043.

**Narise, T., Kobayashi, K., Baba, S., Shimojima, M., Masuda, S., Fukaki, H. and Ohta, H.** (2010) Involvement of auxin signaling mediated by IAA14 and ARF7/19 in membrane lipid remodeling during phosphate starvation. *Plant Mol. Biol.*, **72**, 533–544.

**Navarro, L., Dunoyer, P., Jay, F., Arnold, B., Dharmasiri, N., Estelle, M., Voinnet, O. and Jones, J.D.** (2006) A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science*, **312**, 436–439.

**Noiraud, N., Delrot, S. and Lemoine, R.** (2000) The sucrose transporter of celery. Identification and expression during salt stress. *Plant Physiol.*, **122**, 1447–1455.

**Park, J.-E., Park, J.-Y., Kim, Y.-S., et al.** (2007) GH3-mediated auxin homeostasis links growth regulation with stress adaptation response in Arabidopsis. *J. Biol. Chem.*, **282**, 10036–10046.

**Patade, V.Y., Bhargava, S. and Suprasanna, P.** (2012) Transcript expression profiling of stress responsive genes in response to short-term salt or PEG stress in sugarcane leaves. *Mol. Biol. Rep.*, **39**, 3311–3318.

**Pattanagul, W. and Thitisaksakul, M.** (2008) Effect of salinity stress on growth and carbohydrate metabolism in three rice (Oryza sativa L.) cultivars differing in salinity tolerance. *Indian J. Exp. Biol.*, **46**, 736–742.

**Pekker, I., Alvarez, J.P. and Eshed, Y.** (2005) Auxin response factors mediate Arabidopsis organ asymmetry via modulation of KANADI activity. *Plant Cell*, **17**, 2899–2910.

**Pereira-Netto, A.B. de, Gabriele, A.C. and Pinto, H.S.** (1999) Aspects of leaf anatomy of kudzu (Pueraria lobata, Leguminosae-Faboideae) related to water and energy balance. *Pesqui. Agropecu. Bras.*, **34**, 1361–1365.

**Perl, A., Perl-Treves, R., Galili, S., Aviv, D., Shalgi, E., Malkin, S. and Galun, E.** (1993) Enhanced oxidative-stress defense in transgenic potato expressing tomato Cu,Zn superoxide dismutases. *Theor. Appl. Genet.*, **85**, 568–576.

**Rahman, S.L., Mackay, W.A., Nawata, E., Sakuratani, T., Uddin, A.M. and Quebedeaux, B.** (2004) Superoxide dismutase and stress tolerance of four tomato cultivars. *Hortscience*, **39**, 983–986.

**Rahnama, H., Vakilian, H., Fahimi, H. and Ghareyazie, B.** (2011) Enhanced salt stress
tolerance in transgenic potato plants (Solanum tuberosum L.) expressing a bacterial
mtlD gene. *Acta Physiol. Plant.*, **33**, 1521–1532.

**Reddy, A.R., Chaitanya, K.V. and Vivekanandan, M.** (2004) Drought-induced responses
of photosynthesis and antioxidant metabolism in higher plants. *J. Plant Physiol.*, **161**,
1189–1202.

**Reed, J.W.** (2001) Roles and activities of Aux/IAA proteins in Arabidopsis. *Trends Plant*
*Sci.*, **6**, 420–425.

**Rhizopoulou, S. and Psaras, G.K.** (2003) Development and Structure of Drought☐tolerant
Leaves of the Mediterranean Shrub Capparis spinosa L. *Ann. Bot.*, **92**, 377–383.

**Rodeghiero, M., Niinemets, Ü. and Cescatti, A.** (2007) Major diffusion leaks of clamp-on
leaf cuvettes still unaccounted: How erroneous are the estimates of Farquhar et al.
model parameters? *Plant Cell Environ.*, **30**, 1006–1022.

**Rosa, M., Prado, C., Podazza, G., Interdonato, R., González, J.A., Hilal, M. and Prado,**
**F.E.** (2009) Soluble sugars: Metabolism, sensing and abiotic stress: A complex
network in the life of plants. *Plant Signal. Behav.*, **4**, 388–393.

**Roy, B. and Basu, A.K.** (2009) *Abiotic Stress Tolerance in Crop Plants: Breeding and*
*Biotechnology*, New India Publishing.

**Saddam, S., Bibi, A., Sadaqat, H. and Usman, B.** (2014) Comparison of 10 sorghum
(Sorghum bicolor L.) genotypes under various water stress regimes. *J Anim Plant Sci*,
**24**, 1811–1820.

**Saeidi, M. and Abdoli, M.** (2015) Effect of Drought Stress during Grain Filling on Yield and
Its Components, Gas Exchange Variables, and Some Physiological Traits of Wheat
Cultivars. *J. Agric. Sci. Technol.*, **17**, 885–898.

**Sagar, M., Chervin, C., Mila, I., et al.** (2013) SlARF4, an auxin response factor involved in
the control of sugar metabolism during tomato fruit development. *Plant Physiol.*, **161**,
1362–74.

**Saito, S., Hirai, N., Matsumoto, C., Ohigashi, H., Ohta, D., Sakata, K. and Mizutani, M.**
(2004) Arabidopsis CYP707As encode (+)-abscisic acid 8′-hydroxylase, a key enzyme
in the oxidative catabolism of abscisic acid. *Plant Physiol.*, **134**, 1439–1449.

**Sami, F., Yusuf, M., Faizan, M., Faraz, A. and Hayat, S.** (2016) Role of sugars under
abiotic stress. *Plant Physiol. Biochem.*, **109**, 54–61.

**Schruff, M.C., Spielman, M., Tiwari, S., Adams, S., Fenby, N. and Scott, R.J.** (2005) The
AUXIN RESPONSE FACTOR 2 gene of Arabidopsis links auxin signalling, cell
division, and the size of seeds and other organs. *Development*, **133**.

**Schultz, H.R.** (2003) Differences in hydraulic architecture account for near-isohydric and
anisohydric behaviour of two field-grown Vitis vinifera L. cultivars during drought.
*Plant Cell Environ.*, **26**, 1393–1405.

**Sharma, P. and Shanker Dubey, R.** (2005) Modulation of nitrate reductase activity in rice seedlings under aluminium toxicity and water stress: role of osmolytes as enzyme protectant. *J. Plant Physiol.*, **162**, 854–864.

**Siddique, M., Hamid, A. and Islam, M.** (1999) Drought stress effects on photosynthetic rate and leaf gas exchange of wheat. *Bot. Bull. Acad. Sin.*, **40**.

**Smart, R.E. and Bingham, G.E.** (1974) Rapid Estimates of Relative Water Content. *Plant Physiol.*, **53**, 258–260.

**Su, L., Bassa, C., Audran, C., Mila, I., Cheniclet, C., Chevalier, C., Bouzayen, M., Roustan, J.-P. and Chervin, C.** (2014) The auxin Sl-IAA17 transcriptional repressor controls fruit size via the regulation of endoreduplication-related cell expansion. *Plant Cell Physiol.*, **55**, 1969–1976.

**Sui, N., Yang, Z., Liu, M. and Wang, B.** (2015) Identification and transcriptomic profiling of genes involved in increasing sugar content during salt stress in sweet sorghum leaves. *BMC Genomics*, **16**.

**Szemenyei, H., Hannon, M. and Long, J.A.** (2008) TOPLESS mediates auxin-dependent transcriptional repression during Arabidopsis embryogenesis. *Science*, **319**, 1384–1386.

**Szőllősi, R.** (2014) Superoxide Dismutase (SOD) and Abiotic Stress Tolerance in Plants: An Overview. In *Oxidative Damage to Plants*. Elsevier, pp. 89–129.

**Tan, B.C., Schwartz, S.H., Zeevaart, J.A. and McCarty, D.R.** (1997) Genetic control of abscisic acid biosynthesis in maize. *Proc. Natl. Acad. Sci.*, **94**, 12235–12240.

**Thompson, A.J., Jackson, A.C., Symonds, R.C., Mulholland, B.J., Dadswell, A.R., Blake, P.S., Burbidge, A. and Taylor, I.B.** (2000) Ectopic expression of a tomato 9‐cis‐epoxycarotenoid dioxygenase gene causes over‐production of abscisic acid. *Plant J.*, **23**, 363–374.

**Tombesi, S., Nardini, A., Frioni, T., Soccolini, M., Zadra, C., Farinelli, D., Poni, S. and Palliotti, A.** (2015) Stomatal closure is induced by hydraulic signals and maintained by ABA in drought-stressed grapevine. *Sci. Rep.*, **5**, 12449.

**Ulmasov, T., Hagen, G. and Guilfoyle, T.J.** (1999) Activation and repression of transcription by auxin-response factors. *Proc. Natl. Acad. Sci.*, **96**, 5844–5849.

**Ulmasov, T., Hagen, G. and Guilfoyle, T.J.** (1997) ARF1, a Transcription Factor That Binds to Auxin Response Elements. *Science*, **276**.

**Umezawa, T., Okamoto, M., Kushiro, T., et al.** (2006) CYP707A3, a major ABA 8′‐hydroxylase involved in dehydration and rehydration response in Arabidopsis thaliana. *Plant J.*, **46**, 171–182.

**Van Ha, C., Le, D.T., Nishiyama, R., et al.** (2013) The auxin response factor transcription factor family in soybean: genome-wide identification and expression analyses during development and water stress. *DNA Res.*, **20**, 511–524.

**Vanneste, S. and Friml, J.** (2009) Auxin: A Trigger for Change in Plant Development. *Cell*, **136**, 1005–1016.

**Volkenburgh, E.V.** (1999) Leaf expansion–an integrating plant behaviour. *Plant Cell Environ.*, **22**, 1463–1473.

**Wan, X.-R. and Li, L.** (2006) Regulation of ABA level and water-stress tolerance of Arabidopsis by ectopic expression of a peanut 9-cis-epoxycarotenoid dioxygenase gene. *Biochem. Biophys. Res. Commun.*, **347**, 1030–1038.

**Wang, F.-Z., Wang, Q.-B., Kwon, S.-Y., Kwak, S.-S. and Su, W.-A.** (2005) Enhanced drought tolerance of transgenic rice plants expressing a pea manganese superoxide dismutase. *J. Plant Physiol.*, **162**, 465–472.

**Wang, H., Jones, B., Li, Z., et al.** (2005) The tomato Aux/IAA transcription factor IAA9 is involved in fruit development and leaf morphogenesis. *Plant Cell*, **17**, 2676–2692.

**Wang, S., Bai, Y., Shen, C., Wu, Y., Zhang, S., Jiang, D., Guilfoyle, T.J., Chen, M. and Qi, Y.** (2010) Auxin-related gene families in abiotic stress response in Sorghum bicolor. *Funct. Integr. Genomics*, **10**, 533–546.

**Wang, Y., Ying, Y., Chen, J. and Wang, X.** (2004) Transgenic Arabidopsis overexpressing Mn-SOD enhanced salt-tolerance. *Plant Sci.*, **167**, 671–677.

**Wong, S., Cowan, I. and Farquhar, G.** (1979) Stomatal conductance correlates with photosynthetic capacity. *Nature*, **282**, 424.

**Yang, R., Yang, T., Zhang, H., et al.** (2014) Hormone profiling and transcription analysis reveal a major role of ABA in tomato salt tolerance. *Plant Physiol. Biochem.*, **77**, 23–34.

**Yang, S.H. and Zeevaart, J.A.** (2006) Expression of ABA 8′□hydroxylases in relation to leaf water relations and seed development in bean. *Plant J.*, **47**, 675–686.

**Yang, Y., Han, C., Liu, Q., Lin, B. and Wang, J.** (2008) Effect of drought and low light on growth and enzymatic antioxidant system of Picea asperata seedlings. *Acta Physiol. Plant.*, **30**, 433–440.

**Yong, B., Wang, X., Xu, P., et al.** (2017) Isolation and Abiotic Stress Resistance Analyses of a Catalase Gene from Ipomoea batatas (L.) Lam. *BioMed Res. Int.*, **2017**.

**Zarafshar, M., Akbarinia, M., Askari, H., Mohsen Hosseini, S., Rahaie, M., Struve, D. and Gabriel Striker, G.** (2014) Morphological, physiological and biochemical responses to soil water deficit in seedlings of three populations of wild pear tree (Pyrus boisseriana). *Base*.

**Zhang, X., Yan, F., Tang, Y., Yuan, Y., Deng, W. and Li, Z.** (2015) Auxin response gene SlARF3 plays multiple roles in tomato development and is involved in the formation of epidermal cells and trichomes. *Plant Cell Physiol.*, **56**, 2110–2124.

**Zhang, Y., Wang, Z., Wu, Y. and Zhang, X.** (2006) Stomatal characteristics of different green organs in wheat under different irrigation regimes. *Zuo Wu Xue Bao*, **32**, 70–75.

**Zlatev, Z., Lidon, F., Ramalho, J. and Yordanov, I.** (2006) Comparison of resistance to drought of three bean cultivars. *Biol. Plant.*, **50**, 389–394.

**Zouine, M., Fu, Y., Chateigner-Boutin, A.-L.L., Mila, I., Frasse, P., Wang, H., Audran, C., Roustan, J.-P.P. and Bouzayen, M.** (2014) Characterization of the tomato ARF gene family uncovers a multi-levels post-transcriptional regulation including alternative splicing. *PLoS ONE*, **9**, e84203.

**Tables**

**Table 1**. **Characterization of photosynthetic parameters in tomato cv. Micro-Tom (WT) and isogenic *ARF4* antisense transgenic line (*ARF4*-as).** A: photosynthetic $CO_2$ assimilation rate; gs: stomatal conductance; $V_{cmax}$: *maximum Rubisco carboxylation rate* ; $J_{max}$: maximum electron transport rate; TPU: triose phosphate utilization.. Mean values (*n=5*) ± s.e.m. Significant differences by *t*-test at *0.01 and **0.001

|  | **WT** | ***ARF4*-as** |
|---|---|---|
| *A* (**µmol CO$_2$ m$^{-2}$ s$^{-1}$**) | 18.30 ± 0.81 | 11.59** ± 0.59 |
| *g*$_s$ (**µmol m$^{-2}$ s$^{-1}$**) | 0.18 ± 0.02 | 0.09** ± 0.01 |
| *A*/*g*$_s$ | 101.52 ± 4.62 | 122.76* ± 3.16 |
| *V*$_{cmax}$ (**µmol m$^{-2}$ s$^{-1}$**) | 80.37 ± 2.34 | 70.43* ± 3.01 |
| *J*$_{max}$ (**µmol m$^{-2}$ s$^{-1}$**) | 156.40 ± 7.74 | 140.70* ± 6.30 |
| **TPU (µmol m$^{-2}$ s$^{-1}$)** | 11.84 ± 0.57 | 10.82 ± 0.53 |

1160 **Table 2: ABA content in WT and ARF4-as plants in response to salt or drought stress**
1161 **conditions.** Values presented are mean ± SD of three biological replicates (except for data
1162 related to *ARF4*-as plants in drought stress conditions, where only two biological replicates
1163 were done).

1164

| | Salt stress | | Drought stress | |
|---|---|---|---|---|
| | 0 mM NaCl | 150 mM NaCl | 0% PEG | 15% PEG |
| **WT** | 0,1272±0 | 0,1454±0,007 | 0,0681±0,007 | 0,05±0,007 |
| ***ARF4*-as** | 0,1151±0,015 | 0,2636±0,02 | 0,0636±0,014 | 0,1272±0,012 |

1165

1166

1167

**Article figures**

**Figure 1. Phenotype of cv. Micro-Tom tomato plants (wild-type, WT) and isogenic *ARF4***

**antisense transgenic line (*ARF4*-as), 35 days after germination**. (a) WT and *ARF4*-as

plants (b) at the same stage of development. (c) Upward-curled leaf phenotype of *ARF4*-as

compared to WT (d) number of leaves to first inflorescence, (e) plant height and (f) plant

internode length. Bars are mean values (n=7) ± s.e.m. Asterisks indicate values that were

determined by Student's *t* test to be significantly different (P < 0.05) from WT.

**Figure 2. Dry weight and parameters related with leaf area for Micro-Tom (WT) and**

**isogenic *ARF4* antisense transgenic line (*ARF4*-as), 45 days after germination.** (a) leaf

dry weight. (b) stem dry weight. (C) root dry weight. (d) total dry weight. (e) total leaf area.

(f) specific leaf area (SLA). Values are means ± s.e.m (n=8). Asterisks indicate values that

were determined by Student's *t* test to be significantly different (P < 0.05) from the Micro-

tom (WT).

**Figure 3. Leaf anatomy and morphology for Micro-Tom (WT) and *ARF4*-as transgenic**

**line**. Representative leaf cross-sections of Micro-Tom (WT, a) and *ARF4*-antisense line

(*ARF4*-as, b). Scale bars= 20 µm.  Total thickness of the leaf blade (c)Thickness of palisade

(d), spongy (e) and ratio of palisade to spongy mesophyll (f), proportion of intercellular air

spaces (g) and adaxial (h) and abaxial (i) and mesophyll (j) thickness in WT and *ARF4*-as

plants. Values are means ± s.e.m (n=6).

1187

1188 **Figure 4**. **Photosynthetic assimilation rate and stomatal conductance in Micro-Tom**

1189 **(WT) and *ARF4*-antisense silencing line.** (a) Net photosynthesis (*A*) curves in response to

1190 sub-stomatal (*C*$_i$) $CO_2$ concentration in WT and *ARF4*-as plants. Values are presented as

1191 means ± s.e.m. (n=5) obtained using the fully expanded fifth leaf. Two-branch curves: the

biochemically based leaf photosynthesis model (Farquhar et al., 1980) was fitted to the data

based on $C_i$, values of $A/C_i$ for five plants of WT (gray) and *ARF4-as* (black). (b) Response of stomatal conductance ($g_s$) to changes in leaf-to-air vapor pressure deficit (VPD). Vapor pressure difference was varied by changing the humidity of air, keeping leaf temperature constant. Each point represents one measurement on an individual plant.

**Figure 5**. *SlARF4* **expression in response to salt and drought stresses.** Gene expression in leaves (a) and roots (b) of WT plants exposed to salt stress, in leaves (c) and roots (d) of WT plants exposed to drought stress. GUS activity in pARF4::GUS tomato lines in salt (e) or drought (f) stress conditions. Stars (*) indicate the statistical significance (p<0,05) according to Student's t-test.

**Figure 6. Growth parameters of tomato wildtype (WT),** *ARF4-***as and** *ARF4-***ox in response to salt and drought stresses.** (a) and (b) shoot fresh weight in salt and drought stress conditions respectively, (c) and (d) root fresh weight in salt and drought stress conditions respectively, (e) and (f) primary root length root in salt and drought stress conditions respectively, (g) and (h) root density in salt and drought stress conditions respectively. Salt and drought stresses were performed on three weeks tomato plants for two weeks by adding 100mM of NaCl or 150 mM of NaCl for salt stress or 5% or 15% of PEG 20 000 for drought. Values are mean ± SD of three biological replicates. Bars with different letters indicate the statistical significance (p<0,05) according to Student Newman-Keuls test.

**Figure 7. Stomatal conductance and relative water content in WT and** *ARF4-***as plants.** (a) and (b) stomatal conductance in salt and drought stress conditions, (c) relative water content in response to dehydration. Bars with different letters indicate the statistical significance (p<0,05) according to Student Newman-Keuls test.

**Figure 8. Expression of** *SlNCED1***,** *SlNCED2***,** *SlCYP707A1***,** *SlCYP707A2* **and** *SlCYP707A3* **genes in WT and** *ARF4-***as leaves and roots after 2H and 24H of salt stress**

1216     **and 48H of drought stress application.** ΔΔCt refers to fold differences in gene expression

1217     relative to untreated plants. Values presented are mean ± SD of three biological replicates.

1218     Asterisks (*) indicate the statistical significance (p<0,05) according to Student's t-test.

1219

**Figure 1. Phenotype of cv. Micro-Tom tomato plants (wild-type, WT) and isogenic *ARF4* antisense transgenic line (*ARF4*-as), 35 days after germination**. (a) WT and *ARF4*-as plants (b) at the same stage of development. (c) Upward-curled leaf phenotype of *ARF4*-as compared to WT (d) number of leaves to first inflorescence, (e) plant weight and (f) plant internode length. Bars are mean values (n=7) ± s.e.m. Asterisks indicate values that were determined by Student's *t* test to be significantly different (P < 0.05) from WT.

**Figure 2. Dry weight and parameters related with leaf area for Micro-Tom (WT) and isogenic *ARF4* antisense transgenic line (*ARF4*-as), 45 days after germination.** (a) leaf dry weight. (b) stem dry weight. (C) root dry weight. (d) total dry weight. (e) total leaf area. (f) specific leaf area (SLA). Values are means ± s.e.m (n=8). Asterisks indicate values that were determined by Student's *t* test to be significantly different (P < 0.05) from the Micro-tom (WT).

**Figure 3. Leaf anatomy and morphology for Micro-Tom (WT) and *ARF4*-as transgenic line**. Representative leaf cross-sections of Micro-Tom (WT, a) and *ARF4*-antisense line (*ARF4*-as, b). Scale bars= 20 µm. Total thickness of the leaf blade (a)Thickness of palisade (d), spongy (e) and ratio of palisade to spongy mesophyll (f), proportion of intercellular air spaces (g) and adaxial (h) and abaxial (i) and mesophyll (j) thickness in WT and *ARF4*-as plants. Values are means ± s.e.m (n=6).

**Figure 4**. **Photosynthetic assimilation rate and stomatal conductance in Micro-Tom (WT) and *ARF4*-antisense silencing line.** (a) Net photosynthesis (*A*) curves in response to sub-stomatal ($C_i$) $CO_2$ concentration in WT and *ARF4*-as plants. Values are presented as means ± s.e.m. (n=5) obtained using the fully expanded fifth leaf. Two-branch curves: the biochemically based leaf photosynthesis model (Farquhar et al., 1980) was fitted to the data based on $C_i$, values of $A/C_i$ for five plants of WT (gray) and *ARF4-as* (black). (b) Response of stomatal conductance ($g_s$) to changes in leaf-to-air vapor pressure deficit (VPD). Vapor pressure difference was varied by changing the humidity of air, keeping leaf temperature constant. Each point represents one measurement on an individual plant.

**Figure 5**. *SlARF4* **expression in response to salt and drought stresses.** Gene expression in leaves (a) and roots (b) of WT plants exposed to salt stress, in leaves (c) and roots (d) of WT plants exposed to drought stress. GUS activity in pARF4::GUS tomato lines after 24 hours of salt exposure (e) or after 5 days of drought (f) stress application. Stars (*) indicate the statistical significance (p<0,05) according to Student's t-test.

**Figure 6. Growth parameters of tomato wildtype (WT), *ARF4*-as and *ARF4*-ox in response to salt and drought stresses.** (a) and (b) shoot fresh weight in salt and drought stress conditions respectively, (c) and (d) root fresh weight in salt and drought stress conditions respectively, (e) and (f) primary root length root in salt and drought stress conditions respectively, (g) and (h) root density in salt and drought stress conditions respectively. Salt and drought stresses were performed on three weeks tomato plants for two weeks by adding 100mM of NaCl or 150 mM of NaCl for salt stress or 5% or 15% of PEG 20 000 for drought. Values are mean ± SD of three biological replicates. Bars with different letters indicate the statistical significance (p<0,05) according to Student Newman-Keuls test.

**Figure 7. Stomatal conductance and relative water content in WT and *ARF4*-as plants.** (a) and (b) stomatal conductance in salt and drought stress conditions, (c) relative water content in response to dehydration. Bars with different letters indicate the statistical significance ($p<0,05$) according to Student Newman-Keuls test.

**Figure 8. Expression of *SlNCED1*, *SlNCED2*, *SlCYP707A1*, *SlCYP707A2* and *SlCYP707A3* genes in WT and *ARF4*-as leaves and roots after 2H and 24H of salt stress and 48H of drought stress application.** ΔΔCt refers to fold differences in gene expression relative to untreated plants. Values presented are mean ± SD of three biological replicates. Asterisks (*) indicate the statistical significance ($p < 0,05$) according to Student's t-test.