Elevated exposure to prenatal thyroid hormones affects embryonic mortality but

has no effects into adulthood

Tom Sarraude*[1,2], Bin-Yan Hsu[1], Ton G.G. Groothuis[2], Suvi Ruuskanen[1]

[1] Department of Biology, University of Turku, Finland

[2] Groningen Institute for Evolutionary Life Sciences, University of Groningen, the

Netherlands

[*] Corresponding author: Tom Sarraude, t.sarraude@rug.nl

Keywords: maternal hormones; thyroid hormones; avian growth; hatching success; Japanese

quails; life-history strategies.

Summary statement: Thyroid hormones are important hormones in all vertebrates, although

overlooked in the context of maternal effect. We found short-term effects of prenatal THs but

no evidence of programming effects.

Abstract

Maternal thyroid hormones (THs) are known to be crucial in embryonic development in
humans, but their influence on other, especially wild, animals remains poorly understood. So
far, the studies that experimentally investigated the consequences of maternal THs focused on
short-term effects, while long-term organisational effects, as shown for other prenatal
hormones, could also be expected. In this study, we aimed at investigating both the short- and
long-term effects of prenatal THs in a bird species, the Japanese quail *Coturnix japonica*. We
experimentally elevated yolk TH content (the prohormone T_4, and its active metabolite T_3, as
well as a combination of both hormone) We analysed hatching success, embryonic
development, offspring growth and oxidative stress as well as their potential organisational
effects on reproduction, moult, and oxidative stress in adulthood. We found that eggs injected
with both hormones had a higher hatching success compared with control eggs, suggesting
conversion of T_4 into T_3 by the embryo. We detected no other short-term or organisational
effects of yolk THs. Unfortunately, sex-specific responses could not be properly tested due to
low sample sizes. These results suggest that yolk thyroid hormones are important in the
embryonic stage of precocial birds, but may not have other short and long-term consequences,
at least not in captivity. Research on maternal thyroid hormones will greatly benefit from
studies investigating how embryos use and respond to this maternal signalling. Long-term
studies on prenatal THs in other taxa in the wild are needed for a better understanding of this
hormone-mediated maternal pathway.

Introduction

Maternal effects represent all the non-genetic influences of a mother on her offspring and have received increasing attention in evolutionary and behavioural ecology. Through maternal effects, mothers can influence the fitness of their progeny by adapting their phenotype to expected environmental conditions ("adaptive maternal effects" in Marshall and Uller, 2007; Mousseau and Fox, 1998), and this view is now also incorporated in the human disease literature (Gluckman et al., 2005). Maternal hormones transferred to the offspring can mediate important maternal effects. Historically, research on maternal hormones has mostly focused on steroid hormones (Groothuis et al., 2005; von Engelhardt and Groothuis, 2011). Recently, thyroid hormones (THs) also received attention in the context of hormone-mediated maternal effects (Ruuskanen and Hsu, 2018).

Thyroid hormones (THs) are metabolic hormones produced by the thyroid gland and are present in two main forms: the prohormone thyroxine (T_4) and the biologically active form triiodothyronine (T_3). THs play a crucial role in various aspects of an individual's life, e.g. development, metabolism and reproduction, across vertebrates, including humans (Krassas et al., 2010; Morreale de Escobar et al., 2004). In humans, physiological variation of maternal THs (i.e. no clinical symptoms in both mothers and foetuses) is found to be associated with infant birth weight and IQ in older children (Korevaar et al., 2016; Medici et al., 2013). In birds as well, THs play a role in brain development and neuronal turnover (reviewed in McNabb, 2007). THs control the endothermic heat production, and are therefore important in thermoregulation in juveniles and adults (McNabb and Darras, 2015).

THs can act, in concert with other hormonal axes, as mediators of life stage transitions across vertebrates (e.g., reviewed in Watanabe et al., 2016). The interaction between thyroid hormones and corticosteroids on amphibian metamorphosis is a well-known example of such effect on life stage transition (Kikuyama et al., 1993; Wada, 2008). THs are involved in gonadal development, and hyperthyroidism tends to fasten maturation (Holsberger and Cooke, 2005), and coordinate the transition between reproduction and moult (McNabb and Darras, 2015). Administration of exogenous THs is known to stop egg laying and induce moult in birds (Keshavarz and Quimby, 2002; Sekimoto et al., 1987). THs are also involved in photoperiodic control in seasonal breeding (Dardente et al., 2014). For example, thyroidectomised starlings transferred to long photoperiods became insensitive to future changes in photoperiod, and short photoperiod did not induce gonadal regression (Dawson, 1993).

While there has been recent research effort on the influence of maternal THs on
offspring traits across vertebrate taxa, there are still substantial knowledge gaps. First,
research on maternal thyroid hormones up to date has mainly investigated the short-term
effects of prenatal THs on developing fish (Brown et al., 1988; Raine et al., 2004) and
amphibians (Duarte-Guterman et al., 2010; Fini et al., 2012) and pre-fledging birds (Hsu et
al., 2017; Hsu et al., 2019; Ruuskanen et al., 2016). So far, only a study on rock pigeons has
looked at the influence of yolk THs on post-fledging survival and found no effect (Hsu et al.,
2017). None of these studies in any taxa investigated the potential organisational effects of
prenatal THs on life-history stage transitions in adult life.

Second, previous studies on prenatal THs in birds focused only on altricial species
(great tits, Ruuskanen et al., 2016; rock pigeons, Hsu et al., 2017; collared flycatchers, Hsu et
al., 2019). Embryonic development differs substantially between altricial and precocial
species. In the latter, embryonic development is more advanced than in the former. In
addition, precocial embryos start their endogenous production of TH around mid-incubation,
considerably earlier than their altricial counterparts, in which endogenous TH production
begins only after hatching (McNabb et al., 1998). While embryonic hormone production may
limit the influence of maternal hormones, prenatal hormones have been shown to affect chick
endogenous production and sensitivity (Pfannkuche et al., 2011). Overall, exposure to
maternal hormones may be of different importance in these two developmental modes.

Third, previous research has studied the effects of T_3 only (Fini et al., 2012; Raine et
al., 2004; Walpita et al., 2007) or a combination of T_3 and T_4 (Hsu et al., 2017; Hsu et al.,
2019; Ruuskanen et al., 2016), where the effects of the two forms cannot be separated.
Although T_3 is the biologically active form that binds to the receptors, both are deposited in
eggs (Prati et al., 1992) and T_4 may be converted to T_3 via deiodinases from the mother or the
developing embryo (Van Herck et al., 2015) or may still exert non-genomic actions (reviewed
in Davis et al., 2016). Manipulating yolk T_4 and T_3 independently would help understanding
the relative contribution of these two hormones.

In this study, we aimed at assessing the effects of maternal THs on development and
life-history traits in a precocial bird species, the Japanese quail (*Coturnix japonica*). To do so,
manipulated eggs received either an injection of T_4 or T_3 separately, a combination of both
hormones, or a control injection of the vehicle saline solution. This design allowed us to
explore the effects of T_4 and T_3 separately. The elevation in yolk THs remained within the
natural range of this species, a crucial condition to obtain relevant results for an eco-
evolutionary context. We measured traits known to be influenced by circulating and yolk THs:

hatching success, age at embryonic mortality, growth, transition between life-history stages

(i.e., reproductive state and moult) and oxidative stress. First, we hypothesise that elevation of

yolk THs in Japanese quails positively affects hatching success, as found in two studies on

collared flycatchers and rock pigeons (Hsu et al., 2017; Hsu et al., 2019, but see Ruuskanen et

al., 2016). Second, elevation of yolk THs is predicted to increase the proportion of well-

developed embryos before hatching, as found in rock pigeons (Hsu et al., 2017). We therefore

looked at the age at mortality in unhatched eggs. Third, we expect elevated yolk THs to affect

chick growth (in body mass, tarsus and wing length) either positively (Hsu et al., 2019;

Wilson and McNabb, 1997), negatively (Hsu et al., 2017), or in a sex-specific manner

(Ruuskanen et al., 2016). Fourth, we predict that yolk THs will have organisational effects on

life-history stage transitions; that is, age at sexual maturity and male gonadal regression

(using cloacal gland size as a proxy), and moult when birds are exposed to short photoperiod.

Based on the literature mentioned above we expect elevated yolk THs to advance the timing

of puberty, gonadal regression and moult. The rate of moult should also be influenced, with

birds receiving experimental TH elevation moulting faster. We also explored the effects of

yolk THs on reproductive investment in females, another important fitness aspect. Finally,

yolk THs may increase oxidative stress due to their stimulating effects on metabolism.

Material and Methods

Parental generation and egg collection

The parental generation was composed of adult Japanese quails provided by Finnish private

local breeders that were kept in two acclimated rooms. Twenty-four breeding pairs were

formed by pairing birds from different breeders. Individuals were identified using metal leg

bands. The floor was covered with 3–5cm sawdust bedding. A hiding place, sand and calcium

grit were provided. Each pair was housed in a 1 m² pen. The temperature was set to 20°C with

a 16L:8D photoperiod (light from 06.00 to 22.00). Food (Poultry complete feed, "Kanan Paras

Täysrehu", Hankkija, Finland) was provided *ad libitum* and water was changed every day.

Pairs were monitored every morning to collect eggs. Eggs were individually marked

(non-toxic marker), weighed and stored for 7 days in a climate-controlled chamber at 15°C

and 50% relative humidity. On the last day of collection, an average of 6.6 eggs per pair

(range = 4–8 eggs) were injected with a solution (see next section).

Preparation of the solution, injection procedure and incubation

The preparation of hormone solution and the procedure of injection were based on previous
studies (Hsu et al., 2017; Ruuskanen et al., 2016). In brief, crystal T_4 (L-thyroxine, $\geq$ 98%
HPCL, CAS number 51-48-9, Sigma-Aldrich) and T_3 (3,3',5-triiodo-L-thyronine, > 95%
HPCL, CAS number 6893-02-3, Sigma-Aldrich) were first dissolved in 0.1M NaOH and then
diluted in 0.9% NaCl. The injection of thyroid hormones resulted in an increase of two
standard deviations (T_4 = 8.9 ng/egg; T_3 = 4.7 ng/egg), a recommended procedure for
hormone manipulation within the natural range (Hsu et al., 2017; Podmokła et al., 2018;
Ruuskanen et al., 2016). The control solution (CO) was a saline solution (0.9% NaCl). The
concentrations of the hormone solutions were based on previous measurements of 15 eggs
from the same flock (T_4 content per egg (SD) = 15.3 (4.4) ng, T_3 content per egg (SD) = 7.6
(2.3) ng).

Hormone injections were performed at room temperature in a laminar hood. Eggs
were put sideways, allowing yolks to float up to the middle position. Before injection, the
shell was disinfected with a cotton pad dipped in 70% EtOH. We used a 27G needle (BD
Microlance ™) to pierce the eggshell and then used a 0.3 ml syringe to deliver 50 µl of the
respective hormone solution or control. After injection, the hole was sealed with a sterile
plaster (OPSITE Flexigrid, Smith&Nephew).

In total, 158 eggs were injected and divided as follows over the treatments: T_3
treatment (N = 39); T_4 treatment (N = 39); T_3+T_4 treatment (N = 40); and control, CO (N =
40). To balance the genetic background of the parents and the effect of storage, each egg laid
by the same female was sequentially assigned to a different treatment and the order of
treatments was rotated among females. After injection, eggs were placed in an incubator at
37.8°C and 55% relative humidity. Until day 14 after starting incubation, eggs were
automatically tilted every hour by 90°. On day 14, tilting was halted and each egg was
transferred to an individual container to monitor which chick hatched from which egg. On day
16 after injection, (normal incubation time = 17 days), the temperature was set to 37.5°C and
the relative humidity to 70%. Eggs were checked for hatching every 4 hours from day 16
onwards. Four days after the first egg hatched, all unhatched eggs were stored in a freezer and
dissected to determine the presence of an embryo. The age of developed embryos was
assessed according to Ainsworth et al. (2010).

Rearing conditions of the experimental birds

In total, 66 chicks hatched (N = 10 CO, 15 T_3, 20 T_4 and 21 T_3T_4). The overall hatching

success was rather low (ca. 40%), partly due to the injection procedure itself (Groothuis and

von Engelhardt, 2005), although low hatching success in quails has also been reported in

unmanipulated conditions previously (e.g. Okuliarová et al., 2007). Among the unhatched

eggs, 33.7% (31 out of 92) had no developed embryos. Twelve hours after hatching, the

chicks were marked by a unique combination of coloured rings and nail coding and

transferred to two 1 m² cages (ca. 30 chicks/cage, sex and treatments mixed together). The

chicks were provided with heating mats and lamps as extra heat sources for the first two

weeks. The chicks were fed with sieved commercial poultry feed ("Punaheltta paras

poikanen", Hankkija, Finland), and provided with Calcium and bathing sand. Two weeks after

hatching, the chicks were separated in four 1 m² cages of about 16 individuals. Around 3

weeks after hatching, coloured rings were replaced by unique metal rings. On week 4 after

hatching, birds were transferred to eight 1 m² pens (average of 7.1 birds/pen, range = 4–9),

under the same conditions as the parents. Around the age of sexual maturity (ca. 6–8 weeks

after hatching), the birds were separated by sex in twelve 1 m² pens (average of 4.8 birds/pen,

range = 4–5).

Monitoring of growth and reproductive maturation

Body mass and wing length were measured twelve hours after hatching. Tarsus was not

measured because it bends easily, resulting in inaccurate measures and potential harm for the

young. From day 3 to day 15, these three traits were monitored every 3 days. From day 15 to

day 78 (ca. 12 weeks), chicks were measured once a week. Body mass was recorded using a

digital balance to the nearest 0.1 g. Wing and tarsus lengths were respectively measured with

a ruler and a calliper to the nearest 0.5 mm and 0.1 mm. From week 6 to week 10, we

monitored cloacal gland development and foam production in 28 males. Cloacal glands were

measured every other day with a calliper to the nearest 0.1 mm as a proxy for testes

development and sexual maturation (Biswas et al., 2007). Foam production (by gently

squeezing the cloacal gland) was assessed at the same time and coded from 0 (no foam) to 3

(high production of foam), as a proxy of cloacal gland function (Cheng et al., 1989a; Cheng et

al., 1989b). The same observer performed all measurements. We collected eggs produced by

10-week-old females over a 6-day period, and measured the short and long axes of the eggs

with a calliper to the nearest 0.01 mm and record their mass to the nearest 0.1 g. We collected

on average 5.7 eggs (range = 4–7) per female from 28 females.

Monitoring of cloacal gland regression and moult

In Japanese quails, exposure to short photoperiod and cold temperature triggers reproductive inhibition and postnuptial moulting (Tsuyoshi and Wada, 1992). Thyroid hormones are known to coordinate these two responses (see introduction). When the birds reached the age of ca. 7 months, we exposed birds to short photoperiod (8L:16D, i.e., light from 08.00 to 16.00) with a 12:12-h cycle of normal (20°C) and low (9°C) temperature (low temperature was effective from 18.00 to 06.00). Cloacal gland regression (as a proxy for testes regression) was monitored every other day for 2 weeks with a calliper by measuring the width and length to obtain the area of the gland to the nearest 0.1 mm² (N = 26 males). Primary moult was recorded from a single wing by giving a score to each primary from 0 (old feather) to 5 (new fully-grown feather) following Ginn and Melville (1983) (N = 54 males and females). The total score of moulting was obtained by adding the score of all feathers.

Oxidative status biomarker analyses

Two blood samples were drawn, when birds were 2 weeks (N = 51 chicks) and 4 months old (N = 49 adults), respectively. 200 µl of blood was collected from the brachial vein in heparinized capillaries and directly frozen in liquid nitrogen. Then, the samples were stored at -80°C until analyses. We measured various biomarkers of antioxidant status; the antioxidant glutathione (tGSH), the ratio of reduced and oxidised glutathione (GSH:GSSG) and activity of the antioxidant enzymes glutathione peroxidase (GPx), catalase (CAT) and superoxide dismutase (SOD) from the blood. It is important to measure multiple biomarkers of oxidative and antioxidant status for a broader understanding of the mechanism. Also, the interpretation of the results is more reliable if multiple markers show similar patterns. Of the measured biomarkers, the ratio of GSH:GSSG represents the overall oxidative state of cells and consequently, deviations in this ratio is often used as an indicator of oxidative stress (Halliwell and Gutteridge, 2015; Hoffman, 2002; Isaksson et al., 2005; Lilley et al., 2013; Rainio et al., 2013). GPx enzymes catalyse the glutathione cycle, whereas CAT and SOD directly regulate the level of reactive oxygen species (ROS) (Ercal et al., 2001; Halliwell and Gutteridge, 2015). The methodology for measuring each biomarker is described in detail in Rainio et al. (2015). All analyses were conducted blindly of the treatment. Briefly, the samples were analyzed using a microplate reader (EnVision, PerkinElmer-Wallac, Finland). All antioxidant and enzyme activities were measured in triplicate (intra-assay coefficient of variability [CV] < 10% in all cases) using 96-(CAT) or 384-well (GPx, SOD, tGSH and GSH:GSSG) microplates. Three control samples were used with each plate, to be able to

correct inter-assay precision with the ratio specific to the particular plate. Overall protein

concentration (mg/ml) was measured according to the Bradford method (Bradford, 1976)

using BioRad stock (BioRad, Finland) diluted with dH_2O (1:5) and BSA (bovine serum

albumin, 1 mg/ml) (Sigma Chemicals, USA) as a standard. GPx-assay was conducted using

Sigma CGP1 kit, CAT-assay using SigmaCAT100 kit and SOD-assay using Fluka 19160 SOD

determination kit. Total GSH and the ratio of GSH:GSSG were measured with the ThioStar®

glutathione detection reagent (Arbor Assays, USA) according to kit instructions, using

reduced glutathione as a standard (Sigma Chemicals, USA).

Ethics

The study complied with Finnish regulation and was approved by the Finnish Animal

Experiment Board (ESAVI/1018/04.10.07/2016).

Statistical analysis

Data were analysed with the software R version 3.5.3 (R core team, 2019). In this study, two

different statistical approaches were used: null-hypothesis testing with Generalised Linear

Mixed Models (GLMMs) and Linear Mixed Models (LMMs), and multimodel inference with

Generalised Additive Mixed Models (GAMMs). GAMMs were used to analyse the data on

body and cloacal gland growth to account for its non-linear pattern (see *Growth*). In this

analysis, we preferred multimodel inference as GAMMs generate many candidate models that

cannot be directly compared (e.g., by the Kenward-Roger approach). Instead, candidate

models were ranked based on their Akaike Information Criterion (AIC) values. Models with a

$\Delta AIC \leq 2$ from the top-ranked model were retained in the set of best models. Akaike weights

of all models were calculated following (Burnham and Anderson, 2002), and evidence ratios

of the top-ranked models were calculated as the weight of a model divided by the weight of

the null model (Burnham et al., 2011). To estimate the effect of the predictors, we computed

the 95% confidence intervals from the best models using the *nlme* package (Pinheiro et al.,

2018). GLMMs and LMMs were fitted using the R package *lme4* (Bates et al., 2015), and

GAMMs were fitted using the package *mgcv* (Wood, 2017). P-values for GLMMs were

obtained by parametric bootstrapping with 1,000 simulations and p-values for LMMs were

calculated by model comparison using Kenward-Roger approximation, using the package

pbkrtest in both cases (Halekoh and Højsgaard, 2014). Post-hoc Tukey analyses were

conducted with the package *multcomp* (Hothorn et al., 2008). Due to our experiment design,

eggs injected with both hormones received a higher absolute amount of hormones than eggs

injected with T_4 or T_3 only. Therefore, we also tested a potential dose-dependent effect of the

treatment on the response variables when treatment groups showed significant differences.

Model residuals were checked visually for normality and homoscedasticity. Covariates and

interactions were removed when non-significant ($\alpha = 0.05$).

Hatching success

To analyse hatching success, each egg was given a binary score: 0 for unhatched egg and 1 for

hatched egg. A series of GLMMs were fitted with a binomial error distribution (logit link) and

mother identity as a random intercept. The first model included the 4-level treatment

(treatments: CO, T_3, T_4 and T_3T_4) as the predictor, while in a second model treatment was

converted into an ordered variable, following the increasing levels (i.e. CO, T_3, T_4 and T_3T_4).

The second model was meant to test for a potential dose-dependent effect as the eggs received

an increasing amount of total THs as a potential source of T_3, the most potent hormone. Egg

mass might affect hatchability and was therefore added as a covariate in both models. The

potential effect of storage duration on hatchability (Reis et al., 1997) was accounted for by

including laying order as a covariate in both models.

Duration of embryonic period, age at embryonic mortality and mass at hatching

Duration of embryonic period and mass at hatching were modelled with LMMs. Treatment,

sex of the individuals and egg mass were included as fixed factors. Laying order was added as

a covariate to account for potential effects of storage duration on hatching time and on chick

weight (Reis et al., 1997). Mother identity was included as a random intercept. In the model

for mass at hatching, duration of embryonic period was further added as a covariate.

286   The data for embryonic age had a skewed distribution and residuals were not normally

distributed and heterogenous, which violated LMM assumptions on residual distribution. We

therefore performed a simple Kruskal-Wallis test.

Growth

As growth curves typically reach an asymptote, we fitted non-linear GAMMs to these curves.

Growth in body mass, tarsus and wing length were analysed in separate GAMMs. Growth

was analysed until week 10 after hatching as all birds appeared to have reached their

maximum body mass and tarsus and wing length. The data are composed of repeated

measurements of the same individuals over time; therefore, we first corrected for temporal

autocorrelation between the measurements using an ARMA(1,1) model for the residuals (Zuur

et al., 2009). Second, as mothers produced several eggs, the models included nested random

effects, with measured individuals nested into mother identity, allowing for random intercepts.

GAMMs allow modelling the vertical shift of the curves (i.e., changes in intercepts) and their

shape. Treatment and sex were included as predictors. A smoothing function for the age of the

birds was included to model the changes in the growth curves, and was allowed to vary by sex

or treatment only, or none of these predictors. The interaction between sex and treatment was

not analysed due to low statistical power. Additive effect of treatment and sex was tested for

the intercept but could not be computed for curve shape. All combinations of the relevant

predictors were tested for both shape parameters (i.e., intercept and curve shape).

Reproductive maturation, regression and investment

Due to low sample sizes in sex-specific responses, we could not perform robust statistical

analyses. We therefore present these analyses and results in the supplementary material and

only briefly discuss them.

Oxidative stress

A principal component analysis (PCA) was first performed on measured antioxidant markers

(SOD, CAT, GPx, tGSH and GST), to reduce the number of metrics for subsequent analyses.

The first and the second principal components (PCs) explained together 60.2% of the variance

(Table 1). PC1 and PC2 were then used as dependent variables in separate LMMs. LMMs

included the treatment, sex and age of individuals (2 weeks and 4 months old) as fixed factors

and the 2-way interactions between treatment and sex, and treatment and age. Mother and

individual identities, to account for repeated measures, were added as random intercepts.

Malondialdehyde (MDA) is a marker of oxidative damage, which is a different measure from

antioxidant activity, and was therefore analysed in a separate LMM using the same parameters

as for PC1 and PC2, adding the batch of the assay as an additional random intercept. The

marker of cell oxidative status (GSH:GSSG ratio) was analysed with the same model used for

PC1 and PC2.

Moult

Two parameters of moult were analysed in separate LMMs: the timing of moult (i.e., the

moult score after one week of short photoperiod), and the rate of moult (i.e., how fast birds

moulted). Both models included treatment and sex as fixed factors, and mother identity as a

random intercept. The rate of moult was tested by fitting an interaction between treatment and

age. This model also included the main effect of age and individual identity, nested within

mother identity, as a random intercept to account for repeated measures. Estimated marginal

means and standard errors (EMMs ± SE) were derived from the model using the package

emmeans (Lenth, 2019).

Results

Effects of prenatal THs on hatching success and age of embryo mortality

There was a significant effect of elevated prenatal THs on hatching success (GLMM, p =

0.05, Fig. 1). Tukey post-hoc analysis revealed that hatching success in the T_3T_4 group was

significantly higher than in the CO group (Estimate±SE = 1.24±0.50, Tukey z = 2.46, p =

0.05). The other groups (T_3 and T_4) were not different from the control group (all $z < 2.22$ and

p > 0.09). The data suggested a dose-dependent effect that we tested by changing the

treatment factor to an ordered variable, as the eggs received an increasing amount of TH from

T_3 to T_3T_4 injections ($T_3 < T_4 < T_3T_4$). We found a linear positive dose-dependent effect of

yolk TH elevation (GLMM, Estimate±SE = 0.96±0.36, z = 2.69, p = 0.007), but no quadratic

or cubic effects (p > 0.53).

Dissection of the unhatched eggs showed that age of embryo mortality did not differ

between the treatments (Kruskal-Wallis χ^2 = 7.22, df = 3, p = 0.07). Finally, the manipulation

of yolk THs did not affect the duration of embryonic period (LMM, $F_{3,42.0}$ = 0.57, p = 0.64,

Fig. S1). Sex of the embryo or egg mass (LMM sex, $F_{1,49.7}$ = 2.63, p = 0.11; LMM egg mass,

$F_{1,19.3}$ = 0.01, p = 0.92) were also not associated with the duration of the embryonic period.

Effects of prenatal THs on growth

Mass at hatching was not influenced by the elevation of prenatal THs (LMM, $F_{3,35.0}$ = 0.81, p

= 0.50, Fig. S2). Mass at hatching was positively correlated with egg mass (LMM,

Estimate±SE = 0.72±0.10 g, $F_{1,24.1}$ = 46.9, p < 0.001), while duration of embryonic period

was negatively correlated with mass at hatching (LMM, Estimate±SE = -0.008±0.003 g,

$F_{1,46.7}$ = 4.49, p = 0.04).

Regarding body mass growth, the top-ranked model showed that the curve shape and

the intercept differ according to sex (Table 2). After 10 weeks, females had a larger body mass

than males (mean±SE females = 214.4±5.7 g, males = 172.4±4.5 g, Fig. 2), which was

supported by the 95% CIs (Table 3). Based on model selection we conclude that the treatment

had no effect on body mass growth (Table 2).

For wing length, the top-ranked model (ΔAIC $\leq$ 2) included sex in the intercept, while

treatment was not included in the best supported model (Table S1). The 95% CIs (Table 3)

confirmed that males had a lower wing length than females (Fig. S3).

Concerning tarsus length, the models within ΔAIC $\leq$ 2 included no predictors for the

curve shape but included treatment for the intercept (Table S2). The 95% CIs of the parameter

estimates from these models suggested that there was a slight negative effect of T_3T_4

treatment on tarsus growth (Table 3, Fig. S4). However, as the estimates were close to 0

(Table 3) and evidence ratios showed that the model with treatment as a predictor was only

3.5 times more supported than the null model (Table S2), we conclude that the effect of THs

on tarsus length is likely to be very small. Likewise, the second model for tarsus length

included sex as a predictor for the intercept, but its 95% CIs overlapped with 0 (Table 3). We

therefore conclude that sex had no effect on tarsus growth.

Effects of prenatal THs on postnuptial moult

As expected, birds started to moult soon after being exposed to short photoperiod, with an

average increase of moult score by 6 per week (SE = 0.2, $F_{1,254.0}$ = 827.4, p < 0.001, Fig. 3).

The first moult score (assessed one week after switching to short photoperiod) was not

affected by the treatment (LMM, $F_{3,42.7}$ = 0.36, p = 0.78), but was influenced by sex, with

females having a higher score than male (EMMs $\pm$ SE: female = 21.4 $\pm$ 1.6, male = 7.2 $\pm$ 1.7;

LMM $F_{1,45.3}$ = 41.9, p < 0.001). Yolk TH elevation did not affect the rate of moult (LMM

interaction treatment $\times$ time, $F_{3,251.0}$ = 0.59, p = 0.62, Fig. 3).

Effects of prenatal THs on oxidative stress

The elevation of yolk THs had no effect on PC1 or PC2 of antioxidants at either 2 weeks

("chicks") or 4 months ("adults") old (LMM on PC1, $F_{3,40.3}$ = 2.40 , p = 0.08; LMM on PC2,

$F_{3,42.2}$ = 0.92, p = 0.44, treatment $\times$ age, F < 0.91, p > 0.44). The age of the birds had a highly

significant effect on PC1, with chicks generally having higher antioxidant capacities (CAT,

GST and tGSH) than adults (LMM, Estimate$\pm$SE = -1.34$\pm$0.19, $F_{1,49.2}$ = 52.1, p < 0.0001). All

the other predictors had no effect on either PC1 or PC2 (all F < 2.93 and all p > 0.09).

The marker of oxidative damage, MDA, was affected by the elevation of yolk THs

(LMM, $F_{3,43.6}$ = 3.08, p = 0.04, Fig. 4). Tukey post-hoc analysis showed that the T4 group had

higher MDA values than the T3 group (Estimate$\pm$SE = 0.01$\pm$0.004, Tukey contrast p = 0.01),

but none of the groups differed from the control (Tukey p-values > 0.19). However, this result

became non-significant when removing the outlier in the T_4 group (LMM, $F_{3,43.1}$ = 2.68, p =

0.06). MDA levels were not affected by the age or the sex of individuals (LMM age, $F_{1,54.4}$ =

0.30, p = 0.59; LMM sex, $F_{1,42.0}$ = 1.47, p = 0.23).

The marker of cell oxidative balance, GSH:GSSG, was not influenced by the yolk THs

nor by the sex of the birds (LMM treatment, $F_{3,33.0} = 0.85$, p = 0.48; LMM sex, $F_{1,40.6} = 0.57$,

p = 0.45). However, chicks had a higher GSH:GSSG ratio than adults (LMM, Estimate±SE =

0.17±0.04, $F_{1,50.0} = 18.3$, p < 0.0001).

Discussion

The aim of this experimental study was to investigate the short-term and organisational effects

of maternal thyroid hormones (THs) in a precocial species, the Japanese quail, by

experimental elevation of THs in eggs. Our study is the first to investigate the effects of yolk

T_3 and T_4 separately, within the natural range of the study model. In addition we studied both

short and long term effects on embryonic development, growth, life stage transitions and

oxidative stress. We only detected a positive effect of yolk THs on hatching success. All other

response variables studied were not affected by elevated prenatal THs.

Effects of prenatal THs on hatching success and embryonic development

We found that hatching success increased when the eggs received an injection of both T_4 and

T_3. Previous similar studies reported comparable effects of yolk THs in rock pigeons (Hsu et

al., 2017) and in collared flycatchers (Hsu et al., 2019). In these studies, injections consisted

of a mixture of both T_3 and T_4. Our results point towards a dose-dependent effect of yolk THs,

as found previously of androgen hormones (e.g., Muriel et al., 2015). Importantly, given that

only T_3 binds to receptors, these results also suggest that embryos must express deiodinase

enzymes to convert T_4 to T_3, and/or yolk may contain maternally derived deiodinase mRNA.

Indeed, precocial embryos start to produce endogenous T_3 and deiodinase expression has

previously been characterised in chicken embryos (Darras et al., 2009; Van Herck et al.,

2012). In contrast with our study, a similar study in great tits detected no increased hatching

success due to the injection of THs (Ruuskanen et al., 2016). The dissimilarities between the

studies may come from inter-specific differences in terms of utilisation of yolk THs by the

embryos or from context-dependent effects (e.g. due to other egg components). Further

comparative and mechanistic studies could help understanding the dynamic of yolk THs

during incubation.

Increased yolk THs did not improve age of embryo mortality. Similar to our study,

Ruuskanen et al. (2016) did not find any difference in the timing of mortality in great tit

embryos. Conversely, the study on rock pigeons found that yolk THs increased the proportion

of well-developed embryos (Hsu et al., 2017). Similarly to our result on hatching success,

yolk THs' effects on embryonic development may differ in a species-specific manner.

Effects of prenatal THs on growth

We found no influence of yolk THs on growth, contrary to our expectations based on the
recent literature. Other comparable studies found either a positive (Hsu et al., 2019), a
negative (Hsu et al., 2017) or a sex-specific effect (Ruuskanen et al., 2016) of yolk THs on
growth. This notable difference may be due to the captive conditions experienced by the
Japanese quails in our study, with unrestricted access to food and water. Although the pigeon
study also provided ad libitum food, parents still needed to process food before feeding their
nestlings in the form of crop milk, whereas precocial quails have no such limitation. In
addition, the Japanese quail has been domesticated for many generations, and probably
selected for rapid growth for economic reasons. Whole-genome sequencing in chickens
showed that domestication induced a strong positive selection on genes associated with
growth (Rubin et al., 2010). Interestingly, that study also found a strong selection for a locus
associated with thyroid stimulating hormone (TSH) receptor. TSH controls most of the TH
production by the thyroid gland (McNabb and Darras, 2015), and this artificial selection may
overshadow the effects of natural variations of prenatal THs on growth. Besides, the low
number of individuals in the control and T_3 groups (7 and 11, respectively) may have limited
statistical power to detect differences between the treatments. Repeating the study with a
larger sample size may allow us to ascertain the effects of yolk THs on growth in precocial
study models. Research on the influences of prenatal THs on growth will also benefit from
experimental studies on wild precocial species.

Effects of prenatal THs on postnuptial moult

Short photoperiod in combination with cold temperature triggered primary moult, as expected.
However, we detected no effect of yolk THs on the timing or speed of moult. Thyroid
hormones are important in moult and feather growth (reviewed in Dawson, 2015). For
example, thyroidectomised birds fail to moult after being exposed to long photoperiods
(Dawson, 2015). In addition, thyroidectomised nestling starlings failed to grow normal adult
plumage and grown feathers presented an abnormal structure (Dawson et al., 1994). By
removing the thyroid gland, these two studies implemented extreme pharmacological
protocols that differ drastically from our injection of physiological doses. In addition, our
experimental design, increasing TH exposure (vs decreased TH exposure in the above-
mentioned studies), may have different consequences. For example, there may be a threshold

above which any, additional hormones may not affect moult.

Overall, our results show no support for the hypothesis of programming effect of

prenatal THs on life stage transitions. Yet, due to small sample sizes in sex-specific analyses

(i.e., male gonadal maturation and regression, and female reproductive investment), there

remains a relatively high uncertainty about the potential programming effects of prenatal THs.

Replicate studies with larger samples sizes and different study models will reduce this

uncertainty.

Effects of prenatal THs on oxidative stress

In contrast to our predictions, elevated yolk THs did not affect oxidative status during chick or

adult phase. We found no changes in antioxidant activities in relation to yolk THs and no

imbalance in the oxidative cell status. Nevertheless, T_4 birds had a higher level of oxidative

damage on lipids than T_3 birds, but this was a weak effect driven by one outlier. The lack of

effects on chick oxidative status among the treatment groups could be explained by the

absence of treatment effects on growth, given that high growth rates usually result in higher

oxidative stress and damage (e.g. Alonso-Alvarez et al., 2007). In turn, the lack of treatment

effects on adult oxidative status may suggest no organisational effects of prenatal THs on

adult metabolism. A recent study in an altricial species also found no influence of yolk THs on

nestling oxidative stress (Hsu et al., 2019). Our study shows for the first time that prenatal

THs have no influence on adult oxidative stress either. The previous study focused on a

limited set of biomarkers: one antioxidant enzyme, oxidative damage on lipids and oxidative

balance. In the present study, we measured 7 biomarkers, thus providing broader support to

the absence of effects of prenatal THs on oxidative stress.

Conclusion

To our knowledge, this study is the first one to experimentally investigate the consequences of

natural variations of maternal THs not only early but also in adult physiology and postnuptial

moult in any vertebrate. Furthermore, this study explored for the first time the effects of

maternal T_3 and T_4 separately. We found no evidence for differential effects of maternal T_4

and T_3, while a dose-dependent effect on hatching success suggests that T_4 is converted into

T_3, the biologically active form during embryonic development. Contrary to similar studies on

wild altricial species, we found no influence of maternal THs on growth. Further research on

embryos utilisation of maternal THs may help understand the differences observed between

precocial and altricial species. Studies in other vertebrates are urgently needed to understand

the potential for long-term organising effects of maternal THs.

List of symbols and abbreviations

- CAT: catalase
- CO: control treatment
- GP: glutathione peroxidase
- tGSH: oxidised glutathione
- GSSG: reduced glutathione
- GST: Glutathione S-transferase
- MDA: malonaldehyde
- RMR: resting metabolic rate
- SOD: super-oxide dismutase
- T_3: triiodothyronine
- T_4: thyroxine
- THs: thyroid hormones

Acknowledgements

We thank Sophie Michon for her help on setting up the parental generation. We also thank Ido Pen for consultation and help with statistical analysis, and Esther Chang for her help throughout the writing phase.

Competing interests

We declare no competing interests.

Funding

The study was funded by the Academy of Finland (grant no. 286278 to SR), the Finnish National Agency for Education (grant no. TM-15-9960 to TS) and the University of Groningen (grant to TS).

References

Ainsworth, S. J., Stanley, R. L. and Evans, D. J. R. (2010). Developmental stages of the Japanese quail. *J. Anat.* **216**, 3–15.

Alonso-Alvarez, C., Bertrand, S., Faivre, B. and Sorci, G. (2007). Increased susceptibility to oxidative damage as a cost of accelerated somatic growth in zebra finches. *Funct. Ecol.* **21**, 873–879.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **67**, 1–48.

Biswas, A., Ranganatha, O. S., Mohan, J. and Sastry, K. V. H. (2007). Relationship of cloacal gland with testes, testosterone and fertility in different lines of male Japanese quail. *Anim. Reprod. Sci.* **97**, 94–102.

Brown, C. L., Doroshov, S. I., Nunez, J. M., Hadley, C., Vaneenennaam, J., Nishioka, R. S. and Bern, H. A. (1988). Maternal triiodothyronine injections cause increases in swimbladder inflation and survival rates in larval striped bass, *Morone saxatilis*. *J. Exp. Zool.* **248**, 168–176.

Burnham, K. P. and Anderson, D. R. (2002). *Model selection and multimodel inference: a practical information-theoretic approach*. 2nd ed. New York: Springer.

Burnham, K. P., Anderson, D. R. and Huyvaert, K. P. (2011). AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behav. Ecol. Sociobiol.* **65**, 23–35.

Cheng, K. M., McIntyre, R. F. and Hickman, A. R. (1989a). Proctodeal Gland Foam Enhances Competitive Fertilization in Domestic Japanese Quail. *The Auk* **106**, 286–291.

Cheng, K. M., Hickman, A. R. and Nichols, C. R. (1989b). Role of the proctodeal gland foam of male Japanese Quail in natural copulations. *The Auk* **106**, 279–285.

Dardente, H., Hazlerigg, D. G. and Ebling, F. J. P. (2014). Thyroid Hormone and Seasonal Rhythmicity. *Front. Endocrinol.* **5**, 19.

Darras, V. M., Van Herck, S. L. J., Geysens, S. and Reyns, G. E. (2009). Involvement of thyroid hormones in chicken embryonic brain development. *Gen. Comp. Endocrinol.* **163**, 58–62.

Davis, P. J., Goglia, F. and Leonard, J. L. (2016). Nongenomic actions of thyroid hormone. *Nat. Rev. Endocrinol.* **12**, 111–121.

Dawson, A. (1993). Thyroidectomy progressively renders the reproductive system of starlings (Sturnus vulgaris) unresponsive to changes in daylength. *J. Endocrinol.* **139**, 51–55.

Dawson, A. (2015). Chapter 38 - Avian Molting. In *Sturkie's Avian Physiology (Sixth Edition)* (ed. Scanes, C. G.), pp. 907–917. San Diego: Academic Press.

Dawson, A., McNaughton, F. J., Goldsmith, A. R. and Degen, A. A. (1994). Ratite-like

neoteny induced by neonatal thyroidectomy of European starlings, *Sturnus vulgaris. J. Zool.* **232**, 633–639.

Duarte-Guterman, P., Langlois, V. S., Pauli, B. D. and Trudeau, V. L. (2010). Expression and T3 regulation of thyroid hormone- and sex steroid-related genes during Silurana (Xenopus) tropicalis early development. *Gen. Comp. Endocrinol.* **166**, 428–435.

Ercal, N., Gurer-Orhan, H. and Aykin-Burns, N. (2001). Toxic Metals and Oxidative Stress Part I: Mechanisms Involved in Metal-induced Oxidative Damage. *Curr. Top. Med. Chem.* **1**, 529–539.

Fini, J. B., Mével, S. L., Palmier, K., Darras, V. M., Punzon, I., Richardson, S. J., Clerget-Froidevaux, M. S. and Demeneix, B. A. (2012). Thyroid Hormone Signaling in the Xenopus laevis Embryo Is Functional and Susceptible to Endocrine Disruption. *Endocrinology* **153**, 5068–5081.

Ginn, H. and Melville, D. (1983). *{Moult in birds (BTO guide)}*. Thetford: British Trust for Ornithology.

Gluckman, P. D., Hanson, M. A. and Spencer, H. G. (2005). Predictive adaptive responses and human evolution. *Trends Ecol. Evol.* **20**, 527–533.

Groothuis, T. G. G. and von Engelhardt, N. (2005). Investigating Maternal Hormones in Avian Eggs: Measurement, Manipulation, and Interpretation. *Ann. N. Y. Acad. Sci.* **1046**, 168–180.

Groothuis, T. G. G., Müller, W., von Engelhardt, N., Carere, C. and Eising, C. (2005). Maternal hormones as a tool to adjust offspring phenotype in avian species. *Neurosci. Biobehav. Rev.* **29**, 329–352.

Halekoh, U., Højsgaard, S. 2014. A Kenward-Roger Approximation and Parametric Bootstrap Methods for Tests in Linear Mixed Models - The *R* Package pbkrtest. J. Stat. Softw. **59**.

Halliwell, B. and Gutteridge, J. M. C. (2015). *Free Radicals in Biology and Medicine*. Oxford University Press.

Hoffman, D. J. (2002). Role of selenium toxicity and oxidative stress in aquatic birds. *Aquat. Toxicol.* **57**, 11–26.

Holsberger, D. R. and Cooke, P. S. (2005). Understanding the role of thyroid hormone in Sertoli cell development: a mechanistic hypothesis. *Cell Tissue Res.* **322**, 133–140.

Hsu, B.-Y., Dijkstra, C., Darras, V. M., de Vries, B. and Groothuis, T. G. G. (2017). Maternal thyroid hormones enhance hatching success but decrease nestling body mass in the rock pigeon (Columba livia). *Gen. Comp. Endocrinol.* **240**, 174–181.

Hsu, B.-Y., Doligez, B., Gustafsson, L. and Ruuskanen, S. (2019). Transient growth-enhancing effects of elevated maternal thyroid hormones at no apparent oxidative cost during early postnatal period. *J. Avian Biol.*

Isaksson, C., Örnborg, J., Stephensen, E. and Andersson, S. (2005). Plasma Glutathione

and Carotenoid Coloration as Potential Biomarkers of Environmental Stress in Great Tits. *EcoHealth* **2**, 138–146.

Keshavarz, K. and Quimby, F. W. (2002). An Investigation of Different Molting Techniques with an Emphasis on Animal Welfare. *J. Appl. Poult. Res.* **11**, 54–67.

Kikuyama, S., Kawamura, K., Tanaka, S. and Yamamoto, K. (1993). Aspects of Amphibian Metamorphosis: Hormonal Control. In *International Review of Cytology*, pp. 105–148. Elsevier.

Korevaar, T. I. M., Muetzel, R., Medici, M., Chaker, L., Jaddoe, V. W. V., de Rijke, Y. B., Steegers, E. A. P., Visser, T. J., White, T., Tiemeier, H., et al. (2016). Association of maternal thyroid function during early pregnancy with offspring IQ and brain morphology in childhood: a population-based prospective cohort study. *Lancet Diabetes Endocrinol.* **4**, 35–43.

Krassas, G. E., Poppe, K. and Glinoer, D. (2010). Thyroid Function and Human Reproductive Health. *Endocr. Rev.* **31**, 702–755.

Lenth, R. (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.3.2.

Lilley, T. M., Ruokolainen, L., Meierjohann, A., Kanerva, M., Stauffer, J., Laine, V. N., Atosuo, J., Lilius, E.-M. and Nikinmaa, M. (2013). Resistance to oxidative damage but not immunosuppression by organic tin compounds in natural populations of Daubenton's bats (Myotis daubentonii). *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* **157**, 298–305.

Marshall, D. J. and Uller, T. (2007). When is a maternal effect adaptive? *Oikos* **116**, 1957–1963.

McNabb, F. M. A. (2007). The Hypothalamic-Pituitary-Thyroid (HPT) Axis in Birds and Its Role in Bird Development and Reproduction. *Crit. Rev. Toxicol.* **37**, 163–193.

McNabb, F. M. A. and Darras, V. M. (2015). Thyroids. In *Sturkie's Avian Physiology*, pp. 535–547. Elsevier.

McNabb, F. M. A., Scanes, C. G. and Zeman, M. (1998). Endocrine control of development. In *Avian Growth and Development: Evolution Within the Altricial-precocial Spectrum*, pp. 174–202. New York: Starcj, J.M., Ricklefs, R.E.

Medici, M., Timmermans, S., Visser, W., de Muinck Keizer-Schrama, S. M. P. F., Jaddoe, V. W. W., Hofman, A., Hooijkaas, H., de Rijke, Y. B., Tiemeier, H., Bongers-Schokking, J. J., et al. (2013). Maternal Thyroid Hormone Parameters during Early Pregnancy and Birth Weight: The Generation R Study. *J. Clin. Endocrinol. Metab.* **98**, 59–66.

Morreale de Escobar, G., Obregon, M. and Escobar del Rey, F. (2004). Role of thyroid hormone during early brain development. *Eur. J. Endocrinol.* U25–U37.

Mousseau, T. A. and Fox, C. W. (1998). *Maternal Effects As Adaptations*. New York: Oxford University Press.

Muriel, J., Perez-Rodriguez, L., Puerta, M. and Gil, D. (2015). Diverse dose-response effects of yolk androgens on embryo development and nestling growth in a wild passerine. *J. Exp. Biol.* **218**, 2241–2249.

Okuliarová, M., Škrobánek, P. and Zeman, M. (2007). Effect of Increasing Yolk Testosterone Levels on Early Behaviour in Japanese Quail Hatchlings. *Acta Vet. Brno* **76**, 325–331.

Pfannkuche, K. A., Gahr, M., Weites, I. M., Riedstra, B., Wolf, C. and Groothuis, T. G. G. (2011). Examining a pathway for hormone mediated maternal effects – Yolk testosterone affects androgen receptor expression and endogenous testosterone production in young chicks (Gallus gallus domesticus). *Gen. Comp. Endocrinol.* **172**, 487–493.

Pinheiro, J. C., Bates, D. M., Sarkar, D. and R Core Team (2018). nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-137.

Podmokła, E., Drobniak, S. M. and Rutkowska, J. (2018). Chicken or egg? Outcomes of experimental manipulations of maternally transmitted hormones depend on administration method - a meta-analysis: Maternal hormones and manipulation methods. *Biol. Rev.* **93**, 1499–1517.

Prati, M., Calvo, R., Morreale, G. and Morreale de Escobar, G. (1992). L-thyroxine and 3,5,3'-triiodothyronine concentrations in the chicken egg and in the embryo before and after the onset of thyroid function. *Endocrinology* **130**, 2651–2659.

Raine, J. C., Cameron, C., Vijayan, M. M., Lamarre, J. and Leatherland, J. F. (2004). The effect of elevated oocyte triiodothyronine content on development of rainbow trout embryos and expression of mRNA encoding for thyroid hormone receptors. *J. Fish Biol.* **65**, 206–226.

Rainio, M. J., Kanerva, M., Salminen, J.-P., Nikinmaa, M. and Eeva, T. (2013). Oxidative status in nestlings of three small passerine species exposed to metal pollution. *Sci. Total Environ.* **454–455**, 466–473.

Rainio, M. J., Eeva, T., Lilley, T., Stauffer, J. and Ruuskanen, S. (2015). Effects of early-life lead exposure on oxidative status and phagocytosis activity in great tits (Parus major). *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* **167**, 24–34.

Reis, L. H., Gama, L. and Soares, M. (1997). Effects of short storage conditions and broiler breeder age on hatchability, hatching time, and chick weights. *Poult. Sci.* **76**, 1459–1466.

Rubin, C.-J., Zody, M. C., Eriksson, J., Meadows, J. R. S., Sherwood, E., Webster, M. T., Jiang, L., Ingman, M., Sharpe, T., Ka, S., et al. (2010). Whole-genome resequencing reveals loci under selection during chicken domestication. *Nature* **464**, 587–591.

Ruuskanen, S. and Hsu, B.-Y. (2018). Maternal Thyroid Hormones: An Unexplored Mechanism Underlying Maternal Effects in an Ecological Framework. *Physiol. Biochem. Zool.* **91**, 904–916.

Ruuskanen, S., Darras, V. M., Visser, M. E. and Groothuis, T. G. G. (2016). Effects of experimentally manipulated yolk thyroid hormone levels on offspring development in a wild bird species. *Horm. Behav.* **81**, 38–44.

Sekimoto, K., Imai, K., Suzuki, M., Takikawa, H., Hoshino, N. and Totsuka, K. (1987). Thyroxine-Induced Molting and Gonadal Function of Laying Hens. *Poult. Sci.* **66**, 752–756.

Tsuyoshi, H. and Wada, M. (1992). Termination of LH secretion in Japanese quail due to high- and low-temperature cycles and short daily photoperiods. *Gen. Comp. Endocrinol.* **85**, 424–429.

Van Herck, S. L. J., Geysens, S., Delbaere, J., Tylzanowski, P. and Darras, V. M. (2012). Expression profile and thyroid hormone responsiveness of transporters and deiodinases in early embryonic chicken brain development. *Mol. Cell. Endocrinol.* **349**, 289–297.

Van Herck, S. L. J., Delbaere, J., Bourgeois, N. M. A., McAllan, B. M., Richardson, S. J. and Darras, V. M. (2015). Expression of thyroid hormone transporters and deiodinases at the brain barriers in the embryonic chicken: Insights into the regulation of thyroid hormone availability during neurodevelopment. *Gen. Comp. Endocrinol.* **214**, 30–39.

von Engelhardt, N. and Groothuis, T. G. G. (2011). Maternal Hormones in Avian Eggs. In *Hormones and Reproduction of Vertebrates*, pp. 91–127. Elsevier.

Wada, H. (2008). Glucocorticoids: Mediators of vertebrate ontogenetic transitions. *Gen. Comp. Endocrinol.* **156**, 441–453.

Walpita, C. N., Van der Geyten, S., Rurangwa, E. and Darras, V. M. (2007). The effect of 3,5,3'-triiodothyronine supplementation on zebrafish (Danio rerio) embryonic development and expression of iodothyronine deiodinases and thyroid hormone receptors. *Gen. Comp. Endocrinol.* **152**, 206–214.

Watanabe, Y., Grommen, S. V. H. and De Groef, B. (2016). Corticotropin-releasing hormone: Mediator of vertebrate life stage transitions? *Gen. Comp. Endocrinol.* **228**, 60–68.

Wilson, C. M. and McNabb, F. M. A. (1997). Maternal Thyroid Hormones in Japanese Quail Eggs and Their Influence on Embryonic Development. *Gen. Comp. Endocrinol.* **107**, 153–165.

Wood, S. N. (2017). Generalized Additive Models: An Introduction with R, Second Edition. *CRC Press*.

Zuur, A. F., Ieno, E. N., Walker, N. J., Savaliev, A. A. and Smith, G. M. (2009). *Mixed effects models and extensions in ecology with R*. New York, NY: Springer.

Figure legends

Figure 1: Percentage of hatching success according to yolk TH manipulation treatments: CO (N = 40), T_3 (N = 39), T_4 (N = 39), T_3T_4 (N = 40). CO = control, T_4 (thyroxine) = injection of T_4, T_3 (triiodothyronine) = injection of T_3, T_3T_4 = injection of T_3 and T_4.

Figure 2: Growth curves in body mass of Japanese quails hatching from eggs treated with either T_3, T_4, a combination of both hormones, or a control solution. See Fig. 1 for a description of the treatments. Each line represents an individual bird, while thick coloured lines represent mean values. A: Growth curve according to yolk TH manipulation. N = 7 CO, 11 T_3, 18 T_4 and 21 T_3T_4. B: Growth curve according to sex. N = 29 females and 28 males.

Figure 3: Primary moult score in 7-month old Japanese quails according to yolk TH manipulation treatments: CO (N = 7), T_3 (N = 11), T_4 (N = 16), T_3T_4 (N = 20). See Fig. 1 for a description of the treatments. Measures were taken once a week after switching from long photoperiod (16L:8D) to short photoperiod (8L:16D, switch = time point 0 on x-axis). Each line represents an individual bird, while thick coloured lines represent group mean values.

Figure 4: MDA concentration according to yolk TH manipulation treatments, samples from two ages pooled: CO (N = 7 individuals), T_3 (N = 11), T_4 (N = 17), T_3T_4 (N = 20). See Fig. 1 for a description of the treatments.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Tables

Table 1: Loadings of the different antioxidant biomarkers on the principal components 1 and 2.

Factor loadings	PC1 (34.0%)	PC2 (26.2%)
CAT	-0.49	0.14
SOD	0.20	-0.71
GST	-0.65	-0.10
GP	0.04	-0.63
tGSH	-0.60	-0.26

Table 2: Results of the Generalised Additive Mixed Models (GAMMs) on body mass growth, with sex and treatment fitted either as intercept, curve shape or both (all combinations tested). A total of 12 GAMMs were fitted and ranked based on their AIC, from the lowest to the highest. Weight: Akaike's weights.

Model	Intercept	Curve shape	ΔAIC	df	Weight
1	Sex	Sex	0.0	11	0.8430
8	Treatment + sex	Sex	3.5	14	0.1497
3	-	Sex	9.9	10	0.0061
2	Treatment	Sex	13.2	13	0.0012
11	Sex	-	77.6	9	<0.001
9	Treatment + sex	-	81.6	12	<0.001
12	-	-	91.2	8	<0.001
10	Treatment	-	95.0	11	<0.001
5	Sex	Treatment	147.9	15	<0.001
7	Treatment + sex	Treatment	151.7	18	<0.001
6	-	Treatment	161.2	14	<0.001
4	Treatment	Treatment	165.5	17	<0.001

Table 3: 95% confidence intervals of the predictors in the top-ranked models according to AIC values (see Tables 2, S1 and S2). Predictors in bold have confidence intervals that do not overlap with 0. For the intercept, the reference groups are female and CO for the predictors sex and treatment, respectively.

Curve parameter	Predictors	Lower limit	Estimate	Upper limit
(A) Body mass (Model 1)				
Intercept	**Sex (M)**	-19.7	-12.6	-5.5
Curve shape	**Sex (F)**	9.9	20.0	30.0
Curve shape	**Sex (M)**	14.3	24.5	34.7
(B) Wing length (Model 11)				
Intercept	**Sex (M)**	-2.3	-1.2	-0.1
Curve shape	**Age**	26.4	28.7	31.0
(C) Tarsus length (Model 10)				
Intercept	Treatment (T_3)	-0.8	0.02	0.8
Intercept	**Treatment (T_3T_4)**	-1.5	-0.8	-0.1
Intercept	Treatment (T_4)	-1.3	-0.6	0.2
Curve shape	**Age**	10.5	11.1	11.8
Tarsus length (Model 9)				
Intercept	Treatment (T_3)	-0.9	-0.07	0.7
Intercept	**Treatment (T_3T_4)**	-1.5	-0.8	-0.1
Intercept	Treatment (T_4)	-1.4	-0.6	0.1
Intercept	Sex (M)	-0.8	-0.3	0.3
Curve shape	**Age**	10.5	11.1	11.7