Elevated exposure to prenatal thyroid hormones affects embryonic mortality but has no

clear effects into adulthood

Tom Sarraude*[1,2], Bin-Yan Hsu[1], Ton G.G. Groothuis[2], Suvi Ruuskanen[1]

[1] Department of Biology, University of Turku, Turku, Finland

[2] Groningen Institute for Evolutionary Life Sciences, University of Groningen,

Groningen, the Netherlands

[*] Corresponding author: Tom Sarraude, t.sarraude@rug.nl

Keywords: maternal hormones; thyroid hormones; avian growth; hatching success; Japanese
quails; life-history strategies.

Abstract

Maternal thyroid hormones (THs) are known to be crucial in embryonic development in humans,

but their influence on other, especially wild, animals remains poorly understood. So far, the

studies that experimentally investigated the consequences of maternal THs focused on short-term

effects, while early organisational effects with long-term consequences, as shown for other

prenatal hormones, could also be expected. In this study, we aimed at investigating both the

short- and long-term effects of prenatal THs in a bird species, the Japanese quail *Coturnix*

japonica. We experimentally elevated yolk TH content (the prohormone T_4, and its active

metabolite T_3, as well as a combination of both hormones). We analysed hatching success,

embryonic development, offspring growth and oxidative stress as well as their potential

organisational effects on reproduction, moult, and oxidative stress in adulthood. We found that

eggs injected with both hormones had a higher hatching success compared with control eggs,

suggesting conversion of T_4 into T_3 by the embryo. We detected no other clear short-term or long-

term effects of yolk THs. These results suggest that yolk thyroid hormones are important in the

embryonic stage of precocial birds, but other short- and long-term consequences remain unclear.

Research on maternal thyroid hormones will greatly benefit from studies investigating how

embryos use and respond to this maternal signalling. Long-term studies on prenatal THs in other

taxa in the wild are needed for a better understanding of this hormone-mediated maternal

pathway.

Introduction

Maternal effects represent all the non-genetic influences of a mother on her offspring and have received increasing attention in evolutionary and behavioural ecology. Through maternal effects, mothers can influence the fitness of their progeny by adapting their phenotype to expected environmental conditions ("adaptive maternal effects" in Marshall and Uller, 2007; Mousseau and Fox, 1998), and this view is now also incorporated in the human disease literature (Gluckman, Hanson & Spencer, 2005). Maternal hormones transferred to the offspring can mediate important maternal effects. Historically, research on maternal hormones has mostly focused on steroid hormones (Groothuis et al., 2005; von Engelhardt & Groothuis, 2011). Recently, thyroid hormones (THs) also received attention in the context of hormone-mediated maternal effects (Ruuskanen & Hsu, 2018).

Thyroid hormones (THs) are metabolic hormones produced by the thyroid gland and are present in two main forms: the prohormone thyroxine (T_4) and the biologically active form triiodothyronine (T_3). THs play a crucial role in various aspects of an individual's life, e.g. development, metabolism and reproduction, across vertebrates, including humans (Morreale de Escobar, Obregon & Escobar del Rey, 2004; Krassas, Poppe & Glinoer, 2010). In humans, physiological variation of maternal THs (i.e. no clinical symptoms in both mothers and foetuses) is found to be associated with infant birth weight and IQ in older children (Medici et al., 2013; Korevaar et al., 2016). In birds as well, THs in general play a role in brain development and neuronal turnover (reviewed in McNabb, 2007). THs control the endothermic heat production, and are therefore important in thermoregulation in juveniles and adults (McNabb & Darras, 2015).

THs can act, in concert with other hormonal axes, as mediators of life stage transitions

across vertebrates (reviewed in Watanabe et al., 2016). The interaction between thyroid hormones

and corticosteroids on amphibian metamorphosis is a well-known example of such effect on life

stage transition (Kikuyama et al., 1993; Wada, 2008). THs are involved in gonadal development,

and hyperthyroidism tends to fasten maturation (Holsberger & Cooke, 2005), and coordinate the

transition between reproduction and moult (McNabb and Darras, 2015). Administration of

exogenous THs is known to stop egg laying and induce moult in birds (Sekimoto et al., 1987;

Keshavarz & Quimby, 2002). THs are also involved in photoperiodic control in seasonal breeding

(Dardente, Hazlerigg & Ebling, 2014). For example, thyroidectomised starlings transferred to

long photoperiods became insensitive to future changes in photoperiod, and short photoperiod did

not induce gonadal regression (Dawson, 1993).

While there has been recent research effort on the influence of maternal THs on offspring

traits across vertebrate taxa, there are still substantial knowledge gaps. First, research on maternal

thyroid hormones up to date has mainly investigated the short-term effects of prenatal THs on

developing fish (Brown et al., 1988; Raine et al., 2004) and amphibians (Duarte-Guterman et al.,

2010; Fini et al., 2012) and pre-fledging birds (Ruuskanen et al., 2016; Hsu et al., 2017, 2019;

Sarraude et al., 2020, *in press*). So far, only a study on rock pigeons has looked at the influence of

yolk THs on post-fledging survival and found no effect (Hsu et al., 2017). None of these studies

in any taxa investigated the potential organisational effects of prenatal THs on life-history stage

transitions in adult life. Early exposure to elevated THs may affect the hypothalamic-pituitary-

thyroid (HPT) axis (humans and mice: Alonso et al., 2007; Srichomkwun et al., 2017; Anselmo et

al., 2019), via epigenetic modifications for example, such as those induced by adverse early life

conditions (Jimeno et al., 2019) or yolk testosterone (Bentz, Becker & Navara, 2016).

Second, previous studies on prenatal THs in birds focused only on altricial species (great

tits, Ruuskanen et al., 2016; rock pigeons, Hsu et al., 2017; collared flycatchers, Hsu et al., 2019,

pied flycatchers, Sarraude et al., 2020, *in press*). Embryonic development differs substantially

between altricial and precocial species. In the latter, embryonic development is more advanced

than in the former. In addition, precocial embryos start their endogenous production of TH

around mid-incubation, considerably earlier than their altricial counterparts, in which endogenous

TH production begins only after hatching (McNabb, Scanes & Zeman, 1998). While embryonic

hormone production may limit the influence of maternal hormones, prenatal hormones have been

shown to affect chick endogenous production and sensitivity (Pfannkuche et al., 2011). Overall,

exposure to maternal hormones may be of different importance in these two developmental

modes.

Third, previous research has studied the effects of T_3 only (Raine et al., 2004; Walpita et

al., 2007; Fini et al., 2012) or a combination of T_3 and T_4 (Ruuskanen et al., 2016; Hsu et al.,

2017, 2019; Sarraude et al., 2020, *in press*), where the effects of the two forms cannot be

separated. Although T_3 is the biologically active form that binds to the receptors, both T_3 and T_4

are deposited in eggs (Prati et al., 1992) and T_4 may be converted to T_3 via deiodinases from the

mother or the developing embryo (Van Herck et al., 2015) or may still exert non-genomic actions

(reviewed in Davis et al., 2016). Manipulating yolk T_4 and T_3 independently would help

understanding the relative contribution of these two hormones.

In this study, we aimed at assessing the effects of maternal THs on development and life-

history traits in a precocial bird species, the Japanese quail (*Coturnix japonica*). To do so,

manipulated eggs received either an injection of T_4 or T_3 separately, a combination of both

hormones, or a control injection of the vehicle saline solution. This design allowed us to explore

the effects of T_4 and T_3 separately. The elevation in yolk THs remained within the natural range

of this species, a crucial condition to obtain relevant results for an eco-evolutionary context. We

measured traits known to be influenced by circulating and yolk THs: hatching success, age at

embryonic mortality, growth, transition between life-history stages (i.e., reproductive state and

moult) and oxidative stress. First, we hypothesise that elevation of yolk THs in Japanese quails

positively affects hatching success, as found in two studies on collared flycatchers and rock

pigeons (Hsu et al., 2017; Hsu et al., 2019, but see Ruuskanen et al., 2016 and Sarraude et al.,

2020, *in press*). Second, elevation of yolk THs is predicted to increase the proportion of well-

developed embryos before hatching, as found in rock pigeons (Hsu et al., 2017). We therefore

looked at the age at mortality in unhatched eggs. Third, we expect elevated yolk THs to affect

chick growth (in body mass, tarsus and wing length) either positively (Wilson & McNabb, 1997;

Hsu et al., 2019; weak effect in Sarraude et al., 2020, *in press*), negatively (Hsu et al., 2017), or in

a sex-specific manner (Ruuskanen et al., 2016). Prenatal THs may exert most of their effects in

the offspring early life; this is why we separately tested both early morphological traits and the

growth curve. Similarly, we also independently analysed morphological traits at adulthood, as

these traits may affect the fitness of an individual. For example, small adult females may lay

smaller eggs and larger males may be more dominant. Fourth, we predict that yolk THs will have

organisational effects on life-history stage transitions; that is, age at sexual maturity and male

gonadal regression (using cloacal gland size as a proxy), and moult when birds are exposed to

short photoperiod. Based on the literature mentioned above we expect elevated yolk THs to

advance the timing of puberty, gonadal regression and moult. The rate of moult should also be

influenced, with birds receiving experimental TH elevation moulting faster. Previous studies have

reported that gravid female three-spined sticklebacks (*Gasterosteus aculeatus*) exposed to

predatory cues produced eggs with higher corticosterone (Giesing et al., 2011), disturbed

embryonic transcriptome (Mommer & Bell, 2014), offspring with altered anti-predator behaviour

(Giesing et al., 2011) and modified cortisol response in adulthood (Mommer & Bell, 2013). We

may therefore expect elevated yolk THs to similarly induce long-term behavioural changes in

response to environmental cues (i.e. photoperiod), via organising effects during the embryonic

development. We also explored the effects of yolk THs on reproductive investment in females,

another important fitness aspect. Finally, yolk THs may increase oxidative stress due to their

stimulating effects on metabolism.

Material and Methods

Parental generation and egg collection

The parental generation was composed of adult Japanese quails provided by Finnish private local

breeders that were kept in two acclimated rooms. Twenty-four breeding pairs were formed by

pairing birds from different breeders. Individuals were identified using metal leg bands. The floor

was covered with 3–5cm sawdust bedding. A hiding place, sand and calcium grit were provided.

Each pair was housed in indoor aviary dividing into pens of 1 m² floor area. The temperature was

set to 20°C with a 16L:8D photoperiod (light from 06.00 to 22.00). Food (Poultry complete feed,

"Kanan Paras Täysrehu", Hankkija, Finland) was provided *ad libitum* and water was changed

every day.

Pairs were monitored every morning to collect eggs for 7 days. Eggs were individually

marked (non-toxic marker), weighed and stored in a climate-controlled chamber at 15°C and 50%

relative humidity. On the last day of collection, an average of 6.6 eggs per pair (range = 4–8 eggs)

were injected with a solution (see next section).

Preparation of the solution, injection procedure and incubation

The preparation of hormone solution and the procedure of injection were based on previous

studies (Ruuskanen et al., 2016; Hsu et al., 2017). In brief, crystal T_4 (L-thyroxine, $\geq$ 98% HPCL,

CAS number 51-48-9, Sigma-Aldrich) and T_3 (3,3',5-triiodo-L-thyronine, > 95% HPCL, CAS

number 6893-02-3, Sigma-Aldrich) were first dissolved in 0.1M NaOH and then diluted in 0.9%

NaCl. The injection of thyroid hormones resulted in an increase of two standard deviations (T_4 =

8.9 ng/egg; T_3 = 4.7 ng/egg), a recommended procedure for hormone manipulation within the

natural range (Ruuskanen et al., 2016; Hsu et al., 2017; Podmokła, Drobniak & Rutkowska,

2018). The control solution (CO) was a saline solution (0.9% NaCl). The concentrations of the

hormone solutions were based on previous measurements of 15 eggs from the same flock (T_4

content per egg (SD) = 15.3 (4.4) ng, T_3 content per egg (SD) = 7.6 (2.3) ng).

Hormone injections were performed at room temperature in a laminar hood. Eggs were

put sideways, allowing yolks to float up to the middle position. Before injection, the shell was

disinfected with a cotton pad dipped in 70% EtOH. We used a 27G needle (BD Microlance ™) to

pierce the eggshell and then used a 0.3 ml syringe to deliver 50 µl of the respective hormone

solution or control. After injection, the hole was sealed with a sterile plaster (OPSITE Flexigrid,

Smith&Nephew).

In total, 158 eggs were injected and divided as follows over the treatments: T_3 treatment

(N = 39); T_4 treatment (N = 39); T_3+T_4 treatment (N = 40); and control, CO (N = 40). To balance

the genetic background of the parents and the effect of storage, each egg laid by the same female

was sequentially assigned to a different treatment and the order of treatments was rotated among

females. After injection, eggs were placed in an incubator at 37.8°C and 55% relative humidity.

Until day 14 after starting incubation, eggs were automatically tilted every hour by 90°. On day

14, tilting was halted and each egg was transferred to an individual container to monitor which

chick hatched from which egg. On day 16 after injection, (normal incubation time = 17 days), the

temperature was set to 37.5°C and the relative humidity to 70%. Eggs were checked for hatching

every 4 hours from day 16 onwards. Four days after the first egg hatched, all unhatched eggs

were stored in a freezer and dissected to determine the presence of an embryo. The age of

developed embryos was assessed according to Ainsworth et al. (2010).

Rearing conditions of the experimental birds

In total, 66 chicks hatched (N = 10 CO, 15 T_3, 20 T_4 and 21 T_3T_4), yielding a rather low

overall hatching success (ca. 40%). Among the unhatched eggs, 33.7% (31 out of 92) had no

developed embryos, thus discarding the unfertilised eggs gives an overall hatching success of ca.

51%. Previous studies on Japanese quails have reported comparable hatching success, even in

unmanipulated eggs (e.g. 40% in Okuliarová, Škrobánek & Zeman, 2007; ca. 60% in Pick et al.,

2016 and in Stier, Metcalfe & Monaghan, 2019). In addition, the injection procedure itself is also

known to reduce hatching success to some extent (Groothuis & von Engelhardt, 2005). Twelve

hours after hatching, the chicks were marked by a unique combination of coloured rings and nail

coding and transferred to two cages of 1 m² floor area and ca. 30 cm height (ca. 30 chicks/cage,

sex and treatments mixed together). The chicks were provided with heating mats and lamps as

extra heat sources for the first two weeks. The chicks were fed with sieved commercial poultry

feed ("Punaheltta paras poikanen", Hankkija, Finland), and provided with Calcium and bathing

sand. Two weeks after hatching, the chicks were separated in four 1 m² cages (ca. 30 cm high) of

about 16 individuals. Around 3 weeks after hatching, coloured rings were replaced by unique

metal rings. On week 4 after hatching, birds were transferred to eight pens of 1 m² floor area

(average of 7.1 birds/pen, range = 4–9), under the same conditions as the parents. Around the age

of sexual maturity (ca. 6–8 weeks after hatching), the birds were separated by sex in twelve 1 m²

pens (average of 4.8 birds/pen, range = 4–5). The chicks were under the same photoperiod as the

adults (i.e. 16L:8D).

Monitoring of growth and reproductive maturation

Body mass and wing length were measured twelve hours after hatching. Tarsus was not measured

because it bends easily, resulting in inaccurate measures and potential harm for the young. From

day 3 to day 15, these three traits were monitored every 3 days. From day 15 to day 78 (ca. 12

weeks), chicks were measured once a week. Body mass was recorded using a digital balance to

the nearest 0.1 g. Wing and tarsus lengths were respectively measured with a ruler and a calliper

to the nearest 0.5 mm and 0.1 mm. The sample size for the growth analysis was 7 CO, 11 T_3, 18

T_4 and 21 T_3T_4. From week 6 to week 10, we monitored cloacal gland development and foam

production in 28 males. Cloacal glands were measured every other day with a calliper to the

nearest 0.1 mm as a proxy for testes development and sexual maturation (Biswas et al., 2007).

Foam production (by gently squeezing the cloacal gland) was assessed at the same time and

coded from 0 (no foam) to 3 (high production of foam), as a proxy of cloacal gland function

(Cheng et al., 1989a; Cheng et al., 1989b). The same observer performed all measurements. We

collected eggs produced by 10-week-old females over a 6-day period and recorded their mass to

the nearest 0.1 g. We collected on average 5.7 eggs (range = 4–7) per female from 28 females.

Monitoring of cloacal gland regression and moult

In Japanese quails, exposure to short photoperiod and cold temperature triggers reproductive

inhibition and postnuptial moulting (Tsuyoshi & Wada, 1992). Thyroid hormones are known to

coordinate these two responses (see introduction). When the birds reached the age of ca. 7

months, we exposed birds to short photoperiod (8L:16D, i.e., light from 08.00 to 16.00) with a

12:12-h cycle of normal (20°C) and low (9°C) temperature (low temperature was effective from

18.00 to 06.00). Cloacal gland regression (as a proxy for testes regression) was monitored every

other day for 2 weeks with a calliper by measuring the width and length to obtain the area of the

gland to the nearest 0.1 mm² (N = 26 males). Primary moult was recorded from a single wing by

giving a score to each primary from 0 (old feather) to 5 (new fully-grown feather) following Ginn

and Melville (1983) (N = 54 males and females). The total score of moulting was obtained by

adding the score of all feathers. The sample size for the moult analysis was 7 CO, 11 T_3, 16 T_4

and 20 T_3T_4.

Oxidative status biomarker analyses

Two blood samples were drawn, when birds were 2 weeks (N = 51 chicks) and 4 months old (N =

49 adults), respectively. The sample size per treatment was 7 CO, 11 T_3, 17 T_4 and 20 T_3T_4. 200

μl of blood was collected from the brachial vein in heparinized capillaries and directly frozen in

liquid nitrogen. Then, the samples were stored at -80°C until analyses. We measured various

biomarkers of antioxidant status; the antioxidant glutathione (tGSH), the ratio of reduced and

oxidised glutathione (GSH:GSSG) and activity of the antioxidant enzymes glutathione

peroxidase (GPx), catalase (CAT) and superoxide dismutase (SOD) from the blood. It is

important to measure multiple biomarkers of oxidative and antioxidant status for a broader

understanding of the mechanism. Also, the interpretation of the results is more reliable if multiple

markers show similar patterns. Of the measured biomarkers, the ratio of GSH:GSSG represents

the overall oxidative state of cells and consequently, deviations in this ratio is often used as an

indicator of oxidative stress (Hoffman, 2002; Isaksson et al., 2005; Lilley et al., 2013; Rainio et

al., 2013; Halliwell & Gutteridge, 2015). GPx enzymes catalyse the glutathione cycle, whereas

CAT and SOD directly regulate the level of reactive oxygen species (ROS) (Ercal, Gurer-Orhan

& Aykin-Burns, 2001; Halliwell & Gutteridge, 2015). The methodology for measuring each

biomarker is described in detail in Rainio et al. (2015). All analyses were conducted blindly of

the treatment. Briefly, the samples were analysed using a microplate reader (EnVision,

PerkinElmer-Wallac, Finland). All antioxidant and enzyme activities were measured in triplicate

(intra-assay coefficient of variability [CV] < 10% in all cases) using 96-(CAT) or 384-well (GPx,

SOD, tGSH and GSH:GSSG) microplates. Three control samples were used with each plate, to

be able to correct for inter-assay precision with the ratio specific to the particular plate. Overall

protein concentration (mg/ml) was measured according to the Bradford method (Bradford, 1976)

using BioRad stock (BioRad, Finland) diluted with dH_2O (1:5) and BSA (bovine serum albumin,

1 mg/ml) (Sigma Chemicals, USA) as a standard. GPx-assay was conducted using Sigma CGP1

kit, CAT-assay using SigmaCAT100 kit and SOD-assay using Fluka 19160 SOD determination

kit. Total GSH and the ratio of GSH:GSSG were measured with the ThioStar® glutathione

detection reagent (Arbor Assays, USA) according to kit instructions, using reduced glutathione as

a standard (Sigma Chemicals, USA).

Ethics

The study complied with Finnish regulation and was approved by the Finnish Animal Experiment

Board (ESAVI/1018/04.10.07/2016). In case of signs of harassment or disease, birds were placed

in quarantine and monitored daily until they had recovered. Criteria for humane endpoints were

defined as follow: passive behaviour, loss of appetite, loss of 30% of body weight, moving

abnormally, trouble breathing. If we observed no clear improvement after two days, we would

consult the veterinarian. A bird would be euthanised if it does not show signs of improvement in

the next two days, though some judgement can be applied based on the alleged cause. One male

was euthanised before the end of the experiment due to severe head injury. At the end of the

experiment, all birds were euthanised by decapitation for collection of tissue samples (not used in

this study).

Statistical analysis

Data were analysed with the software R version 3.5.3 (R core team, 2019). In this study, two

different statistical approaches were used: null-hypothesis testing with Generalised Linear Mixed

Models (GLMMs) and Linear Mixed Models (LMMs), and multimodel inference with

Generalised Additive Mixed Models (GAMMs). GAMMs were used to analyse the data on body

and cloacal gland growth to account for its non-linear pattern (see *Growth*). In this analysis, we

preferred multimodel inference as GAMMs generate many candidate models that cannot be

directly compared (e.g., by the Kenward-Roger approach). Instead, candidate models were

ranked based on their Akaike Information Criterion (AIC) values. Models with a $\Delta AIC \leq 2$ from

the top-ranked model were retained in the set of best models. Akaike weights of all models were

calculated following (Burnham & Anderson, 2002), and evidence ratios of the top-ranked models

were calculated as the weight of a model divided by the weight of the null model (Burnham,

Anderson & Huyvaert, 2011). To estimate the effect of the predictors, we computed the 95%

confidence intervals from the best models using the *nlme* package (Pinheiro et al., 2018).

GLMMs and LMMs were fitted using the R package *lme4* (Bates et al., 2015), and GAMMs were

fitted using the package *mgcv* (Wood, 2017). P-values for GLMMs were obtained by parametric

bootstrapping with 1,000 simulations and p-values for LMMs were calculated by model

comparison using Kenward-Roger approximation, using the package *pbkrtest* in both cases

(Halekoh & Højsgaard, 2014). Post-hoc Tukey analyses were conducted with the package

multcomp (Hothorn et al., 2008). Model residuals were checked visually for normality and

homoscedasticity. Covariates and interactions were removed when non-significant ($\alpha = 0.05$).

Effect size calculations and statistical power analyses were performed using t-tests for

independent means with GPower (Faul et al., 2009). Estimated marginal means and standard

errors (EMMs ± SE) were extracted from the models, using the package *emmeans* (Lenth, 2019)

and pooled standard deviations were calculated based on the standard errors of the EMMs and

group sample sizes. These parameters (EMMs and pooled SD) were entered in the GPower

software to calculate Cohen's d index and achieved power.

Hatching success

To analyse hatching success, each egg was given a binary score: 0 for unhatched egg and 1 for

hatched egg. A GLMM was fitted with a binomial error distribution (logit link) and mother

identity as a random intercept and the 4-level treatment as the predictor. Egg mass might affect

hatchability and was therefore added as a covariate in both models. The potential effect of storage

duration on hatchability (Reis, Gama & Soares, 1997) was accounted for by including laying

order as a covariate in both models. This covariate allowed us to control for the age of the egg as

well.

Duration of embryonic period, age at embryonic mortality and early morphological traits

Duration of embryonic period and early morphological traits (mass and wing length at hatching,

and tarsus length at day 3) were modelled with separate LMMs. Treatment, sex of the individuals

and egg mass were included as fixed factors. Laying order was added as a covariate to account

for potential effects of storage duration on hatching time and on chick weight (Reis, Gama &

Soares, 1997). Mother identity was included as a random intercept.

The data for embryonic age had a skewed distribution and residuals were not normally

distributed and heterogenous, which violated LMM assumptions on residual distribution. We

therefore performed a simple Kruskal-Wallis test.

Growth

As growth curves typically reach an asymptote, we fitted non-linear GAMMs to these curves.

Growth in body mass, tarsus and wing length were analysed in separate GAMMs. Growth was

analysed until week 10 after hatching as all birds appeared to have reached their maximum body

mass and tarsus and wing length. The data are composed of repeated measurements of the same

individuals over time; therefore, we first corrected for temporal autocorrelation between the

measurements using an ARMA(1,1) model for the residuals (Zuur et al., 2009). Second, as

mothers produced several eggs, the models included nested random effects, with measured

individuals nested into mother identity, allowing for random intercepts. GAMMs allow modelling

the vertical shift of the curves (i.e., changes in intercepts) and their shape. Treatment and sex

were included as predictors. A smoothing function for the age of the birds was included to model

the changes in the growth curves, and was allowed to vary by sex or treatment only, or none of

these predictors. The interaction between sex and treatment was not analysed due to low

statistical power. Additive effect of treatment and sex was tested for the intercept but could not be

computed for curve shape. All combinations of the relevant predictors were tested for both shape

parameters (i.e., intercept and curve shape).

Prenatal THs may exert most of their effects in the offspring early life; this is why we

additionally tested early morphological traits apart from the growth curve. Likewise, we also

analysed separately morphological traits at adulthood (ca. 9 weeks old), as these traits may

condition the fitness of an individual. Because of sex differences and low sex-specific sample

sizes, we standardised the measures within sex and regressed the standardised responses against

treatment in a linear regression.

Reproductive maturation, regression and investment

Due to low sample sizes in these sex-specific responses, we could not perform robust statistical

analyses. We therefore present these analyses and results in the supplementary material and only

briefly discuss them.

Oxidative stress

A principal component analysis (PCA) was first performed on measured antioxidant markers

(SOD, CAT, GPx, tGSH and GST), to reduce the number of metrics for subsequent analyses. The

first and the second principal components (PCs) explained together 60.2% of the variance (Table

1). PC1 and PC2 were then used as dependent variables in separate LMMs. LMMs included the

treatment, sex and age of individuals (2 weeks and 4 months old) as fixed factors and the 2-way

interactions between treatment and sex, and treatment and age. Mother and individual identities,

to account for repeated measures, were added as random intercepts. Malondialdehyde (MDA) is a

marker of oxidative damage, which is a different measure from antioxidant activity, and was

therefore analysed in a separate LMM using the same parameters as for PC1 and PC2, adding the

batch of the assay as an additional random intercept. The marker of cell oxidative status

(GSH:GSSG ratio) was analysed with the same model used for PC1 and PC2.

Moult

Two parameters of moult were analysed in separate LMMs: the timing of moult (i.e., the moult

score after one week of short photoperiod), and the rate of moult (i.e., how fast birds moulted).

Both models included treatment and sex as fixed factors, and mother identity as a random

intercept. The rate of moult was tested by fitting an interaction between treatment and age. This

model also included the main effect of age and individual identity, nested within mother identity,

as a random intercept to account for repeated measures. Estimated marginal means and standard

errors (EMMs ± SE) were derived from the model using the package *emmeans* (Lenth, 2019).

Results

Effects of prenatal THs on hatching success and age of embryo mortality

There was a clear effect of elevated prenatal THs on hatching success (GLMM, $p = 0.05$, Fig. 1).

Tukey post-hoc analysis revealed that hatching success in the T_3T_4 group was significantly higher

than in the CO group ($T_3T_4 = 53\%$, $CO = 25\%$, Tukey $z = 2.46$, p = 0.05). The other groups ($T_3 =$

38% and $T_4 = 51\%$) were not different from the control group (all $z < 2.22$ and p > 0.09).

Dissection of the unhatched eggs showed that age of embryo mortality did not differ

between the treatments (Kruskal-Wallis $\chi^2 = 7.22$, df = 3, p = 0.07). Finally, the manipulation of

yolk THs did not affect the duration of embryonic period (LMM, $F_{3,42.0} = 0.57$, p = 0.64, Fig. S1).

Sex of the embryo or egg mass (LMM sex, $F_{1,49.7} = 2.63$, p = 0.11; LMM egg mass, $F_{1,19.3} = 0.01$,

p = 0.92) were also not associated with the duration of the embryonic period. Laying order (i.e.

the effect of storage duration) was not correlated with any of the responses (all p ≥ 0.25)

Effects of prenatal THs on growth

Mass at hatching was not influenced by the elevation of prenatal THs (LMM, $F_{3,35.0} = 0.81$, p =

0.50, Fig. S2). Mass at hatching was positively correlated with egg mass (LMM, Estimate±SE =

0.72±0.10 g, $F_{1,24.1} = 46.9$, p < 0.001). Although we detected no clear differences on early

morphological traits (body mass, wing and tarsus length) due to prenatal THs (all p > 0.12),

calculated effect sizes (Cohen's d) ranged from small to moderate (body mass, range = 0.06–0.26;

wing length, range = 0.20–0.43; tarsus length, range = 0.35–0.38). However, the achieved power

was very low (range = 0.05–0.15). Similarly, adult morphology was not affected by the treatment

(all p > 0.13), but effect sizes indicate small to moderate effects of prenatal THs (body mass,

range = 0.13–0.29; wing length, range = 0.33–0.66; tarsus length, range = 0.09–0.33). Yet again,

achieve statistical power was low (range = 0.05–0.25).

Regarding body mass growth, the top-ranked model showed that the curve shape and the

intercept differ according to sex (Table 2). After 10 weeks, females had a larger body mass than

males (mean±SE females = 214.4±5.7 g, males = 172.4±4.5 g, Fig. 2), which was supported by

the 95% CIs (Table 3). Based on model selection we conclude that the treatment had no effect on

body mass growth (Table 2).

For wing growth, the top-ranked model (ΔAIC $\leq$ 2) included sex in the intercept, while

treatment was not included in the best supported model (Table S1). The 95% CIs (Table 3)

confirmed that males had a lower wing length than females (Fig. S3).

Concerning tarsus growth, the models within ΔAIC $\leq$ 2 included no predictors for the

curve shape but included treatment for the intercept (Table S2). The 95% CIs of the parameter

estimates from these models suggested that there was a slight negative effect of T_3T_4 treatment on

tarsus growth (Table 3, Fig. S4). However, as the estimates were close to 0 (Table 3) and

evidence ratios showed that the model with treatment as a predictor was only 3.5 times more

supported than the null model (Table S2), we conclude that the effect of THs on tarsus length is

likely to be very small. Likewise, the second model for tarsus length included sex as a predictor

for the intercept, but its 95% CIs overlapped with 0 (Table 3). We therefore conclude that sex had

no effect on tarsus growth.

Effects of prenatal THs on postnuptial moult

As expected, birds started to moult soon after being exposed to short photoperiod, with an

average increase of moult score by 6 per week (SE = 0.2, $F_{1,254.0}$ = 827.4, p < 0.001, Fig. 3). The

first moult score (assessed one week after switching to short photoperiod) was not affected by the

treatment (LMM, $F_{3,42.7}$ = 0.36, p = 0.78), but was influenced by sex, with females having a

higher score than male (EMMs $\pm$ SE: female = 21.4 $\pm$ 1.6, male = 7.2 $\pm$ 1.7; LMM $F_{1,45.3}$ = 41.9,

p < 0.001). Yolk TH elevation did not affect the rate of moult (LMM interaction treatment $\times$ time,

$F_{3,251.0}$ = 0.59, p = 0.62, Fig. 3).

Effects of prenatal THs on oxidative stress

The elevation of yolk THs had no effect on PC1 or PC2 of antioxidants at either 2 weeks

("chicks") or 4 months ("adults") old (LMM on PC1, $F_{3,40.3} = 2.40$, p = 0.08; LMM on PC2,

$F_{3,42.2} = 0.92$, p = 0.44, treatment × age, $F < 0.91$, p > 0.44). The age of the birds had a highly

significant effect on PC1, with chicks generally having higher antioxidant capacities (CAT, GST

and tGSH) than adults (LMM, Estimate±SE = -1.34±0.19, $F_{1,49.2} = 52.1$, p < 0.001). All the other

predictors had no effect on either PC1 or PC2 (all $F < 2.93$ and all p > 0.09).

The marker of oxidative damage, MDA, was affected by the elevation of yolk THs

(LMM, $F_{3,43.6} = 3.08$, p = 0.04, Fig. 4). Tukey post-hoc analysis showed that the T4 group had

higher MDA values than the T3 group (Estimate±SE = 0.01±0.004, Tukey contrast p = 0.01), but

none of the groups differed from the control (Tukey p-values > 0.19). However, this result

became non-significant when removing the outlier in the T_4 group (LMM, $F_{3,43.1} = 2.68$, p =

0.06). MDA levels were not affected by the age or the sex of individuals (LMM age, $F_{1,54.4} =$

0.30, p = 0.59; LMM sex, $F_{1,42.0} = 1.47$, p = 0.23).

The marker of cell oxidative balance, GSH:GSSG, was not influenced by the yolk THs

nor by the sex of the birds (LMM treatment, $F_{3,33.0} = 0.85$, p = 0.48; LMM sex, $F_{1,40.6} = 0.57$, p =

0.45). However, chicks had a higher GSH:GSSG ratio than adults (LMM, Estimate±SE =

0.17±0.04, $F_{1,50.0} = 18.3$, p < 0.001).

Discussion

The aim of this experimental study was to investigate the potential short-term and organisational

effects (with long-term consequences) of maternal thyroid hormones (THs) in a precocial species,

the Japanese quail, by experimental elevation of THs in eggs. Our study is the first to investigate

the effects of yolk T_3 and T_4 separately, within the natural range of the study model. In addition

we studied both short- and long-term effects on embryonic development, growth, life stage

transitions and oxidative stress. We detected a positive effect of yolk THs on hatching success.

All other response variables studied were not affected by elevated prenatal THs.

Effects of prenatal THs on hatching success and embryonic development

We found that hatching success increased when the eggs received an injection of both T_4 and T_3.

Previous similar studies reported comparable effects of yolk THs in rock pigeons (Hsu et al.,

2017) and in collared flycatchers (Hsu et al., 2019). In these studies, injections consisted of a

mixture of both T_3 and T_4. Given that only T_3 binds to receptors, these results suggest that

embryos must express deiodinase enzymes to convert T_4 to T_3, and/or yolk may contain

maternally derived deiodinase mRNA, as injection with T_3 only did not differ from control.

Indeed, precocial embryos start to produce endogenous T_3 and deiodinase expression has

previously been characterised in chicken embryos (Darras et al., 2009; Van Herck et al., 2012). In

contrast with our study, two similar studies in altricial species detected no increased hatching

success due to the injection of THs (Ruuskanen et al., 2016; Sarraude et al., 2020, *in press*). The

dissimilarities between the studies may come from inter-specific differences in terms of

utilisation of yolk THs by the embryos or from context-dependent effects (e.g. due to other egg

components). Further comparative and mechanistic studies could help understanding the dynamic

of yolk THs during incubation.

Increased yolk THs did not improve age of embryo mortality. Similar to our study,

Ruuskanen et al. (2016) did not find any difference in the timing of mortality in great tit embryos.

Conversely, the study on rock pigeons found that yolk THs increased the proportion of well-

developed embryos (Hsu et al., 2017). Similarly to our result on hatching success, yolk THs'

effects on embryonic development may differ in a species-specific manner.

Effects of prenatal THs on growth

We found no apparent influence of yolk THs on growth, contrary to our expectations based on the

recent literature. Other comparable studies found either a positive (Hsu et al., 2019; weak effect

in Sarraude et al., 2020, *in press*), a negative (Hsu et al., 2017) or a sex-specific effect

(Ruuskanen et al., 2016) of yolk THs on growth. This notable difference may be due to the

captive conditions experienced by the Japanese quails in our study, with unrestricted access to

food and water. Although the pigeon study also provided ad libitum food, parents still needed to

process food before feeding their nestlings in the form of crop milk, whereas precocial quails

have no such limitation. In addition, the Japanese quail has been domesticated for many

generations, and probably selected for rapid growth for economic reasons. Whole-genome

sequencing in chickens showed that domestication induced a strong positive selection on genes

associated with growth (Rubin et al., 2010). Interestingly, that study also found a strong selection

for a locus associated with thyroid stimulating hormone (TSH) receptor. TSH controls most of the

TH production by the thyroid gland (McNabb & Darras, 2015), and this artificial selection may

overshadow the effects of natural variations of prenatal THs on growth. Besides, the low number

of individuals in the control and T_3 groups (7 and 11, respectively) limited the statistical power to

detect differences between the treatments. Indeed, we were able to detect small to moderate

negative effects of yolk THs on morphological traits at hatching and in adulthood. Such negative

effects, although small, may still be biologically relevant. Repeating the study with a larger

sample size may allow us to ascertain the effects of yolk THs on growth in precocial study

models. Research on the influences of prenatal THs on growth will also benefit from

experimental studies on wild precocial species.

Effects of prenatal THs on postnuptial moult

Short photoperiod in combination with cold temperature triggered primary moult, as expected.

However, we detected no effect of yolk THs on the timing or speed of moult. Thyroid hormones

are important in moult and feather growth (reviewed in Dawson, 2015). For example,

thyroidectomised birds fail to moult after being exposed to long photoperiods (Dawson, 2015). In

addition, thyroidectomised nestling starlings failed to grow normal adult plumage and grown

feathers presented an abnormal structure (Dawson et al., 1994). By removing the thyroid gland,

these two studies implemented extreme pharmacological protocols that differ drastically from our

injection of physiological doses. In addition, our experimental design, increasing TH exposure

(vs decreased TH exposure in the above-mentioned studies), may have different consequences.

For example, there may be a threshold above which any, additional hormones may not affect

moult.

Overall, our results show no support for the hypothesis of organising effect of prenatal

THs on life stage transitions. Yet, due to small sample sizes in sex-specific analyses (i.e., male

gonadal maturation and regression, and female reproductive investment), there remains a

relatively high uncertainty about the potential organising effects of prenatal THs. Replicate

studies with larger samples sizes and different study models will reduce this uncertainty.

Effects of prenatal THs on oxidative stress

In contrast to our predictions, elevated yolk THs did not affect oxidative status during chick or

adult phase. We found no changes in antioxidant activities in relation to yolk THs and no

imbalance in the oxidative cell status. Nevertheless, T_4 birds had a higher level of oxidative

damage on lipids than T_3 birds, but this was a weak effect driven by one outlier. The lack of

effects on chick oxidative status among the treatment groups could be explained by the absence

of treatment effects on growth, given that high growth rates usually result in higher oxidative

stress and damage (e.g. Alonso-Alvarez et al., 2007). In turn, the lack of treatment effects on

adult oxidative status may suggest no organisational effects of prenatal THs on adult metabolism.

Two recent studies in altricial species also found no influence of yolk THs on nestling oxidative

stress (Hsu et al., 2019; Sarraude et al., 2020, *in press*), yet telomere length, a biomarker of aging

was affected (A Stier et al., 2020, unpublished data). Our study shows for the first time that

prenatal THs have no influence on adult oxidative stress either. The previous study focused on a

limited set of biomarkers: one antioxidant enzyme, oxidative damage on lipids and oxidative

balance. In the present study, we measured 7 biomarkers, thus providing broader support to the

absence of effects of prenatal THs on post-natal/hatching oxidative stress.

Conclusion

To our knowledge, this study is the first one to experimentally investigate the consequences of

natural variations of maternal THs not only early but also in adult physiology and postnuptial

moult in any vertebrate. Furthermore, this study explored for the first time the effects of maternal

T_3 and T_4 separately. We found no evidence for differential effects of maternal T_4 and T_3, while an

additive effect of T_3 and T_4 on hatching success suggests that T_4 is converted into T_3, the

biologically active form during embryonic development. Contrary to similar studies on wild

altricial species, we found no influence of maternal THs on growth. Further research on embryos

utilisation of maternal THs may help understand the differences observed between precocial and

altricial species. Studies in other vertebrates are urgently needed to understand the potential

organising effects of maternal THs with long-term consequences.

List of symbols and abbreviations

- CAT: catalase

- CO: control treatment

- GP: glutathione peroxidase

- tGSH: oxidised glutathione

- GSSG: reduced glutathione

- GST: Glutathione S-transferase

- MDA: malonaldehyde

- RMR: resting metabolic rate

- SOD: super-oxide dismutase

- T_3: triiodothyronine

- T_4: thyroxine

- THs: thyroid hormones

Acknowledgements

We thank Sophie Michon for her help on setting up the parental generation. We also thank Ido Pen for consultation and help with statistical analysis, and Esther Chang for her help throughout the writing phase.

References

Ainsworth SJ, Stanley RL, Evans DJR. 2010. Developmental stages of the Japanese quail. *Journal of Anatomy* 216:3–15. DOI: 10.1111/j.1469-7580.2009.01173.x.

Alonso M, Goodwin C, Liao X, Page D, Refetoff S, Weiss RE. 2007. Effects of Maternal Levels of Thyroid Hormone (TH) on the Hypothalamus-Pituitary-Thyroid Set Point: Studies in TH Receptor β Knockout Mice. *Endocrinology* 148:5305–5312. DOI: 10.1210/en.2007-0677.

Alonso-Alvarez C, Bertrand S, Faivre B, Sorci G. 2007. Increased susceptibility to oxidative damage as a cost of accelerated somatic growth in zebra finches. *Functional Ecology* 21:873–879. DOI: 10.1111/j.1365-2435.2007.01300.x.

Anselmo J, Scherberg NH, Dumitrescu AM, Refetoff S. 2019. Reduced Sensitivity to Thyroid Hormone as a Transgenerational Epigenetic Marker Transmitted Along the Human Male Line. *Thyroid* 29:778–782. DOI: 10.1089/thy.2019.0080.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67:1–48. DOI: 10.18637/jss.v067.i01.

Bentz AB, Becker DJ, Navara KJ. 2016. Evolutionary implications of interspecific variation in a maternal effect: a meta-analysis of yolk testosterone response to competition. *Royal Society Open Science* 3:160499. DOI: 10.1098/rsos.160499.

Biswas A, Ranganatha OS, Mohan J, Sastry KVH. 2007. Relationship of cloacal gland with testes, testosterone and fertility in different lines of male Japanese quail. *Animal Reproduction Science* 97:94–102. DOI: 10.1016/j.anireprosci.2005.12.012.

Brown, C. L., Doroshov, S. I., Nunez, J. M., Hadley, C., Vaneenennaam, J., Nishioka, R. S. and Bern, H. A. (1988). Maternal triiodothyronine injections cause increases in swimbladder

inflation and survival rates in larval striped bass, *Morone saxatilis*. *J. Exp. Zool.* 248, 168–176.

Burnham KP, Anderson DR. 2002. *Model selection and multimodel inference: a practical information-theoretic approach*. New York: Springer.

Burnham KP, Anderson DR, Huyvaert KP. 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology* 65:23–35. DOI: 10.1007/s00265-010-1029-6.

Cheng KM, Hickman AR, Nichols CR. 1989. Role of the proctodeal gland foam of male Japanese Quail in natural copulations. *The Auk* 106:279–285.

Cheng KM, McIntyre RF, Hickman AR. 1989. Proctodeal Gland Foam Enhances Competitive Fertilization in Domestic Japanese Quail. *The Auk* 106:286–291.

Dardente H, Hazlerigg DG, Ebling FJP. 2014. Thyroid Hormone and Seasonal Rhythmicity. *Frontiers in Endocrinology* 5:19. DOI: 10.3389/fendo.2014.00019.

Darras VM, Van Herck SLJ, Geysens S, Reyns GE. 2009. Involvement of thyroid hormones in chicken embryonic brain development. *General and Comparative Endocrinology* 163:58–62. DOI: 10.1016/j.ygcen.2008.11.014.

Davis PJ, Goglia F, Leonard JL. 2016. Nongenomic actions of thyroid hormone. *Nature Reviews Endocrinology* 12:111–121. DOI: 10.1038/nrendo.2015.205.

Dawson A. 1993. Thyroidectomy progressively renders the reproductive system of starlings (Sturnus vulgaris) unresponsive to changes in daylength. *Journal of Endocrinology* 139:51–55. DOI: 10.1677/joe.0.1390051.

Dawson A. 2015. Chapter 38 - Avian Molting. In: Scanes CG ed. *Sturkie's Avian Physiology (Sixth Edition)*. San Diego: Academic Press, 907–917. DOI: 10.1016/B978-0-12-407160-5.00038-5.

Dawson A, McNaughton FJ, Goldsmith AR, Degen AA. 1994. Ratite-like neoteny induced by

    neonatal thyroidectomy of European starlings, *Sturnus vulgaris*. *Journal of Zoology*

    232:633–639. DOI: 10.1111/j.1469-7998.1994.tb04618.x.

Duarte-Guterman P, Langlois VS, Pauli BD, Trudeau VL. 2010. Expression and T3 regulation of

    thyroid hormone- and sex steroid-related genes during Silurana (Xenopus) tropicalis early

    development. *General and Comparative Endocrinology* 166:428–435. DOI:

    10.1016/j.ygcen.2009.12.008.

von Engelhardt N, Groothuis TGG. 2011. Maternal Hormones in Avian Eggs. In: *Hormones and*

    Reproduction of Vertebrates. Elsevier, 91–127. DOI: 10.1016/B978-0-12-374929-

    1.10004-6.

Ercal N, Gurer-Orhan H, Aykin-Burns N. 2001. Toxic Metals and Oxidative Stress Part I:

    Mechanisms Involved in Metal-induced Oxidative Damage. *Current Topics in Medicinal*

    Chemistry 1:529–539. DOI: 10.2174/1568026013394831.

Faul F, Erdfelder E, Buchner A, Lang A-G. 2009. Statistical power analyses using G*Power 3.1:

    Tests for correlation and regression analyses. *Behavior Research Methods* 41:1149–1160.

    DOI: 10.3758/BRM.41.4.1149.

Fini JB, Mével SL, Palmier K, Darras VM, Punzon I, Richardson SJ, Clerget-Froidevaux MS,

    Demeneix BA. 2012. Thyroid Hormone Signaling in the Xenopus laevis Embryo Is

    Functional and Susceptible to Endocrine Disruption. *Endocrinology* 153:5068–5081.

    DOI: 10.1210/en.2012-1463.

Giesing ER, Suski CD, Warner RE, Bell AM. 2011. Female sticklebacks transfer information via

    eggs: effects of maternal experience with predators on offspring. *Proceedings of the Royal*

    Society B: Biological Sciences 278:1753–1759. DOI: 10.1098/rspb.2010.1819.

Ginn H, Melville D. 1983. *{Moult in birds (BTO guide)}*. Thetford: British Trust for Ornithology.

Gluckman PD, Hanson MA, Spencer HG. 2005. Predictive adaptive responses and human evolution. *Trends in Ecology & Evolution* 20:527–533. DOI: 10.1016/j.tree.2005.08.001.

Groothuis TGG, von Engelhardt N. 2005. Investigating Maternal Hormones in Avian Eggs: Measurement, Manipulation, and Interpretation. *Annals of the New York Academy of Sciences* 1046:168–180. DOI: 10.1196/annals.1343.014.

Groothuis TGG, Müller W, von Engelhardt N, Carere C, Eising C. 2005. Maternal hormones as a tool to adjust offspring phenotype in avian species. *Neuroscience & Biobehavioral Reviews* 29:329–352. DOI: 10.1016/j.neubiorev.2004.12.002.

Halekoh, U., Højsgaard, S. 2014. A Kenward-Roger Approximation and Parametric Bootstrap Methods for Tests in Linear Mixed Models - The *R* Package pbkrtest. J. Stat. Softw. 59.

Halliwell B, Gutteridge JMC. 2015. *Free Radicals in Biology and Medicine*. Oxford University Press.

Hoffman DJ. 2002. Role of selenium toxicity and oxidative stress in aquatic birds. *Aquatic Toxicology* 57:11–26. DOI: 10.1016/S0166-445X(01)00263-6.

Holsberger DR, Cooke PS. 2005. Understanding the role of thyroid hormone in Sertoli cell development: a mechanistic hypothesis. *Cell and Tissue Research* 322:133–140. DOI: 10.1007/s00441-005-1082-z.

Hsu B-Y, Dijkstra C, Darras VM, de Vries B, Groothuis TGG. 2017. Maternal thyroid hormones enhance hatching success but decrease nestling body mass in the rock pigeon (Columba livia). *General and Comparative Endocrinology* 240:174–181. DOI: 10.1016/j.ygcen.2016.10.011.

Hsu B-Y, Doligez B, Gustafsson L, Ruuskanen S. 2019. Transient growth-enhancing effects of elevated maternal thyroid hormones at no apparent oxidative cost during early postnatal period. *Journal of Avian Biology*. DOI: 10.1111/jav.01919.

Isaksson C, Örnborg J, Stephensen E, Andersson S. 2005. Plasma Glutathione and Carotenoid
Coloration as Potential Biomarkers of Environmental Stress in Great Tits. *EcoHealth*
2:138–146. DOI: 10.1007/s10393-005-3869-5.

Jimeno B, Hau M, Gómez-Díaz E, Verhulst S. 2019. Developmental conditions modulate DNA
methylation at the glucocorticoid receptor gene with cascading effects on expression and
corticosterone levels in zebra finches. *Scientific Reports* 9:15869. DOI: 10.1038/s41598-
019-52203-8.

Keshavarz K, Quimby FW. 2002. An Investigation of Different Molting Techniques with an
Emphasis on Animal Welfare. *The Journal of Applied Poultry Research* 11:54–67. DOI:
10.1093/japr/11.1.54.

Kikuyama S, Kawamura K, Tanaka S, Yamamoto K. 1993. Aspects of Amphibian
Metamorphosis: Hormonal Control. In: *International Review of Cytology*. Elsevier, 105–
148. DOI: 10.1016/S0074-7696(08)60426-X.

Korevaar TIM, Muetzel R, Medici M, Chaker L, Jaddoe VWV, de Rijke YB, Steegers EAP,
Visser TJ, White T, Tiemeier H, Peeters RP. 2016. Association of maternal thyroid
function during early pregnancy with offspring IQ and brain morphology in childhood: a
population-based prospective cohort study. *The Lancet Diabetes & Endocrinology* 4:35–
43. DOI: 10.1016/S2213-8587(15)00327-7.

Krassas GE, Poppe K, Glinoer D. 2010. Thyroid Function and Human Reproductive Health.
Endocrine Reviews 31:702–755. DOI: 10.1210/er.2009-0041.

Lenth R. 2019. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package
version 1.3.2.

Lilley TM, Ruokolainen L, Meierjohann A, Kanerva M, Stauffer J, Laine VN, Atosuo J, Lilius E-
M, Nikinmaa M. 2013. Resistance to oxidative damage but not immunosuppression by

organic tin compounds in natural populations of Daubenton's bats (Myotis daubentonii). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 157:298–305. DOI: 10.1016/j.cbpc.2013.01.003.

Marshall DJ, Uller T. 2007. When is a maternal effect adaptive? *Oikos* 116:1957–1963. DOI: 10.1111/j.2007.0030-1299.16203.x.

McNabb FMA. 2007. The Hypothalamic-Pituitary-Thyroid (HPT) Axis in Birds and Its Role in Bird Development and Reproduction. *Critical Reviews in Toxicology* 37:163–193.

McNabb FMA, Darras VM. 2015. Thyroids. In: *Sturkie's Avian Physiology (Sixth Edition)*. London: Elsevier, 535–547. DOI: 10.1016/B978-0-12-407160-5.00024-5.

McNabb FMA, Scanes CG, Zeman M. 1998. Endocrine control of development. In: *Avian Growth and Development: Evolution Within the Altricial-precocial Spectrum*. New York: Starcj, J.M., Ricklefs, R.E., 174–202.

Medici M, Timmermans S, Visser W, de Muinck Keizer-Schrama SMPF, Jaddoe VWW, Hofman A, Hooijkaas H, de Rijke YB, Tiemeier H, Bongers-Schokking JJ, Visser TJ, Peeters RP, Steegers EAP. 2013. Maternal Thyroid Hormone Parameters during Early Pregnancy and Birth Weight: The Generation R Study. *The Journal of Clinical Endocrinology & Metabolism* 98:59–66. DOI: 10.1210/jc.2012-2420.

Mommer BC, Bell AM. 2013. A test of maternal programming of offspring stress response to predation risk in threespine sticklebacks. *Physiology & Behavior* 122:222–227. DOI: 10.1016/j.physbeh.2013.04.004.

Mommer BC, Bell AM. 2014. Maternal Experience with Predation Risk Influences Genome-Wide Embryonic Gene Expression in Threespined Sticklebacks (Gasterosteus aculeatus). *PLoS ONE* 9:e98564. DOI: 10.1371/journal.pone.0098564.

Morreale de Escobar G, Obregon M, Escobar del Rey F. 2004. Role of thyroid hormone during

early brain development. *European Journal of Endocrinology*:U25–U37. DOI: 10.1530/eje.0.151u025.

Mousseau TA, Fox CW. 1998. *Maternal Effects As Adaptations*. New York: Oxford University Press.

Okuliarová M, Škrobánek P, Zeman M. 2007. Effect of Increasing Yolk Testosterone Levels on Early Behaviour in Japanese Quail Hatchlings. *Acta Veterinaria Brno* 76:325–331. DOI: 10.2754/avb200776030325.

Pfannkuche KA, Gahr M, Weites IM, Riedstra B, Wolf C, Groothuis TGG. 2011. Examining a pathway for hormone mediated maternal effects – Yolk testosterone affects androgen receptor expression and endogenous testosterone production in young chicks (Gallus gallus domesticus). *General and Comparative Endocrinology* 172:487–493. DOI: 10.1016/j.ygcen.2011.04.014.

Pick JL, Ebneter C, Hutter P, Tschirren B. 2016. Disentangling Genetic and Prenatal Maternal Effects on Offspring Size and Survival. *The American Naturalist* 188:628–639. DOI: 10.1086/688918.

Pinheiro JC, Bates DM, Sarkar D, R Core Team. 2018. nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-137.

Podmokła E, Drobniak SM, Rutkowska J. 2018. Chicken or egg? Outcomes of experimental manipulations of maternally transmitted hormones depend on administration method - a meta-analysis: Maternal hormones and manipulation methods. *Biological Reviews* 93:1499–1517. DOI: 10.1111/brv.12406.

Prati M, Calvo R, Morreale G, Morreale de Escobar G. 1992. L-thyroxine and 3,5,3'-triiodothyronine concentrations in the chicken egg and in the embryo before and after the onset of thyroid function. *Endocrinology* 130:2651–2659. DOI:

10.1210/endo.130.5.1572286.

Raine JC, Cameron C, Vijayan MM, Lamarre J, Leatherland JF. 2004. The effect of elevated

oocyte triiodothyronine content on development of rainbow trout embryos and expression

of mRNA encoding for thyroid hormone receptors. *Journal of Fish Biology* 65:206–226.

DOI: 10.1111/j.0022-1112.2004.00445.x.

Rainio MJ, Eeva T, Lilley T, Stauffer J, Ruuskanen S. 2015. Effects of early-life lead exposure on

oxidative status and phagocytosis activity in great tits (Parus major). *Comparative*

Biochemistry and Physiology Part C: Toxicology & Pharmacology 167:24–34. DOI:

10.1016/j.cbpc.2014.08.004.

Rainio MJ, Kanerva M, Salminen J-P, Nikinmaa M, Eeva T. 2013. Oxidative status in nestlings of

three small passerine species exposed to metal pollution. *Science of The Total*

Environment 454–455:466–473. DOI: 10.1016/j.scitotenv.2013.03.033.

Reis LH, Gama L, Soares M. 1997. Effects of short storage conditions and broiler breeder age on

hatchability, hatching time, and chick weights. *Poultry Science* 76:1459–1466. DOI:

10.1093/ps/76.11.1459.

Rubin C-J, Zody MC, Eriksson J, Meadows JRS, Sherwood E, Webster MT, Jiang L, Ingman M,

Sharpe T, Ka S, Hallböök F, Besnier F, Carlborg Ö, Bed'hom B, Tixier-Boichard M,

Jensen P, Siegel P, Lindblad-Toh K, Andersson L. 2010. Whole-genome resequencing

reveals loci under selection during chicken domestication. *Nature* 464:587–591. DOI:

10.1038/nature08832.

Ruuskanen S, Darras VM, Visser ME, Groothuis TGG. 2016. Effects of experimentally

manipulated yolk thyroid hormone levels on offspring development in a wild bird species.

Hormones and Behavior 81:38–44. DOI: 10.1016/j.yhbeh.2016.03.006.

Ruuskanen S, Hsu B-Y. 2018. Maternal Thyroid Hormones: An Unexplored Mechanism

Underlying Maternal Effects in an Ecological Framework. *Physiological and Biochemical Zoology* 91:904–916. DOI: 10.1086/697380.

Sarraude T, Hsu B-Y, Groothuis T, Ruuskanen S. 2020. Manipulation of prenatal thyroid hormones does not affect growth or physiology in nestling pied flycatchers. *Physiological and Biochemical Zoology*, in press.

Sekimoto K, Imai K, Suzuki M, Takikawa H, Hoshino N, Totsuka K. 1987. Thyroxine-Induced Molting and Gonadal Function of Laying Hens. *Poultry Science* 66:752–756. DOI: 10.3382/ps.0660752.

Srichomkwun P, Anselmo J, Liao X-H, Hönes GS, Moeller LC, Alonso-Sampedro M, Weiss RE, Dumitrescu AM, Refetoff S. 2017. Fetal Exposure to High Maternal Thyroid Hormone Levels Causes Central Resistance to Thyroid Hormone in Adult Humans and Mice. *The Journal of Clinical Endocrinology & Metabolism* 102:3234–3240. DOI: 10.1210/jc.2017-00019.

Stier A, Metcalfe NB, Monaghan P. 2019. Ageing before birth: pace and stability of prenatal growth affect telomere dynamics. *bioRxiv*:809087. DOI: 10.1101/809087.

Tsuyoshi H, Wada M. 1992. Termination of LH secretion in Japanese quail due to high- and low-temperature cycles and short daily photoperiods. *General and Comparative Endocrinology* 85:424–429. DOI: 10.1016/0016-6480(92)90087-Z.

Van Herck SLJ, Delbaere J, Bourgeois NMA, McAllan BM, Richardson SJ, Darras VM. 2015. Expression of thyroid hormone transporters and deiodinases at the brain barriers in the embryonic chicken: Insights into the regulation of thyroid hormone availability during neurodevelopment. *General and Comparative Endocrinology* 214:30–39. DOI: 10.1016/j.ygcen.2015.02.021.

Van Herck SLJ, Geysens S, Delbaere J, Tylzanowski P, Darras VM. 2012. Expression profile and

thyroid hormone responsiveness of transporters and deiodinases in early embryonic chicken brain development. *Molecular and Cellular Endocrinology* 349:289–297. DOI: 10.1016/j.mce.2011.11.012.

Wada H. 2008. Glucocorticoids: Mediators of vertebrate ontogenetic transitions. *General and Comparative Endocrinology* 156:441–453. DOI: 10.1016/j.ygcen.2008.02.004.

Walpita CN, Van der Geyten S, Rurangwa E, Darras VM. 2007. The effect of 3,5,3′-triiodothyronine supplementation on zebrafish (Danio rerio) embryonic development and expression of iodothyronine deiodinases and thyroid hormone receptors. *General and Comparative Endocrinology* 152:206–214. DOI: 10.1016/j.ygcen.2007.02.020.

Watanabe Y, Grommen SVH, De Groef B. 2016. Corticotropin-releasing hormone: Mediator of vertebrate life stage transitions? *General and Comparative Endocrinology* 228:60–68. DOI: 10.1016/j.ygcen.2016.02.012.

Wilson CM, McNabb FMA. 1997. Maternal Thyroid Hormones in Japanese Quail Eggs and Their Influence on Embryonic Development. *General and Comparative Endocrinology* 107:153–165. DOI: 10.1006/gcen.1997.6906.

Wood SN. 2017.Generalized Additive Models: An Introduction with R, Second Edition. *Available at https://www.crcpress.com/Generalized-Additive-Models-An-Introduction-with-R-Second-Edition/Wood/p/book/9781498728331* (accessed November 8, 2018).

Zuur AF, Ieno EN, Walker NJ, Savaliev AA, Smith GM. 2009. *Mixed effects models and extensions in ecology with R*. New York, NY: Springer.

Figure legends

Figure 1: Percentage of hatching success according to yolk TH manipulation treatments: N = 40 CO, 39 T_3, 39 T_4 and 40 T_3T_4. CO control, T_4 (thyroxine) = injection of T_4, T_3 (triiodothyronine) = injection of T_3, T_3T_4 = injection of T_3 and T_4.

Figure 2: Growth curves in body mass of Japanese quails hatching from eggs treated with either T_3, T_4, a combination of both hormones, or a control solution. See Fig. 1 for a description of the treatments. Each line represents an individual bird, while thick coloured lines represent mean values. A: Growth curve according to yolk TH manipulation. N = 7 CO, 11 T_3, 18 T_4 and 21 T_3T_4. B: Growth curve according to sex. N = 29 females and 28 males.

Figure 3: Primary moult score in 7-month old Japanese quails according to yolk TH manipulation treatments: N = 7 CO, 11 T_3, 16 T_4 and 20 T_3T_4. See Fig. 1 for a description of the treatments. Measures were taken once a week after switching from long photoperiod (16L:8D) to short photoperiod (8L:16D, switch = time point 0 on x-axis). Each line represents an individual bird, while thick coloured lines represent group mean values.

Figure 4: MDA concentration according to yolk TH manipulation treatments, samples from two ages pooled: N = 7 CO, 11 T_3, 17 T_4 and 20 T_3T_4. See Fig. 1 for a description of the treatments.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Tables

Table 1: Loadings of the different antioxidant biomarkers on the principal components 1 and 2.

Factor loadings	PC1 (34.0%)	PC2 (26.2%)
CAT	-0.49	0.14
SOD	0.20	-0.71
GST	-0.65	-0.10
GP	0.04	-0.63
tGSH	-0.60	-0.26

Table 2: Results of the Generalised Additive Mixed Models (GAMMs) on body mass growth, with sex and treatment fitted either as intercept, curve shape or both (all combinations tested). A total of 12 GAMMs were fitted and ranked based on their AIC, from the lowest to the highest. Weight: Akaike's weights.

Model	Intercept	Curve shape	ΔAIC	df	Weight
1	Sex	Sex	0.0	11	0.8430
8	Treatment + sex	Sex	3.5	14	0.1497
3	-	Sex	9.9	10	0.0061
2	Treatment	Sex	13.2	13	0.0012
11	Sex	-	77.6	9	<0.001
9	Treatment + sex	-	81.6	12	<0.001
12	-	-	91.2	8	<0.001
10	Treatment	-	95.0	11	<0.001
5	Sex	Treatment	147.9	15	<0.001
7	Treatment + sex	Treatment	151.7	18	<0.001
6	-	Treatment	161.2	14	<0.001
4	Treatment	Treatment	165.5	17	<0.001

Table 3: 95% confidence intervals of the predictors in the top-ranked models according to AIC values (see Tables 2, S1 and S2). Predictors in bold have confidence intervals that do not overlap with 0. For the intercept, the reference groups are female and CO for the predictors sex and treatment, respectively.

Curve parameter	Predictors	Lower limit	Estimate	Upper limit
(A) Body mass (Model 1)				
Intercept	**Sex (M)**	-19.7	-12.6	-5.5
Curve shape	**Sex (F)**	9.9	20.0	30.0
Curve shape	**Sex (M)**	14.3	24.5	34.7
(B) Wing length (Model 11)				
Intercept	**Sex (M)**	-2.3	-1.2	-0.1
Curve shape	**Age**	26.4	28.7	31.0
(C) Tarsus length (Model 10)				
Intercept	Treatment (T_3)	-0.8	0.02	0.8
Intercept	**Treatment (T_3T_4)**	-1.5	-0.8	-0.1
Intercept	Treatment (T_4)	-1.3	-0.6	0.2
Curve shape	**Age**	10.5	11.1	11.8
Tarsus length (Model 9)				
Intercept	Treatment (T_3)	-0.9	-0.07	0.7
Intercept	**Treatment (T_3T_4)**	-1.5	-0.8	-0.1
Intercept	Treatment (T_4)	-1.4	-0.6	0.1
Intercept	Sex (M)	-0.8	-0.3	0.3
Curve shape	**Age**	10.5	11.1	11.7