METHOD

Unified inference of missense variant effects and gene constraints in the human genome

Yi-Fei Huang

Correspondence: yuh371@psu.edu
Department of Biology and Huck
Institutes of the Life Sciences,
Pennsylvania State University,
University Park, PA 16802, USA
Full list of author information is
available at the end of the article

Abstract

Numerous statistical methods have been developed to predict deleterious missense variants or constrained genes in the human genome, but unified prioritization methods that utilize both variant- and gene-level information are underdeveloped. Here we present UNEECON, an evolution-based deep learning framework for unified variant and gene prioritization. UNEECON outperforms existing methods in predicting variants and genes associated with dominant disorders. Furthermore, based on UNEECON we show that disordered proteins are tolerant to missense mutations but not to loss-of-function mutations. The genes under strong selective constraints at both missense and loss-of-function levels are likely to play key roles in brain development and autism.

Keywords: Variant prioritization; Gene constraint; Negative selection; Deep learning

Background

A fundamental question in biology is to understand how genomic variation contributes to phenotypic variation and disease risk. While millions of protein-altering variants have been identified in the human genome, it is challenging to assess the functional and clinical significance of these variants. In particular, a large fraction of missense variants have been annotated as "variants of uncertain significance" (VUS) [1, 2], forming a major hurdle for both basic research and medical practice. This problem is further exacerbated by the difficulty of experimentally validating the function of large numbers of missense variants *in vivo*. Therefore, there is a tremendous need for accurate computational tools to prioritize deleterious missense variants [3].

Since early-onset, severe genetic disorders are often associated with a reduction of evolutionary fitness, signatures of negative (purifying) selection, such as sequence conservation, have been widely used to predict deleterious variants associated with Mendelian disorders [4, 5, 6, 7, 8, 9, 10, 11, 12]. Among existing evolutionary approaches for variant prioritization, recently developed integrative methods are particularly powerful [13, 8, 14, 9, 10, 11, 12, 15]. By learning a linear or nonlinear mathematical function from predictive variant features, such as sequence conservation scores and protein structural features, to the strength of negative selection, these statistical methods estimate negative selection on observed and potential mutations in the human genome. The estimated strength of negative selection can then be utilized to prioritize deleterious variants associated with severe genetic disorders. Because these evolutionary approaches are trained on tremendous natural polymorphisms observed in healthy individuals instead of sparsely annotated pathogenic

variants, they have shown good performance in predicting pathogenic variants, frequently outperforming or on par with supervised machine learning models trained on disease data [8, 10, 11, 12].

Despite the success of evolution-based metrics of variant effects, the existing methods nevertheless suffer from a few critical limitations. First, most existing methods focus on learning a shared mathematical function from predictive variant features to negative selection and assume that this function is equally applicable to all protein-coding genes. Instead, a subset of genes can depart from the genome-wide trend between variant features and negative selection, possibly due to enhanced or relaxed purifying selection on these genes in the human lineage [12]. Second, most existing methods are trained on common genetic variants, making it challenging to distinguish strong negative selection associated with severe genetic diseases from moderate negative selection without clinical implications. Third, these methods typically are agnostic to the mode of inheritance and, therefore, may be suboptimal in the prediction of deleterious variants associated with dominant disorders on which we focus in this work.

In parallel with the development of variant-level interpretation methods, several complementary, gene-centric methods have been proposed to predict protein-coding genes associated with dominant genetic disorders [16, 17, 18, 19, 20, 21, 22, 23]. Unlike variant-level predictors, the gene-level prioritization methods seek to identify constrained genes that are intolerant to heterozygous nonsynonymous mutations. These gene-level constraint metrics have been shown to provide complementary information on variant effects and have been successfully used to prioritize variants associated with dominant genetic disorders [16, 17, 24]. However, these methods typically assume that all the missense mutations in a gene or a genic region have identical effects and, therefore, may not be able to distinguish pathogenic missense variants from proximal benign missense variants.

Since variant-level and gene-level prioritization methods leverage complementary signatures of negative selection, unifying the two lines of research for joint inference of variant effects and gene constraints should be beneficial. Recently, a couple of studies have tried to address this question [25, 26, 27], but all of these methods are supervised machine learning models trained on disease variants. Therefore, we lack an evolution-based statistical framework to combine predictive variant features and gene-level selective constraints for variant and gene prioritization. Based on a novel deep learning framework, *i.e.*, deep mixed-effects model, we develop **UNEECON** (**UN**ified inferenc**E** of variant **E**ffects and gene **CON**straints), an evolution-based framework to predict deleterious variants and constrained genes from both variant features and gene-level intolerance to missense mutations. By integrating 30 predictive variant features and genomic variation from 141,456 human genomes [28, 29], UNEECON outperforms existing variant effect predictors and gene constraint scores in predicting pathogenic missense variants with a dominant mode of inheritance. In addition, deleterious *de novo* variants predicted by UNEECON are strongly enriched in individuals affected by severe development disorders [30, 31], highlighting its power for interpreting the effects of *de novo* mutations. Furthermore, UNEECON provides estimates of gene-level selective constraints (UNEECON-G scores) for all protein-coding genes. In the setting of gene prioritization, UNEECON-G scores

show unmatched performance in predicting human essential genes [32], mouse essential genes [33, 34], autosomal dominant disease genes [35, 36], and haploinsufficient genes [37]. UNEECON is a powerful framework for both variant and gene prioritization.

Results

UNEECON integrates variant- and gene-level information to predict detrimental variants and constrained genes

The key idea of UNEECON is to combine variant-level predictive features, such as sequence conservation scores and protein structural features, and gene-level signatures of selective constraints to infer negative selection on every potential missense mutation in the human genome. Inspired by classical sequence conservation models [38, 39, 40, 41, 42], which use site-specific substitution rate as a proxy of negative selection, we utilize the relative probability of the occurrence of a potential missense mutation in human populations, compared to neutral mutations, as an allele-specific predictor of negative selection.

We first fit a mutation model to calculate μ_{ij}, *i.e.*, the probability of the occurrence of mutation i in gene j when selection is absent. Our mutation model is trained on putatively neutral mutations in the gnomAD data [29] and captures the impact of multiple neutral and technical factors, including the 7-mer sequence context centered on the focal site, the identity of the alternative allele, the local mutation rate, and the average sequencing coverage, on the probability of occurrence of a neutral mutation. At the gene level, the number of synonymous mutations predicted by the mutation model was nearly perfectly correlated with the observed number of synonymous mutations (Spearman's $\rho = 0.976$; Fig. S1), suggesting that the mutation model can serve as a proper baseline for inferring negative selection.

Given the mutation model, UNEECON employs a novel deep learning framework, deep mixed-effects model, to infer the relative probability of the occurrence of each potential missense mutation (Fig. 1). In more detail, we denote η_{ij} as the relative probability of the occurrence of missense mutation i in gene j with respect to the neutral occurrence probability, μ_{ij}. η_{ij} captures the impact of natural selection, instead of mutation and genetic drift, on the occurrence of a missense mutation. Analogous to generalized linear mixed models, we assume that η_{ij} depends on the sum of a variant-level fixed effect, z_{ij}, and a gene-level random effect, u_j. We assume that z_{ij} captures the contribution of predictive variant features to negative selection. Denoting $\mathbf{X}_{ij}$ as the vector of predictive features associated with mutation i in gene j, we use a feedforward neural network to model the relationship between $\mathbf{X}_{ij}$ and z_{ij} (Fig. 1). Furthermore, we assume that the random-effect term, u_j, is a Gaussian (normal) random variable which models the variation of gene-level constraints that is not predictable from feature vector $\mathbf{X}_{ij}$. We then perform a logistic transformation on the sum of z_{ij} and u_j to obtain η_{ij}. Finally, we multiply η_{ij} and μ_{ij} to obtain the occurrence probability of missense mutation i in the gnomAD exome sequencing data.

We estimate the parameters of the UNEECON model by minimizing the discrepancy between the predicted occurrence probability, $\eta_{ij} \cdot \mu_{ij}$, and the observed presence/absence of each potential missense mutation, which is equivalent to maximizing a Bernoulli likelihood function. We also employ regularization techniques,

including dropout [43] and early stopping [44], to avoid overfitting. After training, we calculate variant-effect scores (UNEECON scores) and gene-level intolerance to missense mutations (UNEECON-G scores) defined as the expected reduction of the occurrence probability at variant and gene levels, respectively. Both UNEECON and UNEECON-G scores range from 0 to 1, with higher scores suggesting stronger negative selection.

UNEECON scores capture variation of negative selection within and across genes

We trained the UNEECON model with 30 missense variant features, including conservation scores, protein structural features, and regulatory features (Table S1), and rare missense variants with a minor allele frequency (MAF) lower than 0.1% in the gnomAD dataset [29]. After the training process, we first compared the distributions of UNEECON scores across potential missense mutations in haploinsufficient genes [37], autosomal dominant disease genes [35, 36], autosomal recessive disease genes [35, 36], and olfactory receptor genes [45]. In agreement with previous studies [35, 16, 28], missense mutations in autosomal dominant disease genes had higher UNEECON scores than those in autosomal recessive disease genes (Fig. 2a), suggesting a stronger selection on heterozygous missense mutations in the genes associated with autosomal dominant disorders. Also, we observed that missense mutations in olfactory receptor genes had much lower UNEECON scores (Fig. 2a), potentially due to relaxed purifying selection or enhanced positive selection in the human lineage [46].

Second, we investigated whether the distributions of UNEECON scores varied across different types of protein regions annotated by UniProt [47, 48] and MobiDB [49]. We observed that the distributions of UNEECON scores were similar among α-helices, β-strands, and hydrogen-bonded turns (Fig. 2b). In contrast, UNEECON scores were significantly lower in disordered protein regions. As expected, UNEECON scores were significantly higher in functional protein sites, including enzyme active sites and ligand binding sites (Fig. 2b), suggesting much stronger purifying selection on these critical residues.

Furthermore, we used the human CDKL5 protein as an example to illustrate UNEECON's capability to capture regional variation of negative selection within a gene. Similar to a previously published metric of regional missense constraints [25], UNEECON predicted that the N-terminus of CDKL5 was under strong negative selection (Fig. 2c). In agreement with the evidence of strong selection, this region was also enriched with known pathogenic missense variants and depleted with benign missense variants (Fig. 2c). Interestingly, UNEECON also predicted that the C-terminus of CDKL5 was under much weaker selection compared with the rest of the protein (Fig. 2c). Accordingly, this unconstrained region was enriched with benign missense variants and depleted with pathogenic missense variants.

UNEECON scores accurately predict missense mutations associated with dominant genetic disorders

Since UNEECON measures the strength of strong negative selection on heterozygous missense mutations, we hypothesize that UNEECON scores are predictive of missense variants associated with dominant Mendelian disorders. To test this hypothesis, we compared the performance of UNEECON with eight existing methods

in the setting of predicting autosomal dominant disease variants in ClinVar [30]. The eight previously published methods can be classified into three categories: 1) variant-level prediction methods, including LASSIE [12], CADD [8], Eigen [50], and PrimateAI [11]; 2) gene-level and region-level prediction methods, including RVIS [16], pLI [17], and CCR [22]; and 3) MPC [25], a supervised machine learning method utilized both variant features and regional constraint scores as input features. It is worth noting that several aforementioned methods, including UNEECON, used the variants from the gnomAD dataset as a part of their training data. If the same gnomAD variants are also used as negative controls in the evaluation of performance, we will overestimate the power of these methods trained on the gnomAD data. To avoid this problem, we used benign missense variants in ClinVar as negative controls and removed any ClinVar variants that are also present in gnomAD. Then, we matched the numbers of positives and negatives by random sampling without replacement.

Overall, UNEECON outperformed previous methods in predicting ClinVar missense variants associated with autosomal dominant disorders (Fig. 3a; Table S2). Among the previously published methods, variant-level predictors performed better than gene-level and region-level constraint scores. To test the robustness of these results, we constructed an alternative set of dominant pathogenic variants defined as all the pathogenic missense variants located in 709 genes associated with autosomal dominant diseases [35, 36]. UNEECON again outperformed the other methods in this dataset, even though the difference of performance between UNEECON and LASSIE was not statistically significant (Fig. S2; Table S2).

While UNEECON was highly powerful in predicting autosomal dominant disease variants, it might not be as accurate when applied to predict recessive disease variants which are only detrimental in the homozygous state. To test this hypothesis, we evaluated UNEECON's performance in predicting autosomal recessive disease variants in ClinVar [30]. As expected, UNEECON was outperformed by multiple methods, such as LASSIE [12] and Eigen [50], in this setting (Fig. S3). Similar results were reached with an alternative set of recessive pathogenic variants defined as all the pathogenic missense variants located in 1,183 autosomal recessive genes [35, 36] (Fig. S4).

As an orthogonal benchmark of UNEECON's performance in predicting dominant disease variants, we investigated whether UNEECON was able to predict *de novo* missense mutations identified in individuals affected by severe developmental disorders [31]. We obtained *de novo* missense variants identified in affected individuals and healthy individuals from denovo-db [51]. Then, for multiple percentile rank cutoff values (top 10%, 20%, 30%, and 40%), we evaluated the enrichment of deleterious variants predicted by each method in the affected individuals. Overall, missense variants predicted by UNEECON and MPC showed the highest enrichments in the affected individuals (Fig. 3b), suggesting that these two methods were more powerful in predicting de novo risk mutations. The performance gaps between UNEECON/MPC and the other methods were highest at the most stringent cutoff of 10% (Fig. 3b).

UNEECON-G scores perform favorably in predicting disease genes and essential genes

The UNEECON model also provided UNEECON-G scores which measure gene-level intolerance to missense mutations. We compared the performance of UNEECON-G scores with alternative gene constraint scores, including RVIS [16], pLI [17], mis-z [17], and GDI [19], in the setting of predicting disease genes and essential genes. The disease gene sets included 709 autosomal dominant disease genes [35, 36] and 294 haploinsufficient genes [37]. The essential gene sets included 2,454 human orthologs of mouse essential genes [33, 34] and 683 human essential genes identified by CRISPR knockout experiments in human cell lines [32]. For each set of autosomal dominant disease genes, haploinsufficient genes, and mouse essential genes, we constructed a negative gene set by sampling a matched number of genes from the other genes in the human genome. For the essential genes identified by CRISPR in cell lines, we constructed a negative gene set by sampling a matched number of genes from the nonessential genes reported in the same study [32]. Overall, UNEECON-G significantly outperformed alternative gene-level metrics in predicting the four sets of essential and disease genes (Fig. 4; Table S3).

Both variant-level features and gene-level constraints are important for variant interpretation

To gain more insights into which components of the UNEECON model are critical for predicting detrimental variants, we characterized the contributions of the 30 variant features and the gene-level random effect to negative selection. Unfortunately, it is challenging to directly interpret the UNEECON model due to the nonlinearity introduced by the feedforward artificial neural network. To bypass this difficulty, we used a linear version of the UNEECON model as an approximation of the nonlinear UNEECON model. The linear UNEECON model has no hidden units and can be interpreted as a Generalized linear mixed model. Then, we evaluated the importance of the variant-level features and the gene-level random effect in this linear surrogate model. The variant score predicted by the linear UNEECON model was nearly perfectly correlated with the UNEECON score predicted by the original nonlinear UNEECON model (Spearman's $\rho = 0.95$), supporting the use of the linear UNEECON model as a surrogate for model interpretation.

In the linear surrogate model, the relative probability of the occurrence of mutation i in gene j, η_{ij}, is a linear combination of variant features, $\mathbf{X}_{ij}$, and the gene-level random effect, u_j, up to a logistic transformation. Analogous to the interpretation of canonical linear models, we defined the contribution score of a variant feature as the negative value of the weight (regression coefficient) associated with this feature, and similarly defined the contribution score of the gene-level random effect as the negative value of its standard deviation. Positive and negative contribution scores suggest that the corresponding variables are positively and negatively correlated with negative selection, respectively, and contribution scores near zero indicate that the corresponding variables are not important for predicting detrimental mutations.

As shown in Fig. S5, the gene-level random effect was the most important feature for predicting deleterious variants, highlighting the importance of gene-level constraints for variant interpretation. In addition, several conservation scores, such

as the qualitative predictions from LRT, PROVEAN, and SIFT, and a subset of protein structural features, such as the predicted probabilities of forming various protein secondary structures, were also important for accurate prediction of deleterious variants. Even though these variant features individually were less important than the gene-level random effect, accumulatively they explained a significant faction of the variation of negative selection across missense mutations. Therefore, both variant features and gene-level constraints are important for predicting deleterious variants.

Missense constraints and loss-of-function constraints are weakly correlated across genes

The UNEECON-G score represents gene-level intolerance to missense mutations in human populations. Alternatively, selective constraint on a gene can be defined as its degree of intolerance to loss-of-function mutations, such as stop-gained, frameshift, and splice-site mutations [17, 28, 29]. We investigated the correlation between the UNEECON-G score and the pLI score, a metric of gene-level intolerance to loss-of-function mutations [17]. Surprisingly, the UNEECON-G score was only moderately correlated with the pLI score (Spearman's $\rho = 0.59$; Fig. 5a), suggesting a discrepancy between gene-level intolerance to missense mutations and that to loss-of-function mutations in a subset of protein-coding genes.

Intrinsically disordered proteins play key roles in a multitude of biological processes [52], but their sequences are poorly conserved across species [53]. Therefore, missense mutations in intrinsically disordered proteins may be under weak negative selection even if loss-of-function mutations in these proteins are deleterious. To test this hypothesis, we investigated the distributions of protein disorder content, *i.e.*, the fraction of disordered regions in a protein [49], across 1,912 protein-coding genes intolerant to heterozygous loss-of-function mutations (pLI score ≥ 0.9). We split the 1,912 genes into two equal-size groups based on their UNEECON-G scores, and defined the 956 genes with higher UNEECON-G scores as the gene set intolerant to both missense and loss-of-function mutations. Accordingly, we defined the 956 genes with lower UNEECON-G scores as the gene set tolerant to missense but not to loss-of-function mutations. Compared with the 956 genes intolerant to both missense and loss-of-function mutations, the disorder contents were significantly higher in the 956 genes tolerant to missense mutations but not to loss-of-function mutations (Fig. 5b). Therefore, disordered protein regions can at least partially explain the observed discrepancy between missense constraints and loss-of-function constraints.

We further investigated the enrichment of Reactome pathways [54] and Gene Ontology terms [55, 56] in the 956 genes intolerant to both missense and loss-of-function mutations, using the 956 genes tolerant to missense but not to loss-of-function mutations as a background. Several key pathways associated with central nervous system, gene regulation, cell cycle, and innate immune response were overrepresented in the genes intolerant to both missense and loss-of-function mutations (Fig. 5c; Table S4). We observed an enrichment of similar terms based on the Gene Ontology (Fig. S5 & S6). In agreement with the enrichment of pathways associated with central nervous system, the genes intolerant to both missense and loss-of-function mutations

were strongly enriched in the gene set implicated in autism spectrum disorder [57] (Fig. 5d).

Discussion

Here we present UNEECON for a unified prediction of deleterious missense mutations and highly constrained genes in the human genome. Compared with previously published methods, UNEECON shows superior performance in predicting missense variants and protein-coding genes associated with dominant genetic disorders. Therefore, UNEECON is a promising framework for both variant and gene prioritization. Furthermore, unlike supervised machine learning approaches, such as MPC [25], UNEECON integrates variant features and gene constraints based on signatures of negative selection instead of labeled disease variants. Therefore, UNEECON is unlikely to suffer from the circularity and the inflated performance commonly found in supervised methods [58].

It is worth noting that UNEECON is different from existing prioritization methods in multiple important aspects. First, unlike classical sequence conservation metrics [38, 39, 40, 41, 42] and integrative variant scores [13, 8, 14, 9, 10, 11, 12, 15], UNEECON infers the strength of negative selection based on rare genetic variants. Therefore, UNEECON predicts strongly deleterious variants with a dominant/semidominant mode of inheritance [59]. Second, unlike previously published gene constraint metrics that typically assign the same score to all missense variants within a gene [16, 17, 18, 19, 20, 21, 22, 23], UNEECON assigns different scores to the variants in the same gene in a feature-dependent manner, providing high-resolution maps of variant effects within genes. Third, UNEECON is able to adjust the distribution of variant scores within a gene according to the degree of depletion of missense variants in this gene, allowing for assigning different scores to missense mutations with similar variant features but located in different genes.

Similar to other metrics of negative selection, the performance of UNEECON strongly depends on the correlation between variant penetrance and negative selection. Our analysis of pathogenic variants suggests that UNEECON can predict deleterious variants associated with dominant genetic disorders but not necessarily those associated with recessive disorders. Also, because common variants are unlikely to be under strong negative selection [12], UNEECON scores might not be able to predict common variants associated with complex traits or late-onset diseases. In contrast, a recent study suggests that rare variants associated with complex traits are strongly enriched in coding regions and tend to be under negative selection [60], implying that UNEECON could be a useful tool for rare variant association studies.

UNEECON is based on a novel machine learning framework, deep mixed-effects model, to integrate variant features and gene-level constraints. Our analysis of the linear UNEECON model suggests that both the predictive variant features and the gene-level constraints are important for variant interpretation. In particular, gene-level evidence of selective constraints is the single most important predictor of negative selection on missense mutations. Also, predictive variant features accumulatively explain a large fraction of the variation of negative selection across missense variants. We expect that the combination of variant-level predictors and gene-level constraints will be an essential component in the future development of variant and gene prioritization methods.

By contrasting UNEECON-G scores against pLI scores, we observe an unexpected low correlation between gene-level intolerance to missense mutations and that to loss-of-function mutations. The prevalence of disordered protein regions in the human proteome is a key biological factor contributing to the low correlation between missense and loss-of-function constraints. Furthermore, we observe that the genes intolerant to both loss-of-function mutations and missense mutations may play key roles in central nervous system and autism spectrum disorder, highlighting the needs to investigate the function of these genes using state-of-the-art experimental techniques. By combing powerful variant and gene prioritization tools, such as UNEECON, and high-throughput mutagenesis and genome editing techniques [61, 62], we will obtain more insights into the function of these strongly constrained genes in the future.

Methods

Predictive variant features

UNEECON was trained on 30 missense variant features previously used to infer fitness effects of coding variants in the human genome ([12]; Table S1). These features can be classified into three categories: sequence conservation scores, protein structural features, and functional genomic features. The sequence conservation scores included SIFT [5], PROVEAN [7], SLR [63], Grantham [64], PSIC [65], LRT [66], MutationAssessor [4], HMMEntropy [67], and phyloP scores [40]. The protein structural features included predicted secondary structures, B-factors, contributions to protein stability, and relative solvent accessibilities from SNVBox [67]. The functional genomic features included the non-commercial version of SPIDEX splicing scores [68] and the maximum RNA-seq signals from the Roadmap Epigenomics Project [69]. Following a common practice in machine learning and statistics, we standardized continuous features by subtracting the mean and dividing by the standard deviation. All the features were based on the hg19 (GRCh37) assembly.

Population genomic data

We dowloaded whole genome variation data, exome variation data, and corresponding sequencing coverage data from the gnomAD browser [28, 29] (version 2.1.0). We only retained rare SNVs (MAF < 0.1%) that passed gnomAD's built-in quality filter for downstream analysis.

Context-dependent mutation model

We trained a context-dependent mutation model to capture the impact of 7-mer sequence context, local mutation rate, and sequencing coverage on the probability of the occurrence of each mutation in the gnomAD exome sequencing data. Because of the intrinsic sparsity of putatively neutral variants in coding regions, it is difficult to build a mutation model solely based on the gnomAD exome sequencing data. Therefore, we first built a mutation model based on neutral noncoding variants in the gnomAD whole genome sequencing (WGS) data. Then, we recalibrated the WGS-based mutation model in the gnomAD exome sequencing data to adjust for the differences in population sample size and sequencing coverage between the WGS and the exome sequencing data.

To build the WGS-based mutation model, we first compiled a list of putatively neutral noncoding regions following a strategy described in previous studies [70, 71, 10]. We removed coding exons [72], conserved phastCons elements [73], nucleotide sites within 1000 bp of any coding exons, and nucleotide sites within 100 bp of any phastCons elements. We assumed that the remaining noncoding regions were largely depleted with functional elements and, therefore, mutations in these regions were putatively neutral. It is worth noting that we also removed any nucleotide sites with an average sequencing coverage below 20, due to the difficulty of variant calling in low coverage regions, and any sites overlapping CpGs, due to the high mutation rates in these sites.

Then, we fit a WGS-based mutation model to the gnomAD WGS data in putatively neutral noncoding regions. First, for each possible combination of mutation l and sequence context k (the focal nucleotide with 3 flanking nucleotides on each side), we calculated its mutability, $f_{k \to l}$, defined as the proportion of 7-mer sequences of k with observed rare variant l in the gnomAD WGS data. We defined $F_{k \to l} = \mathrm{logit}(f_{k \to l}) \equiv \log(\frac{f_{k \to l}}{1 - f_{k \to l}})$, which represents the contribution of sequence context and mutation type to the probability of variant occurrence in logit scale. Second, we fit a genome-wide logistic regression model to estimate the contributions of sequencing depth and context-dependent mutability to the probability of the occurrence of a mutation. This logistic regression model assumed

$$\mathrm{logit}(Y_i^{\mathrm{WGS}}) = \alpha_0 + \alpha_1 \log(d_i) + \alpha_2 F_{k_i \to l_i}, \tag{1}$$

where Y_i^{WGS} is a binary indicator of the occurrence of neutral noncoding mutation i in the gnomAD WGS data. d_i is the average sequencing depth at the nucleotide position of mutation i. k_i and l_i are the sequence context and the mutation type of mutation i, respectively. We then fit a local logistic regression model for each exon m in the human genome with

$$\mathrm{logit}(Y_i^{\mathrm{WGS}}) = \alpha_3^m + \hat{\alpha}_1 \log(d_i) + \hat{\alpha}_2 F_{k_i \to l_i}, \text{ for all mutation } i's$$
$$\text{within 60 kb of exon } m \tag{2}$$

where α_3^m is an exon specific intercept independently estimated for each exon m, and $\hat{\alpha}_1$ and $\hat{\alpha}_2$ are the estimates of regression coefficients in equation 1. The local regression model effectively added an exon-specific, multiplicative scaling factor to adjust for the variation of local mutation rates across exons. We defined

$$q_i = \hat{\alpha}_3^m + \hat{\alpha}_1 \log(d_i) + \hat{\alpha}_2 F_{k_i \to l_i} \tag{3}$$

as the logit of the occurrence probability of mutation i in exon m in the WGS-based mutation model.

Finally, we recalibrated the WGS-based mutation model in the gnomAD exome sequencing data with a logistic regression model,

$$\mathrm{logit}(Y_i^{\mathrm{exome}}) = \beta_0 + q_i, \tag{4}$$

where Y_i^{exome} is a binary indicator of the presence of synonymous mutation i in the gnomAD exome sequencing data. The exome-based logistic regression model effectively added a multiplicative scaling factor to accommodate for the differences in population sample size and sequencing coverage between the WGS and the exome sequencing data. In the final exome mutation model, we defined the probability of the occurrence of missense mutation i in gene j as

$$\mu_{ij} = \text{logistic}(\hat{\beta}_0 + q_i) \equiv \frac{\exp(\hat{\beta}_0 + q_i)}{1 + \exp(\hat{\beta}_0 + q_i)}, \tag{5}$$

where $\hat{\beta}_0$ is the maximum likelihood estimate of β_0 in equation 4. We fit all the logistic regression models using the glm function in R [74].

Details of the UNEECON model

We assume that negative selection on a potential mutation is a mathematical function of the sum of a variant-level fixed effect, z_{ij}, and a gene-level random effect, u_j (Fig. 1). Denoting $\mathbf{X}_{ij}$ as the vector of variant features associated with mutation i in gene j, we assume that the variant-level fixed effect, z_{ij}, can be modeled by a feedforward neural network. At the bottom of the neural network, a nonlinear hidden layer and a dropout layer are employed to transform $\mathbf{X}_{ij}$ to a hidden vector,

$$\mathbf{H}_{ij} = \text{droput}(\text{ReLU}(\mathbf{X}_{ij} \cdot \mathbf{W}_{\text{hidden}} + \mathbf{B}_{\text{hidden}})), \tag{6}$$

where **ReLU** and **dropout** are the rectified linear layer [75] and the dropout layer [43], respectively. $\mathbf{W}_{\text{hidden}}$ and $\mathbf{B}_{\text{hidden}}$ are the weight matrix and bias vector of the hidden layer, respectively. Then, we assume the fixed effect term, z_{ij}, is a linear combination of the hidden vector $\mathbf{H}_{ij}$,

$$z^{ij} = \mathbf{H}_{ij} \cdot \mathbf{W}_{\text{output}} + b_{\text{output}}, \tag{7}$$

where $\mathbf{W}_{\text{output}}$ and b_{output} are the weight vector and the bias term, respectively. We use the Glorot method [76] to initialize the weights, $\mathbf{W}_{\text{hidden}}$ and $\mathbf{W}_{\text{output}}$, and initialize the bias terms, $\mathbf{B}_{\text{hidden}}$ and b_{output}, with zeros. In the dropout layer, we use a fixed dropout rate of 0.5. It is worth noting that, if a linear version of the UNEECON model is used, equation 6 will be replaced by an identity function $\mathbf{H}_{ij} \equiv \mathbf{X}_{ij}$.

To capture the variation of gene-level selective constraints that cannot be predicted from feature vector $\mathbf{X}_{ij}$, we introduce a gene-level random-effect term following a Gaussian distribution,

$$u_j \sim \mathcal{N}(0, \sigma), \tag{8}$$

where σ is the standard deviation of the Gaussian distribution. Given the fixed and random effect terms, we assume η_{ij}, the relative probability of the occurrence of mutation i in gene j, follows

$$\eta_{ij} = \text{logistic}(z_{ij} + u_j) \equiv \frac{\exp(z_{ij} + u_j)}{1 + \exp(z_{ij} + u_j)}. \tag{9}$$

We further assume that the total probability of the occurrence of potential missense mutation i in gene j is equal to the product of η_{ij} and μ_{ij}, the probability of variant occurrence under the neutral mutation model described in equation 5. Accordingly, the likelihood function for the data associated with gene j is defined as

$$\mathcal{L}_j = \prod_i (\eta_{ij}\mu_{ij})^{Y_{ij}}(1 - \eta_{ij}\mu_{ij})^{1-Y_{ij}}, \tag{10}$$

where Y_{ij} is a binary indicator of the occurrence of missense mutation i in gene j in the gnomAD exome sequencing data.

Training the UNEECON model

We trained UNEECON with the Adam optimization algorithm [77]. In each iteration, UNEECON loaded a mini-batch of data consisting of all the potential missense mutations in a single gene. Then, we calculated the log likelihood of the mini-batch of data using equation 10 and numerically integrated out u_j with a 20-point Gaussian Quadrature rule. The negative log value of equation 10 was used as the objective function in the Adam optimizer.

We randomly split the data into a training set (80% genes), a validation set (10% genes), and a test set (10% genes). After each epoch of training, UNEECON evaluated the objective function in the validation set and stopped training when the objective function did not improve over 5 successive epochs to avoid overfitting (early stopping). Furthermore, we performed a grid search to optimize two hyperparameters, $i.e.$, the learning rate of the Adam algorithm (10^{-2}, 10^{-3}, and 10^{-4}) and the number of hidden units (64, 128, 256, 512, and a linear model without hidden units). The model was evaluated on the test dataset to choose optimal hyperparameters.

Calculating UNEECON and UNEECON-G scores

After training, we fixed the free parameters, $i.e.$, $\mathbf{W}_{\text{hidden}}$, $\mathbf{W}_{\text{output}}$, $\mathbf{B}_{\text{hidden}}$, b_{output}, and σ, to the estimated values. Then, we calculated the UNEECON score for all the potential missense mutations in the human genome. For each gene j, UNEECON first calculated the posterior distribution of the random effect, $\mathbb{P}(u_j|\text{Data}_j)$, where $\text{Data}_j = \{\mathbf{X}_{1j}, \mathbf{X}_{2j}, \cdots, \mathbf{X}_{Nj}, Y_{1j}, Y_{2j}, \cdots, Y_{Nj}\}$ are the features and indicators of occurrence of all the potential missense mutations in gene j. Denoting $\mathbb{E}_{u_j|\text{Data}_j}(\eta_{ij}|u_j) \equiv \int \eta_{ij}\mathbb{P}(u_j|\text{Data}_j)du_j$ as expected relative probability of the occurrence of mutation i in gene j, we define the UNEECON score of mutation i as $1 - \mathbb{E}_{u_j|\text{Data}}(\eta_{ij}|u_j)$. The UNEECON-G score is defined as the average UNEECON score of all the missense mutations in a gene.

Distributions of UNEECON scores across gene categories and protein regions

We investigated the distributions of UNEECON scores across different gene categories and protein regions. We obtained lists of haploinsufficient genes [37] ($n = 294$), autosomal dominant disease genes [35, 36] ($n = 709$), autosomal recessive disease genes [35, 36] ($n = 1,183$), and olfactory receptor genes [45] ($n = 371$) from the GitHub repository for the MacArthur Lab at the Broad Institute (https://github.com/macarthur-lab/gene_lists). We obtained annotations of

α-helices, β-strands, hydrogen-bonded turns, enzyme active sites, and binding sites from UniProt [48]. We obtained annotated disordered protein regions from MobiDB 3.0 [49]. We plotted the distributions of UNEECON scores using R [74].

Comparison with other methods in the prediction of ClinVar pathogenic variants

We evaluated the performance of UNEECON, compared with eight previously published methods, in the setting of predicting missense pathogenic variants in the ClinVar database downloaded on Feb 25, 2019 [30]. We obtained MPC [25], PrimateAI [11], Eigen [50], CADD [8], RVIS [16], and pLI [17] scores from dbNSFP (version 4.0b2a) [78]. We obtained LASSIE scores [12] from the LASSIE GitHub repository (https://github.com/CshlSiepelLab/LASSIE) and CCR score (version 2) [22] from https://s3.us-east-2.amazonaws.com/ccrs/ccrs/ccrs.autosomes.v2.20180420.bed.gz. It is worth noting that we used the version of pLI scores trained on the gnomAD exome sequencing data [29].

We considered autosomal missense variants with annotations of "pathogenic/likely pathogenic" as positives and the ones with annotations of "benign/likely benign" as negative controls. We further removed the variants with conflict pathogenicity annotations and the variants present in the gnomAD exome sequencing dataset [29]. For all the comparisons, we only included the variants scored by all methods. We plotted the receiver operating characteristic (ROC) curves and calculated the areas under the receiver operating characteristic curves (AUCs) using the ROCR package [79]. Because the AUC metrics is sensitive to label imbalance, we matched the numbers of positives and negatives using random sampling without replacement.

Comparison with alternative methods in predicting *de novo* mutations associated with developmental disorders

We downloaded *de novo* mutations identified in 4,293 individuals affected by developmental disorders [31] and 2,278 healthy individuals from denovo-db (version v1.6.1). The healthy controls included individuals enrolled in previous studies of autism spectrum disorder [80, 81, 82, 83, 84], severe non-syndromic sporadic intellectual disability [85], schizophrenia [86], and healthy populations [87, 88, 89, 90]. We removed redundant variants and any variants presented in the gnomAD exome sequencing data. Then, for various percentile rank cutoffs (top 10%, 20%, 30%, and 40%), we calculated the $\log_2$ odds ratio of the enrichment of predicted deleterious variants in affected individuals using the fisher.test function in R [74].

Comparison with alternative methods in predicting disease genes and essential genes

We obtained four sets of disease genes and essential genes from the GitHub repository for the MacArthur Lab at the Broad Institute (https://github.com/macarthur-lab/gene_lists). These gene sets included 294 haploinsufficient genes [37], 709 autosomal dominant disease genes [35, 36], 2,454 human orthologs of mouse essential genes [33, 34], and 683 essential genes based on CRISPR screen data from human cell lines [32]. For each set of the haploinsufficient, autosomal dominant, and mouse essential genes, we utilized the other autosomal genes as negative controls and constructed a negative gene set with matched gene length and gene number using MatchIt [91]. Similarly, for the 683 human essential genes based on CRISPR

screen data, we used the 913 nonessential genes from the same study [32] as negative controls and constructed a negative gene set with matched gene length and gene number. We plotted the ROC curves and calculated the AUC metrics using the ROCR package [79].

Comparison of UNEECON-G and pLI scores

To evaluate the relationship between gene-level intolerance to missense mutations and that to loss-of-function mutations, we compared the UNEECON-G scores with the pLI scores trained on the gnomAD data [17, 78]. We identified 1,912 genes intolerant to loss-of-function mutations based on a threshold of pLI score of 0.9. Then, we split the 1,912 genes into 956 genes tolerant to missense genes and 956 genes intolerant to missense mutations based on the median UNEECON-G score. We used PANTHER [92] to calculate the enrichment of Reactome pathways and Gene Ontology categories in the set of genes intolerant to both missense and loss-of-function mutations, compared with those tolerant to missense constraints but not to loss-of-function mutations. We also downloaded 1,054 autism related genes from the SFARI database on March 9, 2019 [57], and evaluated the enrichment of autism genes in the two gene sets.

Availability of data and materials

The UNEECON program and precomputed UNEECON/UNEECON-G scores are available at https://github.com/yifei-lab/UNEECON.

Competing interests
The author declares that he has no competing interests.

Author's contributions
Y.-F.H. conceived the study, implemented the UNEECON model, analyzed the data, and wrote the manuscript.

Acknowledgements
The author thanks Noah Dukler for comments on the manuscript. This work was supported by start-up fund from the Pennsylvania State University.

References
1. Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., Grody, W.W., Hegde, M., Lyon, E., Spector, E., Voelkerding, K., Rehm, H.L.: Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the american college of medical genetics and genomics and the association for molecular pathology. Genetics in Medicine **17**(5), 405–423 (2015)
2. Maxwell, K., Hart, S., Vijai, J., Schrader, K., Slavin, T., Thomas, T., Wubbenhorst, B., Ravichandran, V., Moore, R., Hu, C., Guidugli, L., Wenz, B., Domchek, S., Robson, M., Szabo, C., Neuhausen, S., Weitzel, J., Offit, K., Couch, F., Nathanson, K.: Evaluation of acmg-guideline-based variant classification of cancer susceptibility and non-cancer-associated genes in families affected by breast cancer. The American Journal of Human Genetics **98**(5), 801–817 (2016)
3. Eilbeck, K., Quinlan, A., Yandell, M.: Settling the score: variant prioritization and mendelian disease. Nature Reviews Genetics **18**(10), 599–612 (2017)
4. Reva, B., Antipin, Y., Sander, C.: Predicting the functional impact of protein mutations: application to cancer genomics. Nucleic Acids Research **39**(17), 118 (2011)
5. Ng, P.C., Henikoff, S.: SIFT: predicting amino acid changes that affect protein function. Nucleic Acids Research **31**(13), 3812–3814 (2003)
6. Cooper, G.M., Shendure, J.: Needles in stacks of needles: finding disease-causal variants in a wealth of genomic data. Nature Reviews Genetics **12**(9), 628–640 (2011)
7. Choi, Y., Sims, G.E., Murphy, S., Miller, J.R., Chan, A.P.: Predicting the functional effect of amino acid substitutions and indels. PLOS ONE **7**(10), 1–13 (2012)
8. Kircher, M., Witten, D.M., Jain, P., O'Roak, B.J., Cooper, G.M., Shendure, J.: A general framework for estimating the relative pathogenicity of human genetic variants. Nature Genetics **46**(3), 310–315 (2014)
9. Gulko, B., Hubisz, M.J., Gronau, I., Siepel, A.: A method for calculating probabilities of fitness consequences for point mutations across the human genome. Nature Genetics **47**(3), 276–283 (2015)
10. Huang, Y.-F., Gulko, B., Siepel, A.: Fast, scalable prediction of deleterious noncoding variants from functional and population genomic data. Nature Genetics **49**(4), 618–624 (2017)

11. Sundaram, L., Gao, H., Padigepati, S.R., McRae, J.F., Li, Y., Kosmicki, J.A., Fritzilas, N., Hakenberg, J., Dutta, A., Shon, J., Xu, J., Batzloglou, S., Li, X., Farh, K.K.-H.: Predicting the clinical impact of human mutation with deep neural networks. Nature Genetics, (2018)

12. Huang, Y.-F., Siepel, A.: Estimation of allele-specific fitness effects across human protein-coding sequences and implications for disease. Genome Research (2019). doi:10.1101/gr.245522.118

13. Khurana, E., Fu, Y., Colonna, V., Mu, X.J., Kang, H.M., Lappalainen, T., Sboner, A., Lochovsky, L., Chen, J., Harmanci, A., Das, J., Abyzov, A., Balasubramanian, S., Beal, K., Chakravarty, D., Challis, D., Chen, Y., Clarke, D., Clarke, L., Cunningham, F., Evani, U.S., Flicek, P., Fragoza, R., Garrison, E., Gibbs, R., Gumus, Z.H., Herrero, J., Kitabayashi, N., Kong, Y., Lage, K., Liluashvili, V., Lipkin, S.M., MacArthur, D.G., Marth, G., Muzny, D., Pers, T.H., Ritchie, G.R.S., Rosenfeld, J.A., Sisu, C., Wei, X., Wilson, M., Xue, Y., Yu, F., Consortium, .G.P., Dermitzakis, E.T., Yu, H., Rubin, M.A., Tyler-Smith, C., Gerstein, M.: Integrative annotation of variants from 1092 humans: Application to cancer genomics. Science **342**(6154) (2013)

14. Fu, Y., Liu, Z., Lou, S., Bedford, J., Mu, X., Yip, K., Khurana, E., Gerstein, M.: FunSeq2: a framework for prioritizing noncoding regulatory variants in cancer. Genome Biology **15**(10), 480 (2014)

15. Gulko, B., Siepel, A.: An evolutionary framework for measuring epigenomic information and estimating cell-type-specific fitness consequences. Nature Genetics **51**(2), 335–342 (2019)

16. Petrovski, S., Wang, Q., Heinzen, E.L., Allen, A.S., Goldstein, D.B.: Genic intolerance to functional variation and the interpretation of personal genomes. PLOS Genetics **9**(8), 1003709 (2013)

17. Samocha, K.E., Robinson, E.B., Sanders, S.J., Stevens, C., Sabo, A., McGrath, L.M., Kosmicki, J.A., Rehnstrom, K., Mallick, S., Kirby, A., Wall, D.P., MacArthur, D.G., Gabriel, S.B., DePristo, M., Purcell, S.M., Palotie, A., Boerwinkle, E., Buxbaum, J.D., Cook Jr, E.H., Gibbs, R.A., Schellenberg, G.D., Sutcliffe, J.S., Devlin, B., Roeder, K., Neale, B.M., Daly, M.J.: A framework for the interpretation of de novo mutation in human disease. Nature Genetics **46**(9), 944–950 (2014)

18. Petrovski, S., Gussow, A.B., Wang, Q., Halvorsen, M., Han, Y., Weir, W.H., Allen, A.S., Goldstein, D.B.: The intolerance of regulatory sequence to genetic variation predicts gene dosage sensitivity. PLoS Genet **11**(9), 1005492 (2015)

19. Itan, Y., Shang, L., Boisson, B., Patin, E., Bolze, A., Moncada-Vélez, M., Scott, E., Ciancanelli, M.J., Lafaille, F.G., Markle, J.G., Martinez-Barricarte, R., de Jong, S.J., Kong, X.-F., Nitschke, P., Belkadi, A., Bustamante, J., Puel, A., Boisson-Dupuis, S., Stenson, P.D., Gleeson, J.G., Cooper, D.N., Quintana-Murci, L., Claverie, J.-M., Zhang, S.-Y., Abel, L., Casanova, J.-L.: The human gene damage index as a gene-level approach to prioritizing exome variants. Proceedings of the National Academy of Sciences (2015)

20. Gussow, A., Petrovski, S., Wang, Q., Allen, A., Goldstein, D.: The intolerance to functional genetic variation of protein domains predicts the localization of pathogenic mutations within genes. Genome Biology **17**(1), 9 (2016)

21. Pérez-Palma, E., May, P., Iqbal, S., Niestroj, L.-M., Du, J., Heyne, H., Castrillon, J., O'Donnell-Luria, A., Nürnberg, P., Palotie, A., Daly, M., Lal, D.: Identification of pathogenic variant enriched regions across genes and gene families. bioRxiv, 641043 (2019)

22. Havrilla, J.M., Pedersen, B.S., Layer, R.M., Quinlan, A.R.: A map of constrained coding regions in the human genome. Nature Genetics **51**(1), 88–95 (2019)

23. Silk, M., Petrovski, S., Ascher, D.B.: MTR-Viewer: identifying regions within genes under purifying selection. Nucleic Acids Research **47**(W1), 121–126 (2019)

24. Iossifov, I., Levy, D., Allen, J., Ye, K., Ronemus, M., Lee, Y.-h., Yamrom, B., Wigler, M.: Low load for disruptive mutations in autism genes and their biased transmission. Proceedings of the National Academy of Sciences **112**(41), 5600–5607 (2015)

25. Samocha, K.E., Kosmicki, J.A., Karczewski, K.J., O'Donnell-Luria, A.H., Pierce-Hoffman, E., MacArthur, D.G., Neale, B.M., Daly, M.J.: Regional missense constraint improves variant deleteriousness prediction. bioRxiv (2017). doi:10.1101/148353

26. Jagadeesh, K.A., Wenger, A.M., Berger, M.J., Guturu, H., Stenson, P.D., Cooper, D.N., Bernstein, J.A., Bejerano, G.: M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. Nature Genetics **48**(12), 1581–1586 (2016)

27. Evans, P., Wu, C., Lindy, A., McKnight, D.A., Lebo, M., Sarmady, M., Abou Tayoun, A.N.: Genetic variant pathogenicity prediction trained using disease-specific clinical sequencing data sets. Genome Research **29**(7), 1144–1151 (2019)

28. Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., ODonnell-Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., Tukiainen, T., Birnbaum, D.P., Kosmicki, J.A., Duncan, L.E., Estrada, K., Zhao, F., Zou, J., Pierce-Hoffman, E., Berghout, J., Cooper, D.N., Deflaux, N., DePristo, M., Do, R., Flannick, J., Fromer, M., Gauthier, L., Goldstein, J., Gupta, N., Howrigan, D., Kiezun, A., Kurki, M.I., Moonshine, A.L., Natarajan, P., Orozco, L., Peloso, G.M., Poplin, R., Rivas, M.A., Ruano-Rubio, V., Rose, S.A., Ruderfer, D.M., Shakir, K., Stenson, P.D., Stevens, C., Thomas, B.P., Tiao, G., Tusie-Luna, M.T., Weisburd, B., Won, H.-H., Yu, D., Altshuler, D.M., Ardissino, D., Boehnke, M., Danesh, J., Donnelly, S., Elosua, R., Florez, J.C., Gabriel, S.B., Getz, G., Glatt, S.J., Hultman, C.M., Kathiresan, S., Laakso, M., McCarroll, S., McCarthy, M.I., McGovern, D., McPherson, R., Neale, B.M., Palotie, A., Purcell, S.M., Saleheen, D., Scharf, J.M., Sklar, P., Sullivan, P.F., Tuomilehto, J., Tsuang, M.T., Watkins, H.C., Wilson, J.G., Daly, M.J., MacArthur, D.G., Consortium, E.A.: Analysis of protein-coding genetic variation in 60,706 humans. Nature **536**(7616), 285–291 (2016)

29. Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alfoldi, J., Wang, Q., Collins, R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., Gauthier, L.D., Brand, H., Solomonson, M., Watts, N.A., Rhodes, D., Singer-Berk, M., Seaby, E.G., Kosmicki, J.A., Walters, R.K., Tashman, K., Farjoun, Y., Banks, E., Poterba, T., Wang, A., Seed, C., Whiffin, N., Chong, J.X., Samocha, K.E., Pierce-Hoffman, E., Zappala, Z., ODonnell-Luria, A.H., Vallabh Minikel, E., Weisburd, B., Lek, M., Ware, J.S., Vittal, C., Armean, I.M., Bergelson, L., Cibulskis, K., Connolly, K.M., Covarrubias, M., Donnelly, S., Ferriera, S., Gabriel, S., Gentry, J., Gupta, N., Jeandet, T., Kaplan, D., Llanwarne, C., Munshi, R., Novod, S., Petrillo, N., Roazen, D., Ruano-Rubio, V., Saltzman, A.,

Schleicher, M., Soto, J., Tibbetts, K., Tolonen, C., Wade, G., Talkowski, M.E., Neale, B.M., Daly, M.J., MacArthur, D.G.: Variation across 141,456 human exomes and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding genes. bioRxiv (2019). doi:10.1101/531210

30. Landrum, M.J., Lee, J.M., Riley, G.R., Jang, W., Rubinstein, W.S., Church, D.M., Maglott, D.R.: ClinVar: public archive of relationships among sequence variation and human phenotype. Nucleic Acids Research **42**(D1), 980–985 (2014)

31. Deciphering Developmental Disorders Study, McRae, J.F., Clayton, S., Fitzgerald, T.W., Kaplanis, J., Prigmore, E., Rajan, D., Sifrim, A., Aitken, S., Akawi, N., Alvi, M., Ambridge, K., Barrett, D.M., Bayzetinova, T., Jones, P., Jones, W.D., King, D., Krishnappa, N., Mason, L.E., Singh, T., Tivey, A.R., Ahmed, M., Anjum, U., Archer, H., Armstrong, R., Awada, J., Balasubramanian, M., Banka, S., Baralle, D., Barnicoat, A., Batstone, P., Baty, D., Bennett, C., Berg, J., Bernhard, B., Bevan, A.P., Bitner-Glindzicz, M., Blair, E., Blyth, M., Bohanna, D., Bourdon, L., Bourn, D., Bradley, L., Brady, A., Brent, S., Brewer, C., Brunstrom, K., Bunyan, D.J., Burn, J., Canham, N., Castle, B., Chandler, K., Chatzimichali, E., Cilliers, D., Clarke, A., Clasper, S., Clayton-Smith, J., Clowes, V., Coates, A., Cole, T., Colgiu, I., Collins, A., Collinson, M.N., Connell, F., Cooper, N., Cox, H., Cresswell, L., Cross, G., Crow, Y., DAlessandro, M., Dabir, T., Davidson, R., Davies, S., de Vries, D., Dean, J., Deshpande, C., Devlin, G., Dixit, A., Dobbie, A., Donaldson, A., Donnai, D., Donnelly, D., Donnelly, C., Douglas, A., Douzgou, S., Duncan, A., Eason, J., Ellard, S., Ellis, I., Elmslie, F., Evans, K., Everest, S., Fendick, T., Fisher, R., Flinter, F., Foulds, N., Fry, A., Fryer, A., Gardiner, C., Gaunt, L., Ghali, N., Gibbons, R., Gill, H., Goodship, J., Goudie, D., Gray, E., Green, A., Greene, P., Greenhalgh, L., Gribble, S., Harrison, R., Harrison, L., Harrison, V., Hawkins, M., He, L., Hellens, S., Henderson, A., Hewitt, S., Hildyard, L., Hobson, E., Holden, S., Holder, M., Holder, S., Hollingsworth, G., Homfray, T., Humphreys, M., Hurst, J., Hutton, B., Ingram, S., Irving, M., Islam, L., Jackson, A., Jarvis, J., Jenkins, L., Johnson, D., Jones, E., Josifova, D., Joss, S., Kaemba, B., Kazembe, S., Kelsell, R., Kerr, B., Kingston, H., Kini, U., Kinning, E., Kirby, G., Kirk, C., Kivuva, E., Kraus, A., Kumar, D., Kumar, V.K.A., Lachlan, K., Lam, W., Lampe, A., Langman, C., Lees, M., Lim, D., Longman, C., Lowther, G., Lynch, S.A., Magee, A., Maher, E., Male, A., Mansour, S., Marks, K., Martin, K., Maye, U., McCann, E., McConnell, V., McEntagart, M., McGowan, R., McKay, K., McKee, S., McMullan, D.J., McNerlan, S., McWilliam, C., Mehta, S., Metcalfe, K., Middleton, A., Miedzybrodzka, Z., Miles, E., Mohammed, S., Montgomery, T., Moore, D., Morgan, S., Morton, J., Mugalaasi, H., Murday, V., Murphy, H., Naik, S., Nemeth, A., Nevitt, L., Newbury-Ecob, R., Norman, A., OShea, R., Ogilvie, C., Ong, K.-R., Park, S.-M., Parker, M.J., Patel, C., Paterson, J., Payne, S., Perrett, D., Phipps, J., Pilz, D.T., Pollard, M., Pottinger, C., Poulton, J., Pratt, N., Prescott, K., Price, S., Pridham, A., Procter, A., Purnell, H., Quarrell, O., Ragge, N., Rahbari, R., Randall, J., Rankin, J., Raymond, L., Rice, D., Robert, L., Roberts, E., Roberts, J., Roberts, P., Roberts, G., Ross, A., Rosser, E., Saggar, A., Samant, S., Sampson, J., Sandford, R., Sarkar, A., Schweiger, S., Scott, R., Scurr, I., Selby, A., Seller, A., Sequeira, C., Shannon, N., Sharif, S., Shaw-Smith, C., Shearing, E., Shears, D., Sheridan, E., Simonic, I., Singzon, R., Skitt, Z., Smith, A., Smith, K., Smithson, S., Sneddon, L., Splitt, M., Squires, M., Stewart, F., Stewart, H., Straub, V., Suri, M., Sutton, V., Swaminathan, G.J., Sweeney, E., Tatton-Brown, K., Taylor, C., Taylor, R., Tein, M., Temple, I.K., Thomson, J., Tischkowitz, M., Tomkins, S., Torokwa, A., Treacy, B., Turner, C., Turnpenny, P., Tysoe, C., Vandersteen, A., Varghese, V., Vasudevan, P., Vijayarangakannan, P., Vogt, J., Wakeling, E., Wallwark, S., Waters, A., Weber, A., Wellesley, D., Whiteford, M., Widaa, S., Wilcox, S., Wilkinson, E., Williams, D., Williams, N., Wilson, L., Woods, G., Wragg, C., Wright, M., Yates, L., Yau, M., Nellaker, C., Parker, M., Firth, H.V., Wright, C.F., FitzPatrick, D.R., Barrett, J.C., Hurles, M.E.: Prevalence and architecture of de novo mutations in developmental disorders. Nature **542**, 433 (2017)

32. Hart, T., Brown, K.R., Sircoulomb, F., Rottapel, R., Moffat, J.: Measuring error rates in genomic perturbation screens: gold standards for human functional genomics. Mol Syst Biol **10**(7), 733 (2014)

33. Blake, J.A., Bult, C.J., Kadin, J.A., Richardson, J.E., Eppig, J.T., the Mouse Genome Database Group: The Mouse Genome Database (MGD): premier model organism resource for mammalian genomics and genetics. Nucleic Acids Research **39**(suppl1), 842–848 (2010)

34. Georgi, B., Voight, B.F., Bucan, M.: From mouse to human: Evolutionary genomics analysis of human orthologs of essential genes. PLOS Genetics **9**(5), 1003484 (2013)

35. Blekhman, R., Man, O., Herrmann, L., Boyko, A.R., Indap, A., Kosiol, C., Bustamante, C.D., Teshima, K.M., Przeworski, M.: Natural selection on genes that underlie human disease susceptibility. Current Biology **18**(12), 883–889 (2008)

36. Berg, J.S., Adams, M., Nassar, N., Bizon, C., Lee, K., Schmitt, C.P., Wilhelmsen, K.C., Evans, J.P.: An informatics approach to analyzing the incidentalome. Genetics In Medicine **15**, 36 (2012)

37. Rehm, H.L., Berg, J.S., Brooks, L.D., Bustamante, C.D., Evans, J.P., Landrum, M.J., Ledbetter, D.H., Maglott, D.R., Martin, C.L., Nussbaum, R.L., Plon, S.E., Ramos, E.M., Sherry, S.T., Watson, M.S.: ClinGen – the clinical genome resource. N Engl J Med **372**(23), 2235–2242 (2015)

38. Armon, A., Graur, D., Ben-Tal, N.: ConSurf: an algorithmic tool for the identification of functional regions in proteins by surface mapping of phylogenetic information. Journal of Molecular Biology **307**(1), 447–463 (2001)

39. Cooper, G.M., Stone, E.A., Asimenos, G., Green, E.D., Batzoglou, S., Sidow, A.: Distribution and intensity of constraint in mammalian genomic sequence. Genome Research **15**(7), 901–913 (2005)

40. Pollard, K.S., Hubisz, M.J., Rosenbloom, K.R., Siepel, A.: Detection of nonneutral substitution rates on mammalian phylogenies. Genome Research **20**(1), 110–121 (2010)

41. Huang, Y.-F., Golding, G.B.: Phylogenetic Gaussian process model for the inference of functionally important regions in protein tertiary structures. PLoS Computational Biology **10**(1), 1003429 (2014)

42. Huang, Y.-F., Golding, G.B.: Funcpatch: a web server for the fast bayesian inference of conserved functional patches in protein 3d structures. Bioinformatics **31**(4), 523–531 (2015)

43. Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., Salakhutdinov, R.: Dropout: A simple way to prevent neural networks from overfitting. Journal of Machine Learning Research **15**, 1929–1958 (2014)

44. Bengio, Y.: Practical recommendations for gradient-based training of deep architectures. In: Neural Networks: Tricks of the Trade: Second Edition, pp. 437–478. Springer, Berlin, Heidelberg (2012)

45. Mainland, J.D., Li, Y.R., Zhou, T., Liu, W.L.L., Matsunami, H.: Human olfactory receptor responses to odorants. Scientific Data **2**, 150002 (2015)
46. Gilad, Y., Bustamante, C.D., Lancet, D., Pääbo, S.: Natural selection on the olfactory receptor gene family in humans and chimpanzees. The American Journal of Human Genetics **73**(3), 489–501 (2003)
47. The UniProt Consortium: UniProt: a worldwide hub of protein knowledge. Nucleic Acids Research **47**(D1), 506–515 (2018)
48. McGarvey, P.B., Nightingale, A., Luo, J., Huang, H., Martin, M.J., Wu, C., Consortium, U.: Uniprot genomic mapping for deciphering functional effects of missense variants. Human mutation (2019)
49. Piovesan, D., Tabaro, F., Paladin, L., Necci, M., Mičetić, I., Camilloni, C., Davey, N., Dosztányi, Z., Mészáros, B., Monzon, A.M., Parisi, G., Schad, E., Sormanni, P., Tompa, P., Vendruscolo, M., Vranken, W.F., Tosatto, S.C.E.: Mobidb 3.0: more annotations for intrinsic disorder, conformational diversity and interactions in proteins. Nucleic Acids Research **46**(D1), 471–476 (2017)
50. Ionita-Laza, I., McCallum, K., Xu, B., Buxbaum, J.D.: A spectral approach integrating functional genomic annotations for coding and noncoding variants. Nature Genetics **48**(2), 214–220 (2016)
51. Turner, T.N., Yi, Q., Krumm, N., Huddleston, J., Hoekzema, K., F. Stessman, H.A., Doebley, A.-L., Bernier, R.A., Nickerson, D.A., Eichler, E.E.: denovo-db: a compendium of human de novo variants. Nucleic Acids Research **45**(D1), 804–811 (2016)
52. Wright, P.E., Dyson, H.J.: Intrinsically disordered proteins in cellular signalling and regulation. Nature Reviews Molecular Cell Biology **16**, 18 (2015)
53. Brown, C.J., Takayama, S., Campen, A.M., Vise, P., Marshall, T.W., Oldfield, C.J., Williams, C.J., Keith Dunker, A.: Evolutionary rate heterogeneity in proteins with long disordered regions. Journal of Molecular Evolution **55**(1), 104–110 (2002)
54. Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P., Haw, R., Jassal, B., Korninger, F., May, B., Milacic, M., Roca, C.D., Rothfels, K., Sevilla, C., Shamovsky, V., Shorser, S., Varusai, T., Viteri, G., Weiser, J., Wu, G., Stein, L., Hermjakob, H., D'Eustachio, P.: The Reactome pathway knowledgebase. Nucleic Acids Research **46**(D1), 649–655 (2018)
55. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G.: Gene ontology: tool for the unification of biology. the gene ontology consortium. Nature Genetics **25**(1), 25–29 (2000)
56. The Gene Ontology Consortium: The gene ontology resource: 20 years and still going strong. Nucleic Acids Research **47**(D1), 330–338 (2019)
57. Abrahams, B.S., Arking, D.E., Campbell, D.B., Mefford, H.C., Morrow, E.M., Weiss, L.A., Menashe, I., Wadkins, T., Banerjee-Basu, S., Packer, A.: SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). Molecular Autism **4**(1), 36 (2013)
58. Grimm, D.G., Azencott, C.-A., Aicheler, F., Gieraths, U., MacArthur, D.G., Samocha, K.E., Cooper, D.N., Stenson, P.D., Daly, M.J., Smoller, J.W., Duncan, L.E., Borgwardt, K.M.: The evaluation of tools used to predict the impact of missense variants is hindered by two types of circularity. Human Mutation **36**(5), 513–523 (2015)
59. Fuller, Z.L., Berg, J.J., Mostafavi, H., Sella, G., Przeworski, M.: Measuring intolerance to mutation in human genetics. Nature Genetics (2019)
60. Wainschtein, P., Jain, D.P., Yengo, L., Zheng, Z., Cupples, L.A., Shadyab, A.H., McKnight, B., Shoemaker, B.M., Mitchell, B.D., Psaty, B.M., Kooperberg, C., Roden, D., Darbar, D., Arnett, D.K., Regan, E.A., Boerwinkle, E., Rotter, J.I., Allison, M.A., McDonald, M.-L.N., Chung, M.K., Smith, N.L., Ellinor, P.T., Vasan, R.S., Mathias, R.A., Rich, S.S., Heckbert, S.R., Redline, S., Guo, X., Chen, Y.-D.I., Liu, C.-T., de Andrade, M., Yanek, L.R., Albert, C.M., Hernandez, R.D., McGarvey, S.T., North, K.E., Lange, L.A., Weir, B.S., Laurie, C.C., Yang, J., Visscher, P.M.: Recovery of trait heritability from whole genome sequence data. bioRxiv (2019)
61. Starita, L.M., Ahituv, N., Dunham, M.J., Kitzman, J.O., Roth, F.P., Seelig, G., Shendure, J., Fowler, D.M.: Variant interpretation: Functional assays to the rescue. The American Journal of Human Genetics **101**(3), 315–325 (2017)
62. Kinney, J.B., McCandlish, D.M.: Massively parallel assays and quantitative sequence-function relationships. Annu. Rev. Genom. Hum. Genet. (2019)
63. Massingham, T., Goldman, N.: Detecting amino acid sites under positive selection and purifying selection. GENETICS **169**(3), 1753–1762 (2005). doi:10.1534/genetics.104.032144
64. Grantham, R.: Amino acid difference formula to help explain protein evolution. Science **185**(4154), 862–864 (1974)
65. Adzhubei, I.A., Schmidt, S., Peshkin, L., Ramensky, V.E., Gerasimova, A., Bork, P., Kondrashov, A.S., Sunyaev, S.R.: A method and server for predicting damaging missense mutations. Nature Methods **7**, 248–249 (2010)
66. Chun, S., Fay, J.C.: Identification of deleterious mutations within three human genomes. Genome Research **19**(9), 1553–1561 (2009)
67. Wong, W.C., Kim, D., Carter, H., Diekhans, M., Ryan, M.C., Karchin, R.: CHASM and SNVBox: toolkit for detecting biologically important single nucleotide mutations in cancer. Bioinformatics **27**(15), 2147–2148 (2011)
68. Xiong, H.Y., Alipanahi, B., Lee, L.J., Bretschneider, H., Merico, D., Yuen, R.K.C., Hua, Y., Gueroussov, S., Najafabadi, H.S., Hughes, T.R., Morris, Q., Barash, Y., Krainer, A.R., Jojic, N., Scherer, S.W., Blencowe, B.J., Frey, B.J.: The human splicing code reveals new insights into the genetic determinants of disease. Science **347**, 1254806 (2015)
69. Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M., Yen, A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., Ziller, M.J., Amin, V., Whitaker, J.W., Schultz, M.D., Ward, L.D., Sarkar, A., Quon, G., Sandstrom, R.S., Eaton, M.L., Wu, Y.-C., Pfenning, A.R., Wang, X., Claussnitzer, M., Liu, Y., Coarfa, C., Harris, R.A., Shoresh, N., Epstein, C.B., Gjoneska, E., Leung, D., Xie, W., Hawkins, R.D., Lister, R., Hong, C., Gascard, P., Mungall, A.J., Moore, R., Chuah, E., Tam, A., Canfield, T.K.,

Hansen, R.S., Kaul, R., Sabo, P.J., Bansal, M.S., Carles, A., Dixon, J.R., Farh, K.-H., Feizi, S., Karlic, R., Kim, A.-R., Kulkarni, A., Li, D., Lowdon, R., Elliott, G., Mercer, T.R., Neph, S.J., Onuchic, V., Polak, P., Rajagopal, N., Ray, P., Sallari, R.C., Siebenthall, K.T., Sinnott-Armstrong, N.A., Stevens, M., Thurman, R.E., Wu, J., Zhang, B., Zhou, X., Beaudet, A.E., Boyer, L.A., De Jager, P.L., Farnham, P.J., Fisher, S.J., Haussler, D., Jones, S.J.M., Li, W., Marra, M.A., McManus, M.T., Sunyaev, S., Thomson, J.A., Tlsty, T.D., Tsai, L.-H., Wang, W., Waterland, R.A., Zhang, M.Q., Chadwick, L.H., Bernstein, B.E., Costello, J.F., Ecker, J.R., Hirst, M., Meissner, A., Milosavljevic, A., Ren, B., Stamatoyannopoulos, J.A., Wang, T., Kellis, M., Consortium, R.E.: Integrative analysis of 111 reference human epigenomes. Nature **518**(7539), 317–330 (2015)

70. Arbiza, L., Gronau, I., Aksoy, B.A., Hubisz, M.J., Gulko, B., Keinan, A., Siepel, A.: Genome-wide inference of natural selection on human transcription factor binding sites. Nature Genetics **45**(7), 723–729 (2013)
71. Gronau, I., Arbiza, L., Mohammed, J., Siepel, A.: Inference of natural selection from interspersed genomic elements based on polymorphism and divergence. Molecular Biology and Evolution **30**(5), 1159–1171 (2013)
72. Harrow, J., Frankish, A., Gonzalez, J.M., Tapanari, E., Diekhans, M., Kokocinski, F., Aken, B.L., Barrell, D., Zadissa, A., Searle, S., Barnes, I., Bignell, A., Boychenko, V., Hunt, T., Kay, M., Mukherjee, G., Rajan, J., Despacio-Reyes, G., Saunders, G., Steward, C., Harte, R., Lin, M., Howald, C., Tanzer, A., Derrien, T., Chrast, J., Walters, N., Balasubramanian, S., Pei, B., Tress, M., Rodriguez, J.M., Ezkurdia, I., van Baren, J., Brent, M., Haussler, D., Kellis, M., Valencia, A., Reymond, A., Gerstein, M., Guigo, R., Hubbard, T.J.: GENCODE: The reference human genome annotation for the encode project. Genome Research **22**(9), 1760–1774 (2012)
73. Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson, H., Spieth, J., Hillier, L.W., Richards, S., Weinstock, G.M., Wilson, R.K., Gibbs, R.A., Kent, W.J., Miller, W., Haussler, D.: Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Research **15**(8), 1034–1050 (2005)
74. Team, R.D.C.: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria (2008). R Foundation for Statistical Computing. ISBN 3-900051-07-0. http://www.R-project.org
75. Nair, V., Hinton, G.E.: Rectified linear units improve restricted boltzmann machines. In: Proceedings of the 27th International Conference on International Conference on Machine Learning. ICML'10, pp. 807–814. Omnipress, USA (2010)
76. Glorot, X., Bengio, Y.: Understanding the difficulty of training deep feedforward neural networks. In: Teh, Y.W., Titterington, M. (eds.) Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics. Proceedings of Machine Learning Research, vol. 9, pp. 249–256. PMLR, Chia Laguna Resort, Sardinia, Italy (2010)
77. Kingma, D.P., Ba, J.: Adam: A method for stochastic optimization. arXiv:1412.6980 (2014)
78. Liu, X., Jian, X., Eric, B.: dbNSFP v2.0: A database of human non-synonymous snvs and their functional predictions and annotations. Human Mutation **34**(9), 2393–2402 (2013)
79. Sing, T., Sander, O., Beerenwinkel, N., Lengauer, T.: ROCR: visualizing classifier performance in R. Bioinformatics **21**(20), 3940–3941 (2005)
80. Iossifov, I., O'Roak, B.J., Sanders, S.J., Ronemus, M., Krumm, N., Levy, D., Stessman, H.A., Witherspoon, K.T., Vives, L., Patterson, K.E., Smith, J.D., Paeper, B., Nickerson, D.A., Dea, J., Dong, S., Gonzalez, L.E., Mandell, J.D., Mane, S.M., Murtha, M.T., Sullivan, C.A., Walker, M.F., Waqar, Z., Wei, L., Willsey, A.J., Yamrom, B., Lee, Y.-h., Grabowska, E., Dalkic, E., Wang, Z., Marks, S., Andrews, P., Leotta, A., Kendall, J., Hakker, I., Rosenbaum, J., Ma, B., Rodgers, L., Troge, J., Narzisi, G., Yoon, S., Schatz, M.C., Ye, K., McCombie, W.R., Shendure, J., Eichler, E.E., State, M.W., Wigler, M.: The contribution of de novo coding mutations to autism spectrum disorder. Nature **515**, 216 (2014)
81. Krumm, N., Turner, T.N., Baker, C., Vives, L., Mohajeri, K., Witherspoon, K., Raja, A., Coe, B.P., Stessman, H.A., He, Z.-X., Leal, S.M., Bernier, R., Eichler, E.E.: Excess of rare, inherited truncating mutations in autism. Nature Genetics **47**, 582 (2015)
82. Turner, T., Hormozdiari, F., Duyzend, M., McClymont, S., Hook, P., Iossifov, I., Raja, A., Baker, C., Hoekzema, K., Stessman, H., Zody, M., Nelson, B., Huddleston, J., Sandstrom, R., Smith, J., Hanna, D., Swanson, J., Faustman, E., Bamshad, M., Stamatoyannopoulos, J., Nickerson, D., McCallion, A., Darnell, R., Eichler, E.: Genome sequencing of autism-affected families reveals disruption of putative noncoding regulatory dna. The American Journal of Human Genetics **98**(1), 58–74 (2016)
83. Yuen, R.K.C., Merico, D., Bookman, M., L Howe, J., Thiruvahindrapuram, B., Patel, R.V., Whitney, J., Deflaux, N., Bingham, J., Wang, Z., Pellecchia, G., Buchanan, J.A., Walker, S., Marshall, C.R., Uddin, M., Zarrei, M., Deneault, E., D'Abate, L., Chan, A.J.S., Koyanagi, S., Paton, T., Pereira, S.L., Hoang, N., Engchuan, W., Higginbotham, E.J., Ho, K., Lamoureux, S., Li, W., MacDonald, J.R., Nalpathamkalam, T., Sung, W.W.L., Tsoi, F.J., Wei, J., Xu, L., Tasse, A.-M., Kirby, E., Van Etten, W., Twigger, S., Roberts, W., Drmic, I., Jilderda, S., Modi, B.M., Kellam, B., Szego, M., Cytrynbaum, C., Weksberg, R., Zwaigenbaum, L., Woodbury-Smith, M., Brian, J., Senman, L., Iaboni, A., Doyle-Thomas, K., Thompson, A., Chrysler, C., Leef, J., Savion-Lemieux, T., Smith, I.M., Liu, X., Nicolson, R., Seifer, V., Fedele, A., Cook, E.H., Dager, S., Estes, A., Gallagher, L., Malow, B.A., Parr, J.R., Spence, S.J., Vorstman, J., Frey, B.J., Robinson, J.T., Strug, L.J., Fernandez, B.A., Elsabbagh, M., Carter, M.T., Hallmayer, J., Knoppers, B.M., Anagnostou, E., Szatmari, P., Ring, R.H., Glazer, D., Pletcher, M.T., Scherer, S.W.: Whole genome sequencing resource identifies 18 new candidate genes for autism spectrum disorder. Nature Neuroscience **20**, 602 (2017)
84. Werling, D.M., Brand, H., An, J.-Y., Stone, M.R., Zhu, L., Glessner, J.T., Collins, R.L., Dong, S., Layer, R.M., Markenscoff-Papadimitriou, E., Farrell, A., Schwartz, G.B., Wang, H.Z., Currall, B.B., Zhao, X., Dea, J., Duhn, C., Erdman, C.A., Gilson, M.C., Yadav, R., Handsaker, R.E., Kashin, S., Klei, L., Mandell, J.D., Nowakowski, T.J., Liu, Y., Pochareddy, S., Smith, L., Walker, M.F., Waterman, M.J., He, X., Kriegstein, A.R., Rubenstein, J.L., Sestan, N., McCarroll, S.A., Neale, B.M., Coon, H., Willsey, A.J., Buxbaum, J.D., Daly, M.J., State, M.W., Quinlan, A.R., Marth, G.T., Roeder, K., Devlin, B., Talkowski, M.E., Sanders, S.J.: An analytical framework for whole-genome sequence association studies and its implications for autism spectrum disorder. Nature Genetics **50**(5), 727–736 (2018)

85. Rauch, A., Wieczorek, D., Graf, E., Wieland, T., Endele, S., Schwarzmayr, T., Albrecht, B., Bartholdi, D., Beygo, J., Di Donato, N., Dufke, A., Cremer, K., Hempel, M., Horn, D., Hoyer, J., Joset, P., Röpke, A., Moog, U., Riess, A., Thiel, C.T., Tzschach, A., Wiesener, A., Wohlleber, E., Zweier, C., Ekici, A.B., Zink, A.M., Rump, A., Meisinger, C., Grallert, H., Sticht, H., Schenck, A., Engels, H., Rappold, G., Schröck, E., Wieacker, P., Riess, O., Meitinger, T., Reis, A., Strom, T.M.: Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: an exome sequencing study. The Lancet **380**(9854), 1674–1682 (2012)

86. Gulsuner, S., Walsh, T., Watts, A., Lee, M., Thornton, A., Casadei, S., Rippey, C., Shahin, H., Braff, D., Cadenhead, K., Calkins, M., Dobie, D., Freedman, R., Green, M., Greenwood, T., Gur, R., Gur, R., Lazzeroni, L., Light, G., Nuechterlein, K., Olincy, A., Radant, A., Ray, A., Schork, N., Seidman, L., Siever, L., Silverman, J., Stone, W., Sugar, C., Swerdlow, N., Tsuang, D., Tsuang, M., Turetsky, B., Aduroja, T., Allen, T., Bradford, L.D., Calkins, M.E., Devlin, B., Edwards, N.B., Ganguli, R., Go, R.C.P., Gur, R.E., Gur, R.C., Kwentus, J., Lahti, A.C., Lyons, P., Mathos, K., May, R., McLeod-Bryant, S., McEvoy, J.P., Montgomery-Barefield, L., Nimgaonkar, V.L., O'Jile, J., Santos, A., Savage, R.M., Swanson, C.L., Wilson, W., Nimgaonkar, V.L., Go, R.C.P., Savage, R.M., Swerdlow, N.R., Gur, R.E., Braff, D.L., King, M.-C., McClellan, J.M.: Spatial and temporal mapping of de novo mutations in schizophrenia to a fetal prefrontal cortical network. Cell **154**(3), 518–529 (2013)

87. The 1000 Genomes Project, Conrad, D.F., Keebler, J.E.M., DePristo, M.A., Lindsay, S.J., Zhang, Y., Casals, F., Idaghdour, Y., Hartl, C.L., Torroja, C., Garimella, K.V., Zilversmit, M., Cartwright, R., Rouleau, G.A., Daly, M., Stone, E.A., Hurles, M.E., Awadalla, P.: Variation in genome-wide mutation rates within and between human families. Nature Genetics **43**, 712 (2011)

88. Ramu, A., Noordam, M.J., Schwartz, R.S., Wuster, A., Hurles, M.E., Cartwright, R.A., Conrad, D.F.: Denovogear: de novo indel and point mutation discovery and phasing. Nature Methods **10**, 985 (2013)

89. The Genome of the Netherlands Consortium: Whole-genome sequence variation, population structure and demographic history of the dutch population. Nature Genetics **46**, 818–825 (2014)

90. Besenbacher, S., Liu, S., Izarzugaza, J.M.G., Grove, J., Belling, K., Bork-Jensen, J., Huang, S., Als, T.D., Li, S., Yadav, R., Rubio-García, A., Lescai, F., Demontis, D., Rao, J., Ye, W., Mailund, T., Friborg, R.M., Pedersen, C.N.S., Xu, R., Sun, J., Liu, H., Wang, O., Cheng, X., Flores, D., Rydza, E., Rapacki, K., Damm Sørensen, J., Chmura, P., Westergaard, D., Dworzynski, P., Sørensen, T.I.A., Lund, O., Hansen, T., Xu, X., Li, N., Bolund, L., Pedersen, O., Eiberg, H., Krogh, A., Børglum, A.D., Brunak, S., Kristiansen, K., Schierup, M.H., Wang, J., Gupta, R., Villesen, P., Rasmussen, S.: Novel variation and de novo mutation rates in population-wide de novo assembled danish trios. Nature Communications **6**, 5969 (2015)

91. Ho, D.E., Imai, K., King, G., Stuart, E.A.: MatchIt: Nonparametric preprocessing for parametric causal inference. Journal of Statistical Software **42**(8), 1–28 (2011)

92. Mi, H., Huang, X., Muruganujan, A., Tang, H., Mills, C., Kang, D., Thomas, P.D.: PANTHER version 11: expanded annotation data from gene ontology and reactome pathways, and data analysis tool enhancements. Nucleic Acids Research **45**(D1), 183–189 (2017)

Figures

Figure 1 Overview of the UNEECON model. UNEECON estimates negative selection on missense mutation i in gene j based on the relative probability of the occurrence of the missense mutation, η_{ij}, compared to the occurrence probability of neutral mutations, μ_{ij}. η_{ij} depends on the sum of a variant-level fixed effect, z_{ij}, and a gene-level random effect, u_j. We assume that z_{ij} captures the contribution of variant-level features, $\mathbf{X}_{ij}$, to negative selection, and model the relationship between $\mathbf{X}_{ij}$ and z_{ij} with a feedforward neural network. We assume that u_j is a Gaussian random variant modeling the gene-level variation of selective constraints that cannot be predicted from variant features. The sum of z_{ij} and u_j is then sent to a logistic function to obtain η_{ij}. The neutral occurrence probability, μ_{ij}, is from a context-dependent mutation model trained on putatively neutral mutations. Free parameters of the UNEECON model are estimated by minimizing the discrepancy between the predicted occurrence probability, $\eta_{ij} \cdot \mu_{ij}$, and the observed occurrence of each potential missense mutation in the gnomAD exome sequencing data [29].

Figure 2 Distributions of UNEECON scores across potential missense mutations. (a) Distributions of UNEECON scores estimated for potential missense mutations in haploinsufficient (HI) genes [37], autosomal dominant disease genes [35, 36], autosomal recessive disease genes [35, 36], and olfactory receptor genes [45]. **(b)** Distributions of UNEECON scores estimated for potential missense mutations in various protein regions. The functional sites and protein secondary structures are based on UniProt annotations [48]. The predicted disordered protein regions are from MobiDB [49]. **(c)** Average UNEECON scores estimated for all codon positions in the CDKL5 protein. Each grey dot represents the UNEECON score averaged over all missense mutations in a codon position. Blue curve represents the locally estimated scatterplot smoothing (LOESS) fit. Blue and red dots represent pathogenic and benign missense variants from ClinVar [30], respectively. The horizontal line represents a constrained region reported in a previous study [25].

Figure 3 Predictive power of various methods for distinguishing pathogenic missense variants from benign missense variants. (a) Performance in predicting autosomal pathogenic variants from ClinVar [30]. True positive and true negative rates correspond to the fractions of pathogenic and benign variants exceeding various thresholds, respectively. AUC corresponds to the area under the receiver operating characteristic curve. **(b)** Enrichment of predicted deleterious *de novo* variants in individuals affected by developmental disorders [31]. The y-axis corresponds to the $\log_2$ odds ratio of the enrichment of predicted deleterious variants in the affected individuals for a given percentile threshold. The x-axis corresponds to the various percentile threshold values used in the enrichment analysis. Error bars represent the standard error of the $\log_2$ odds ratio.

Figure 4 Predictive power of various methods for distinguishing disease and essential genes from genes not likely to have strong phenotypic effects. (a) Performance in predicting autosomal dominant disease genes [35, 36]. **(b)** Performance in predicting haploinsufficient genes [37]. **(c)** Performance in predicting human orthologs of mouse essential genes [33, 34]. **(d)** Performance in predicting human essential genes in cell lines [32]. True positive and true negative rates correspond to the fractions of positive and negative genes exceeding various thresholds, respectively. AUC corresponds to the area under the receiver operating characteristic curve.

Figure 5 Distributions of gene-level intolerance to missense and to loss-of-function mutations.
(a) Correlation between gene-level intolerance to missense mutations (UNEECON-G score) and that to loss-of-function (LOF) mutations (pLI score). Blue dots represent 956 genes intolerant to both missense and LOF mutations. Red dots represent 956 genes tolerant to missense but not to loss-of-function mutations. **(b)** Distribution of protein disorder content in the gene sets intolerant to loss-of-function mutations. **(c)** Enrichment of Reactome pathways in the gene set intolerant to both missense and loss-of-function mutations. The gene set tolerant to missense but not to loss-of-function mutations is used as a background. Only the highest-level Reactome terms from the PANTHER hierarchy view are included in the visualization. **(d)** Enrichment of autism genes in the gene sets intolerant to loss-of-function mutations. Error bars represent the standard error of the $\log_2$ odds ratio.

Additional Files
Additional file 1 — Supplementary Material
Supplementary figures and tables.

Additional file 2 — Supplementary Dataset
List of 1,912 genes intolerant to loss-of-function mutations.