Unified inference of missense variant effects and gene constraints in the human genome

Yi-Fei Huang[1,2*],

1 Department of Biology, Pennsylvania State University, University Park, PA, USA
2 Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA, USA

* yuh371@psu.edu

Abstract

A challenge in medical genomics is to identify variants and genes associated with severe genetic disorders. Based on the premise that severe, early-onset disorders often result in a reduction of evolutionary fitness, several statistical methods have been developed to predict pathogenic variants or constrained genes based on the signatures of negative selection in human populations. However, we currently lack a statistical framework to jointly predict deleterious variants and constrained genes from both variant-level features and gene-level selective constraints. Here we present such a unified approach, UNEECON, based on deep learning and population genetics. UNEECON treats the contributions of variant-level features and gene-level constraints as a variant-level fixed effect and a gene-level random effect, respectively. The sum of the fixed and random effects is then combined with an evolutionary model to infer the strength of negative selection at both variant and gene levels. Compared with previously published methods, UNEECON shows unmatched performance in predicting missense variants and protein-coding genes associated with autosomal dominant disorders, and feature importance analysis suggests that both gene-level selective constraints and variant-level predictors are important for accurate variant prioritization. Furthermore, based on UNEECON, we observe an unexpected low correlation between gene-level intolerance to missense mutations and that to loss-of-function mutations, which can be partially explained by the prevalence of disordered protein regions that are highly tolerant to missense mutations. Finally, we show that genes intolerant to both missense and loss-of-function mutations play key roles in the central nervous system and the autism spectrum disorders. Overall, UNEECON is a promising framework for both variant and gene prioritization.

Author summary

Numerous statistical methods have been developed to predict deleterious missense variants or constrained genes in the human genome, but unified prioritization methods that utilize both variant- and gene-level information are underdeveloped. Here we present UNEECON, an evolution-based deep learning framework for unified variant and gene prioritization. By integrating variant-level predictors and gene-level selective constraints, UNEECON outperforms existing methods in predicting missense variants and protein-coding genes associated with dominant disorders. Based on UNEECON, we show that disordered proteins are tolerant to missense mutations but not to

loss-of-function mutations. In addition, we find that genes under strong selective constraints at both missense and loss-of-function levels are strongly associated with the central nervous system and the autism spectrum disorders, highlighting the need to investigate the function of these highly constrained genes in future studies.

Introduction 1

A fundamental question in biology is to understand how genomic variation contributes to phenotypic variation and disease risk. While millions of protein-altering variants have been identified in the human genome, it is challenging to assess the functional and clinical significance of these variants. In particular, a large fraction of missense variants have been annotated as "variants of uncertain significance" (VUS) [1, 2], forming a major hurdle for both basic research and medical practice. This problem is further exacerbated by the difficulty of experimentally validating the function of large numbers of missense variants *in vivo*. Therefore, there is a tremendous need for accurate computational tools to prioritize deleterious missense variants [3].

Since early-onset, severe genetic disorders are often associated with a reduction of evolutionary fitness, signatures of negative (purifying) selection, such as sequence conservation, have been widely used to predict deleterious variants associated with Mendelian disorders [4–12]. Among existing evolutionary approaches for variant prioritization, recently developed integrative methods are particularly powerful [8–15]. By learning a linear or nonlinear mathematical function from predictive variant features, such as sequence conservation scores and protein structural features, to the strength of negative selection, these statistical methods estimate negative selection on observed and potential mutations in the human genome. The estimated strength of negative selection can then be utilized to prioritize deleterious variants associated with severe genetic disorders. Because these evolutionary approaches are trained on tremendous natural polymorphisms observed in healthy individuals instead of sparsely annotated pathogenic variants, they have shown good performance in predicting pathogenic variants, frequently outperforming or on par with supervised machine learning models trained on disease data [8, 10–12].

Despite the success of evolution-based metrics of variant effects, the existing methods nevertheless suffer from a few critical limitations. First, most existing methods focus on learning a shared mathematical function from predictive variant features to negative selection and assume that this function is equally applicable to all protein-coding genes. Instead, a subset of genes can depart from the genome-wide trend between variant features and negative selection, possibly due to enhanced or relaxed purifying selection on these genes in the human lineage [12]. Second, most existing methods are trained on common genetic variants, making it challenging to distinguish strong negative selection associated with severe genetic diseases from moderate negative selection without clinical implications. Third, these methods typically are agnostic to the mode of inheritance and, therefore, may be suboptimal in the prediction of deleterious variants associated with dominant disorders on which we focus in this work.

In parallel with the development of variant-level interpretation methods, several complementary, gene-centric methods have been proposed to predict protein-coding genes associated with dominant genetic disorders [16–23]. Unlike variant-level predictors, the gene-level prioritization methods seek to identify constrained genes that are intolerant to heterozygous nonsynonymous mutations. These gene-level constraint metrics have been shown to provide complementary information on variant effects and have been successfully used to prioritize variants associated with dominant genetic disorders [16, 17, 24]. However, these methods typically assume that all the missense mutations in a gene or a genic region have identical effects and, therefore, may not be

able to distinguish pathogenic missense variants from proximal benign missense variants.

Since variant-level and gene-level prioritization methods leverage complementary signatures of negative selection, unifying the two lines of research for joint inference of variant effects and gene constraints should be beneficial. Recently, a couple of studies have tried to address this question [25–27], but all of these methods are supervised machine learning models trained on disease variants. Therefore, we lack an evolution-based statistical framework to combine predictive variant features and gene-level selective constraints for variant and gene prioritization. Based on a novel deep learning framework, *i.e.*, deep mixed-effects model, we develop **UNEECON** (**UN**ified inferenc**E** of variant **E**ffects and gene **CON**straints), an evolution-based framework to predict deleterious variants and constrained genes from both variant features and gene-level intolerance to missense mutations. By integrating 30 predictive variant features and genomic variation from 141,456 human genomes [28, 29], UNEECON outperforms existing variant effect predictors and gene constraint scores in predicting pathogenic missense variants with a dominant mode of inheritance. In addition, deleterious *de novo* variants predicted by UNEECON are strongly enriched in individuals affected by severe development disorders [30, 31], highlighting its power for interpreting the effects of *de novo* mutations. Furthermore, UNEECON provides estimates of gene-level selective constraints (UNEECON-G scores) for all protein-coding genes. In the setting of gene prioritization, UNEECON-G scores show unmatched performance in predicting human essential genes [32], mouse essential genes [33, 34], autosomal dominant disease genes [35, 36], and haploinsufficient genes [37]. UNEECON is a powerful framework for both variant and gene prioritization.

Results

UNEECON integrates variant- and gene-level information to predict detrimental variants and constrained genes

The key idea of UNEECON is to combine variant-level predictive features, such as sequence conservation scores and protein structural features, and gene-level signatures of selective constraints to infer negative selection on every potential missense mutation in the human genome. Inspired by classical sequence conservation models [38–42], which use site-specific substitution rate as a proxy of negative selection, we utilize the relative probability of the occurrence of a potential missense mutation in human populations, compared to neutral mutations, as an allele-specific predictor of negative selection.

We first fit a mutation model to calculate μ_{ij}, *i.e.*, the probability of the occurrence of mutation i in gene j when selection is absent. Our mutation model is trained on putatively neutral mutations in the gnomAD data [29] and captures the impact of multiple neutral and technical factors, including the 7-mer sequence context centered on the focal site, the identity of the alternative allele, the local mutation rate, and the average sequencing coverage, on the probability of occurrence of a neutral mutation. At the gene level, the number of synonymous mutations predicted by the mutation model was nearly perfectly correlated with the observed number of synonymous mutations (Spearman's $\rho = 0.976$; Fig. S1), suggesting that the mutation model can serve as a proper baseline for inferring negative selection.

Given the mutation model, UNEECON employs a novel deep learning framework, deep mixed-effects model, to infer the relative probability of the occurrence of each potential missense mutation (Fig. 1). In more detail, we denote η_{ij} as the relative probability of the occurrence of missense mutation i in gene j with respect to the neutral occurrence probability, μ_{ij}. η_{ij} captures the impact of natural selection, instead of mutation and genetic drift, on the occurrence of a missense mutation. Analogous to

generalized linear mixed models, we assume that η_{ij} depends on the sum of a 96
variant-level fixed effect, z_{ij}, and a gene-level random effect, u_j. We assume that z_{ij} 97
captures the contribution of predictive variant features to negative selection. Denoting 98
$\mathbf{X}_{ij}$ as the vector of predictive features associated with mutation i in gene j, we use a 99
feedforward neural network to model the relationship between $\mathbf{X}_{ij}$ and z_{ij} (Fig. 1). 100
Furthermore, we assume that the random-effect term, u_j, is a Gaussian (normal) 101
random variable which models the variation of gene-level constraints that is not 102
predictable from feature vector $\mathbf{X}_{ij}$. We then perform a logistic transformation on the 103
sum of z_{ij} and u_j to obtain η_{ij}. Finally, we multiply η_{ij} and μ_{ij} to obtain the 104
occurrence probability of missense mutation i in the gnomAD exome sequencing data. 105

Fig 1. Overview of the UNEECON model. UNEECON estimates negative
selection on missense mutation i in gene j based on the relative probability of the
occurrence of the missense mutation, η_{ij}, compared to the occurrence probability of
neutral mutations, μ_{ij}. η_{ij} depends on the sum of a variant-level fixed effect, z_{ij}, and a
gene-level random effect, u_j. We assume that z_{ij} captures the contribution of
variant-level features, $\mathbf{X}_{ij}$, to negative selection, and model the relationship between
$\mathbf{X}_{ij}$ and z_{ij} with a feedforward neural network. We assume that u_j is a Gaussian
random variant modeling the gene-level variation of selective constraints that cannot be
predicted from variant features. The sum of z_{ij} and u_j is then sent to a logistic function
to obtain η_{ij}. The neutral occurrence probability, μ_{ij}, is from a context-dependent
mutation model trained on putatively neutral mutations. Free parameters of the
UNEECON model are estimated by minimizing the discrepancy between the predicted
occurrence probability, $\eta_{ij} \cdot \mu_{ij}$, and the observed occurrence of each potential missense
mutation in the gnomAD exome sequencing data [29].

We estimate the parameters of the UNEECON model by minimizing the discrepancy 106
between the predicted occurrence probability, $\eta_{ij} \cdot \mu_{ij}$, and the observed 107
presence/absence of each potential missense mutation, which is equivalent to 108
maximizing a Bernoulli likelihood function. We also employ regularization techniques, 109
including dropout [43] and early stopping [44], to avoid overfitting. After training, we 110
calculate variant-effect scores (UNEECON scores) and gene-level intolerance to missense 111
mutations (UNEECON-G scores) defined as the expected reduction of the occurrence 112
probability at variant and gene levels, respectively. Both UNEECON and UNEECON-G 113
scores range from 0 to 1, with higher scores suggesting stronger negative selection. 114

UNEECON scores capture variation of negative selection within and across genes

We trained the UNEECON model with 30 missense variant features, including conservation scores, protein structural features, and regulatory features (Table S1), and rare missense variants with a minor allele frequency (MAF) lower than 0.1% in the gnomAD dataset [29]. After the training process, we first compared the distributions of UNEECON scores across potential missense mutations in haploinsufficient genes [37], autosomal dominant disease genes [35, 36], autosomal recessive disease genes [35, 36], and olfactory receptor genes [45]. In agreement with previous studies [16, 28, 35], missense mutations in autosomal dominant disease genes had higher UNEECON scores than those in autosomal recessive disease genes (Fig. 2a), suggesting a stronger selection on heterozygous missense mutations in the genes associated with autosomal dominant disorders. Also, we observed that missense mutations in olfactory receptor genes had much lower UNEECON scores (Fig. 2a), potentially due to relaxed purifying selection or enhanced positive selection in the human lineage [46].

Second, we investigated whether the distributions of UNEECON scores varied across different types of protein regions annotated by UniProt [47, 49] and MobiDB [48]. We observed that the distributions of UNEECON scores were similar among α-helices, β-strands, and hydrogen-bonded turns (Fig. 2b). In contrast, UNEECON scores were significantly lower in disordered protein regions. As expected, UNEECON scores were significantly higher in functional protein sites, including enzyme active sites and ligand binding sites (Fig. 2b), suggesting much stronger purifying selection on these critical residues.

Furthermore, we used the human CDKL5 protein as an example to illustrate UNEECON's capability to capture regional variation of negative selection within a gene. Similar to a previously published metric of regional missense constraints [25], UNEECON predicted that the N-terminus of CDKL5 was under strong negative selection (Fig. 2c). In agreement with the evidence of strong selection, this region was also enriched with known pathogenic missense variants and depleted with benign missense variants (Fig. 2c). Interestingly, UNEECON also predicted that the C-terminus of CDKL5 was under much weaker selection compared with the rest of the protein (Fig. 2c). Accordingly, this unconstrained region was enriched with benign missense variants and depleted with pathogenic missense variants.

UNEECON scores accurately predict missense mutations associated with dominant genetic disorders

Since UNEECON measures the strength of strong negative selection on heterozygous missense mutations, we hypothesize that UNEECON scores are predictive of missense variants associated with dominant Mendelian disorders. To test this hypothesis, we compared the performance of UNEECON with eight existing methods in the setting of predicting autosomal dominant disease variants in ClinVar [30]. The eight previously published methods can be classified into three categories: 1) variant-level prediction methods, including LASSIE [12], CADD [8], Eigen [50], and PrimateAI [11]; 2) gene-level and region-level prediction methods, including RVIS [16], pLI [17], and CCR [22]; and 3) MPC [25], a supervised machine learning method utilized both variant features and regional constraint scores as input features. It is worth noting that several aforementioned methods, including UNEECON, used the variants from the gnomAD dataset as a part of their training data. If the same gnomAD variants are also used as negative controls in the evaluation of performance, we will overestimate the power of these methods trained on the gnomAD data. To avoid this problem, we used benign missense variants in ClinVar as negative controls and removed any ClinVar variants that

Fig 2. Distributions of UNEECON scores across potential missense mutations. **(a)** Distributions of UNEECON scores estimated for potential missense mutations in haploinsufficient (HI) genes [37], autosomal dominant disease genes [35, 36], autosomal recessive disease genes [35, 36], and olfactory receptor genes [45]. **(b)** Distributions of UNEECON scores estimated for potential missense mutations in various protein regions. The functional sites and protein secondary structures are based on UniProt annotations [47]. The predicted disordered protein regions are from MobiDB [48]. **(c)** Average UNEECON scores estimated for all codon positions in the CDKL5 protein. Each grey dot represents the UNEECON score averaged over all missense mutations in a codon position. Blue curve represents the locally estimated scatterplot smoothing (LOESS) fit. Blue and red dots represent pathogenic and benign missense variants from ClinVar [30], respectively. The horizontal line represents a constrained region reported in a previous study [25].

are also present in gnomAD. Then, we matched the numbers of positives and negatives by random sampling without replacement.

Overall, UNEECON outperformed previous methods in predicting ClinVar missense variants associated with autosomal dominant disorders (Fig. 3a; Table S2). Among the previously published methods, variant-level predictors performed better than gene-level and region-level constraint scores. To test the robustness of these results, we constructed an alternative set of dominant pathogenic variants defined as all the pathogenic missense variants located in 709 genes associated with autosomal dominant diseases [35, 36]. UNEECON again outperformed the other methods in this dataset, even though the difference of performance between UNEECON and LASSIE was not

Fig 3. Predictive power of various methods for distinguishing pathogenic missense variants from benign missense variants. (a) Performance in predicting autosomal pathogenic variants from ClinVar [30]. True positive and true negative rates correspond to the fractions of pathogenic and benign variants exceeding various thresholds, respectively. AUC corresponds to the area under the receiver operating characteristic curve. (b) Enrichment of predicted deleterious *de novo* variants in individuals affected by developmental disorders [31]. The y-axis corresponds to the $\log_2$ odds ratio of the enrichment of predicted deleterious variants in the affected individuals for a given percentile threshold. The x-axis corresponds to the various percentile threshold values used in the enrichment analysis. Error bars represent the standard error of the $\log_2$ odds ratio.

While UNEECON was highly powerful in predicting autosomal dominant disease   176
variants, it might not be as accurate when applied to predict recessive disease variants   177
which are only detrimental in the homozygous state. To test this hypothesis, we   178
evaluated UNEECON's performance in predicting autosomal recessive disease variants   179
in ClinVar [30]. As expected, UNEECON was outperformed by multiple methods, such   180
as LASSIE [12] and Eigen [50], in this setting (Fig. S3). Similar results were reached   181
with an alternative set of recessive pathogenic variants defined as all the pathogenic   182
missense variants located in 1,183 autosomal recessive genes [35, 36] (Fig. S4).   183

As an orthogonal benchmark of UNEECON's performance in predicting dominant   184
disease variants, we investigated whether UNEECON was able to predict *de novo*   185
missense mutations identified in individuals affected by severe developmental   186
disorders [31]. We obtained *de novo* missense variants identified in affected individuals   187
and healthy individuals from denovo-db [51]. Then, for multiple percentile rank cutoff   188
values (top 10%, 20%, 30%, and 40%), we evaluated the enrichment of deleterious   189
variants predicted by each method in the affected individuals. Overall, missense variants   190
predicted by UNEECON and MPC showed the highest enrichments in the affected   191

individuals (Fig. 3b), suggesting that these two methods were more powerful in ₁₉₂
predicting de novo risk mutations. The performance gaps between UNEECON/MPC ₁₉₃
and the other methods were highest at the most stringent cutoff of 10% (Fig. 3b). ₁₉₄

UNEECON-G scores perform favorably in predicting disease genes and essential genes ₁₉₅ ₁₉₆

The UNEECON model also provided UNEECON-G scores which measure gene-level ₁₉₇
intolerance to missense mutations. We compared the performance of UNEECON-G ₁₉₈
scores with alternative gene constraint scores, including RVIS [16], pLI [17], mis-z [17], ₁₉₉
and GDI [19], in the setting of predicting disease genes and essential genes. The disease ₂₀₀
gene sets included 709 autosomal dominant disease genes [35,36] and 294 ₂₀₁
haploinsufficient genes [37]. The essential gene sets included 2,454 human orthologs of ₂₀₂
mouse essential genes [33,34] and 683 human essential genes identified by CRISPR ₂₀₃
knockout experiments in human cell lines [32]. For each set of autosomal dominant ₂₀₄
disease genes, haploinsufficient genes, and mouse essential genes, we constructed a ₂₀₅
negative gene set by sampling a matched number of genes from the other genes in the ₂₀₆
human genome. For the essential genes identified by CRISPR in cell lines, we ₂₀₇
constructed a negative gene set by sampling a matched number of genes from the ₂₀₈
nonessential genes reported in the same study [32]. Overall, UNEECON-G significantly ₂₀₉
outperformed alternative gene-level metrics in predicting the four sets of essential and ₂₁₀
disease genes (Fig. 4; Table S3). ₂₁₁

Both variant-level features and gene-level constraints are important for variant interpretation ₂₁₂ ₂₁₃

To gain more insights into which components of the UNEECON model are critical for ₂₁₄
predicting detrimental variants, we characterized the contributions of the 30 variant ₂₁₅
features and the gene-level random effect to negative selection. Unfortunately, it is ₂₁₆
challenging to directly interpret the UNEECON model due to the nonlinearity ₂₁₇
introduced by the feedforward artificial neural network. To bypass this difficulty, we ₂₁₈
used a linear version of the UNEECON model as an approximation of the nonlinear ₂₁₉
UNEECON model. The linear UNEECON model has no hidden units and can be ₂₂₀
interpreted as a Generalized linear mixed model. Then, we evaluated the importance of ₂₂₁
the variant-level features and the gene-level random effect in this linear surrogate model. ₂₂₂
The variant score predicted by the linear UNEECON model was nearly perfectly ₂₂₃
correlated with the UNEECON score predicted by the original nonlinear UNEECON ₂₂₄
model (Spearman's $\rho = 0.95$), supporting the use of the linear UNEECON model as a ₂₂₅
surrogate for model interpretation. ₂₂₆

In the linear surrogate model, the relative probability of the occurrence of mutation ₂₂₇
i in gene j, η_{ij}, is a linear combination of variant features, $\mathbf{X}_{ij}$, and the gene-level ₂₂₈
random effect, u_j, up to a logistic transformation. Analogous to the interpretation of ₂₂₉
canonical linear models, we defined the contribution score of a variant feature as the ₂₃₀
negative value of the weight (regression coefficient) associated with this feature, and ₂₃₁
similarly defined the contribution score of the gene-level random effect as the negative ₂₃₂
value of its standard deviation. Positive and negative contribution scores suggest that ₂₃₃
the corresponding variables are positively and negatively correlated with negative ₂₃₄
selection, respectively, and contribution scores near zero indicate that the corresponding ₂₃₅
variables are not important for predicting detrimental mutations. ₂₃₆

As shown in Fig. S5, the gene-level random effect was the most important feature for ₂₃₇
predicting deleterious variants, highlighting the importance of gene-level constraints for ₂₃₈
variant interpretation. In addition, several conservation scores, such as the qualitative ₂₃₉

Fig 4. Predictive power of various methods for distinguishing disease and essential genes from genes not likely to have strong phenotypic effects. (a) Performance in predicting autosomal dominant disease genes [35, 36]. **(b)** Performance in predicting haploinsufficient genes [37]. **(c)** Performance in predicting human orthologs of mouse essential genes [33, 34]. **(d)** Performance in predicting human essential genes in cell lines [32]. True positive and true negative rates correspond to the fractions of positive and negative genes exceeding various thresholds, respectively. AUC corresponds to the area under the receiver operating characteristic curve.

predictions from LRT, PROVEAN, and SIFT, and a subset of protein structural features, such as the predicted probabilities of forming various protein secondary structures, were also important for accurate prediction of deleterious variants. Even though these variant features individually were less important than the gene-level random effect, accumulatively they explained a significant faction of the variation of negative selection across missense mutations. Therefore, both variant features and gene-level constraints are important for predicting deleterious variants.

Missense constraints and loss-of-function constraints are weakly correlated across genes

The UNEECON-G score represents gene-level intolerance to missense mutations in human populations. Alternatively, selective constraint on a gene can be defined as its degree of intolerance to loss-of-function mutations, such as stop-gained, frameshift, and splice-site mutations [17, 28, 29]. We investigated the correlation between the UNEECON-G score and the pLI score, a metric of gene-level intolerance to loss-of-function mutations [17]. Surprisingly, the UNEECON-G score was only moderately correlated with the pLI score (Spearman's $\rho = 0.59$; Fig. 5a), suggesting a

discrepancy between gene-level intolerance to missense mutations and that to   256
loss-of-function mutations in a subset of protein-coding genes.   257

Fig 5. Distributions of gene-level intolerance to missense and to loss-of-function mutations. (a) Correlation between gene-level intolerance to missense mutations (UNEECON-G score) and that to loss-of-function (LOF) mutations (pLI score). Blue dots represent 956 genes intolerant to both missense and LOF mutations. Red dots represent 956 genes tolerant to missense but not to loss-of-function mutations. **(b)** Distribution of protein disorder content in the gene sets intolerant to loss-of-function mutations. **(c)** Enrichment of Reactome pathways in the gene set intolerant to both missense and loss-of-function mutations. The gene set tolerant to missense but not to loss-of-function mutations is used as a background. Only the highest-level Reactome terms from the PANTHER hierarchy view are included in the visualization. **(d)** Enrichment of autism genes in the gene sets intolerant to loss-of-function mutations. Error bars represent the standard error of the $\log_2$ odds ratio.

Intrinsically disordered proteins play key roles in a multitude of biological   258
processes [52], but their sequences are poorly conserved across species [53]. Therefore,   259
missense mutations in intrinsically disordered proteins may be under weak negative   260
selection even if loss-of-function mutations in these proteins are deleterious. To test this   261
hypothesis, we investigated the distributions of protein disorder content, *i.e.*, the   262
fraction of disordered regions in a protein [48], across 1,912 protein-coding genes   263
intolerant to heterozygous loss-of-function mutations (pLI score ≥ 0.9). We split the   264
1,912 genes into two equal-size groups based on their UNEECON-G scores, and defined   265
the 956 genes with higher UNEECON-G scores as the gene set intolerant to both   266

missense and loss-of-function mutations. Accordingly, we defined the 956 genes with ²⁶⁷
lower UNEECON-G scores as the gene set tolerant to missense but not to ²⁶⁸
loss-of-function mutations. Compared with the 956 genes intolerant to both missense ²⁶⁹
and loss-of-function mutations, the disorder contents were significantly higher in the 956 ²⁷⁰
genes tolerant to missense mutations but not to loss-of-function mutations (Fig. 5b). ²⁷¹
Therefore, disordered protein regions can at least partially explain the observed ²⁷²
discrepancy between missense constraints and loss-of-function constraints. ²⁷³

We further investigated the enrichment of Reactome pathways [54] and Gene ²⁷⁴
Ontology terms [55,56] in the 956 genes intolerant to both missense and loss-of-function ²⁷⁵
mutations, using the 956 genes tolerant to missense but not to loss-of-function ²⁷⁶
mutations as a background. Several key pathways associated with central nervous ²⁷⁷
system, gene regulation, cell cycle, and innate immune response were overrepresented in ²⁷⁸
the genes intolerant to both missense and loss-of-function mutations (Fig. 5c; Table S4). ²⁷⁹
We observed an enrichment of similar terms based on the Gene Ontology (Table S5 & ²⁸⁰
S6). In agreement with the enrichment of pathways associated with the central nervous ²⁸¹
system, the genes intolerant to both missense and loss-of-function mutations were ²⁸²
strongly enriched in the gene set implicated in the autism spectrum disorders [57] ²⁸³
(Fig. 5d). ²⁸⁴

Discussion ²⁸⁵

Here we present UNEECON for a unified prediction of deleterious missense mutations ²⁸⁶
and highly constrained genes in the human genome. Compared with previously ²⁸⁷
published methods, UNEECON shows superior performance in predicting missense ²⁸⁸
variants and protein-coding genes associated with dominant genetic disorders. ²⁸⁹
Therefore, UNEECON is a promising framework for both variant and gene ²⁹⁰
prioritization. Furthermore, unlike supervised machine learning approaches, such as ²⁹¹
MPC [25], UNEECON integrates variant features and gene constraints based on ²⁹²
signatures of negative selection instead of labeled disease variants. Therefore, ²⁹³
UNEECON is unlikely to suffer from the circularity and the inflated performance ²⁹⁴
commonly found in supervised methods [58]. ²⁹⁵

It is worth noting that UNEECON is different from existing prioritization methods ²⁹⁶
in multiple important aspects. First, unlike classical sequence conservation ²⁹⁷
metrics [38–42] and integrative variant scores [8–15], UNEECON infers the strength of ²⁹⁸
negative selection based on rare genetic variants. Therefore, UNEECON predicts ²⁹⁹
strongly deleterious variants with a dominant/semidominant mode of inheritance [59]. ³⁰⁰
Second, unlike previously published gene constraint metrics that typically assign the ³⁰¹
same score to all missense variants within a gene [16–23], UNEECON assigns different ³⁰²
scores to the variants in the same gene in a feature-dependent manner, providing ³⁰³
high-resolution maps of variant effects within genes. Third, UNEECON is able to adjust ³⁰⁴
the distribution of variant scores within a gene according to the degree of depletion of ³⁰⁵
missense variants in this gene, allowing for assigning different scores to missense ³⁰⁶
mutations with similar variant features but located in different genes. ³⁰⁷

Similar to other metrics of negative selection, the performance of UNEECON ³⁰⁸
strongly depends on the correlation between variant penetrance and negative selection. ³⁰⁹
Our analysis of pathogenic variants suggests that UNEECON can predict deleterious ³¹⁰
variants associated with dominant genetic disorders but not necessarily those associated ³¹¹
with recessive disorders. Also, because common variants are unlikely to be under strong ³¹²
negative selection [12], UNEECON scores might not be able to predict common variants ³¹³
associated with complex traits or late-onset diseases. In contrast, a recent study ³¹⁴
suggests that rare variants associated with complex traits are strongly enriched in ³¹⁵
coding regions and tend to be under negative selection [60], implying that UNEECON ³¹⁶

could be a useful tool for rare variant association studies. ₃₁₇

UNEECON is based on a novel machine learning framework, deep mixed-effects ₃₁₈
model, to integrate variant features and gene-level constraints. Our analysis of the linear ₃₁₉
UNEECON model suggests that both the predictive variant features and the gene-level ₃₂₀
constraints are important for variant interpretation. In particular, gene-level evidence of ₃₂₁
selective constraints is the single most important predictor of negative selection on ₃₂₂
missense mutations. Also, predictive variant features accumulatively explain a large ₃₂₃
fraction of the variation of negative selection across missense variants. We expect that ₃₂₄
the combination of variant-level predictors and gene-level constraints will be an essential ₃₂₅
component in the future development of variant and gene prioritization methods. ₃₂₆

By contrasting UNEECON-G scores against pLI scores, we observe an unexpected ₃₂₇
low correlation between gene-level intolerance to missense mutations and that to ₃₂₈
loss-of-function mutations. The prevalence of disordered protein regions in the human ₃₂₉
proteome is a key biological factor contributing to the low correlation between missense ₃₃₀
and loss-of-function constraints. Furthermore, we observe that the genes intolerant to ₃₃₁
both loss-of-function mutations and missense mutations may play key roles in the ₃₃₂
central nervous system and the autism spectrum disorders, highlighting the needs to ₃₃₃
investigate the function of these genes using state-of-the-art experimental techniques. ₃₃₄
By combing powerful variant and gene prioritization tools, such as UNEECON, and ₃₃₅
high-throughput mutagenesis and genome editing techniques [61,62], we will obtain ₃₃₆
more insights into the function of these strongly constrained genes in the future. ₃₃₇

Materials and methods ₃₃₈

Predictive variant features ₃₃₉

UNEECON was trained on 30 missense variant features previously used to infer fitness ₃₄₀
effects of coding variants in the human genome ([12]; Table S1). These features can be ₃₄₁
classified into three categories: sequence conservation scores, protein structural features, ₃₄₂
and functional genomic features. The sequence conservation scores included SIFT [5], ₃₄₃
PROVEAN [7], SLR [63], Grantham [64], PSIC [65], LRT [66], MutationAssessor [4], ₃₄₄
HMMEntropy [67], and phyloP scores [40]. The protein structural features included ₃₄₅
predicted secondary structures, B-factors, contributions to protein stability, and relative ₃₄₆
solvent accessibilities from SNVBox [67]. The functional genomic features included the ₃₄₇
non-commercial version of SPIDEX splicing scores [68] and the maximum RNA-seq ₃₄₈
signals from the Roadmap Epigenomics Project [69]. Following a common practice in ₃₄₉
machine learning and statistics, we standardized continuous features by subtracting the ₃₅₀
mean and dividing by the standard deviation. All the features were based on the hg19 ₃₅₁
(GRCh37) assembly. ₃₅₂

Population genomic data ₃₅₃

We dowloaded whole genome variation data, exome variation data, and corresponding ₃₅₄
sequencing coverage data from the gnomAD browser [28,29] (version 2.1.0). We only ₃₅₅
retained rare SNVs (MAF < 0.1%) that passed gnomAD's built-in quality filter for ₃₅₆
downstream analysis. ₃₅₇

Context-dependent mutation model ₃₅₈

We trained a context-dependent mutation model to capture the impact of 7-mer ₃₅₉
sequence context, local mutation rate, and sequencing coverage on the probability of the ₃₆₀
occurrence of each mutation in the gnomAD exome sequencing data. Because of the ₃₆₁

intrinsic sparsity of putatively neutral variants in coding regions, it is difficult to build a mutation model solely based on the gnomAD exome sequencing data. Therefore, we first built a mutation model based on neutral noncoding variants in the gnomAD whole genome sequencing (WGS) data. Then, we recalibrated the WGS-based mutation model in the gnomAD exome sequencing data to adjust for the differences in population sample size and sequencing coverage between the WGS and the exome sequencing data.

To build the WGS-based mutation model, we first compiled a list of putatively neutral noncoding regions following a strategy described in previous studies [10, 70, 71]. We removed coding exons [72], conserved phastCons elements [73], nucleotide sites within 1000 bp of any coding exons, and nucleotide sites within 100 bp of any phastCons elements. We assumed that the remaining noncoding regions were largely depleted with functional elements and, therefore, mutations in these regions were putatively neutral. It is worth noting that we also removed any nucleotide sites with an average sequencing coverage below 20, due to the difficulty of variant calling in low coverage regions, and any sites overlapping CpGs, due to the high mutation rates in these sites.

Then, we fit a WGS-based mutation model to the gnomAD WGS data in putatively neutral noncoding regions. First, for each possible combination of mutation l and sequence context k (the focal nucleotide with 3 flanking nucleotides on each side), we calculated its mutability, $f_{k \to l}$, defined as the proportion of 7-mer sequences of k with observed rare variant l in the gnomAD WGS data. We defined $F_{k \to l} = \text{logit}(f_{k \to l}) \equiv \log(\frac{f_{k \to l}}{1 - f_{k \to l}})$, which represents the contribution of sequence context and mutation type to the probability of variant occurrence in logit scale. Second, we fit a genome-wide logistic regression model to estimate the contributions of sequencing depth and context-dependent mutability to the probability of the occurrence of a mutation. This logistic regression model assumed

$$\text{logit}(Y_i^{\text{WGS}}) = \alpha_0 + \alpha_1 \log(d_i) + \alpha_2 F_{k_i \to l_i}, \tag{1}$$

where Y_i^{WGS} is a binary indicator of the occurrence of neutral noncoding mutation i in the gnomAD WGS data. d_i is the average sequencing depth at the nucleotide position of mutation i. k_i and l_i are the sequence context and the mutation type of mutation i, respectively. We then fit a local logistic regression model for each exon m in the human genome with

$$\text{logit}(Y_i^{\text{WGS}}) = \alpha_3^m + \hat{\alpha}_1 \log(d_i) + \hat{\alpha}_2 F_{k_i \to l_i}, \text{ for all mutation } i\text{'s} \\ \text{within 60 kb of exon } m \tag{2}$$

where α_3^m is an exon specific intercept independently estimated for each exon m, and $\hat{\alpha}_1$ and $\hat{\alpha}_2$ are the estimates of regression coefficients in Eq. 1. The local regression model effectively added an exon-specific, multiplicative scaling factor to adjust for the variation of local mutation rates across exons. We defined

$$q_i = \hat{\alpha}_3^m + \hat{\alpha}_1 \log(d_i) + \hat{\alpha}_2 F_{k_i \to l_i} \tag{3}$$

as the logit of the occurrence probability of mutation i in exon m in the WGS-based mutation model.

Finally, we recalibrated the WGS-based mutation model in the gnomAD exome sequencing data with a logistic regression model,

$$\text{logit}(Y_i^{\text{exome}}) = \beta_0 + q_i, \tag{4}$$

where Y_i^{exome} is a binary indicator of the presence of synonymous mutation i in the gnomAD exome sequencing data. The exome-based logistic regression model effectively added a multiplicative scaling factor to accommodate for the differences in population

sample size and sequencing coverage between the WGS and the exome sequencing data. In the final exome mutation model, we defined the probability of the occurrence of missense mutation i in gene j as

$$\mu_{ij} = \text{logistic}(\hat{\beta}_0 + q_i) \equiv \frac{\exp(\hat{\beta}_0 + q_i)}{1 + \exp(\hat{\beta}_0 + q_i)}, \tag{5}$$

where $\hat{\beta}_0$ is the maximum likelihood estimate of β_0 in Eq. 4. We fit all the logistic regression models using the glm function in R [74].

Details of the UNEECON model

We assume that negative selection on a potential mutation is a mathematical function of the sum of a variant-level fixed effect, z_{ij}, and a gene-level random effect, u_j (Fig. 1). Denoting $\mathbf{X}_{ij}$ as the vector of variant features associated with mutation i in gene j, we assume that the variant-level fixed effect, z_{ij}, can be modeled by a feedforward neural network. At the bottom of the neural network, a nonlinear hidden layer and a dropout layer are employed to transform $\mathbf{X}_{ij}$ to a hidden vector,

$$\mathbf{H}_{ij} = \text{droput}(\text{ReLU}(\mathbf{X}_{ij} \cdot \mathbf{W}_{\text{hidden}} + \mathbf{B}_{\text{hidden}})), \tag{6}$$

where **ReLU** and **dropout** are the rectified linear layer [75] and the dropout layer [43], respectively. $\mathbf{W}_{\text{hidden}}$ and $\mathbf{B}_{\text{hidden}}$ are the weight matrix and bias vector of the hidden layer, respectively. Then, we assume the fixed effect term, z_{ij}, is a linear combination of the hidden vector $\mathbf{H}_{ij}$,

$$z^{ij} = \mathbf{H}_{ij} \cdot \mathbf{W}_{\text{output}} + b_{\text{output}}, \tag{7}$$

where $\mathbf{W}_{\text{output}}$ and b_{output} are the weight vector and the bias term, respectively. We use the Glorot method [76] to initialize the weights, $\mathbf{W}_{\text{hidden}}$ and $\mathbf{W}_{\text{output}}$, and initialize the bias terms, $\mathbf{B}_{\text{hidden}}$ and b_{output}, with zeros. In the dropout layer, we use a fixed dropout rate of 0.5. It is worth noting that, if a linear version of the UNEECON model is used, Eq. 6 will be replaced by an identity function $\mathbf{H}_{ij} \equiv \mathbf{X}_{ij}$.

To capture the variation of gene-level selective constraints that cannot be predicted from feature vector $\mathbf{X}_{ij}$, we introduce a gene-level random-effect term following a Gaussian distribution,

$$u_j \sim \mathcal{N}(0, \sigma), \tag{8}$$

where σ is the standard deviation of the Gaussian distribution. Given the fixed and random effect terms, we assume η_{ij}, the relative probability of the occurrence of mutation i in gene j, follows

$$\eta_{ij} = \text{logistic}(z_{ij} + u_j) \equiv \frac{\exp(z_{ij} + u_j)}{1 + \exp(z_{ij} + u_j)}. \tag{9}$$

We further assume that the total probability of the occurrence of potential missense mutation i in gene j is equal to the product of η_{ij} and μ_{ij}, the probability of variant occurrence under the neutral mutation model described in Eq. 5. Accordingly, the likelihood function for the data associated with gene j is defined as

$$\mathcal{L}_j = \prod_i (\eta_{ij}\mu_{ij})^{Y_{ij}} (1 - \eta_{ij}\mu_{ij})^{1-Y_{ij}}, \tag{10}$$

where Y_{ij} is a binary indicator of the occurrence of missense mutation i in gene j in the gnomAD exome sequencing data.

Training the UNEECON model

We trained UNEECON with the Adam optimization algorithm [77]. In each iteration, UNEECON loaded a mini-batch of data consisting of all the potential missense mutations in a single gene. Then, we calculated the log likelihood of the mini-batch of data using Eq. 10 and numerically integrated out u_j with a 20-point Gaussian Quadrature rule. The negative log value of Eq. 10 was used as the objective function in the Adam optimizer.

We randomly split the data into a training set (80% genes), a validation set (10% genes), and a test set (10% genes). After each epoch of training, UNEECON evaluated the objective function in the validation set and stopped training when the objective function did not improve over 5 successive epochs to avoid overfitting (early stopping). Furthermore, we performed a grid search to optimize two hyperparameters, *i.e.*, the learning rate of the Adam algorithm (10^{-2}, 10^{-3}, and 10^{-4}) and the number of hidden units (64, 128, 256, 512, and a linear model without hidden units). The model was evaluated on the test dataset to choose optimal hyperparameters.

Calculating UNEECON and UNEECON-G scores

After training, we fixed the free parameters, *i.e.*, $\mathbf{W}_{\text{hidden}}$, $\mathbf{W}_{\text{output}}$, $\mathbf{B}_{\text{hidden}}$, b_{output}, and σ, to the estimated values. Then, we calculated the UNEECON score for all the potential missense mutations in the human genome. For each gene j, UNEECON first calculated the posterior distribution of the random effect, $\mathbb{P}(u_j|\text{Data}_j)$, where $\text{Data}_j = \{\mathbf{X}_{1j}, \mathbf{X}_{2j}, \cdots, \mathbf{X}_{Nj}, Y_{1j}, Y_{2j}, \cdots, Y_{Nj}\}$ are the features and indicators of occurrence of all the potential missense mutations in gene j. Denoting $\mathbb{E}_{u_j|\text{Data}_j}(\eta_{ij}|u_j) \equiv \int \eta_{ij} \mathbb{P}(u_j|\text{Data}_j) du_j$ as expected relative probability of the occurrence of mutation i in gene j, we define the UNEECON score of mutation i as $1 - \mathbb{E}_{u_j|\text{Data}}(\eta_{ij}|u_j)$. The UNEECON-G score is defined as the average UNEECON score of all the missense mutations in a gene.

Distributions of UNEECON scores across gene categories and protein regions

We investigated the distributions of UNEECON scores across different gene categories and protein regions. We obtained lists of haploinsufficient genes [37] ($n = 294$), autosomal dominant disease genes [35, 36] ($n = 709$), autosomal recessive disease genes [35, 36] ($n = 1,183$), and olfactory receptor genes [45] ($n = 371$) from the GitHub repository for the MacArthur Lab at the Broad Institute (https://github.com/macarthur-lab/gene_lists). We obtained annotations of α-helices, β-strands, hydrogen-bonded turns, enzyme active sites, and binding sites from UniProt [47]. We obtained annotated disordered protein regions from MobiDB 3.0 [48]. We plotted the distributions of UNEECON scores using R [74].

Comparison with other methods in the prediction of ClinVar pathogenic variants

We evaluated the performance of UNEECON, compared with eight previously published methods, in the setting of predicting missense pathogenic variants in the ClinVar database downloaded on Feb 25, 2019 [30]. We obtained MPC [25], PrimateAI [11], Eigen [50], CADD [8], RVIS [16], and pLI [17] scores from dbNSFP (version 4.0b2a) [78]. We obtained LASSIE scores [12] from the LASSIE GitHub repository (https://github.com/CshlSiepelLab/LASSIE) and CCR score (version 2) [22] from https://s3.us-east-2.amazonaws.com/ccrs/ccrs/ccrs.autosomes.v2.20180420.bed.gz. It is

worth noting that we used the version of pLI scores trained on the gnomAD exome sequencing data [29].

We considered autosomal missense variants with annotations of "pathogenic/likely pathogenic" as positives and the ones with annotations of "benign/likely benign" as negative controls. We further removed the variants with conflict pathogenicity annotations and the variants present in the gnomAD exome sequencing dataset [29]. For all the comparisons, we only included the variants scored by all methods. We plotted the receiver operating characteristic (ROC) curves and calculated the areas under the receiver operating characteristic curves (AUCs) using the ROCR package [79]. Because the AUC metrics is sensitive to label imbalance, we matched the numbers of positives and negatives using random sampling without replacement.

Comparison with alternative methods in predicting *de novo* mutations associated with developmental disorders

We downloaded *de novo* mutations identified in 4,293 individuals affected by developmental disorders [31] and 2,278 healthy individuals from denovo-db (version v1.6.1). The healthy controls included individuals enrolled in previous studies of autism [80–84], severe non-syndromic sporadic intellectual disability [85], schizophrenia [86], and healthy populations [87–90]. We removed redundant variants and any variants presented in the gnomAD exome sequencing data. Then, for various percentile rank cutoffs (top 10%, 20%, 30%, and 40%), we calculated the $\log_2$ odds ratio of the enrichment of predicted deleterious variants in affected individuals using the fisher.test function in R [74].

Comparison with alternative methods in predicting disease genes and essential genes

We obtained four sets of disease genes and essential genes from the GitHub repository for the MacArthur Lab at the Broad Institute (https://github.com/macarthur-lab/gene_lists). These gene sets included 294 haploinsufficient genes [37], 709 autosomal dominant disease genes [35, 36], 2,454 human orthologs of mouse essential genes [33, 34], and 683 essential genes based on CRISPR screen data from human cell lines [32]. For each set of the haploinsufficient, autosomal dominant, and mouse essential genes, we utilized the other autosomal genes as negative controls and constructed a negative gene set with matched gene length and gene number using MatchIt [91]. Similarly, for the 683 human essential genes based on CRISPR screen data, we used the 913 nonessential genes from the same study [32] as negative controls and constructed a negative gene set with matched gene length and gene number. We plotted the ROC curves and calculated the AUC metrics using the ROCR package [79].

Comparison of UNEECON-G and pLI scores

To evaluate the relationship between gene-level intolerance to missense mutations and that to loss-of-function mutations, we compared the UNEECON-G scores with the pLI scores trained on the gnomAD data [17, 78]. We identified 1,912 genes intolerant to loss-of-function mutations based on a threshold of pLI score of 0.9. Then, we split the 1,912 genes into 956 genes tolerant to missense genes and 956 genes intolerant to missense mutations based on the median UNEECON-G score. We used PANTHER [92] to calculate the enrichment of Reactome pathways and Gene Ontology categories in the set of genes intolerant to both missense and loss-of-function mutations, compared with those tolerant to missense constraints but not to loss-of-function mutations. We also

downloaded 1,054 autism related genes from the SFARI database on March 9, 2019 [57], and evaluated the enrichment of autism genes in the two gene sets.

Availability of data and materials

The UNEECON program and precomputed UNEECON/UNEECON-G scores are available at https://github.com/yifei-lab/UNEECON.

Supporting information

S1 File.   Supplementary material. Supplementary figures and tables.

S2 File.   Supplementary dataset. List of 1,912 genes intolerant to loss-of-function mutations.

Acknowledgments

The author thanks Noah Dukler for comments on the manuscript. This work was supported by start-up fund from the Pennsylvania State University.

References

1. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genetics in Medicine. 2015;17(5):405–423.

2. Maxwell K, Hart S, Vijai J, Schrader K, Slavin T, Thomas T, et al. Evaluation of ACMG-Guideline-Based Variant Classification of Cancer Susceptibility and Non-Cancer-Associated Genes in Families Affected by Breast Cancer. The American Journal of Human Genetics. 2016;98(5):801–817.

3. Eilbeck K, Quinlan A, Yandell M. Settling the score: variant prioritization and Mendelian disease. Nature Reviews Genetics. 2017;18(10):599–612.

4. Reva B, Antipin Y, Sander C. Predicting the functional impact of protein mutations: application to cancer genomics. Nucleic Acids Research. 2011;39(17):e118.

5. Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein function. Nucleic Acids Research. 2003;31(13):3812–3814.

6. Cooper GM, Shendure J. Needles in stacks of needles: finding disease-causal variants in a wealth of genomic data. Nature Reviews Genetics. 2011;12(9):628–640.

7. Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the Functional Effect of Amino Acid Substitutions and Indels. PLOS ONE. 2012;7(10):1–13.

8. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. Nature Genetics. 2014;46(3):310–315.

9. Gulko B, Hubisz MJ, Gronau I, Siepel A. A method for calculating probabilities of fitness consequences for point mutations across the human genome. Nature Genetics. 2015;47(3):276–283.

10. Huang YF, Gulko B, Siepel A. Fast, scalable prediction of deleterious noncoding variants from functional and population genomic data. Nature Genetics. 2017;49(4):618–624.

11. Sundaram L, Gao H, Padigepati SR, McRae JF, Li Y, Kosmicki JA, et al. Predicting the clinical impact of human mutation with deep neural networks. Nature Genetics. 2018;50(8):1161–1170.

12. Huang YF, Siepel A. Estimation of allele-specific fitness effects across human protein-coding sequences and implications for disease. Genome Research. 2019;29(8):1310–1321.

13. Khurana E, Fu Y, Colonna V, Mu XJ, Kang HM, Lappalainen T, et al. Integrative Annotation of Variants from 1092 Humans: Application to Cancer Genomics. Science. 2013;342(6154):1235587.

14. Fu Y, Liu Z, Lou S, Bedford J, Mu X, Yip K, et al. FunSeq2: a framework for prioritizing noncoding regulatory variants in cancer. Genome Biology. 2014;15(10):480.

15. Gulko B, Siepel A. An evolutionary framework for measuring epigenomic information and estimating cell-type-specific fitness consequences. Nature Genetics. 2019;51(2):335–342.

16. Petrovski S, Wang Q, Heinzen EL, Allen AS, Goldstein DB. Genic Intolerance to Functional Variation and the Interpretation of Personal Genomes. PLOS Genetics. 2013;9(8):e1003709.

17. Samocha KE, Robinson EB, Sanders SJ, Stevens C, Sabo A, McGrath LM, et al. A framework for the interpretation of de novo mutation in human disease. Nature Genetics. 2014;46(9):944–950.

18. Petrovski S, Gussow AB, Wang Q, Halvorsen M, Han Y, Weir WH, et al. The Intolerance of Regulatory Sequence to Genetic Variation Predicts Gene Dosage Sensitivity. PLoS Genet. 2015;11(9):e1005492.

19. Itan Y, Shang L, Boisson B, Patin E, Bolze A, Moncada-Vélez M, et al. The human gene damage index as a gene-level approach to prioritizing exome variants. Proceedings of the National Academy of Sciences. 2015;112(44):13615–13620.

20. Gussow A, Petrovski S, Wang Q, Allen A, Goldstein D. The intolerance to functional genetic variation of protein domains predicts the localization of pathogenic mutations within genes. Genome Biology. 2016;17(1):9.

21. Pérez-Palma E, May P, Iqbal S, Niestroj LM, Du J, Heyne H, et al. Identification of pathogenic variant enriched regions across genes and gene families. bioRxiv. 2019;doi:https://doi.org/10.1101/641043.

22. Havrilla JM, Pedersen BS, Layer RM, Quinlan AR. A map of constrained coding regions in the human genome. Nature Genetics. 2019;51(1):88–95.

23. Silk M, Petrovski S, Ascher DB. MTR-Viewer: identifying regions within genes under purifying selection. Nucleic Acids Research. 2019;47(W1):W121–W126.

24. Iossifov I, Levy D, Allen J, Ye K, Ronemus M, Lee Yh, et al. Low load for disruptive mutations in autism genes and their biased transmission. Proceedings of the National Academy of Sciences. 2015;112(41):E5600–E5607.

25. Samocha KE, Kosmicki JA, Karczewski KJ, O'Donnell-Luria AH, Pierce-Hoffman E, MacArthur DG, et al. Regional missense constraint improves variant deleteriousness prediction. bioRxiv. 2017;doi:https://doi.org/10.1101/148353.

26. Jagadeesh KA, Wenger AM, Berger MJ, Guturu H, Stenson PD, Cooper DN, et al. M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. Nature Genetics. 2016;48(12):1581–1586.

27. Evans P, Wu C, Lindy A, McKnight DA, Lebo M, Sarmady M, et al. Genetic variant pathogenicity prediction trained using disease-specific clinical sequencing data sets. Genome Research. 2019;29(7):1144–1151.

28. Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al. Analysis of protein-coding genetic variation in 60,706 humans. Nature. 2016;536(7616):285–291.

29. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alfoldi J, Wang Q, et al. Variation across 141,456 human exomes and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding genes. bioRxiv. 2019;doi:10.1101/531210.

30. Landrum MJ, Lee JM, Riley GR, Jang W, Rubinstein WS, Church DM, et al. ClinVar: public archive of relationships among sequence variation and human phenotype. Nucleic Acids Research. 2014;42(D1):D980–D985.

31. Deciphering Developmental Disorders Study, McRae JF, Clayton S, Fitzgerald TW, Kaplanis J, Prigmore E, et al. Prevalence and architecture of de novo mutations in developmental disorders. Nature. 2017;542:433–438.

32. Hart T, Brown KR, Sircoulomb F, Rottapel R, Moffat J. Measuring error rates in genomic perturbation screens: gold standards for human functional genomics. Mol Syst Biol. 2014;10(7):733.

33. Blake JA, Bult CJ, Kadin JA, Richardson JE, Eppig JT, the Mouse Genome Database Group. The Mouse Genome Database (MGD): premier model organism resource for mammalian genomics and genetics. Nucleic Acids Research. 2010;39(suppl1):D842–D848.

34. Georgi B, Voight BF, Bucan M. From Mouse to Human: Evolutionary Genomics Analysis of Human Orthologs of Essential Genes. PLOS Genetics. 2013;9(5):e1003484.

35. Blekhman R, Man O, Herrmann L, Boyko AR, Indap A, Kosiol C, et al. Natural Selection on Genes that Underlie Human Disease Susceptibility. Current Biology. 2008;18(12):883–889.

36. Berg JS, Adams M, Nassar N, Bizon C, Lee K, Schmitt CP, et al. An informatics approach to analyzing the incidentalome. Genetics In Medicine. 2012;15:36.

37. Rehm HL, Berg JS, Brooks LD, Bustamante CD, Evans JP, Landrum MJ, et al. ClinGen – The Clinical Genome Resource. N Engl J Med. 2015;372(23):2235–2242.

38. Armon A, Graur D, Ben-Tal N. ConSurf: an algorithmic tool for the identification of functional regions in proteins by surface mapping of phylogenetic information. Journal of Molecular Biology. 2001;307(1):447–463.

39. Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglou S, Sidow A. Distribution and intensity of constraint in mammalian genomic sequence. Genome Research. 2005;15(7):901–913.

40. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. Genome Research. 2010;20(1):110–121.

41. Huang YF, Golding GB. Phylogenetic Gaussian Process Model for the Inference of Functionally Important Regions in Protein Tertiary Structures. PLoS Computational Biology. 2014;10(1):e1003429.

42. Huang YF, Golding GB. FuncPatch: a web server for the fast Bayesian inference of conserved functional patches in protein 3D structures. Bioinformatics. 2015;31(4):523–531.

43. Srivastava N, Hinton G, Krizhevsky A, Sutskever I, Salakhutdinov R. Dropout: A Simple Way to Prevent Neural Networks from Overfitting. Journal of Machine Learning Research. 2014;15:1929–1958.

44. Bengio Y. Practical Recommendations for Gradient-Based Training of Deep Architectures. In: Neural Networks: Tricks of the Trade: Second Edition. Berlin, Heidelberg: Springer Berlin Heidelberg; 2012. p. 437–478.

45. Mainland JD, Li YR, Zhou T, Liu WLL, Matsunami H. Human olfactory receptor responses to odorants. Scientific Data. 2015;2:150002.

46. Gilad Y, Bustamante CD, Lancet D, Pääbo S. Natural Selection on the Olfactory Receptor Gene Family in Humans and Chimpanzees. The American Journal of Human Genetics. 2003;73(3):489–501.

47. McGarvey PB, Nightingale A, Luo J, Huang H, Martin MJ, Wu C, et al. UniProt genomic mapping for deciphering functional effects of missense variants. Human mutation. 2019;40(6):694–705.

48. Piovesan D, Tabaro F, Paladin L, Necci M, Mičetić I, Camilloni C, et al. MobiDB 3.0: more annotations for intrinsic disorder, conformational diversity and interactions in proteins. Nucleic Acids Research. 2017;46(D1):D471–D476.

49. The UniProt Consortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Research. 2018;47(D1):D506–D515.

50. Ionita-Laza I, McCallum K, Xu B, Buxbaum JD. A spectral approach integrating functional genomic annotations for coding and noncoding variants. Nature Genetics. 2016;48(2):214–220.

51. Turner TN, Yi Q, Krumm N, Huddleston J, Hoekzema K, F Stessman HA, et al. denovo-db: a compendium of human de novo variants. Nucleic Acids Research. 2016;45(D1):D804–D811.

52. Wright PE, Dyson HJ. Intrinsically disordered proteins in cellular signalling and regulation. Nature Reviews Molecular Cell Biology. 2015;16:18.

53. Brown CJ, Takayama S, Campen AM, Vise P, Marshall TW, Oldfield CJ, et al. Evolutionary Rate Heterogeneity in Proteins with Long Disordered Regions. Journal of Molecular Evolution. 2002;55(1):104–110.

54. Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The Reactome Pathway Knowledgebase. Nucleic Acids Research. 2018;46(D1):D649–D655.

55. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nature Genetics. 2000;25(1):25–29.

56. The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing strong. Nucleic Acids Research. 2019;47(D1):D330–D338.

57. Abrahams BS, Arking DE, Campbell DB, Mefford HC, Morrow EM, Weiss LA, et al. SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). Molecular Autism. 2013;4(1):36.

58. Grimm DG, Azencott CA, Aicheler F, Gieraths U, MacArthur DG, Samocha KE, et al. The Evaluation of Tools Used to Predict the Impact of Missense Variants Is Hindered by Two Types of Circularity. Human Mutation. 2015;36(5):513–523.

59. Fuller ZL, Berg JJ, Mostafavi H, Sella G, Przeworski M. Measuring intolerance to mutation in human genetics. Nature Genetics. 2019;51(5):772–776.

60. Wainschtein P, Jain DP, Yengo L, Zheng Z, Cupples LA, Shadyab AH, et al. Recovery of trait heritability from whole genome sequence data. bioRxiv. 2019;doi:https://doi.org/10.1101/588020.

61. Starita LM, Ahituv N, Dunham MJ, Kitzman JO, Roth FP, Seelig G, et al. Variant Interpretation: Functional Assays to the Rescue. The American Journal of Human Genetics. 2017;101(3):315–325.

62. Kinney JB, McCandlish DM. Massively Parallel Assays and Quantitative Sequence-Function Relationships. Annu Rev Genom Hum Genet. 2019;20:99–127.

63. Massingham T, Goldman N. Detecting amino acid sites under positive selection and purifying selection. GENETICS. 2005;169(3):1753–1762.

64. Grantham R. Amino Acid Difference Formula to Help Explain Protein Evolution. Science. 1974;185(4154):862–864.

65. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. Nature Methods. 2010;7:248–249.

66. Chun S, Fay JC. Identification of deleterious mutations within three human genomes. Genome Research. 2009;19(9):1553–1561.

67. Wong WC, Kim D, Carter H, Diekhans M, Ryan MC, Karchin R. CHASM and SNVBox: toolkit for detecting biologically important single nucleotide mutations in cancer. Bioinformatics. 2011;27(15):2147–2148.

68. Xiong HY, Alipanahi B, Lee LJ, Bretschneider H, Merico D, Yuen RKC, et al. The human splicing code reveals new insights into the genetic determinants of disease. Science. 2015;347:1254806.

69. Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al. Integrative analysis of 111 reference human epigenomes. Nature. 2015;518(7539):317–330.

70. Arbiza L, Gronau I, Aksoy BA, Hubisz MJ, Gulko B, Keinan A, et al. Genome-wide inference of natural selection on human transcription factor binding sites. Nature Genetics. 2013;45(7):723–729.

71. Gronau I, Arbiza L, Mohammed J, Siepel A. Inference of Natural Selection from Interspersed Genomic Elements Based on Polymorphism and Divergence. Molecular Biology and Evolution. 2013;30(5):1159–1171.

72. Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, et al. GENCODE: The reference human genome annotation for The ENCODE Project. Genome Research. 2012;22(9):1760–1774.

73. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Research. 2005;15(8):1034–1050.

74. Team RDC. R: A Language and Environment for Statistical Computing; 2008. Available from: `http://www.R-project.org`.

75. Nair V, Hinton GE. Rectified Linear Units Improve Restricted Boltzmann Machines. In: Proceedings of the 27th International Conference on International Conference on Machine Learning. ICML'10. USA: Omnipress; 2010. p. 807–814.

76. Glorot X, Bengio Y. Understanding the difficulty of training deep feedforward neural networks. In: Teh YW, Titterington M, editors. Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics. vol. 9 of Proceedings of Machine Learning Research. Chia Laguna Resort, Sardinia, Italy: PMLR; 2010. p. 249–256.

77. Kingma DP, Ba J. Adam: A Method for Stochastic Optimization. arXiv:14126980. 2014;.

78. Liu X, Jian X, Eric B. dbNSFP v2.0: a database of human non-synonymous SNVs and their functional predictions and annotations. Human Mutation. 2013;34(9):E2393–E2402.

79. Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCR: visualizing classifier performance in R. Bioinformatics. 2005;21(20):3940–3941.

80. Iossifov I, O'Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, et al. The contribution of de novo coding mutations to autism spectrum disorder. Nature. 2014;515:216–221.

81. Krumm N, Turner TN, Baker C, Vives L, Mohajeri K, Witherspoon K, et al. Excess of rare, inherited truncating mutations in autism. Nature Genetics. 2015;47(6):582–588.

82. Turner T, Hormozdiari F, Duyzend M, McClymont S, Hook P, Iossifov I, et al. Genome Sequencing of Autism-Affected Families Reveals Disruption of Putative Noncoding Regulatory DNA. The American Journal of Human Genetics. 2016;98(1):58–74.

83. Yuen RKC, Merico D, Bookman M, L Howe J, Thiruvahindrapuram B, Patel RV, et al. Whole genome sequencing resource identifies 18 new candidate genes for autism spectrum disorder. Nature Neuroscience. 2017;20:602–611.

84. Werling DM, Brand H, An JY, Stone MR, Zhu L, Glessner JT, et al. An analytical framework for whole-genome sequence association studies and its implications for autism spectrum disorder. Nature Genetics. 2018;50(5):727–736.

85. Rauch A, Wieczorek D, Graf E, Wieland T, Endele S, Schwarzmayr T, et al. Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: an exome sequencing study. The Lancet. 2012;380(9854):1674–1682.

86. Gulsuner S, Walsh T, Watts A, Lee M, Thornton A, Casadei S, et al. Spatial and Temporal Mapping of De Novo Mutations in Schizophrenia to a Fetal Prefrontal Cortical Network. Cell. 2013;154(3):518–529.

87. The 1000 Genomes Project, Conrad DF, Keebler JEM, DePristo MA, Lindsay SJ, Zhang Y, et al. Variation in genome-wide mutation rates within and between human families. Nature Genetics. 2011;43(7):712–714.

88. Ramu A, Noordam MJ, Schwartz RS, Wuster A, Hurles ME, Cartwright RA, et al. DeNovoGear: de novo indel and point mutation discovery and phasing. Nature Methods. 2013;10(10):985–987.

89. The Genome of the Netherlands Consortium. Whole-genome sequence variation, population structure and demographic history of the Dutch population. Nature Genetics. 2014;46:818–825.

90. Besenbacher S, Liu S, Izarzugaza JMG, Grove J, Belling K, Bork-Jensen J, et al. Novel variation and de novo mutation rates in population-wide de novo assembled Danish trios. Nature Communications. 2015;6:5969.

91. Ho DE, Imai K, King G, Stuart EA. MatchIt: Nonparametric Preprocessing for Parametric Causal Inference. Journal of Statistical Software. 2011;42(8):1–28.

92. Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, et al. PANTHER version 11: expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool enhancements. Nucleic Acids Research. 2017;45(D1):D183–D189.