Title: Movement and space-use responses of a large herbivore to experimental alteration of

resource availability

List of Authors: Nathan Ranc[1,2], Paul R. Moorcroft[1], K. Whitney Hansen[1,2a], Federico Ossi[2],

Tobia Sforna[2], Enrico Ferraro[3], Alessandro Brugnoli[3] and Francesca Cagnacci[1,2]

Affiliations

[1]Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford Street,

Cambridge MA02138, USA.

[2]Department of Biodiversity and Molecular Ecology, Research and Innovation Centre,

Fondazione Edmund Mach, Via E. Mach 1, 38010 San Michele all'Adige, Italy.

[3]Trentino Hunting Association, Via Guardini 41, 38121 Trento, Italy.

Corresponding Author: Nathan Ranc

[a] Current affiliation: Environmental Studies Department, University of California Santa Cruz, USA

Abstract

Resource acquisition is a key factor governing patterns of animal movement. While the link
between spatio-temporal resource patterns and movement behaviour has been widely
documented for migration, our understanding of home ranging behaviour in relation to resource
dynamics has been limited by challenges of quantifying resource heterogeneity at fine spatio-
temporal scales. In this study, we addressed this issue by analysing the movement responses of a
resident large herbivore in response to an *in situ* manipulation of a high-quality, concentrated
food resource. Specifically, we fitted roe deer (*Capreolus capreolus*) with GPS collars in the
Eastern Italian Alps and recorded their fine-scale movement responses to an alteration of feeding
site accessibility, for a total of 25 animal-years. The experiment involved identifying, for each
animal, a familiar feeding site, whose accessibility was transitorily restricted (i.e., closed) while
maintaining the availability of forage at alternative feeding sites. First, we tested whether
individuals altered their spatial behaviour to track dynamical changes in resource availability.
Experimental closure led to larger, spatially-shifted home ranges, resulting from more
exploratory movements. Individual roe deer compensated the loss of their familiar feeding site
by using alternative ones, and by doing so maintained their overall use of these concentrated
resources. Our results demonstrate that roe deer actively track resource dynamics, and rapidly
adjust their space-use and movement behaviour in response to changes in resource availability.
Second, we showed that individual resource preferences mediate these behavioural adjustments:
individuals characterized by a high preference for feeding sites exhibited larger space-use shifts
and stronger changes in resource use in response to the resource manipulation. In contrast, sex
had relatively minor influence on the observed patterns. Third, we investigated the role of site
familiarity in roe deer foraging decisions. We found that space-use and resource use during the

38 post-closure period were restored to pre-closure patterns, thereby providing evidence for the

39 inherent benefits of site familiarity to animals maintaining a home range. Together, our results

40 establish the connections between movement, space-use, individual preference, and the spatio-

41 temporal pattern of resources in deer home ranging behaviour.

Introduction

Animals move to change the environmental context they experience (Van Moorter,

Rolandsen, Basille, & Gaillard, 2016), including abiotic conditions, the presence of predators

and competitors, and the availability of resources. Because foraging efficiency can be linked to

individual fitness (Stephens & Krebs, 1986), food acquisition is thought to be a primary driver

underlying animal movements (e.g., Mueller & Fagan, 2008). Consequently, space-use

represents the geographic realization of optimizing fitness as a function of resource availability

and acquisition costs (Mitchell & Powell, 2004).

Food resources are usually dynamic in both space and time (Wiens, 1976). In the case of

herbivores, animals typically feed on resources distributed in patches, which are characterized by

important temporal variations in quantity and quality (Owen-Smith, Fryxell, & Merrill, 2010). In

this context, Mueller et al. (2011) have shown that strong spatio-temporal gradients in resource

availability at either landscape or regional scales appear to drive migration and nomadism tactics.

In many herbivore populations, however, individuals show a high year-round fidelity to a

spatially-localized home range. It has been suggested that the foraging benefits of site

familiarity, where resources are constant or predictable, are responsible for the formation of a

stable home range (see Fagan et al., 2013 for a review). While the home range has traditionally

been perceived as a relatively static space-use tactic, recent evidence suggests that animals have

sub-seasonal home ranges (Benhamou 2014), i.e. focus their movements into particular areas in

response to seasonal variation in local resource availability. For example, Couriot et al. (2018)

have shown that in roe deer (*Capreolus capreolus*) sub-seasonal home ranges are a ubiquitous

behavioural tactic across a wide ecological gradient.

The link between movement behaviour and resource dynamics is less clear when

observing home ranging behaviour than migration or nomadism (Mueller & Fagan, 2008),

because of the difficulty to quantify spatio-temporal variability in resource heterogeneity at small

spatial scales (Couriot et al., 2018). In this study, we address this issue by experimentally

manipulating the spatio-temporal patterns of food availability within home ranges. *In situ* food

manipulation experiments have a long history in the study of population dynamics, with a

primary focus on understanding the numerical response to food supplementation (e.g., Krebs,

1971), and of animal communities (e.g., Brown & Munger, 1985). Although these field

experiments have provided fundamental insights in animal ecology, they have seldom been

combined with the emerging technological capabilities of animal tracking (Cagnacci, Boitani,

Powell, & Boyce, 2010) to investigate the implications of food availability on individual

movements and space-use. In a rare example of field experiment in large herbivores, Kilpatrick

and Stober (2002) showed that white-tailed deer (*Odocoileus virginianus*) shifted their core-area

i.e., familiar areas of use, in response to novel food supplementation. In a second experiment,

van Overveld and Matthysen (2010) demonstrated that the individual variability in responses to

an alteration of resource distribution are personality-dependent in great tits (*Parus major*). Our

research builds upon these two studies by investigating the spatial responses of a large herbivore,

roe deer, to an experimental *in situ* manipulation of a high-quality, concentrated food resource in

relation to both individual resource preferences and site familiarity.

Observational studies have suggested that, as browsers with limited fat reserves

(Andersen, Gaillard, Linnell, & Duncan, 2000), roe deer exhibit a tight association between

movement and resource dynamics (Ossi et al. 2017) with a strong plasticity to adapt its resource

acquisition at different spatio-temporal scales (De Groeve et al., 2019; Morellet et al., 2013;

Peters et al., 2017). In contrast to group-living ungulates, the foraging decisions of roe deer are

expected to be clearly expressed at the level of individuals. The experiment was conducted

during winter, when food scarcity limits roe deer foraging performance, and individuals are most

inclined to adjust their spatial behaviour to continue meeting their energy requirements (Ossi,

Gaillard, Hebblewhite, & Cagnacci, 2015).

We tagged roe deer in the Eastern Italian Alps with GPS units and followed their

movements during transitory alterations of food availability at supplemental feeding sites (FS),

i.e. discrete resource patches with an identifiable resource value distinguishable from the

vegetation matrix (Mitchell & Powell, 2004; Wiens, 1976). The experiment therefore mimics –

on free-ranging animals – the variation in the availability of concentrated, high-reward resources

akin to watering holes for savannah ungulates (Owen-Smith, 2004) and feral horses (*Equus*

ferus; Rozen-Rechels et al. 2015), or fruit trees for hornbills (*Ceratogymna atrata* and *C.*

cylindricus; Holbrook et al. 2002), forest chimpanzees (*Pan troglodytes*; Normand et al. 2009)

and other frugivorous primates.

In large herbivores, and roe deer in particular, our initial hypothesis states that individuals

alter their movement behaviours and consequently space-use patterns to track dynamics in

resource availability (H1; Table 1). We predicted that the loss of a key foraging resource should

lead to larger (P1.1), and spatially-shifted (P1.2) home ranges, resulting from more explorative

movements (P1.3). Furthermore, we predicted that roe deer reduced the intensity of use of the

familiar FS when food accessibility was prevented (P1.4a) and compensated for this loss by

using other accessible FS (P1.4b).

We further hypothesized that the behavioural adjustments to changes in resource

availability would vary between individuals (H2; Table 1). In particular, because roe deer males

have been shown to maintain a high year-round fidelity to their summer territory (Linnell &

Andersen, 1998), we predicted that they would respond less markedly to the experiment than

females (P2.1). We also predicted the responsiveness of roe deer to be positively influenced by

the individual's prior preference for FS (P2.2).

If the spatial patterns of roe deer home ranges result from the foraging benefits of site

familiarity (Riotte-lambert, Benhamou, & Chamaillé-Jammes, 2015; Van Moorter et al., 2009),

animals should strive to use familiar areas and resources when accessible (H3; Table 1).

Accordingly, we predicted that when initial conditions of food accessibility are re-established

after perturbation, the initial space-use patterns would be restored (P3.1), following a return to

high use of the familiar FS (P3.2).

Materials and methods

Study area

Roe deer is the primary large herbivore in the study area (7-8 individuals km^{-2}), located in

the north-eastern Italian Alps (Argentario range, Autonomous Province of Trento). The area is

characterized by a continental climate (mean temperature of January: 1.0 °C; July: 21.0 °C; mean

annual rainfall: 966 m) with occasional snow cover, and is largely forested (80%). It

encompasses four hunting reserves in which selective hunting occurs between September and

December (see Appendix S1 for further details).

Supplemental feeding management of roe deer is conducted year-round at > 50 distinct

feeding sites (FS; Appendix S2: Fig. S2.1). FS are typically wooden hopper dispensers that

provide a continuous supply of corn accessible through a tray (Fig. 1a). They are managed by

private hunters for roe deer but are also attended sporadically by red deer (*Cervus elaphus*).

Experimental design

We took advantage of roe deer use of a focal, identifiable resource – the FS – to design an

in situ experimental manipulation of resource availability. We created three successive

experimental phases of the availability of this resource – pre-closure, closure and post-closure –

by physically managing the accessibility of food at the FS. During the closure phase, access to

forage at FS was transitorily restricted by placing wooden boards obstructing the tray; these were

then removed again in the post-closure phase (Fig. 1).

The experiment was conducted between January and April, when roe deer use of

supplemental feeding is the most intense (Ossi et al., 2017), for three consecutive winters (2017,

2018 and 2019). We implemented the experiment on 18 individuals, including five recaptures

and two deployments spanning two winters, leading to a total of 25 animal-years (21 adults: 15

females, 6 males; 4 yearlings: 2 females, 2 males; sample size *n*=4, 11 and 10 in 2017, 2018 and

2019 respectively; see Appendix S2 for details). The animal-year was our sampling unit, on the

assumption that the same individual may respond independently to manipulations in different

years. Roe deer were captured using baited box traps (n=16) or net drives (n=3), and were fitted

with GPS-GSM radio collars programmed to acquire hourly GPS locations for a year, after

which they were released via a drop-off mechanism. Radio-collared roe deer moved an average

of 61.2 m per hour. This value of the average hourly movement distance *(l)* was subsequently

utilized in the analyses described below.

For all captured animals, we assumed a post-capture response in ranging behaviour. We

therefore considered the first re-visitation of the capture location as the sign of resettlement in

the original range and we used this time as the earliest possible onset of the experimental pre-

closure phase. Although not all the individuals were manipulated at the same time, we avoided

interference between capture operations and FS manipulations and between co-occurring

different manipulation phases (i.e. ensuring that co-occurring manipulations occurred in separate

areas).

During the *pre-closure phase*, we assessed the use of FS by radio-collared roe deer. We

identified the "main" FS (M) for each individual as the site with the largest number of locations

within a radius l during this initial phase, and considered it as the FS to which an individual is

most familiar. During the *closure phase*, corn was made inaccessible at M for a duration of about

15 days, depending on personnel availabilities (min=14.0 days, max=18.1, mean=15.5). M was

then re-opened, thereby initiating the *post-closure phase*. During both pre- and post-closure

phases, corn was available *ad libitum* at M. All "other" (O) managed FS – i.e., that were

provisioned at least once in the month prior to the experiment – had corn available *ad libitum*

throughout the experiment.

Data preparation

To ensure meaningful comparisons between animal-years, we homogenized the durations

of each experimental phase to the minimum length of the closure phase (i.e., 14 days).

Specifically, we truncated the movement data by removing initial excess positions for the pre-

closure and closure phases, and terminal excess positions for the post-closure phase. GPS

acquisition success was extremely high (99.57 % during the experiment) and we did not

interpolate missing fixes in the collected data.

The analyses of space-use and movement behaviour were based on spatially-explicit, raw

movement trajectories. The analyses of resource use relied on spatially-implicit, state time series

derived from the underlying movement data. To this end, we created an initial time series, for

each animal, by intersecting the relocations with three spatial domains: vegetation (the matrix;

V), main FS (M) and other FS (O). We converted FS locations (M and O) into areas by buffering

them. To investigate the sensitivity of buffer choice we considered six buffer sizes: l (i.e., 61.2

m) multiplied by 0.5, 1, 1.5, 2, 3 and 4. We associated all locations falling outside M and O to

the state V. The three-state time series was then converted into three single-state

presence/absence time series.

Preference for feeding sites

We calculated each individual's preference for FS (h_{FS}) as the relative use of FS over

natural vegetation during the pre-closure phase (i.e., the proportion of GPS fixes classified as

either M or O). Because preference is considered to be temporally dynamic (Beyer et al., 2010),

we chose to evaluate h_{FS} for each year separately in case individuals were manipulated in two

separate years. This reasoning allowed for the influence of individual condition and of the

relative quality and quantity of vegetation resources on h_{FS}. We included h_{FS} in all space-use,

movement, and resource use analyses described below.

The variability of h_{FS} across animal-years was maximal when FS attendance was defined

as a GPS location within a 61.2 m distance (i.e., the population mean hourly step length) from

the FS (interquartile range=0.278, mean=0.343; Appendix S3: Table S3.1). Accordingly, the

results described below are based on this definition (see Appendix S7 for sensitivity analysis). At

this scale, h_{FS} did not differ consistently between sex (mean for females=0.346; mean for

males=0.336; t-test: p-value=0.901).

Analysis

We analysed how the experimental manipulation, and its interaction with both preference

for FS and sex, affected roe deer space-use, movement behaviour, and resource use.

Space-use: We assessed the changes of home range and core area sizes (P1.1), and of

space-use overlap (P1.2, P3.1) between experimental phases. We calculated utilization

distributions, (UD; *sensu* Worton 1989) for each animal-year and experimental phase using a

Gaussian kernel density estimation. After visual inspection, we chose to compute the UDs at a

spatial resolution of 10 m and with a fixed bandwidth, set to half the average hourly movement

distance (i.e. $l/2$=30.6 m).

For home range and core area sizes, we calculated the area (in hectares) corresponding to

the 95% and 50% UD contours, respectively, for each experimental phase (*Phase*; three levels;

reference level: *Pre-closure*). We then analysed the log-transformed areas using a linear mixed-

effect model (LMM) with five fixed effects: *Phase*, h_{FS}, *Sex* (categorical predictor; reference

level: *Female*), and two interaction terms (*Phase:h_{FS}* and *Phase:Sex*). We included animal-year

(*ind*) as random effect (intercept). In all analyses, interaction terms were dropped when

statistically non-significant (p-value>0.05).

For space-use patterns, we estimated the overlaps for three pairs of UDs – pre- and post-

closure, pre-closure and closure, and closure and post-closure (*Contrast*; three levels; reference

level: Pre-/Closure) – using the volume of intersection statistic (VI; Fieberg and Kochany 2005).

VI ranges between 0 (no overlap) and 1 (complete overlap). We then analysed the logit-

transformed overlaps using an LMM with *Contrast*, h_{FS}, *Sex*, *Contrast:h_{FS}* and *Contrast:Sex* as

fixed effects, and *ind* as random intercept.

Movement behaviour: We investigated the movement responses of roe deer to the

experiment (P1.3) by analysing the changes in hourly step length (Euclidean distance between

two successive relocations) and turning angle θ_t (angle between two successive movement

steps). Turning angles range between $-\pi$ and π, and were symmetric around 0. We analysed the

log-transformed step length, s_t, and the logit-transformed absolute turning angle, $\varphi_t =$

$log \left(\frac{|\theta_t|}{1-|\theta_t|} \right)$ using LMMs with *Phase*, h_{FS}, *Sex*, *Phase:h_{FS}* and *Phase:Sex* as fixed effects, and

ind as random intercept. Because step length was characterized by strong serial autocorrelation at

short lags and at circadian periodicities (a common pattern in movement trajectories; Fieberg,

Matthiopoulos, Hebblewhite, Boyce, & Frair, 2010), we also included step length measured at

lags 1, 2 and 24 h (i.e., $s_{t-1}, s_{t-2}, s_{t-24}$) as fixed effects to reduce the autocorrelation of the

model residuals.

Resource use: To test whether the experiment led to a transitory change in resource use

(P1.4a-b, P3.2), we fitted separate mixed-effect logistic regression models to the three single-

state presence/absence time series ($u_{M,t}$, $u_{O,t}$ and $u_{V,t}$) using *Phase*, h_{FS}, *Sex*, *Phase:h_{FS}* and

Phase:Sex as fixed effects, and *ind* as random intercept. The pre-closure level for *Phase* was

dropped for u_V to avoid circularity ($h_{FS} = 1 - \bar{u}_{V,t\,Pre-closure}$). We also included the response

variables measured at lags 1, 2 and 24 h (e.g., $u_{M,t-1}, u_{M,t-2}, u_{M,t-24}$) as fixed effects to reduce

the autocorrelation of the model residuals. However, for the sake of conciseness and clarity, we

omit these response lags when visualizing resource use predictions. Because the model results

were consistent regardless of the inclusion of the response lags (Appendix S6: Table S6.2), this

decision had no impact on the interpretation. Two animal-years were excluded from the analyses

of resource use due to the absence of suitable O-state: F4-2017 did not seem to have visited any

other FS (O) prior to the experiment; and F16-2016 had two distinct, highly-used FS during pre-

closure, but only the second most visited FS could be manipulated (due to stakeholder

acceptance). While the use of O was more variable when including these two outliers, the

general patterns remained unchanged (Appendix S6: Table S6.3).

Software: All analyses were conducted in the R environment (R Development Core

Team, 2016). We used the packages adehabitatLT and adehabitatHR (Calenge 2006) for the

spatial analyses, fitted all mixed-effect models via Maximum Likelihood with lme4 (Bates,

Mächler, Bolker, & Walker, 2015), and obtained the coefficients of determination using MuMin

(Bartón, 2018).

Results

Space-use and movement responses to alteration of resource availability

Roe deer space-use changed significantly during the experiment: the size of both home

ranges (95% UD isopleth; Appendix S4: Table S4.1) and core areas (50% isopleth; Table 2)

increased significantly during the experimental closure (Fig. 2; P1.1). On average, home range

size increased from 27.99 ha (σ=11.02) during pre-closure to 34.97 ha (σ=10.17) during closure,

and settled to 29.40 ha (σ=9.27) during post-closure. Core area size followed a similar trend with

averages of 4.23 ha (σ=2.34), 5.85 ha (σ=2.33) and 4.98 ha (σ=2.09), respectively.

Home range and core area sizes were influenced by individual preference for FS (h_{FS})

and there was an interaction between h_{FS} and experimental phase: individuals with a high h_{FS}

had smaller home ranges during the pre-closure and, overall, smaller core areas than those with

lower h_{FS}, but stronger increases following the experimental closure (Fig. 2; Table 2; Appendix

S4: Table S4.1; P2.2). There was no significant effect of sex or interactions between sex and

experimental phase on home range size (Appendix S4: Table S4.2; P2.1), but a marginally

significant interaction between sex and experimental phase on core area size (Appendix S4:

Table S4.3) with responses to closure tending to be slightly larger for males. Overall, the models

quantifying the changes in observed home range and core area sizes accounted for a high

proportion of the total variance (conditional coefficient of determination, cR^2: 0.62 and 0.51,

respectively).

Alongside home range size, the spatial pattern of roe deer home ranges shifted

dramatically following the experimental closure (Fig. 3): the degree of space-use overlap

between pre-closure and closure phases was significantly lower (mean=0.370, CI=0.301-0.405;

P1.2) than the overlap between the temporally-separated pre- and post-closure phases

(mean=0.535, CI=0.475-0.594; P3.1). Space-use overlap was significantly affected by h_{FS} (Fig.

3; Table 3; P2.2), with higher h_{FS} being associated to larger space-use shifts. However, there was

no apparent influence of sex in the space-use patterns (Appendix S4: Table S4.4; P2.1). The

model predicting space-use overlap accounted for an important proportion of the variance

(cR^2=0.50).

Underpinning these changes in home range size and space-use patterns were significant

changes in roe deer movement behaviour during the experiment. Average hourly step length

during the pre-closure phase was 60.32 m (σ=85.79); during closure it increased to 74.26 m

(σ=108.11); and during post-closure it decreased to 68.18 m (σ=96.61, P1.3). In general, males

(Appendix S5: Fig. S5.1, right-hand panels, Table S5.1; P2.1), and individuals associated with

high h_{FS} values (Appendix S5: Fig. S5.1, top panels, Table S5.1; P2.2) were characterized by

stronger increases in step length during the closure phase. In addition, roe deer movements were

more persistent during the closure phase, as shown by a significant decrease in the mean absolute

turning angle for males with a high h_{FS} (Appendix S5: Fig. S5.2, top-right panel, Tables S5.2,

5.3; P1.3, P2.1, P2.2).

Resource use responses to alteration of availability

The spatio-temporal dynamics of resources availability during the experiment led to

important shifts in FS resource use (Fig. 4; Table 4). On average, the proportion of use of the

main FS (M) dropped from 31% during the pre-closure phase to 4% during closure (P1.4a), and

then rebounded to 19% in the post-closure phase (P3.2). This decrease in the use of M during the

closure phase was partially compensated by elevated use of the other FS (O) – which increased

from 3% to 16% following closure (P1.4b), and an increase of the use of vegetation (V) from

66% to 80% following closure. During the post closure, use of O and V declined to 9% and 72%,

respectively. The shifts in resource use were very consistent among animal-years for M and O

but were more variable for V (Fig. 4, top panels).

Roe deer preference for FS significantly influenced how animals used the three resource

types and, in particular, interacted with experimental phase for M and O (Fig. 4, bottom panels;

Table 4). Roe deer characterized by a high h_{FS} had significantly higher use of M during pre-

closure (by definition) and post-closure, as well as consistently lower use of V. High h_{FS} animals

were associated with stronger decreases in use of M and larger increases in the use of O during

closure (P2.2). This compensation for O during closure was stronger for females (Table 4;

Appendix S6: Fig. S6.1; P2.1). However, sex did not influence the use of M or V (Appendix S6:

Table S6.1; P2.1). Overall, the fitted models accounted for a high proportion of the variance in

resource use (cR^2: 0.35, 0.21 and 0.31 for M, O and V, respectively).

Discussion

The results of this field resource manipulation experiment provide direct evidence for the

tight coupling between the spatio-temporal distribution of resources and consequently spatially-

restricted movements of a large herbivore. Specifically, we show that roe deer track resource

dynamics (Fig. 4; H1), which leads to changes in their space-use (Figs. 2, 3) and underpinning

movements (Appendix S5: Figs S5.1, S5.2), and that individual traits, especially resource

preference, mediate these behavioural adjustments (H2). In additional, we show that roe deer

exhibit a high attraction to familiar locations, a process which leads to site fidelity (H3). As far

as we are aware, this is the first experimental demonstration of these interdependencies in a large

mammalian herbivore.

Roe deer alter space-use and movement behaviour to track resources

The experimental alterations of food availability led to larger (Fig. 2), spatially-shifted

home ranges (Fig. 3), and more explorative movements by roe deer (Appendix S5: Figs S5.1,

S5.2), thereby directly establishing the connections between movement, space-use and the

spatio-temporal patterns of resources.

In a previous observational study, Fryxell et al. (2008) showed that elk alternated

between two movement modes: a low speed and high sinuosity mode thought to be within-patch

area-restricted search, and a high speed and low sinuosity mode between resource patches. In our

experimental study, we can directly link these movement modes to changing resources: the

exploratory movements of roe deer (high velocity and low sinuosity) observed during the closure

phase (P1.3) suggested that the animals were motivated to find alternative resource patches when

their familiar feeding site (FS) became inaccessible, thereby increasing (P1.1), but mainly

shifting, their home range (P1.2). While changes in home range size and location following

resource manipulation have been found in studies of lizards (Eifler 1996), birds (van Overveld

and Matthysen 2010) and voles (Ims 1987), to date, there have been few experimental

investigations of the connections between space-use and the spatio-temporal distribution of

resources in large herbivores.

In an earlier study, Kilpatrick and Stober (2002) showed that white-tailed deer shifted

their home range core towards the vicinity of newly deployed FS. Our study builds upon these

results by demonstrating multiple, successive responses to resource manipulation, linking

measured changes in underlying fine-scale movement behaviour of individuals to resulting

patterns of space-use that indicate dynamic resource tracking behaviour by roe deer (H1).

Although roe deer increased their use of the vegetation matrix during the closure phase,

individuals compensated the loss of their main FS (M), to a large degree, by shifts in their

movements and space-use towards alternative FS (O; Figs 4; P1.4a-b). Consequently, individuals

maintained a high overall use of FS throughout the experiment.

While resource tracking behaviour may be expected under the optimal foraging theory,

other individual- and population-based factors, such as social fences due to territoriality or

density-dependent resource competition, can constrain the movement responses of individuals to

changes in the spatial distribution of resources (e.g., Mysterud et al. 2011). The marked

responses shown here are likely to be explained by three factors. First, with the exception of

adult males during spring and summer (Liberg, Johansson, Andersen, & Linnell, 1998), roe deer

do not generally defend territories, and consequently their spatial distribution can approximate

that of an ideal free distribution (Walhström & Kjellander, 1995). Territorial tenure (Linnell &

Andersen, 1998) may explain the marginally different response of males (P2.1), specifically their

tendency to have larger core areas (Appendix S4: Table S4.3), more explorative movements

(Appendix S5: Table S5.1, S5.2) and lower resource compensation than females following

closure (Table 4; Appendix S6: Fig. S6.1). In fact, except for one individual (M11) showing high

spatial instability during its two-year monitoring, the space-use patterns of males tended to be

less influenced by the experimental closure than females.

Second, the rapidity and magnitude of the observed responses are likely linked to the

inability of roe deer to buffer transitory shortages of food availability: as income breeders with

little fat reserves, roe deer need a constant, high-nutritional intake (Andersen et al., 2000). Third,

while intra- and inter-specific competition in herbivores is largely linked to resource depletion

(Owen-Smith et al., 2010), this density-dependent constraint of food availability was prevented

by providing *ad libitum* forage at the FS.

Resource tracking is mediated by individual preference

This study moves beyond the findings of Kilpatrick and Stober (2002) by demonstrating

that inter-individual variation in preference for FS strongly mediated the responses of roe deer

movement patterns, space-use and resource use to changes in the spatio-temporal distribution of

resources (H2). During the closure phase, the changes in all measured variables were of larger

magnitude for individuals associated with a high FS preference (P2.1). The influence of FS

preference was particularly striking in the shifts of space-use (Fig. 3) and in the compensating

use of alternative FS following loss of the familiar resource (Fig. 4).

In this study, FS preference varied across individuals (Appendix S3: Table S3.1). These

inter-individual differences may be linked to either the environment the individuals were

exposed to, or a property of the individuals themselves. In our experimental setting, all roe deer

had access to at least one FS provided with *ad libitum* food where use was not prevented by

inter-individual competition (Ossi et al. 2017; Ossi et al. in prep). Hence, we could quantitatively

approximate FS preference as the relative use of FS over natural vegetation. Preference for FS

was derived from the relative use of FS, and so our definition implies a conditionality on both the

physiological state of the individual and on the relative nutritional value of the provisioned food

(corn) over natural vegetation in a given season. For example, the body condition at the onset of

winter is likely to vary between years. In addition, the dynamics in the quality and quantity of

natural browse – either spatial (e.g., between home ranges) and/or temporal (e.g., between years)

– could lead to variations in FS preference. Indeed, FS preference was higher in 2017 than in

2018 for all roe deer manipulated in these consecutive years (Appendix S3: Table S3.1).

Preference can therefore be considered a dynamic variable (Beyer et al., 2010) that we evaluated

at the individual level over a short period of relative stability (pre-closure phase in each winter).

We considered the temporal extent of our experiment (*ca* 6 weeks) short enough to consider FS

preference for each animal-year to be relatively constant, because in this time period the

physiological conditions and vegetation nutritional value would not vary substantially or

consistently.

Individuals attending a FS benefit from exploiting a forage-rich location, but risk elevated

intra- and inter-specific contacts (Ossi et al. in prep), high anthropogenic disturbance, and an

increased susceptibility to predation by humans. Because FS are associated with such risks,

individuals in poorer condition may be more willing to trade on predation risk to access a forage-

rich patch (McNamara & Houston, 1987). Alternatively, FS preference could be linked to

differences in personality. In roe deer, Bonnot et al. (2015) investigated the association between

individual behavioural profile (e.g., body temperature at capture) and the use of risky but

profitable habitats (open areas), and suggested that variations in personality could lead to

individual differences in habitat use. Similarly, FS preference could be associated to bold or risk-

taking personalities. Interestingly, preference for FS tended to correlate with individual body

temperature at capture (Pearson's $r = - 0.37$ p-value $= 0.084$; Appendix S8), with bolder

individuals (lower temperature) using FS more intensely.

The underlying drivers of preference listed above may not only lead individuals to use

resources to a different extent, but also condition their tendency to track spatio-temporal resource

dynamics (Spiegel, Leu, Bull, & Sih, 2017). For instance, risk-taking personalities tend to be

associated with explorations of novel environments (Cote, Clobert, Brodin, Fogarty, & Sih,

2010), as shown experimentally in great tits following the loss of a familiar foraging area (van

Overveld & Matthysen, 2010).

The role of site familiarity

During the post-closure phase of the experiment, roe deer increased their use of familiar

FS (M), whose food accessibility had been restored after a transitory restriction (Fig. 4, left-hand

panels; P3.1), and home ranges shifted back to pre-closure patterns, as suggested by the high

overlap between temporally-disjointed pre- and post-closure space-use (Fig. 3; P3.2). The

restoration of these pre-manipulation patterns supports the hypothesis that site familiarity

provides inherent benefits to animals maintaining a home range (H3; Piper 2011). These results

are coherent with published literature demonstrating that ungulates tend to select for previously

visited locations i.e., site familiarity (e.g., Dalziel, Morales, & Fryxell, 2008; Merkle, Fortin, &

Morales, 2014).

Unlike most observational studies, our experimental approach allowed us to contrast two

alternative, concentrated resources (M and O) of equal nutritional value but of distinct

familiarity, and hence to separate the effects of resource tracking from those of familiarity.

Resource tracking can explain roe deer use of other FS (O) to compensate the inaccessibility of

the familiar, main FS (M) during closure; however, the systematic return of roe deer to M during

the post-closure phase, while both alternative resources were accessible, cannot be explained by

resource tracking alone. Our experiment suggests that roe deer were actively selecting for

familiar areas and that site familiarity has an inherent value.

In observational studies of animal movement, a spurious familiarity effect (*sensu* Van

Moorter et al. 2013) can occur when an important factor influencing animal behaviour is not

considered, and the re-visitation of particular locations is interpreted as an evidence for site

familiarity selection. However, this confounding effect is unlikely to affect the results of this

experiment. First, corn was delivered *ad libitum* across all FS (M or O) i.e., homogeneous

foraging value. Second, the FS were located in comparable environments with close proximity to

cover, a factor that largely influence roe deer movements and space-use (e.g., Tufto et al. 1996,

Bongi et al. 2008). Third, and most importantly, the specific identities of M and O varied

interchangeably between individuals. Hence, we conclude that the return to pre-closure patterns

of foraging behaviour and space-use are unlikely to be result of variations in the characteristics

of specific FS, but rather of an inherent familiarity effect. In roe deer, site familiarity could allow

a more profitable exploitation of natural forage, as seen in bison (*Bison bison*; Merkle et al.,

2014) or reduce intraspecific competition for such resource (see Riotte-lambert et al. 2015 for a

theoretical argument). Alternatively, the attraction to familiar areas could be related to a predator

(natural or human) avoidance tactic (Gehr et al., in prep; Stamps, 1995).

Our results imply that when resource patterns are changing, individual behavioural

decisions probably reflect a trade-off between the advantages of site familiarity and resource

tracking. In some cases, finding an alternative resource patch of high nutritional value may cause

an individual not to return to familiar sites (e.g., Ims 1987, Eifler 1996, van Overveld and

Matthysen 2010), as seen in this study during the closure phase. The relative influence of

resource tracking and site familiarity likely depends on species traits and ecology. For example,

it is possible that the greater capacity of capital breeding species to buffer temporal dynamics in

resource compared to income breeders such as roe deer alters the trade-off between tracking

resources and site fidelity.

Ultimately, site familiarity is the manifestation of an animal's ability to acquire spatial

information, in particular by means of spatial memory (Fagan et al., 2013). Large herbivores are

capable of memorizing resource location (e.g., Merkle et al., 2014). In this study, it is likely that

the variations in roe deer responses to resource changes that are not explained by preference for

FS may be the result of individual prior experience and knowledge of the status and distribution

of alternative FS. An interesting avenue for further studies will be to evaluate the role of these

cognitive processes on individual foraging decisions.

Acknowledgements

N. Ranc was supported by Harvard University Graduate Fellowship and Fondazione

Edmund Mach International Doctoral Programme Fellowship. F. Cagnacci was supported by the

Sarah and Daniel Hrdy Fellowship 2015-2016 at Harvard University OEB during part of the

development of this ms. We are grateful to all the people involved in the roe deer captures and

data collection, and especially M.B. Almeida, P. Bonanni, A. Corradini, J. De Groove, G. Lilli,

S. Mumme, S. Nicoloso, D. Righetti, M. Rocca, M. Salvatori, M. Sanchez and P. Semenzato. We

thank J.W. Cain, B. Ölveczky, N. Pierce, all the members of the Cagnacci lab and of the

Moorcroft lab, and the graduate students of the Harvard Statistics Consulting Service for their

valuable suggestions. We also thank the Wildlife and Forestry Service of the Autonomous

Province of Trento and the Trentino Hunting Association (ACT) for their support. Finally, we

express our gratitude to the hunters of Albiano, Civezzano, Fornace and Trento Nord without

whose help this study could not be conducted.

Author's contributions

NR, FC and PM conceived the ideas and designed the methodology; FC supervised the

field study; NR, FO, WH and TS collected the data; NR analysed the data with support from WH

and TS; NR led the writing of the manuscript, together with FC and PM. All authors contributed

critically to the drafts and gave final approval for publication.

Data availability statement

Data will be made available from the Dryad Digital Repository upon acceptance.

References

Aarts, G., MacKenzie, M., McConnell, B., Fedak, M., & Matthiopoulos, J. (2008). Estimating

space-use and habitat preference from wildlife telemetry data. *Ecography*, *31*(1), 140–160.

doi: 10.1111/j.2007.0906-7590.05236.x

Andersen, R., Gaillard, J.-M., Linnell, J. D. C., & Duncan, P. (2000). Factors affecting maternal

care in an income breeder, the European roe deer. *Journal of Animal Ecology*, *69*, 672–682.

doi: 10.1046/j.1365-2656.2000.00425.x

Bartón, K. (2018). *MuMIn: Multi-model inference. R Package version 1.42.1. http//CRANR-*

project.org/package=MuMIn.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software*, *67*(1), 1–48. doi: 10.18637/jss.v067.i01

Benhamou, S. (2014). Of scales and stationarity in animal movements. *Ecology Letters*, *17*(3),

261–272. doi: 10.1111/ele.12225

Beyer, H. L., Haydon, D. T., Morales, J. M., Frair, J. L., Hebblewhite, M., Mitchell, M., &

Matthiopoulos, J. (2010). The interpretation of habitat preference metrics under use-

availability designs. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *365*(1550), 2245–2254. doi: 10.1098/rstb.2010.0083

Bongi, P., Ciuti, S., Grignolio, S., Del Frate, M., Simi, S., Gandelli, D., & Apollonio, M. (2008).

Anti-predator behaviour, space use and habitat selection in female roe deer during the

fawning season in a wolf area. *Journal of Zoology*, *276*(3), 242–251. doi: 10.1111/j.1469-

7998.2008.00481.x

Bonnot, N., Verheyden, H., Blanchard, P., Cote, J., Debeffe, L., Cargnelutti, B., … Morellet, N.

518   (2015). Interindividual variability in habitat use: Evidence for a risk management syndrome

519   in roe deer? *Behavioral Ecology*, *26*(1), 105–114. doi: 10.1093/beheco/aru169

520   Brown, J. H., & Munger, J. C. (1985). Experimental manipulation of a desert rodent community:

521   food addition and species removal. *Ecology*, *66*(5), 1545–1563. doi: 10.2307/1938017

522   Cagnacci, F., Boitani, L., Powell, R. A., & Boyce, M. S. (2010). Animal ecology meets GPS-

523   based radiotelemetry: A perfect storm of opportunities and challenges. *Philosophical*

524   *Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2157–2162. doi:

525   10.1098/rstb.2010.0107

526   Calenge, C. (2006). The package "adehabitat" for the R software: A tool for the analysis of space

527   and habitat use by animals. *Ecological Modelling*, *197*, 516–519. doi:

528   10.1016/j.ecolmodel.2006.03.017

529   Cote, J., Clobert, J., Brodin, T., Fogarty, S., & Sih, A. (2010). Personality-dependent dispersal:

530   characterization, ontogeny and consequences for spatially structured populations.

531   *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*, 4065–4076.

532   doi: 10.1098/rstb.2010.0176

533   Couriot, O., Hewison, A. J. M., Saïd, S., Cagnacci, F., & Chamaillé, S. (2018). Truly sedentary ?

534   The multi-range tactic as a response to resource heterogeneity and unpredictability in a

535   large herbivore. *Oecologia*, *187*(1), 47–60. doi: 10.1007/s00442-018-4131-5

536   Dalziel, B. D., Morales, J. M., & Fryxell, J. M. (2008). Fitting probability distributions to animal

537   movement trajectories: using artificial neural networks to link distance, resources, and

538   memory. *The American Naturalist*, *172*(2), 248–258. doi: 10.1086/589448

De Groeve, J., Cagnacci, F., Ranc, N., Bonnot, N. C., Gehr, B., Heurich, M., … Van De Weghe,

N. (2019). Individual Movement - Sequence Analysis Method (IM-SAM): characterizing

spatio-temporal patterns of animal habitat use across landscapes. *International Journal of*

Geographical Information Science, *0*(0), 1–22. doi: 10.1080/13658816.2019.1594822

Eifler, D. A. (1996). Experimental manipulation of spacing patterns in the widely foraging lizard

Cnemidophorus uniparens. *Herpetologica*, *52*(4), 477–486.

Fagan, W. F., Lewis, M. A., Auger-Méthé, M., Avgar, T., Benhamou, S., Breed, G., … Mueller,

T. (2013). Spatial memory and animal movement. *Ecology Letters*, *16*, 1316–1329. doi:

10.1111/ele.12165

Fieberg, J., & Kochany, C. O. (2005). Quantifying home-range overlap: the importance of the

utilization distribution. *The Journal of Wildlife Management*, *69*(4), 1346–1359. doi:

10.2193/0022-541X(2005)69[1346:QHOTIO]2.0.CO;2

Fieberg, J., Matthiopoulos, J., Hebblewhite, M., Boyce, M. S., & Frair, J. L. (2010). Correlation

and studies of habitat selection: problem, red herring or opportunity? *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *365*, 2233–2244. doi:

10.1098/rstb.2010.0079

Fryxell, J. M., Hazell, M., Börger, L., Dalziel, B. D., Haydon, D. T., Morales, J. M., … Rosatte,

R. C. (2008). Multiple movement modes by large herbivores at multiple spatiotemporal

scales. *PNAS*, *105*(49), 19114–19119. doi: 10.1073/pnas.0801737105

Holbrook, K. M., Smith, T. B., & Hardesty, B. D. (2002). Implications of long-distance

movements of frugivorous rain forest hornbills. *Ecography*, *25*(6), 745–749. doi:

10.1034/j.1600-0587.2002.250610.x

Ims, R. A. (1987). Responses in spatial organization and behaviour to manipulations of the food

resouce in the vole Clethrionomys rufocanus. *Journal of Animal Ecology*, *56*, 585–596. doi:

10.2307/5070

Johnson, D. H. (1980). The comparison of usage and availability measurements for evaluating

resource preference. *Ecology*, *61*(1), 65–71. doi: 10.2307/1937156

Kilpatrick, H. J., & Stober, W. A. (2002). Effects of temporary bait sites on movements of

suburban white-tailed deer. *Wildlife Society Bulletin*, *30*(3), 760–766.

Krebs, J. R. (1971). Territory and breeding density in the great tit, Parus Major L. *Ecology*,

52(1), 2–22. doi: 10.2307/1934734

Liberg, O., Johansson, A., Andersen, R., & Linnell, J. D. C. (1998). Mating system, mating

tactics and the function of male territory in roe deer. In R. Andersen, P. Duncan, & J. D. C.

Linnell (Eds.), *The European roe deer: the biology of success* (pp. 221–256). Oslo, Norway:

Scandinavian University Press.

Linnell, J. D. C., & Andersen, R. (1998). Territorial fidelity and tenure in roe deer bucks. *Acta*

Theriologica, *43*(1), 67–75. doi: 10.4098/AT.arch.98-5

McNamara, J. M., & Houston, A. I. (1987). Starvation and predation as factors limiting

population size. *Ecology*, *68*(5), 1515–1519. doi: 10.2307/1939235

Merkle, J. A., Fortin, D., & Morales, J. M. (2014). A memory-based foraging tactic reveals an

adaptive mechanism for restricted space use. *Ecology Letters*, *17*, 924–931. doi:

10.1111/ele.12294

Mitchell, M. S., & Powell, R. A. (2004). A mechanistic home range model for optimal use of

spatially distributed resources. *Ecological Modelling*, *177*(1–2), 209–232. doi:

10.1016/j.ecolmodel.2004.01.015

Morellet, N., Bonenfant, C., Börger, L., Ossi, F., Cagnacci, F., Heurich, M., … Mysterud, A.

(2013). Seasonality, weather and climate affect home range size in roe deer across a wide

latitudinal gradient within Europe. *Journal of Animal Ecology*, *82*(6), 1326–1339. doi:

10.1111/1365-2656.12105

Mueller, T., & Fagan, W. F. (2008). Search and navigation in dynamic environments behaviors

to population distributions. *Oikos*, *117*, 654–664. doi: 10.1111/j.0030-1299.2008.16291.x

Mueller, T., Olson, K. A., Dressler, G., Leimgruber, P., Fuller, T. K., Nicolson, C., … Fagan, W.

F. (2011). How landscape dynamics link individual- to population-level movement patterns:

A multispecies comparison of ungulate relocation data. *Global Ecology and Biogeography*,

20(5), 683–694. doi: 10.1111/j.1466-8238.2010.00638.x

Mysterud, A., Loe, L. E., Zimmermann, B., Bischof, R., Veiberg, V., & Meisingset, E. (2011).

Partial migration in expanding red deer populations at northern latitudes – a role for density

dependence? *Oikos*, *120*, 1817–1825. doi: 10.1111/j.1600-0706.2011.19439.x

Normand, E., Ban, S. D., & Boesch, C. (2009). Forest chimpanzees (Pan troglodytes verus)

remember the location of numerous fruit trees. *Animal Cognition*, *12*(6), 797–807. doi:

10.1007/s10071-009-0239-7

Ossi, F., Gaillard, J. M., Hebblewhite, M., & Cagnacci, F. (2015). Snow sinking depth and forest

canopy drive winter resource selection more than supplemental feeding in an alpine

population of roe deer. *European Journal of Wildlife Research*, *61*(1), 111–124. doi:

10.1007/s10344-014-0879-z

Ossi, F., Gaillard, J. M., Hebblewhite, M., Morellet, N., Ranc, N., Sandfort, R., … Cagnacci, F.

(2017). Plastic response by a small cervid to supplemental feeding in winter across a wide

environmental gradient. *Ecosphere*, *8*(1). doi: 10.1002/ecs2.1629

Owen-Smith, N., Fryxell, J. M., & Merrill, E. H. (2010). Foraging theory upscaled: The

608 behavioural ecology of herbivore movement. *Philosophical Transactions of the Royal*

609 *Society B: Biological Sciences*, *365*(1550), 2267–2278. doi: 10.1098/rstb.2010.0095

610 Owen-Smith, Norman. (2004). Functional heterogeneity in resources within landscapes and

611 herbivore population dynamics. *Landscape Ecology*, *19*, 761–771. doi: 10.1007/s10980-

612 005-0247-2

613 Peters, W., Hebblewhite, M., Mysterud, A., Spitz, D., Focardi, S., Urbano, F., … Cagnacci, F.

614 (2017). Migration in geographic and ecological space by a large herbivore. *Ecological*

615 *Monographs*, *87*(2), 297–320. doi: 10.1002/ecm.1250

616 Piper, W. H. (2011). Making habitat selection more "familiar": a review. *Behavioral Ecology*

617 *and Sociobiology*, *65*(7), 1329–1351. doi: 10.1007/s00265-011-1195-1

618 R Development Core Team. (2016). *R: a language and environment for statistical computing. R*

619 *Foundation for Statistical Computing, Vienna, Austria. www.r-project.org.*

620 Riotte-lambert, L., Benhamou, S., & Chamaillé-Jammes, S. (2015). How memory-based

621 movement leads to nonterritorial spatial segregation. *The American Naturalist*, *185*(4),

622 E103-116. doi: 10.1086/680009

623 Rozen-Rechels, D., van Beest, F. M., Richard, E., Uzal, A., Medill, S. A., & Mcloughlin, P. D.

624 (2015). Density-dependent, central-place foraging in a grazing herbivore: Competition and

625 tradeoffs in time allocation near water. *Oikos*, *124*(9), 1142–1150. doi: 10.1111/oik.02207

626 Spiegel, O., Leu, S. T., Bull, C. M., & Sih, A. (2017). What's your move? Movement as a link

627 between personality and spatial dynamics in animal populations. *Ecology Letters*, *20*(1), 3–

628 18. doi: 10.1111/ele.12708

629 Stamps, J. (1995). Motor learning and the value of familiar space. *The American Naturalist*,

630 *146*(1), 41–58. doi: 10.1086/285786

Stephens, D. W., & Krebs, J. R. (1986). *Foraging theory*. Princeton, New Jersey, USA:

Princeton University Press.

Tufto, J., Andersen, R., & Linnell, J. (1996). Habitat use and ecological correlates of home range

size in a small cervid: the roe deer. *Journal of Animal Ecology*, *65*(6), 715–724. doi:

10.2307/5670

Van Moorter, B., Rolandsen, C. M., Basille, M., & Gaillard, J. M. (2016). Movement is the glue

connecting home ranges and habitat selection. *Journal of Animal Ecology*, *85*(1), 21–31.

doi: 10.1111/1365-2656.12394

Van Moorter, B., Visscher, D., Benhamou, S., Börger, L., Boyce, M. S., & Gaillard, J.-M.

(2009). Memory keeps you at home: a mechanistic model for home range emergence.

Oikos, *118*(5), 641–652. doi: 10.1111/j.1600-0706.2008.17003.x

Van Moorter, B., Visscher, D., Herfindal, I., Basille, M., & Mysterud, A. (2013). Inferring

behavioural mechanisms in habitat selection studies getting the null-hypothesis right for

functional and familiarity responses. *Ecography*, *36*(3), 323–330. doi: 10.1111/j.1600-

0587.2012.07291.x

van Overveld, T., & Matthysen, E. (2010). Personality predicts spatial responses to food

manipulations in free-ranging great tits (Parus major). *Biology Letters*, *6*, 187–190. doi:

10.1098/rsbl.2009.0764

Walhström, L. K., & Kjellander, P. (1995). Ideal free distribution and natal dispersal in female

roe deer. *Oecologia*, *103*(3), 302–308. doi: 10.1007/BF00328618

Wiens, J. A. (1976). Population responses to patchy environments. *Annual Review of Ecology*

and Systematics, *7*(1976), 81–120. doi: 10.1146/annurev.es.07.110176.000501

Worton, B. J. (1989). Kernel methods for estimating the utilization cistribution in home-range

studies. *Ecology, 70*(1), 164–168. doi: 10.2307/1938423

Tables

Table 1. Hypotheses and corresponding predictions.

Hypotheses	Predictions
H1: roe deer alter their space-use patterns and movement behaviour to track dynamics in resource availability.	*The closure of the familiar feeding site (FS) leads to…* - P1.1: an increase in HR size - P1.2: a spatial shift in space-use - P1.3: more explorative movements - P1.4a: a reduced use of the familiar FS - P1.4b: a compensation by using other, alternative FS
H2: these behavioural adjustments are mediated by sex and FS preference.	- P2.1: males respond less markedly to the experimental transitions - P2.2: individuals with a high preference for FS respond more markedly to the experimental transitions
H3: roe deer strive to use familiar resources when accessible.	*The re-opening of the familiar FS leads to…* - P3.1: the return of space-use to pre-closure patterns - P3.2: the return of resource use to pre-closure patterns

659   Table 2. Summary of the final model for core area size (50% UD).

	Estimate	Std. Error	df	t value	p-value
(Intercept)	1.930	0.181	65.921	10.684	<0.001***
PhaseClosure	-0.106	0.223	48.722	-0.478	0.635
PhasePost-closure	-0.101	0.224	48.970	-0.450	0.655
h_{FS}	-2.096	0.464	65.921	-4.514	<0.001***
PhaseClosure:h_{FS}	1.572	0.572	48.722	2.747	0.008**
PhasePost-closure:h_{FS}	1.179	0.591	49.914	1.997	0.051(*)

	Std. Dev		R^2	
Random effect	0.209	Marginal	0.351
Residual	0.373	Conditional	0.506

661   Table 3. Summary of the final model for space-use overlap.

	Estimate	Std. Error	df	t value	p-value
(Intercept)	0.096	0.225	66.109	0.428	0.670
ContrastClosure/Post-c.	0.119	0.286	47.785	0.414	0.680
ContrastPre-c./Post-c.	-0.045	0.286	47.785	-0.156	0.877
h_{FS}	-2.073	0.578	66.109	-3.590	0.001**
ContrastClosure/Post-c.:h_{FS}	1.179	0.753	49.192	1.565	0.124
ContrastPre-c./Post-c.:h_{FS}	2.366	0.753	49.192	3.141	0.003**

	Std. Dev			R^2	
Random effect	0.238		Marginal	0.372	
Residual	0.476		Conditional	0.498	

664   Table 4. Summary of the final models for the use of the main feeding site ($u_{M,t}$), other feeding

665   sites ($u_{O,t}$) and vegetation ($u_{V,t}$).

	Main feeding site (M)			
	Estimate	Std. Error	z value	p-value
(Intercept)	-3.325	0.112	-29.804	<0.001***
PhaseClosure	-0.645	0.172	-3.740	<0.001***
PhasePost-closure	-0.102	0.126	-0.809	0.418
h_{FS}	1.723	0.288	5.990	<0.001***
PhaseClosure:h_{FS}	-1.657	0.459	-3.606	<0.001***
PhasePost-closure:h_{FS}	-0.473	0.325	-1.456	0.145
$u_{M,t-1}$	3.155	0.065	48.555	<0.001***
$u_{M,t-2}$	0.900	0.068	13.286	<0.001***
$u_{M,t-24}$	0.739	0.062	11.917	<0.001***
	Std. Dev			R^2
Random effect	0.161		Marginal	0.346
Residual	1.000		Conditional	0.349

	Other feeding sites (O)			
	Estimate	Std. Error	z value	p-value
(Intercept)	-6.032	0.379	-15.913	<0.001***
PhaseClosure	2.191	0.297	7.38	<0.001***
PhasePost-closure	1.663	0.308	5.395	<0.001***
h_{FS}	3.869	0.829	4.668	<0.001***

	Estimate	Std. Error	z value	p-value
Sex	-0.917	0.372	-2.464	0.014*
PhaseClosure:h_{FS}	-1.726	0.571	-3.022	0.003**
PhasePost-closure:h_{FS}	-1.831	0.594	-3.081	0.002**
PhaseClosure:SexM	0.529	0.292	1.815	0.069(*)
PhasePost-closure:SexM	0.855	0.302	2.835	0.005**
$u_{O,t-1}$	2.993	0.081	36.935	<0.001***
$u_{O,t-2}$	1.175	0.086	13.611	<0.001***
$u_{O,t-24}$	0.394	0.087	4.521	<0.001***

	Std. Dev			R^2
Random effect	0.551		Marginal	0.188
Residual	1.000		Conditional	0.208

Vegetation (V)

	Estimate	Std. Error	z value	p-value
(Intercept)	-0.496	0.187	-2.655	0.008**
PhasePost-closure	-0.252	0.055	-4.593	<0.001***
h_{FS}	-1.870	0.443	-4.223	<0.001***
$u_{V,t-1}$	2.597	0.061	42.240	<0.001***
$u_{V,t-2}$	0.853	0.064	13.327	<0.001***
$u_{V,t-24}$	0.382	0.061	6.256	<0.001***

	Std. Dev			R^2
Random effect	0.361		Marginal	0.298
Residual	1.000		Conditional	0.314

Figure 1. Schematic representation of the experiment. (a) The manipulation consists of a transitory

alteration of resource accessibility at a main (M) feeding site. (b) In response to the experiment,

roe deer are expected to shift from M (green/red dot, change of colour denoting the alteration of

accessibility) towards alternative resources – other feeding sites (O; black triangles) or the

vegetation (V; underlying matrix). In particular, this can lead to spatio-temporal dynamics in space-

use (utilization distribution: colour gradient; 95% and 50% contour lines: thick and thin white lines,

respectively; data from roe deer F5-2018).

Figure 2. Roe deer home range (y-axis, left panel) and core area (y-axis, right panel) sizes as a

function of preference for feeding sites (x-axis) and experimental phase. Observations are

represented as dots (females) and squares (males), (points are jittered slightly), and model

predictions as solid lines (95% confidence intervals: ribbons).

Figure 3. Roe deer space-use overlap (y-axis) as a function of preference for feeding sites (x-axis)

and experimental phase contrast. Observations are represented as dots (females) and squares

(males) and model predictions as solid lines (95% confidence intervals: ribbons).

Figure 4. Roe deer shifts in resource use during the experiment – main feeding site (M, left), other

feeding sites (O, centre) and vegetation (V, right). Top panel: mean proportional use (dots and

lines) as a function of the experiment phase (x-axis) and preference for feeding sites (colour scale).

Bottom panel: model predictions for $u_{M,t}$, $u_{O,t}$ and $u_{V,t}$ (estimate: solid lines; 95% confidence

interval: ribbon) and mean relative use (females: dots; males: squares) as a function of the

experiment phase (colour) and preference for feeding sites (x-axis). The model predictions do not

consider resource lags at 1, 2 and 24h nor the influence of *Sex* (although selected in the final model

for $u_{O,t}$).