Causes and consequences of facultative sea crossing in a soaring migrant

Paolo Becciu[1], Shay Rotics[2,†], Nir Horvitz[2,‡], Michael Kaatz[3], Wolfgang Fiedler[4,5], Damaris Zurell[6,7], Andrea Flack[4,5], Florian Jeltsch[8,9], Martin Wikelski[4,5], Ran Nathan[2] and Nir Sapir[1]

[1] Animal Flight Laboratory, Department of Evolutionary and Environmental Biology, and Institute of Evolution, University of Haifa, 199 Aba Khoushy Ave. Mount Carmel, 3498838 Haifa, Israel;

[2] Movement Ecology Laboratory, Department of Ecology, Evolution and Behavior, Alexander Silberman Institute of Life Sciences, The Hebrew University of Jerusalem, 91904 Jerusalem, Israel;

[3] Vogelschutzwarte Storchenhof Loburg e.V., Chausseestr. 18, D-39279 Loburg, Germany;

[4] Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany;

[5] Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany;

[6] Geography Department, Humboldt-Universität Berlin, D-10099 Berlin, Germany;

[7] Land Change Science, Swiss Federal Research Institute WSl, CH-8903 Birmensdorf, Switzerland;

[8] Plant Ecology and Conservation Biology, Institute for Biochemistry and Biology, University of Potsdam, Am Mühlenberg 3, 14476 Potsdam, Germany;

[9] Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), D-14195 Berlin, Germany;

[†] Present address: Department of Zoology, University of Cambridge, UK;

[‡] Present address: Department of Environmental Science, Policy, and Management, University of California, Berkeley, CA 94720, USA.

Correspondence author: Paolo Becciu, 199 Aba Khoushy Ave. Mount Carmel, 3498838 Haifa, Israel, e-mail: pbecciu89@gmail.com

Abstract

1. Studying the causes and consequences of route selection in animal migration is important for understanding the evolution of migratory systems and how they may be affected by environmental factors at various spatial and temporal scales. One key decision during migration is whether to cross "high transport cost" areas, or to circumvent them. Soaring birds may face this choice when encountering waterbodies where convective updrafts are weak or scarce. Thus, crossing waterbodies requires flying using metabolically costly flapping flight, while circumnavigating them over land permits energetically cheap soaring.

2. We tested how several atmospheric factors (e.g., wind, thermal uplift) and geographic, seasonal and state-related factors (sex and age) affected route selection in migrating white storks (*Ciconia ciconia*). We used 196 GPS tracks of 70 individuals either crossing or circumventing the north-easternmost section of the Mediterranean Sea, over Iskenderun Bay in southern Turkey.

3. We found that westward and southward winds promoted a cross-bay journey in spring and autumn, respectively, acting as tailwinds. Also, overall weaker winds promoted a sea crossing in spring. Sea crossing was associated with flapping flight and higher values of Overall Dynamic Body Acceleration (ODBA) and resulted in higher groundspeed than travel over land.

4. Notably, daily travel distances of spring migrants crossing the bay when returning to their nesting locations were 60 kilometres and 2 hours longer than those of land-detouring birds. No such benefit was found during autumn.

5. Our findings confirm that atmospheric conditions can strongly affect bird route selection. Consequently, migration timing, speed and movement-related energy expenditure differed considerably between the two migratory seasons and the two route choices, highlighting a time-energy trade-off in the migration of white storks.

Key words: environmental effects, flight modes, geographical barriers, land detour, route selection, season-related behavioural responses, time-energy trade-offs, white stork

1. Introduction

Environmental conditions during long-distance bird migration are known to affect migration timing, flight performance and energy expenditure (Becciu et al., 2019; Shamoun-Baranes, Liechti, & Vansteelant, 2017). Still, how migration route is influenced by those is much less clear. Route selection over ecological barriers such as large waterbodies and mountainous reliefs may depend on weather and geographical features (Alerstam, 2001; Becciu et al., 2019; Eisaguirre et al., 2018; Nourani, Yamaguchi, Manda, & Higuchi, 2016), affecting migration time and energy expenditures, with consequences for animal fitness (Shamoun-Baranes, Bouten, & Van Loon, 2010; Shamoun-Baranes et al., 2017). Large terrestrial soaring birds depend on local atmospheric conditions during their flight, since they utilize thermal uplifts to gain height and later glide towards their destination (Norberg, 1990). During soaring flight, the birds stretch and do not flap their wings, allowing them to save energy while covering large distances. Yet, in some cases, such as low availability of thermals, soaring birds are forced to switch to the metabolically demanding flapping flight (Hedenstrom, 1993; Norberg, 1990; Sapir et al., 2011). Usually, soaring birds avoid flying over waterbodies where thermals are typically weaker (but see Duriez, Peron, Gremillet, Sforzi, & Monti, 2018). Flying across a waterbody may increase the soaring birds' transport cost due the use of flapping flight, characterizing such zones as "high transport cost" areas (Alerstam, 2001). We note that differences in transport cost may be caused by additional factors, such as variable wind conditions experienced by the birds when travelling over different areas (Alerstam, 2001). Besides increasing the transport cost, barrier crossing versus barrier circumvention (i.e., facultative barrier crossing) may shorten migration distance, with possible consequences for migration time saving (Alerstam, 2001).

In cases of obligatory sea crossing during migration, weather conditions may affect the timing and the location of the crossing (Agostini, Panuccio, & Pasquaretta, 2015; Bildstein, 2006; Bildstein, Bechard, Farmer, & Newcomb, 2009; Meyer, Spaar, & Bruderer, 2000; Nourani et al., 2016). Several studies on Oriental honey-buzzards (*Pernis ptilorhynchus*) found that seasonal wind conditions and the geography of the study area (sea crossing between the mainland and Japan) affect the properties of bird sea crossing. In fact, honey-buzzards adjusted their path before crossing the sea to exploit favourable air currents at different locations on the coast and then over sea (Nourani et al., 2016; Yamaguchi, Arisawa, Shimada, & Higuchi, 2012). Furthermore, a multi-species radar study suggested that wind conditions affect the decision of soaring migrants that are about to cross the Mediterranean sea between Spain and Morocco, to leave the coast before reaching the point where they could find the shortest distance to travel over the strait of Gibraltar (Meyer et al., 2000). Compared to the

latter situations of obligatory sea crossing, causes and consequences of a facultative cross-sea

decision in soaring migrants were rarely studied to date (Kerlinger, 1984).

We investigated the flight behaviour of the white stork, *Ciconia ciconia*, a large soaring bird

and a long-distant migrant, as it passes through the Iskenderun Bay ("the bay" hereafter) in

the north-eastern corner of the Mediterranean Sea. We found that about half of the birds

crossed the bay over water while the other storks chose to circumvent it. We examined how

meteorological conditions affected migration route selection (bay crossing vs. overland

detour) and furthermore explored the consequences of route selection for migration distance

and movement-related energetics due to changes in the prevalence of the two flight modes

(soaring vs. flapping) used by the birds. Large differences in flight metabolism between the

two flight modes (Hedenstrom, 1993; Sapir, Wikelski, Mccue, Pinshow, & Nathan, 2010)

imply a possible trade-off between different benefits and costs of facultative sea-crossing

behaviour. For example, over-sea shortcutting may involve high prevalence of metabolically

expensive flapping flight whereas the longer overland detour might be associated with

energetically cheap soaring flight. We consequently hypothesize that cross-sea flight is

selected only when specific meteorological conditions prevail, such as tailwinds, which

increase the benefit-(shorter travel time)-to-cost (high metabolism due to flapping) ratio of

crossing the bay. We therefore tested how wind speed and direction, temperature and thermal

availability affected the decision of the storks to cross the bay. We furthermore considered the

sex and the age of the individuals because intrinsic individual attributes may play an

important role in determining movement decisions (Nathan et al., 2008). Specifically in white

storks, males were found to arrive earlier than females on the nesting grounds (Rotics et al.,

2018b) and juveniles were found to be slower and to have higher energy costs than adults

while migrating (Rotics et al., 2016b). Also, we considered the timing of bird passage through

the study area within the season. Time and energy consequences of route selection were

additionally explored. We hypothesize that sea-crossing behaviour is not random and depends

on both extrinsic and intrinsic factors that could affect individual fitness. We specifically

predict that (a) tailwinds will facilitate sea-crossing flight, and increase the speed of migration

(Becciu, Panuccio, Catoni, Dell'Omo, & Sapir, 2018; Nourani et al., 2016); (b) late-spring

migrants will show higher sea-crossing prevalence due to time pressure to arrive earlier at

their breeding grounds (Rotics et al., 2018b); (c) sea-crossing flight will be beneficial to the

migrants, shortening their migration distance and time compared with land detour, and

consequently allowing them to cover more distance after the barrier; (d) sea-crossing flight

will require flapping as opposed to soaring during land detour and consequently will be

metabolically more costly (Hedenstrom, 1993; Sapir et al., 2010; Wilson et al., 2019).

Therefore, we suggest that facultative sea-crossing behaviour could be the outcome of a time-

energy trade-off during white stork migration, in which the birds may trade off energy

expenditure for migration speed.

2. Materials and Methods

2.1 Bird tagging and study area

The white stork is a large long-distance migrant that breeds mainly in Europe and Western

Asia, and the majority of its population over-winters in sub-Saharan Africa. The study took

place at the area of Iskenderun Bay, Turkey (36.6330 N, 35.8786 E). White storks that

migrate along the eastern Mediterranean flyway pass regularly over the study area twice a

year. When encountering the bay, storks may choose to cross the bay, which is 30-45 km

wide, or to circumvent it over land (Figure 1). From 2011 to 2014, we fitted solar-charged

GPS transmitters with tri-axial acceleration (ACC) sensors (e-obs GmbH, Munich, Germany)

to 62 adult and 84 immature white storks in the state of Saxony-Anhalt, Germany (see Rotics

et al., 2016, 2017). Immature storks are birds in their first, second and third year of life. A

total of eight immature individuals survived to the following years. Since the behaviour of 1^{st}-

year birds was similar to that of 2^{nd}- and 3^{rd}-year birds, in terms of sea crossing choice and

day of passage over the study area (Figure S2), we considered them in the same age class

(juvenile) in the statistical analysis. Bird sex was determined by molecular methods (Rotics et

al., 2018b). The transmitters recorded GPS fixes every 5 minutes when solar conditions were

good (95% of the time) or otherwise every 20 minutes. Every five minutes an ACC burst of

3.8 seconds was recorded at 10.54 Hz for each of three perpendicular axes. Data were stored

on-board and were downloaded via a VHF radio link upon locating the stork (Rotics et al.,

2016b). We excluded from the analysis tracks that did not present a clear route choice, those

birds wintering at higher latitudes in the northern hemisphere (Rotics et al., 2017) and those

that did not cross the bay in one day (e.g. stopping over at the area of the bay). Consequently,

we used data from 70 storks (39 adults and 31 immatures) that provided a total of 196 tracks

(153 from adult and 43 from immature storks, 83 for spring and 113 for autumn migration).

Figure 1. Map of the study area showing the white storks' routes in (a) spring and (b) autumn. Blue routes depict
sea crossing and red ones represent land detour. The shaded area is the bay-crossing stage named "ACROSS",
which is considered in the analysis of land-detouring birds (see Methods for details). The topography is depicted
by a colour gradient from sea level (dark green) to mountains of about 3000 m a.s.l. (dark brown).

2.2 Environmental data annotation

Each GPS fix was annotated with environmental data using the Env-DATA track annotation
tool of MoveBank (Dodge et al., 2013). These data included (1) E-W wind (also known as the
U wind component, with positive values indicating eastward wind), (2) N-S wind (the V
component of the wind, with positive values indicating northward wind) that were
downloaded for both 10 m above the surface and at the height of the storks' flight with
temporal resolution of 3 hours, (3) thermal uplift velocity (6 hours temporal resolution), and
(4) surface air temperature (6 hours temporal resolution) from the European Centre for
Medium-Range Weather Forecasts (ECMWF) re-analysis data archive at 0.75 degrees of
spatial resolution. From the U and V components of the wind we calculated wind speed (W_s)
and direction (blowing towards θ) as follows: $W_s = \sqrt{U^2 + V^2}$, and $\theta = \tan^{-1}(U, V)$. For
each track segment between two positions we additionally calculated wind support (W_p),
which represents the component of the wind vector towards the direction of the bird (positive
values) or against it (negative values), by applying the following formula: $W_p = W_s \times \cos(\theta - \sigma_i)$, where σ is the angle of each segment i per individual (see below). Also, we
calculated the crosswind W_c, which is the lateral component of the wind relative to the birds'
tracks for each segment: $W_c = W_s \times \sin(\theta - \sigma_i)$. We used the U and V wind components at

10 meters above the surface because they were highly correlated with those estimated at the

storks' heights (Spearman-$\rho > 0.9$, $p < 0.001$).

2.3 Movement parameters

We calculated ground speed (V_g) based on the time interval between two consecutive

locations and additionally calculated the angle (σ_i) of each such segment relative to the

previous segment. These parameters were calculated using the package "move" in R

(Kranstauber, Smolla, & Scharf, 2018). Ground speed was consequently averaged for the

entire day during which the bay crossing took place. For every ACC burst we calculated the

birds' Overall Dynamic Body Acceleration (ODBA), a valid proxy for activity-related energy

expenditure (Wilson et al., 2019), and their flight mode (either flapping or soaring-gliding

flight; see (Rotics et al., 2018b) for details). ODBA and the proportion of flapping flight are

highly correlated (Spearman-$\rho = 0.92$, p < 0.001). Flight height above ground was calculated

by subtracting ground elevation (obtained from ASTER DEM data set (Dodge et al., 2013))

and geoid height (the elevation difference between ellipsoid and geoid earth models) from the

ellipsoid height recorded by the transmitter. Air speed (V_a) was calculated for each segment of

the individual tracks following Safi et al. (2013): $V_a = \sqrt{(V_g - W_p)^2 + (W_c)^2}$.

2.4 Data analysis

We divided our dataset into different subsets depending on the position of the birds with

respect to the bay on the day of crossing the study area: 1) BEFORE (from take-off to the

"bay area" – see below), 2) ACROSS (over the "bay area" or its projection over land), 3)

AFTER (from the "bay area" until landing). The "bay area" is considered as the water body

itself plus its projection over land in a direction perpendicular to the GPS tracks (shaded area

in Figure 1, see also Figure S1). We used averaged movement and environmental data per day

and per each bay-crossing stage (depending on the analysis) to avoid spatial and temporal

correlation on the day when the storks passed over the study area. We assigned bird tracks to

two categories, namely LAND and SEA, for land-detour and sea-crossing routes,

respectively.

We tested bird route choice before arriving at the bay (using a minimum average of 3

locations per individual in the BEFORE stage) in relation to environmental factors by using

Generalized Linear Mixed Models (hereafter GLMMs) with a binomial response variable

(route choice: 0=LAND; 1=SEA) that is similar to logistic regression but allows the inclusion

of random variables (calendar year and bird ID, separately for autumn and spring migration).

To avoid multicollinearity issues, we chose the most biologically meaningful variable from

pairs of variables with a Spearman rank correlation | ρ | > 0.6. This ensured that all the

predictors in the GLMMs had a Variance Inflation Factor (VIF) < 3 (Zuur, Ieno, & Elphick, 2010). We then tested all combinations of remaining variables in the global model and ranked the selected models according to the Akaike information criterion (Burnham & Anderson, 2002) using an automated stepwise model selection where models are fitted through repeated evaluation of modified calls extracted from the model containing all the meaningful variables, corrected for small sample sizes (AIC_c) (Sugiura, 1978). Furthermore, we averaged all models with $\Delta AIC_c < 7$ (Burnham, Anderson, & Huyvaert, 2011) and used the Akaike weights (w_i) (Anderson, Burnham, & Thompson, 2000; Anderson, Link, Johnson, & Burnham, 2001) to assess the relative importance of the different variables. We used two global models, the first including E-W and N-S winds (but not W_p and W_c), and the second with W_p and W_c (without E-W and N-S winds), and then used the one with lowest AIC_c among them. We used 10-fold cross-validation with 10 repetitions, where the best model was trained on 70% of the data and then applied to the remaining 30% of the data. These subsets of data were chosen randomly for each repetition (Hastie, Tibshirani, & Friedman, 2009; Meijer & Goeman, 2013). From the repeated cross-validation we reported the ability of the best model to distinguish between land/sea-crossing decisions using the area under the curve (AUC) of the receiver-operating characteristic curve (with standard deviation), the logistic regression accuracy (defined as the ratio between the sum of correct predicted cases of sea crossing and land detour and the sum of correct and non-correct predicted cases), sensitivity (proportion of land-detour choices correctly classified) and specificity (proportion of sea-crossing choices correctly classified) (Fawcett, 2006). Individual consistency in route choice was examined by calculating repeatability across years (Intraclass correlation) using the rptR package (Stoffel, Nakagawa, & Schielzeth, 2017). We used linear mixed models (LMMs) to test the effects of route choice and environmental factors on the daily ground and air speeds. We found the optimal structure of the fixed component as described above for GLMMs, using AIC_c in a multi-model selection framework. Also, we inspected GLMMs and LMMs residuals and considered the dispersion of the data (Zuur, 2009) using a simulation-based approach to create readily interpretable scaled (quantile) residuals for fitted (generalized) LMMs with the package DHARMa (Hartig, 2019). Additionally, we used LMMs to examine whether route selection affected the time spent flying, distance covered, ground and air speeds, ODBA and daily proportion of flapping flight in all the segments of the tracks during the day of crossing (BEFORE, ACROSS and AFTER the bay), and also using the mean of the entire day. We report differences between route choices among the path segments using the *lsmeans()* R function of package lsmeans (Lenth, 2016). Model fitting and multi-model inference were carried out in the statistical environment R 3.5.1 (R Core Team, 2018) by the packages lme4 (Bates, Mächler, Bolker, & Walker, 2015) and MuMIn (Barton, 2019), while the cross-validation was done using the package caret (Kuhn, 2019).

3. Results

3.1 Route selection

Migrating white storks crossed the Iskenderun Bay more often in spring (61.5%) than in autumn (39.8%). During the spring seasons of 2011 to 2015 storks crossed the Iskenderun Bay 51 times and detoured it 32 times. Adults preferred crossing the bay rather than detouring it (N_{LAND} = 24, N_{SEA} = 50; χ^2 = 9.13, p < 0.01), while an opposite trend was found in juveniles (N_{LAND} = 8, N_{SEA} = 1). Juveniles travelled mostly over land also in autumn, (N_{LAND} = 25, N_{SEA} = 9; χ^2 = 7.53, p < 0.01), whereas adults did not show any route selection preference in this season (N_{LAND} = 43, N_{SEA} = 36; χ^2 = 0.62, p = 0.43).

In spring, wind speed, E-W wind speed and the ordinal date were ranked as the most important variables influencing route selection such that sea crossing was facilitated by decreasing wind speed, increasing westward wind speed and earlier passage date (Figure 2a). The average (± SD) logistic regression accuracy of the best ranked model following the testing of the data subsets was 0.86 (± 0.13), with sensitivity$_{LAND}$ = 0.76 (± 0.30) and specificity$_{SEA}$ = 0.92 (± 0.14). The average (± SD) AUC was 0.96 (± 0.09). Route choice of individual birds was not consistent among years (repeatability: r = 0.05 ± 0.09, p = 0.34).

In autumn, N-S wind and ordinal date were the most influential factors affecting route selection (Figure 2b). The probability of sea crossing increased with southward wind speed and when passing over the area relatively late in the season. The average (± SD) logistic regression accuracy of the best ranked model was 0.74 (± 0.14), with sensitivity$_{LAND}$ = 0.84 (± 0.17) and specificity$_{SEA}$ = 0.60 (± 0.27). The average AUC was 0.85 (± 0.15). Also in autumn, route choice of individual birds was not consistent among years (repeatability: r = 0.0001 ± 0.07, p = 0.626). Tables with model and variable rankings as well as the selected models are reported in the electronic supplementary material. In both seasons the best models with the lowest AIC_c values were those that included E-W and N-S winds, but not W_p and W_c (ΔAIC_c = 3.41 in spring and 5.98 in autumn).

Figure 2. Summary of predictor averaged coefficients (β) ranked according to their predictive importance (Σw) in models with $\Delta AIC_c < 7$. Only results with a minimal $\Sigma w = 0.2$ are presented. Asterisks represent the *p*-value ranges: $p < 0.001$ (***), $p < 0.01$ (**), $p < 0.05$ (*), $p > 0.05$ (n.s.). Dependent variables are: probability of sea crossing (a, b), ground speed (c, d) and air speed (e, f). The baseline levels of the binomial variables "Route choice", "Age" and "Sex" are land-detour (*LAND*), *adult* and *female*, respectively.

3.2 Flight speed

3.2.1 Ground speed

The storks' ground speed was 7% faster on average in autumn than in spring (LMM: $\beta = -0.77 \pm 0.18$, $t_{192} = -4.3$, $p < 0.001$). Considering their daily track and regardless of the season, they were 8% faster on average when crossing the sea than when flying over land (LMM: $\beta = 0.7 \pm 0.17$, $t_{192} = 3.99$, $p < 0.001$). No difference in ground speed was found between adults and juveniles when data from the two migration seasons were pooled. In spring, white storks flew faster in tailwinds and slower under headwinds in general, but their route choice

modulated their response (Figure 2c). Over land, they increased their ground speed in tailwind
(and decreasing it under headwinds), but during sea crossing they maintained a rather steady
ground speed regardless of wind support (Figure S3). In autumn, storks flew faster under
stronger tailwinds (W_p), thermal uplifts, crosswinds (W_c) and when crossing the bay (which
was ranked as the most important variable in the model), compared to over-land flight. Also,
adults flew faster than juveniles in this season (Figure 2d, Table S26). In both seasons, the
best models with the lowest (or equal) AIC_c values included W_p and W_c, but not E-W and N-S
winds (ΔAIC_c = 11.45 in spring and 0 in autumn).

3.2.2 Airspeed
Overall, the storks' daily airspeed was 7% higher on average in spring than in autumn (LMM:
β = 0.55 ± 0.19, t_{192} = 2.89, p = 0.004), and adults were 9% faster on average than juveniles
(LMM: β = -0.5 ± 0.22, t_{192} = -2.25, p = 0.025). Notably, no significant difference in bird
airspeed was found between detouring and bay-crossing storks. In spring, bay-crossing storks
adjusted their airspeed to wind support (Figure 2e), decreasing it with tailwinds and
increasing it with headwinds, while land detouring storks did not adjust their airspeed to wind
conditions (Figure S3). Also, storks generally increased their airspeed with increasing wind
speed (Figure 2e). In autumn, stork airspeed was higher under stronger headwinds,
crosswinds and thermal uplifts and when crossing the bay (Figure 2f). In both seasons the best
models with the lowest AIC_c values were those that included W_p and W_c, but not E-W and N-
S winds (ΔAIC_c = 11.89 in spring and 40.81 in autumn).

3.3 Route choice and flight time, distance, energy and speed
We tested for differences in several flight parameters – namely time spent flying, distance
covered, ODBA, proportion of flapping flight – between the two route choices (LAND or
SEA) BEFORE, ACROSS and AFTER crossing the bay area, as well as over the entire daily
path of the birds (Figure 3, Table S1). We found that the distance covered and the time spent
flying depended on route choice. In spring, white storks that crossed the bay spent on average
2 h more time flying and covered 60 km more distance, with the main difference found after
crossing the bay, while in autumn the distance covered and the time spent flying were similar
between the two route choices (Figure 3a,b). The average distance covered over the bay was
55.47 km (range: 28.12 – 144.33) in spring, and 70.50 km (range: 25.36 – 182.81) in autumn
(Figure S4). ODBA and proportion of flapping flight were about 40% higher in storks that
crossed the bay in both seasons (Figure 3c,d and Figure 4) in the day that included the cross-
bay journey.

Figure 3. Summary statistics of (a) time spent flying, (b) distance covered, (c) Overall Dynamic Body Acceleration (ODBA), and (d) proportion of flapping flight of migrating storks flying over Iskenderun Bay area, according to the section of the flight path with respect to the bay (before, across or after) and the entire daily path, and by season. Colours represent the two route choices: land-detour (red) and sea-crossing (blue). Horizontal grey lines are averages per section and season (a, b) and overall average regardless of season and section (c, d). Dots are mean values and the shapes represent the distributions of the data. Asterisks indicate the *p*-value ranges: $p < 0.001$ (***), $p < 0.01$ (**), $p < 0.05$ (*). See also Figure S6 and S7 for explanations regarding differences in time and distance between the two route choices over the bay area.

Figure 4. Visualization of cumulative sum of Overall Dynamic Body Acceleration (ODBA) in relation to cumulative distance covered (a) and cumulative time spent flying (b). The curves show the average relationships recorded from white storks that crossed the bay (blue) and those that circumvented it (red). Closed circles of these colours represent the mean (± SD) of each selected route choice (sea-crossing and over-land). Sequences of open coloured (see details below) circles depict data from white storks such that each sequence represents data from a single track across the bay area. The circles' colour depicts the section over which they were recorded, with respect to the Iskenderun Bay: BEFORE (violet), ACROSS (yellow) and AFTER (green) the bay (see also electronic supplementary material, Figure S1).

4. Discussion

We highlight how important and consequential the choice of migration route is for soaring birds that either cross a sea barrier or fly around it. Our findings uncover how migration route selection over a shorter path that is nonetheless characterized by a "high transport cost" is undertaken. In our case the birds must flap over the sea while flying a longer over land detour route is associated with a lower transport cost because the birds are able to soar over it. We explore the factors that modulate route selection and explore its consequences for flight behaviour, migration speed and flight energetics in a large soaring migrant, the white stork. Specifically, wind influenced route selection which in turn affected bird ground and air speed, as well as the birds' flight mode selection (soaring-gliding flight over land and flapping flight over the sea). Furthermore, the combined effects of route selection and the environmental conditions influencing it are responsible for the increase in the time spent flying and distance

travelled of migrating storks that decided to cross the bay during spring. Yet, this longer daily

migration distance came with a likely energetic flight cost, highlighting a likely time-energy

trade-off in the migration of white storks. Yet, this benefit of sea crossing was found only in

spring. It is possible that the lower propensity of over-sea flights in autumn was based on

lower, or lack thereof of such benefits in this season. Because route selection was strongly

related to the local wind conditions at the day of passage, and was characterized by low

repeatability, we hypothesize that route choice is not based on a fixed strategy of each

individual but rather on a flexible selection based on local atmospheric conditions when

arriving to the bay area. It is also noteworthy that the storks migrate in flocks, thus the route

selection might not be an individual decision but rather a decision taken by the flock leaders

(Flack, Nagy, Fiedler, Couzin, & Wikelski, 2018), possibly masking individual-related

attributes. The lower rates of sea crossing in juveniles compared with adults could be related

to their lower migratory experience (Rotics et al., 2016b), or to the higher metabolic costs

associated with sea crossing. Possibly, juvenile birds trade-off time and energy in a different

manner than adults by responding more strongly to the negative aspects of the cross-bay

flight.

Overall, we tested four predictions, how (a) tailwinds and (b) time pressure could affect route

choice, and how (c) sea-crossing could save time, allowing extending the daily migration

distance, which may result in (d) higher metabolic costs due to flapping flight mode while

passing over the bay. Our first prediction (a) that tailwinds facilitate sea-crossing decision

was confirmed. Decreasing wind speed, increasing westward winds in spring and increasing

southward winds in autumn promoted sea crossing. The N-S and the E-W winds have a likely

supporting effect in each season, since the crossing of the bay took place mostly from north to

south in autumn and from east to west in spring. Similar results were reported by Meyer et al.

(2000) for fall migration of soaring migrants crossing the Strait of Gibraltar with favourable

southward and westward winds. In the same area also Griffon vultures (*Gyps fulvus*) were

observed to cross the Strait of Gibraltar with weak variable winds or similar tailwinds

(Bildstein et al., 2009). A facultative sea crossing for migration route shortcutting under

tailwind conditions was hypothesized in a radar study recording spring migrating soaring

birds in Southern Italy (Becciu et al., 2018).

Notably, we found contrasting responses to tailwinds between birds that selected the two

routes (see Figure S3). Specifically, birds that travelled over land adjusted their airspeed but

kept a steady airspeed in both tailwind and headwind conditions, increasing their ground

speed under tailwinds and decreasing it under headwinds. On the contrary, over the sea, when

the birds employed flapping flight (see below), the birds adjusted their airspeed and

maintained a quasi-steady ground speed, as observed in several studies of flapping birds and

bats (Liechti, 1995; Sapir, Horvitz, Dechmann, Fahr, & Wikelski, 2014).

Our second prediction (b) was nonetheless not supported by our exploration since early-

spring migrants commonly crossed the bay while relatively late migrants mostly detoured

over land (Figure 2a). We note that the higher tendency to cross the sea in spring and mostly

with westward winds may be related to less suitable thermal conditions over land in spring

that hindered soaring flight, compared to autumn. Moreover, soaring conditions likely

improved with ordinal date in spring, possibly explaining the increasing tendency for a land

detour with the progression of the spring (Figure S5). Notably, early-returning spring

migrants are typically considered as 'higher-quality' individuals, with better body condition

(Dittmann & Becker, 2003), breeding success (Smith & Moore, 2005), and flight performance

(Matyjasiak, 2013), which might explain their higher rates of selecting the shorter but energy

demanding sea-crossing route.

Our findings partially support our third prediction (c) that sea-crossing flight is beneficial as it

saves travelling time (see also Figure S7), and extends the daily distance travelled. Our data

suggest that this was the case only for spring but not for autumn. The results support our last

prediction (d) that sea-crossing is associated with higher movement-related metabolic costs,

since sea-crossing birds mostly used flapping flight and had higher ODBA (and thus likely

higher flight metabolism) compared with overland detouring birds.

In autumn, choosing one route or the other had no benefits in terms of more distance covered

after the bay, but due to the use of flapping flight when crossing the bay, the birds that flew

over the sea likely had higher flight metabolic costs compared with land detouring birds that

mostly flew using soaring flight. However, one has to bear in mind that due to the effects of

meteorological conditions on the decision to select a specific route, we could not compare

storks that used the two alternative routes under similar weather conditions. Hence, our

comparison of migration performance between overland versus sea-crossing tracks are limited

by the specific weather conditions that prevailed in the area in which the storks decided about

the selection of their route.

Understanding how atmospheric processes impact migration movements is of fundamental

importance in a time of climate change (La Sorte, Horton, Nilsson, & Dokter, 2018; Winkler

et al., 2014). In Turkey, and mostly in the area of Iskenderun Bay, wind speed and

specifically the E-W component of wind speed decreased over the last decades (Dadaser-

Celik & Cengiz, 2014), partially following changes in global circulation patterns and

increasing surface roughness (Vautard, Cattiaux, Yiou, Thépaut, & Ciais, 2010). Our results

indicate that migrants are sensitive to the dynamics of their aerial environment and their

behaviour and movement properties are strongly affected by local meteorological conditions. Changing atmospheric patterns due to climate change may thus result in changes in migration route selection of migrating white stroks, with possible implications for population dynamics (La Sorte, Fink, & Johnston, 2019) and conservation (Wilcove & Wikelski, 2008).

Acknowledgements

We thank T. Schaffer, H.G. Benecke and W. Sender and his crew in the Drömling Nature Park for their essential help in the field work in Germany, H. Schmid, H. Eggers, G. Sterzer and N. Aljadeff for their help in the data downloading, O. Hatzofe from the Israel Nature and Parks Authority and B. Keeves from the Max Planck Institute of Animal Behavior for their help in retrieving lost transmitters and W. Heidrich and F. Kuemmeth from e-obs GmbH for their dedicated technical support. We thank I. A. Lazagabaster for his help with the graphic visualizations. We acknowledge the generous funding of DIP grants (DFG) NA 846-1-1 and WI 3576-1-1 to R.N., F.J. and M.W. This study was also supported by the Minerva Center for Movement Ecology granted to R.N. and the Max Planck Institute of Animal Behavior. S.R. was supported by a doctoral bird study scholarship of the Ministry of Science and Technology, Israel.

Authors' contributions

P.B. and N.S. conceived the study and designed the methodology with the help of S.R., N.H. and R.N. S.R. and M.K. carried out the field work with the help of M.W. and D.Z. P.B. analysed the data and compiled all figures. P.B. and N.S. led the writing of the manuscript and all authors contributed to the revisions of the draft. S.R., N.H. and R.N. further contributed to data interpretation and manuscript revision. All authors contributed critically to the drafts and gave final approval for publication.

Data availability statement

The GPS-ACC data that was used in this study is available in Movebank Data Repository (movebank.org): https://doi.org/10.5441/001/1.v8d24552 (Rotics et al., 2018a), and https://doi.org/10.5441/001/1.hn1bd23k (Rotics et al., 2016a).

References

Agostini, N., Panuccio, M., & Pasquaretta, C. (2015). Morphology, flight performance, and water crossing tendencies of Afro-Palearctic raptors during migration. *Current Zoology*, *61*(6), 951–958. doi: 10.1093/czoolo/61.6.951

Alerstam, T. (2001). Detours in bird migration. *Journal of Theoretical Biology*, *209*(3), 319–331. doi: 10.1006/jtbi.2001.2266

Anderson, D. R., Burnham, K. P., & Thompson, W. L. (2000). Null Hypothesis Testing: Problems, Prevalence, and an Alternative. *Journal of Wildlife Management*, *64*(4), 912–923.

Anderson, D. R., Link, W. A., Johnson, D. H., & Burnham, K. P. (2001). Suggestions for Presenting the Results of Data Analyses. *Journal of Wildlife Management*, *65*(3), 373–378. Retrieved from http://www.jstor.org/stable/3803088

Barton, K. (2019). MuMIn: multi-model inference. Retrieved from https://cran.r-project.org/package=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). doi: 10.18637/jss.v067.i01

Becciu, P., Menz, M. H. M., Aurbach, A., Cabrera-Cruz, S. A., Wainwright, C. E., Scacco, M., … Sapir, N. (2019). Environmental effects on flying migrants revealed by radar. *Ecography*, *42*(5), 942–955. doi: 10.1111/ecog.03995

Becciu, P., Panuccio, M., Catoni, C., Dell'Omo, G., & Sapir, N. (2018). Contrasting aspects of tailwinds and asymmetrical response to crosswinds in soaring migrants. *Behavioral Ecology and Sociobiology*, *72*, 28. doi: 10.1007/s00265-018-2447-0

Bildstein, K. L. (2006). *Migrating raptors of the world: Their ecology and conservation*. Ithaca: Cornell University Press.

Bildstein, K. L., Bechard, M. J., Farmer, C., & Newcomb, L. (2009). Narrow sea crossings present major obstacles to migrating Griffon Vultures Gyps fulvus. *Ibis*, *151*(2), 382–391. doi: 10.1111/j.1474-919X.2009.00919.x

Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information- theoretic approach* (2nd ed.). New York: Springer-Verlag Inc.

Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and multimodel inference in behavioral ecology: Some background, observations, and comparisons. *Behavioral Ecology and Sociobiology*, *65*(1), 23–35. doi: 10.1007/s00265-010-1029-6

Dadaser-Celik, F., & Cengiz, E. (2014). Wind speed trends over Turkey from 1975 to 2006. *International Journal of Climatology*, *34*(6), 1913–1927. doi: 10.1002/joc.3810

Dittmann, T., & Becker, P. H. (2003). Sex, age, experience and condition as factors affecting arrival date in prospecting common terns, Sterna hirundo. *Animal Behaviour*, *65*(5), 981–986. doi: 10.1006/anbe.2003.2128

Dodge, S., Bohrer, G., Weinzierl, R., Davidson, S., Kays, R., Douglas, D., … Wikelski, M. (2013). The environmental-data automated track annotation (Env-DATA) system: linking animal tracks with environmental data. *Movement Ecology*, *1*(1), 3. doi: 10.1186/2051-3933-1-3

Duriez, O., Peron, G., Gremillet, D., Sforzi, A., & Monti, F. (2018). Migrating ospreys use thermal uplift over the open sea. *Biology Letters*, (December). doi: 10.1098/rsbl.2018.0687

Eisaguirre, J. M., Booms, T. L., Barger, C. P., McIntyre, C. L., Lewis, S. B., & Breed, G. A. (2018).
Local meteorological conditions reroute a migration. *Proceedings of the Royal Society B:*
Biological Sciences, *285*(1890). doi: 10.1098/rspb.2018.1779

Fawcett, T. (2006). An introduction to ROC analysis. *Pattern Recognition Letters*, *27*(8), 861–874. doi:
10.1016/j.patrec.2005.10.010

Flack, A., Nagy, M., Fiedler, W., Couzin, I. D., & Wikelski, M. (2018). From local collective behavior
to global migratory patterns in white storks. *Science*, *360*, 911–914.

Hartig, F. (2019). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression*
Models. Retrieved from http://florianhartig.github.io/DHARMa/ BugReports

Hastie, T., Tibshirani, R., & Friedman, J. (2009). *The Elements of Statistical Learning: Data Mining,*
Inference, and Prediction (2nd ed). doi: 10.1007/b94608

Hedenstrom, A. (1993). Migration by soaring or flapping flight in birds: The relative importance of
energy cost and speed. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
342(1302), 353–361. doi: 10.1098/rstb.1993.0164

Kerlinger, P. (1984). Flight behaviour of Sharp-shinned Hawks during migration. II: Over water.
Animal Behaviour, *32*, 1029–1034.

Kranstauber, B., Smolla, M., & Scharf, A. K. (2018). move: Visualizing and Analyzing Animal Track
Data Version. Retrieved from R package version 3.1.0. website: https://cran.r-
project.org/package=move

Kuhn, M. (2019). *caret: Classification and Regression Training*. Retrieved from
https://github.com/topepo/caret/

La Sorte, F. A., Fink, D., & Johnston, A. (2019). Time of emergence of novel climates for North
American migratory bird populations. *Ecography*, *42*, 1079–1091. doi: 10.1111/ecog.04408

La Sorte, F. A., Horton, K. G., Nilsson, C., & Dokter, A. M. (2018). Projected changes in wind
assistance under climate change for nocturnally migrating bird populations. *Global Change*
Biology, (July). doi: 10.1111/gcb.14531

Lenth, R. V. (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software*,
69(1). doi: 10.18637/jss.v069.i01

Liechti, F. (1995). Modelling optimal heading and airspeed of migrating birds in relation to energy
expenditure and wind influence. *Journal of Avian Biology*, *26*(4), 330–336. Retrieved from
http://www.jstor.org/stable/3677049

Matyjasiak, P. (2013). Timing of arrival from spring migration is associated with flight performance in
the migratory barn swallow. *Behavioral Ecology and Sociobiology*, *67*(1), 91–100. doi:
10.1007/s00265-012-1429-x

Meijer, R. J., & Goeman, J. J. (2013). Efficient approximate k-fold and leave-one-out cross-validation
for ridge regression. *Biometrical Journal*, *55*(2), 141–155. doi: 10.1002/bimj.201200088

Meyer, S. K., Spaar, R., & Bruderer, B. (2000). To cross the sea or to follow the coast? Flight
directions and behaviour of migrating raptors approaching the Mediterranean Sea in autumn.
Behaviour, *137*(3), 379–399.

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E. (2008). A

movement ecology paradigm for unifying organismal movement research. *Proceedings of the*
National Academy of Sciences, *105*(49), 19052–19059.

Norberg, U. M. (1990). *Vertebrate flight: mechanisms, physiology, morphology, ecology and evolution*.
doi: 10.1007/978-3-642-83848-4

Nourani, E., Yamaguchi, N. M., Manda, A., & Higuchi, H. (2016). Wind conditions facilitate the
seasonal water-crossing behaviour of Oriental Honey-buzzards Pernis ptilorhynchus over the
East China Sea. *Ibis*, *158*, 506–518. doi: 10.1111/ibi.12383

R Core Team. (2018). R: A language and environment for statistical computing. Retrieved from R
Foundation for Statistical Computing website: http://www.r-project.org/

Rotics, S., Kaatz, M., Resheff, Y. S., Turjeman, S. F., Zurell, D., Sapir, N., … Nathan, R. (2016a). Data
from: The challenges of the first migration: movement and behavior of juvenile versus adult
white storks with insights regarding juvenile mortality. *Movebank Data Repository*. doi:
10.5441/001/1.hn1bd23k.

Rotics, S., Kaatz, M., Resheff, Y. S., Turjeman, S. F., Zurell, D., Sapir, N., … Nathan, R. (2016b). The
challenges of the first migration: movement and behavior of juvenile versus adult white storks
with insights regarding juvenile mortality. *Journal of Animal Ecology*, (June). doi: 10.1111/1365-
2656.12525

Rotics, S., Kaatz, M., Turjeman, S., Zurell, D., Wikelski, M., Sapir, N., … Nathan, R. (2018a). Data
from: Early arrival at breeding grounds: causes, costs and a trade-off with overwintering latitude.
Movebank Data Repository. doi: https://doi.org/10.5441/001/1.v8d24552

Rotics, S., Kaatz, M., Turjeman, S., Zurell, D., Wikelski, M., Sapir, N., … Nathan, R. (2018b). Early
arrival at breeding grounds: Causes, costs and a trade-off with overwintering latitude. *Journal of*
Animal Ecology, *87*(6), 1627–1638. doi: 10.1111/1365-2656.12898

Rotics, S., Turjeman, S., Kaatz, M., Resheff, Y. S., Zurell, D., Sapir, N., … Nathan, R. (2017).
Wintering in Europe instead of Africa enhances juvenile survival in a long-distance migrant.
Animal Behaviour, *126*, 79–88. doi: 10.1016/j.anbehav.2017.01.016

Safi, K., Kranstauber, B., Weinzierl, R., Griffin, L., Rees, E. C., Cabot, D., … Bohrer, G. (2013).
Flying with the wind: Scale dependency of speed and direction measurements in modelling wind
support in avian flight. *Movement Ecology*, *1*(1), 1–13. doi: 10.1186/2051-3933-1-4

Sapir, N., Horvitz, N., Dechmann, D. K. N., Fahr, J., & Wikelski, M. (2014). Commuting fruit bats
beneficially modulate their flight in relation to wind. *Proceedings of the Royal Society B*, *281*,
20140018. doi: 10.1098/rspb.2014.0018

Sapir, N., Horvitz, N., Wikelski, M., Avissar, R., Mahrer, Y., & Nathan, R. (2011). Migration by
soaring or flapping: numerical atmospheric simulations reveal that turbulence kinetic energy
dictates bee-eater flight mode. *Proceedings. Biological Sciences / The Royal Society*, *278*(1723),
3380–3386. doi: 10.1098/rspb.2011.0358

Sapir, N., Wikelski, M., Mccue, M. D., Pinshow, B., & Nathan, R. (2010). Flight modes in migrating
European bee-eaters: Heart rate may indicate low metabolic rate during soaring and gliding.
PLoS ONE, *5*(11), 1–13. doi: 10.1371/journal.pone.0013956

Shamoun-Baranes, J., Bouten, W., & Van Loon, E. E. (2010). Integrating meteorology into research on
migration. *Integrative and Comparative Biology*, *50*(3), 280–292. doi: 10.1093/icb/icq011

Shamoun-Baranes, J., Liechti, F., & Vansteelant, W. M. G. (2017). Atmospheric conditions create

freeways, detours and tailbacks for migrating birds. *Journal of Comparative Physiology A*, *203*(6–7), 509–529. doi: 10.1007/s00359-017-1181-9

Smith, R. J., & Moore, F. R. (2005). Arrival timing and seasonal reproductive performance in a long-distance migratory landbird. *Behavioral Ecology and Sociobiology*, *57*(3), 231–239. doi: 10.1007/s00265-004-0855-9

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *8*(11), 1639–1644. doi: 10.1111/2041-210X.12797

Sugiura, N. (1978). Further analysts of the data by akaike' s information criterion and the finite corrections. *Communications in Statistics,Theory and Methods*, *A7*, 13–27.

Vautard, R., Cattiaux, J., Yiou, P., Thépaut, J. N., & Ciais, P. (2010). Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness. *Nature Geoscience*, *3*(11), 756–761. doi: 10.1038/ngeo979

Wilcove, D. S., & Wikelski, M. (2008). Going, Going, Gone: Is Animal Migration Disappearing. *PLoS Biology*, *6*(7), e188. doi: 10.1371/journal.pbio.0060188

Wilson, R. P., Börger, L., Holton, M. D., Scantlebury, D. M., Quintana, F., Rosell, F., … Shepard, E. L. C. (2019). Estimates for energy expenditure in free-living animals using acceleration proxies; a reappraisal. *Journal of Animal Ecology*, *in press*. doi: 10.1111/1365-2656.13040

Winkler, D. W., Jørgensen, C., Both, C., Houston, A. I., McNamara, J. M., Levey, D. J., … Piersma, T. (2014). Cues, strategies, and outcomes: how migrating vertebrates track environmental change. *Movement Ecology*, *2*(1), 10. doi: 10.1186/2051-3933-2-10

Yamaguchi, N. M., Arisawa, Y., Shimada, Y., & Higuchi, H. (2012). Real-time weather analysis reveals the adaptability of direct sea-crossing by raptors. *Journal of Ethology*, *30*(1), 1–10. doi: 10.1007/s10164-011-0301-1

Zuur, A. F. (2009). Mixed effects models and extensions in ecology with R. In *Statistics for biology and health*. doi: 10.1007/978-0-387-87458-6

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, *1*(1), 3–14. doi: 10.1111/j.2041-210X.2009.00001.x