Title: Intrinsic base substitution patterns in diverse species reveal links to cancer

and metabolism

One Sentence Summary:

Energy metabolism produces DNA damage leading to similar patterns of base

substitutions in many species and in human cancers.

Authors:

Suzana P. Gelova[1,2], Kassidy N. Doherty[3†‡], Salma Alasmar[3†§], and Kin Chan[1,2*]

Affiliations:

[1] Department of Biochemistry, Microbiology and Immunology, University of Ottawa

[2] Ottawa Institute of Systems Biology, University of Ottawa

[3] Biopharmaceutical Sciences Undergraduate Program, University of Ottawa

15   * Corresponding Author: kin.chan@uottawa.ca

16   † Equal Contribution

17   Current affiliations:

18   ‡ Common Law Section, University of Ottawa

19   § Department of Chemistry and Biomolecular Sciences, University of Ottawa

Abstract:

Analyses of large-scale sequencing data reveal that mutagenic processes create distinctive

patterns of base substitutions, called mutational signatures. Here, we analyzed

substitution patterns from seven model species and single nucleotide polymorphisms

(SNPs) in 42 species, totalling >1.9 billion variants. We found that the base substitution

patterns for many species most closely match mutational signature 5 in cancers.

Signature 5 is also ubiquitous in cancers and normal human cells, suggesting similar

patterns of mutation across species are likely due to conserved biochemistry. Finally, we

present evidence from yeast that sugar metabolism is directly linked to this form of DNA

damage. We propose that conserved metabolic processes in cells are coupled to

continuous generation of mutations, which are acted upon by genetic selection to drive

the evolution of species, and cancers.

Main Text:

Mutations in genomic DNA are the starting material for evolution by natural

selection. Mutagenesis is often thought of as a stochastic process, wherein the temporal

and positional occurrence of each mutational event is not predictable (*1*). And yet, sets of

mutations induced by a given DNA damaging agent or process do create reproducible,

characteristic patterns, called mutational signatures (*2*). Mutational signatures derived

from cancer genomics data have been published, providing a composite picture of the

recurrent patterns of base substitutions found in cancers, and in some cases, normal

tissues as well (*3*). These observations in humans led us to investigate whether similar

patterns of base substitutions are found when comparing genetic variation within and

between other species.

We analyzed mutational data (784,102,480 single base substitutions) from seven

model species, comprising: 1,011 varieties of budding yeast (*Saccharomyces cerevisiae*)

(*4*); 161 varieties of fission yeast (*Schizosaccharomyces pombe*) (*5*); 152 isolates of

roundworm (*Caenorhabditis elegans*) (*6*); 85 isolates of fruit fly (*Drosophila*

melanogaster) (*7*); 36 strains of mouse (*Mus musculus*) (*8*); 42 strains of rat (*Rattus*

norvegicus) (*9*); and 49 canines (*Canis familiaris*) (*10*). When the data were plotted

using the convention for reporting mutational signatures (i.e., six substitution types, each

subdivided into 16 contexts with one flanking base on either side, 5′ and 3′) (*11*), it

became clear that the majority of mutations in all seven species were either C→T or

T→C transitions (see Figure 1A). To assess the similarity of these base substitution

patterns to known mutational signatures, we computed cosine similarity values vs. a set

of published COSMIC (Catalog of Somatic Mutations in Cancer) signatures (see Figure

1B) (*2*). For context, a cosine similarity value of 1 is obtained when comparing identical

signatures. The substitution patterns for all seven model species were each most similar

to COSMIC signature 5 (see Figure 1C, referred to hereafter as COSMIC 5), which is one

of two 'clock-like' mutational signatures found in human, as the numbers of COSMIC 5

mutations essentially always increase monotonically with age (*12*). COSMIC 5 also

accounts for ~75% of de novo mutations in healthy human children (*13*). These results

show that, just as COSMIC 5 is found ubiquitously in human, intrinsic base substitution

patterns similar to COSMIC 5 are found in multiple model species.

Given that COSMIC 5-like base substitution patterns are intrinsic to these model

species, we then investigated if similar patterns are observed among SNPs in a larger,

more diverse set of species. We downloaded sets of available VCF (variant call format)

files from the NCBI SNP database for 44 non-human species (*14*). Two data sets were

obviously invalid, as they completely lacked two substitution types (C→G and T→A,

likely due to data processing error(s)), while another data set was dropped from analysis

because of probable sequencing artefacts (see Figure S1). Among the remaining 41

species (yielding 603,722,210 SNPs, see Figure S2), the large majority showed the

highest cosine similarity to COSMIC 5 (see Figure 2). We also analyzed 554,168,376

human SNPs from NCBI and found that they produced a pattern very similar to COSMIC

5 as well (cosine similarity = 0.957, see Figures 3A and S3). Taken altogether,

observations from prior studies in human and our results here strongly suggest that

conserved biochemical processes in many species are responsible for the set of COSMIC

5-like intrinsic base substitution patterns.

To quantify and visualize the extent of similarity among base substitution

patterns, we generated profiles showing means and 95% confidence intervals of relative

frequency for each mutation type among the seven model species (see Figure 3B). The

95% confidence intervals for most substitution types were relatively small, i.e., indicating

low variance (see Figure S4A). The most highly variable trinucleotides with respect to

differences in frequencies of mutations were TTA and ATT for T→A transversions (see

also Figure 1A). Despite these instances of (relatively) higher variability, the mean

values do provide a consensus COSMIC 5-like pattern among the seven model species.

The cosine similarity between this consensus pattern and COSMIC 5 itself is 0.928 (see

Figure S5).

Likewise, we applied this analysis to the SNP data from the NCBI database (see

Figure 3C). The 95% confidence intervals for the means were more compact, reflecting

the larger number of species examined and yielding a lower variance. Again, the

substitution types with higher mean relative frequencies showed more variability (see

Figure S4B). The consensus pattern from SNP data very closely matched the

corresponding pattern from mutations among the seven model organisms, with a cosine

similarity of 0.981 (see Figure S6). All four NCG motifs exhibit more common C→T

transitions in the SNP data, suggestive of possible contributions from deamination of 5-

methylcytosine at those motifs in a subset of mammals (e.g., goat, horse, macaque, sheep,

etc.), similar to COSMIC signature 1 (*3*). Also similar to the model organism data, the

consensus SNP pattern has a cosine similarity of 0.931 compared to COSMIC 5 (see

Figure S7). Taken together, we propose that for many species, their respective sets of

intrinsic base substitutions form a range of related patterns, and COSMIC 5 is simply the

corresponding human counterpart in this continuum. The close relatedness of COSMIC 5

and the human SNPs pattern (see Figure S3) is consistent with this model.

In principle, many species can be used as models to study the molecular

mechanisms underlying these COSMIC 5-like mutational patterns, as these mechanisms

are likely to be well conserved. We previously used a budding yeast system which

features controlled generation of genomic single-stranded DNA (ssDNA) to elucidate the

relative contributions of APOBEC3A and APOBEC3B cytidine deaminases to cancer

mutagenesis (*15*). This system relies on the *cdc13-1* temperature sensitive mutation (*16*),

such that shifting to 37°C causes telomere uncapping, extensive DNA end resection, and

cell cycle arrest in G_2 via the DNA damage checkpoint (*17*). Since exposed ssDNA is

much more prone to damage than double-stranded DNA, this system can acquire

mutations for analysis much more readily than more conventional approaches. When we

sequenced the genomes of cells that were allowed to accumulate mutations, we obtained

a pattern of base substitutions that resembled the consensus base substitution pattern in

the seven model species, the consensus pattern from SNPs of 41 species, and COSMIC 5

(cosine similarities = 0.941, 0.921, and 0.906, respectively, see Figures 4A and Figure

S8-S10). This corroborates the generality of COSMIC 5-like intrinsic base substitution

patterns.

We then carried out additional experiments to investigate mechanisms responsible

for the mutational pattern. Mutagenesis is dependent on the translesion DNA synthesis

(TLS) polymerases zeta and Rev1 (*18*), i.e., on the error-prone bypass of damaged bases

(*19*) (see Figure 4B). In contrast, deletion of *RAD30* (encoding polymerase eta) had no

effect, since Rad30 specializes in error-free bypass of ultraviolet light-induced

cyclobutane dimer photoproducts (*20*) and apparently plays no role in mutagenesis

opposite DNA damage that gives rise to the COSMIC 5-like pattern. Deletion of genes

important for error-free bypass also had no significant effect (see Figure S11). Moreover,

we found that energy metabolism is key to this form of mutagenesis. Cells maintained in

glucose (a fermentative, glycolytic energy source (*21*)) acquired mutations most

frequently, while cells in acetate (a carbon source that can be introduced directly into the

tricarboxylic acid cycle, and must be utilized by aerobic respiration (*22*)) had the fewest

acquired mutations (see Figure S12). Moreover, increasing the concentration of glucose

from 2% (the amount normally used in yeast media) to 8% led to a five-fold increase in

the frequency of inactivating mutations in the *CAN1* reporter gene (see Figure 4C).

Blocking the last step of glycolysis using auxin-induced degron alleles of *CDC19*

(encoding pyruvate kinase) caused a large decrease in mutagenesis (see Figure 4D),

further supporting the central role of energy metabolism to generate the DNA damage

which produces a COSMIC 5-like base substitution pattern. Finally, the pattern of

mutations acquired in 8% glucose is essentially the same pattern as in 2% glucose (cosine

similarity = 0.970, see Figure S13); there is simply more DNA damage overall in 8%

glucose.

Taken altogether, our results suggest the intrinsic base substitution patterns of

many organisms are linked to sugar metabolism. This would provide a plausible

explanation for why such similar patterns are found across many species, including

human, since the main pathways of sugar metabolism are well conserved (*23*). This

model also implies that processes of sugar metabolism will inevitably produce DNA

damage, which leads to mutations that fit a similar pattern across many species. In turn,

these mutations are starting material for evolution by natural selection. Since energy

generation is a prerequisite for living, the continuous basal level of mutagenesis which

results would ensure that natural selection will have an ever-renewing repertoire of

genetic variants to act upon. We propose the actual DNA damage itself is likely very

similar between species, but the resulting patterns are more or less divergent depending

on the similarity of the respective sets of error-prone TLS polymerases when comparing

between species (see Figures S14 and S15). Species with polymerases that are more

closely related would have more similar patterns than those from more diverged species.

Finally, a coupling between metabolism and COSMIC 5-like mutational patterns could

help explain why the extent of that type of mutagenesis can be highly variable in cancers,

when comparing between different patients (*12*) or temporally within the same patient

(*24*): namely, such variability would be linked to differences in sugar metabolism

between individuals or over the course of disease progression in particular patients.

References and Notes:

1. N. Timoféeff-Ressovsky, K. Zimmer, M. Delbrück, The nature of genetic mutations and structure of the gene. *Nachrichten Aus Biol. Ges. Wiss. Gött.* **1**, 189–245 (1935).

2. L. B. Alexandrov, S. Nik-Zainal, D. C. Wedge, P. J. Campbell, M. R. Stratton, Deciphering Signatures of Mutational Processes Operative in Human Cancer. *Cell Rep.* **3**, 246–259 (2013).

3. L. B. Alexandrov, S. Nik-Zainal, D. C. Wedge, S. A. J. R. Aparicio, S. Behjati, A. V. Biankin, G. R. Bignell, N. Bolli, A. Borg, A.-L. Borresen-Dale, S. Boyault, B. Burkhardt, A. P. Butler, C. Caldas, H. R. Davies, C. Desmedt, R. Eils, J. E. Eyfjord, J. A. Foekens, M. Greaves, F. Hosoda, B. Hutter, T. Ilicic, S. Imbeaud, M. Imielinsk, N. Jager, D. T. W. Jones, D. Jones, S. Knappskog, M. Kool, S. R. Lakhani, C. Lopez-Otin, S. Martin, N. C. Munshi, H. Nakamura, P. A. Northcott, M. Pajic, E. Papaemmanuil, A. Paradiso, J. V. Pearson, X. S. Puente, K. Raine, M. Ramakrishna, A. L. Richardson, J. Richter, P. Rosenstiel, M. Schlesner, T. N. Schumacher, P. N. Span, J. W. Teague, Y. Totoki, A. N. J. Tutt, R. Valdes-Mas, M. M. van Buuren, L. van /'t Veer, A. Vincent-Salomon, N. Waddell, L. R. Yates, Australian Pancreatic Cancer Genome Initiative, ICGC Breast Cancer Consortium, ICGC MMML-Seq Consortium, ICGC PedBrain, J. Zucman-Rossi, P. Andrew Futreal, U. McDermott, P. Lichter, M. Meyerson, S. M. Grimmond, R. Siebert, E. Campo, T. Shibata, S. M. Pfister, P. J. Campbell, M. R. Stratton, Signatures of mutational processes in human cancer. *Nature.* **500**, 415–421 (2013).

4. J. Peter, M. De Chiara, A. Friedrich, J.-X. Yue, D. Pflieger, A. Bergström, A.

Sigwalt, B. Barre, K. Freel, A. Llored, C. Cruaud, K. Labadie, J.-M. Aury, B. Istace, K. Lebrigand, P. Barbry, S. Engelen, A. Lemainque, P. Wincker, G. Liti, J. Schacherer, Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature*. **556**, 339–344 (2018).

5.   A. Lock, K. Rutherford, M. A. Harris, J. Hayles, S. G. Oliver, J. Bähler, V. Wood, PomBase 2018: user-driven reimplementation of the fission yeast database provides rapid and intuitive access to diverse, interconnected information. *Nucleic Acids Res*. **47**, D821–D827 (2018).

6.   D. E. Cook, S. Zdraljevic, J. P. Roberts, E. C. Andersen, CeNDR, the Caenorhabditis elegans natural diversity resource. *Nucleic Acids Res*. **45**, D650–D657 (2016).

7.   J. K. Grenier, J. R. Arguello, M. C. Moreira, S. Gottipati, J. Mohammed, S. R. Hackett, R. Boughton, A. J. Greenberg, A. G. Clark, Global diversity lines - a five-continent reference panel of sequenced Drosophila melanogaster strains. *G3 Bethesda Md*. **5**, 593–603 (2015).

8.   T. M. Keane, L. Goodstadt, P. Danecek, M. A. White, K. Wong, B. Yalcin, A. Heger, A. Agam, G. Slater, M. Goodson, N. A. Furlotte, E. Eskin, C. Nellåker, H. Whitley, J. Cleak, D. Janowitz, P. Hernandez-Pliego, A. Edwards, T. G. Belgard, P. L. Oliver, R. E. McIntyre, A. Bhomra, J. Nicod, X. Gan, W. Yuan, L. van der Weyden, C. A. Steward, S. Bala, J. Stalker, R. Mott, R. Durbin, I. J. Jackson, A. Czechanski, J. A. Guerra-Assunção, L. R. Donahue, L. G. Reinholdt, B. A. Payseur, C. P. Ponting, E. Birney, J. Flint, D. J. Adams, Mouse genomic variation and its effect on phenotypes and gene regulation. *Nature*. **477**, 289 (2011).

9. R. Hermsen, J. de Ligt, W. Spee, F. Blokzijl, S. Schäfer, E. Adami, S. Boymans, S. Flink, R. van Boxtel, R. H. van der Weide, T. Aitman, N. Hübner, M. Simonis, B. Tabakoff, V. Guryev, E. Cuppen, Genomic landscape of rat strain and substrain variation. *BMC Genomics*. **16**, 357–357 (2015).

10. E. A. Ostrander, G.-D. Wang, G. Larson, B. M. vonHoldt, B. W. Davis, V. Jagannathan, C. Hitte, R. K. Wayne, Y.-P. Zhang, Dog10K Consortium, Dog10K: an international sequencing effort to advance studies of canine domestication, phenotypes and health. *Natl. Sci. Rev.* (2019), doi:10.1093/nsr/nwz049.

11. F. Blokzijl, R. Janssen, R. van Boxtel, E. Cuppen, MutationalPatterns: comprehensive genome-wide analysis of mutational processes. *Genome Med*. **10**, 33 (2018).

12. L. B. Alexandrov, P. H. Jones, D. C. Wedge, J. E. Sale, P. J. Campbell, S. Nik-Zainal, M. R. Stratton, Clock-like mutational processes in human somatic cells. *Nat. Genet*. **47**, 1402 (2015).

13. R. Rahbari, A. Wuster, S. J. Lindsay, R. J. Hardwick, L. B. Alexandrov, S. Al Turki, A. Dominiczak, A. Morris, D. Porteous, B. Smith, M. R. Stratton, UK10K Consortium, M. E. Hurles, Timing, rates and spectra of human germline mutation. *Nat. Genet*. **48**, 126 (2015).

14. A. Kitts, L. Phan, M. Ward, J. B. Holmes, *The Database of Short Genetic Variation (dbSNP)* (National Center for Biotechnology Information (US), 2014; https://www.ncbi.nlm.nih.gov/books/NBK174586/).

15. K. Chan, S. A. Roberts, L. J. Klimczak, J. F. Sterling, N. Saini, E. P. Malc, J. Kim, D. J. Kwiatkowski, D. C. Fargo, P. A. Mieczkowski, G. Getz, D. A. Gordenin, An

APOBEC3A hypermutation signature is distinguishable from the signature of background mutagenesis by APOBEC3B in human cancers. *Nat Genet*. **47**, 1067–1072 (2015).

16.   L. H. Hartwell, R. K. Mortimer, J. Culotti, M. Culotti, Genetic Control of the Cell Division Cycle in Yeast: V. Genetic Analysis of cdc Mutants. *Genetics*. **74**, 267–286 (1973).

17.   B. Garvik, M. Carson, L. Hartwell, Single-stranded DNA arising at telomeres in cdc13 mutants may constitute a specific signal for the RAD9 checkpoint. *Mol*. *Cell*. *Biol*. **15**, 6128–6138 (1995).

18.   J. F. Lemontt, Mutants of yeast defective in mutation induced by ultraviolet light. *Genetics*. **68**, 21–33 (1971).

19.   S. Prakash, R. E. Johnson, L. Prakash, Eukaryotic translesion synthesis DNA polymerases: specificity of structure and function. *Annu*. *Rev*. *Biochem*. **74**, 317–353 (2005).

20.   R. E. Johnson, S. Prakash, L. Prakash, Efficient Bypass of a Thymine-Thymine Dimer by Yeast DNA Polymerase, Polη. *Science*. **283**, 1001 (1999).

21.   R. H. Hopkins, The selective fermentation of glucose and fructose by brewer's yeast. *Biochem*. *J*. **22**, 1145–1156 (1928).

22.   M. S. Esposito, R. E. Esposito, M. Arnaud, H. O. Halvorson, Acetate utilization and macromolecular synthesis during sporulation of yeast. *J*. *Bacteriol*. **100**, 180–186 (1969).

23.   J. M. Peregrín-Alvarez, C. Sanford, J. Parkinson, The conservation and evolutionary modularity of metabolism. *Genome Biol*. **10**, R63 (2009).

24.   S. C. Dentro, I. Leshchiner, K. Haase, M. Tarabichi, J. Wintersinger, A. G.

Deshwar, K. Yu, Y. Rubanova, G. Mcintyre, I. Vazquez-Garcia, K. Kleinheinz, D. G. Livitz, S. Malikic, N. Donmez, S. Sengupta, J. Demeulemeester, P. Anur, C. Jolly, M. Cmero, D. Rosebrock, S. Schumacher, Y. Fan, M. Fittall, R. M. Drews, X. Yao, J. Lee, M. Schlesner, H. Zhu, D. J. Adams, G. Getz, P. C. Boutros, M. Imielinski, R. Beroukhim, S. C. Sahinalp, Y. Ji, M. Peifer, I. Martincorena, F. Markowetz, V. Mustonen, K. Yuan, M. Gerstung, P. T. Spellman, W. Wang, Q. D. Morris, D. C. Wedge, P. Van Loo, Portraits of genetic intra-tumour heterogeneity and subclonal selection across cancer types. *bioRxiv* (2018), doi:10.1101/312041.

25.   H. Pagès, *BSgenome: Software infrastructure for efficient representation of full genomes and their SNPs* (https://doi.org/doi:10.18129/B9.bioc.BSgenome), *R package*.

26.   S. Arora, M. Morgan, M. Carlson, H. Pagès, *GenomeInfoDb: Utilities for manipulating chromosome names, including modifying them to follow a particular naming style* (https://doi.org/doi:10.18129/B9.bioc.GenomeInfoDb), *R package*.

27.   H. Li, A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*. **27**, 2987–2993 (2011).

28.   P. Danecek, A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, R. Durbin, 1000 Genomes Project Analysis Group, The variant call format and VCFtools. *Bioinformatics*. **27**, 2156–2158 (2011).

29.   B. J. Knaus, N. J. Grünwald, vcfr: a package to manipulate and visualize variant call format data in R. *Mol. Ecol. Resour.* **17**, 44–53 (2017).

30.   J. Kim, K. W. Mouw, P. Polak, L. Z. Braunstein, A. Kamburov, G. Tiao, D. J.

Kwiatkowski, J. E. Rosenberg, E. M. Van Allen, A. D. D'Andrea, G. Getz, Somatic ERCC2 mutations are associated with a distinct genomic signature in urothelial tumors. *Nat. Genet.* **48**, 600 (2016).

31.   A. Mayakonda, D.-C. Lin, Y. Assenov, C. Plass, H. P. Koeffler, Maftools: efficient and comprehensive analysis of somatic variants in cancer. *Genome Res.* **28**, 1747–1756 (2018).

32.   K. Chan, J. F. Sterling, S. A. Roberts, A. S. Bhagwat, M. A. Resnick, D. A. Gordenin, Base Damage within Single-Strand DNA Underlies In Vivo Hypermutability Induced by a Ubiquitous Environmental Agent. *PLoS Genet.* **8**, e1003149 (2012).

33.   R. J. Rothstein, in *Methods in Enzymology* (Academic Press, 1983; http://www.sciencedirect.com/science/article/pii/0076687983010150), vol. 101, pp. 202–211.

34.   A. Papagiannakis, B. Niebel, E. C. Wit, M. Heinemann, Autonomous Metabolic Oscillations Robustly Gate the Early and Late Cell Cycle. *Mol. Cell.* **65**, 285–295 (2017).

35.   Nishimura Kohei, Kanemaki Masato T., *Curr. Protoc. Cell Biol.*, in press, doi:10.1002/0471143030.cb2009s64.

36.   K. Chan, in *Genome Instability: Methods and Protocols*, M. Muzi-Falconi, G. W. Brown, Eds. (Springer New York, New York, NY, 2018; https://doi.org/10.1007/978-1-4939-7306-4_4), pp. 33–42.

37.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Meth*. **9**, 357–359 (2012).

38.   H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, 1000 Genome Project Data Processing Subgroup, The Sequence

1   Alignment/Map format and SAMtools. *Bioinformatics*. **25**, 2078–2079 (2009).

Acknowledgments:

We thank all members of the Chan laboratory and numerous other colleagues for their

critical feedback on this work. Funding was provided by a Tier 2 Canada Research

Chair, an NSERC Discovery Grant, an Ontario Early Researcher Award, and uOttawa

startup funding to K.C. Author contributions are as follows: All authors performed

experiments and discussed the data; S.P.G. and K.C. analyzed data; K.C. conceived and

supervised the study, and wrote the manuscript. The authors declare there are no

competing interests. Genomics data are available online as described in the Data

Acquisition section within Materials and Methods. Yeast strains are available on request.

Figure 1. Intrinsic base substitution patterns of seven model species resemble one

another and COSMIC 5. (A) 96-bar graphs showing the intrinsic base substitution

patterns of seven model species, derived by analysis of data from (*4–10*). The patterns

tended to show mostly C→T or T→C substitutions. Interestingly, both invertebrate

species and fission yeast also had relatively prominent T→A peaks at ATT and TTA

motifs. (B) Heatmap showing cosine similarity values to 30 COSMIC signatures. For all

seven species, the highest cosine similarity values suggested closest resemblance to

COSMIC 5. (C) COSMIC signature 5.

Figure 2. Heatmap depicting cosine similarity values comparing sets of SNPs from 41

non-human species vs. the 30 COSMIC signatures. Most species had the highest cosine

similarity values compared to COSMIC signature 5. See Figure S2 for the 96-bar

substitution patterns of these species.

Figure 3. Base substitution patterns for NCBI human SNPs, model organisms, and

NCBI SNPs from 41 non-human species. (A) The pattern obtained by plotting the entire

collection of human SNPs from NCBI. (B) 96-bar graph showing mean and 95%

confidence interval for relative frequencies of each substitution type among the seven

model species. (C) 96-bar graph showing mean and 95% confidence interval for relative

frequencies of each SNP type among the 41 non-human species from NCBI.

2 **Figure 4.** Corroborating genetic and genomic data from yeast model system. (A) Base

3 substitution pattern resulting from damage to exposed single-stranded DNA in yeast

4 metabolizing glucose. This is most closely related to the model species consensus

5 substitution pattern in Figure 3B, with cosine similarity of 0.941. (B) *CAN1* reporter

6 gene inactivation frequency was dependent on *REV3* (encoding the catalytic subunit of

polymerase zeta) and *REV1*, i.e., on error-prone translesion DNA synthesis. Deletion of

RAD30 (encoding polymerase eta) had no effect. Cells were temperature shifted to 37°C

for 6 hours. (C) Increasing glucose concentration to 8% resulted in a significant increase

in mutagenesis within single-stranded DNA, when compared to baseline conditions in 2%

glucose. Cells in 0.5% and 1% glucose both experienced significantly less mutagenesis

than controls in 2%. Cells were shifted to 37°C for 24 hours. (D) Mutagenesis was

dependent on completion of glycolysis. Inducible degradation of Cdc19 (pyruvate

kinase) blocked the last step of the glycolytic pathway and caused over a ten-fold

decrease in mutagenesis. Cells were shifted to 37°C for 24 hours in the presence or

absence of 1 mM auxin. * denotes $P < 0.05$; ** is $P < 0.01$; and *** is $P < 0.001$ by t-

test.

Materials and Methods

Data Acquisition

VCF files were downloaded from the following sources: *Saccharomyces*

cerevisiae, http://1002genomes.u-strasbg.fr/files/; *Schizosaccharomyces pombe*,

ftp://ftp.pombase.org/pombe/external_datasets/DJeffares_Diversity/; *Caenorhabditis*

elegans, https://www.elegansvariation.org/data/release/20160408; *Drosophila*

melanogaster, https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?study=SRP050151; *Mus*

musculus, ftp://ftp-mouse.sanger.ac.uk/REL-1807-SNPs_Indels/; *Rattus norvegicus*,

ftp://ftp.rgd.mcw.edu/pub/strain_specific_variants/Hermsen_et_al_40Genomes_Variants/

; *Canis familiaris*,

ftp://download.big.ac.cn/dogsd/dog10k/variations/58indiv.unifiedgenotyper.recalibrated_

95.5_filtered.pass_snp.vcf.gz; NCBI non-human SNP data,

ftp://ftp.ncbi.nih.gov/snp/organisms/archive/; and human SNP data,

ftp://ftp.ncbi.nih.gov/snp/latest_release/VCF/GCF_000001405.38.gz. Illumina WGS

reads from ssDNA system in either 2% or 8% glucose are here:

https://www.ncbi.nlm.nih.gov/sra/PRJNA608117.

Mutational Analyses of Downloaded VCFs

Where possible, we used the MutationalPatterns R package (version 1.10.0) to

extract mutational and trinucleotide contexts from VCF files, to plot mutational

signatures, to calculate cosine similarities, and to plot heatmaps (*11*). We constructed

BSgenome (*25*) reference genome packages in order to use MutationalPatterns to process

VCF files, if none of the stock reference genomes were appropriate. In cases where an

organism's reference genome proved to be incompatible with BSgenome 1.52.0 (or more

precisely, its dependency GenomeInfoDB 1.20.0 (*26*)), we used a script that extracted

mutational and trinucleotide context data. This script used a combination of Unix shell

functions, bcftools 1.9 (*27*), VCFtools 0.1.15 (*28*), base R (*29, 30*), and the MafTools

2.0.16 R package (*31*). Summed matrices of mutational counts were loaded into

MutationalPatterns for graphics rendering, in order to produce a consistent style for

figures. We noticed that sometimes, GenomeInfoDB was unable to resolve the

appropriate reference genome. A workaround for this issue is to ensure that only one file,

specifying the chromosome names of the species being analyzed, is present in the

~/Library/R/3.6/library/GenomeInfoDb/extdata/dataFiles folder. When the irrelevant

files for other species are moved (temporarily) to a different folder, GenomeInfoDB is

not given the option to use an incorrect reference. Most analyses were run on a custom

built PC operating Ubuntu version 18.04, with the remaining analyses run on a MacBook

Pro operating macOS version 10.14.5 or 10.14.6. R version 3.6.0 was used on both

computing platforms.

Reagents and Consumables

Bacto peptone (product code 211677) and yeast extract (212750) were purchased

from Becton, Dickinson and Co. (Franklin Lakes, New Jersey). Auxin (3-indoleacetic

acid, I3750), canavanine (C9758), and adenine sulfate dihydrate (AD0028) were

purchased from MilliporeSigma (St. Louis, Missouri). Agar (FB0010), carbon sources

(i.e., glucose [GB0219], galactose [GB0215], raffinose [RJ392], and potassium acetate

[PB0438]), hygromycin (BS725), PCR purification spin column kit (BS654), agarose

(D0012), and Tris-Borate-EDTA (TBE) buffer (A0026) were purchased from BioBasic

(Markham, Ontario). G418 sulfate (450-130) was purchased from Wisent (St-Bruno,

Québec). Q5 PCR kit were purchased from New England Biolabs Canada (Whitby,

Ontario). Plasmid pOsTIR1w/oGFP (for integrating the *Oryza sativa TIR1* gene into the

HO locus in yeast) was purchased from Addgene (Watertown, Massachusetts). pMK152

plasmid (for degron tagging *CDC19*) was purchased from the National BioResource

Project Yeast Genetic Resource Center (NBRP YGRC, Osaka, Japan). Petri dishes

(82.1473.001) were purchased from Sarstedt (Montréal, Québec).

Yeast Genetics

All yeast strains were constructed in the ySR127 background, a *MATα* haploid

bearing the *cdc13-1* temperature sensitive allele. In addition, there is a cassette of three

reporter genes (*CAN1*, *URA3*, and *ADE2*) near the de novo left telomere of chromosome

V. Details about ySR127 were described previously (*32*).

Gene knockouts were constructed by one step gene replacement using antibiotic

resistance marker cassettes (*33*). To construct the CDC19 degron strains, we first

introduced the *O. sativa TIR1* gene cassette (OsTIR1) into the *HO* locus of ySR127,

using the protocol described in (*34*). We then inserted the 3mAID degron tag (containing

three tandem repeats of the core domain of the IAA17 protein from *Arabidopsis*

thaliana), using the protocol described in (*35*). Genetic constructions were verified by

diagnostic replica plating and PCR.

Mutagenesis experiments were initiated by inoculating single colonies each into 5

mL of rich media (2% Bacto peptone, 1% Bacto yeast extract, specified percentage of a

carbon source, supplemented with 0.001% adenine sulfate) in round bottom glass tubes.

Cells were grown at permissive temperature (23°C) for three days. Then, cultures were

diluted ten-fold into fresh media and shifted to restrictive temperature (37°C). Auxin was

added to 1 mM final concentration for degron experiments. After temperature shift, cells

were collected by light centrifugation, washed in water, and plated (using a turntable and

cell spreader) onto synthetic complete media to assess survival and onto canavanine-

containing media to select for mutants. Further details of this procedure are described in

(*36*).

<u>Illumina Whole Genome Sequencing</u>

To generate mutants for sequencing, four replicate ySR127 colonies were each suspended

into a small volume of water (e.g., 50 μL). Half of each cell suspension was used to

inoculate 5 mL of YPDA liquid media (2% Bacto peptone, 1% Bacto yeast extract, 2%

glucose, supplemented with 0.001% adenine sulfate). The other half was used to

inoculate 5 mL of YPDA-8 (2% Bacto peptone, 1% Bacto yeast extract, 8% glucose,

supplemented with 0.001% adenine sulfate). Cultures were grown for three days at 23°C.

500 μL of each culture was then diluted with 4.5 mL of fresh media and shifted to 37°C.

Cells from the remainder of each 23°C culture were collected by centrifugation and

pellets were frozen at -20°C. Can^r Ade$^-$ mutants were collected and reporter gene loss of

function phenotypes were verified as described (*36*). Genomic DNA from mutants of

interest were extracted and sequenced as described (*15*). Illumina whole genome

sequencing was outsourced to Genome Québec (McGill University, Montréal). Eight

parental cultures (actually four pairs of matched 2% and 8% glucose cultures) and 87

mutants derived from those parentals were sequenced. Bowtie2 version 2.3.5.1 (*37*),

SAMtools 1.9 (*38*), and bcftools 1.9 (*27*) were used to map reads and call variants.

Variants with quality score < 30 and/or with sequencing coverage < 10 were filtered out.

Variants found in parentals were filtered from the sets of variants identified in the

corresponding derivative mutants. The resulting VCF files, containing mutant-specific

variants, were passed to MutationalPatterns for further analysis and visualization.

1 Supplementary Figures

A

B

C

3 **Fig. S1.**

4 SNP patterns from NCBI database that were disqualified from further analysis. (A)

5 Plotting of the SNP dataset for apple (*Malus domestica*) revealed an apparent data

processing issue that deleted all C→G and T→A substitutions. (B) The SNP dataset for

flycatcher (*Ficedula albicollis*) was also affected by this issue. (C) The SNP pattern for

Nomascus leucogenys (the northern white-cheeked gibbon, but referred to simply as

"primates" on NCBI) was also suspiciously unusual with very high peaks at the

reciprocal motifs ACA and ATA, as well as ACG and ATG, for both C→T and T→C

substitutions. As this was very much an outlier, the *Nomascus leucogenys* dataset was

dropped for further consideration due to suspected sequencing artefact.

A

B

C

D

E

Fig. S2.

96-bar profiles showing the intrinsic base substitution patterns inferred from SNPs of 41

non-human species downloaded from NCBI. (A) Patterns for *Aspergillus fumigatus*

(ascomycetes_746128), *Apis mellifera* (bee_7460), *Daucus carota* (carrot_79200), *Felis*

catus (cat_9685), *Gallus gallus* (chicken_9031), *Cicer arietinum* (chickpea_3827), and

Pan troglodytes (chimpanzee_9598). (B) Patterns for *Callithrix jacchus*

(common_marmoset_9483), *Zea mays* (corn_4577), *Gossypium hirsutum* (cotton_3635),

Bos taurus (cow_9913), *Canis familiaris* (dog_9615), and *Brassica oleracea*

(eudicots_3712). (C) Patterns for *Drosophila melanogaster* (fruitfly_7227), *Capra hircus*

(goat_9925), *Vitis vinifera* (grape_29760), *Chlorocebus sabaeus* (green_monkey_60711),

Equus caballus (horse_9796), *Macaca fascicularis* (macaque_9541), and *Macaca*

mulatta (macaque_9544). (D) Patterns for *Anopheles gambiae* (mosquito_7165), *Mus*

musculus (mouse_10090), *Monodelphis domestica* (opossum_13616), *Sus scrofa*

(pig_9823), *Ornithorhynchus anatinus* (platypus_9258), *Coturnis japonica* (quail_93934)

and *Brassica napus* (rapeseeds_3708). (E) Patterns for *Rattus norvegicus* (rat_10116),

Ovis orientalis (red_sheep_469796), *Oryza sativa* (rice_4530), *Salmo salar*

(salmon_8030), *Ovis aries* (sheep_9940), *Sorghum bicolor* (sorghum_4558), and *Glycine*

max (soybean_3847). (F) Patterns for *Solanum lycopersicum* (tomato_4081), *Meleagris*

gallopavo (turkey_9103), *Capra hircus* (wild_goat_9923), *Taeniopygia guttata*

(zebra_finch_59729), *Danio rerio* (zebrafish_7955), and *Bos indicus* (zebu_9915). Most

patterns showed, to varying extents, a preponderance of C→T and T→C substitutions

where these substitution types were similar in overall proportion, forming a range of

COSMIC 5-like patterns. Notable exceptions to this overarching meta-pattern were

mosquito, bee, grape, and corn.

Fig. S3.

Comparison of the substitution profile from 554,168,376 human SNPs vs. COSMIC

signature 5. The high cosine similarity value of 0.957 suggested that the mutagenic

processes happening in cancers and over evolutionary time in humans are most likely the

same, but the rate of mutation acquisition is significantly higher in cancers.

A

B

4   **Fig. S4.**

The most frequently mutated motifs tended to show the most variability in mutation

frequency when comparing among species. (A) Mean relative frequencies of mutations

from the 96-bar graph for the consensus pattern among seven model species (see Figure

3B) were plotted with their corresponding 95% confidence intervals. (B) Mean relative

frequencies of mutations from the 96-bar graph for the consensus pattern among 41

species with NCBI SNP data (see Figure 3C) were plotted with their corresponding 95%

confidence intervals. For both groups, frequency of mutation at a trinucleotide motif was

positively correlated with variability (i.e., larger 95% confidence intervals).

Fig. S5.

Comparison of the consensus substitution profile mean from seven model species (see

Figure 3B) vs. COSMIC signature 5. The cosine similarity value of 0.928 suggested that,

despite the evolutionary distance between these model species and human, the underlying

biochemical mechanisms creating the intrinsic DNA damage are likely to be well

conserved.

Fig. S6.

Comparison of the consensus substitution profile mean from 41 (non-human) species

with NCBI SNP data (see Figure 3C) vs. the analogous pattern from seven model species

(see Figure 3B). The very high cosine similarity value of 0.981 strongly implied that the

underlying biochemical mechanisms creating the intrinsic DNA damage are likely to be

well conserved across many species.

Fig. S7.

Comparison of the consensus substitution profile mean from 41 (non-human) species

with NCBI SNP data (see Figure 3C) vs. COSMIC signature 5. Similar to the model

species data, the high cosine similarity value of 0.931 suggested that, despite significant

evolutionary divergence vs. human in many species, the underlying biochemical

mechanisms which damage the DNA to produce these related substitution patterns are

likely to be well conserved.

Fig. S8.

Comparison of the mutational profile observed in yeast single-stranded DNA vs. the

consensus pattern among seven model species. Cosine similarity = 0.941.

2 **Fig. S9.**

3 Comparison of the mutational profile observed in yeast single-stranded DNA vs. the

4 consensus pattern from 41 (non-human) species with NCBI SNP data. Cosine similarity

5 = 0.921.

2 **Fig. S10.**

3 Comparison of the mutational profile observed in yeast single-stranded DNA vs.

4 COSMIC signature 5. Cosine similarity = 0.906.

Fig. S11.

CAN1 reporter gene inactivation frequency as a function of DNA damage tolerance or

DNA repair genotype. Deletion of *REV3* or *REV1* had the largest effect on mutagenesis,

as they encode for the catalytic subunit of translesion DNA synthesis (TLS) polymerase

zeta and for the sole subunit of a second TLS polymerase, respectively. Deletion of

genes involved in error-free damage bypass (*RAD5*, *RAD18*, and *MMS2*) had only minor

effects which were not statistically significant, after correction for false discovery for

multiple t-tests. Deletion of *RAD30* (TLS polymerase eta) had no effect, nor did deletion

of *RAD51* or *RAD52*, both encoding key players in homologous recombination.

Fig. S12.

CAN1 reporter gene inactivation frequency as a function of carbon source. Frequencies

at Day 0 denote the background gene inactivation when cells were simply grown at 23°C.

Glucose was the most mutagenic, leading to a gene inactivation frequency of $\sim 10^3$ after

one day at 37°C to damage ssDNA. Acetate, a non-fermentable carbon source that is

directly fed into the tricarboxylic acid cycle, then into oxidative phosphorylation, was the

least mutagenic. Galactose and raffinose, which have mixed "respiro-fermentative"

metabolism, were intermediate in mutagenicity.

Fig. S13.

Comparison of the patterns from metabolizing 2% glucose vs. 8% glucose revealed a

very high cosine similarity (0.970), indicating that in 8% glucose, cells simply acquired

more of the same pattern of mutations.

4   **Fig. S14.**

5   Cosine similarity, when comparing base substitution patterns in (non-human) NCBI SNP

6   species vs. NCBI human SNPs pattern, was plotted as a function of amino acid identity to

the two key translesion DNA synthesis polymerases REV1 (A) and REV3 (B). 95%

confidence intervals are shaded in gray. Species whose REV1 and REV3 homologs are

more similar to the human polymerases tended to have base substitution patterns with

higher cosine similarity. Remarkably, many species with relatively low amino acid

identity (less than 0.5) still had cosine similarity values > 0.9, underscoring the extensive

conservation of COSMIC 5-like patterns throughout evolution.

Fig. S15.

Model explaining how a continuum of COSMIC 5-like patterns arises among diverse

species. DNA damage from metabolic source(s) are similar among species. Translesion

DNA synthesis polymerases (TLS Pols) insert bases in an error-prone manner opposite

the base damage. The species with TLS polymerases which are most similar to human

(shown in orange) produce intrinsic base substitution patterns with the highest cosine

similarity values (~ 0.97) when compared to COSMIC 5. Species with TLS polymerases

that are most diverged from those in human (shown in blue) yield base substitution

patterns in the lower end of the cosine similarity range (~ 0.90) vs. COSMIC 5. Species

1 with TLS polymerases that are intermediate also tend to produce base substitution

2 patterns that are intermediate in cosine similarity. Notably, even the most diverged TLS

3 polymerases can still yield cosine similarity values > 0.9 vs. COSMIC 5.