Learning improves decoding of odor identity with phase-referenced oscillations

in the olfactory bulb

Justin Losacco[1,2,*], Daniel Ramirez-Gordillo[2,*], Jesse Gilmer[1,3] and Diego Restrepo[1,2,#]

[1] Neuroscience Graduate Program

[2] Department of Cell and Developmental Biology

[3] Department of Physiology and Biophysics

University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA

[*] J.L and D.R.G. contributed equally to this work

[#] Corresponding author: Diego Restrepo, diego.restrepo@cuanschutz.edu

Abstract

Local field potential oscillations reflect temporally coordinated neuronal ensembles—

coupling distant brain regions, gating processing windows, and providing a reference for

spike timing-based codes. In phase amplitude coupling (PAC), the amplitude of the

envelope of a faster oscillation is larger within a phase window of a slower carrier wave.

Here, we characterized PAC, and the related theta phase-referenced high gamma and

beta power (PRP), in the olfactory bulb of mice learning to discriminate odorants. PAC

changes throughout learning, and odorant-elicited changes in PRP increase for

rewarded and decrease for unrewarded odorants. Contextual odorant identity (is the

odorant rewarded?) can be decoded from peak PRP in animals proficient in odorant

discrimination, but not in naïve mice. As the animal learns to discriminate the odorants

the dimensionality of PRP decreases. Therefore, modulation of phase-referenced

chunking of information in the course of learning plays a role in early sensory

processing in olfaction.

Significance

Early processing of olfactory information takes place in circuits undergoing slow

frequency theta oscillations generated by the interplay of olfactory input modulated by

sniffing and centrifugal feedback from downstream brain areas. Studies in the

hippocampus and cortex suggest that different information "chunks" are conveyed at

different phases of the theta oscillation. Here we show that in the olfactory bulb, the first

processing station in the olfactory system, the amplitude of high frequency gamma

oscillations encodes for information on whether an odorant is rewarded when it is

observed at the peak phase of the theta oscillation. Furthermore, encoding of

information by the theta phase-referenced gamma oscillations becomes more accurate

as the animal learns to differentiate two odorants.

Introduction

Animals must modulate early sensory processing to optimize navigation of their

environment (Gire et al., 2013a; Pakan et al., 2018). This experience-dependent

shaping involves an interplay between sensory input, behavioral state, arousal and

motor activity. Neuronal activity, temporally organized by local field potential (LFP)

oscillations, carries distinct information at different phases of each cycle (Jensen, 2001;

Kepecs et al., 2006; Lisman and Jensen, 2013). For example, in the hippocampus

different phases of the theta LFP (2-12 Hz) preferentially encode present and future

spatial locations, illustrating temporal "chunking" of information (Amemiya and Redish,

2018; Siegle and Wilson, 2014). Place cell preferred fields are encoded temporally

through phase precession in relation to hippocampal theta oscillations (Buzsaki and

Draguhn, 2004; Lisman, 2005; O'Keefe and Recce, 1993; Skaggs et al., 1996).

Additionally, LFP oscillations in different frequency bandwidths can be related through

cross frequency coupling (CFC), which can further enhance information carrying

capacity (Canolty and Knight, 2010; Engel et al., 1999; Florin and Baillet, 2015; Fries,

2005).

One example of CFC is PAC, wherein the amplitude of a faster oscillation (e.g.

gamma, 35-95 Hz) is related to the phase of a slower oscillation (e.g. theta, 2-12 Hz)

(Chrobak and Buzsaki, 1998; Soltesz et al., 1993). PAC occurs during task engagement

in rodent hippocampus (Bragin et al., 1995; Lisman and Idiart, 1995; Soltesz et al.,

1993), rodent orbitofrontal cortex (van Wingerden et al., 2014), macaque auditory cortex

(Marton et al., 2019), human hippocampus (Lega et al., 2016) and human visual cortex

(Daume et al., 2017; Seymour et al., 2017). Interestingly, Van Wingerden et al. (van

Wingerden et al., 2014) described how theta-gamma PAC strengthens in rat

orbitofrontal cortex with learning for trials in which the correct decision was made in the

olfactory go no-go task.

The study of PAC in the earliest sensory processing brain areas is virtually non-

existent. The olfactory bulb (OB) is an ideal location to study whether PAC is relevant to

sensory processing; its intrinsic oscillatory activity is linked to odorant signal processing

(Doucette et al., 2011; Gschwend et al., 2012; Li et al., 2015; Smear et al., 2011). The

rhythmic activation of olfactory sensory neurons (OSNs) through breathing generates a

theta-frequency (2-12 Hz) respiratory-related LFP in the target region of the OSNs, the

OB, that is coherent with oscillations in downstream processing areas such as piriform

cortex (PC) and hippocampus (Gourevitch et al., 2010; Heck et al., 2019; Nguyen Chi et

al., 2016; Tort et al., 2018). Interestingly, PAC is present in human PC, and the

respiratory phase for presentation of stimuli modulates the ability to retrieve episodic

memory (Zelano et al., 2016). Downstream neural networks receiving olfactory

information from the OB could be tuned to respond to sensory input at certain phases of

this theta LFP (Buzsaki, 2010; Jensen, 2001) and could send back modulatory signals

during other phases (Kay, 2015; Kepecs et al., 2006).

Decades of work have characterized LFP activity in the OB in various states of

wakefulness (Adrian, 1950; Chery et al., 2014; Li et al., 2012) and olfactory task

engagement (Gourevitch et al., 2010; Martin and Ravel, 2014). Surprisingly, this LFP

signal changes in response to prior experience (Courtiol and Wilson, 2017; Gire et al.,

2013b). Work in the go no-go operant conditioning task where the animal learns to

respond to the reinforced odorant for a water reward has revealed broadband LFP

power changes for the rewarded versus the unrewarded odorant with discrimination

learning (Beshel et al., 2007; Martin et al., 2004; Ramirez-Gordillo et al., 2018; Stopfer

and Laurent, 1999). Additionally, changes in the balance of sensory and corticofugal

inputs to OB interneurons can alter timing of beta and gamma oscillatory states,

suggesting that information transfer from the OB is influenced by top-down signaling

(David et al., 2015). Also, these investigators posited that slower theta oscillations gate

temporal windows for sensory and centrifugal inputs to ascend and descend,

respectively, suggesting an important role for PAC in gating information transfer. Since

PAC exists in the OB in anesthetized (Buonviso et al., 2003) and awake, idle rats

(Rojas-Libano et al., 2014), we wondered whether PAC changes with learning,

reflecting unsupervised Bayesian learning or supervised learning through top-down

modulation sensory information (Hiratani and Latham, 2019), and whether encoding this

information by the power of high frequency oscillations differs as a function of the phase

of the theta LFP.

Here, thirsty mice implanted with tetrodes in the OB learned to associate an

odorant with water reward in the go no-go task, discriminating between two

monomolecular odorants (e.g. between isoamyl acetate vs acetophenone, IAAP) or

between a monomolecular odorant and an odorant mixture (between ethyl acetate and

a mixture of ethyl acetate and propyl acetate, EAPA). LFP was analyzed over task

learning for changes in PAC and for the ability of theta phase-referenced high frequency

oscillatory power—termed phase referenced power (PRP) hereafter—to encode for the

perceived contextual identity of the odorant.

We demonstrate that the strength of PAC, quantified as the modulation index

(Tort et al., 2010), changes for the rewarded odorant (S+) as the animals learn.

Interestingly, the variance of the angle of theta with the strongest gamma oscillation

(peak angle) increases for the unrewarded odorant (S-) over the course of learning.

These changes in PAC raised the question whether a downstream observer focused on

high frequency oscillations at different theta phases would find a difference in stimulus

prediction with learning. We found that the ability to decode stimulus identity by PRP

improves with task proficiency and the dimensionality, a measure of separable

representation within signals, decreased as the animal learned to differentiate the

odorants, suggesting that a stable and unambiguous representation of odor identity

arises in PRP after learning.

Results

The experiments were designed to study encoding of odorant identity by cross-

frequency coupling. We report data obtained in two different sets of experiments that we

name Exp1 and Exp2. As shown in Table S1 the two experiments differed in electrode

locations, mouse genotype, device implantation (tetrode vs. optetrode), and odorant

pairs used (see also Methods). Mice learned to discriminate between rewarded (S+)

and unrewarded odorants (S-) presented in pseudorandomized order in the go no-go

olfactory task (Figure 1A-C). The dataset is comprised of 119 recording sessions in 30

mice. The odorant pairs tested were different volatile compounds (or mixtures) whose

nomenclature addresses the odorant names and the experimental set (e.g. IAAPexp1,

see Table S1 for the nomenclature). Table S2 enumerates the total number of sessions

per odorant pair, mouse, and experiment.

In the go-no go task mice start the trial spontaneously by poking their nose into

the odor spout and licking on the lick port. The odorant is delivered at a random time 1-

1.5 sec after nose poke (the time course for odorant concentration is shown in Figure 1-

figure supplement 1). The mice must decide to either lick a waterspout at least once

during each 0.5-second bin in the 2-second response area (green blocks in Figure 1B)

to obtain a water reward for the rewarded odorant or refrain from licking for the

unrewarded odorant (Figure 1C). We did not punish for licking for the unrewarded

odorant, and mice refrain because of the effort it takes to lick. Behavioral performance

was termed naïve or proficient when their performance estimated in a 20-trial window

was ≤ 65% for naïve and ≥ 80% for proficient. After three 20-trial blocks of proficient

performance, the session was ended; valence was reversed the next day (not shown) or

another odorant was tested. We recorded the LFP using four tetrodes (16 electrodes)

implanted in the OB and we analyzed the data to determine whether information carried

by cross-frequency coupling can encode for the contextual identity of the odorant for

naïve or proficient mice. Importantly, reversal experiments from previous studies where

we switched the rewarded and unrewarded odorants have shown that the power of the

LFP in the OB and mitral/tufted cell spiking encodes for the *contextual* identity of the

odorant (is the odorant rewarded?)(Doucette et al., 2011; Doucette and Restrepo, 2008;

Li et al., 2015; Ramirez-Gordillo et al., 2018). Therefore, when we refer to identity in this

publication we do not mean the chemical identity of the odorant, we mean whether the

odorant is rewarded or unrewarded. Finally, to evaluate the statistical significance of

differences in oscillatory parameters estimated in this study the estimates were either

averaged in the time period of odorant application (0.5 to 2.5 sec after diverting the

odorant to the odor delivery spout) or the statistical significance was evaluated for the

entire time course with time points every 0.1 sec using a generalized linear model

(GLM, see Methods). Furthermore, we complement testing of statistical significance

using p values with estimation of bootstrapped confidence intervals (Halsey et al.,

2015).

Phase amplitude coupling analysis of the LFP recorded in the go no-go

behavioral task

In order to understand whether odorant identity is encoded by by cross-frequency

coupling in the OB we characterized PAC, which is a cross-frequency coupling

mechanism where high frequency oscillation bursts take place at specific phases of low

frequency theta oscillations (Tort et al., 2010). PAC has been reported in the OB

(Buonviso et al., 2003; Rojas-Libano et al., 2014), but has not been thoroughly

characterized. Figures 1D-E show our approach to quantify the strength of PAC using

the modulation index, a measure of how localized high frequency firing is within the

phase of theta oscillations. Figures 1Di and ii show an example of the extracellular LFP

sampled at 20 kHz and filtered between 1-750 Hz. The raw signal (Figure 1Di) was

filtered with a 20th order Butterworth filter into different LFP frequency bands (Figures

1Dii, iii and v) (theta, 6-14 Hz, adapted from (Nguyen Chi et al., 2016); beta, 15-30 Hz;

or high gamma, 65-95 Hz). Figures 1Diii and 1Dv show that the filtered theta and high

gamma LFP appear to change amplitude in a coordinated manner. We used PAC

analysis (Tort et al., 2010) to evaluate the degree of coupling of the amplitude of the

envelope of the beta or high gamma LFP on the phase of the theta LFP. Figure 1Div

shows the theta phase and the red line in Figure 1Dv shows the envelope for the

amplitude of the high gamma LFP, both calculated with the Hilbert transform as detailed

in Tort et al. (Tort et al., 2010). Figure 1E shows the phase amplitude plot of the

normalized distribution of high gamma amplitudes in different theta phase bins. For this

example there are two maxima, and the vertical red line corresponds to the phase of

theta at which the gamma amplitude is highest (at ~95 degrees); this is called the "peak

phase". Similarly, the phase with the smallest normalized amplitude is called the "trough

phase" (vertical blue line at ~5 degrees). The strength of PAC was estimated as the

modulation index (MI), a measure computed as the Kullback–Leibler (KL) distance to

the uniform phase distribution normalized to range from 0 (uniform distribution) to 1

(sharp gamma bursts at a specific theta phase) (Tort et al., 2010). The MI for the

distribution in Figure 1E is 0.018. Figure 1-figure supplement 2 shows the LFP power

spectrogram for simulated data with low gamma oscillations that are either uniformly

distributed in the theta phase (MI=0.0004, Figure 1-figure supplement 2A) or with

gamma bursts at 180 degrees of the theta oscillation (MI=0.01, Figure 1-figure

supplement 2B). Recordings in hippocampus and cortex yield MI values in the range of

0.005-0.03.

Peak angle variance for theta/high gamma phase amplitude coupling increases

for S- when the animals learn to differentiate odorants in the go-no go task

We proceeded to ask whether the strength of PAC, quantified by MI, and the

peak angle change as the animal learns to differentiate odorants in the go-no go task.

Figure 2A-F illustrates theta/high gamma PAC for all S+ and S- odorant trials in one

session for a mouse proficient in differentiating the odorants. The phase amplitude plots

for all the trials are shown in pseudocolor in Figure 2A, behavioral performance is

shown in Figure 2B, and the corresponding normalized phase amplitude plots are

shown in Figure 2C, while the mean average theta LFP is shown for one cycle in Figure

2D. There is clear (biphasic) theta phase coupling for high gamma amplitude with

maxima at ~120° and 270° for S+ (Figure 2Ai and red line in 2C). This strong PAC is

reflected by an MI with a mean of 0.02 with small trial to trial variance (Figure 2E, red

circles), and small variation in peak phase (Figure 2F, red polar histogram). In contrast,

PAC displays variable strength and phase for S- in this session (Figure 2Aii, blue line in

2E and blue polar histogram in 2F). For S- the average MI is smaller and displays large

variance ranging from 0 to 0.06 (mean MI 0.014, Figure 2E, blue circles). The peak

angle for S+ takes place slightly above 90 degrees and varies significantly for S- (Figure

2F). The MI and peak angle variance differ significantly between S+ and S- (MI:

ranksum p value < 0.001, peak angle variance: ranksum p value < 0.05, n = 50 S- trials,

49 S+ trials).

Browsing through PAC data for different sessions, plotted as in the example

shown in Figures 2A-F, raised the question whether PAC changes over the course of

learning in the go-no go olfactory discrimination task—as reported by mean MI and

peak angle variance. We noticed that in sessions where the mouse learned to

differentiate the odorants the phase of the peak angle became highly variable when the

animal became proficient. An example is shown in Figure 2-figure supplement 1 where

a mouse started the session performing close to 50% correct, and became proficient

during the last portion of the session. The phase of the peak angle is highly variable for

the last 20 S- trials. We proceeded to estimate mean MI and peak angle variance for all

mice. The mean MI and peak angle variance was estimated as the mean in all trials

within either the naïve or proficient periods for S+ and S-, calculated between 0 and 2.5

sec after odorant application. Figure 2G illustrates cumulative histograms for the mean

MI (i,iii) and peak angle variance (ii,iv) for theta/beta (i,ii) and theta/high gamma PAC

(iii,iv) for the APEBexp1 odorant for S+ and S- odorants for trial ranges when the mice

were naïve or proficient. The cumulative histograms for mean MI measured per

electrode (Figure 2Gi,iii) show small yet statistically significant changes in PAC strength

for both the theta/beta and theta/high gamma as the animal learns to discriminate

between odorants (GLM analysis, indicates that the differences in mean MI in terms of

proficiency and the event type are significant $p<0.001$, 896 degrees of freedom, d.f., for

theta/beta and theta/high gamma, 16 electrodes per mouse, 14 mice). In addition, the

circles in Figure 2Gi,iii show the per mouse mean MI. A GLM analysis did not yield

significant changes for the per mouse average MI as a function of either event type or

proficiency for theta/beta PAC ($p > 0.05$, 52 d.f., 14 mice). Per mouse average MI was

significant for both event type and proficiency for theta/high gamma PAC ($p < 0.05$, 52

d.f., 14 mice). Finally, for the APEBexp1 odorant pair we found a substantial decrease

in peak angle variance for S+ and an increase in peak angle variance for S- for both the

theta/beta and theta/high gamma PAC (Figure 2Gii,iv, GLM analysis for changes in

peak angle variance for both proficiency and event type yield $p < 0.001$, 892 d.f. for both

theta/beta and theta/high gamma, 16 electrodes per mouse, 14 mice). The changes

were also statistically significant when a GLM was computed for the average peak angle

variance per mouse (circles in Figures 2Gii,iv, GLM p value < 0.01 for theta/beta and p

< 0.001 for theta/high gamma for both event type and proficiency with 52 d.f.,14 mice).

Similar differences between S+ and S- and changes when the animal became

proficient were found for peak angle variance for the other odorant pairs (Supplemental

Information and Figure 2-figure supplement 2). Furthermore, we evaluated the changes

in PAC strength quantified as MI and peak angle variance by analyzing per-experiment

mean values (averaged over odorant pairs, Figure 2H). GLM yielded significant p values

for peak angle variance for theta/high gamma PAC for S+ vs. S- (< 0.01, 16 d.f.) and for

the interaction between S+ vs. S- and proficiency (p<0.05, 16 d.f.) (Figure 2Hiv). For a

downstream observer focused on the LFP power at the peak of PAC for the rewarded

odorant these changes in peak angle variance may reflect a mechanism by which the

unrewarded stimulus is devalued while the rewarded odorant is valued. To our

knowledge, this is the first demonstration of dynamic cross frequency interactions

changing with learning for early processing in sensory systems.

Peak phase referenced power increases for S+ and decreases for S- over learning

The existence and evolution of PAC over learning alone does not demonstrate its

potential utility to the animal in its decision-making process. In order to assess whether

high frequency LFP power at the peak of theta oscillations changes as the animal learns

we performed wavelet analysis to extract the power of high gamma (and beta) LFP

referenced to specific PAC phases of theta—peak and trough (as defined in Figure 1E).

We term this analysis phase-referenced (high-frequency) power, or PRP. PRP analysis

allows answering the question whether the power of burst high frequency LFP

evaluated at a specific phase for PAC changes as the animal learns to differentiate the

odorants in the go-no go task.

In Figures 3A-C, we show how this PRP analysis was performed. Figure 3A

shows Morlet wavelet LFP power analysis (Chery et al., 2014; David et al., 2015) for a

single S+ trial characterized by strong, persistent theta power during and after odor

delivery (related to respiration and olfactory investigation) accompanied by higher

frequency power bursts whose periodicity reflects PAC. Figure 3B shows the theta (6-14

Hz)-filtered waveform (Figure 3Bi) and the Hilbert-extracted phase (Figure 3Bii, (Tort et

al., 2010)) co-registered to the one-second epoch extracted from the wavelet LFP

power plot shown in Figure 3Aii. We proceeded to extract the value of the power (in

decibels) referenced to the peak or the trough of PAC measured in S+ trials (peak and

trough are defined in Figure 1E). The red line in Figure 3Ci shows a clear odor-induced

increase in the peak-referenced wavelet power for high gamma/theta PAC in this single

trial example (Figure 3Ci, red line). In contrast, there was little change in power

referenced to the trough of PAC (Figure 3Ci, blue line). Figure 3Cii shows the time

course for the average peak (red) and trough (blue) PRP calculated for all trials in this

go-no go session for a mouse that was performing proficiently in differentiation of the

two odorants.

In order to assess whether PRP changes with learning we performed this

analysis for all sessions for naïve and proficient periods. Figure 3D shows cumulative

histograms for the peak and trough PRP for all mice for the odor period for all go-no go

sessions with the IAAPexp2 odorant pair for theta/high gamma PAC. Over the course of

learning, peak-related high gamma power increases for S+ and decreases for S- (Figure

3D top). In contrast, the changes elicited by learning in trough PRP are smaller (Figure

3D bottom). The differences were statistically significant between peak and trough, S+

and S- and naïve and proficient (GLM p values < 0.001, 1016 d.f.). The summary of

PRP for peak and trough for theta/high gamma PAC for the odor period are shown for

all odorant pairs in Figure 3E (peak PRP) and 3F (trough PRP). For all odorant pairs

peak PRP increases for S+ over learning and decreases for S-. Trough PRP is weaker

yet follows the same trend as the animal learns to discriminate the odorant as peak

PRP (GLM p value < 0.001, 6386 d.f. for S+ vs. S-, peak vs. trough, naïve vs.

proficient). In addition, the learning-induced changes in PRP are larger for Exp2 (GLM p

value < 0.001 for experiment, 6386 d.f.). Similar changes were found for PRP for

theta/beta PAC, but there was no difference between peak and trough (not shown).

Finally, we found that theta peak phase-referenced power decreases for S- and

increases for S+ regardless of the chemical identity of the odorant. Figure 4A shows

that after the valence of the odorant is reversed the animal learns to respond to the new

rewarded odorant. We asked the question whether odorant-induced changes in peak

PRP would change when odorant valence was reversed. Figures 4B and C show that in

proficient mice peak PRP increases when the mouse responds to the rewarded odorant

regardless of the chemical identity of the odorant, and PRP decreases when the mouse

responds to the unrewarded odorant. GLM analysis yields significant changes for

odorant, reversal and the interaction of reversal and odorant (p<0.001, 1051 d.f.). These

data show that as the animal learns to differentiate between odorants in the go-no go

task there is an increase in rewarded odorant-induced peak PRP and a decrease in

unrewarded-odorant peak PRP.

Linear discriminant analysis classifies stimuli using PRP

The finding that LFP power referenced to the peak of the theta oscillation

increases for S+ and decreases for S- when the animal learns to differentiate the odors

(naïve vs. proficient) raises the question whether a downstream reader evaluating peak

power could decode which stimulus is presented. To assess decoding of olfactory

stimuli using PRP, a linear discriminant analysis (LDA) was used to set a decision

boundary hyper-plane between binary stimulus classes (S+ versus S-) (Vizcay et al.,

2015) using PRP data referenced to the peak (or the trough) of theta/beta or theta/high

gamma PAC. LDA was trained with PRP from each electrode for each mouse (16

electrodes per mouse) for all trials except one (the training dataset) and then the

missing trial (test data) was classified as S+ or S- using the PRP data from that trial.

This was performed separately for trials where the mouse was naïve or proficient to

odorant discrimination, for each 0.1 sec time point throughout the trial and was repeated

for all trials. As a control we shuffled the identity of trials in the training set. In addition,

we performed a complementary principal component analysis (PCA) of the PRP to

visualize the odor-induced divergence in the time course of the first principal component

(PC1).

Figure 5A shows for theta/beta (i and ii) and theta/high gamma (iii and iv) the

time course for PC1 calculated for S+ and S- odorants for naïve (i and iii) and proficient

(ii and iv) mice calculated with the PRP for 16 electrodes for one example odorant pair

(APEBexp1). In the naïve state PC1 diverged between S+ and S- shortly after water

reinforcement for peak and trough theta/beta PRP (Figure 5Ai) and for peak theta/high

gamma PRP (Figure 5Aiii, with a smaller change for trough). However, when the animal

became proficient PC1 diverged between S+ and S- shortly after odorant addition for

theta/beta PRP (Figure 5Aii) and theta/high gamma PRP (Figure 5Aiv). This analysis

suggests that for the proficient animal PRP LFP carries information on the rewarded

odorant.

In order to quantify the accuracy for decoding odorant identity using PRP LFP we

performed LDA analysis (Figure 5B). For naïve animals the LDA could not classify

stimulus identity using peak or trough PRP for either theta/beta (Figure 5Bi) or

theta/high gamma (Figure 5Biii). In contrast, when the animal was proficient the

precision of the LDA classifier increased shortly after odorant application from chance to

values above 70%, significantly higher than the 95% confidence interval of the LDA

performed with shuffled trials for both theta/beta (Figure 5Bii) and theta/gamma (Figure

5Biv)(GLM analysis p value for learning < 0.001, 104 d.f.). For theta/beta, both peak and

trough PRP LDA performed significantly better than shuffled data for proficient mice

(Figure 5Bii, GLM p value < 0.001, 104 d.f.). However, for theta/high gamma in

proficient animals (Figure 5Biv, bottom) LDA discriminated the identity of the stimulus

more effectively using peak PRP (GLM p value < 0.001, 104 d.f.).

The difference in LDA performance between peak and trough PRP was further

evaluated by calculating the area under the curve (AUC) for LDA performance during

the odor application window (0.5-2.5 sec after odorant valve opening). AUC is a

statistical method to quantify differences between the distributions of values of two

variables (Green and Swets, 1988). The AUC was normalized and it ranged from zero

(random decoding at 50%, Figure 5-figure supplement 1 Ai) to 1 (100% decoding during

the odor period, Figure 5-figure supplement 1 Aii). AUC is plotted for all odorant pairs

for peak and trough PRP in Figure 5C for theta/beta (i and ii) and theta/high gamma (iii

and iv) for naïve (i,iii) and proficient (ii,iv) conditions. In the proficient state, AUC for

peak PRP LDA for high gamma was significantly higher than AUC for trough PRP LDA

(Figure 5Civ, GLM for AUC peak vs. trough p value < 0.001, 104 d.f.). AUC for beta

PRP LDA did not differ between peak and trough (Figure 5Cii). Furthermore, Figure 5D

shows that the AUC was larger for Exp2 (GLM p value < 0.001 for experiment, 6386

d.f.).

Finally, we asked the question whether decoding performance differs when mice

make mistakes in the go-no go task. If PRP encodes for the chemical identity of the

odorant we would expect that decoding performance would be similar between Hit/Miss

and CR/FA during the first second of the trial when the animal keeps their head in the

odor port. If PRP encodes for contextual odorant identity decoding performance would

differ between correct responses (Hit and CR) and mistakes (Miss, FA) shortly after the

animal detects the odorant. Figure 5E shows the results of the per trial analysis of

decoding performance for LDA with theta/beta PRP (i) and theta/high gamma PRP (ii)

sorted by behavioral outcome. Interestingly, shortly after addition of the odorant there is

a divergence in decoding performance between trials with correct decision (Hit, CR) and

mistakes (Miss, FA). Shortly after addition of the odorant decoding performance for FA

drops below 50% (it tends to encode for the opposite odor), and for Miss trials the

decoder performs like the shuffled trials. Decoder performance increases above 50% for

both FA and Miss towards the end of the trial. Taken together with the shift of PRP in

the reversal experiments (Figure 4) the difference in LDA encoding between correct

trials and mistakes is further evidence that PRP encodes for the contextual identity of

the odorant (as opposed to the chemical identity). A GLM analysis indicates that there

are significant differences between all behavioral outcomes except for Hit vs. CR for

theta/high gamma (p<0.001 for all outcome pairs, except for Hit vs CR for theta/beta

PRP that has p<0.01 and Hit vs. CR for theta/high gamma that has p>0.05, 2110 d.f.).

There is no significant difference between experiments (GLM p>0.05, 2110 d.f.).

To our knowledge, this is the first demonstration of high gamma phase-

referenced LFP power developing differential information carrying capacity in early

sensory systems with learning.

Decision-making takes place at the same time for peak PRP LDA and licks

After learning that using peak PRP LDA can decode stimulus identity, we

wondered how decision-making times estimated based on either the behavioral

response of the animal in the go-no go task (licks) or classification of the stimulus by

PRP LDA relate to each other. In addition, we asked whether the decision-making time

calculated with peak PRP LDA differed from decision-making time estimated using

trough PRP LDA. Decision-making time was defined as the time when there is a

statistically significant difference between S+ and S- in odorant classification (estimated

separately for licks and LDA).

Figures 6A-B show examples of licks by a mouse in the go-no go task for the

EAPAexp2 odorant pair. Figure 6A shows the licks detected at the waterspout for 25 S+

trials when the animal was performing proficiently while Figure 6B shows the lick traces

for 25 S- trials in the same session. As in previous studies (Doucette and Restrepo,

2008), we estimated the p value for lick decision-making by computing in each 0.1 sec

time bin the ranksum test for the difference between licks (scored 0 for no lick vs. 1 for

lick). An example of the time course of the p value for licks for mice performing the go-

no go task for EAPAexp2 is shown by the green line in Figure 6Ci. The p value drops

below 0.05 ~250 msec after odorant addition. We defined the lick decision-making time

as the time point when the p value for difference in licks between S+ and S- falls and

stays below 0.05 after odor application.

LDA decision-making time was estimated in a similar manner by computing the p

value for a ranksum test for the difference between predictions (scored 1 for correct

prediction vs. 0 for incorrect prediction) for PRP LDA decisions evaluated after pooling

data for all mice for each odor pair. As explained under supplemental information

(Figure 6-figure supplement 1) pooled animal data was used as opposed to per animal

PRP LDA because this resulted in consistent estimation of earlier decision-making

times. Examples of the time courses of the p values for PRP LDA performing the go-no

go task for EAPAexp2 are displayed in Figure 6Ci for theta/high gamma PAC (red line:

peak, blue line: trough). Before odorant presentation the p value fluctuates and

sometimes falls below 0.05 transiently, while after odor presentation the p values for

peak PRP fall below 0.05 sharply and continuously at approximately the same time as

lick p value (~250 msec). Trough PRP p value for LDA decoding performance drops

below 0.05 later (~950 msec) and the p values are higher than peak p values (compare

red and blue lines in Figure 6Ci). Thus, in this example lick decision-making time is

similar for licks and peak PRP LDA, and is slower for trough PRP LDA.

We proceeded to calculate decision-making times for licks and peak PRP LDA

for all odorant pairs. Decision-making times for LDA decoding performance via peak

PRP and lick for all odorant pairs for pooled mouse data are shown in Figure 6D for

naïve (Figure 6Di) an proficient (Figure 6Dii) mice. Decision times were reduced

significantly with task learning (GLM p value < 0.001, 44 d.f.). However, decision-

making times did not differ between lick and peak PRP LDA (GLM p value > 0.05, 44

d.f.).

Dimensionality of PRP LFP space decreases when the animal learns to

discriminate the odorants

Figure 7A shows the change in the scatter pattern of the first two principal

components (PCs) of the theta/high gamma peak PRP calculated for pooled data from

all mice as the animals learn to differentiate the two odorants for the EAPAexp1 odorant

pair. Figure 7Ai shows that when the mice are naïve S+ (red) and S- (blue) points are

mixed in the 2D PC space before and during odorant application. In contrast, Figure 7Aii

shows that the 2D PC patterns become clearly separable when the animals are

proficient. There appears to be a change in the geometrical arrangement of the points in

principal component space suggesting that there is a change in dimensionality as the

animal learns to differentiate the odorants (Litwin-Kumar et al., 2017).

In order to provide a more nuanced analysis of whether the number of

independent signals in the PRP LFP data measured by the 16 electrodes changes as

the animals learn to discriminate between odorants we calculated a quantitative

measure that characterizes dimensionality in the PRP LFP space (see Methods). The

16 electrodes are closely spaced (12.5 μm diameter electrodes in the same tetrode are

directly abutted against each other and tetrodes are separated by 100-200 μm in both

preparations: Exp1 with optetrodes and Exp2 with tetrodes). Because of this close

proximity the extracellular field potentials would not be expected to differ greatly (Gold

et al., 2006) and therefore the dimensionality of the PRP LFP space would be expected

to be close to one. Interestingly, when the dimensionality was calculated on a per

mouse basis where the maximum dimensionality (M) is 16 we found that the average

dimensionality for theta/high gamma PRP LFP for electrodes in Exp2 was indeed close

to one, but it was significantly higher (2-4) in Exp1, with the highest dimensionality for

PRP LFP trough (Figure 7-figure supplement 1 Ai, GLM p value < 0.001 for peak vs.

trough and Exp1 vs. Exp2, 379 d.f.). We found similar differences in dimensionality

between Exp1 and Exp2 for theta/beta PRP LFP, but in this case there was no

difference between peak and trough dimensionality (Figure 7-figure supplement 1 Aii,

GLM p value < 0.001 for Exp1 vs. Exp2 and >0.05 for peak vs. trough, 379 d.f.). This

indicates that even when measured using electrodes located within 100-200 μm the

LFP carries information on multiple independent components.

Interestingly, the dimensionality for peak PRP LFP for theta/high gamma

calculated on a per mouse basis decreased after addition of the odorant for both

experiments and the decrease in dimensionality was larger for proficient mice.

Examples of the dimensionality time course for peak PRP LFP for theta/high gamma

calculated on a per mouse basis are shown for two odorant pairs in Figure 7-figure

supplement 1 Bi (Exp1) and Figure 7-figure supplement 1 Bii (Exp2). A GLM analysis of

the dimensionality for peak PRP LFP calculated on a per mouse basis for all odorant

pairs indicates that there are significant differences in theta/high gamma peak PRP LFP

dimensionality for proficient vs. naïve, time course, experiments, and for interactions of

these factors (except for the interaction of experiments x proficient vs. naïve) (GLM p

values < 0.001, 6098 d.f.). For theta/beta peak PRP LFP dimensionality we found

similar results with the exception that there was no difference between peak and trough

(Figure 7-figure supplement 1 Ci and Cii, GLM p values < 0.001, 6098 d.f.).

Finally, when the dimensionality for peak PRP LFP was calculated for LFPs

recorded from all mice the difference in the decrease in dimensionality between naïve

and proficient became more evident. Time courses for pooled mouse peak PRP LFP

dimensionality for theta/high gamma are shown in Figure 7B for Exp1 (7Bi) and Exp2

(7Bii). Peak theta/high gamma PRP LFP dimensionality decreases markedly when the

odorant is delivered for both Exp1 and Exp2 for proficient mice and this decrease is

larger for proficient compared to naïve mice (compare naïve vs. proficient in Figures 7Bi

and ii). A GLM analysis indicates that there are significant differences for time course,

proficient vs. naïve, experiments and peak vs. trough, and for interactions of these

factors (GLM p values < 0.001, 1688 d.f.). For theta/beta peak PRP LFP dimensionality

for pooled mice we found similar results (Figures 7Ci and 7Cii), but there was no

difference between peak and trough. GLM analysis found statistically significant

differences for time course, and proficient vs. naïve, and for interaction of these factors

(GLM p values < 0.001, 1688 d.f.), and a statistically significant difference between

experiments (GLM p value < 0.01, 1688 d.f.). However, was no difference between peak

and trough (GLM p value >0.05, 1688 d.f). Therefore, for both the per animal and the

pooled animal PRP LFP space there is a decrease in dimensionality as the animal

learns to discriminate between odorants, but this difference is more evident for the

pooled animal data. This is evidence for a learned tightening of dimensionality for the

odor representation manifold (defined as the independent neural components

representing the odor).

Discussion

We asked whether information carried by high frequency oscillations in brain

regions involved in early sensory processing changes when observed at different

phases of theta frequency oscillations. Specifically, we asked whether information on

contextual odorant identity conveyed by LFP power of high gamma and beta frequency

oscillations of the OB differs when the downstream observer filters the input through

different theta LFP phase windows. We focused on the OB where it has been

suggested that amplitude modulation may be used by higher-order processing centers

such as PC to decode stimulus identity (Freeman and Schneider, 1982; Heck et al.,

2019). We found that identity could not be decoded from beta or high gamma OB LFP

power when the mouse was naïve to the identity of the reinforced odorant (left panels in

Figure 5B). However, the contextual identity of the odorant could be decoded from the

PRP after the animal learned to discriminate between the odorants (right panels in

Figure 5B). Furthermore, decoding was more accurate when the high gamma LFP

power was observed at the peak of the theta oscillation, but it was similar at both the

peak and trough phases for beta LFP oscillations. LDA correctly identified the stimulus

with a time course for decision-making similar to the behavioral readout (licking on the

water-delivery spout, Figure 6). Thus, stimulus representation by theta phase-

referenced beta and high gamma oscillations of the OB evolves over the course of

learning. Importantly, our findings were replicated in two different experiments (Exp1

and Exp2) differing in genotype, electrode location, device implanted (tetrode or

electrode) and odorant pairs used (see Table S1). We did find differences in measured

parameters between experiments, and these are likely due to methodological

differences between Exp1 and Exp2 such as electrode location and genotype.

Regardless of the reason for differences between experiments it is important for this

study that the findings on PRP are found for both experiments under substantially

different experimental conditions. To our knowledge this is the first demonstration that

decoding of stimulus identity from theta phase-referenced high gamma oscillations,

reflecting synchronized neuronal firing, changes with learning in brain areas involved in

early sensory processing.

The importance of PAC in information processing is based on the notion that

different information is carried at different phases of the slow carrier oscillation. This

discretization of information is a potential mechanism for encoding short-term memories

(Lisman and Idiart, 1995) and more broadly as a neural communication protocol, the

theta-gamma neural code (Hopfield, 1995; Lisman and Jensen, 2013). Circuit activation

at theta frequency was discovered to be essential for long-term potentiation in the rabbit

hippocampus (Bliss and Lomo, 1973), arguing that neural information transfer may go

beyond mean rate codes and involve spike timing with respect to oscillatory

timekeepers. Subsequent studies indicated that temporal windows set by theta cycles

allow for local circuit interactions and a considerable degree of computational

independence in subdivisions of the entorhinal cortex-hippocampal loop (Mizuseki et al.,

2009). For CA1, it was hypothesized that theta partitions the encoding of new

information versus the retrieval of stored information (Hasselmo et al., 2002). Indeed,

optogenetic recruitment of fast spiking inhibition in CA1 delivered at specific theta

phases altered murine performance in a spatial navigation task (Siegle and Wilson,

2014). When the mice were encoding a location, triggering of inhibition of CA1 at theta

peaks improved performance, whereas inhibition at theta troughs improved

performance during the retrieval phase. Recently, age related cognitive decline in

working memory in humans was reversed temporarily via transcranial alternating-

current stimulation frequency-tuned to the endogenous theta peak frequency of each

individual's frontotemporal network (Reinhart and Nguyen, 2019). It appears that

working memory is undergirded by theta-gamma PAC and that inter-regional

communication is facilitated by theta phase coherence (Daume et al., 2017; Reinhart

and Nguyen, 2019). Thus, the theta LFP which has been cited as one of the most global

oscillations in the brain acts as a timekeeper (Siegle and Wilson, 2014) that is coherent

across numerous cortical and subcortical structures arguing for its role in transfer of

discrete chunks of information (Buzsaki, 2002).

Here we asked whether odor identity information was carried at different phases

of theta. Our study of PAC in the OB involved learning of odor discrimination in the go

no-go task. In orbitofrontal cortex, Van Wingerden et al. (van Wingerden et al., 2014)

showed that this task elicited strongest theta/gamma PAC during odor sampling

preceding correct decision-making, which they attributed to temporal coordination

between cell assemblies responsible for stimulus representation and reward

association. The OB circuit is characterized by beta and gamma oscillations nested in a

slower theta carrier wave pertaining to respiration (Buonviso et al., 2003; Heck et al.,

2019; Kay, 2005; Kepecs et al., 2006; Rebello et al., 2014; Rojas-Libano et al., 2014).

However, Granger directionality analysis shows that the theta oscillation in the OB is

also influenced by hippocampal slow oscillations indicating that the origin of OB slow

carrier oscillations is complex, involving both respiration and centrifugal input from

downstream brain regions whose relative contribution likely changes under different

behavioral states (Nguyen Chi et al., 2016). Cyclical activation of OSNs through

respiration (Adrian, 1950; Bressler, 1988; Rosero and Aylwin, 2011; Zhuang et al.,

2019) provides input to mitral/tufted (M/T) cells, and coordinated activity by M/T cells in

the OB caused by mitral cell-granule cell reciprocal dendrodendritic interactions

establishes gamma oscillations (Arnson and Strowbridge, 2017; Osinski and Kay, 2016;

Pouille and Schoppa, 2018; Schoppa, 2006; Stopfer et al., 1997)(see review by (Heck

et al., 2019)). Gamma LFP is thought to represent the stimulus identity (Bathellier et al.,

2006; Beshel et al., 2007; Li and Cleland, 2017; Rojas-Libano and Kay, 2008) and beta

LFP oscillations (15-35 Hz) have been linked to learned reward (Martin et al., 2006).

However, the classification of these two bandwidths as carrying learning vs. odor

identity information is not always appropriate, and Frederick et al. (Frederick et al.,

2016) proposed that they represent different cognitive states that vary depending on

particular behavioral demands. PAC is present in the OB (Buonviso et al., 2003; Rojas-

Libano et al., 2014) and it may be of great albeit relatively unexplored utility. We

contributed to the understanding of which information is carried by different frequency

oscillations by asking whether the high gamma and beta oscillations carry different

information when they are observed within particular phase windows of the theta carrier

wave (Figure 5). We hypothesized that PAC may be used to convey olfactory stimulus

information to theta phase-synchronized downstream regions like PC (Kepecs et al.,

2006; Lisman and Buzsaki, 2008).

We demonstrate that PAC changes in the OB during associative learning (Figure

2). Peak angle variance increased for the unrewarded odorants with learning. The theta

phase-referenced PRP of the high gamma and beta LFPs also follows this trend (Figure

3), increasing for S+ and decreasing with S- over learning. The changes in PRP

correlating with learning may reflect contrast enhancement that facilitates the

discrimination between similar stimuli by a downstream observer. Neuronal adaptation

to facilitate contrast enhancement has been shown at the level of the OSNs (Haney et

al., 2018), the glomerular layer (Cavarretta et al., 2016), and the granule cell layer

(Adams et al., 2019) of the OB. We speculate that the concurrent increase in PRP for

S+ and decrease in PRP for S- reflects adaptational mechanisms in the OB to enhance

contrast between similar stimuli with different associated values.

To quantify whether these changes in PAC increased discriminability between

odorants using PRP we asked whether the stimulus that was delivered could be

decoded from PRP data. Previous investigations have decoded olfactory stimuli from

spike counts aligned to sniffing in the mouse (Cury and Uchida, 2010; Gschwend et al.,

2012), in the antennal lobe of the locust (Stopfer and Laurent, 1999), and M/T neural

activity assayed through calcium transients in mice (Chu et al., 2016), but trial

classification via PAC remains unexplored. Here, we performed LDA on PRP data to

determine whether a single hyperplane could differentiate between stimuli as the

animals learned the task. Figure 5B indicates that the identity of the stimuli cannot be

decoded from PRP recorded in the naïve mice, but that the stimuli can be decoded

when the mice learn to discriminate the odorants. Furthermore, decoding the stimuli

from peak high gamma PRP is significantly better than trough PRP. Thus, decoding of

the odorant identity by downstream observer using a phase window for the slow carrier

oscillation could detect the identity of the odorants after the animal learns to differentiate

between odors in the go no-go task. Finally, this study contributes to the mounting

evidence that the OB is an important locus for learning-related activity changes

(Abraham et al., 2014; Doucette et al., 2011; Doucette and Restrepo, 2008; Koldaeva et

al., 2019; Li et al., 2015).

We wondered how timing for decision-making would relate comparing stimulus

decoding with PRP with decision-making estimated from the behavioral readout of the

olfactory-driven decision in the go no-go task (differential licking between the two

odorants). The time course for the behavioral decision, reflected in the graph of p value

for differential licks, is overlaid by the p value time course for peak PRP decoding

(Figure 6C, right) with trough PRP lagging behind. Neither beta nor high gamma peak

PRP p values are discriminable from lick p values in terms of decision time in this task.

As shown in Figure 6D, for proficient animals licking differs between odorants within

about 0.5 seconds from stimulus onset, at the same time that PRP decoding becomes

significant. This is slower than discrimination times for odorant responses recorded in

mice performing two alternative forced choice odorant discrimination or receiving odors

passively where discrimination takes place within the first respiratory cycle (~250 ms)

after stimulus onset (Cury and Uchida, 2010; Short and Wachowiak, 2019; Shusterman

et al., 2011). However, the 0.5 sec decision-making time is consistent with studies with

the go no-go task where there is not a strong motivation for the animal to stop licking for

the unrewarded odorant (Abraham et al., 2004; Doucette and Restrepo, 2008; Jordan et

al., 2018; Rinberg et al., 2006)—with the exception of rapid detection of optogenetic

activation of glomerular input in the go no-go task (Smear et al., 2011). Thus, we

conclude that PRP is an accurate reflection of decision-making in olfactory

discrimination.

Interestingly, in contrast with theta/high gamma PRP where the decoding

performance for peak and through differ substantially, for theta/beta PRP there is no

difference in decoding performance between peak and trough (compare Figures 5Bii

and 5Biv). Increases in OB beta LFP for rodents engaged in odorant discrimination

tasks take place after the onset of changes in gamma LFP and it has been postulated

that beta LFP represents systemwide coherent states perhaps conveying centrifugal

modulation from downstream brain areas to the OB (David et al., 2015; Frederick et al.,

2016; Kay, 2014). If beta oscillations indeed represent centrifugal feedback our findings

suggest that centrifugal modulation does not differ through phases of the theta

oscillation and would therefore excerpt modulation on high frequency circuit activity

occurring at different theta phases. Here we show that peak-referenced high gamma

oscillation power encodes for the contextual identity of the odorant (is the odorant

rewarded?). What information could be conveyed at other theta phases is an open

question. Perhaps asynchronous M/T firing carries chemical odorant information in the

theta trough.

Finally, we turn to dimensionality. The measurements in the experiments

presented here are inherently low dimensional because: 1) The animal is discriminating

between two odors, and the behavior is constrained by the go no-go task (Gao et al.,

2017) and 2) While the number of independent degrees of freedom for inputs to the OB

is potentially large (~2000 glomeruli), the four tetrodes are close together and they

should sample a small subset of these input dimensions. Indeed, because the tetrodes

are within 200 µm of each other (and electrodes are adjacent within tetrodes), the LFP

recorded from each electrode is likely to be similar, assuming that the cellular milieu

around the electrodes does not separate the inputs (Gold et al., 2006). Interestingly, the

dimensionality measured from the 16 electrodes (four tetrodes) varies between 1 and 6

suggesting that subsets of electrodes are detecting different signals (Figure 7-figure

supplement 1 A). However, when the mice learn to discriminate the dimensionality

decreases in the presence of the relevant odorant (Figures 7 and Figure 7-figure

supplement 1 B and C), suggesting that population representation is a modifiable

aspect of OB processing. This likely reflects experience-dependent, attention (Canas

and Jones, 2010; Niv et al., 2015). In the naïve state, relevant (rewarded) dimensions

are unknown to the mouse, and a downstream structure sampling from the recorded

population would receive ambiguous signaling in the presence of the rewarded odorant

without decoding or reducing the input information. We find a large decrease in

dimensionality post-learning, suggesting that OB now enhances a relevant and invariant

signal to allow for unambiguous odor detection. Attention, guided by reward prediction

error, likely biases/constrains the learning to the relevant dimensions (Leong et al.,

2017)—in this task, odor identity.

The changes in signal processing elicited by learning, habituation, and passive

odorant exposure in the OB, the earliest processing center in the olfactory system, are

remarkable. Why is processing in this circuit so plastic and influenced by centrifugal

input from downstream targets, such as PC, and from a variety of neuromodulatory

centers (Gire et al., 2013b; Linster and Cleland, 2016)? Likely the reason is that this

processing center receives massive parallel glomerular input and projects M/T output of

high dimensionality to downstream brain processing areas (Chae et al., 2019).

Furthermore, unsupervised learning modifying cellular interactions in the OB would

increase dimensionality (Hiratani and Latham, 2019). Therefore, odors must be

identified in the midst of a high dimensionality cocktail party (Rokni et al., 2014). Based

on the present data showing a large increase in the accuracy for decoding of contextual

odor identity from the PRP as the animal learns to differentiate the odorants we

speculate that when a subset of the degrees of freedom of the sensory input become

behaviorally relevant, centrifugal feedback elicits supervised learning that results in a

decrease in dimensionality of the output from the OB to downstream brain areas. Thus,

the fact that peak theta-referenced high gamma power encodes information on whether

the odorant is rewarded may simply reflect the fact that the transfer of sensory

information on the rewarded odorant is being prioritized in the OB circuit. This would

result in more accurate discrimination of the input channels that are relevant to the

current task. Whether (and how) supervised learning-induced OB circuit plasticity

results in a decrease in dimensionality of the output remains to be determined.

Materials and Methods

Key Resources Table

REAGENT TYPE	REAGENT or RESOURCE	SOURCE	IDENTIFIER	ADDITIONAL INFORMATION
Chemical compound, drug	Isoamyl acetate	Sigma-Aldrich	Cat#123-92-2	Odorant
Chemical compound, drug	Ethyl acetate	Sigma-Aldrich	Cat# 141-78-6	Odorant
Chemical compound, drug	Propyl acetate	Sigma-Aldrich	Cat# 109-60-4	Odorant
Chemical compound, drug	Mineral oil	Sigma-Aldrich	Cat# 8042-47-5	Odorant
Chemical compound, drug	Acetophenone	Sigma-Aldrich	Cat# 98-86-2	Odorant
Chemical compound, drug	Ethyl benzoate	Sigma-Aldrich	Cat# 93-89-0	Odorant
Other	Nickel-chrome wire	Sandvik	Cat#PX000002	Tetrode fabrication
Other	16 Channel Electrode Interface Board	Neuralynx	EIB-16	Tetrode fabrication
Software, algorithm	Analysis code	This paper	Available on github and by request	https://github.com/restrepd/drgMaster
Strain, strain background	Mouse: C57BL/6J	Jackson Lab	RRID: IMSR_JAX:000664	Male mice
Strain, strain background	Mouse: Tg(Dbh-cre)KH212Gsat/Mmucd	Mutant Mouse Resource and Research Center	032081-UCD, RRID:MMRRC	All sexes used
Strain, strain background	Mouse: eNpHR3.0, 129S-Gt(ROSA)26Sortm39(CAG-hop/EYFP) Hze/J	Jackson Lab	014539, RRID:SCR_004633	All sexes used
Software, algorithm	MATLAB_R2018a	Mathworks	RRID: SCR_001622	
Software, algorithm	Illustrator	Adobe	RRID: SCR_010279	
Software, algorithm	Photoshop	Adobe	RRID: SCR_014199	

Animals

The study was performed in two sets of experiments. For the first set of

experiments (Exp1), DBH-eNpHR3.0 mice were used as per Ramirez-Gordillo et al.,

2018. These mice express halorhodopsin in cells expressing Cre under the *Dbh*

promoter (DBH-Cre eNpHR3.0) in LC-NA neurons and were produced by crossing

DBH-Cre mice (032081-UCD, Mutant Mouse Resource and Research Center) with

mice expressing halorhodopsin in a Cre-dependent fashion [eNpHR3.0,

129S Gt(ROSA)26Sortm39(CAG-hop/EYFP) Hze/J, Jackson labs]. C57BL/6 mice

(Jackson stock number: 000664) were used for the second set of experiments (Exp2).

All animal procedures were performed under approval from the Institutional Animal Care

and Use Committee (IACUC) at the University of Colorado Anschutz Medical Campus

under guidelines from the National Institutes of Health. The number of mice used in

each experiment is shown in Table S1.

(Op)tetrode implantation

Surgery was performed under approval from the Institutional Animal Care and

Use Committee (IACUC) at the University of Colorado Anschutz Medical Campus, using

aseptic technique. As per Li et al. (2015), tetrode boards (EIB-16, Neuralynx) with

optional fiber optic cannula for photostimulation (optetrodes) were populated with four

tetrodes consisting of four 12.5 μm nichrome wires coated with polyimide (Sandvik

RO800). For optetrode fabrication the tetrode boards received the addition of a fiber

optic cannula 105 μm in diameter (Thorlabs FG105UCA) with 1.25 mm OD ceramic

ferrule (Precision Fiber Products) for coupling to a DPSS laser (473 nm, Shanghai

Laser and Optics Century). The optogenetic experiments of Ramirez-Gordillo et al.

(2018) are not included in this study. In the rest of the text we refer to the inserts as

(op)tetrode with the understanding that optetrodes were implanted for Exp1 while

tetrodes were implanted for Exp2. Electrode tips were electroplated to 0.2-0.4 MΩ

impedance.

Two-month-old male mice were anesthetized with 5% isoflurane in oxygen.

Intraperitoneal ketamine/xylazine (100 mg/kg and 10 mg/kg respectively) was then

administered along with 100 µl of 2% lidocaine injected subcutaneously over the skull.

After the mouse was found to be unresponsive to a toe pinch, the animal's head was

then secured in the stereotaxic apparatus (Narishige SR-5M-HT) and the skull was

leveled (≤ 50 µm difference DV between bregma and lambda). Gentamycin ophthalmic

ointment was applied to the eyes to maintain hydration. After incising the skin overlaying

the skull, the periosteum was cleared with 15% H_2O_2. A manipulator (Sutter MP-285)

was zeroed at bregma and midline and the target location for OB implantation was

marked with respect to bregma (Exp1: AP +4.28 mm, ML +0.4 mm, Exp2: AP +4.28

mm, ML +0.5 mm).

A craniotomy performed at this site (Marathon III drill) exposed dura mater which

was removed prior to implantation. Another craniotomy was performed more caudally

for implantation of one ground screw (Plastics1 00-96 x 1/16). The (op)tetrode was

positioned above the craniotomy over the OB while the ground wire was wrapped

around the ground screw with the connection coated in silver paint (SPI Flash-Dry silver

conductive paint). After securing the ground screw to the skull, the (op)tetrodes were

lowered into position at the rate of 1 mm/minute (Exp1: AP +4.28 mm, ML +0.4 mm, DV

0.53 mm, Exp2: AP +4.28 mm, ML +0.5 mm, DV 1.0 mm). After reaching the target

depth, the optetrode was adhered to the skull with C&B Metabond, followed by Teets

'Cold Cure' dental cement. After curing (10 minutes), the (op)tetrode was detached from

the manipulator, the animal was removed from the stereotax and received

subcutaneous injections of carprofen (10 mg/kg) and buprenorphine (0.05 mg/kg) and

recovered on a heating pad kept at 37°C. The mice were monitored daily and received

additional carprofen injections daily for the first two days postoperatively.

Go-no go behavioral task

Mice were water deprived until they reached 80% normal body mass. Then they

were placed into the Slotnick olfactometer (Bodyak and Slotnick, 1999; Li et al., 2015)

chamber where they could move freely. All mice were first trained to lick the water spout

to obtain water in the presence of odor (1% isoamyl acetate in mineral oil, v/v) in the

"begin" task (Slotnick and Restrepo, 2005). Subsequently they learned to discriminate

1% isoamyl acetate (S+) versus mineral oil (S-) in the "go no-go" task (Doucette et al.,

2011; Li et al., 2015), followed by learning to discriminate other odorant pairs (Table

S1). Data were analyzed for the go-no go discrimination task, but not for the begin task.

Mice self-initiated trials by poking their head into the odor delivery port, breaking

a photodiode beam (Figure 1A). During reinforced odorant delivery (lasting 2.5 seconds)

they must lick a water delivery spout at least once during each of four 0.5 second-long

response areas in order to register the decision as a Hit (Figure 1B). Licks were

detected as electrical connectivity between the water spout and the ground plate on

which they stand (Slotnick and Restrepo, 2005). If the mice licked during a rewarded

odorant trial, they received ~10 µl water reinforcement (Figure 1C). The mice learn to

refrain from licking for the unrewarded odorant due to the unrewarded effort of

sustained licking. For correct rejections mice leave the spout shortly after the last lick

that takes place 0.5-1.8 sec after odorant onset (Figure 6B). Performance was

evaluated in blocks of 20 trials, with 10 S+ and 10 S- trials presented at random.

Animals performed as many as 10 blocks per session. Sessions were terminated when

animals demonstrated satiety/disengagement from the task or when they performed at

or above 80% correct discrimination in three or more blocks in a session. For

EAPAexp1, IAAPexp2 and EAPAexp2 once criterion was reached, the next session had

reversed valence, meaning that the previous S+ became S- (and the previous S-

became S+). This reversal disambiguates the identity of the odor from its valence. Data

were analyzed for all odorant pair sessions, including isoamyl acetate vs. mineral oil.

Data were analyzed within two performance windows: when the animal was performing

below 65% (naïve) or above 80% (proficient). The data were also analyzed in the 65-

80% performing window, and the results fell between the naïve and proficient windows

(data not shown).

Odor stimulus delivery time was measured with a photoionization detector

(miniPID, Aurora Scientific). Figure 1-figure supplement 1 shows the time course for

odorant concentration measured at the odor spout. The time difference between valve

opening and detection of odor at the odor port was between 66 – 133 ms, depending on

which olfactometer was used.

Neural recording

Extracellular potentials from the four tetrodes were captured and digitized at 20

kHz on the RHD2216 amplifier of the Intan RHD2000 Evaluation System with a 1-750

Hz bandpass filter. Information on behavioral events (valve times, mouse presence at

the odor port) was sent through a digital line from the Slotnick olfactometer to the Intan

board. Licks detected by the Slotnick olfactometer were recorded as an analog signal by

the Intan board.

Phase Amplitude Coupling

PAC data were processed using the Hilbert transform method described by Tort

et al., 2010. Briefly, data were bandpass filtered with a 20th order Butterworth filter using

Matlab's filtfilt function with zero phase shift to extract LFP in the low frequency

oscillation used for phase estimation (theta, 2-14 Hz, Figure 1Diii) and the high

frequency oscillation used for estimation of the amplitude of the envelope (either beta,

15-30 Hz, or high gamma, 65-95 Hz, Figure 1v). Hilbert transform established the theta

phase (Figure 1Div) and, separately, the envelope for beta or high gamma (red line in

Figure 1Dv). A plot of the distribution of beta/high gamma amplitude in 51 bins of the

phase of theta (PAC) was plotted to gauge the strength of PAC (Figure 1E). To quantify

the strength of PAC we calculated the MI estimating the KL distance to quantify the

difference between the observed beta/high gamma amplitude distribution along the

phase of theta from a uniform distribution. If PAC is non-existent, MI = 0, meaning the

mean amplitude is distributed uniformly over theta phases, and if PAC is a delta function

MI = 1. MI for signals measured in brain areas such as the hippocampus typically fall

between 0-0.03 (Tort et al., 2010).

Phase-referenced LFP power

Beyond the mere existence of PAC that has been documented in numerous brain

regions, we sought to evaluate whether the information carried by beta or high gamma

power at different theta phases could be used to discriminate between olfactory stimuli.

To accomplish this analysis, we developed the PRP approach using custom Matlab

code. Briefly, PAC was calculated using the approach documented by Tort *et al.*, 2010,

as described above and summarized in Figure 1. Peak and trough theta phases are

defined as the phase for maxima and minima of the PAC distribution measured for the

S+ trials (red plot in Figure 2C). A continuous Morlet wavelet transform was used to

estimate the power for the high frequency oscillations (Buonviso et al., 2003). PRP was

estimated as the power of the high frequency oscillations (beta or high gamma)

measured at the peak or trough of PAC (Figure 5). The Matlab code used for data

analysis has been deposited to https://github.com/restrepd/drgMaster.

PAC/PRP Simulation

To validate our analysis of PAC and PRP, we performed simulations with

uncoupled oscillations (no PAC) with a low frequency cosine oscillation at 8 Hz and a

high frequency cosine oscillation at 40 Hz (Figure 1-figure supplement 2A) or with

coupled oscillations where high frequency bursts took place at 180 degrees for the low

frequency cosine oscillation (Figure 1-figure supplement 2B). The bursts were

generated by a cosine oscillation at the high frequency multiplied by gaussians with a

full width half maximum of (1/8 Hz)/2 centered at 180 degrees of the low frequency

cosine wave. Top panels in both S1A and S1B are pseudocolor Morlet wavelet power

time courses, while the bottom panels show the PRP of the 40 Hz oscillation. For the

case with no PAC (Figure 1-figure supplement 2A), peak and trough PRP are equal.

However, for the PAC example (Figure 1-figure supplement 2B) the peak PRP is

significantly higher than trough PRP.

Statistical analysis

Statistical analysis was performed in Matlab. PAC parameters and PRP were

calculated separately per electrode (16 electrodes per mouse) for all electrodes per

mouse. Statistical significance for changes in measured parameters for factors such as

learning and odorant identity was estimated using generalized linear model (GLM)

analysis, with post-hoc tests for all data pairs corrected for multiple comparisons using

false discovery rate (Curran-Everett, 2000). The post hoc comparisons between pairs of

data were performed either with a t-test, or a ranksum test, depending on the result of

an Anderson-Darling test of normality. GLM is a general statistical method that includes

regression and analysis of variance. Degrees of freedom and statistical significance

have the same meaning in GLM as in analysis of variance and regression (Agresti,

2015). In addition, as a complementary assessment of significant differences (Halsey et

al., 2015) we display 95% confidence intervals (CIs) shown in the figures as vertical

black lines were estimated by bootstrap analysis of the mean by sampling with

replacement 1000 times.

Linear discriminant analysis

Classification of trials using PRP was accomplished via LDA in Matlab whereby

PRP for every trial except one were used to train the LDA, and the missing trial was

classified by the LDA prediction. This was repeated for all trials and was performed

separately for peak and trough PRP, and for analysis where the identity of the odorants

was shuffled. LDA and dimensionality analysis were performed either on a per-mouse

basis where the input was the PRP recorded from 16 electrodes, or on pooled mouse

data where the input was the PRP recorded from 16 x N electrodes where N is the

number of mice. For pooled mouse analysis a pooled response vector was therefore

created by concatenating across animals and the number of trials *n*, was determined by

the session with the lowest number of trials for a single odorant (Chu et al., 2016).

Dimensionality

Following Litwin-Kumar et al. (2017) we defined the dimension of the system

(dim) with *M* inputs as the square of the sum of the eigenvalues of the covariance matrix

of the measured PRP LFP (*pLFP*) divided by the sum of each eigenvalue squared:

935
$$\dim (pLFP) = \left(\sum_{i=1}^{M} \lambda_i \right)^2 / \left(\sum_{i=1}^{M} \lambda_i{}^2 \right)$$

where λ_i are the eigenvalues of the covariance matrix of *pLFP* computed over the

distribution of PRP LFP signals measured in the OB. If the components of *pLFP* are

independent and have the same variance, all the eigenvalues are equal and dim(*pLFP*)

= *M*. Conversely, if the *pLFP* components are correlated so that the data points are

distributed equally in each dimension of an *m*-dimensional subspace of the full *M*-

dimensional space, only *m* eigenvalues will be nonzero and dim(*pLFP*) = *m*.

Acknowledgments: We thank Ms. Nicole Arevalo for animal husbandry, Ms. Dnate'

Baxter for laboratory support, Dr. Douglas Curran-Everett for advice on statistical

analysis, Dr. Cecilia Gauthier Umaña for discussions on wavelet analysis and Ms.

Arianna Gentile-Polese for immunohistochemical assays. This research was supported

by NIDCD 1F31DC016483-01A1 to J.L. and NIDCD 5R01DC000566-30 to D.R.

Declaration of interests: The authors declare no competing interests.

References

References

Abraham, N.M., Spors, H., Carleton, A., Margrie, T.W., Kuner, T., and Schaefer, A.T.

(2004). Maintaining accuracy at the expense of speed: stimulus similarity defines odor

discrimination time in mice. Neuron *44*, 865-876.

Abraham, N.M., Vincis, R., Lagier, S., Rodriguez, I., and Carleton, A. (2014). Long term

functional plasticity of sensory inputs mediated by olfactory learning. Elife *3*, e02109.

Adams, W., Graham, J.N., Han, X., and Riecke, H. (2019). Top-down inputs drive

neuronal network rewiring and context-enhanced sensory processing in olfaction. PLoS

Comput Biol *15*, e1006611.

Adrian, E.D. (1950). The electrical activity of the mammalian olfactory bulb.

Electroencephalography and Clinical Neurophysiology *2*, 377-388.

Agresti, A. (2015). Foundations of linear and generalized linear models. In Wiley series

in probability and statistics (Hoboken, New Jersey: John Wiley & Sons Inc.,), pp. 1

online resource.

Amemiya, S., and Redish, A.D. (2018). Hippocampal Theta-Gamma Coupling Reflects

State-Dependent Information Processing in Decision Making. Cell Rep *25*, 3894-3897.

Arnson, H.A., and Strowbridge, B.W. (2017). Spatial Structure of Synchronized

Inhibition in the Olfactory Bulb. J Neurosci *37*, 10468-10480.

Bathellier, B., Lagier, S., Faure, P., and Lledo, P.M. (2006). Circuit properties

generating gamma oscillations in a network model of the olfactory bulb. J Neurophysiol

95, 2678-2691.

Beshel, J., Kopell, N., and Kay, L.M. (2007). Olfactory bulb gamma oscillations are

enhanced with task demands. J Neurosci *27*, 8358-8365.

Bliss, T.V., and Lomo, T. (1973). Long-lasting potentiation of synaptic transmission in

the dentate area of the anaesthetized rabbit following stimulation of the perforant path. J

Physiol *232*, 331-356.

Bodyak, N., and Slotnick, B. (1999). Performance of mice in an automated olfactometer:

odor detection, discrimination and odor memory. ChemSenses *24*, 637-645.

Bragin, A., Jando, G., Nadasdy, Z., Hetke, J., Wise, K., and Buzsaki, G. (1995). Gamma

(40-100 Hz) oscillation in the hippocampus of the behaving rat. J Neurosci *15*, 47-60.

Bressler, S.L. (1988). Changes in electrical activity of rabbit olfactory bulb and cortex to

conditioned odor stimulation. Behav Neurosci *102*, 740-747.

Buonviso, N., Amat, C., Litaudon, P., Roux, S., Royet, J.P., Farget, V., and Sicard, G.

(2003). Rhythm sequence through the olfactory bulb layers during the time window of a

respiratory cycle. EurJNeurosci *17*, 1811-1819.

Buzsaki, G. (2002). Theta oscillations in the hippocampus. Neuron *33*, 325-340.

Buzsaki, G. (2010). Neural syntax: cell assemblies, synapsembles, and readers. Neuron

68, 362-385.

Buzsaki, G., and Draguhn, A. (2004). Neuronal oscillations in cortical networks. Science

304, 1926-1929.

Canas, F., and Jones, M. (2010). Attention and reinforcement learning: constructing

representations from indirect feedback. Paper presented at: Proceedings of the Annual

Meeting of the Cognitive Science Society.

Canolty, R.T., and Knight, R.T. (2010). The functional role of cross-frequency coupling.

Trends Cogn Sci *14*, 506-515.

Cavarretta, F., Marasco, A., Hines, M.L., Shepherd, G.M., and Migliore, M. (2016).

Glomerular and Mitral-Granule Cell Microcircuits Coordinate Temporal and Spatial

Information Processing in the Olfactory Bulb. Front Comput Neurosci *10*, 67.

Chae, H., Kepple, D.R., Bast, W.G., Murthy, V.N., Koulakov, A.A., and Albeanu, D.F.

(2019). Mosaic representations of odors in the input and output layers of the mouse

olfactory bulb. Nat Neurosci *22*, 1306-1317.

Chery, R., Gurden, H., and Martin, C. (2014). Anesthetic regimes modulate the temporal

dynamics of local field potential in the mouse olfactory bulb. J Neurophysiol *111*, 908-

917.

Chrobak, J.J., and Buzsaki, G. (1998). Gamma oscillations in the entorhinal cortex of

the freely behaving rat. J Neurosci *18*, 388-398.

Chu, M.W., Li, W.L., and Komiyama, T. (2016). Balancing the Robustness and

Efficiency of Odor Representations during Learning. Neuron *92*, 174-186.

Courtiol, E., and Wilson, D.A. (2017). The Olfactory Mosaic: Bringing an Olfactory

Network Together for Odor Perception. Perception *46*, 320-332.

Curran-Everett, D. (2000). Multiple comparisons: philosophies and illustrations.

AmJPhysiol RegulIntegrComp Physiol *279*, R1-R8.

Cury, K.M., and Uchida, N. (2010). Robust odor coding via inhalation-coupled transient

activity in the mammalian olfactory bulb. Neuron *68*, 570-585.

Daume, J., Gruber, T., Engel, A.K., and Friese, U. (2017). Phase-Amplitude Coupling

and Long-Range Phase Synchronization Reveal Frontotemporal Interactions during

Visual Working Memory. J Neurosci *37*, 313-322.

David, F., Courtiol, E., Buonviso, N., and Fourcaud-Trocme, N. (2015). Competing

Mechanisms of Gamma and Beta Oscillations in the Olfactory Bulb Based on

Multimodal Inhibition of Mitral Cells Over a Respiratory Cycle. eNeuro *2*.

Doucette, W., Gire, D.H., Whitesell, J., Carmean, V., Lucero, M.T., and Restrepo, D.

(2011). Associative cortex features in the first olfactory brain relay station. Neuron *69*,

1176-1187.

Doucette, W., and Restrepo, D. (2008). Profound context-dependent plasticity of mitral

cell responses in olfactory bulb. PLoS Biol *6*, e258.

Engel, A.K., Fries, P., Konig, P., Brecht, M., and Singer, W. (1999). Temporal binding,

binocular rivalry, and consciousness. Conscious Cogn *8*, 128-151.

Florin, E., and Baillet, S. (2015). The brain's resting-state activity is shaped by

synchronized cross-frequency coupling of neural oscillations. Neuroimage *111*, 26-35.

Frederick, D.E., Brown, A., Brim, E., Mehta, N., Vujovic, M., and Kay, L.M. (2016).

Gamma and Beta Oscillations Define a Sequence of Neurocognitive Modes Present in

Odor Processing. J Neurosci *36*, 7750-7767.

Freeman, W.J., and Schneider, W. (1982). Changes in spatial patterns of rabbit

olfactory EEG with conditioning to odors. Psychophysiology *19*, 44-56.

Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through

neuronal coherence. Trends Cogn Sci *9*, 474-480.

Gao, P., Trautmann, E., Byron, M.Y., Santhanam, G., Ryu, S., Shenoy, K., and Ganguli,

S. (2017). A theory of multineuronal dimensionality, dynamics and measurement.

BioRxiv, 214262.

Gire, D.H., Kapoor, V., Seminara, A., Arrighi-Allisan, A., and Murthy, V.N. (2013a).

Olfactory search behavior in mice: A novel task design that moves beyond a single sniff.

Society for Neuroscience 2013 Online *Program No. 452.28*.

Gire, D.H., Restrepo, D., Sejnowski, T.J., Greer, C., De Carlos, J.A., and Lopez-

Mascaraque, L. (2013b). Temporal processing in the olfactory system: can we see a

smell? Neuron *78*, 416-432.

Gold, C., Henze, D.A., Koch, C., and Buzsaki, G. (2006). On the origin of the

extracellular action potential waveform: A modeling study. JNeurophysiol *95*, 3113-

3128.

Gourevitch, B., Kay, L.M., and Martin, C. (2010). Directional coupling from the olfactory

bulb to the hippocampus during a go/no-go odor discrimination task. J Neurophysiol

103, 2633-2641.

Green, D.M., and Swets, J.A. (1988). Signal detection theory and psychophysics, Vol

Reprint ed (Los Altos, Calif., USA: Peninsula Pub).

Gschwend, O., Beroud, J., and Carleton, A. (2012). Encoding odorant identity by spiking

packets of rate-invariant neurons in awake mice. PloS one *7*, e30155.

Halsey, L.G., Curran-Everett, D., Vowler, S.L., and Drummond, G.B. (2015). The fickle

P value generates irreproducible results. Nature methods *12*, 179-185.

Haney, S., Saha, D., Raman, B., and Bazhenov, M. (2018). Differential effects of

adaptation on odor discrimination. J Neurophysiol *120*, 171-185.

Harno, E., Cottrell, E.C., and White, A. (2013). Metabolic pitfalls of CNS Cre-based

technology. Cell Metab *18*, 21-28.

Hasselmo, M.E., Bodelon, C., and Wyble, B.P. (2002). A proposed function for

hippocampal theta rhythm: separate phases of encoding and retrieval enhance reversal

of prior learning. Neural Comput *14*, 793-817.

Heck, D.H., Kozma, R., and Kay, L.M. (2019). The rhythm of memory: how breathing

shapes memory function. J Neurophysiol *122*, 563-571.

Hiratani, N., and Latham, P.E. (2019). Rapid Bayesian learning in the mammalian

olfactory system. bioRxiv, 706200.

Hopfield, J.J. (1995). Pattern recognition computation using action potential timing for

stimulus representation. Nature *376*, 33-36.

Jensen, O. (2001). Information transfer between rhythmically coupled networks: reading

the hippocampal phase code. Neural Comput *13*, 2743-2761.

Jordan, R., Fukunaga, I., Kollo, M., and Schaefer, A.T. (2018). Active Sampling State

Dynamically Enhances Olfactory Bulb Odor Representation. Neuron *98*, 1214-1228

e1215.

Kay, L.M. (2005). Theta oscillations and sensorimotor performance. Proc Natl Acad Sci

U S A *102*, 3863-3868.

Kay, L.M. (2014). Circuit oscillations in odor perception and memory. Prog Brain Res

208, 223-251.

Kay, L.M. (2015). Olfactory system oscillations across phyla. Curr Opin Neurobiol *31*,

141-147.

Kepecs, A., Uchida, N., and Mainen, Z.F. (2006). The sniff as a unit of olfactory

processing. ChemSenses *31*, 167-179.

Koldaeva, A., Schaefer, A.T., and Fukunaga, I. (2019). Rapid task-dependent tuning of

the mouse olfactory bulb. Elife *8*.

Lega, B., Burke, J., Jacobs, J., and Kahana, M.J. (2016). Slow-Theta-to-Gamma Phase-

Amplitude Coupling in Human Hippocampus Supports the Formation of New Episodic

Memories. Cereb Cortex *26*, 268-278.

Leong, Y.C., Radulescu, A., Daniel, R., DeWoskin, V., and Niv, Y. (2017). Dynamic

Interaction between Reinforcement Learning and Attention in Multidimensional

Environments. Neuron *93*, 451-463.

Li, A., Gire, D.H., and Restrepo, D. (2015). Y spike-field coherence in a population of

olfactory bulb neurons differentiates between odors irrespective of associated outcome.

J Neurosci *35*, 5808-5822.

Li, A., Zhang, L., Liu, M., Gong, L., Liu, Q., and Xu, F. (2012). Effects of different

anesthetics on oscillations in the rat olfactory bulb. J Am Assoc Lab Anim Sci *51*, 458-

463.

Li, G., and Cleland, T.A. (2017). A coupled-oscillator model of olfactory bulb gamma

oscillations. PLoS Comput Biol *13*, e1005760.

Linster, C., and Cleland, T.A. (2016). Neuromodulation of olfactory transformations.

Curr Opin Neurobiol *40*, 170-177.

Lisman, J. (2005). The theta/gamma discrete phase code occuring during the

hippocampal phase precession may be a more general brain coding scheme.

Hippocampus *15*, 913-922.

Lisman, J., and Buzsaki, G. (2008). A neural coding scheme formed by the combined

function of gamma and theta oscillations. Schizophr Bull *34*, 974-980.

Lisman, J.E., and Idiart, M.A. (1995). Storage of 7 +/- 2 short-term memories in

oscillatory subcycles. Science *267*, 1512-1515.

Lisman, J.E., and Jensen, O. (2013). The theta-gamma neural code. Neuron *77*, 1002-

1016.

Litwin-Kumar, A., Harris, K.D., Axel, R., Sompolinsky, H., and Abbott, L.F. (2017).

Optimal Degrees of Synaptic Connectivity. Neuron *93*, 1153-1164 e1157.

Martin, C., Gervais, R., Chabaud, P., Messaoudi, B., and Ravel, N. (2004). Learning-

induced modulation of oscillatory activities in the mammalian olfactory system: the role

of the centrifugal fibres. J Physiol Paris *98*, 467-478.

Martin, C., Gervais, R., Messaoudi, B., and Ravel, N. (2006). Learning-induced

oscillatory activities correlated to odour recognition: a network activity. Eur J Neurosci

23, 1801-1810.

Martin, C., and Ravel, N. (2014). Beta and gamma oscillatory activities associated with

olfactory memory tasks: different rhythms for different functional networks? Front Behav

Neurosci *8*, 218.

Marton, C.D., Fukushima, M., Camalier, C.R., Schultz, S.R., and Averbeck, B.B. (2019).

Signature Patterns for Top-Down and Bottom-Up Information Processing via Cross-

Frequency Coupling in Macaque Auditory Cortex. eNeuro *6*.

Mizuseki, K., Sirota, A., Pastalkova, E., and Buzsaki, G. (2009). Theta oscillations

provide temporal windows for local circuit computation in the entorhinal-hippocampal

loop. Neuron *64*, 267-280.

Nguyen Chi, V., Muller, C., Wolfenstetter, T., Yanovsky, Y., Draguhn, A., Tort, A.B., and

Brankack, J. (2016). Hippocampal Respiration-Driven Rhythm Distinct from Theta

Oscillations in Awake Mice. J Neurosci *36*, 162-177.

Niv, Y., Daniel, R., Geana, A., Gershman, S.J., Leong, Y.C., Radulescu, A., and Wilson,

R.C. (2015). Reinforcement learning in multidimensional environments relies on

attention mechanisms. J Neurosci *35*, 8145-8157.

O'Keefe, J., and Recce, M.L. (1993). Phase relationship between hippocampal place

units and the EEG theta rhythm. Hippocampus *3*, 317-330.

Osinski, B.L., and Kay, L.M. (2016). Granule cell excitability regulates gamma and beta

oscillations in a model of the olfactory bulb dendrodendritic microcircuit. J Neurophysiol,

jn 00988 02015.

Pakan, J.M., Francioni, V., and Rochefort, N.L. (2018). Action and learning shape the

activity of neuronal circuits in the visual cortex. Curr Opin Neurobiol *52*, 88-97.

Pouille, F., and Schoppa, N.E. (2018). Cannabinoid Receptors Modulate Excitation of

an Olfactory Bulb Local Circuit by Cortical Feedback. Front Cell Neurosci *12*, 47.

Ramirez-Gordillo, D., Ma, M., and Restrepo, D. (2018). Precision of Classification of

Odorant Value by the Power of Olfactory Bulb Oscillations Is Altered by Optogenetic

Silencing of Local Adrenergic Innervation. Front Cell Neurosci *12*, 48.

Rebello, M.R., McTavish, T.S., Willhite, D.C., Short, S.M., Shepherd, G.M., and

Verhagen, J.V. (2014). Perception of odors linked to precise timing in the olfactory

system. PLoS Biol *12*, e1002021.

Reinhart, R.M.G., and Nguyen, J.A. (2019). Working memory revived in older adults by

synchronizing rhythmic brain circuits. Nat Neurosci *22*, 820-827.

Rinberg, D., Koulakov, A., and Gelperin, A. (2006). Sparse odor coding in awake

behaving mice. JNeurosci *26*, 8857-8865.

Rojas-Libano, D., Frederick, D.E., Egana, J.I., and Kay, L.M. (2014). The olfactory bulb

theta rhythm follows all frequencies of diaphragmatic respiration in the freely behaving

rat. Front Behav Neurosci *8*, 214.

Rojas-Libano, D., and Kay, L.M. (2008). Olfactory system gamma oscillations: the

physiological dissection of a cognitive neural system. Cogn Neurodyn *2*, 179-194.

Rokni, D., Hemmelder, V., Kapoor, V., and Murthy, V.N. (2014). An olfactory cocktail

party: figure-ground segregation of odorants in rodents. Nat Neurosci *17*, 1225-1232.

Rosero, M.A., and Aylwin, M.L. (2011). Sniffing shapes the dynamics of olfactory bulb

gamma oscillations in awake behaving rats. Eur J Neurosci *34*, 787-799.

Schoppa, N.E. (2006). Synchronization of olfactory bulb mitral cells by precisely timed

inhibitory inputs. Neuron *49*, 271-283.

Seymour, R.A., Rippon, G., and Kessler, K. (2017). The Detection of Phase Amplitude

Coupling during Sensory Processing. Front Neurosci *11*, 487.

Short, S.M., and Wachowiak, M. (2019). Temporal Dynamics of Inhalation-Linked

Activity across Defined Subpopulations of Mouse Olfactory Bulb Neurons Imaged In

Vivo. eNeuro *6*.

Shusterman, R., Smear, M.C., Koulakov, A.A., and Rinberg, D. (2011). Precise olfactory

responses tile the sniff cycle. Nat Neurosci *14*, 1039-1044.

Siegle, J.H., and Wilson, M.A. (2014). Enhancement of encoding and retrieval functions

through theta phase-specific manipulation of hippocampus. Elife *3*, e03061.

Skaggs, W.E., McNaughton, B.L., Wilson, M.A., and Barnes, C.A. (1996). Theta phase

precession in hippocampal neuronal populations and the compression of temporal

sequences. Hippocampus *6*, 149-172.

Slotnick, B.M., and Restrepo, D. (2005). Olfactometry with mice. In Current Protocols in

Neuroscience, J.N. Crawley, C.R. Gerefen, M.A. Rogawski, D.R. Sibley, P. Skolnick,

and S. Wray, eds. (New York: John Wiley and Sons, Inc), pp. 1-24

1181

Smear, M., Shusterman, R., O'Connor, R., Bozza, T., and Rinberg, D. (2011).

Perception of sniff phase in mouse olfaction. Nature *479*, 397-400.

Soltesz, I., Bourassa, J., and Deschenes, M. (1993). The behavior of mossy cells of the

rat dentate gyrus during theta oscillations in vivo. Neuroscience *57*, 555-564.

Stopfer, M., Bhagavan, S., Smith, B.H., and Laurent, G. (1997). Impaired odour

discrimination on desynchronization of odour-encoding neural assemblies. Nature *390*,

70-74.

Stopfer, M., and Laurent, G. (1999). Short-term memory in olfactory network dynamics.

Nature *402*, 664-668.

Tort, A.B., Komorowski, R., Eichenbaum, H., and Kopell, N. (2010). Measuring phase-

amplitude coupling between neuronal oscillations of different frequencies. J

Neurophysiol *104*, 1195-1210.

Tort, A.B.L., Ponsel, S., Jessberger, J., Yanovsky, Y., Brankack, J., and Draguhn, A.

(2018). Parallel detection of theta and respiration-coupled oscillations throughout the

mouse brain. Sci Rep *8*, 6432.

van Wingerden, M., van der Meij, R., Kalenscher, T., Maris, E., and Pennartz, C.M.

(2014). Phase-amplitude coupling in rat orbitofrontal cortex discriminates between

correct and incorrect decisions during associative learning. J Neurosci *34*, 493-505.

Vizcay, M.A., Duarte-Mermoud, M.A., and Aylwin Mde, L. (2015). Odorant recognition

using biological responses recorded in olfactory bulb of rats. Comput Biol Med *56*, 192-

199.

Zelano, C., Jiang, H., Zhou, G., Arora, N., Schuele, S., Rosenow, J., and Gottfried, J.A.

(2016). Nasal Respiration Entrains Human Limbic Oscillations and Modulates Cognitive

Function. J Neurosci *36*, 12448-12467.

Zhuang, L., Zhang, B., Qin, Z., and Wang, P. (2019). Nasal Respiration is Necessary for

the Generation of gamma Oscillation in the Olfactory Bulb. Neuroscience *398*, 218-230.

1208

1209

1210

1211

1212
1213

1214

1215

1216

1217

1218

1219

Figure 1. Behavioral task and PAC analysis.

(A) Behavioral apparatus. Mouse self-initiates trials by breaking a photodiode beam.

Odorants and water are delivered according to the timeline in B and the decision tree in

C.

(B) Timeline for a single trial. When the animal enters the port the air flow is diverted by

turning the final valve output to the exhaust and the odorant valve is turned on. At this

time the odor builds up to steady-state concentration flowing away from the animal for

1-1.5 seconds. At time 0 the final valve air flow is turned towards the odor port for 2.5 s

resulting in odor onset 100 ms after the valve is actuated. In order to trigger water

reward for the rewarded odorant the mouse must lick at least once during each 0.5 s

block for four blocks in the response area. If the stimulus is rewarded and the animal

licked appropriately, a water reward is delivered.

(C) Decision tree for each stimulus. Green check mark means the correct decision was

made. Red "X" mark indicates an incorrect decision. Water reward is represented by the

water droplet symbol in the case of a hit.

(D) PAC data analysis of the LFP. For each electrode, raw signal collected at 20 kHz (i)

is bandpass filtered to 1-100 Hz for broadband LFP (ii) or filtered into different frequency

bands as needed (e.g. theta 6-14 Hz in iii or high gamma 65-95 Hz in v, blue line).

Hilbert transform is used to calculate the theta phase (iv) and the amplitude envelope of

higher oscillations such as high gamma (65-95 Hz) (red line in v). Theta phase and the

envelope of the amplitude of high gamma are then used to calculate the probability for

high gamma amplitude at specific phases of theta (PAC, shown in E).

(E) Probability for high gamma amplitude at specific phases of theta. In this example,

PAC is strongest at ~100° and 270°. The coupling strength is quantified by the

modulation index, MI. Peak phase (the phase of theta with the highest amplitude in high

gamma) is indicated with a red line and trough phase is indicated with a blue line.

1246

1247

1248

1249

Figure 2. The peak angle variance of PAC changes as the animal learns to

discriminate odorants.

(A-F) Example of theta/high gamma phase amplitude coupling for a proficient animal

(IAAPexp1 odorant pair).

(A) Pseudocolor plots showing the phase amplitude relationship for S+ (left) and S-

(right) for an example go no-go session. PAC between theta (6-14 Hz) phase and high

gamma (65-95 Hz) amplitude was more variable for S- compared to S+.

(B) Behavioral performance for this session. Over 100 trials, this trained animal

discriminated between isoamyl acetate and acetophenone with 95-100% accuracy.

(C) Theta phase distribution for high gamma amplitude (mean $\pm$ CI) for S+ (red) and S-

(blue).

(D) Mean theta LFP (mean $\pm$ CI) for one cycle for S+ (red) and S- (blue).

(E) MI per trial for S+ (red) and S- (blue) as a function of trial number. The difference in

MI between S+ and S- is statistically significant (ranksum p value < 0.001, n=50 S-

trials, 49 S+ trials).

(F) Polar histograms for high gamma peak theta angles for S+ (red) and S- (blue).

(G) Cumulative histograms showing the differences in mean MI (i,iii) and peak angle

variance (ii,iv) for theta/beta PAC (i,ii) and theta/high gamma PAC (iii,iv) for the APEB

odorant in Exp1 for the S+ and S- odorants for naïve and proficient trial ranges. Peak

angle variance for S+ displays a substantial decrease with learning, while mean MI

shows small changes. Changes with learning and between events for mean MI and

peak angle variance are statistically significant (GLM p value < 0.001, 896 d.f. for

theta/beta and theta/high gamma, 16 electrodes per mouse, 14 mice). Blue and red

arrows in Giii are MIs derived from the single-session example in C.

(H) Bar graph displaying mean MI (i,iii) and peak angle variance (ii,iv) for naïve and

proficient trial ranges and S+ and S- events for mean values calculated per experiment

for theta/beta (i,ii) and theta/high gamma PAC (iii,iv). GLM yielded significant p values

for peak angle variance for theta/high gamma PAC for S+ vs. S- (< 0.01, 16 d.f.) and

for the interaction between S+ vs. S- and proficiency (p<0.05, 16 d.f.). GLM yielded

significant p values for peak angle variance for theta/beta PAC for S+ vs. S- and

experiment and for the interaction between S+ vs. S- and experiment (< 0.01, 16 d.f.).

For mean MI theta/high gamma PAC GLM yields significant difference for experiments

(p<0.01, 16 d.f.). Finally, there are no significant effects for mean MI for theta/beta PAC

(p>0.05, 16 d.f.) and no post hoc tests yield significant differences (p>pFDR).

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

Figure 3. PRP increases for S+ and decreases for S- over learning.

(A) Example wavelet broadband LFP spectrogram for one S+ trial. Right: Pseudocolor

scale for LFP power (in dB). Black bar represents the odor stimulation epoch (2.5 s). i:

Full trial length demonstrating pre-odor baseline LFP, odor driven LFP, and reward

epoch. ii: One-second epoch during odor stimulation.

(B) i: Theta waveform (6-14 Hz) during odor stimulation. ii: Theta phase, extracted with

Hilbert transform.

(C) i: Single trial wavelet power referenced to either peak (108°, red) or trough (0°, blue)

of the high gamma amplitude distribution along the theta phase. ii: PRP (± CI) averaged

across the entire session for S+ and S-. Peak PRP for S+ displays the largest odor-

induced increase in power that is sustained after the end of odor stimulation.

(D) The thin lines are cumulative histograms for PRP for the LFP recorded in each

electrode averaged for theta/high gamma for all proficient or naïve trials for mice

performing to differentiate odorants in the IAAPexp2 odorant pair. i and ii are peak PRP

and iii and iv are trough PRP. i and iii are S+; ii and iv are S-. As the animals learn the

task, peak PRP (i and ii) increases for S+ and decreases for S-, whereas trough PRP (iii

and iv) is relatively stable. The differences were statistically significant between peak

and trough, S+ and S- and naïve and proficient (GLM p values < 0.001, 1016 d.f.).

Circles are per mouse averages for PRP.

(E-F) Odor-induced changes in high gamma PRP averages for all odorant pairs

evaluated per electrode during the odor delivery epoch (0.5 – 2.5 s). Peak PRP (E)

increases for S+ and decreases for S- when the mice become proficient. Trough PRP

changes (F) are markedly weaker, yet they follow a similar trend (GLM p value < 0.001,

6386 d.f. for S+ vs. S-, peak vs. trough, naïve vs. proficient). In addition, the learning-

induced changes in PRP are larger in Exp2 (GLM p value < 0.001 for experiment, 6386

d.f.).

1320

1321

Figure 4. Peak theta-phase referenced power switches when odorant valence is reversed.

(A) Behavioral performance in three go-no go sessions for a mouse learning to

differentiate the IAAPexp1 odorant pair. In the first session the rewarded odorant (S+)

was isoamyl acetate (IA) and the unrewarded odorant (S-) was acetophenone (AP). In

the other two sessions the valence of the odorant was reversed.

(B-C) Theta/beta (B) and theta/high gamma (C) peak PRP, calculated for trails when the

mouse is proficient, switches when odorant valence is reversed. The data are shown for

three odor pairs: EAPAexp1 (11 mice), EAPAexp2 (9 mice), IAAPexp2 (3 mice). GLM

analysis yields significant changes for odorant, reversal and the interaction of reversal

1333   and odorant ($p<0.001$, 1051 d.f.). Asterisks denote post hoc significant differences

1334   ($p<p\text{FDR}=0.04$).

Figure 5. Linear discriminant analysis classifies stimuli using peak PRP.

(A-B) PCA (in panel A) and linear discriminant analysis (LDA; in panel B) of averaged

PRP data for one example odorant pair, APEBexp1.

(A) PCA of PRP data from naïve (i,iii) to proficient (ii,iv)—i,ii: theta/beta, iii,iv: theta/high

gamma. Both PRPs for S+ and S- are discriminable with learning.

(B) Decoding performance for the LDA of PRP data over learning (i,iii: naïve, ii,iv:

proficient). i,ii: theta/beta, iii,iv: theta/high gamma. In the proficient state, LDA can

decode stimulus identity using both PRPs (theta/beta PRP GLM naïve-proficient p value

< 0.001, 104 d.f.; theta/high gamma PRP GLM naïve-proficient p value < 0.001, 104

d.f.). Peak and trough carry similar information for theta/beta; both peak- and trough-

referenced beta power can be used to decode stimulus identity with learning. In

theta/high gamma, peak PRP is significantly better at decoding the stimulus identity

than trough PRP (GLM p value < 0.001, 104 d.f.), which is still better than shuffled data.

(C) Area under the curve (AUC) for peak versus trough PRP over task learning (i,iii:

naïve, ii,iv: proficient) for all odorant pairs and experiments. i,ii: Theta/beta PRP. iii,iv:

For theta/high gamma PRP GLM statistics show that peak AUC is significantly higher

than trough (p value < 0.001 for 6 odorant pairs, 184 d.f.) and that there are significant

changes over learning (p value < 0.001 for 6 odorant pairs, 184 d.f.). These data

suggest that peak PRP decodes stimulus identity better than trough PRP.

(D) Average peak AUC ($\pm$CI) for all odorant pairs for theta/beta and theta/high gamma

PRP for proficient mice for the two experimental settings. The learning-induced changes

in PRP are larger in Exp2 (GLM p value < 0.05 for experiment, 102 d.f., *post hoc

p<pFDR=0.03).

(E) Decoding performance for the LDA of PRP data for proficient mice calculated for the

different behavioral outcomes of the go-no go task (Hit: red, Miss: cyan, CR: blue, FA:

magenta, shuffled: black). i: Theta/beta PRP, ii: Theta/high gamma PRP. A GLM

analysis indicates that there are significant differences between all behavioral outcomes

except for Hit vs. CR for theta/high gamma (p<0.001 for all outcome pairs, except for Hit

vs CR for theta/beta PRP that has p<0.01 and Hit vs. CR for theta/high gamma that has

p>0.05, 2110 d.f.). There is no significant difference between experiments (GLM

p>0.05, 2110 d.f.).

The lines in A,B and E are bounded by the bootstrapped 95% CI.

Figure 6. Decision-making time does not differ between licks and peak PRP LDA

decoding of odorant identity

(A) Lick time course for 25 S+ trials in one session of EAPAexp2 for one proficient

mouse. Horizontal black bar indicates odor delivery epoch for all time course subpanels.

(B) Lick time course for 25 S- trials in the same session as in A. Mouse refrains from

licking in the presence of the unrewarded odorant.

(C) i: p value time courses for difference between S+ and S- for licks and peak PRP

LDA predictions. ii and iii: decoding performance for PRP LDA. Decision making times

for both licks and PRP LDA were calculated using data pooled for all mice performing

odorant discrimination with the EAPAexp2 odorant pair. i: p value for the ranksum test

compares the time-course for decision-making using licks for S+ vs. S- (green) or LDA

prediction of odorant identity with peak (red) or trough (blue) PRP. This example

illustrates that that peak PRP LDA prediction and lick divergence perform similarly,

while LDA prediction with trough PRP is significantly slower in stimulus decoding.

Horizontal red line indicates p = 0.05. ii and iii: LDA decoding performance appears to

be faster and more accurate using peak PRP (ii) vs. trough PRP (iii).

(D) Decision times for licks and for pooled animal peak theta/beta and theta/high

gamma PRP LDA in naïve (i) and proficient (ii) mice. Decision times were significantly

reduced with task learning (GLM p value < 0.001, 44 d.f.). Decision times for lick vs.

PRP LDA and theta/beta vs. theta/high gamma did not differ (GLM p values > 0.05, 44

d.f.).

1390

1391

Figure 7. Dimensionality changes with learning and during the time course of the trial.

(A) Scatter plot showing the first two principal components for twenty trials of a PCA for peak theta/high gamma PRP recorded in mice learning to differentiate the odorants in the EAPAexp1 odorant pair. The input to the PCA was the peak PRP recorded in 16 electrodes pooled for all mice. Principal components are intermingled between S+ (red)

and S- (blue) trials in the naïve state (left) in pre-odor and odor-delivery epochs, yet

they separate during odor delivery upon task acquisition (right). i. Naïve animals. ii.

Proficient animals.

(B) Within-trial peak (red) and trough (blue) theta/high gamma PRP dimensionality time

course for data pooled across mice is shown for Exp1 (i) and Exp2 (ii), normalized to

pre-trial dimensionality. A GLM analysis indicates that there are significant differences

for time course, proficient vs. naïve, experiments and peak vs. trough, and for pair

interactions of these factors (GLM p values < 0.001, 1688 d.f.).

(C) Within-trial peak (red) and trough (blue) theta/beta PRP dimensionality time course

for data pooled across mice is shown for Exp1 (i) and Exp2 (ii), normalized to pre-trial

dimensionality. GLM analysis found statistically significant differences for time course,

and proficient vs. naïve, and for interaction of these factors (GLM p values < 0.001,

1688 d.f.), and there was a statistically significant difference between experiments (GLM

p value < 0.01, 1688 d.f.). However, there was no difference between peak and trough

(GLM p value >0.05, 1688 d.f).

1414

1415

SUPPLEMENTAL INFORMATION

Detailed evaluation of phase amplitude coupling on a per odorant pair basis.

Figure 2-figure supplement 2 A and B show the summary of changes with learning for S+ and S- odorants for MI (Ai, Bi) and the peak angle variance (Aii,Bii) for theta/beta (A) and theta/high gamma (B) for per electrode PAC for all odorant pairs. When evaluated on a per odorant pair basis theta/beta and theta/high gamma PAC strength, quantified as the MI calculated for LFPs recorded in each electrode, was stronger for recordings in Exp2 compared to Exp1, and there was a statistically significant increase in the MI for S+ with learning (naïve vs. proficient) (GLM analysis p value < 0.001, for event, proficiency and experiment, with the exception that p < 0.01 for proficiency for theta/high gamma, 3255 d.f., number of mice is in Table S1). Interestingly, there were decreases in peak angle variance for S+, and/or increases in peak angle variance for S- for theta/high gamma PAC when the animals became proficient (GLM analysis p values < 0.001 for event type, proficiency and experiment for both theta/beta and theta/high gamma, 3255 d.f., number of mice is in Table S1).

Comparison of decision-making times calculated using pooled animal PRP LDA data vs. per animal data.

Figure 6-figure supplement 1 B and C show that for both lick (B) and PRP peak LDA (C) the decision-making time is faster when data are pooled for all mice. Lick decision-making times were significantly faster with pooled data (GLM p value < 0.01, 21 d.f.) as were decision making times for peak PRP LDA (GLM p value < 0.01, 21 d.f.).

1438   Because pooled animal data yield faster decision times we did the comparison of

1439   decision times in Figure 6 using pooled animal data.

1440

1441

Table S1. Odorant pairs, implant device, mouse genotype and electrode locations

for each experiment

Experiment	Implant	Coordinates	Genotype	Odorant pair	Odorant name	Odorant dilution (v/v)	Number of mice
Exp1	Optetrodes	+4.25 mm AP, +0.4 mm ML, -0.53 mm DV	DBH-Cre eNpHR3.0 (see Methods)	APEBexp1	Acetophenone	1%	14
					Ethyl benzoate	1%	
				EAPAexp1	Ethyl acetate (EA)	0.1%	11
					EA/Propyl acetate (PA)	0.05% EA + 0.05% PA	
				IAMOexp1	Isoamyl acetate	1%	9
					Mineral oil	100%	
Exp2	Tetrodes	+4.28 mm AP, +0.05 mm ML, -1.0 mm DV	C57BL/6	EAPAexp2	Ethyl acetate	1%	9
					Propyl acetate	1%	
				IAAPexp2	Isoamyl acetate	1%	6
					Acetophenone	1%	
				IAMOexp2	Isoamyl acetate	1%	4
					Mineral oil	100%	

Odorant identities for each pair are listed along with concentrations represented as

volume/volume dilutions in mineral oil (MO). In the case of EAPAexp1, EA is 0.1% ethyl

acetate and PA is a mixture of 0.05% ethyl acetate + 0.05% propyl acetate. The two

recording locations (Exp1 and Exp2) are listed in millimeters with respect to bregma.

Note: Cre expression can affect physiological parameters and therefore, the differences

found between experiments in this publication can be due to Cre expression (Harno et

al., 2013). However, it is not evident to us which of the methodological differences

between experiments contributes to statistical differences in measured parameters.

Experiment	Mouse ID	Odorant pair				
		APEB	EAPA	IAAP	IAMO	
1	261365	2	2	N/A	2	
1	261366	2	2	N/A	2	
1	261367	2	2	N/A	2	
1	261368	2	2	N/A	2	
1	262641	2	2	N/A	2	
1	293851	0	2	N/A	0	
1	294213	0	2	N/A	0	
1	254071	2	2	N/A	2	
1	261032	2	2	N/A	2	
1	294211	0	2	N/A	0	
1	294212	0	2	N/A	0	
1	159868	2	0	N/A	0	
1	159863	2	0	N/A	0	
1	159865	2	0	N/A	0	
1	159866	2	0	N/A	0	
1	159867	2	0	N/A	0	
1	159869	2	0	N/A	0	
1	254072	2	0	N/A	2	
1	261036	0	0	N/A	2	Experiment 1 total
	Total per odorant pair	28	22	N/A	18	68
2	M4	N/A	0	5	0	
2	M5	N/A	0	2	0	
2	T0	N/A	1	3	0	
2	T2	N/A	1	4	0	
2	R2	N/A	2	0	2	
2	R3	N/A	3	0	0	
2	R4	N/A	3	0	3	
2	R5	N/A	4	0	4	
2	R6	N/A	2	0	2	
2	R8	N/A	1	3	0	
2	R9	N/A	2	4	0	Experiment 2 total
	Total per odorant pair	N/A	19	21	11	51
						Grand total
						119

Table S2. Total sessions per mouse, experiment, and odorant pair.
Odorant pair abbreviations derived from Table S1. The number of sessions are in the "odorant pair" columns. Each row represents one animal.

1455

Figure 1-figure supplement 1. Time course for odorant concentration measured at the odor spout in the olfactometer. A miniPID (Aurora Scientific) photoionization detector was used to measure the concentration of odorant at the odor spout of the olfactometer. The odorant was air equilibrated with Sharpie black ink. The red line is the concentration of the odorant in arbitrary units bounded by the bootstrapped confidence interval (n=10 traces). Time was aligned to the onset of the increase in odorant concentration that took place 104.5±7 msec (mean±SD, n=10) after the final valve was diverted to deliver the odorant to the odorant spout. The black lines are regression fits of exponential increase and decrease lines with time constants of 0.12 and 2.16 sec

1465 (increase, two exponential terms) and 0.19 and 1.28 sec (decrease, two exponential

1466 terms).

1467

Figure 1-figure supplement 2. Simulation: PRP wavelet analysis for theta-gamma waves with/without PAC.

LFP data were generated by sinusoids supplemented with gaussian random noise. The data was analyzed for PRP as described in the text.

(A) Example with no theta/gamma PAC generated by input of two continuous cosine waves of 8 Hz and 40 Hz. Top: Morlet wavelet spectrogram shows a continuous increase in power at the two frequencies. Bottom: peak (red) and trough (blue) PRP do not differ. MI = 0.0004.

(B) Example of theta/gamma PAC generated with input of 40 Hz gamma bursts placed at the 180° phase of an 8 Hz theta sine wave (the bursts take place at the peak of PAC). Top: Morlet wavelet spectrogram shows discrete increases of gamma power and a continuous increase in power at 8 Hz. Bottom: peak (red) PRP shows a large increase in power while trough PRP (blue) does not increase. MI = 0.01.

1487

Figure 2-figure supplement 1. Example of PAC changes over learning in one

session.

(A) PAC for all S+ trials (left) and S- trials (middle), with high γ amplitude color bar

quantified at right.

(B) Behavioral performance. Mouse starts the session performing at chance, then

learns the task, acquiring proficiency by the end. Blue: naïve, performance ≤ 65%. Red:

proficient, performance ≥ 80%.

 (C) PAC quantified as mean MI for S+ (red) and S- (blue). With learning, PAC

increases for S+ and decreases for S-.

1497

Figure 2-figure supplement 2. Modulation index and peak angle variance shown

for each odorant pair.

Bar graphs displaying for theta/beta (A) and theta/high gamma (B) PAC differences in

mean MI (Ai and Bi) and peak angle variance (Aii and Bii) between naïve and proficient

trial ranges and S+ and S- events for all sessions for all the odorant pairs. GLM yielded

significant p values (< 0.001) for both mean MI and peak angle variance for event,

proficiency and experiment, with the exception that $p < 0.01$ for proficiency for theta/high

gamma (3255 d.f., number of mice is in Table S1). Asterisks denote a significant post-

hoc difference evaluated with either a t-test or a ranksum test ($p < pFDR$, for the

number of mice per odorant, see Table S1). Changes in mean MI were variable for

different odorants. MI displays differences between events and learning states that differ

in magnitude between odorant pairs. Peak angle variance (at right) increases with task

proficiency for S- and decreases for S+ for all odorants.

1512

1513

1514

Figure 5-figure supplement 1. Simulated LDA performance data illustrating AUC

zero and one.

(A) Simulated data illustrating the time course for decoding performance for LDA for

discrimination of odor identity with: i, The area under the curve (AUC) = 0, LDA was not

able to identify the stimuli, and ii, AUC = 1, perfect identification of the stimulus after

odor application. AUC was calculated as the area under the curve during odorant

stimulation normalized to 1.

1522

1523

Figure 6-figure supplement 1. Decision times are faster for peak PRP LFP LDA

computed when data are pooled across mice.

(A) Comparison of lick decision times between per mouse (light green) vs. pooled

mouse (dark green) p value estimates. GLM p value for the difference in lick decision

times between per-mouse vs. pooled GLM < 0.01, 20 d.f.

(B) Decision times for peak PRP decoding of stimulus identity for the per mouse (light

red) vs. pooled mouse (red) LDA for naïve (i) vs. proficient (ii) mice. For both theta/beta

and theta/high gamma decoding of stimulus identity via peak PRP is faster using data

pooled across mice than the per mouse estimate (GLM p values < 0.001, 20 d.f.). There

is no difference in decision-making times between theta/beta and theta/high gamma

(GLM p values > 0.05, 20 d.f.).

Vertical black bars represent the 95% bootstrap confidence interval of the data in each

condition; they are shifted to the left to better display the mean and individual odorant

pair values.

Figure 7-figure supplement 1. PRP dimensionality estimated per mouse.

(A) Per mouse theta/high gamma (i) and theta/beta (ii) PRP dimensionality summary

for all six odorant pairs calculated before odorant delivery. GLM p value < 0.001 for

Exp1 vs. Exp2 and > 0.05 for peak vs. trough, 379 d.f.

(B) Per mouse examples of the time course of theta/high gamma PRP dimensionality

for APEBexp1 (i) and IAMOexp2 (ii). A GLM analysis of the dimensionality for peak

PRP LFP calculated on a per mouse basis for all odorant pairs indicates that there are

significant differences in theta/high gamma peak PRP LFP dimensionality for proficient

vs. naïve, time, Exp1 vs. Exp2 and for interactions of these factors (except for the

interaction of Exp1 vs. Exp2 x proficient vs. naïve) (GLM p values < 0.001, 6098 d.f.).

(C) Per mouse examples of the time course of theta/beta PRP dimensionality for

APEBexp1 (i) and IAMOexp2 (ii). A GLM analysis of the dimensionality for theta/beta

peak PRP LFP calculated on a per mouse basis for all odorant pairs indicates that there

are significant differences for proficient vs. naïve, time, Exp1 vs. Exp2 and for

interactions of these factors (GLM p values < 0.01, 6098 d.f.).

For theta/high gamma peak PRP LFP dimensionality there was a statistically significant

difference between peak and trough (GLM p value < 0.0001, 12196 d.f.). and for

theta/beta peak PRP LFP dimensionality there was no difference between peak and

trough (GLM p value > 0.05, 12196 d.f.).