AR phosphorylation and CHK2 kinase activity regulates IR stabilized AR – CHK2

interaction and prostate cancer survival

Huy Q Ta[1,#], Natalia Dworak[1,#], Melissa L Ivey[1], Devin G. Roller[1], and Daniel

Gioeli[1,2,*]

1 Department of Microbiology, Immunology, and Cancer Biology, University of Virginia,

Charlottesville, Virginia, United States of America

2 Cancer Center Member, University of Virginia, Charlottesville, Virginia, United States

of America

Running title: CHK2 and AR interact to regulate PCa

[#] These authors contributed equally to the work.

[*] To whom correspondence should be addressed. Daniel Gioeli, Department of

Microbiology, Immunology, and Cancer Biology, University of Virginia, Charlottesville,

Virginia, 22908, United States of America; Tel: (1) 434-982-4243; Fax: (1) 434-982-

0689; Email: dgioeli@virginia.edu

ABSTRACT

We have previously demonstrated that checkpoint kinase 2 (CHK2) is a critical

negative regulator of androgen receptor (AR) transcriptional activity, prostate cancer

(PCa) cell growth, and androgen sensitivity. We have now uncovered that the AR

directly interacts with CHK2 and ionizing radiation (IR) increases this interaction. This

IR-induced increase in AR–CHK2 interactions requires AR phosphorylation and CHK2

kinase activity. PCa associated CHK2 mutants with impaired kinase activity reduced IR-

induced AR–CHK2 interactions. The destabilization of AR–CHK2 interactions induced

by CHK2 variants impairs CHK2 negative regulation of cell growth. CHK2 depletion

increases transcription of DNAPK and RAD54, increases clonogenic survival, and

increases resolution of DNA double strand breaks. The data support a model where

CHK2 sequesters the AR through direct binding decreasing AR transcription and

suppressing PCa cell growth. CHK2 mutation or loss of expression thereby leads to

increased AR transcriptional activity and survival in response to DNA damage.

INTRODUCTION

Mammalian cells are continuously being bombarded by endogenous and exogenous forces that jeopardize the integrity of DNA. In response to DNA damage, a conserved network of signaling pathways known as the DNA damage response (DDR) is activated to maintain cell viability and genome stability [1]. Prostate cancer (PCa) remains one of the leading causes of death among men of all races (cdc.gov), as castration-resistant prostate cancer (CRPC) is currently incurable. Recently, the DDR has been a focus of PCa research since the androgen receptor (AR), a major driver of PCa, modulates the transcription of DDR genes and DNA repair [2]–[4]. We have previously shown that checkpoint kinase 2 (CHK2) negatively regulates androgen sensitivity and PCa cell growth [5].

CHK2 is a serine/threonine protein kinase that plays a crucial role in sensing DNA damage and initiating the DDR, which is comprised of cell cycle arrest, DNA repair, and apoptosis [6]. CHK2 consists of an amino-terminal SQ/TQ cluster domain (SCD) where threonine 68 serves as a substrate for phosphorylation by ataxia-telangectasia mutated (ATM) kinase [7]; a carboxy-terminal kinase domain (KD) and nuclear localization sequence [8]; and a central forkhead-associated domain (FHA) that provides an interface for interactions with phosphorylated proteins [9]. Currently, there are approximately 24 CHK2 substrates in human cells that have been experimentally validated, including polo-like kinase 1 (PLK1), promyelocytic leukemia protein (PML), E2F1, p53, and cell division cycle 25C (CDC25C) [10]. These studies show that one mechanism CHK2 utilizes to affect cellular function is through direct protein-protein interactions.

CHK2 association with PLK1 leads to its localization at centrosomes where it

regulates mitotic entry [11]. CHK2 autophosphorylation and activation are regulated by

the tumor suppressor PML within nuclear matrix-associated structures called PML-

nuclear bodies, which are nuclear matrix-associated structures [12]. Binding to PML

keeps CHK2 in an inactive state within these PML-nuclear bodies. In return, activated

CHK2 can phosphorylate PML on S117 and induce PML-mediated apoptosis. CHK2

can also modify the transcription of apoptotic genes through direct binding and S364

phosphorylation of the E2F1 transcription factor in response to DNA damage, thereby

stabilizing E2F1 and activating gene transcription [13]. CHK2 also regulates apoptosis

through p53 phosphorylation, and promotion of p53-mediated cell death [14]. The

interaction with the core domain of p53 induces an allosteric change in CHK2 which

permits p53 S20 phosphorylation [15]. Moreover, CHK2 modulates CDC25C localization

by associating with and phosphorylating CDC25 on S216, which creates a binding site

for 14-3-3 proteins [16]. 14-3-3 proteins in turn sequester CDC25C in the cytoplasm and

block the G2/M transition since cyclin dependent kinase 1 (CDK1) cannot be activated.

Finally, our group has shown that CHK2 co-immunoprecipitated with AR in PCa cells

and regulated growth, suggesting that AR may be a novel substrate of CHK2 [5]. Thus,

given the importance of CHK2 and AR to the DDR and prostate cancer growth, a full

understanding of the functional consequences of the CHK2–AR interaction is required,

with the hope of possible clinical applications towards CRPC.

Here, we uncovered novel molecular interactions between CHK2 and AR that

provide mechanistic insight into our observation that CHK2 negatively regulates

prostate cancer growth. We demonstrate that AR directly binds CHK2, and that this

interaction increases with ionizing radiation (IR). The IR-induced increase in AR – CHK2

interaction requires AR phosphorylation on both S81 and S308 and CHK2 kinase

activity, which is impaired in PCa associated CHK2 mutations. Furthermore, these

CHK2 mutants exhibit diminished effects on prostate cancer cell growth. CHK2

depletion increases transcription of DNAPK and RAD54, clonogenic survival, and

resolution of DNA double strand breaks (DSBs) following IR, while altering the kinetics

of IR-induced γH2AX foci.

RESULTS

AR directly binds CHK2

We previously showed that AR co-immunoprecipitated with CHK2 immune

complexes in several prostate cancer cell lines [5]. To determine whether this co-

association was through direct protein-protein interaction, we performed Far Western

blotting (Figure 1, Figure 1-figure supplement 1) [17], [18]. We also used Far Western

blotting to test the impact of AR phosphorylation on CHK2 binding to AR. We focused

on S81 and S308 phosphorylation since the phosphorylation of AR on S81 by CDK1

and CDK9 stimulates AR transcriptional activity and growth of PCa cells [19]–[21]. AR

phosphorylation on S308 by CDK1 represses AR transcriptional activity and alters AR

localization during mitosis [22], [23]. To generate purified protein for the far westerns,

293T cells were transfected with mammalian plasmids expressing FLAG-wtAR, or the

FLAG-AR mutants S81A, S81D, S308A, or S308D, FLAG-ERK2, or V5-wtCHK2 (Figure

1A). We used FLAG-ERK2 as a positive control since it has been reported that CHK2

physically associates with ERK1/2 in cancer cells [24]. FLAG-ERK and FLAG-AR

targets were immunoaffinity purified and resolved by SDS-PAGE. The target proteins

(AR wt and phospho-mutants and ERK) on the membrane were probed with purified V5-

wtCHK2 protein, crosslinked, and stained with V5 antibodies to detect bound V5-

wtCHK2. Membranes were also immunoblotted with AR and ERK1/2 antibodies to

confirm that the molecular weight of AR and ERK1/2 corresponded with the CHK2

signal, which then indicates direct protein-protein interaction. We found that V5-wtCHK2

bound to FLAG-wtAR, as well as the control FLAG-ERK2 (Figure 1A). Only AR S308A

showed a statistically significant reduction in binding to CHK2 (adjusted p=0.035),

although all AR phosphorylation mutants trended to decrease association with CHK2.

112   We observed the reciprocal binding of AR and CHK2 by Far Western.

Expression, purification, and Far Western blotting were performed as described above

using FLAG-wtCHK2 and FLAG-CHK2 mutants K373E or T387N as targets and HA-

wtAR as a probe. This also showed that AR and CHK2 directly bind (Figure 1B). In

these experiments we also determined whether the CHK2 mutants in PCa with reduced

kinase activity bind to AR. The K373E CHK2 mutation impairs CHK2 function

suppressing cell growth and promoting survival in response to IR, as a result of reduced

kinase activity due to the disruption of CHK2 autophosphorylation [25]. Less is known

about the heterozygous missense mutation T387N, but it is reported to diminish kinase

activity [26]. We observed reduced AR – CHK2 binding (adj p=0.026) with the CHK2

K373E mutant. No significant change was observed with the CHK2 T387N mutant.

These data suggest that the AR directly binds CHK2 and that AR phosphorylation and

CHK2 kinase activity may impact that binding.

Radiation increases AR – CHK2 association

126   Since IR is a standard of care for patients with localized advanced PCa, and

CHK2 is a mediator of the DDR, we wanted to assess the impact of IR on AR – CHK2

interactions. Moreover, we wanted to test the influence of AR phosphorylation and

CHK2 mutation on any effect of IR on the AR – CHK2 interaction. LNCaP cells were

transduced with lentiviral particles containing wtAR or AR mutants S81A or S308A

(Figure 2A). The association of AR was analyzed from endogenous CHK2 immune

complexes generated one hour after irradiation. There was a 4-fold increase in AR co-

immunoprecipitating with CHK2 after IR treatment of cells expressing wtAR. However,

neither S81A nor S308A increased in association with CHK2 upon IR treatment

indicating that phosphorylation of AR on S81 and S308 are required for the IR-induced

increase in the interaction between AR and CHK2. Interestingly, in contrast to the AR –

CHK2 association by Far Western, the basal level of CHK2 and AR co-

immunoprecipitation (co-IP) was unaffected by AR phosphorylation.

The requirement for AR phosphorylation for the IR-induced increase in AR –

CHK2 association led us to test if AR phosphorylation on S81 and S308 was increased

in response to IR. AR was immunoprecipitated from irradiated cells, and phospho-S81

and phospho-S308 were measured by western blotting using phospho-specific

antibodies to those sites [20], [27] (Figure 2-figure supplement 1). There were no

significant changes in S81 or S308 phosphorylation in response to IR in either cell line.

Thus, these findings indicate that while the intensity of S81 and S308 phosphorylation

does not markedly change with IR, S81 and S308 phosphorylation is necessary for AR

– CHK2 interactions under certain conditions.

CHK2 kinase activity is required for AR – CHK2 interaction

To further determine if CHK2 kinase activity was necessary for the AR – CHK2

association and what impact IR has on CHK2 mutant association with AR, we tested if

CHK2 mutants that are found in PCa and have impaired kinase activity could interact

with the AR by co-IP. We expressed FLAG-wtCHK2, FLAG-K373E, or FLAG-T387N in

combination with HA-wtAR in LNCaP cells. FLAG-CHK2 immunoprecipitations were

performed and HA-AR association was evaluated (Figure 2B). In response to IR there

was a striking increase in CHK2 – AR co-association in cells expressing FLAG-wtCHK2

and HA-wtAR. However, the amount of IR-induced increase in the AR – CHK2

association in cells expressing either K373E or T387N was dramatically reduced. No

change in basal AR – CHK2 co-IP was observed. Immunoblotting of cell lysates used in

the co-IPs indicate that the differences in co-IPed proteins was not due to protein

expression (Figure 2C). These data indicate that the kinase activity of CHK2 is required

for the optimal induction of the CHK2 and AR interaction in response to IR, and that the

CHK2 mutants associated with PCa have a diminished ability to interact with the AR in

irradiated cells.

PCa associated CHK2 mutants limit suppression of PCa growth

Since CHK2 regulates the cell cycle and PCa cell growth [5], [15], we

investigated the effect of CHK2 mutations found in PCa, which disrupt the AR – CHK2

interaction, on CHK2 regulation of PCa cell growth (Figure 3). PCa cells were

transduced with vector or shRNA specific to CHK2 to deplete endogenous CHK2, and

then CHK2 expression was rescued with wtCHK2, K373E, or T387N. Cell growth was

quantitated in the presence and absence of 0.05nM R1881 (synthetic androgen) seven

days following transduction using CyQUANT to assess cell proliferation as a function of

DNA content. In agreement with previous reports [5], hormone stimulated growth of

vector-expressing cells and knockdown of CHK2 significantly augmented growth in all

PCa cell lines tested (Figure 3). Here growth induced by CHK2 knockdown is likely an

increase in cell proliferation, as the basal apoptotic rate in the PCa lines is low. Re-

expression of wtCHK2, K373E, and T387N in CHK2-depleted cells markedly

suppressed the increase in growth induced by CHK2 knockdown in all PCa cell lines

tested. Interestingly, in LNCaP cells the extent of growth inhibition induced by K373E

and T387N was significantly less than that generated by wtCHK2 (Figure 3A). The

effect of CHK2 re-expression on cell growth in castration-resistant C4-2 (Figure 3B) and

Rv1 (Figure 3C) cells was not significantly different between wtCHK2 and the kinase

deficient K373E and T387N mutants, although the magnitude of inhibition caused by the

mutants was consistently less than that produced by wtCHK2. Expression levels of

wtCHK2, K373E, and T387N do not account for the difference observed between

LNCaP and C4-2 or Rv1 (Figure 3) since the relative expression of wtCHK2, K373E,

and T387N were similar across the cell lines. These observations suggest that CHK2

kinase activity limits the ability of CHK2 to negatively regulate PCa cell growth,

especially in androgen dependent PCa.

CHK2 suppresses IR induction of DNAPK and RAD54

Reports in the literature suggest that the AR is a critical regulator of genes in the

DDR [2]–[4] Therefore we evaluated the impact of CHK2 knockdown on IR induced

transcription of DDR genes in LNCaP cells (Figure 4). In our experiments, androgen or

IR only modestly affected DNAPK and RAD54 transcript levels; we did not observe

androgen or IR induction of XRCC2, XRCC3, XRCC4, XRCC5, MRE11, RAD51,

FANC1 and BRCA1 transcripts as reported by others (data not shown) [2], [3]. This

discrepancy is consistent with the observation that androgen regulation of DDR genes is

specific to the model system, experimental conditions, and disease state examined [4].

Knockdown of CHK2 in LNCaP cells grown in CSS and stimulated with 1nM DHT led to

an increase in transcription of DNAPK and RAD54 in response to IR, suggesting that

CHK2 may suppress AR transcription of DDR genes in response to IR.

CHK2 effect on IR sensitivity and DNA repair

We next examined the impact of CHK2 interactions on cell survival and

sensitivity to IR (Figure 5A). LNCaP cells expressing CHK2 (pLKO) or depleted of CHK2

(CHK2 KD) were exposed to increasing doses of IR, plated, and allowed to grow for 14

days. Clonogenic assays revealed that CHK2 knockdown promoted cell survival

following ionizing radiation. This data, along with the data above indicating that CHK2

suppresses AR transcription of DNA repair genes suggested the hypothesis that loss of

CHK2 could facilitate DNA repair.

To assess the effect of IR-induced DNA damage in the presence or absence of

CHK2 knockdown we performed comet assays to measure DNA breaks [28], and

immunofluorescence staining of γH2AX, a marker for DNA double-strand breaks [29],

[30]. When we examined DNA damage using the comet assay, we observed statistically

significant less DNA breaks with CHK2 knockdown at 24hrs in LNCaP cells (Figure 5B;

Figure 5-figure supplement 1). The comet assay shows less % DNA in tail, shorter DNA

tail length, and lower tail moment with CHK2 knockdown suggesting more efficient DNA

repair when CHK2 is compromised. In order to quantify DSB repair activity in an

unbiased manner, we developed an approach that utilizes automated quantitation of

γH2AX and 53BP1 foci detected with immunofluorescence that enables us to rapidly

determine the signal intensity (Figure 6-figure supplement 1). Rv1 cells were transduced

with vector or CHK2 shRNA 48hrs before IR. Radiation induced more γH2AX foci at

15min when CHK2 was knocked down (Figure 6A, 6B). However, less foci are found in

CHK2 knock-down cells at 2hrs and by 4hrs through 24hrs an equal number of foci are

present in control and CHK2 knock-down cells. H2AX is phosphorylated in response to

inputs in addition to IR induced DNA double strand breaks, including RNA polymerase II

dependent transcription [31]–[33]. Therefore, we also measured 53BP1 foci to assess

DSB repair and found no significant differences when CHK2 was knocked down (Figure

6C). The apparent discrepancy in the 53BP1 and γH2AX foci may be explained by an

increase in transcription when CHK2 is knocked down, as shown in Figure 4 and in our

previous study [5].

Since supraphysiological doses of androgen has been associated with

transcription dependent double strand breaks [34], [35], we also tested if CHK2

knockdown would augment supraphysiological androgen dependent γH2AX foci. LNCaP

and Rv1 cells were transduced with vector or CHK2 shRNA 48 hours before treatment

with a range of the synthetic androgen R1881 up to 100nM for 6 hours (Figure 6-figure

supplement 2). We saw modest hormone induced γH2AX foci and no change with

CHK2 knockdown. The supraphysiologic dose of androgen likely maximally activates

AR dependent transcription negating the increase in AR transcriptional activity typically

observed with CHK2 knockdown.

DISCUSSION

PCa is the most frequently diagnosed cancer and the second leading cause of cancer death among American men, with approximately 88 men dying from PCa every day (pcf.org). While androgen deprivation therapy (ADT) is effective initially, most patients will relapse and develop incurable CRPC. Recently, there has been an emphasis on understanding the link between the DDR and AR, since radiation is a standard of care for locally advanced PCa where the AR is a major driver, and PARP inhibitors may be efficacious in CRPC patients with mutations in DDR genes [2], [3], [5], [36]–[38].

Our study identifies AR as a direct interacting protein with CHK2 in PCa cells. Several studies elucidated the role of DDR protein-AR interactions in modulating AR transcriptional activity. PARP-1 was recruited to AR binding sites, enhancing AR occupancy and transcriptional function [39]. Tandem mass spectroscopy analysis identified Ku70 and Ku80 as direct AR-interacting proteins that positively regulate AR transactivation [40]. Furthermore, BRCA1 physically interacted with the DNA-binding domain (DBD) of AR to enhance AR transactivation and induce androgen-mediated cell death through p21 expression [41]. In contrast, the association of the LBD of AR with hRad9, a crucial member of the checkpoint Rad family, suppressed AR transactivation by preventing the androgen-induced interaction between the n-terminus and c-terminus of AR [42]. Other groups reported non-genomic effects as a result of DDR protein-AR interactions. Mediator of DNA damage checkpoint protein 1 (MDC1), an essential player in the Intra-S phase and G2/M checkpoints, physically associated with FL-AR and AR^{V7} to negatively regulate PCa cell growth and migration [43]. Yin and colleagues, on the

other hand, showed that increased clonogenic survival following IR was a consequence

of DNA-PKc directly complexing with both FL-AR and AR^{V5-7}, with radiation increasing

these interactions and enzalutamide blocking the association with FL-AR but not AR^{V5-7}

[44]. These data support a model where AR is integrated in the DDR, interfacing at

multiple points.

We show that the direct association of CHK2 and AR by Far Western requires

phosphorylation of AR on S308. Since proteins containing FHA domains bind

phosphoproteins [9], we hypothesize that AR interacts with CHK2 through the CHK2

FHA domain. In support of this, the Zhao lab determined that AR physically associated

with the FHA domain of another critical DDR member, MDC1 [43]. Expression of

truncation mutants of different MDC1 domains in LNCaP cells led to the discovery that

AR only co-immunoprecipitated with MDC1 mutants containing the FHA domain in the

absence and presence of dihydrotestosterone. Their results indicated that the FHA

domain of MDC1 mediated the interaction with AR.

Here we report that AR phosphorylation on S81 and S308 is required for the IR

induction of AR – CHK2 binding by co-IP. Interestingly neither of these phosphorylation

sites were altered by IR. AR S81 and S308 can both be phosphorylated by CDK1,

which is downstream of canonical CHK2 signaling [21], [27], and was the motivation for

examining these sites in response to IR. The prediction is that IR would lead to a

decrease in S81 and S308 phosphorylation. However, our previous studies

demonstrated that S81 is predominantly phosphorylated by CDK9, and thus is more

indicative of AR transcriptional activity [20]. We also found that S308 phosphorylation

was restricted to late G2 and M phase of the cell cycle [27]. CDK9 phosphorylation of

S81 and the restriction of S308 phosphorylation to G2/M likely accounts for the lack of

significant change in these phosphorylation sites in response to IR. The disconnect

between these sites being required for the IR induced increase in AR – CHK2

association but not being regulated by IR suggests that the hormone induced activation

state of the AR is a critical determinant in the IR induced increase in AR – CHK2

association. We observed differences when testing the AR – CHK2 association by Far

Western and co-IP. The direct protein-protein interaction is only impacted by loss of

S308 phosphorylation, however, basal co-IP of AR and CHK2 is not affected by AR

phosphorylation. Only the IR induced AR – CHK2 association is impacted by loss of AR

phosphorylation on S81 and S308. This suggests that additional proteins present in

cells may assist and participate in the AR – CHK2 interaction.

We found that CHK2 variants with diminished kinase activity impaired the IR-

induced increase in AR – CHK2 interaction but did not completely block the interaction.

As with AR phosphorylation, there were differences when the CHK2 mutant association

with AR was examined by Far Western or co-IP suggesting a more complicated

dynamic of AR – CHK2 association in cells. The PCa associated CHK2 mutants also

exhibited a reduced inhibition of cell growth. CHK2-depleted cells re-expressing CHK2

variants exhibited an approximate 2-3-fold reduction in growth inhibition in response to

hormone when compared to cells re-expressing wtCHK2. Moreover, the fold change in

growth suppression between wtCHK2 and CHK2 variants was greater in androgen-

dependent LNCaP cells than in castration-resistant C4-2 and Rv1 cells, suggesting that

in hormone sensitive PCa CHK2 variants may play a larger role in regulating growth.

This raises the possibility that the CHK2 mutants found in PCa with diminished AR

binding are selected for, decreasing CHK2 suppression of AR activity and PCa cell

growth. Berge and colleagues discovered numerous CHK2 splice variants in breast

cancer tissue, where all variants were co-expressed with wtCHK2 [45]. Furthermore,

several of these variants reduced kinase activity when simultaneously expressed with

wtCHK2 and displayed a dominant negative effect on wtCHK2. The impact of the CHK2

variants found in PCa on wtCHK2 function has not yet been fully explored.

In our experiments examining AR – CHK2 binding, ERK was used as a positive

control for a protein that interacts with CHK2 [24]. Interestingly, we observed a

significant increase in CHK2–ERK association with IR. This is consistent with IR

increasing CHK2 T68 phosphorylation, which is required for the CHK2–ERK interaction.

These data point to a potentially larger role for CHK2 beyond canonical DDR and cell

cycle checkpoint signaling; consistent with this notion CHK2 has been implicated in

diverse cellular processes [46]–[48]. MEK inhibition is effective in lung tumors with ATM

mutations where CHK2 is inactive [49] providing further support that CHK2 negatively

regulates ERK. Negative regulation of both the AR and ERK by CHK2 suggests the

hypothesis that CHK2 may serve as a general negative regulator of mitogenic signals in

response to IR.

Multiple studies indicate that the AR is a critical regulator of genes in the DDR

[2], [3]. Reports by others demonstrate that androgen and IR increased DNAPK,

XRCC2, and XRCC3 [3]. This concept was supported by more global analysis of

transcripts demonstrating androgen regulation of DDR genes [2]. Another global

analysis identified a different set of DDR genes are regulated by AR [4]. Our test by RT-

qPCR of ten different DDR genes found only a modest induction by IR or androgen

treatment in two DDR genes, DNAPK and RAD54. The discrepancy in the DDR gene

subsets regulated by the AR in these studies and ours may be system dependent.

However, it also indicates that the dogma of AR regulating DDR genes needs further

examination. Our data indicating that CHK2 knockdown increases DNAPK and RAD54

transcript levels is consistent with our earlier observation that CHK2 knockdown led to

the increase in the transcripts of canonical AR target genes [5]. This leads to the

hypothesis that CHK2 binding to the AR suppresses AR transcription of DDR genes

enabling cells to turn off the DDR following DNA repair.

We observed radiation resistance in CHK2 knockdown cells. We also observed

an increase in γH2AX signal when CHK2 was knocked down, and a decrease in DNA

breaks as measured by the comet assay under similar conditions. However, when

53BP1 signal was measured to assess DSB repair we found no significant differences

when CHK2 was knocked down. These conflicting results of γH2AX and 53BP1 signal

may be explained by phosphorylation of H2AX in response to transcription induced DNA

breaks [31]–[33]. These incongruous results may also be due to competing effects of

CHK2 as both a regulator of the cell cycle and apoptosis [46].

The data reported herein along with our previous work [5] indicate that CHK2

acts as a tumor suppressor in PCa, either through loss of expression or mutation. This

raises the concern that CHK2 antagonists in clinical development may paradoxically

lead to enhanced PCa growth and resistance to IR. However, it is important to note that

we have predominately used an RNAi/overexpression approach. Our RNAi approach is

more similar to the CHK2 variants in PCa that have reduced kinase activity. It is

important to consider that pharmacologic inhibition is different than inhibition by RNAi

[50]. A pharmacologic approach that provides a sudden and complete inhibition of

CHK2 kinase activity may impact PCa differently than our RNAi approach, especially

when combined with IR or AR antagonists. However, our work and that in the literature

also suggests that approaches downstream of CHK2 may be more straightforward than

targeting CHK2.

In this study, we presented data that provides mechanistic insight into our

observation that CHK2 negatively regulates PCa growth. We demonstrated that AR

directly bound CHK2, and that IR elevated the AR – CHK2 interaction. Not only did

these AR – CHK2 protein complexes require AR phosphorylation, but CHK2 kinase

activity was also necessary. Furthermore, these CHK2 mutants found in PCa exhibited

a diminished effect on restricting prostate cancer cell growth. We observed that

knockdown of CHK2 led to an increase in PCa cell survival and resolution of DNA DSBs

in response to IR. This suggests that the deregulation of CHK2 in PCa can confer

resistance to radiation. In a previous study, we showed that CHK2 knockdown

hypersensitized PCa cells to castrate levels of androgen and increased AR

transcriptional activity on both androgen-activated and androgen-repressed genes [5].

As part of a feedback loop, AR transcriptionally represses CHK2 levels. Thus, these

data along with our previously published results suggest the following model (Figure 7).

Wild-type CHK2 maintains homeostasis through mediating the balance between DNA

repair and cell death, antagonizes the AR through direct binding and inhibition of

transcription of AR target genes while AR transcriptionally represses CHK2. This

feedback loop enables AR repression of CHK2 to turn off the DDR, while CHK2

negatively regulates AR to dampen growth during repair of damaged DNA. CHK2

mutation or loss of expression in PCa leads to increased AR transcriptional activity,

possibly of DDR genes, and survival in response to DNA damage, all leading to a more

aggressive cancer. Collectively, the work provides a foundation for the continued study

of AR – CHK2 interactions and functional consequences to benefit PCa therapies.

MATERIALS AND METHODS

Cell culture

LNCaP (RRID:CVCL_0395) and C4-2 (RRID:CVCL_4782) cells (a gift from Dr. L. W. K. Chung) were grown in DMEM:F12 (Invitrogen) with 5% non-heat inactivated serum (Gemini) and 1% Insulin-Transferrin-Selenium-Ethanolamine (ITS) (ThermoFisher). CWR22Rv1 (Rv1) (RRID:CVCL_1045) (gift from Drs. Steven Balk) and 293T (RRID:CVCL_0063) cells (gift from Dr. Tim Bender) were grown in DMEM (Invitrogen) with 10% heat-inactivated serum. For growth experiments, phenol-red free DMEM:F12 media with 5% Charcoal-Stripped Serum (CSS) (Sigma) was used. Commercial DNA fingerprinting kits (DDC Medical) verified cell lines. The following STR markers were tested: CSF1PO, TPOX, TH01, Amelogenin, vWA, D16S539, D7S820, D13S317 and D5S818. Allelic score data revealed a pattern related to the scores reported by the ATCC, and consistent with their presumptive identity. Cell lines are tested monthly for mycoplasm or three passages following reconstitution from frozen stocks.

Reagents

Transfection: Fugene 6 (Promega); TransIT-2020 (Mirus Bio). Inhibitors: Enzalutamide (Selleck Chemicals), BML-277 (Santa Cruz Biotech). Antibodies: CHK2 (2G1D5), pCHK2 T68, ERK1/2 (137F5), Actin, FLAG-Tag, V5-Tag, HA-Tag, γH2AX (Cell Signaling); AR, pAR S308 (in-house); pAR S81 (Millipore); Cy3-labeled donkey anti-rabbit (Jackson ImmunoResearch). Western blotting performed as previously described [5].

Far Western Blot

To measure direct protein interactions, the protocol was adapted from Prickett *et*

al [17] and Wu *et al* [18]. 293T cells were transfected with FLAG-wtAR, FLAG-S81A,

S81D, S308A, S308D, HA-wtAR, FLAG-ERK2, V5-wtCHK2, FLAG-CHK2, FLAG-CHK2

mutants K373E or T387N, or empty vector control. Whole cell extracts were made using

Triton-X lysis buffer, sonicated, and immunopurified using anti-FLAG, anti-HA, or anti-

V5 beads (Sigma) for 2hr at 4°C. Protein bound to beads was washed three times with

Triton-X lysis buffer, eluted with 35μl 2X sample buffer, and boiled for 5 min. Proteins

were resolved by SDS-PAGE and transferred to PVDF membrane. Proteins on the

membrane were denatured and renatured in buffers with varying guanidine–HCl

concentrations. Membranes were blocked in 3% blocking buffer (3% bovine serum

albumin in Tris-buffered saline/Tween 20) for 1hr. Probes were diluted in 3% blocking

buffer and incubated overnight at 4°C. Membranes were washed three times with PBS

for 5min followed by fixation using 0.5% paraformaldehyde for 30 min at room

temperature. Membranes were then rinsed quickly twice with PBS and quenched using

2% glycine in PBS for 10 min at room temperature. The membrane was blotted for

FLAG, HA, V5, AR, CHK2, or ERK1/2 and analyzed using the LI-COR Odyssey system

and software.

Immunoprecipitation

CHK2 or AR protein was immunoprecipitated from 1mg cell lysate from LNCaP

cells transiently transfected with FLAG-wtAR/FLAG-S81A/FLAG-S308A or FLAG-

wtCHK2/Flag-K373E/Flag-T387N plus HA-wtAR for 48hrs; treated with radiation.

Immunoprecipitations (IP) were performed with either agarose or magnetic beads,

proteins were separated by 7.5% SDS-PAGE; and immunoblotted with AR, pAR S81,

pAR S308, CHK2, pCHK2 T68, HA, or ERK1/2 antibodies. Non-cropped versions of the

Far Western and IP Western blots are shown in Figure 3-figure supplement 1.

CyQuant Growth Assays

Assay was performed as previously described [5]. Briefly, shCHK2-209, shCHK2-

588, wtCHK2, K373E, T387N or Vector control virus was added to fibronectin-coated

(1μg/ml) 96well plates. Constructs of CHK2 wild-type and variants were verified by

sequencing. Cells were plated in phenol-red free DMEM:F12 or DMEM media with 5%

CSS in the presence or absence of 0.05nM R1881. CyQuant reagent was added on day

7 according to the manufacturer's protocol (ThermoFisher). Quantification was

performed using a BioTek Synergy 2 plate reader.

qPCR

RNA isolation and quantitative real-time PCR (qPCR) were performed as

previously described [51], [52]. RNA concentrations were determined using a NanoDrop

2000 UV-Vis Spectrophotometer (Thermo Scientific). Primer sequences and annealing

temperature: DNAPKc   FW   (ATGAGTACAAGCCCTGAG);   DNAPKc   RV

(ATATCAGAGCGTGAGAGC)   (Tm=60deg).   RAD54B   FW

(ATAACAGAGATAATTGCAGTGG); RAD54B RV (GATCTAATGTTGCCAGTGTAG)

(Tm=60deg).   PSMB6   FW   (CAAACTGCACGGCCATGATA);   PSMB6   RV

(GAGGCATTCACTCCAGACTGG) (Tm=60deg). The relative standard curve method

was used to determine transcriptional fold changes as we have done previously [53]–

[56]. RNA starting quantities (SQ) were determined using a standard curve. The SQ

mean of DNAPKc or RAD54B was then normalized to the reference gene PSMB6.

Clonogenic Survival Assay

LNCaP cells were transduced with lentiviral particles expressing vector or CHK2

shRNAs and treated with 0-6Gy of radiation 72hrs after transduction. Cells were

trypsinized, counted, and appropriate numbers were plated in triplicate with the

appropriate growth media for colony formation assays (100 cells/0Gy, 200 cells/2Gy,

1000 cells/4Gy, and 6000 cells/6Gy). After 10-14 days, colonies consisting of 50-70

cells were counted using crystal violet. All counts were made by a single observer in a

blinded manner using a colony counter pen to maximize consistency and minimize bias.

Plotted is the surviving fraction (number of colonies counted/(number of cells seeded x

PE) where PE = plating efficiency = number of colonies counted/number of cells

seeded) following radiation.

Comet Assay

All Comet Assay steps were performed in the dark. Cells were washed twice with

PBS, scraped and suspended in PBS. Cells were combined with molten LMAgarose

(Trevigen, 4250-050-02) and placed into comet suitable slides. Samples were left at 4°C

for 15 minutes in order to create a flat surface. Slides were immersed in lysis solution

(Trevigen, 4250-050-01) for 40 minutes at 4°C and then in alkaline solution for 30min at

room temperature. Slides were electrophoresed in 200mM NaOH, 1mM EDTA in water;

pH>13 at 21Volts (300mA) for 30 minutes at 4°C. After the electrophoresis, slides were

gently drained, washed twice in dH2O for 10min and immersed in 70% ethanol for 5min.

Samples were left to dry overnight at RT. Samples were stained with SYBR Green at

4°C for 5 minutes and left to dry completely overnight. For quantification, images were

acquired using a fluorescence microscope (Olympus BX51, High-mag) equipped with a

20×, 0.5 NA objective and a camera (DP70). Images were acquired with DPController

software. Images were analyzed by ImageJ software and graphs generated using Prism

(GraphPad Software). All imaging was performed at ~24°C.

Immunofluorescence (IF)

LNCaP cells were transduced with lentivirus expressing vector or CHK2 shRNAs

on 1μg/ml fibronectin-coated coverslips and treated with radiation 48hrs after

transduction. Cells were allowed to recover from IR exposure up to 24hrs. Coverslips

were washed 3X with PBS, permeabilized with 0.2% Triton-X for 5mins, washed 3x

quick and 3x 5min with PBS-Tween 20 at 0.05% (PBS-T), blocked with 2%

FBS/BSA/donkey serum in PBS-T for 2hrs at room temperature, and incubated with

γH2AX or 53BP1 antibodies overnight at 4°C. Coverslips were then washed by rocking

for 3x quick and 3x 10min in PBS-T and incubated with donkey anti-rabbit or anti-mouse

secondary antibody (Invitrogen) for 1hr at room temperature. Coverslips were then

washed by rocking for 3x quick and 3x 10min in PBS-T and mounted with Vectashield

containing DAPI (ThermoFisher). Images were acquired with a LSM 880 confocal

microscope (Carl Zeiss) and signals were measured using ImageJ software.

Scientific Rigor

Each experiment was performed independently a minimum of three times and

each experiment had technical replicates for measuring the endpoint. An independent

experiment is defined as an experiment performed on a different day with a different cell

line passage number. The number of independent experiments is reported in each

figure legend. All data are shown, no outliers were removed. Statistical analysis was

performed using GraphPad Prism 8.2.1 and the test used is reported in each figure

legend.

ACKNOWLEDGEMENTS

We thank the members of the laboratories of Drs. Gioeli, Jameson, Bouton, Dudley, Kashatus, Park, Rutkowski, Smith, and Zong for helpful discussions. We would also like to acknowledge exceptional assistance from UVA's Advanced Microscopy Facility.

FUNDING

This work was supported by the National Cancer Institute [R01 CA178338 to DG]; Paul Mellon Urologic Cancer Institute; and University of Virginia Cancer Center Patient and Friends.

COMPETING INTERESTS

The authors declare no competing financial interests

REFERENCES

[1] N. Rhind and P. Russell, "Chk1 and Cds1: linchpins of the DNA damage and replication checkpoint pathways," *J. Cell. Sci.*, vol. 113 (Pt 22), pp. 3889–3896, Nov. 2000.

[2] W. R. Polkinghorn *et al.*, "Androgen receptor signaling regulates DNA repair in prostate cancers," *Cancer Discov*, vol. 3, no. 11, pp. 1245–1253, Nov. 2013, doi: 10.1158/2159-8290.CD-13-0172.

[3] J. F. Goodwin *et al.*, "A hormone-DNA repair circuit governs the response to genotoxic insult," *Cancer Discov*, vol. 3, no. 11, pp. 1254–1271, Nov. 2013, doi: 10.1158/2159-8290.CD-13-0108.

[4] K. Jividen, K. Z. Kedzierska, C.-S. Yang, K. Szlachta, A. Ratan, and B. M. Paschal, "Genomic analysis of DNA repair genes and androgen signaling in prostate cancer," *BMC Cancer*, vol. 18, no. 1, p. 960, Oct. 2018, doi: 10.1186/s12885-018-4848-x.

[5] H. Q. Ta *et al.*, "Checkpoint Kinase 2 Negatively Regulates Androgen Sensitivity and Prostate Cancer Cell Growth," *Cancer Res.*, vol. 75, no. 23, pp. 5093–5105, Dec. 2015, doi: 10.1158/0008-5472.CAN-15-0224.

[6] S. Matsuoka, M. Huang, and S. J. Elledge, "Linkage of ATM to cell cycle regulation by the Chk2 protein kinase," *Science*, vol. 282, no. 5395, pp. 1893–1897, Dec. 1998.

[7] S. T. Kim, D. S. Lim, C. E. Canman, and M. B. Kastan, "Substrate specificities and identification of putative substrates of ATM kinase family members," *J. Biol. Chem.*, vol. 274, no. 53, pp. 37538–37543, Dec. 1999.

[8] J. Ahn, M. Urist, and C. Prives, "The Chk2 protein kinase," *DNA Repair (Amst.)*, vol. 3, no. 8–9, pp. 1039–1047, Sep. 2004, doi: 10.1016/j.dnarep.2004.03.033.

[9] J. Li *et al.*, "Structural and functional versatility of the FHA domain in DNA-damage signaling by the tumor suppressor kinase Chk2," *Mol. Cell*, vol. 9, no. 5, pp. 1045–1054, May 2002.

[10] C. García-Limones *et al.*, "CHK2 stability is regulated by the E3 ubiquitin ligase SIAH2," *Oncogene*, vol. 35, no. 33, pp. 4289–4301, 18 2016, doi: 10.1038/onc.2015.495.

[11] L. Tsvetkov, X. Xu, J. Li, and D. F. Stern, "Polo-like kinase 1 and Chk2 interact and co-localize to centrosomes and the midbody," *J. Biol. Chem.*, vol. 278, no. 10, pp. 8468–8475, Mar. 2003, doi: 10.1074/jbc.M211202200.

[12] S. Yang, C. Kuo, J. E. Bisi, and M. K. Kim, "PML-dependent apoptosis after DNA damage is regulated by the checkpoint kinase hCds1/Chk2," *Nat. Cell Biol.*, vol. 4, no. 11, pp. 865–870, Nov. 2002, doi: 10.1038/ncb869.

[13] C. Stevens, L. Smith, and N. B. La Thangue, "Chk2 activates E2F-1 in response to DNA damage," *Nat. Cell Biol.*, vol. 5, no. 5, pp. 401–409, May 2003, doi: 10.1038/ncb974.

[14] A. Hirao *et al.*, "DNA damage-induced activation of p53 by the checkpoint kinase Chk2," *Science*, vol. 287, no. 5459, pp. 1824–1827, Mar. 2000.

[15] J. Bartek and J. Lukas, "Chk1 and Chk2 kinases in checkpoint control and cancer," *Cancer Cell*, vol. 3, no. 5, pp. 421–429, May 2003.

554 [16] C. Y. Peng, P. R. Graves, R. S. Thoma, Z. Wu, A. S. Shaw, and H. Piwnica-
555   Worms, "Mitotic and G2 checkpoint control: regulation of 14-3-3 protein binding by
556   phosphorylation of Cdc25C on serine-216," *Science*, vol. 277, no. 5331, pp. 1501–
557   1505, Sep. 1997.
558 [17] T. D. Prickett *et al.*, "TAB4 stimulates TAK1-TAB1 phosphorylation and binds
559   polyubiquitin to direct signaling to NF-kappaB," *J. Biol. Chem.*, vol. 283, no. 28, pp.
560   19245–19254, Jul. 2008, doi: 10.1074/jbc.M800943200.
561 [18] Y. Wu, Q. Li, and X.-Z. Chen, "Detecting protein-protein interactions by Far western
562   blotting," *Nat Protoc*, vol. 2, no. 12, pp. 3278–3284, 2007, doi:
563   10.1038/nprot.2007.459.
564 [19] S. Chen, Y. Xu, X. Yuan, G. J. Bubley, and S. P. Balk, "Androgen receptor
565   phosphorylation and stabilization in prostate cancer by cyclin-dependent kinase 1,"
566   *Proc Natl Acad Sci U S A*, 2006, [Online]. Available:
567   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Ci
568   tation&list_uids=17043241.
569 [20] V. Gordon *et al.*, "CDK9 regulates AR promoter selectivity and cell growth through
570   serine 81 phosphorylation," *Mol Endocrinol*, vol. 24, pp. 2267–80, 2010.
571 [21] S. Chen, S. Gulla, C. Cai, and S. P. Balk, "Androgen receptor serine 81
572   phosphorylation mediates chromatin binding and transcriptional activation," *J. Biol.*
573   *Chem.*, vol. 287, no. 11, pp. 8571–8583, Mar. 2012, doi:
574   10.1074/jbc.M111.325290.
575 [22] D. Gioeli *et al.*, "Androgen receptor phosphorylation. Regulation and identification
576   of the phosphorylation sites," *J Biol Chem*, vol. 277, pp. 29304–14, 2002.
577 [23] H. Zong *et al.*, "Cyclin D3/CDK11p58 complex is involved in the repression of
578   androgen receptor," *Mol Cell Biol*, vol. 27, pp. 7125–42, 2007.
579 [24] B. Dai *et al.*, "Functional and molecular interactions between ERK and CHK2 in
580   diffuse large B-cell lymphoma," *Nat Commun*, vol. 2, p. 402, 2011, doi:
581   10.1038/ncomms1404.
582 [25] M. Higashiguchi *et al.*, "Clarifying the biological significance of the CHK2 K373E
583   somatic mutation discovered in The Cancer Genome Atlas database," *FEBS Lett.*,
584   Oct. 2016, doi: 10.1002/1873-3468.12449.
585 [26] M. Wasielewski *et al.*, "Deleterious CHEK2 1100delC and L303X mutants identified
586   among 38 human breast cancer cell lines," *Breast Cancer Res. Treat.*, vol. 113, no.
587   2, pp. 285–291, Jan. 2009, doi: 10.1007/s10549-008-9942-3.
588 [27] Y. Koryakina, K. E. Knudsen, and D. Gioeli, "Cell-cycle-dependent regulation of
589   androgen receptor function," *Endocr. Relat. Cancer*, vol. 22, no. 2, pp. 249–264,
590   Apr. 2015, doi: 10.1530/ERC-14-0549.
591 [28] A. R. Collins, "The comet assay for DNA damage and repair: principles,
592   applications, and limitations," *Mol. Biotechnol.*, vol. 26, no. 3, pp. 249–261, Mar.
593   2004, doi: 10.1385/MB:26:3:249.
594 [29] J. Vignard, G. Mirey, and B. Salles, "Ionizing-radiation induced DNA double-strand
595   breaks: a direct and indirect lighting up," *Radiother Oncol*, vol. 108, no. 3, pp. 362–
596   369, Sep. 2013, doi: 10.1016/j.radonc.2013.06.013.
597 [30] W. M. Bonner *et al.*, "γH2AX and cancer," *Nature Reviews Cancer*, vol. 8, no. 12,
598   pp. 957–967, Dec. 2008, doi: 10.1038/nrc2523.

599 [31] I. Singh *et al.*, "High mobility group protein-mediated transcription requires DNA
600    damage marker γ-H2AX," *Cell Res.*, vol. 25, no. 7, pp. 837–850, Jul. 2015, doi:
601    10.1038/cr.2015.67.
602 [32] H. Bunch *et al.*, "Transcriptional elongation requires DNA break-induced signalling,"
603    *Nature Communications*, vol. 6, p. 10191, Dec. 2015, doi: 10.1038/ncomms10191.
604 [33] J.-H. Ji *et al.*, "De novo phosphorylation of H2AX by WSTF regulates transcription-
605    coupled homologous recombination repair," *Nucleic Acids Res.*, vol. 47, no. 12, pp.
606    6299–6314, Jul. 2019, doi: 10.1093/nar/gkz309.
607 [34] M. C. Haffner *et al.*, "Androgen-induced TOP2B-mediated double-strand breaks
608    and prostate cancer gene rearrangements," *Nat. Genet.*, vol. 42, no. 8, pp. 668–
609    675, Aug. 2010, doi: 10.1038/ng.613.
610 [35] P. Chatterjee *et al.*, "Supraphysiological androgens suppress prostate cancer
611    growth through androgen receptor-mediated DNA damage," *J. Clin. Invest.*, vol.
612    130, Jul. 2019, doi: 10.1172/JCI127613.
613 [36] J. Mateo *et al.*, "DNA-Repair Defects and Olaparib in Metastatic Prostate Cancer,"
614    *N. Engl. J. Med.*, vol. 373, no. 18, pp. 1697–1708, Oct. 2015, doi:
615    10.1056/NEJMoa1506859.
616 [37] L. Li *et al.*, "Targeting poly(ADP-ribose) polymerase and the c-Myb-regulated DNA
617    damage response pathway in castration-resistant prostate cancer," *Sci Signal*, vol.
618    7, no. 326, p. ra47, May 2014, doi: 10.1126/scisignal.2005070.
619 [38] Y. Yin *et al.*, "Chk1 inhibition potentiates the therapeutic efficacy of PARP inhibitor
620    BMN673 in gastric cancer," *Am J Cancer Res*, vol. 7, no. 3, pp. 473–483, 2017.
621 [39] M. J. Schiewer *et al.*, "Dual roles of PARP-1 promote cancer growth and
622    progression," *Cancer Discov*, vol. 2, no. 12, pp. 1134–1149, Dec. 2012, doi:
623    10.1158/2159-8290.CD-12-0120.
624 [40] G. L. Mayeur, W. J. Kung, A. Martinez, C. Izumiya, D. J. Chen, and H. J. Kung, "Ku
625    is a novel transcriptional recycling coactivator of the androgen receptor in prostate
626    cancer cells," *J Biol Chem*, vol. 280, pp. 10827–33, 2005.
627 [41] S. Yeh *et al.*, "Increase of androgen-induced cell death and androgen receptor
628    transactivation by BRCA1 in prostate cancer cells," *Proc. Natl. Acad. Sci. U.S.A.*,
629    vol. 97, no. 21, pp. 11256–11261, Oct. 2000, doi: 10.1073/pnas.190353897.
630 [42] L. Wang *et al.*, "Human checkpoint protein hRad9 functions as a negative
631    coregulator to repress androgen receptor transactivation in prostate cancer cells,"
632    *Mol. Cell. Biol.*, vol. 24, no. 5, pp. 2202–2213, Mar. 2004.
633 [43] C. Wang *et al.*, "MDC1 functionally identified as an androgen receptor co-activator
634    participates in suppression of prostate cancer," *Nucleic Acids Res.*, vol. 43, no. 10,
635    pp. 4893–4908, May 2015, doi: 10.1093/nar/gkv394.
636 [44] Y. Yin *et al.*, "Androgen Receptor Variants Mediate DNA Repair after Prostate
637    Cancer Irradiation," *Cancer Res.*, vol. 77, no. 18, pp. 4745–4754, 15 2017, doi:
638    10.1158/0008-5472.CAN-17-0164.
639 [45] E. O. Berge, V. Staalesen, A. H. Straume, J. R. Lillehaug, and P. E. Lønning,
640    "Chk2 splice variants express a dominant-negative effect on the wild-type Chk2
641    kinase activity," *Biochim. Biophys. Acta*, vol. 1803, no. 3, pp. 386–395, Mar. 2010,
642    doi: 10.1016/j.bbamcr.2010.01.005.

[46] L. Zannini, D. Delia, and G. Buscemi, "CHK2 kinase in the DNA damage response
and beyond," *J Mol Cell Biol*, vol. 6, no. 6, pp. 442–457, Dec. 2014, doi:
10.1093/jmcb/mju045.
[47] H.-H. Choi *et al.*, "CHK2 kinase promotes pre-mRNA splicing via phosphorylating
CDK11(p110)," *Oncogene*, vol. 33, no. 1, pp. 108–115, Jan. 2014, doi:
10.1038/onc.2012.535.
[48] T. Choudhuri, S. C. Verma, K. Lan, M. Murakami, and E. S. Robertson, "The
ATM/ATR signaling effector Chk2 is targeted by Epstein-Barr virus nuclear antigen
3C to release the G2/M cell cycle block," *J. Virol.*, vol. 81, no. 12, pp. 6718–6730,
Jun. 2007, doi: 10.1128/JVI.00053-07.
[49] M. Smida *et al.*, "MEK inhibitors block growth of lung tumours with mutations in
ataxia-telangiectasia mutated," *Nat Commun*, vol. 7, p. 13701, 06 2016, doi:
10.1038/ncomms13701.
[50] K. J. Jensen, C. B. Moyer, and K. A. Janes, "Network Architecture Predisposes an
Enzyme to Either Pharmacologic or Genetic Targeting," *Cell Syst*, vol. 2, no. 2, pp.
112–121, Feb. 2016, doi: 10.1016/j.cels.2016.01.012.
[51] V. Gordon *et al.*, "CDK9 regulates AR promoter selectivity and cell growth through
serine 81 phosphorylation," *Molecular Endocrinology (Baltimore, Md.)*, vol. 24, no.
12, pp. 2267–2280, Dec. 2010, doi: 10.1210/me.2010-0238.
[52] H. Whitworth *et al.*, "Identification of Kinases Regulating Prostate Cancer Cell
Growth Using an RNAi Phenotypic Screen," *PLoS ONE*, vol. 7, no. 6, Jun. 2012,
doi: 10.1371/journal.pone.0038950.
[53] M. L. Wong and J. F. Medrano, "Real-time PCR for mRNA quantitation,"
BioTechniques, vol. 39, no. 1, pp. 75–85, Jul. 2005, doi: 10.2144/05391RV01.
[54] S. A. Bustin, "Absolute quantification of mRNA using real-time reverse transcription
polymerase chain reaction assays," *J. Mol. Endocrinol.*, vol. 25, no. 2, pp. 169–193,
Oct. 2000, doi: 10.1677/jme.0.0250169.
[55] M. A. Gordon *et al.*, "Synergy between androgen receptor antagonism and
inhibition of mTOR and HER2 in breast cancer," *Mol. Cancer Ther.*, May 2017, doi:
10.1158/1535-7163.MCT-17-0111.
[56] H. Whitworth *et al.*, "Identification of Kinases Regulating Prostate Cancer Cell
Growth Using an RNAi Phenotypic Screen," *PLoS ONE*, vol. 7, no. 6, p. e38950,
2012, doi: 10.1371/journal.pone.0038950.
676

FIGURE LEGEND

Figure 1. CHK2 directly binds AR. 293T cells were transfected with Vector, FLAG-

wtAR, or the FLAG-AR mutants S81A, S81D, S308A, or S308D, FLAG-wtCHK2, FLAG-

CHK2 mutants K373E or T387N, FLAG-ERK2, HA-wtAR, or V5-wtCHK2 using Fugene

6. 48hrs following transfection, FLAG, HA, and V5 were immunoprecipitated, and far

western blotting was performed. Membranes were blotted with the following antibodies:

FLAG, HA, V5, AR, CHK2, and ERK2. Blots were visualized using the Odyssey CLx. (A)

Probe = V5-wild-type CHK2; Targets = FLAG-wtAR, or the FLAG-AR mutants S81A,

S81D, S308A, or S308D, and FLAG-ERK2. Representative blots are shown, n=3. (B)

Probe = HA-wtAR; Targets = FLAG-wtCHK2, FLAG-CHK2 mutants K373E or T387N.

Representative blots are shown, n=4. For (A) and (B), graphs are of relative AR – CHK2

binding. Error bars, SEM. Statistical differences were tested for using ANOVA with

Dunnett's multiple comparisons test.

Figure 2. AR phosphorylation and CHK kinase activity regulates AR – CHK2

association. (A) AR – CHK2 interactions requires AR phosphorylation on S81 and

S308. LNCaP cells were transduced with lentiviral particles expressing wtAR, S81A, or

S308A for 48hrs. Cells were irradiated with 6Gy, and CHK2 was immunoprecipitated

one hour after IR. Representative blots are shown. Blots were quantitated on Odyssey

LICOR imaging system. Plotted is the AR signal normalized to total CHK2 and

compared to untreated cells, n=3. Error bars, SEM. Statistical differences were tested

for using ANOVA with Sidak's multiple comparisons test. (B) Expression of CHK2

variants with reduced kinase activity inhibits the radiation-induced increase in AR –

CHK2 interactions. LNCaP cells were transfected with HA-wtAR, HA-S81A, or HA-S308

in combination with FLAG-wtCHK2 for 48hrs using TransIT-2020 (Mirus). Cells were

irradiated with 6Gy, and FLAG was immunoprecipitated using a magnetic bead system

one hour after IR. Representative blots are shown. Plotted is the HA-AR signal

normalized to total FLAG-CHK2 and compared to untreated cells. Error bars, SEM.

Blots were quantitated on Odyssey LICOR imaging system. Statistical differences were

tested for using ANOVA with Sidak's multiple comparisons test.

Figure 3. Wild-type CHK2 negatively regulates prostate cancer cell growth. (A)

LNCaP, (B) C4-2, and (C) Rv-1 cells were transduced with lentiviral particles expressing

vector, shCHK2-exon 12, or shCHK2-3'UTR in combination with wtCHK2, K373E, or

T387N in the presence or absence of 0.05nM R1881. CyQuant assay was performed 7

days after transduction. Cell growth was compared with untreated vector control and the

values were averaged across biological replicates. Error bars, SEM, n=3. Statistical

analysis was performed using ANOVA and Tukey test, * = p<0.01. Representative blots

of CHK2 expression are shown with ERK as a loading control.

Figure 4. CHK2 knockdown increases the transcription of DDR genes in the

presence and absence of radiation. Transcript levels of DDR genes in LNCaP cells

transduced with CHK2 shRNAs and pLKO vector control and grown in CSS

supplemented with 1nM DHT were measured by qPCR. 48hrs following transduction,

cells were exposed to 2Gy ionizing radiation and RNA was isolated 6 hours later. SQ

means were determined using standard curves, normalized to the reference gene

PSMB6, with the untreated pLKO condition set to 1x. Values were averaged across

biological replicates +/- standard error of the mean, n=3. Shown are the histograms for

(A) DNAPKc and (B) Rad54B in LNCaP cells. Statistical analysis was performed using

one-way ANOVA and Tukey's test. * = p<0.02.

Figure 5. CHK2-depleted cells show increased survival and reduced DNA damage

following radiation. (A) Knockdown of CHK2 desensitizes cells to IR. LNCaP cells

were transduced with lentiviral particles expressing pLKO or CHK2 shRNAs for 48hrs,

treated with 0-6Gy IR, and seeded at the appropriate cell number for colony survival

assays. Results were normalized to untreated pLKO control and fitted to a standard

linear quadratic model. Error bars, SEM. Statistical analysis was performed using the

Student's t-test, n=4-8, p<0.01. (B) LNCaP cells were transduced with lentiviral particles

expressing pLKO or CHK2 shRNAs for 48hrs, irradiated at the indicated doses at the

specified times and processed for comet assays. Shown is the %DNA in the comet tail,

comet tail length, and tail moment (% DNA x tail length), n=3 (209-592 cells measured

per condition), 5-95% confidence intervals. Right most plots are of pLKO and CHK2 KD

at 24hrs to illustrate the statistical difference marked on the graphs. Statistical

differences were tested for using Kruskal-Wallis test followed by Dunn's multiple

comparisons test.

Figure 6. CHK2-depleted cells show altered γH2AX foci kinetics. (A) Representative

images of γH2AX, CHK2, and AR immunostaining quantified in (B and C). (B) Rv1 cells

were transduced with lentiviral particles expressing empty vector or CHK2 shRNAs on

fibronectin-coated coverslips in the appropriate growth media. Cells were irradiated with

6Gy after 48hrs. Coverslips were processed for IF at 0, 15min, 2, 4, and 24hrs following

IR. Plotted is the (B) γH2AX signal and (C) 53BP1 signal, which equals the mean grey

value intensity x number of foci per nucleus, n=3 (378-565 nuclei measured per

condition), 5-95% confidence intervals. Statistical differences were tested for using

Kruskal-Wallis test followed by Dunn's multiple comparisons test.

Figure 7. Model of CHK2-AR. We hypothesize the following model. In response to IR,

CHK2 activation antagonizes AR through direct binding and inhibition of transcription of

AR targets. CHK2 mutation or loss of expression that occurs in PCa leads to sustained

AR transcriptional activity, an increase in DDR gene transcripts, and survival in

response to DNA damage.

Figure 1-figure supplement 1. Far Western Blot schematic. Purified proteins are

separated by SDS-PAGE (target proteins) and then transferred to a membrane.

Proteins on the membrane are renatured and then probed with purified proteins (probe),

cross-linked, and subsequently detected with antibodies.

Figure 2-figure supplement 1. AR S81 and S308 phosphorylation are not altered

with radiation. LNCaP and C4-2 cells were seeded in serum-supplemented growth

media for 48hrs, exposed to 6Gy IR, and AR was immunoprecipitated using a magnetic

bead system 1hr after radiation from 1mg cell extracts. Proteins were separated by

7.5% SDS-PAGE, and blotted with AR, pAR S81, pAR S308, CHK2, and ERK1/2.

Representative blots are shown. Plotted is the pAR signal normalized to total AR and

compared to untreated cells, n=3, no statistical difference between the groups by

ANOVA. Blots were quantitated on Odyssey LICOR imaging system.

Figure 5-figure supplement 1. LNCaP cells were transduced with lentiviral particles

expressing pLKO or CHK2 shRNAs for 48hrs, irradiated at the indicated doses at the

specified times and processed for comet assays. Shown are representative images of

quantitation shown in Figure 5B.

Figure 6-figure supplement 1. Automated quantitation of foci. Confocal images are

captured on a Zeiss LSM 880 at 40x oil using Zeiss Zen digital imaging software and

processed into individual TIFF files for each fluorescence channel using ImageJ. An

outline of nuclei (mask) from DAPI channel is created and applied to all channels.

Macro-enabled image processing to measure multiple fluorescence parameters with

background subtraction using the following parameters: area, IntDen, RawIntDen, Min

and Max, fluorescence, background, foci count, CorDen, CorMean. γH2AX signal is the

product of foci number per nuclei and mean grey value (IntDen/area).

Figure 6-figure supplement 2. CHK2 Knockdown does not alter hormone induced

γ**H2AX foci.** LNCaP and Rv1 cells were transduced with lentiviral particles expressing

empty vector or CHK2 shRNA on fibronectin-coated coverslips in whole media for

48hrs. Media was changed to CSS overnight and cells were treated with hormone for 6

hours. Plotted is the number of γH2AX foci per cell. n=3 for LNCaP and n=2 for Rv1. At

the same dose of hormone there was no statistical difference between vector control

and CHK2 knockdown by Kruskal-Wallis test followed by Dunn's multiple comparisons

test.

Figure 3-figure supplement 1. Non-cropped images of Western blots shown in the

manuscript.

Figure 1

Figure 1-figure supplement 1

Figure 2

Figure 2-figure supplement 1

Figure 3

Figure 1

Figure 2

Figure 3

Figure 3-figure supplement 1

Figure 4

Figure 5

pLKO   CHK2 KD

15'

2 hr

4hr

24 hr

Figure 5 - Figure Supplement 1

Figure 6

1) Create a mask from DAPI channel

2) Apply DAPI mask onto the γH2AX channel

3) Background subtraction

4) Maxima points (foci)   Example of output

Figure 6-figure supplement 1

Figure 6-figure supplement 2

wtCHK2

mutCHK2

Figure 7