Metabolic models of mixed culture fermentation diagnose bottlenecks and

predict strategies for improved medium-chain fatty acid production

Matthew J. Scarborough,[1,2,3]* Joshua J. Hamilton,[4] Elizabeth A. Erb,[5] Timothy J. Donohue[1,6]

and Daniel R. Noguera[1,2]

1. The Great Lakes Bioenergy Research Center, UW-Madison, Madison, WI

2. Department of Civil and Environmental Engineering, UW-Madison, Madison, WI

3. Department of Civil and Environmental Engineering, University of Vermont, Burlington,

VT

4. Department of Biochemistry, UW-Madison, Madison, WI

5. Department of Chemical and Biological Engineering, UW-Madison, Madison, WI

6. Department of Bacteriology, UW-Madison, Madison, WI

*corresponding author e-mail: mscarbor@uvm.edu

ABSTRACT

Multi-species microbial communities determine the fate of materials in the environment

and can be harnessed to produce beneficial products from renewable resources. In a recent

example, fermentations by mixed microbial communities have produced medium-chain fatty acids

(MCFA). Tools to predict, assess, and improve the performance of these communities, however,

are limited. To provide such tools, we constructed two metabolic models of a MCFA-producing

community fed a byproduct from a biofuel fermentation. The first model is a unicellular model

(iFerment156) that contains a diverse set of fermentation pathways, including reverse β-oxidation

for MCFA production while the second model (iFermGuilds564) separates fermentation activities

into functional guilds. Both models predicted an energetic advantage for this community to

produce octanoic acid as a major fermentation product. Simulations with iFermGuilds564

predicted that 6- and 8-carbon MCFA were largely produced by a sugar-consuming guild, while

short-chain fatty acids were mainly produced by a guild consuming lactate. These models represent

novel tools for exploring the role of mixed microbial communities on carbon recycling in the

environment, as well as on beneficial reuse of organic residues.

INTRODUCTION

Mixed microbial fermentations have benefited humanity throughout history.[1] Since the

mid-18th century, mixed culture fermentations have produced food, valuable chemicals, and

recovered energy from wastes.[2] Many mixed microbial fermentations involve the synergistic

activity of several functional groups to convert organic substrates into valuable products. In

anaerobic digestion, fermentation products like acetic acid and H_2 are used to support microbial

production of methane, thus enabling the recovery of a large fraction of the chemical energy

originally present in the complex organic substrates. While anaerobic digestion technologies are

well established, other strategies to expand the range of potential products of mixed culture

fermentations are only beginning to emerge, such as the carboxylate platform, which produces

medium-chain fatty acids (MCFA) as a potentially valuable set of bio-based products.[3, 4]

MCFA are an attractive carboxylate platform product because of their many industrial

uses.[5] Octanoic acid (C8), an 8-carbon linear monocarboxylic acid, is a particularly valuable

MCFA because of its numerous uses, high market value, and relative ease of recovery. Further,

when C8 is reduced to its corresponding alkane, it can substitute for octane in liquid transportation

fuels. Recent research has demonstrated the feasibility of using mixed culture fermentations to

transform renewable resources into MCFA.[6-8] While C8 is produced in some self-assembled

microbial communities that develop in MCFA-producing bioreactors, hexanoic acid (C6) is often

the primary MCFA, along with acetic and butyric (C4) acids.[6-13]

Mathematical models have been established for the design and optimization of methane

production via anaerobic digestion.[14] These models describe the kinetics and stoichiometry of

reactions attributed to the individual members of the community that are responsible for hydrolysis

of complex organics, the fermentation of hydrolysis products, the transformation of fermentation

products to acetic acid and H_2, and finally, the transformation of these intermediates to methane.[14]

The metabolic networks in these models are well defined, with interspecies H_2 transfer known to

regulate critical metabolic steps, and therefore, anaerobic digestion models are predictive of the

behavior of self-assembled methane-producing microbiomes.

In contrast, the microbial metabolic networks involved in converting complex organic

substrates to MCFA are not fully elucidated. Consequently, there is a lack of knowledge on how

to obtain a carboxylate platform microbiome that produces C8 as the main MCFA, or how to

minimize the accumulation of shorter-chain fatty acids. Initial models of mixed culture

fermentations that do not include methanogenesis have focused on simulating metabolic networks

for production of C4, propionic acid, acetic acid, ethanol, lactic acid and H_2 from glucose.[15]

Subsequent models took into account that multiple electron carriers (e.g., ferredoxin, NADH) play

different roles in these metabolic networks, enabling the simulation of energy-conserving electron-

bifurcating reactions during production of C4 by reverse β-oxidation.[16] While a recent model

considered the role of homoacetogenesis as a H_2-consuming member of mixed culture

fermentation, it was limited to producing C4 as the major MCFA.[17]

In addition, existing models of carboxylate platform communities do not test the potential

role of recently discovered energy-conserving mechanisms in anaerobic microbes[18] that may shape

the metabolic networks of these mixed culture fermentations. Therefore, one goal of this study was

to test the ability of different models that considered all the known metabolic networks and energy-

conserving mechanisms to predict bottlenecks and strategies to improve MCFA production. To do

this, we constructed a single unit model (iFerment156) and a guild-based microbial community

model (iFermGuilds564) with expanded capability compared to existing models, and tested their

predictive ability using experimental data from a MCFA-producing bioreactor. We used the

models to diagnose bottlenecks for improved MCFA production, with particular emphasis on

predicting factors and specific enzymes that could alter the amount and type of MCFA produced

by the mixed microbial community.

MATERIALS AND METHODS

A metabolic network of a mixed culture fermentation was assembled that contains

reactions that occur during anaerobic metabolism of a previously described lignocellulsoic

biorefinery residue.[19] The first network, iFerment156, contains 156 reactions and 105 metabolites,

models a mixed culture fermentation as a single unit, and expands the metabolic pathways and

substrate range of previously published mixed culture fermentation networks.[15-17] iFerment156 is

the most comprehensive metabolic model of mixed culture fermentation to date, describing

metabolism of exogenous substrates including glucans, glucose, xylans, xylose, and glycerol,

along with lactate, ethanol, acetate, CO_2 and H_2, which are potential intermediates in this mixed

culture fermentation (**Fig. 1**).

The second network, iFermGuilds564, represents activities of different guilds (i.e.,

functional classes of organisms) within the microbial community. The iFermGuilds564 network

describes the same reactions and metabolites as in the iFerment156 model but these are separated

into six functional guilds with each guild representing a subset of compartmentalized reactions.

iFermGuilds564 also includes additional transport reactions that simulate metabolite exchange

among guilds. Combining the six guilds, the iFermGuilds564 network includes 564 reactions and

390 metabolites (**Fig. 2**). The guilds in the iFermGuilds564 model represent previously described

organisms that perform fatty acyl chain elongation from sugars (SEOs), sugar fermenting

organisms that produce mostly or only lactate, acetate, and ethanol (SFOs) as fermentation

products, hydrogenic sugar fermenters (HSFs), organisms that perform fatty acyl chain elongation

from lactate (LEOs) or ethanol (EEOs), and homoacetogenic organisms (HAOs). SEOs include

organisms that utilize sugars and elongate intermediate products through reverse β-oxidation, such

as *Megasphaera* and *Caproicproducens*.[20, 21] SFOs include genera such as *Lactobacillus* and

Bifidobacterium, which can ferment hexoses and pentoses to lactate, acetate and ethanol.[22, 23] HSFs

include organisms that generate H_2 while producing fermentation products, such as members of

the *Coriobacteriaceae* family.[24] LEOs include organisms that perform reverse β-oxidation with

lactate, such as *Pseudoramibacter*[25] and *Ruminococcaceae* bacterium *CPB6*.[9] EEOs, such as

Clostridium kluyveri,[26] perform reverse β-oxidation with ethanol. HAOs produce acetate from H_2

and CO_2 and include species within the *Clostridium*, *Acetobacterium*, *Eubacterium,* and *Blautia*.[27-

29]

When constructing each metabolic network, reactions were added to capture the exchange

of substrates and products with the reactor (extracellular space).[30] Where possible, standard

metabolite and reaction abbreviations from the Biochemical, Genetic, and Genomic (BiGG)

knowledgebase were used.[31] All reactions were checked for mass and charge balance assuming an

intracellular pH of 7.0 with the dominant ionic form of all metabolites obtained from the BiGG

database.[31] Methods for including transport constraints are discussed in **Supplementary Text 1**.

Both metabolic models (iFerment156 and iFermGuilds564) are available as **Supplementary File**

1 and **Supplementary File 2**, respectively. Instructions for using the models are available at

https://github.com/mscarbor/Mixed-Culture-Fermentation-Models.

The production capabilities of both metabolic networks were simulated using constraint-

based methods.[32] These methods predict the steady-state flux distribution through a metabolic

network that is subject to physiochemical constraints. In this work, we used flux variability

analysis (FVA)[33] and parsimonious flux balance analysis (pFBA)[34] to predict flux distributions

through metabolic networks of mixed culture fermentation. When modeling bioreactors with the

presence of solute gradients across the cell membrane, we considered the ATP required for

transport of solutes as described in **Supplementary Text 1.** The objective function for all

modeling scenarios was to maximize ATP production. When simulating the iFermGuilds564

network for a steady-state condition, we required all individual guilds to achieve the same ATP

hydrolysis flux as a proxy for having the same growth rate, as the models do not include biomass

equations. All simulations were performed in Python 3.5 using the cobrapy package (v 0.13.4).[35]

Throughout this work, we present results in terms of electron equivalents (eeq), which

report the number of electrons potentially transferred to an acceptor if the reduced chemical were

completely oxidized (using factors provided in **Table S1** to convert from molar concentrations to

eeq concentrations). For each scenario, we calculated the overall reaction thermodynamics using

the predicted stoichiometry of the model and the standard free energy of formation ($\Delta G_f^{\prime 0}$) of

metabolites (**Supplementary File 3**) obtained from KBase.[36] We ensured that at least 50 kJ was

available per ATP produced under standard conditions at neutral pH.[37, 38]

RESULTS

Simulation of mixed culture fermentations with iFerment156

By constraining the iFerment156 model to produce defined products, we initially tested its

predictive ability with well-established fermentation conditions (See **Supplementary Text 2;**

Table S2). We also investigated predictions derived from an unconstrained iFerment156 model

(i.e., all pathways are on) under conditions where multiple substrates are simultaneously produced

and consumed within this unit. This modeling analysis is a simplified representation of

fermentation in a mixed microbial community where interspecies transfer of metabolites is

assumed to have no energetic penalty, and is similar to previously published models of mixed

culture fermentations.[15-17] For this analysis, we required that all possible solutions released ≥50 kJ

per mol ATP produced as a condition that simulated microbial growth.[37] Using FVA with this

condition, iFerment156 provided a range of solutions that maximized ATP production depending

on the substrates provided to the simulation (**Fig. 3**). For instance, when we examined the

utilization of glucose as the sole substrate, iFerment156 predicted multiple conditions that would

allow ATP production at 4.5 mol ATP per mol of glucose metabolized. One of these solutions

predicts that 61.1% of the eeq in glucose could be converted to C8 (with 22.2% and 16.7% of the

eeq converted to acetate and formate, respectively) (**Fig. 3; Table S3**). Another solution predicts

66.6% of the eeq converted to C6, with 16.6% eeq converted to acetate and 16.6% to formate

(**Table S3**). Homofermentation to acetate is yet another solution that yields the maximum ATP

production. Another prediction from this analysis is that fermentations that result in ethanol or

lactate production, although feasible and known to occur in nature, do not result in the maximum

possible ATP yields (**Fig. 3**). This prediction of iFerment156 agrees with the practice of using pure

cultures (with limited metabolisms) for industrial ethanologenic fermentations, since mixed

cultures would likely produce other unwanted fermentation products that help maximize ATP

production by the microbial community.

The iFerment156 model predicts similar results when xylose is provided as the sole carbon

source (**Fig. 3, Table S3**), albeit with a lower ATP production compared to glucose (3.75 mol ATP

per mol substrate). We found that providing glycerol as a sole carbon source to iFerment156 yields

multiple solutions that maximize ATP production (**Fig. 3**). On the other hand, a single solution

that maximizes ATP production is predicted by iFerment156 if either lactate or ethanol are sole

carbon sources. When providing lactate, this solution predicts stoichiometric conversion of lactate

to C8, while with ethanol, 92% of the eeq would be converted to C8 and 8% to H_2.

These results predict that microbial communities would energetically benefit from

simultaneously producing multiple fermentation products, which is consistent with experimental

observations of fermentation of carbohydrate-rich feedstocks.[6, 8, 39] Consequently, we used

iFerment156 to evaluate whether having multiple electron donors simultaneously available would

change the predictions of maximum ATP production. In particular, since acetate and H_2 are

ubiquitous fermentation products in mixed microbial fermentations, we used iFerment156 to

simulate co-utilization of either acetate or H_2 with the other substrates analyzed (**Fig. 3**). With

acetate as a co-substrate with glucose, xylose, or glycerol, iFerment156 predicted the same

multiple solutions that maximized ATP production in the absence of acetate (**Table S3**). This

observation predicts, that from the point of view of maximizing microbial growth, acetate is a final

fermentation product. In contrast, co-utilization of lactate and acetate is predicted by iFerment156

to increase the ATP yield and favor reverse β-oxidation with solutions that maximize ATP

production including butyrate, C6 or C8 production (**Fig. 3; Table S3**). This analysis also found

that acetate co-utilization with ethanol is predicted by iFerment156 to double the ATP yield

compared to ethanol as the sole carbon source (**Table S3**). In this case, solutions for optimal ATP

yields were obtained when butyrate, C6, or C8 were the main fermentation product, consistent

with the ability of known carboxylate platform systems to produce a variety of short- and medium-

chain products.[40-42] Overall, these iFerment156 simulations suggest that acetate co-utilization can

increase the range of accumulated products, but that this may be detrimental to processes targeting

production of a single or small set of compounds. Further, these results predict that acetate may be

a dead-end product in mixed culture fermentations unless other compounds, such as lactate or

ethanol, are either present in the growth substrate or accumulated as intermediates.

We also tested the impact of H_2 metabolism on mixed culture fermentation, since this has

previously been proposed as either a fermentation intermediate or endogenous reductant to drive

MCFA production.[19] During utilization of sugars and glycerol, iFerment156 predicted that the

presence of H_2 favors higher ATP yields if C8 is the only fermentation product produced (**Fig. 3**).

In the presence of ethanol, H_2 is not predicted to be consumed by iFerment156 since redox balance

cannot be maintained under these conditions. In the presence of H_2 and acetate, or H_2 and CO_2, C8

production is favored for maximizing ATP production. Thus, iFerment156 predicts that H_2

oxidation should improve C8 production because it would maximize the ATP production of the

overall microbial community.

In sum, iFerment156 can predict substrates that favor accumulation of different products

when a large set of biochemical reactions are available in a mixed culture fermentation (**Fig. 1**).

When sugars and glycerol are the main substrates, mixed culture fermentations could give rise to

a wide range of products since multiple sets of fermentation reactions yield the same maximum

ATP production for the community. When lactate or ethanol are the main substrates, a narrower

product spectrum may be expected. In addition, H_2 co-utilization may result in accumulation of a

narrower range of products from sugars and glycerol as H_2 utilization would increase overall ATP

yields. In a large-scale bioreactor, production of fewer products is often considered advantageous

for downstream processing and to improve overall process economics.

Simulation of bioreactor performance with a single-unit model (iFerment156)

The iFerment156 model predicts that multiple substrates could lead to C8 production (**Fig.**

3), but that ATP could be maximized by co-production of other fermentation products. We wanted

to assess if this single-unit model could simulate the observed product yields of a bioreactor fed

lignocellulosic biorefinery residues, a complex organic substrate.[19] The microbial community in

the bioreactor (derived from wastewater sludge) was fed lignocellulosic biorefinery residues for

96 days, at which time bioreactor performance was evaluated using metagenomic and

metatranscriptomic analyses along with end-product sampling.[19] At this time, the bioreactor

contained 11.7 g L^{-1} of volatile suspended solids, which was assumed to represent microbial

biomass. To represent organic substrate metabolism, the substrate uptake fluxes were set according

to the eeq consumed for compounds in the feedstock (**Table S4**). We constrained iFerment156 to

only contain reactions predicted by transcriptomic analyses to be active in the bioreactor microbial

community[19] and performed FVA to identify the range of end-products that would support

maximum ATP yields. Under these conditions, iFerment156 predicted that 90% of the eeq would

be converted to C8 (**Fig. 4A**), which is much higher than the 3% conversion to C8 observed in the

bioreactor. The total predicted ATP production rate is 1.37 mmol hr^{-1}. While this simulation of

iFerment156 did not correctly predict the range of products (butyrate, C6 and C8) measured in the

bioreactor, it did indicate that C8 production could be advantageous for maximizing ATP

production.

Because C6 and C8 are known biocides,[43, 44] it is possible that their production is limited

by their accumulation. To test this hypothesis, we performed simulations constraining this version

of iFerment156 to produce C6 and C8 at the rates measured in the bioreactor (**Table S3**). Using

these parameters, iFerment156 predicted an ATP production rate of 1.32 mmol hr^{-1} (**Fig. 4B**),

corresponding to 96% of the maximum ATP production rate predicted when C6 and C8 production

were not constrained. Under these conditions, iFerment156 predicted that 50% of the eeq were

converted to butyrate, approximately 1.5-fold higher than was observed in the bioreactor (34% of

the eeq, **Fig. 4B**). Although the discrepancies between predicted and measured MCFA levels

indicate limitations in the ability of iFerment156 to predict performance of this mixed microbial

community, the similar ATP yield predictions when the model was constrained to the observed C6

and C8 concentrations illustrates the difficulty of simulating the sum of all metabolic processes

with a single-unit model. However, energetic losses associated with interspecies transfer of

substrates may influence the predicted ATP yields and the resulting levels of fermentation

products. To investigate this, a functional guild fermentation model was developed, based on the

reactions present in iFerment156, as described in the next section.

Simulation of mixed culture fermentations with a functional guild model (iFermGuilds564)

In iFermGuilds564, the metabolic activities of a microbial community are partitioned among a number of functional guilds. This biological division of labor negates the free exchange of reducing equivalents (e.g., NADH, ferredoxin) and of many pathway intermediates (e.g., acetyl-CoA) between community members, ultimately constraining how individual organisms maintain redox balance, which can potentially impact ATP and product yields. For instance, the single unit model (iFerment156) cannot predict the production of lactate via a NADH-dependent dehydrogenase and re-consumption of lactate via an electron-confurcating dehydrogenase, activities that may happen in a microbial community that includes lactate producers and lactate consumers. Thus, to better represent the compartmentalized nature of microbial activities within microbiomes we developed a functional guild model (iFermGuilds564) to simulate mixed culture fermentations where community members exchange extracellular intermediates. The iFermGuilds564 model contains the fermentation pathways in six guilds (**Fig. 2**), with each guild including predictions on transport reactions to simulate uptake and release of intermediate metabolites.

Initial predictions with iFermGuilds564 were focused at evaluating the role of each community member on the performance of this mixed culture fermentation. First, we used iFermGuilds564 to predict the fermentation of glucose by each of the guilds individually (**Table 1)** using an approach similar to that used with the iFerment156 model. That is, we required a release of ≥50 kJ per mol ATP produced as a condition that simulated microbial growth. As expected, only the guilds with reactions needed to degrade carbohydrates were predicted to produce ATP under these conditions. Specifically, iFermGuilds564 predicted that sugar-elongating organisms (SEOs) are predicted to have the highest ATP yield and produce C8, CO_2,

and H_2 (**Table 1**). In addition, sugar-fermenting organisms (SFOs) are predicted by

iFermGuilds564 to produce both acetate and lactate, but not CO_2. In contrast, iFermGuilds564

predicts that HSFs have a higher ATP yield than SFOs and produce acetate, ethanol CO_2, and H_2,

while no growth was predicted for several guilds that represent other major members of the mixed

microbial community (**Table 1**). In total, simulations of these individual guilds show that they

vary in the amount of ATP that can be produced from the same substrate. In addition, these

simulations predict that all of the guilds produce less ATP than a single organism containing all of

these reactions (iFerment156, 4.5 mol ATP mol^{-1} glucose, **Fig. 3**).

Table 1. iFermGuilds564 predicted overall chemical reactions, ATP yields, thermodynamics, and carbon

capture by individual guilds fed glucose.

Functional Guild	Predicted Reaction	ATP Yield	$\Delta G^{0'} mol^{-1}$ ATP	Carbon Capture
SEOs	$C_6H_{12}O_6 \rightarrow 0.5\ C_8H_{15}O_2^- + H_2O + H_2 + 0.5\ H^+ + 2\ CO_2$	4.0	-69.3	66.7%
SFOs	$C_6H_{12}O_6 \rightarrow C_3H_5O_3^- + 1.5\ C_2H_3O_2^- + 2.5\ H^+$	2.5	-74.1	100%
HSFs	$C_6H_{12}O_6 + 0.5H_2O \rightarrow 2C_2H_6O_3^- + 0.5C_2H_6O + H_2 + 2H^+ + CO_2$	3.0	-87.9	83.3%
LEOs	*No Growth Predicted*			
EEOs	*No Growth Predicted*			
HAOs	*No Growth Predicted*			

To explore whether iFermGuilds564 would better simulate the experimental observations

in the MCFA-producing bioreactor, compared to iFerment156 (**Fig. 4**), we used a version of

iFermGuild564 with only four guilds, which represented the most abundant community members

in the bioreactor as determined by metagenome sequencing,[19] which were a *Lachnospiraceae*

(SEO guild), *Eubacteriaceae* (LEO guild), *Lactobacillus* (SFO guild), and *Coriobacteriaceae*

(HSF guild). Performance of these guilds were simulated with substrate constraints that were based

on a combination of experimental observations and predictions from previous metabolic network

reconstructions[19]: SEOs were constrained to use glucose, xylose and glycerol. SFOs could

consume glucose, xylose, xylans, glucans, and glycerol whereas HSFs were constrained to only

consume glucans, glucose and glycerol, and LEOs were restricted to consuming only lactate and

glycerol (**Table S5**).

To mimic steady-state bioreactor conditions, we constrained each iFermGuilds564 model

so that specific ATP production rates (mmol ATP gDCW^{-1} hr^{-1}) for each guild were equal, assumed

that the guilds were present according to their measured relative DNA abundance,[19] and performed

FVA optimizing for the overall ATP production rate of the community. Under these conditions,

iFermGuilds564 had multiple solutions that predicted a maximum ATP rate of 0.58 mmol ATP hr^{-}

^{1}. The solutions derived from these iFermGuilds564 simulations predict that a range of products

could be produced while maintaining the maximum ATP yield (**Fig. 4C**). This is different from

the predictions derived from the iFerment156 model, where single solutions were found that

maximized ATP yield (**Fig. 4A**). However, both iFerment156 and iFermGuilds564 predict that

maximum ATP yields would be attained if C8 is produced from organics in conversion residue

and they overpredict the percent of eeq that is required to be converted to C8 when compared to

experimental data (30-90%, **Fig. 4C**).

Constraining the iFermGuilds564 simulation of the above four guilds to produce C6 and

C8 based on their observed production in the bioreactor (**Fig. 4D**), improved its predictions, with

only a minor reduction in the predicted rate of ATP production (i.e., 0.57 mol ATP hr^{-1}, ~99% of

the maximum ATP yield predicted when C6 and C8 production was not constrained). This finding

supports the notion that toxicity of individual MFCAs may impact the observed product

composition and that the energetic cost of reduced C8 production is minimal. In these simulations,

the predicted levels of other end-products using iFermGuilds564 were better aligned with those

observed in a bioreactor that is fed with the lignocellulosic biorefinery residues (**Fig. 4D**).

The predicted ATP production rates of 1.32-1.37 mmol ATP hr^{-1} with iFerment156 (**Figs.**

4A and 4B) are about two-fold higher than the 0.57-0.58 mmol ATP hr^{-1} predicted in

iFermGuilds564 when the reaction sets are compartmentalized in functional guilds. This suggests

that compartmentalization of reaction sets into individual guilds (microbes) greatly influences ATP

production by mixed culture fermentation communities. Therefore, our analysis predicts that

functional guild metabolic models have advantages over unicellular models for assessing the

performance of mixed microbial communities like those in a carboxylate platform bioreactor.

Assessing interspecies metabolite flux using iFermGuilds546

Our previous data predict that the individual microbes within the MCFA-producing

microbial community perform different functions, compete for resources, and participate in the

interspecies transfer of metabolites that ultimately allows the lignocellulosic biorefinery residues

to be converted to short- and medium-chain fatty acids. Based on this, we sought to use

iFermGuilds564 to test our ability to model the predicted function of each guild in the

transformation of substrates as well as the production and transfer of intermediate metabolites

between microbes. For this, we constrained iFermGuilds564 to produce end-products at the rates

observed in the MCFA-producing bioreactor (**Table S3**), set the substrates consumed by each guild

as described above, and predicted fluxes using pFBA which calculates a single flux distribution

when the model is constrained to the observed end-product production rates. Under these

conditions, iFermGuilds564 predicted a production rate of 0.54 mmol ATP hr^{-1}, which is 93% of

the maximum ATP production rate derived from the FVA analysis (**Fig. 4C**). The agreement of

the predicted ATP production rates by these methods suggests that, as a whole, the bioreactor

community is functioning near its optimal growth rate.

Under these conditions, iFermGuilds564 predicted that the glycerol would be consumed

by SEOs, xylose consumption would be shared by SEOs and SFOs, glucose would exclusively be

consumed by HSFs, glucans consumed by SFOs and HSFs, and xylans exclusively consumed by

SFOs (**Fig. 5**). In addition, these iFermGuilds564 simulations predicted that SEOs would produce

all the C8, most of the C6, and some H_2; SFOs were predicted to perform homofermentative lactate

production; HSFs were predicted to perform heterofementative production of lactate, acetate,

ethanol, and H_2; LEOs were the predicted lactate consumers who produced a combination of

acetate, butyrate, C6 and H_2. When we modeled how the H_2 partial pressure might impact the

predicted conversion routes, a thermodynamic analysis of each guild indicates that these predicted

reactions are feasible at both high and low H_2 partial pressures (**Table 2**). Thus, these

iFermGuilds564 simulations indicated that, unlike conventional anaerobic digestion, MCFA

production via mixed culture fermentation may proceed under a wide range of H_2 partial pressures.

Table 2. Thermodynamics of the net chemical reactions for each functional guild. ΔG' values are calculated

at neutral pH at the noted H_2 partial pressures assuming all other compound concentrations are present at 1

mM.

Functional Guild	Reactor Population Name	Relative Abundance	Transport - associated ATP (mmol hr^{-1})	Growth - associated ATP (mmol hr^{-1})	Total ATP Production (mmol hr^{-1})	ΔG' mol^{-1} ATP	
						$P_{H2} = 1$ atm	$P_{H2} = 10^{-3}$ atm
SEO	*Lachnospiraceae*	45%	0.00766	0.243	0.250	-67.5	-75.5
SFO	*Lactobacillus*	40%	0.00	0.216	0.216	-73.8	-73.8
HSF	*Coriobacteriaceae*	10%	0.00143	0.0539	0.0554	-81.0	-83.0
LEO	*Eubacteriaceae*	5%	0.0390	0.0270	0.0660	-146	-163

These iFermGuilds564 simulations also predicted that lactate was the only product that

was re-consumed by the community. This prediction suggests that acetate and butyrate are

undesirable fermentation end-products if the goal is to maximize C6 or C8 production in the

bioreactor.

An additional prediction of iFermGuilds564is that LEOs, which only represented 5% of

the microbial community, were predicted to produce 72% of the H_2, 87% of the acetate, 100% of

the butyrate and no significant quantities of C6 and C8 (**Fig. 5A**). This predicted flux of lactate to

end-products that do not contribute to MCFA production suggests that LEOs could be a potential

bottleneck for high levels of MCFA production. This prediction is in apparent contradiction with

those from the iFerment156 simulations (**Fig. 3**) that suggested maximization of ATP production

would occur if C8 is the final product of lactate utilization. One possible explanation for this

discrepancy is that LEOs are growing at sub-optimal conditions in the bioreactor given the

imposed restrictions on growth rate (see below).

Predicted metabolic efficiency of guilds in a MCFA-producing bioreactor

The previous simulations with iFermGuilds564 assumed that the specific ATP production

rates (mmol ATP gDCW^{-1} hr^{-1}) for each guild were equal. This constraint was imposed to simulate

observed steady-state bioreactor conditions where the relative concentration of each guild remains

constant over time. However, the previous iFermGuilds564 simulations did not consider potential

variations in maintenance energy requirements across the different guilds. Since we lack pure

cultures of the organisms in the bioreactor, we used the metabolic models to estimate the efficiency

of ATP production for each guild within the reactor community. To do this, we constrained the

individual guild models to consume the substrates that they are predicted to metabolize in the

mixed culture community (**Fig 5**). Under these conditions, and in the absence of competition for

substrate or restrictions of needing to have a common specific ATP production rate for all the

guilds, we allowed the iFermGuilds564 simulation to predict the maximum yield of ATP used for

growth (**Table 3**). An ATP production efficiency was then calculated based on a comparing the

predicted growth-associated ATP production in the iFermGuilds564 model and the maximum ATP

yield in the single unit iFerment156 simulation (**Table 3**). The predicted ATP production

efficiencies of guilds vary widely from SEOs (96.4%) and SFOs (95.6%) to LEOs (14.1%) (**Table**

3). This result predicts that LEOs grow at sub-optimal conditions in the bioreactor where there is

no selective pressure for them to produce MCFA over shorter chain organic acids or other products.

Table 3. Predicted ATP production efficiency of each guild modeled in the bioreactor.

Guild	Predicted Growth ATP with iFermGuilds564 (mmol hr^{-1})	Maximum predicted Growth ATP (mmol hr^{-1})	Efficiency (%)
SEOs	0.244	0.253	96.4%
SFOs	0.217	0.227	95.6%
HSFs	0.054	0.067	80.6%
LEOs	0.027	0.191	14.1%
LEOs[1]	0.027	0.064	42.4%

[1]LEOs constrained to use electron confurcating lactate dehydrogenase exclusively

Another explanation for these findings is that the metabolic networks used in the simulation

of LEOs are not correct, causing the model to overpredict ATP production by this guild. In order

to test this hypothesis, we evaluated an alternative route for lactate metabolism in LEOs. That is,

LEOs were initially modeled as using two different lactate dehydrogenases for the transformation

of lactate to pyruvate, a NADH-dependent lactate dehydrogenase (LDH) and an electron

confurcating lactate dehydrogenase (ECLDH), in which lactate transformation is coupled to

ferredoxin oxidation as described for the anaerobic lactate-consuming *Acetobacterium woodii*.[45]

With these assumptions, the iFermGuilds564 simulation of the LEOs indicated preferential use of

LDH over ECLDH (**Fig. S1A**). Thus, we also ran simulations in which the LEOs exclusively use

ECLDH (**Fig. S1B**). We found that, if LEOs are forced to use ECLDH, their maximum predicted

ATP production drops from 0.191 to 0.0637 mmol ATP hr^{-1}, predicting an efficiency of 42.4% in

the bioreactor, rather than 14.1% (**Table 3**). While this scenario results in an improvement in the

prediction, it suggests that use of ECLDH alone does not fully explain the inefficiencies of LEOs.

In scenarios where the LEOs exclusive using ECLDH for lactate metabolism, the simulations

predict that the overall flow of carbon through the community is nearly identical to those in other

simulations (**Supplementary File 4**). In addition, while the overall ATP production rate for the

community remains at 0.54 mmol ATP hr^{-1} when lactate is metabolized by ECLDH but the

predicted energy metabolism of LEOs changes in several ways (**Supplementary Text 3**).

In sum, simulating the bioreactor community with iFermGuilds564 suggests that SEOs are

responsible for producing C8 in the bioreactor. While the model of LEOs contain all of the

reactions necessary for producing C8, these microbes are predicted to produce most of the acetate,

butyrate and H$_2$ in the bioreactor. LEOs are predicted to have low growth efficiencies and may

require large amounts of ATP for processes other than growth and transport. While the inclusion

of ECLDH results in a higher overall energetic efficiency for LEOs (42.4%), this is still lower than

the predicted efficiencies of the other guilds in the mixed microbial community (**Table 3**).

Combined these predictions suggest that LEOs use ECLDH and may have other ATP-consuming

processes not currently captured by the metabolic model. The iFermGuilds564 predictions also

suggest that SEOs are a better source for MCFA production than LEOs when fed a carbohydrate-

rich substrate. Further, H$_2$ and acetate were always predicted to be metabolic end-products rather

than intermediates in MCFA production, suggesting that acetate is not a productive intermediate

in MCFA production by this community and that electron transfer between species could be

improved to alter the suite of desired products.

Improved MCFA production by a mixed microbial community

The predictions derived from metabolic models like iFermGuilds564 could also help guide

the engineering of natural or synthetic communities that convert complex organic residues to a

desired set of products. While a reactor containing only SEOs is predicted by iFermGuilds564 to

favor C8 production (**Supplementary File 4**), 26% of the carbon from these organisms is predicted

to be lost as CO_2. To test if a synthetic microbial community could convert 100% of the carbon

and eeq to C8 when fed lignocellulosic biorefinery residues, we used iFermGuilds564 to assemble

an *in silico* community to convert all of the carbon in this feedstock to C8. To limit carbon loss as

CO_2 by the SEOs, we included HAOs in this synthetic community and simulated the addition of

an external electron donor, H_2, to drive homoacetogenic activity to convert endogenous CO_2 to

acetate that could undergo chain elongation. We tested the performance of this synthetic

community, under various conditions, including different ratios of each microbe, partial pressures

of H_2 and others.

One synthetic community that was predicted to convert all of the eeq in lignocellulosic

biorefinery residues and exogenous H_2 to C8 contained 90% SEOs and 10% HAOs, both of which

were predicted to both use exogenous H_2 (**Fig. 6**). In this modeled synthetic community, HAOs

were predicted to use H_2 to fix CO_2 produced by SEOs, and acetate produced by the HAOs was a

major source for chain elongation by the SEOs. Formate produced by SEOs would also be used

by the HAOs as a source of carbon and reducing power. Analyzing the scenarios predicted for

each guild indicate that a H_2 partial pressure $\leq 10^{-6}$ atm would make the predicted ATP produced

by HAOs thermodynamically infeasible. However, at H_2 partial pressures $\geq 10^{-3}$ atm, the

transformations and predicted ATP yields of the two-guild mixed culture fermentation were

sufficient to support C8 production (**Table 4**). Further, metabolic modeling of this synthetic

community predicts that increased C8 production could be achieved if HAOs could more readily

utilize H_2 that is produced or provided to the reactor.

Table 4. Predicted thermodynamics of a synthetic community of SEOs and HAOs converting conversion

residue and H_2 to C8.

Functional Guild	Relative Abundance	Predicted ATP Production (mol hr^{-1})	$\Delta G'$ mol^{-1} ATP	
			$P_{H2} = 1$ atm	$P_{H2} = 10^{-3}$ atm
SEO	90%	0.728	-100	-87.6
HAO	10%	0.081	-160	-86.8
Total		0.809	-106	-87.6

DISCUSSION

Under anaerobic conditions, a diverse set of mixed microbial communities catalyze the

transformation of organic substrates to a variety of products. Some fermentation metabolisms are

well understood, using genetically tractable pure cultures that can be analyzed, modeled and

improved for use in industrial applications.[46, 47] For over a century, anaerobic microbial

communities have been harnessed to recover energy, in the form of methane-rich biogas, from

complex organic matter in anaerobic digestion.[2] The ability to harness the metabolic activity of

mixed microbial communities to produce alternative fermentation products is limited since

metabolic and kinetic models that accurately predict the distribution of fermentation products are

only starting to emerge.[15-17] To address this knowledge gap, we constructed metabolic models of

mixed culture fermentations with the objective of gaining greater insights into how to increase

MCFA production by mixed microbial communities. We focused on producing C6 and C8, linear

medium-length MCFA of industrial value that can be produced from a variety of organic residues.[3]

Since many microbes in self-assembled fermentation communities remain uncultured, our

knowledge and ability to predict important metabolic networks of these mixed cultures relies

largely on information from metagenomics, metatranscriptomics and metabolomics analyses to

build the functional activities in each guild.[19, 48] We show in this work that metabolic modeling

offers additional insight that is not gained by these analyses, but that is critical in their evaluation

and in developing hypothesis of how to direct metabolism towards a specific set of fermentation

products. For instance, while iFermGuilds564 included three guilds capable of MCFA production

(SEOs, LEOs, and EEOs; **Fig. 2**), only SEOs and LEOs were included in many simulations since

reactor performance did not indicate ethanol as a fermentation product.[19] From prior analyses[19, 49]

we anticipated that both SEOs and LEOs participated in MCFA production, but the

iFermGuilds564 simulation of interspecies metabolite flux (**Fig. 5**) predicted that, to maximize the

overall ATP production in the community, C8 was largely produced by SEOs, while LEOs were

responsible for production of butyrate and acetate as fermentation end-products that were not

converted to MCFA. This finding makes the unexpected prediction that minimizing the abundance

or activity of LEOs would maximize MCFA production by this community.

We further showed that a guilds-based fermentation model such as iFermGuilds564 can be

used to evaluate scenarios for how to direct a mixed microbial community to produce specific

fermentation products from more efficient utilization of complex organic substrates (**Fig. 6**). For

example, iFerment156 predicted that H_2 is both a product during MCFA production (**Fig. 5**) and a

source of reducing power that could maximize MCFA production by simplifying the range of

accumulated fermentation products (**Fig. 3**). In addition, the results of this simulation predict a

previously unrealized role of HAOs, a guild that is not abundant in bioreactors transforming

lignocellulosic biorefinery residues to MCFA,[6, 19] but that could be enriched by an exogenous

supply of H_2 (**Fig. 6**). The use of metabolic models to assemble an ideal microbial community

represents an important new step to bioprocess design that include synthetic or self-assembled

communities optimized through the application of selective pressures such as H_2 supply.

iFerment156 and iFermGuilds564 are examples of first-generation predictive models of

mixed culture fermentations. Their utility can be improved by obtaining additional knowledge of

metabolic pathways in individual microbes and genetic studies to test predictions of the metabolic

potential of the community members that compliment multi-omic analyses. Further effort is

needed to simulate poorly appreciated conditions that shape community structure, such as the

toxicity of fermentation products on the specific guilds or the contribution(s) of low abundance

microbes to the overall activity of the mixed culture fermentation. Additionally, ours and many

models are limited to maximizing community ATP production as a proxy for microbial growth, so

computational methods to predict the abundance of specific guilds, rather than relying on relative

abundances as a model input, are needed. Efforts to assemble such models have begun for the

human gut microbiome,[50, 51] and as these approaches are applied to mixed microbial fermentations,

their benefits to the production of MCFA and other chemicals of societal importance will continue

to improve.

ACKNOWLEDGMENTS

This work was funded by the DOE Great Lakes Bioenergy Research Center (DOE BER

Office of Science DE-SC0018409). M.J.S. was supported by the National Science Foundation

Graduate Research Fellowship Program under grant No. DGE-1256259. JJH was supported by the

National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number

2016-67012-24709. The authors extend our gratitude to Robert Aidan Campbell for assistance

with python coding.

REFERENCES

1.   McGovern, P. E.; Zhang, J.; Tang, J.; Zhang, Z.; Hall, G. R.; Moreau, R. A.; Nunez, A.;
Butrym, E. D.; Richards, M. P.; Wang, C. S.; Cheng, G.; Zhao, Z.; Wang, C., Fermented beverages
of pre- and proto-historic China. *Proc. Natl. Acad. Sci. U. S. A.* **2004,** *101*, (51), 17593-8.

2.   Bond, T.; Templeton, M. R., History and future of domestic biogas plants in the developing
world. *Energy for Sustainable Development* **2011,** *15*, (4), 347-354.

3.   Angenent, L. T.; Richter, H.; Buckel, W.; Spirito, C. M.; Steinbusch, K. J. J.; Plugge, C.
M.; Strik, D.; Grootscholten, T. I. M.; Buisman, C. J. N.; Hamelers, H. V. M., Chain Elongation
with Reactor Microbiomes: Open-Culture Biotechnology To Produce Biochemicals. *Environ. Sci.*
Technol. **2016,** *50*, (6), 2796-2810.

4.   Chen, W. S.; Strik, D.; Buisman, C. J. N.; Kroeze, C., Production of Caproic Acid from
Mixed Organic Waste: An Environmental Life Cycle Perspective. *Environ. Sci. Technol.* **2017,**
51, (12), 7159-7168.

5.   Sarria, S.; Kruyer, N. S.; Peralta-Yahya, P., Microbial synthesis of medium-chain
chemicals from renewables. *Nat. Biotechnol.* **2017,** *35*, (12), 1158-1166.

6.   Scarborough, M. J.; Lynch, G.; Dickson, M.; McGee, M.; Donohue, T. J.; Noguera, D. R.,
Increasing the economic value of lignocellulosic stillage through medium-chain fatty acid
production. *Biotechnology for Biofuels* **2018,** *11*, (1), 200.

7.   Nelson, R.; Peterson, D.; Karp, E.; Beckham, G.; Salvachúa, D., Mixed Carboxylic Acid
Production by Megasphaera elsdenii from Glucose and Lignocellulosic Hydrolysate. *Fermentation*
2017, *3*, (1), 10.

8.   Kucek, L. A.; Xu, J. J.; Nguyen, M.; Angenent, L. T., Waste Conversion into n-Caprylate
and n-Caproate: Resource Recovery from Wine Lees Using Anaerobic Reactor Microbiomes and
In-line Extraction. *Front. Microbiol.* **2016,** *7*.

9.   Zhu, X.; Zhou, Y.; Wang, Y.; Wu, T.; Li, X.; Li, D.; Tao, Y., Production of high-
concentration n-caproic acid from lactate through fermentation using a newly isolated
Ruminococcaceae bacterium CPB6. *Biotechnol Biofuels* **2017,** *10*, 102.

10.   Kucek, L. A.; Nguyen, M.; Angenent, L. T., Conversion of L-lactate into n-caproate by a
continuously fed reactor microbiome. *Water Res.* **2016,** *93*, 163-171.

11.   Coma, M.; Vilchez-Vargas, R.; Roume, H.; Jauregui, R.; Pieper, D. H.; Rabaey, K.,
Product Diversity Linked to Substrate Usage in Chain Elongation by Mixed-Culture Fermentation.
Environ. Sci. Technol. **2016,** *50*, (12), 6467-76.

12.   Spirito, C. M.; Marzilli, A. M.; Angenent, L. T., Higher Substrate Ratios of Ethanol to
Acetate Steered Chain Elongation toward n-Caprylate in a Bioreactor with Product Extraction.
Environ. Sci. Technol. **2018,** *52*, (22), 13438-13447.

13.   Ge, S.; Usack, J. G.; Spirito, C. M.; Angenent, L. T., Long-Term n-Caproic Acid
Production from Yeast-Fermentation Beer in an Anaerobic Bioreactor with Continuous Product
Extraction. *Environ. Sci. Technol.* **2015,** *49*, (13), 8012-21.

14.   Batstone, D. J.; Keller, J.; Angelidaki, I.; Kalyuznyi, S. V.; Pavlostathis, S. G.; Rozzi, A.;
Sanders, W. T. M.; Siegrist, H.; Vavilin, V. A., The IWA Anaerobic Digestion Model No 1
(ADM1). *Water Sci. Technol.* **2002,** *45*, (10), 65-73.

15.   Rodriguez, J.; Kleerebezem, R.; Lema, J. M.; van Loosdrecht, M. C. M., Modeling product
formation in anaerobic mixed culture fermentations. *Biotechnol. Bioeng.* **2006,** *93*, (3), 592-606.

16.   Zhang, F.; Zhang, Y.; Chen, M.; van Loosdrecht, M. C. M.; Zeng, R. J., A modified
metabolic model for mixed culture fermentation with energy conserving electron bifurcation
reaction and metabolite transport energy. *Biotechnol. Bioeng.* **2013,** *110*, (7), 1884-1894.

17.   Regueira, A.; Gonzalez-Cabaleiro, R.; Ofiteru, I. D.; Rodriguez, J.; Lema, J. M., Electron
bifurcation mechanism and homoacetogenesis explain products yields in mixed culture anaerobic
fermentations. *Water Res.* **2018,** *141*, 349-356.

18.   Buckel, W.; Thauer, R. K., Flavin-Based Electron Bifurcation, Ferredoxin, Flavodoxin,
and Anaerobic Respiration With Protons (Ech) or NAD(+) (Rnf) as Electron Acceptors: A
Historical Review. *Front. Microbiol.* **2018,** *9*, 401.

19.   Scarborough, M. J.; Lawson, C. E.; Hamilton, J. J.; Donohue, T. J.; Noguera, D. R.,
Metatranscriptomic and thermodynamic insights into medium-chain fatty acid production using
an anaerobic microbiome. *mSystems* **2018**.

20.   Marounek, M.; Fliegrova, K.; Bartos, S., Metabolism and Some Characteristics of Ruminal
Strains of Megasphaera-Elsdenii. *Appl. Environ. Microbiol.* **1989,** *55*, (6), 1570-1573.

21. Kim, B. C.; Seung Jeon, B.; Kim, S.; Kim, H.; Um, Y.; Sang, B. I., Caproiciproducens
galactitolivorans gen. nov., sp. nov., a bacterium capable of producing caproic acid from galactitol,
isolated from a wastewater treatment plant. *Int. J. Syst. Evol. Microbiol.* **2015,** *65*, (12), 4902-8.

22. Ganzle, M. G.; Follador, R., Metabolism of oligosaccharides and starch in lactobacilli: a
review. *Front. Microbiol.* **2012,** *3*, 340.

23. Kandler, O., Carbohydrate metabolism in lactic acid bacteria. *Antonie Van Leeuwenhoek*
1983, *49*, (3), 209-24.

24. Dewhirst, F. E.; Paster, B. J.; Tzellas, N.; Coleman, B.; Downes, J.; Spratt, D. A.; Wade,
W. G., Characterization of novel human oral isolates and cloned 16S rDNA sequences that fall in
the family Coriobacteriaceae: description of olsenella gen. nov., reclassification of Lactobacillus
uli as Olsenella uli comb. nov. and description of Olsenella profusa sp. nov. *Int. J. Syst. Evol.*
Microbiol. **2001,** *51*, (Pt 5), 1797-804.

25. Holdeman, L. V.; Elizabeth, C. P.; Moore, W. E. C., Amended description of
Ramibacterium alactolyticum Prevot and Taffanel with propoosal of a neotype strain. *Int. J. Syst.*
Bacteriol. **1967,** *17*, (4), 323-341.

26. H. A. Barker, M. D. K., and B.T. Borenson, The synthesis of butyric and caproic acid from
ethanol and acetate by Clostridium
Kluyveri. *Proc. Natl. Acad. Sci. U. S. A.* **1945,** *31*, (12), 374-381.

27. Henderson, G.; Naylor, G. E.; Leahy, S. C.; Janssen, P. H., Presence of Novel, Potentially
Homoacetogenic Bacteria in the Rumen as Determined by Analysis of Formyltetrahydrofolate
Synthetase Sequences from Ruminants. *Appl. Environ. Microbiol.* **2010,** *76*, (7), 2058-2066.

28. Hugenholtz, J.; Ljungdahl, L. G., Metabolism and energy generation in homoacetogenic
clostridia. *FEMS Microbiol. Rev.* **1990,** *7*, (3-4), 383-9.

29. Balch, W. E.; Schoberth, S.; Tanner, R. S.; Wolfe, R. S., Acetobacterium, a New Genus of
Hydrogen-Oxidizing, Carbon Dioxide-Reducing, Anaerobic Bacteria. *Int. J. Syst. Bacteriol.* **1977,**
27, (4), 355-361.

30. Covert, M. W.; Schilling, C. H.; Famili, I.; Edwards, J. S.; Goryanin, I. I.; Selkov, E.;
Palsson, B. O., Metabolic modeling of microbial strains in silico. *Trends Biochem. Sci.* **2001,** *26*,
(3), 179-186.

31. King, Z. A.; Lu, J.; Drager, A.; Miller, P.; Federowicz, S.; Lerman, J. A.; Ebrahim, A.;
Palsson, B. O.; Lewis, N. E., BiGG Models: A platform for integrating, standardizing and sharing
genome-scale models. *Nucleic Acids Res.* **2016,** *44*, (D1), D515-22.

32. Bordbar, A.; Monk, J. M.; King, Z. A.; Palsson, B. O., Constraint-based models predict
metabolic and associated cellular functions. *Nat. Rev. Genet.* **2014,** *15*, (2), 107-20.

33. Gudmundsson, S.; Thiele, I., Computationally efficient flux variability analysis. *BMC*
Bioinformatics **2010,** *11*, 489.

34. Lewis, N. E.; Hixson, K. K.; Conrad, T. M.; Lerman, J. A.; Charusanti, P.; Polpitiya, A.
D.; Adkins, J. N.; Schramm, G.; Purvine, S. O.; Lopez-Ferrer, D.; Weitz, K. K.; Eils, R.; Konig,
R.; Smith, R. D.; Palsson, B. O., Omic data from evolved E-coli are consistent with computed
optimal growth from genome-scale models. *Mol. Syst. Biol.* **2010,** *6*.

35. Ebrahim, A.; Lerman, J. A.; Palsson, B. O.; Hyduke, D. R., COBRApy: COnstraints-Based
Reconstruction and Analysis for Python. *BMC Syst. Biol.* **2013,** *7*, 74.

36. Arkin, A. P.; Cottingham, R. W.; Henry, C. S.; Harris, N. L.; Stevens, R. L.; Maslov, S.;
Dehal, P.; Ware, D.; Perez, F.; Canon, S.; Sneddon, M. W.; Henderson, M. L.; Riehl, W. J.;
Murphy-Olson, D.; Chan, S. Y.; Kamimura, R. T.; Kumari, S.; Drake, M. M.; Brettin, T. S.; Glass,
E. M.; Chivian, D.; Gunter, D.; Weston, D. J.; Allen, B. H.; Baumohl, J.; Best, A. A.; Bowen, B.;
Brenner, S. E.; Bun, C. C.; Chandonia, J. M.; Chia, J. M.; Colasanti, R.; Conrad, N.; Davis, J. J.;
Davison, B. H.; DeJongh, M.; Devoid, S.; Dietrich, E.; Dubchak, I.; Edirisinghe, J. N.; Fang, G.;
Faria, J. P.; Frybarger, P. M.; Gerlach, W.; Gerstein, M.; Greiner, A.; Gurtowski, J.; Haun, H. L.;
He, F.; Jain, R.; Joachimiak, M. P.; Keegan, K. P.; Kondo, S.; Kumar, V.; Land, M. L.; Meyer, F.;
Mills, M.; Novichkov, P. S.; Oh, T.; Olsen, G. J.; Olson, R.; Parrello, B.; Pasternak, S.; Pearson,
E.; Poon, S. S.; Price, G. A.; Ramakrishnan, S.; Ranjan, P.; Ronald, P. C.; Schatz, M. C.; Seaver,
S. M. D.; Shukla, M.; Sutormin, R. A.; Syed, M. H.; Thomason, J.; Tintle, N. L.; Wang, D.; Xia,
F.; Yoo, H.; Yoo, S.; Yu, D., KBase: The United States Department of Energy Systems Biology
Knowledgebase. *Nat. Biotechnol.* **2018,** *36*, (7), 566-569.

37. Thauer, R. K.; Jungermann, K.; Decker, K., Energy conservation in chemotrophic
anaerobic bacteria. *Bacteriol. Rev.* **1977,** *41*, (1), 100-80.

38. Alberty, R. A., *Thermodynamics of biochemical reactions*. John Wiley & Sons, Inc. :
Hoboken, New Jersey, 2003.

39. Andersen, S. J.; De Groof, V.; Khor, W. C.; Roume, H.; Props, R.; Coma, M.; Rabaey, K.,
A Clostridium Group IV Species Dominates and Suppresses a Mixed Culture Fermentation by
Tolerance to Medium Chain Fatty Acids Products. *Front Bioeng Biotechnol* **2017,** *5*, 8.

40.   Roghair, M.; Hoogstad, T.; Strik, D. P. B. T. B.; Plugge, C. M.; Timmers, P. H. A.;
Weusthuis, R. A.; Bruins, M. E.; Buisman, C. J. N., Controlling Ethanol Use in Chain Elongation
by CO2 Loading Rate. *Environ. Sci. Technol.* **2018,** *52*, (3), 1496-1505.

41.   Kucek, L. A.; Spirito, C. M.; Angenent, L. T., High n-caprylate productivities and
specificities from dilute ethanol and acetate: chain elongation with microbiomes to upgrade
products from syngas fermentation. *Energy Environ. Sci.* **2016,** *9*, (11), 3482-3494.

42.   Weimer, P. J.; Nerdahl, M.; Brandl, D. J., Production of medium-chain volatile fatty acids
by mixed ruminal microorganisms is enhanced by ethanol in co-culture with Clostridium kluyveri.
Bioresour. Technol. **2015,** *175*, 97-101.

43.   Roghair, M.; Liu, Y.; Adiatma, J. C.; Weusthuis, R. A.; Bruins, M. E.; Buisman, C. J. N.;
Strik, D., Effect of n-Caproate Concentration on Chain Elongation and Competing Processes. *Acs
Sustain Chem Eng* **2018,** *6*, (6), 7499-7506.

44.   Liu, P.; Chernyshov, A.; Najdi, T.; Fu, Y.; Dickerson, J.; Sandmeyer, S.; Jarboe, L.,
Membrane stress caused by octanoic acid in Saccharomyces cerevisiae. *Appl. Microbiol.
Biotechnol.* **2013,** *97*, (7), 3239-51.

45.   Weghoff, M. C.; Bertsch, J.; Muller, V., A novel mode of lactate metabolism in strictly
anaerobic bacteria. *Environ. Microbiol.* **2015,** *17*, (3), 670-7.

46.   Juturu, V.; Wu, J. C., Microbial production of lactic acid: the latest development. *Crit. Rev.
Biotechnol.* **2016,** *36*, (6), 967-977.

47.   Buschke, N.; Schafer, R.; Becker, J.; Wittmann, C., Metabolic engineering of industrial
platform microorganisms for biorefinery applications--optimization of substrate spectrum and
process robustness by rational and evolutive strategies. *Bioresour. Technol.* **2013,** *135*, 544-54.

48.   Han, W.; He, P.; Shao, L.; Lu, F., Metabolic Interactions of a Chain Elongation
Microbiome. *Appl. Environ. Microbiol.* **2018**.

49.   Scarborough, M. J.; Myers, K. S.; Donohue, T. J.; Noguera, D. R., Multi-omic analysis of
medium-chain fatty acid synthesis by Candidatus Weimerbacter bifidus, gen. nov., sp. nov., and
Candidatus Pseudoramibacter fermentans, sp. nov. In University of Wisconsin- Madison: 2019.

50.   Zomorrodi, A. R.; Maranas, C. D., OptCom: a multi-level optimization framework for the
metabolic modeling and analysis of microbial communities. *PLoS Comput. Biol.* **2012,** *8*, (2),
e1002363.

51.   Kumar, M.; Ji, B.; Zengler, K.; Nielsen, J., Modelling approaches for studying the
microbiome. *Nat Microbiol* **2019,** *4*, (8), 1253-1267.
655

Figure 1. Summary of metabolic networks in iFerment156. Open hexagons indicate consumption

of the ATP (orange), NADH (green), and reduced ferredoxin (red). Closed hexagons indicate

production of these compounds and consumption of ADP (orange), NAD+ (green), and oxidized

ferredoxin (red). For reversible reactions, larger arrowheads indicate the direction of the indicated

compounds' consumption or production. Dashed lines indicate multiple reactions. Key reactions

are indicated according to the abbreviation used in iFerment156 (**Supplementary File 3**).

Pathways are identified by number: (1) upper glycolysis; (2) phosphoketolase metabolism of

fructose; (3) pentose phosphate; (4) phosphoketolase metabolism of xylulose; (5) glycerol utilization; (6) lower glycolysis; (7) lactate metabolism via NADH-dependent lactate dehydrogenase; (8) lactate metabolism via electron-confurcating lactate dehydrogenase; (9) pyruvate metabolism via pyruvate dehydrogenase; (10) acetate metabolism; (11) ethanol metabolism; (12) pyruvate metabolism by flavodoxin oxidoreductase; (13) pyruvate metabolism by formate lyase; (14) propionate production via the acryloyl-CoA pathway; (15) propionate production via the methylmalonyl-CoA pathway; (16) even-chain reverse β-oxidation; (17) odd-chain reverse β-oxidation; (18) ferredoxin-dependent hydrogenase; (19) proton-translocating hydrogenase; (20) electron confurcating hydrogenase; (21) the RNF complex; (22) homoacetogenesis via the Wood Ljungdahl pathway. More information on the reactions and metabolites contained in iFerment156 is provided in **Supplementary File 3**.

Figure 2. Summary of metabolic networks for six functional guilds contained in iFermGuilds564.

Refer to Fig. 1 for details on pathways included in the iFermGuilds564 guild models. Information

on the reactions and metabolites contained in the iFermGuilds564 guild models is provided in

Supplementary File 3.

Figure 3. Flux variability analysis results for iFerment156 maximizing ATP production from the indicated substrates. The maximum production of each end-product that can maintain the maximum ATP yield is indicated. For instance, when consuming glucose, iFerment156 predicts that production of formate, acetate, butyrate, hexanoate, octanoate, and H_2 can each sustain maximum ATP yields. However, if lactate is the sole substrate, only octanoate production can maintain the maximum ATP yield. The color intensity represents the percent of eeq that is predicted by iFerment156 to be converted to the noted end-product while maintaining maximum ATP production. The chart above the heatmap indicates the mol ATP generated per mol of substrate utilized. For scenarios with two growth substrates, the ATP yield is relative to the substrate indicated in bold text. Example model solutions are provided in Table S3.

Figure 4. Predicted and observed product formation for a mixed culture fermentation bioreactor

fed lignocellulosic biorefinery residues. Blue bars represent the range of products consistent with

maximum ATP production by the microbial community. Products represented by black lines

indicate there is a unique percent of eeq that must be converted to that compound. Closed black

circles represent the observed level of product accumulated in a bioreactor fed lignocellulosic

biorefinery residues (A) predictions with iFerment156; (B) predictions with iFerment156 with

constrained hexanoate and octanoate production; (C) predictions with iFermGuilds564; (D)

predictions with iFermGuilds564 with constrained hexanoate and octanoate production.

Figure 5. Predicted substrate flow through a mixed culture fermentation producing a mixture of MCFA from a complex feedstock using iFermGuilds564. Values were obtained using pFBA while constraining the products according to observed bioreactor performance. The relative abundance of each functional guild assumed based on reactor DNA sbundance is given in parentheses. The relative amount of eeq in the substrates, products, and intermediates are also provided. Line width indicates the relative amount of each substrate consumed and product produced by each of the functional guilds. H_2 was not measured, so for these simulations it was assumed that the difference between eeq in the substrate and eeq in the measured soluble and insoluble products represented H_2 production. In this simulation, 64% of the eeq in conversion residue was predicted to be directed to lactate as an intermediate.

716   **Figure 6.** iFermGuilds564 predicted fluxes through a community of SEOs and HAOs converting

717   lignocellulosic biorefinery residue and exogenous H_2 into octanoate. SEOs provide CO_2 and

718   formate to the HAOs, while the HAOs provide acetate to SEOs.